Matreex: compact and interactive visualisation of large gene families provides evidence for loss of intraflagellar transport in a myxozoan

Victor Rossier[1,2,3], Clement Train[1], Yannis Nevers[1,2], Marc Robinson-Rechavi[2,3] and Christophe Dessimoz[1,2,*]

[1]Department of Computational Biology, University of Lausanne, Switzerland; [2]SIB Swiss Institute of Bioinformatics, Lausanne, Switzerland; [3]Department of Ecology and Evolution, University of Lausanne, Switzerland;

*Corresponding author: Christophe.Dessimoz@unil.ch

Abstract

Studying gene family evolution strongly benefits from insightful visualisations. However, the ever-growing number of sequenced genomes is leading to increasingly larger gene families, which challenges existing gene tree visualisations. Indeed, most of them present users with a dilemma: display complete but intractable gene trees, or collapse subtrees, thereby hiding their children's information. Here, we introduce Matreex, a new dynamic tool to scale-up the visualisation of gene families. Matreex' key idea is to use "phylogenetic" profiles, which are dense representations of gene repertoires, to minimise the information loss when collapsing subtrees. We illustrate Matreex usefulness with three biological applications. First, using the textbook example of visual opsins, we show Matreex' potential to create easily interpretable figures for teaching and outreach. Secondly, by displaying 22 intraflagellar transport gene families across 622 species cumulating 5'500 representatives, we show how Matreex can be used to automate large-scale analyses of gene presence-absence. Notably, we report for the first time the complete loss of intraflagellar transport in the myxozoan *Thelohanellus kitauei*. Finally, we demonstrate on the MutS family the power of combining gene trees and phylogenetic profiles to delve into precise evolutionary analyses of large multi-copy gene families. Matreex is available from the Python Package Index (pip install matreex) with the source code and documentation available at https://github.com/DessimozLab/matreex.

Introduction

Studying the evolutionary dynamics of gene families strongly benefits from appropriate visualisation tools. For example, we can draw evolutionary and functional hypotheses by visually correlating gene repertoires with adaptations or between families. Moreover, visualising the evolutionary history of a gene family provides the framework to generalise classical pairwise gene relationships (*e.g.* orthology and paralogy) to multiple species (Dunn and Munro 2016). However, the growing number of genomes sequenced and processed by comparative genomic pipelines results in increasingly larger gene families. For example, the OMA database provides families with more than 100'000 members across more than 2'500 species (Altenhoff et al. 2021). Gene family visualisation tools able to integrate this large data volume are needed.

Gene trees labelled with duplications and speciations are typically used to depict the evolutionary history of gene families. However, existing gene tree viewers are not equipped to provide overviews of evolutionary trajectories required to study large gene families spanning thousands of taxa and dozens of subfamilies. To keep gene trees interpretable, most viewers merely rely on collapsing or trimming subtrees, by letting users dynamically expand the relevant ones, while collapsing others (Herrero et al. 2016; Mi et al. 2017; Nguyen et al. 2018; Fuentes et al. 2021). For example, the GeneView of Ensembl collapses by default all subtrees lying outside the lineage of the query gene and provides the option to collapse all nodes at a given taxonomic rank (Herrero et al. 2016). Similarly, the PhyloView of Genomicus displays the gene tree at a user-defined taxon and provides many customization features such as trimming outgroups (relative to the query gene) or duplication nodes (Nguyen et al. 2018). However, a collapsed or trimmed subtree is mostly uninformative, as its gene content and topology are not shown. Therefore, users can only choose between keeping a complete and often intractable gene tree, or collapsing nodes and hiding the information of its children, with no middle ground. Moreover, these viewers are limited by their slow reactivity, which makes the exploration of large gene trees cumbersome. For example, a couple of seconds is needed to collapse a node in Ensembl GeneView or PhylomeDB, while any action brings the user back to the top of the page in Genomicus PhyloView. Faster and more scalable web-based tools have been introduced to visualise large phylogenies of species or of viral genomes (Robinson et al. 2016; Turakhia et al. 2020), but they are not tailored to display gene families and also lack a way to summarise relevant information contained in the different relevant parts of the phylogenies.

Alternatively, gene families can be represented as vectors of gene copy numbers across species or phylogenetic profiles. Although these were initially developed to infer gene functions, as repeated co-occurrences provide evidence of interaction (Pellegrini et al. 1999), visualising these profiles has

proven useful to illustrate the gene content of extant species (Musilova et al. 2021; Horn et al. 2022)

or to compare likely coevolving families (van Dam et al. 2013; Nevers et al. 2017). Indeed, displaying

the full gene repertoire of a species in the same column (or row) and all gene family members in the

same row (or column) enables rapid visual identification of repeated and correlated gene presence

and absences. The relevance of this kind of compact representation of gene families is evidenced by

the large number of tools developed for that task (Sadreyev et al. 2015; Cromar et al. 2016; Tran et al.

2018; Tremblay et al. 2021; Ilnitskiy et al. 2022). However, unlike gene trees, phylogenetic profiles do

not show evolutionary relationships among the genes; for instance, it is not possible to deduct from a

profile alone whether two gene absences are the result of independent losses, or a single loss in a

common ancestor.

Here, we introduce Matreex, an innovative viewer for large gene families that bridges the gap

between these two typical representations of gene families: gene trees that provide their complete

evolutionary picture but can be cumbersome to read and phylogenetic profiles that efficiently depict

the distribution of genes across species but lack the evolutionary component. Matreex builds on the

reactive framework from the Phylo.IO viewer (Robinson et al. 2016) and integrates phylogenetic

profiles to summarise collapsed subtrees. Thus, it simplifies gene tree visualisation while reducing the

information loss. The resulting highly compact and reactive visualisation of evolution enables Matreex

to scale-up to the ongoing deluge of genomic data. Moreover, it provides the opportunity for new

biological discoveries, for the production of paper figures, and for didactic support for teaching in

evolutionary biology. We illustrate Matreex with three biological applications.

New Approach

Fig 1. Matreex' layout consists of a gene tree, a species tree and a matrix of phylogenetic profiles. Branch thickness increases with the collapsed subtree size and cell colour darkness with the number of gene copies in the cell. The taxonomic levels of collapsed subtrees and phylogenetic profiles are annotated on the left. "Auto-collapse at depth" enables the automatic collapse of the species tree at a given depth from the root. "Show species thumbnail" enables displaying a taxon image (at present from Wikipedia) when hovering over a taxon. "Collapse All" and "Smart collapse" are two default views described in the main text. The examples shown are red-sensitive visual opsins (data from OMA, All.Dec2021 release). Italic annotations do not belong to the Matreex layout but are added for figure clarity.

To enable compact and reactive visualisation of large gene families, Matreex complements the gene tree with a matrix of phylogenetic profiles and a species tree (Fig. 1). Thus, when collapsing a subtree to simplify the gene tree, the distribution of gene copy numbers across species remains available in the corresponding row. This keeps information about ancestral events such as gene loss, duplication or transfer. Moreover, the species tree displayed orthogonally from the gene tree provides what is often a good proxy for the topology of these subtrees (Morel et al. 2020). Remarkably, the extreme case, where all subtrees without duplications are collapsed, provides the same information as a fully extended gene tree but much more compact. This is achieved by removing most of the species tree redundancy from the gene tree. We provide rapid access to this view with Matreex' "Smart Collapse" option.

For genes families with a high number of duplication events, collapsing only subtrees without duplication is not enough and summarising them requires also collapsing subtrees with subfamilies (children of duplication nodes). In that case, the resulting phylogenetic profiles depict the combined

gene content of each subfamily per species. For example, the profile of the collapsed *Sarcopterygii*
subtree of the red-sensitive visual opsin (LWS) family shows four copies for humans due to the
existence of primates-specific subfamilies (Fig. 1). To deal with large gene families, Matreex includes
the option of collapsing all subfamilies, including the root node (Matreex' "Collapse All"), as manually
collapsing many nodes can be tedious. Starting from the family phylogenetic profile, the user can then
unfold more and more specific subfamilies, thus revealing their species distributions and gene copy
number variations. In particular, unfolding a node will reveal the gene tree topology until the next
duplication nodes, which define the child subfamilies. Other subtrees will remain collapsed, as their
topology is redundant with the species tree. This approach is user-friendly because it begins with a
highly summarised view of the family before zooming into more specific subfamilies of interest.

Two main processes increase the size of gene families in practice: gene duplications and the increase
in the number of species. The latter increases both with the number of available genomes and with
the progress of orthology assessment methods and resources in handling a growing number of species
(*e.g.* (Kriventseva et al. 2019; Altenhoff et al. 2021; Cantalapiedra et al. 2021; Rossier et al. 2021)).
Thus, the ability to control which species (or taxa) to show and which to hide is key to achieve high
levels of gene family customisation and compactness. For that task, Matreex provides control over
which taxa are displayed through its interactive species tree. When collapsing a taxon in the species
tree, all corresponding gene tree nodes are also collapsed, and the phylogenetic profiles are
summarised. This is done by averaging the numbers of gene copies of the species collapsed. For
example, collapsing the *Oryzias* node in the LWS family automatically merges *Oryzias*-specific
subfamilies and averages the copy numbers of *Oryzias* species (Fig. 1). Moreover, to facilitate the
exploration of large species trees, Matreex provides the option to collapse every taxon after a given
node depth from the root.

Finally, Matreex implements several other design features to further facilitate the user experience.
First, as scientific names can be quite obscure, images are displayed when hovering over taxon labels;
at present Wikipedia images are used but other sources could be easily implemented. Second, to
highlight lineage-specific expansions, the matrix of phylogenetic profiles is displayed as a heatmap for
which custom colours can be used to highlight specific clades.

Availability and implementation

Matreex is available from the Python Package Index (pip install matreex) with the source code, the
documentation   and   code   to   reproduce   the   following   figures   at
https://github.com/DessimozLab/matreex. Matreex is implemented in JavaScript with the D3 library

and wrapped in a Python module that supports the APIs of OMA and PANTHER (Kaleb et al. 2019; Mi
et al. 2021), from which only the gene family or HOG identifier is required as input. Moreover, custom
gene trees in JSON are supported when associated with a consistent species tree (gene tree-species
tree conflicts are not supported by Matreex). The specifics of these formats are described in the
GitHub.

Applications

In this section, we illustrate how Matreex facilitates the analysis of gene families on three different
use-cases with real biological applications.

The visual opsin gene repertoire correlates with adaptations in vertebrates

Figure 2. Visual opsin families in vertebrates (data from OMA All.Dec2021). Adaptations involved in textbook
correlations with patterns of gene losses and duplications are annotated with separate colours. Nocturnality in
snakes and mammals: loss of blue- and green-sensitive opsins. Frugivory in old-world primates: duplications of
red-sensitive opsin. Deep-water: loss of violet- and red-sensitive opsins, duplications of blue- and green-sensitive
opsins. Turbid water: duplications of red-sensitive opsins. Benthic: duplications of green-sensitive opsins. Italic
annotations do not belong to the Matreex layout but are added for figure clarity.

Matreex enables to quickly identify correlations between adaptations, or phenotypes, and variations
in gene copy numbers. Indeed, the gene repertoire of a species or taxon is shown in a single column
instead of being scattered across the gene tree. Thus, repeated losses or expansions associated with
an adaptation are easy to identify. Matreex further facilitates the task by depicting losses on a white
background and expansions on a darker one. Clades can also be coloured by adaptation to highlight
the correlation. Here, we illustrate this use-case with the textbook example of visual opsins (Graur and
Li 1999; Paul G. Higgs and Teresa K. Attwood 2005). Because Matreex representation facilitates an

intuitive interpretation of the data, we expect this usage to become particularly popular in outreach
tasks including teaching and conference presentations.

The vertebrate ancestor had one rod opsin (Rhodopsin) for dim light vision and four cone opsins for a
tetrachromatic vision, each sensitive to a specific range of light wavelength (Musilova et al. 2021).
Specifically, the shortest wavelengths are absorbed by the violet-sensitive opsin (SWS1), followed by
the blue- (SWS2) and green-sensitive (RH2) opsins for intermediate wavelengths. The red-sensitive
(LWS) opsins absorb for the largest ones. By contrast, mammals and snakes lack the blue- and green-
sensitive opsins, likely due to the nocturnal lifestyle of their ancestors (Borges et al. 2018; Katti et al.
2019). However, old-world primates (*Catarrhini*, including humans) regained a more complex colour
vision by co-opting a red-sensitive opsin duplicate to absorb green wavelengths, which possibly helped
primates to identify edible fruits (Carvalho et al. 2017). Matreex shows clearly and at a glance both
the losses in mammals and snakes (Fig. 2, pink), as series of white background zeroes, and the
secondary amplification in *Homo sapiens* and *Pongo abelii* (Fig. 2, green), as darker cells with larger
numbers of genes.

By contrast, the visual opsin repertoire of fishes is much more variable, likely due to the diversity of
underwater light environments (Musilova et al. 2021) and this is immediately visible in the Matreex
representation. In deep water, the light spectrum is shrinked to absorb only blue and green. Thus,
deeper-living species are expected to lose red- and violet- sensitive opsins, while duplicating the
green- and blue-sensitive ones to compensate for the lower photon abundance. Here, such an
evolutionary pattern was detected in the cod (*Gadus morhua*, depth: 150-200m, max. 600m), the
sunfish (*Mola mola*, depth 30-70m, max. 480m) and the coelacanth (*Latimeria chalumnae*, depth: 180-
250m, max. 700m) (Fig. 2, purple). Moreover, we found the most green-sensitive opsins (five) in the
turbot flatfish (*Scophthalmus Maximus*), which could be an adaptation to deep benthic life (Wang et
al. 2021) (Fig. 2, orange). Conversely, the light spectrum is shifted toward longer wavelengths in turbid
water, thus favouring red-opsin duplications (Musilova et al. 2021). The present gene tree supports
this assumption as we found the most red-opsin copies in fishes that live in the turbid freshwater and
brackish habitats (Fig. 2, brown). In particular, five copies were detected in the brown trout (*Salmo
trutta*) and four in the red piranha (*Pygocentrus nattereri*), the Atlantic salmon (*Salmo salar*), the
northern pike (*Esox lucius*), the guppy (*Poecilia reticulata*) and the pupfish (*Cyprinodon variegatus*).

Coevolution of the intraflagellar transport genes

Figure 3. Intraflagellar transport gene families (data from OMA All.Dec2021). Coloured clades display partial and complete IFT losses that fit the "last-in, first-out" hypothesis for gene module evolution. ✓ highlight partial IFT losses reported by van Dam et al. (2013) and **?**, the ones reported here. We also report the first evidence to our knowledge of a complete loss of IFT in the mixozoan *Thelohanellus kitauei*. Italic annotations, brackets, ✓ and **?** do not belong to the Matreex layout but are added for figure clarity.

Matreex enables to perform gene presence-absence analyses for dozens of non-homologous families spanning hundreds of species in a few minutes. Indeed, similarly to phylogenetic profile viewers, lists of gene families are valid inputs for Matreex. This is useful to visualise the result of a phylogenetic profile search (Altenhoff et al. 2021) or to study coevolving gene families (*e.g.* involved in the same pathway). In this second application, we illustrate the latter by generalising a study on eukaryotic intraflagellar transport (IFT) genes from 622 species, compared to the 52 used originally (van Dam et al. 2013). Specifically, we used Matreex to simplify the task of contrasting our results with the literature and to propose new biological hypotheses.

Eukaryotic flagella (cilia) are involved in cell motility and sensory detection (Nevers et al. 2017). Their dysfunction is the cause of ciliopathies in humans (Badano et al. 2006). The IFT complex is essential to build and maintain the flagella. From an evolutionary perspective, IFT is a great example of the "last-in, first-out" hypothesis (van Dam et al. 2013), whereby modules added last are more dispensable and thus, lost first. Indeed, of the three IFT modules (IFT-α, IFT-β and BBSome), BBSome and IFT-α emerged from IFT-β duplications and their loss often precedes the complete loss of IFT and cilia. Thus, studying

how ciliated eukaryotes cope with partial IFT loss is promising for the treatment of IFT-related human ciliopathies such as the Bardet–Biedl syndrome caused by BBSome alterations (Badano et al. 2006).

Repeated and correlated losses are visible at a glance in Matreex with columns of zeros on white backgrounds (Fig 3). As expected, complete loss of IFT complexes were detected in the main non-ciliated taxa (*e.g. Spermatophyta*, *Dikarya* or *Amoebozoa*). Moreover, due to the sheer number of used genomes, summarised in one easy to read figure, we were able to identify many other complete IFT losses. Although most were already established (*e.g. Fonticula alba*, *Creolimax fragrantissima*, *Capsaspora owczarzaki* (Torruella et al. 2015), *Entamoeba* (Wickstead and Gull 2007)), we report the first evidence to our knowledge of a complete loss of IFT in the mixozoan *Thelohanellus kitauei*, likely indicating the loss of the organelle in this species.

Then, we could first quickly confirm all established patterns of BBSome and IFT-α losses in species closely related to non-ciliated clades with complete IFT loss from (van Dam et al. 2013). Specifically, we detected the loss of BBSome in basal plants (*Selaginella moellendorffii* and the moss *Physcomitrella patens*) close to seed plants (*Spermatophyta*), in the apicomplexa *Sarcocystidae* (*Toxoplasma gondii* clade) close to *Aconoidasida* (*Plasmodium falciparum* clade) and in the basal fungi *Chytridiomycetes* (*Batrachochytrium dendrobatidis* clade) close to *Dikarya* and *Mucoromycota*. We also recovered the loss of BBSome and IFT-α in the diatoms *Thalassiosira* close to *Phaeodactylum tricornutum*. Secondly, we could identify other independent losses supporting the "last-in, first-out" hypothesis. In particular, we found two losses of BBSome in the basal plants *Marchantia polymorpha* and *Klebsormidium flaccidum*. We also identified complete IFT losses in another three apicomplexa clades (*Eimeria*, *Cryptosporidium* and *Gregarina niphandrodes*) and two basal fungi clades (*Microsporidia* and *Conidiobolus coronatus*). Moreover, we found evidence for losses of BBSome and IFT-α in four basal fungi. While *Rozella allomycis* lacks all BBSome and IFT-α genes, *Piromyces sp.* and *Gonapodya prolifera* were found with merely one IFT-α and two BBSome genes, respectively. *Allomyces macrogynus* lacked BBSome. Finally, the presence of one IFT-α and two IFT-β genes in the chlorophytes *Chlorella variabilis* close to *Ostreococcus* provides a new candidate replicate for this "last-in, first-out" hypothesis. Its low number of IFT genes, which indicates dysfunctional cilia, could be due to the endosymbiont nature of *Chlorella variabilis* (Blanc et al. 2010).

When many gene families underwent duplications in the same species, the column attracts the eye as it becomes darker in Matreex. Thus, we identified four species with many duplicates of IFT-α and IFT-β genes. Although *Paramecium tetraurelia* and *Trichomonas vaginalis* have undergone whole genome duplications (Aury et al. 2006; Carlton et al. 2007), *Paramecium tetraurelia* IFT57 copies show evidence of subfunctionalization (Shi et al. 2018), while *Trichomonas vaginalis* displays specialised cilia that

could have required the recruitment of additional IFT copies. Finally, to explain the retention of *S. moellendorffii* and *Allomyces macrogynus* duplicates, that have lost BBSome, we may speculate whether these extra copies could have been co-opted to replace the BBSome functions.

Origin and evolution of eukaryotic MutS genes

Figure 4. Detailed evolutionary analysis of the MutS family (data from PANTHER v.17). Established hypotheses on MutS evolution are annotated with a ✓ when supported by the data at hand and with a ✗ otherwise. Ongoing and new hypotheses are annotated with **?** and **??**, respectively. Italic annotations do not belong to the Matreex layout.

Matreex enables users to precisely analyse the gene repertoire evolution of large multi-copy families. First, subfamily gene repertoires can be correlated among themselves or with adaptations similarly to the previous applications. Secondly, the gene tree enables to study the evolutionary relationships between phylogenetic profiles. This can be useful, for instance, to differentiate orthologous from paralogous profiles and to diagnose the underlying gene tree. In this last application, we performed a detailed analysis of the MutS family, whose evolutionary history remains largely under debate. Specifically, we used Matreex to simplify the task of systematically contrasting existing knowledge with the data at hand (Fig. 4). First, we used established hypotheses to diagnose the underlying gene tree. Secondly, we assessed which ongoing hypotheses were supported by it and, thirdly, we drew new hypotheses from visual patterns. Finally, we contextualised the results with functional and evolutionary knowledge from the literature to highlight the importance of such an approach.

MutS genes are involved in the DNA mismatch repair pathway (Liu et al. 2017; Mi et al. 2021). Although
bacteria have multiple MutS genes, only MutS1 and MutS2 are found in both bacteria and eukaryotes.
MutS2 is found only in photosynthetic eukaryotes and its transfer from cyanobacteria through the
chloroplast endosymbiosis is well established (Lin et al. 2007). Matreex clearly shows the absence of
MutS2 in other eukaryotes and *Archaea*, as well as its two copies in plants (*Streptophyta*). However,
PANTHER predicts a vertical origin of MutS2 from a pre-LUCA duplication followed by independent
losses in *Archaea* and non-photosynthetic eukaryotes. Matreex shows this evolutionary trajectory
with a fully connected phylogenetic profile for MutS2 and losses instead of empty cells for *Archaea*
and most eukaryotes.

In contrast to MutS2, the origin of MutS1 remains under debate. Eukaryotic MutS1 genes (MSH2-7)
were first thought to originate from the mitochondria endosymbiosis of an α-*Proteobacteria* (Lin et al.
2007) until the Asgard *Archaea* MutS1 was found to be more closely related to Eukaryotes than to α-
Proteobacteria (Hofstatter and Lahr 2021). This implies the vertical origin of MSH2-7 from *Archaea*.
Matreex clearly shows these orthologous relationships between archeal MutS1 and eukaryotic MSH2-
7 because they form a monophyletic clade in the gene tree and their profiles do not overlap.

MSH2-6 genes originated from duplications in the eukaryote ancestor, while MSH7 arose from a plant-
specific duplication of MSH6 (Lin et al. 2007). Matreex clearly represents this radiation with a compact
block of subfamily profiles, although PANTHER misclassified MSH7 as another eukaryote subfamily.
This radiation presents a striking example of functional evolution through gene duplications. Indeed,
while bacterial MutS genes form homodimers, the eukaryotic DNA mismatch repair pathway recruits
three heterodimers that have specialised to bind specific DNA mismatches. Precisely, the mammal
MSH2/6 and MSH2/3 bind up to 3 and 13 nucleotide indels, respectively (Muthye and Lavrov 2021),
while the plant MSH2/7 prefers mismatches containing T and/or G/T, A/C, T/C, G/A, T/T, or A/A
(Karthika et al. 2020). The functional specialisation of these MutS1 homologs, both at the subunit and
heterodimer levels, probably improved the DNA mismatch repair system in eukaryotes. Moreover,
MSH4 and MSH5 have lost the DNA repair function and are involved in meiotic recombination, thus
presenting a striking example of neofunctionalization (Manhart and Alani 2016).

However, the origins of the plant- and yeast-MSH1 remain unclear. Although originally thought to
descend from the same MutS ancestor as MSH2-7 (Lin et al. 2007), an acquisition of yeast-MSH1 in
fungi along the mitochondrial endosymbiosis has been suggested (Hofstatter and Lahr 2021). Similarly,
the plant-MSH1 could have been acquired from giant viruses or along the chloroplast endosymbiosis
(Wu et al. 2020; Hofstatter and Lahr 2021). The underlying gene tree supports the original hypothesis
as we found the plant- and yeast-MSH1 genes in the same eukaryotic subfamily. Matreex simplified

drawing this conclusion as both plant- and yeast-MSH1 belong to the same collapsed subtree and

phylogenetic profile, indicating a monophyletic origin. Moreover, we recovered the absence of MSH1

in animals (with the exception of the tick), which has been recently linked with the exceptionally high

evolutionary rates of their mitochondrial genes, as MSH1 is involved in repairing their sequences (Wu

et al. 2020).

Matreex simplifies the identification of gene repertoire evolutionary patterns. Thus, we observed

unexpected expansions of MSH6 in eukaryotes (*e.g. Alveolata*, *Trypanosomatidae*), yeast

(*Saccharomyces cerevisiae*), nematodes and fruitfly (*Drosophila Melanogaster*). Then, by expanding

the gene tree, we noticed many MSH3 genes misclassified as MSH6, which coincides with predicted

MSH3 losses. For example, of the five *Trypanosomatidae* copies, two were surely misclassified MSH3

and MSH5 genes, and one was undefined. Thus, although the loss of MSH3 in nematodes and insects

and the trypanosome-specific MSH8 subfamily are documented (Bell et al. 2004; Muthye and Lavrov

2021), we hypothesised that MSH6 is artefactually attracting other genes, in particular MSH3 ones,

during phylogenetic reconstruction.

Finally, we observed repeated and correlated losses of MSH4 and MSH5 in fungi, fruit fly and other

eukaryotes (annotated in pink). While losses in the latter are likely artefactual (Rzeszutek et al. 2022),

Schizosaccharomyces and *D. Melanogaster* are known to have lost and replaced MSH4 and MSH5 for

meiotic recombination (Kohl et al. 2012; Manhart and Alani 2016). Moreover, given that these two

genes form an obligate complex, other correlated losses in fungi are plausible and could provide good

candidates to study alternative meiotic recombination mechanisms.

Conclusion

At a time when the goal of sequencing all eukaryotic species before 2030 has been set (Lewin et al.

2022), it has become critical to develop new methods to represent this huge volume of upcoming

data. Here, we introduced an innovative tool to scale the visualisation of gene families and illustrate

its usefulness with three biological applications. First, using the textbook example of visual opsins, we

demonstrated Matreex' potential to create easily interpretable figures for outreach tasks. Secondly,

by displaying 22 Intraflagellar gene families across 622 species cumulating 5'500 representatives, we

showed how Matreex can be used for analyses of gene presence-absence and reported for the first

time the complete loss of IFT in the mixozoan *Thelohanellus kitauei*. Finally, we demonstrated

Matreex' usefulness in delving into precise evolutionary analyses of multi-copy gene families by

combining the gene tree with phylogenetic profiles. Thus, we hope Matreex will become a valuable

tool to gain insights into the evolution of increasingly large gene families.

Funding

This work was supported by the Swiss National Foundation [167276] as part of the National Research Program 75 'Big Data', as well as Swiss National Foundation [205085].

References

Altenhoff AM, Train C-M, Gilbert KJ, Mediratta I, Mendes de Farias T, Moi D, Nevers Y, Radoykova H-S, Rossier V, Warwick Vesztrocy A, et al. 2021. OMA orthology in 2021: website overhaul, conserved isoforms, ancestral gene order and more. *Nucleic Acids Res.* 49:D373–D379.

Aury J-M, Jaillon O, Duret L, Noel B, Jubin C, Porcel BM, Ségurens B, Daubin V, Anthouard V, Aiach N, et al. 2006. Global trends of whole-genome duplications revealed by the ciliate Paramecium tetraurelia. *Nature* 444:171–178.

Badano JL, Mitsuma N, Beales PL, Katsanis N. 2006. The ciliopathies: an emerging class of human genetic disorders. *Annu. Rev. Genomics Hum. Genet.* 7:125–148.

Bell JS, Harvey TI, Sims A-M, McCulloch R. 2004. Characterization of components of the mismatch repair machinery in Trypanosoma brucei. *Mol. Microbiol.* 51:159–173.

Blanc G, Duncan G, Agarkova I, Borodovsky M, Gurnon J, Kuo A, Lindquist E, Lucas S, Pangilinan J, Polle J, et al. 2010. The Chlorella variabilis NC64A genome reveals adaptation to photosymbiosis, coevolution with viruses, and cryptic sex. *Plant Cell* 22:2943–2955.

Borges R, Johnson WE, O'Brien SJ, Gomes C, Heesy CP, Antunes A. 2018. Adaptive genomic evolution of opsins reveals that early mammals flourished in nocturnal environments. *BMC Genomics* 19:121.

Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021. eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* [Internet]. Available from: http://dx.doi.org/10.1093/molbev/msab293

Carlton JM, Hirt RP, Silva JC, Delcher AL, Schatz M, Zhao Q, Wortman JR, Bidwell SL, Alsmark UCM, Besteiro S, et al. 2007. Draft genome sequence of the sexually transmitted pathogen Trichomonas vaginalis. *Science* 315:207–212.

Carvalho LS, Pessoa DMA, Mountford JK, Davies WIL, Hunt DM. 2017. The Genetic and Evolutionary Drives behind Primate Color Vision. *Frontiers in Ecology and Evolution* 5:34.

353   Cromar GL, Zhao A, Xiong X, Swapna LS, Loughran N, Song H, Parkinson J. 2016. PhyloPro2.0: a

354   database for the dynamic exploration of phylogenetically conserved proteins and their domain

355   architectures across the Eukarya. *Database* [Internet] 2016. Available from:

356   http://dx.doi.org/10.1093/database/baw013

357   van Dam TJP, Townsend MJ, Turk M, Schlessinger A, Sali A, Field MC, Huynen MA. 2013. Evolution of

358   modular intraflagellar transport from a coatomer-like progenitor. *Proc. Natl. Acad. Sci. U. S. A.*

359   110:6943–6948.

360   Dunn CW, Munro C. 2016. Comparative genomics and the diversity of life. *Zool. Scr.* 45:5–13.

361   Fuentes D, Molina M, Chorostecki U, Capella-Gutiérrez S, Marcet-Houben M, Gabaldón T. 2021.

362   PhylomeDB V5: an expanding repository for genome-wide catalogues of annotated gene

363   phylogenies. *Nucleic Acids Res.* [Internet]. Available from:

364   http://dx.doi.org/10.1093/nar/gkab966

365   Graur, Li. 1999. Fundamentals of Molecular Evolution, 2nd edn Sinauer Associates. *Inc, Sunderland,*

366   *Massachusetts, USA*.

367   Herrero J, Muffato M, Beal K, Fitzgerald S, Gordon L, Pignatelli M, Vilella AJ, Searle SMJ, Amode R,

368   Brent S, et al. 2016. Ensembl comparative genomics resources. *Database* [Internet] 2016.

369   Available from: http://dx.doi.org/10.1093/database/bav096

370   Hofstatter PG, Lahr DJG. 2021. Complex Evolution of the Mismatch Repair System in Eukaryotes is

371   Illuminated by Novel Archaeal Genomes. *J. Mol. Evol.* 89:12–18.

372   Horn T, Narov KD, Panfilio KA. 2022. Persistent parental RNAi in the beetle *Tribolium castaneum*

373   involves maternal transmission of long double-stranded RNA. *Advanced Genetics*:2100064.

374   Ilnitskiy IS, Zharikova AA, Mironov AA. 2022. OrthoQuantum: visualizing evolutionary repertoire of

375   eukaryotic proteins. *Nucleic Acids Res.* [Internet]. Available from:

376   http://dx.doi.org/10.1093/nar/gkac385

377   Kaleb K, Vesztrocy AW, Altenhoff A, Dessimoz C. 2019. Expanding the Orthologous Matrix (OMA)

378   programmatic interfaces: REST API and the OmaDB packages for R and Python. *F1000Res.* 8:42.

379   Karthika V, Babitha KC, Kiranmai K, Shankar AG, Vemanna RS, Udayakumar M. 2020. Involvement of

380   DNA mismatch repair systems to create genetic diversity in plants for speed breeding programs.

381   *Plant Physiology Reports* 25:185–199.

Katti C, Stacey-Solis M, Coronel-Rojas NA, Davies WIL. 2019. The Diversity and Adaptive Evolution of

Visual Photopigments in Reptiles. *Frontiers in Ecology and Evolution* [Internet] 7. Available from:

https://www.frontiersin.org/article/10.3389/fevo.2019.00352

Kohl KP, Jones CD, Sekelsky J. 2012. Evolution of an MCM complex in flies that promotes meiotic

crossovers by blocking BLM helicase. *Science* 338:1363–1365.

Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, Zdobnov EM. 2019. OrthoDB

v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for

evolutionary and functional annotations of orthologs. *Nucleic Acids Res.* 47:D807–D811.

Lewin HA, Richards S, Lieberman Aiden E, Allende ML, Archibald JM, Bálint M, Barker KB,

Baumgartner B, Belov K, Bertorelle G, et al. 2022. The Earth BioGenome Project 2020: Starting

the clock. *Proc. Natl. Acad. Sci. U. S. A.* [Internet] 119. Available from:

http://dx.doi.org/10.1073/pnas.2115635118

Lin Z, Nei M, Ma H. 2007. The origins and early evolution of DNA mismatch repair genes--multiple

horizontal gene transfers and co-evolution. *Nucleic Acids Res.* 35:7591–7603.

Liu D, Keijzers G, Rasmussen LJ. 2017. DNA mismatch repair and its many roles in eukaryotic cells.

Mutat. Res. - Rev. Mut. Res. 773:174–187.

Manhart CM, Alani E. 2016. Roles for mismatch repair family proteins in promoting meiotic crossing

over. *DNA Repair* 38:84–93.

Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, Thomas PD. 2021. PANTHER

version 16: a revised family classification, tree-based classification tool, enhancer regions and

extensive API. *Nucleic Acids Res.* 49:D394–D403.

Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, Thomas PD. 2017. PANTHER version 11:

expanded annotation data from Gene Ontology and Reactome pathways, and data analysis tool

enhancements. *Nucleic Acids Res.* 45:D183–D189.

Morel B, Kozlov AM, Stamatakis A, Szöllősi GJ. 2020. GeneRax: A tool for species tree-aware

maximum likelihood based gene family tree inference under gene duplication, transfer, and

loss. *Mol. Biol. Evol.* [Internet]. Available from: http://dx.doi.org/10.1093/molbev/msaa141

Musilova Z, Salzburger W, Cortesi F. 2021. The Visual Opsin Gene Repertoires of Teleost Fishes:

Evolution, Ecology, and Function. *Annu. Rev. Cell Dev. Biol.* 37:441–468.

Muthye V, Lavrov DV. 2021. Multiple Losses of MSH1, Gain of mtMutS, and Other Changes in the

MutS Family of DNA Repair Proteins in Animals. *Genome Biol. Evol.* [Internet] 13. Available

from: http://dx.doi.org/10.1093/gbe/evab191

Nevers Y, Prasad MK, Poidevin L, Chennen K, Allot A, Kress A, Ripp R, Thompson JD, Dollfus H, Poch

O, et al. 2017. Insights into Ciliary Genes and Evolution from Multi-Level Phylogenetic Profiling.

Mol. Biol. Evol. 34:2016–2034.

Nguyen NTT, Vincens P, Roest Crollius H, Louis A. 2018. Genomicus 2018: karyotype evolutionary

trees and on-the-fly synteny computing. *Nucleic Acids Res.* 46:D816–D822.

Paul G. Higgs and Teresa K. Attwood. 2005. Bioinformatics and Molecular Evolution. BLACKWELL

PUBLISHING

Pellegrini M, Marcotte EM, Thompson MJ, Eisenberg D, Yeates TO. 1999. Assigning protein functions

by comparative genome analysis: protein phylogenetic profiles. *Proc. Natl. Acad. Sci. U. S. A.*

96:4285–4288.

Robinson O, Dylus D, Dessimoz C. 2016. Phylo.io: Interactive Viewing and Comparison of Large

Phylogenetic Trees on the Web. *Mol. Biol. Evol.* 33:2163–2166.

Rossier V, Vesztrocy AW, Robinson-Rechavi M, Dessimoz C. 2021. OMAmer: tree-driven and

alignment-free protein assignment to subfamilies outperforms closest sequence approaches.

Bioinformatics [Internet]. Available from: http://dx.doi.org/10.1093/bioinformatics/btab219

Rzeszutek I, Swart EC, Pabian-Jewuła S, Russo A, Nowacki M. 2022. Early developmental, meiosis-

specific proteins - Spo11, Msh4-1, and Msh5 - Affect subsequent genome reorganization in

Paramecium tetraurelia. *Biochim. Biophys. Acta Mol. Cell Res.* 1869:119239.

Sadreyev IR, Ji F, Cohen E, Ruvkun G, Tabach Y. 2015. PhyloGene server for identification and

visualization of co-evolving proteins using normalized phylogenetic profiles. *Nucleic Acids Res.*

43:W154–W159.

Shi L, Koll F, Arnaiz O, Cohen J. 2018. The Ciliary Protein IFT57 in the Macronucleus of Paramecium. *J.*

Eukaryot. Microbiol. 65:12–27.

Torruella G, de Mendoza A, Grau-Bové X, Antó M, Chaplin MA, del Campo J, Eme L, Pérez-Cordón G,

Whipps CM, Nichols KM, et al. 2015. Phylogenomics Reveals Convergent Evolution of Lifestyles

in Close Relatives of Animals and Fungi. *Curr. Biol.* 25:2404–2410.

Tran N-V, Greshake Tzovaras B, Ebersberger I. 2018. PhyloProfile: dynamic visualization and

exploration of multi-layered phylogenetic profiles. *Bioinformatics* 34:3041–3043.

Tremblay BJ-M, Lobb B, Doxey AC. 2021. PhyloCorrelate: inferring bacterial gene-gene functional

associations through large-scale phylogenetic profiling. *Bioinformatics* [Internet]. Available

from: http://dx.doi.org/10.1093/bioinformatics/btaa1105

Turakhia Y, De Maio N, Thornlow B, Gozashti L, Lanfear R, Walker CR, Hinrichs AS, Fernandes JD,

Borges R, Slodkowicz G, et al. 2020. Stability of SARS-CoV-2 phylogenies. *PLoS Genet.*

16:e1009175.

Wang Y, Zhou L, Wu L, Song C, Ma X, Xu S, Du T, Li X, Li J. 2021. Evolutionary ecology of the visual

opsin gene sequence and its expression in turbot (Scophthalmus maximus). *BMC Ecol Evol*

21:114.

Wickstead B, Gull K. 2007. Dyneins across eukaryotes: a comparative genomic analysis. *Traffic*

8:1708–1721.

Wu Z, Waneka G, Broz AK, King CR, Sloan DB. 2020. MSH1 is required for maintenance of the low

mutation rates in plant mitochondrial and plastid genomes. *Proc. Natl. Acad. Sci. U. S. A.*

117:16448–16455.