# Machine learning inference of continuous single-cell state transitions during myoblast differentiation and fusion

Amit Shakarchy[1*], Giulia Zarfati[2*], Adi Hazak[2*], Reut Mealem[1], Karina Huk[2], Tamar Ziv[3], Ori Avinoam[2&], Assaf Zaritsky[1&]

[1] Department of Software and Information Systems Engineering, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel

[2] Department of Biomolecular Sciences, Weizmann Institute of Science, Rehovot 761001, Israel

[3] The Smoler Proteomics Center, Lorry I. Lokey Interdisciplinary Center for Life Sciences and Engineering, Technion Israel Institute of Technology, Haifa 3200003, Israel

* Equal contribution

& Co-corresponding authorship

Assaf Zaritsky, assafzar@gmail.com

Ori Avinoam, ori.avinoam@weizmann.ac.il

## **Abstract**

Cells modifying their internal organization during continuous state-transitions, supporting functions from cell division to differentiation. However, tools to measure dynamic physiological states of individual transitioning cells are lacking. We combined live-cell imaging and machine learning to monitor ERK1/2-inhibited primary murine skeletal muscle precursor cells, that transition rapidly and robustly from proliferating myoblasts to post-mitotic myocytes and then fuse, forming multinucleated myotubes. Our model, trained using motility and actin intensity features from single cell tracking data, effectively tracked real-time continuous differentiation, revealing that differentiation occurs 7.5-14.5 hours post-induction, followed by fusion ~3 hours later. Co-inhibition of ERK1/2 and p38 led to differentiation without fusion. Our model inferred co-inhibition leads to terminal differentiation, indicating that p38 is specifically required for transitioning from terminal differentiation to fusion. Our model also predicted that co-inhibition leads to changes in actin dynamics. Mass spectrometry supported these in silico predictions and suggested novel fusion and maturation regulators downstream of differentiation. Collectively, this approach can be adapted to various biological processes to uncover novel links between dynamic single-cell states and their functional outcomes.

## Introduction

Single-cell transitions via dynamic changes in protein expression, intracellular organization, morphology, and function, drive many important biological processes, such as the progression through the phases of the cell cycle, cellular differentiation, the transition from an immotile to a motile state, or from a living to an apoptotic state. Aberrant cell state transitions lead to various diseases, including cancer and neuromuscular disorders. As such, single-cell state transitions play an inherent role in physiological processes such as embryonic development, tissue regeneration, and in various pathologies.

Obtaining a holistic mechanistic understanding of these processes relies on the ability to continuously measure the physiological state of a cell through time. However, technical limitations, such as the number of live fluorescent state transition reporters that can be simultaneously imaged, hinder the elucidation of cell state transitions as continuous processes. Moreover, state markers that could provide a continuous description are unknown for many biological processes. Consequently, we are currently limited to studying discrete cell states with missing intermediate states, which are often critical (Stumpf et al., 2017; Szkalisity et al., 2021).

Attempts to quantitatively follow cell state dynamics have focused on computational construction of "pseudo-time" trajectories from the integration of fixed cell images (Eulenberg et al., 2017; Gut et al., 2015; Rappez et al., 2020; Stallaert et al., 2022; Szkalisity et al., 2021; Yang et al., 2020). However, the capacity to identify single cell trajectories that deviate from the most common progression, is limited in this approach, due to heterogeneity (Schroeder, 2011). Live-cell imaging offers a solution to this challenge by enabling dynamic monitoring and extraction of temporal information at the single-cell resolution, with the caveat that unsupervised modeling may consider extrinsic factors that are unrelated to the state transition, which may confound proper modeling of the continuous process (Copperman et al., 2021; Wang et al., 2022; Wu et al. 2022).

The formation of multinucleated muscle fibers is an essential state-transition for vertebrate muscle development and regeneration. Following injury or growth stimuli, quiescent muscle progenitors called Satellite cells become activated to augment the muscle. At the onset of this process, activated satellite cells (myoblasts) express myogenic regulatory factors such as MyoD and proliferate to generate the myogenic progenitors needed for muscle regeneration (Bischoff, 1986;

Hurme & Kalimo, 1992; Schmidt et al., 2019). Next, myoblasts upregulate the expression of

factors such as Myogenin (MyoG) to exit the cell cycle and initiate terminal differentiation

(Hernández-Hernández et al., 2017; Lepper et al., 2011; Singh & Dilworth, 2013). Myoblasts

initially differentiate into elongated fusion-competent myocytes that migrate, adhere, and fuse with

the regenerating muscle fibers (Abmayr & Pavlath, 2012). Newly formed myofibers are

characterized by the expression of myosin heavy chain (MyHC; (Bentzinger et al., 2012; Lepper

et al., 2011; Yin et al., 2013).

Although significant progress has been made in understanding muscle development, myoblast

differentiation and fusion, remain incompletely understood at the molecular and cellular levels

owing to several technical challenges. First, myoblasts differentiation and fusion are complex

heterogeneous events, confounding systematic investigation. Second, while proliferating and

terminally differentiated cells are relatively easy to distinguish morphologically, there are no

markers for intermediate stages of differentiation or means for correlating between differentiation

state and specific markers or functions such as motility, morphology, or signaling.

In this study, we combined live cell imaging with supervised machine learning to quantitatively

monitor the differentiation state of individual skeletal muscle precursor cells as they differentiate

and fuse to form multinucleated muscle fibers *ex vivo*. Leveraging single-cell trajectories from

time-series data, we trained machine-learning models to evaluate the differentiation state of these

cells. Specifically, differentiation was initiated by pharmacologically inhibiting extracellular

signal-regulated kinases (ERK1/2), leading to differentiation and myotube formation within 18-24

hours (Eigler et al., 2021). Our models, which were trained using cell motility and/or actin intensity

time-series derived features, distinguished between proliferating myoblasts and terminally

differentiated myocytes. We hypothesized that the models' score can serve as a continuous readout

for the cells' differentiation state and proceeded to validate this hypothesis and explore its potential

for uncovering novel biological insights. We found that the average differentiation score

maintained a steady increase between 7.5 and 14.5 hours post-induction, mirroring the continuous

state transition from post-mitotic undifferentiated myoblasts to terminally differentiated myocytes.

Additionally, the predicted differentiation state at the single-cell level correlated with the time of

fusion, suggesting that differentiation and fusion are sequential and coordinated yet distinct

processes.

Building on this foundation, we employed a combined pharmacological approach, inhibiting both
ERK1/2 and p38. This led to an apparent accumulation of terminally differentiated, yet unfused
cells. Our model deduced that these cells have transitioned to terminal differentiation and implied
that co-inhibition leads to changes in actin dynamics. This inference by our model was
experimentally validated using both immunofluorescence and mass spectrometry. Mass
spectrometry also revealed a group of proteins significantly downregulated in co-inhibited unfused
cells compared to those differentiated by ERK1/2 inhibition. Notably, the expression of the fusion
specific protein, Myomixer (Zhang et al., 2017, Quinn et al., 2017, Bi et al., 2017) was abolished
in co-inhibited cells. In addition, several actin regulators were up-regulated in ERK1/2 inhibited
compared to cell where both ERK1/2 and p38 were inhibited, confirming our model's prediction
regarding alteration in the actin machinery. Our findings demonstrate the potential to
mechanistically uncouple differentiation from fusion, emphasizing the pivotal role of p38 in
orchestrating the transition from differentiation to fusion, and revealing potential novel regulators
of the late differentiation and fusion. Collectively, our findings demonstrate a method to quantify
cellular state transitions that can be adapted to other continuous processes.

# Results

**Differentiation correlates with reduced motility and increased actin intensity**

We previously established that ERK1/2 inhibition induced robust, faster and less temporally variable differentiation and fusion of primary myoblasts isolated from chick and mice, leading to the rapid formation of myotubes *ex vivo* within 24 hours post-induction (Eigler et al., 2021). Immunofluorescence staining of the differentiation markers MyoG and MyHC at different time points in cultures treated with the ERK1/2 inhibitor SCH772984 (ERKi, 1 μM) or with DMSO as control (Morris et al., 2013) show that differentiation into fusion-competent myocytes is accompanied by the upregulation of MyoG, which initiates terminal differentiation (Fig. 1A and Fig. S1 and Fig. S2). MyHC expression begins 16h post-induction and peaks in the multinucleated myotubes at 24hr (Fig. 1A and Fig. S2). The number of MyoG-expressing cells increases over time, stabilizing at 14 hours post-induction (Fig. 1B, Fig. S1 and Fig. S2).

To characterize the dynamic behavior of differentiating myoblasts, we isolated primary myoblasts from mice co-expressing the nuclear marker tdTomato fused to a nuclear localization signal (tDTomato-NLS) (Prigge et al., 2013) and the F-actin marker LifeAct-EGFP (Riedl et al., 2010 (Fig. 1C). Actin governs a range of processes during differentiation, from cell motility, elongation, and alignment to fusion and fibrillogenesis, ultimately leading to sarcomere formation post-fusion and the subsequent maturation of myotubes into myofibers (Rubinstein et al., 1976, Otey et al., 1988, Nowak et al., 2009, Luo et al., 2022).

To collect dynamic information on the continuous transition from proliferation to terminal differentiation we performed time-lapse widefield microscopy of large fields of view each containing approximately 3000 cells (Fig. 1C and Video S1). Cultures were imaged for 23 hours, starting 1.5 hours after ERKi or DMSO (control) treatment. We observed that differentiation was accompanied by a decrease in cell motility, consistent with previous studies showing that myocytes are less motile than myoblasts *ex vivo* (Powell, 1973; Griffin et al., 2010) (Fig. 1D and Video S1). The regulated reduction in cellular motility is beneficial for enhancing the cell-cell interactions that would initiate differentiation and subsequently promote fusion (Krauss et al., 2005; Buggenthin et al., 2017, Nowak et al., 2009, Lou et al. 2022). Similarly, we observed an increase in the fluorescence intensity of the F-actin marker most likely corresponding to the expression of

muscle-specific actin isoforms (Fig. 1C and E). Cumulatively, these experiments demonstrate that the transition of myoblasts from the proliferative to the terminally differentiated state is accompanied by dynamic changes in motility and actin intensity.

**Machine learning applied to time series data generates quantitative single-cell differentiation trajectories**

Following the association between differentiation and the population scale changes in actin intensity and motility, we hypothesized that the information encoded in single-cell migration trajectories and actin dynamics might be sufficient to computationally estimate a continuous score reflecting a myoblast's gradual transition from an undifferentiated proliferative state to a terminally differentiated fusion competent state. To test this hypothesis, we took a machine learning approach: (1) extracting features from the motility/actin time-series, (2) training machine learning classification models (aka *classifiers*) to discriminate between the undifferentiated and differentiated states, and (3) using the confidence of these models as a quantitative measurement for cell state.

The first step in designing our machine-learning model was determining which cells and timeframe can be considered as undifferentiated or differentiated. Myocytes must differentiate to become fusion competent (Abmayr & Pavlath, 2012). Hence, we defined the cultures as differentiated for classification 2.5 hours before the first fusion event was observed in the field of view. Undifferentiated cells were defined from the population grown in proliferation medium in the presence of DMSO, which continue to proliferate and remain undifferentiated, except a small fraction that begins to differentiate stochastically towards the end of the experiment due to the increase in cell density (Eigler et al., 2021). To enable continuous scoring along single-cell differentiation trajectories, we performed semi-manual single-cell tracking, where each trajectory was manually verified and corrected when necessary (Video S2). Single cell analysis confirmed our population-based results that differentiation was accompanied by a decrease in cell motility and an increase in the F-actin marker's fluorescence intensity (Fig. S3). We partitioned trajectories of undifferentiated and differentiated myoblasts to overlapping temporal segments of 2.5 hours each, for an overall 16,636 undifferentiated and 47,819 differentiated temporal segments, extracted from 310 and 538 cells correspondingly, that were used for model training (Fig. 2A - top). From each temporal segment, we extracted the corresponding single cell motility (dx/dt, dy/dt) and actin

intensity time series. Single-cell motility/actin time series features were extracted using the Python

package "Time Series FeatuRe Extraction on basis of Scalable Hypothesis tests" (tsfresh) that

derives properties such as temporal peaks, descriptive statistics (e.g., mean, standard deviation)

and autocorrelations (Christ et al., 2018). The extracted single-cell feature vectors and their

corresponding undifferentiated/differentiated labels were used to train random forest classifiers

(Breiman, 2001), which surpassed other machine learning algorithms (Fig. S4). The entire process

is depicted in Fig. 2A and detailed in the Methods.

We applied the trained motility and actin classifiers on single-cell trajectories from an experiment

that was not used for training and attained a continuous quantification following the differentiation

process by using overlapping temporal segments. At the population level, the single cell state

classification performance gradually increased from an area under the receiver operating

characteristic (ROC) curve (AUC) of ~0.6 to ~0.85 at 7.5-14.5 hours from experimental onset (Fig.

2B-C). A classifier trained on features derived from both motility and actin time-series surpassed

each of the motility/actin classifiers, suggesting that motility and actin dynamics contain

complementary information regarding the cells' state (Fig. S5A-C). The AUC values of all models

were well beyond the random value of 0.5, indicating that our classifiers can discriminate between

undifferentiated and differentiated cells at the population level before appreciable cell

morphological changes occur.

Next, we wondered whether we can use these classifiers to predict the differentiation state of a

single cell. For a given temporal segment of a given cell, the classifier outputs a "confidence score"

(i.e., differentiation score) that reflects the model's certainty in its prediction. Low differentiation

scores indicate that the cells are predicted as undifferentiated, while high scores indicate predicted

differentiation. To interpret what temporal features were the most important for the models'

prediction, we applied SHapley Additive exPlanations (SHAP) (Lundberg & Lee, 2017) and used

random forest's feature importance algorithms (Breiman, 2001). Both interpretability methods

highlighted temporal features related to high variance of acceleration rate or high complexity of

actin intensity time series as dominant features driving the models' prediction (Fig. S6). We

hypothesized that the differentiation score could be used as a continuous readout for the cell state.

At the critical time frame of 7.5-14.5 hours, at the population level, the differentiation scores of

ERKi-treated cells gradually increased for the motility (Fig. 2D), the actin-based (Fig. 2E), and

the combined (Fig. S5D-E) models while maintaining low scores for experiments of DMSO-treated cells. For the rest of the manuscript we focused on analyzing the motility- and actin-based models, because showing that each of two independent models trained with different temporal readouts (motility, actin) can quantitatively monitor the cell differentiation process, and thus strengthening our goal of evaluating whether the confidence score of a classifier trained for a binary classification task can be used to continuously measure a biological process that evolves over time.

We conducted a single-cell analysis by measuring the Spearman correlation between single-cell differentiation score and time at the critical time interval of 7.5-14.5 hours when differentiation occurs. This analysis indicated that most cells underwent a monotonic increase in differentiation scores over time (Fig. 2F). A similar gradual increase in differentiation score at 7.5-14.5 was observed when flipping the experiments used for training and testing (Fig. S7), the differentiation score was not sensitive to the size of the temporal segment (Fig. S8), nor to the window size used to measure actin (Fig. S9), and was consistent across multiple independent trainings (Fig. S10). Visualizing single-cell trajectories showed that most trajectories followed a gradual increase in their differentiation scores (Fig. 2G). Measuring the distribution of the per-cell differentiation scores' temporal derivative (Fig. S11A-B), their integration over time (Fig. S11C-F), the predicted onset (Fig. S12), and the predicted duration (Fig. 2H) of the differentiation process, suggested that the progression in single-cell differentiation is highly heterogeneous (Fig. 2H). These results suggest a heterogeneous gradual transition from an undifferentiated to a differentiated state within a typical timeframe.

The motility- and actin-based classifiers' predictions were mostly consistent at their single cell predictions, showing high correlation beyond 0.5 for 62% of cells, and negative correlation for less than 4%, and providing further support that both models measure the continuous state transition (Fig. S13A-B). Analysis of cells sub-groups partitioned according to the agreement between the motility and actin models, showed that lower agreement between the classifiers was associated with lower monotonicity of the differentiation scores of both models (Fig. S13C). In most cases of disagreement between the motility and actin models, we noticed a deviation in the actin-based model when the cell entered a crowded region and/or crawled below other cells (Fig. S14). High agreement between the two models was associated with the single cells' motility

persistence, the ratio between the direct translation (i.e., distance from start to end) and the overall

distance traveled (Fig. S13C). Qualitative and quantitative association between single cells'

motility persistence and motility-based (but not actin-based) differentiation score identified

persistent motility as a functional marker for the intermediate states of myoblast differentiation

(Fig. S15). The lack of association between the actin model and persistence suggests that the actin

model encodes different dynamic properties that are linked to the differentiation. Moreover, this

lack of association highlights the potential to use deviations between these models to discover

mechanisms that uncouple the link between motility, actin dynamics and myoblast differentiation.

Altogether, our data suggest that machine learning can transform motility and actin dynamics to a

quantitative readout characterizing the myoblast differentiation process at single-cell resolution

describing a continuous myoblast state transition from an undifferentiated to the terminally

differentiated states at 7.5-14.5 hours post induction.

**Models that discriminate between undifferentiated and differentiated states are not**

**sufficient for the quantitative characterization of the continuous differentiation processes**

Using the simplest readouts to quantify and delineate different biological conditions/states is

always preferred because it provides more direct insight regarding the underlying mechanisms. Is

it possible that our approach is overly complicated and exceeds what is required to quantitatively

describe the differentiation process? Are straight-forward single-cell measurements sufficient to

discriminate between undifferentiated and differentiated cells and follow the differentiation

process? To test this possibility, we evaluated the discriminative performance of single-cell

properties that are expected to deviate between the undifferentiated and differentiated cells. These

included cell speed, actin intensity, temporal derivative of actin intensity, migration persistence,

and local density. The local density dramatically increased over time for cells grown in

proliferation medium due to continued proliferation throughout the experiment (Fig. 3A). The

mean speed and actin intensity in proliferating cultures slightly decreased and increased

correspondingly over time, perhaps due to the increased density (Fig. 3B-C), and the mean

temporal derivative of actin intensity fluctuated around zero for both differentiating and

proliferating cells (Fig. 3D). Persistent migration of proliferating cells was lower compared to

differentiating cells without a clear trend over time (Fig. 3E). Each of these four discriminative

readouts, as well as their integration, could be generalized across experiments as demonstrated by

using each feature to train a machine learning model and applying this model to discriminate

between the two experimental conditions in an independent experiment (Fig. 3F).

The model trained with the local density and the model trained with all four features surpassed the

discrimination performance of the time-series motility and actin models (also reported in Fig. 2B-

C). However, discrimination does not necessarily imply that these readouts can be used to

quantitatively describe the differentiation process. Indeed, the differentiation score of each of these

classifiers could not capture the differentiation process as measured by single-cell monotonic

increase at the critical differentiation time interval of 7.5-14.5 hours. The single cell correlations

between the differentiation score and time were low for all the single-feature classifiers, as well as

for the integrated classifier (Fig. 3G). This is in contrast to our classifiers that generalized to

effectively quantify the differentiation process leading to a higher correlation between the

differentiation score and time (Fig. 3G - motility, actin intensity, same data as in Fig. 2G). A

plausible explanation for why these effective discriminating models could not capture the

continuous differentiation process is that the discriminating features captured properties attributed

to the undifferentiated state. For example, the increased local cell density of proliferating cells can

be used to effectively discriminate between the two populations but does not provide any

information regarding the progression through differentiation. Indeed, training models that

included temporal features extracted from single-cell local density dynamics showed the same or

deteriorated correlation between the differentiation score and time compared to models that were

not trained with local density (Fig. S16). Motility persistence, which was the most monotonic

among the single feature, and was earlier identified to associate with differentiation (Fig. S15),

was still far behind the integrated models in terms of monotonicity, suggesting that there is further

discriminative information beyond persistence in the cells' trajectories. These results indicate that

effective discrimination between the discrete extreme states is insufficient for the quantitative

characterization of continuous state transitions. Specifically, using machine learning for

quantitative characterization requires extracting features that can capture the state transition and

avoiding features that may confound the quantitative characterization of the process (e.g., avoiding

local cell density in characterizing the differentiation process). In our case, and in agreement with

other studies (Copperman et al., 2021; Wang et al., 2020; Wu et al., 2022), integration of multiple

dynamic features encoding the temporal changes were necessary to continuously measure a
biological process.

**The transition from terminal differentiation to fusion is controlled by p38**

We next aimed at harnessing our single myoblasts continuous differentiation scores to investigate
the relationship between cell differentiation and fusion. We manually annotated the fusion time of
68 myoblasts that fused to 6 myofibers (Fig. 4A) and used the continuous differentiation score to
determine an estimated time of the terminal differentiation state. Both the distributions of the single
cells' terminal differentiation and fusion times followed a normal-like distribution, where the
variability in the predicted differentiation time was higher than that of fusion time (Fig. 4B). The
time duration between terminal differentiation and fusion also followed a normal-like distribution,
indicating a typical duration of ~3 hours between terminal differentiation and fusion at the
population scale (Fig. 4C). These results suggest that cells undergo fusion within a typical time
interval from their terminal differentiation. This coupling was validated by measuring a correlation
between single-cell differentiation and fusion times (Fig. 4D) and was not sensitive to the threshold
used to determine the terminal differentiation time (Fig. S17). These results suggest that myoblasts
must reach a differentiation checkpoint before fusion can proceed.

Previous studies have shown that co-inhibition of p38, a family of MAP kinases that play a critical
role in the initiation of the differentiation program, together with a promyogenic factor, overcomes
the early block in differentiation but not the later impairment of muscle cell fusion imposed by the
p38 inhibitor, leading to differentiated unfused cells (Gardner et al., 2015). Following this logic,
we treated primary myoblasts with the promyogenic ERKi and the p38 inhibitor BIRB 796 (p38i;
5 µM) and performed primary myoblasts live imaging experiments. There was little appreciable
difference between cells treated with p38i and controls treated with DMSO, consistent with
previous studies showing that p38i maintains myoblasts in a proliferative undifferentiated state
(Zetser et al., 1999). Myoblasts co-treated with ERKi and p38i appeared differentiated but failed
to fuse, leading to the complete absence of multinucleated myofibers, reinforcing the notion that
p38 is essential for a transition from differentiation to fusion and maturation (Fig. 5A-B).
Immunofluorescence staining validated that the fraction of MyoG-positive cells remained low for
p38i treated cells and increased in cultures cotreated with p38i and ERKi, indicating that co-

inhibition of p38 and ERK1/2 leads to *bona fide* differentiation (Fig. 5C, Fig. S2, Video S3).
Moreover, cells started fusing once the p38 inhibitor was washed out indicating that they were
stalled at a "ready to fuse" state (Fig. S18).

However, it was not clear whether the differentiation process was altered with respect to ERKi-
treated cells. Thus, we quantitatively described the differentiation process of cells co-treated with
p38i and ERKi by applying our motility and actin models trained with proliferating and
differentiating cells. As a control, we validated that the differentiation score profile of p38i-treated
cells resembled that of proliferating cells treated with DMSO alone (Fig. 5D-E, Fig. S19). The
motility classifier showed that the differentiation profile of p38i-ERKi-treated cells followed a
trend strikingly similar to the one obtained for ERKi-treated cells and specifically included the
gradual transition at the critical time of 7.5-14.5 hours (Fig. 5D). The mean actin classifier score
maxed at a value of ~0.5, which was not sufficient to distinguish undifferentiated from
differentiated cells, suggesting that p38i alters actin dynamics (Fig. 5E). Nevertheless, the actin
model predicted a monotonically increasing trend, suggesting that the cells were becoming more
differentiated over time.

To test the model's prediction that co-inhibition of p38 and ERK1/2 leads to properly differentiated
cells that are ready to fuse and that actin dynamics are altered under these conditions, we acquired
mass spectrometry data. We extracted proteins from primary myoblasts treated with ERKi,
ERKi+p38i, p38i, and DMSO control for 24 hours and used mass spectrometry-based proteomics
for unbiased, high-throughput identification of differentially expressed proteins. In total 4319
proteins were identified across all experiments. Unsupervised hierarchical clustering analysis
showed that p38i treated cultures cluster with the DMSO proliferation control, consistent with
previous studies showing that p38 inhibition maintains myoblasts in the proliferative
undifferentiated state (Fig. 6A). The analysis also showed that ERK1/2 and p38 co-inhibited
cultures cluster with the differentiating ERKi treated cultures, suggesting that differentiation
occurs under co-inhibition conditions (Fig. 6A).

The actin cytoskeleton is essential for fusion and for the subsequent formation of sarcomeres
(Abmayr & Pavlath, 2012; Deng et al., 2015; Gruenbaum-Cohen et al., 2012; Onel & Renkawitz-
Pohl, 2009; Richardson et al., 2008). Gene Ontology (GO) annotation enrichment analysis showed
a significant up regulation of proteins involved in the process of actin cytoskeleton organization

and down regulation in the process of skeletal muscle contraction and maturation in the ERKi-
p38i co-inhibition cultures compared to ERKi alone, consistent with an arrest prior to fusion and
maturation (Fig. 6B). These results supported our conclusion that co-inhibited cells arrest at
terminal differentiation before fusion and maturation and explain the deviation in the
differentiation score of the actin, but not the motility classifier, upon ERKi-p38i co-inhibition.

Strikingly, only 18 out of 504 proteins that were significantly differentially expressed between the
differentiated (ERKi) and the differentiated unfused cells (ERKi-p38i) were completely missing
from the differentiated unfused cells (Fig. 6C). One of these being Myomixer, which is known to
be essential for fusion (Zhang et al., 2017, Quinn et al., 2017, Bi et al., 2017). These results suggest
that cell differentiation and fusion can be uncoupled and that p38 is essential for the transition
from terminal differentiation to fusion and maturation supporting the notion that differentiation
and fusion are coordinated processes that occur in tandem.

## Discussion

We combined live cell imaging and machine learning to infer the differentiation state of single cells during the process of muscle precursor cell differentiation. Many studies highlight the rich information encapsulated in single-cell dynamics that, with the aid of supervised or unsupervised machine learning, enable effective identification of sub-populations and discrimination of perturbations (Choi et al., 2021; Goglia et al., 2020; Jacques et al., 2021; Jena et al., 2022; Kimmel et al., 2018; Valls & Esposito, 2022), that cannot be inferred from static snapshot images (Copperman et al., 2021; Wang et al., 2020; Wu et al., 2022). For example, approaches that rely on static snapshots make it extremely hard to infer trajectories that deviate from the mainstream cell state progression because they are confounded by cell-to-cell variability. The ability to measure a single cell state as it transitions through time during a physiological process, along with careful experimental-computational interplay, enabled us to quantitatively follow the process and derive biological insight. Specifically, identify the key time frame where myoblasts gradually undergo differentiation (Fig. 2D-E), link single-cell differentiation to fusion (Fig. 4D), associate persistent migration with differentiation (Fig. S15), validate that co-inhibition of p38 and ERK1/2 arrests the fusion process without altering the differentiation process, and confirm our model's prediction that actin regulators are altered (Fig. 5-6). Ultimately, our experiments show that p38 regulates the transition from differentiation to fusion, implying that there is a differentiation checkpoint that must be reached before fusion occurs, and opening new avenues to identify novel regulators of this process.

The ability to infer the differentiation state of individual myoblasts can further enable the identification of novel myogenic factors, high throughput screening for pro-regenerative compounds, and the definition and subsequent examination of distinct intermediate steps in the differentiation process. Moreover, this approach of harnessing temporal dynamics by machine learning, without explicit state markers, can be generalized beyond terminal differentiation. Such a computational estimation of the cell state may have wide applications in characterizing other single-cell dynamic functions such as transitioning during the cell cycle, epithelial to mesenchymal transition, immotile to motile, disease progression, and cell death. The dynamic state readout can be correlated to other, independently measured cell readouts to systematically characterize the full spectrum of heterogeneities in complex biological processes.

Unsupervised approaches for cell state inference traverse from an initial to a final state through
steps that rely on similarity in cell appearance (Gut et al., 2015). These trajectories can be distorted
by batch effects or cell phenotypes unrelated to the state transition. In our approach, the supervised
component forces the trajectory to follow the phenotypic axes most relevant to the state transition
under investigation. This approach is similar to the approaches taken by (Szkalisity et al., 2021),
which rely on the manual assignment of cells to discrete states in 2D that are then inferred by
regression analysis, or by (Stallaert et al., 2022) that uses a supervised model to select features
predictive of the cell state before constructing cell state trajectories.

Our approach uses the physiological cell state (undifferentiated vs. differentiated) as the ground
truth, optimizes binary classification, and uses the classification's confidence score as the cell state
measurement. However, there is no guarantee that the classification's confidence score has linear
properties. For example, whether the difference in scores between 0.3 and 0.4 has the same
phenotypic magnitude as between scores of 0.6 and 0.7. This limitation is also common to
approaches that use non-linear dimensionality reduction (Copperman et al., 2021; Eulenberg et al.,
2017; Jacques et al., 2021; Rappez et al., 2020; Stallaert et al., 2022; Wang et al., 2022; Wu et al.,
2022) and could also limit unsupervised state representations that can be dominated by features
that do not relate to the cell state (Copperman et al., 2021; Jacques et al., 2021; Wang et al., 2022;
Wu et al., 2022). Still, the monotonicity property holds, e.g., a differentiation score of 0.4 is
predicted to be more advanced along the differentiation trajectory than a differentiation score of
0.3. This implies that the machine learning model captures more phenotypic evidence for the
advancement along the state transition axis. This monotonicity property is sufficient for comparing
different trajectories and calculating temporal correlations between cell state and other properties,
as demonstrated here, with motility persistence, and elsewhere (e.g., (Mayr et al., 2021; Zaritsky
et al., 2021)).

## Methods

### Mouse lines

We used Actin and nuclear reporter mice (LifeAct-GFP/ nTnG$^{+/+}$) (Eigler et al., 2021). Fluorescence expression validated using visual inspection. All experiments were approved by the Animal Care and Use Committee of the Weizmann Institute of Science (IACUC application #07690920-3).

### Isolation and treatment of primary myoblasts

Primary mouse myoblasts were isolated from gastrocnemius muscle using mechanical tissue dissociation as in (Eigler et al., 2021). Briefly, after cutting the muscle tissue into small pieces, they were incubated in Trypsin EDTA Solution B (0.25%Tripsin and 0.5% EDTA, Biological Industries Israel) and subjected to mechanical dissociation with a serological pipet. Supernatants were strained (FALCON REF no.352340) and centrifuged. Cell pellets were resuspended in BioAMF$^{TM}$-2 media (Biological Industries, Israel), plated on 10% Matrigel® Matrix (Corning REF no. 354234) coated plates, and grown at 37° in a 5% $CO_2$ incubator. Bio-AMF$^{TM}$-2 was used in all experiments (Biological Industries Israel).

### Microscopy

For live imaging, 40.000 cells were plated in a Slide 8-well chamber (Ibidi GmbH, cat.no.80826) coated with 10% Matrigel® Matrix. 15hr after cell seeding, the different treatments were added to the cells cultured in proliferation medium Bio-AMF$^{TM}$-2 (Biological Industries Israel). To induce myoblasts differentiation, cells were treated with 1µM ERK inhibitor (SCH 772984 Cayman Chemical Company). The inhibitors are dissolve in Dimethyl Sulfoxide (DMSO, MP Biomedicals cat.no 196055, 1.10g/ml stock concentration). Therefore, in the control sample of proliferation, DMSO treatment was added in a concentration of 1 µg/ml (equal to 1 µl, the volume added of each inhibitor). In the samples treated with p38 inhibitor, were used 5 µM (BIRB 796, AXON 1358) either alone or together with ERKi.

Live imaging (37°C, with 5% CO2) was performed using Cell discoverer 7-Zaiss inverted in widefield mode with Zeiss Axiocam 506 camera Carl Zeiss Ltd. Images were acquired using a ZEISS Plan-APOCHROMAT 20x / 0.70 Autocorr Objective (Working distance 2.20 mm). Excitation 470nm for GFP signals (LifeAct) and 567nm for tdTomato (nuclei). ZEN blue

software 3.1 was used for image acquisition. If necessary, linear adjustments to brightness and

contrast were applied using ImageJ v1.52 software (Schindelin et al., 2012). Cells were imaged

1.5 hours after adding the treatments, with 5 min intervals and at a pixel size of 0.462 µm.

<u>Fixed samples</u> were imaged using a ZEISS Plan-APOCHROMAT 5x / 0.35 Autocorr Objective

(Working distance 5.10 mm), 1.178 µm/px. Excitation 470nm for GFP (MyHC) and 567nm for

Alexa Fluor® 568 (MyoG) and 405nm for nuclei stained with Hoechst 33342.

**Immunofluorescence staining of MyoG-MyHC**

Primary myoblasts were seeded in a 96-well culture dish, coated in Matrigel® Matrix at 8,000

cells per well cultured in BioAMF-2 media. After 15h incubation at 37°C in a 5% $CO_2$ incubator,

the cells were treated with 1µM ERK inhibitor (SCH 772984 Cayman Chemical Company) and

5 µM p38 inhibitor (BIRB 796, AXON 1358) in the needed samples. The inhibitors are dissolve

in Dimethyl Sulfoxide (DMSO, MP Biomedicals cat.no 196055, 1.10g/ml stock concentration).

Therefore, in the control sample of proliferation, DMSO treatment was added in a concentration

of 1 µg/ml (equal to 1 µl, the volume added of each inhibitor). Cells were fixed at specific time

points (0h-6h-8h-10h-12h-16h-24h) with 3.7% PFA in PBS for 15 minutes at room temperature.

The cells were then quenched with 40mM ammonium chloride for 5 min, washed with PBS 3

times, permeabilized in PBS with 0.01% Triton x-100 for 10 min, and blocked in 10% FBS in

PBS (blocking buffer) for 1h at room temperature. Primary antibody incubation was done in a

blocking buffer overnight at 4 degrees, with the following antibodies: Anti-Fast Myosin Heavy

Chain antibody [MY-32] (Ab51263), Abcam) 1:400, Anti-Myogenin antibody [EPR4789]

(ab124800) 1:500. Cells were washed 3 times in PBS and then incubated with secondary

antibodies: Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) (ab150117) 1:600, Donkey Anti-

Rabbit IgG H&L (Alexa Fluor® 647) (ab150067) 1:600, Donkey Anti-Rabbit IgG H&L (Alexa

Fluor® 568) (ab175692) 1:600. The cells were washed 3 times in PBS, incubated with Hoechst

33342 (Thermo Scientific cat no.62249, 1:1000) for 5 min and washed in PBS.

<u>Quantification</u>: The percentage of expressing cells was calculated by dividing the number of

nuclei labeled by the MyoG antibody by the total amount of cells given by the Hoechst staining

in 3 independent replicates of each experimental condition. The nuclei were segmented and

counted using the Cellpose software (Stringer et al., 2021).

Quantification of fusion index: First, the nuclei were segmented and counted using the Cellpose

software (Stringer et al., 2021) together with a homemade python script to gain the total number

of nuclei. Then, the fusion index was quantified by manually identifying the number of nuclei

found in cells with at least two nuclei. The values were expressed as a percentage of the total

number of nuclei per field of view.

**Actin intensity quantification in a field of view**

The quantification was made using the ImageJ v1.52 software (Schindelin et al., 2012). We

measured the fluorescence intensity signal of the entire field of view every hour and plotted the

mean intensity with stdDev calculated over all the pixel values of every field of view.

**Protein sample preparation for mass spectrometry**

Primary myoblasts were grown on 10cm plates coated in Matrigel® Matrix at $1.2 \times 10^6$ cells per

plate, cultured in BioAMF-2 media. After 15 hours the cells were treated with: (1) 1µM ERK

inhibitor (SCH 772984 Cayman Chemical Company) (2) 5 µM p38 inhibitor (BIRB 796, AXON

1358) (3) 1µM ERK inhibitor with 5µM p38 inhibitor and (4) 0.1% DMSO in the control samples.

24 hours after treatment cells were washed with PBS, harvested by scraping and pelleted by

centrifugation at 2,000 g for 5 min. We lysed each pellet in 1% SDS lysis buffer: 1% SDS in 50

mM Tris (pH 8) containing protease inhibitor cocktail (Sigma Aldrich, catalog no. P8849) and 1

mM PMSF. The samples were boiled at 95°C for 5 min, then diluted into 2X RIPA lysis buffer to

give 1x RIPA lysis buffer (50 mM Tris-HCl [pH 8.0], 150 mM NaCl, 0.5% SDS, 0.5% sodium

deoxycholate, 1% Triton X-100). The lysates were cleared by centrifugation at 16,000g for 10 min

at 4°C. protein concentration analysis was done using BCA kit (Pierce). A total of 12 individual

cell pellets were prepared for LC MS\MS analysis - four treatments with three biological repeats.

Protein samples were frozen in liquid nitrogen and stored at-80°C.

**Proteolysis and Mass Spectrometry Analysis**

The proteins were dissolved in RIPA buffer, sonicated and precipitated in 80% acetone. The

protein pellets were dissolved in in 9M Urea and 400mM ammonium bicarbonate than reduced

with 3mM DTT (60ºC for 30 min), modified with 10mM iodoacetamide in 100mM ammonium

bicarbonate (room temperature 30 min in the dark) and digested in 2M Urea, 25mM ammonium

bicarbonate with modified trypsin (Promega), overnight at 37°C in a 1:50 (M/M) enzyme-to-

substrate ratio. The tryptic peptides were desalted using C18 (Top tip, Glygen) tip, dried and re-

suspended in 0.1% Formic acid**.**

The peptides were resolved by reverse-phase chromatography on 0.075 X 300-mm fused silica

capillaries (J&W) packed with Reprosil reversed phase material (Dr Maisch GmbH, Germany).

The peptides were eluted with linear 180 minutes gradient of 5 to 28% acetonitrile with 0.1%

formic acid in water, 15 minutes gradient of 28 to 95% and 25 minutes at 95% acetonitrile with

0.1% formic acid in water, at flow rates of 0.15 µl/min. Mass spectrometry was performed by Q

Exactive HFX mass spectrometer (Thermo) in a positive mode (m/z 300–1800, resolution

120,000 for MS1 and 15,000 for MS2) using repetitively full MS scan followed by high collision

induces dissociation (HCD, at 27 normalized collision energy) of the 30 most dominant ions (>1

charges) selected from the first MS scan. The AGC settings were $3\times10^6$ for the full MS and

$1\times10^5$ for the MS/MS scans. The intensity threshold for triggering MS/MS analysis was $1\times10^4$.

A dynamic exclusion list was enabled with exclusion duration of 20 s.

The mass spectrometry data was analyzed using the MaxQuant software 1.5.2.8 (1) for peak

picking and identification using the Andromeda search engine, searching against the mouse

proteome from the Uniprot database with mass tolerance of 6 ppm for the precursor masses and

20 ppm for the fragment ions. Oxidation on methionine and protein N-terminus acetylation were

accepted as variable modifications and carbamidomethyl on cysteine was accepted as static

modifications. Minimal peptide length was set to six amino acids and a maximum of two mis

cleavages was allowed. Peptide- and protein-level false discovery rates (FDRs) were filtered to

1% using the target-decoy strategy. The data was quantified by label free analysis using the same

software with "match between runs" option. Protein tables were filtered to eliminate the

identifications from the reverse database, and common contaminants. Statistical analysis of the

identification and quantization results was done using Perseus 1.6.10.43 software. Annotation

enrichment was done by the string tool (https://string-db.org/) and the David bioinformatics

package (https://david.ncifcrf.gov/).

**Automated single-cell tracking and quantification**

Automatic nuclei speed was performed using the commercial software Imaris (v9.7.2, Oxford

Instruments). We created a new "spots" layer on the nuclei label channel using the default

Favorite Creation Parameters to track the spots over time, classify the spots, and object-object

statistics. Next, we estimated the diameter of 8 µm and enabled background subtraction. These

analyses allowed us to collect a large number of single-nuclei trajectories. While trajectories

frequently fragment using this approach, they were sufficient to quantify the mean nuclei speed

over time.

**Semi-manual single-cell tracking**

Semi-manual single-cell tracking was performed to obtain accurate trajectories for training and

evaluating our machine-learning models. The time-lapse images were first converted to

XML/hdf5 format using the BigDataViewer (v.6.2.1) FIJI plugin (Pietzsch et al., 2015;

Schindelin et al., 2012). We then used the Mastodon FIJI plugin (Mastodon – a large-scale

tracking and track-editing framework for large, multi-view images), for single-cell tracking and

manual correction. We tracked cells that resided within the field of view throughout the entire

experiment and included cells that fused into multinucleated fibers and cells that did not fuse

within the experimental timeframe. To reduce the manual annotation load, tracks that contained

less tracking errors were prioritized for manual correction. Altogether, we collected 848 tracks

for training (538 ERKi-treated cells; 310 DMSO-treated cells), 789 tracks, from an independent

experiment, for testing (686 ERKi-treated cells; 103 DMSO-treated cells), and 410 tracks, from

the perturbation experiment (202 p38i-treated cells; 208 ERKi+p38i-treated cells).

**Preprocessing trajectories**

We used OpenCV's CalcOpticalFlowFarneback, based on Gunner Farneback's method

(Farnebäck, 2003), for image registration to correct erroneous offsets of the tracked cells'

trajectories. For each pair of frames, we calculated the average offset and used the corresponding

translation for registration.

**Models training**

The training pipeline implements the following steps.

1. Determining labels for training. We assigned ERKi-treated cells with the

"differentiated" label in a time segment of 2.5 hours (hours 12.3-14.8) before the first

fusion event was observed in the field of view. We decided not to label ERKi-treated

cells as "undifferentiated" at the onset of the experiment because we did not know how

early differentiation phenotypic signs appear. The increase in MyoG-positive cells during

the first 6 hours of the experiment supports this decision. We assigned time segments of

DMSO-treated cells with the "undifferentiated" label because their differentiation begins

after more than the 23 hours of the experiment.

2. Partitioning single-cell trajectories to temporal segments. We partitioned trajectories of

DMSO- and ERKi-treated cells to overlapping temporal segments (overlap lag = 5

minutes) in equal lengths of 2.5 hours each. Temporal segments' length was determined

to match the time frame where we consider ERKi-treated cells as "differentiated". This

step resulted in 16,636 DMSO-treated cells and 47,819 ERKi-treated cells temporal

segments. For training, we labeled as "differentiated" ERKi-treated cells in the temporal

segment of hours 12.3-14.8 and labeled as "undifferentiated" DMSO-treated cells in non-

overlapping temporal segments throughout the experiment. Overall, we extracted 468

undifferentiated and 268 differentiated temporal segments for training.

3. Extracting motility and actin features. We extracted single-cell motility and actin

intensity time series from each temporal segment:

● Motility: We calculated the displacement of a single cell for each time point t,

creating a two-dimensional vector: $displacement(t) = (x_t - x_{t-1}, y_t - y_{t-1})$

● Actin: We cropped a quantification window of size 32X32 µm around the center

of each nucleus at each time point and calculated the minimum, maximum, mean,

median, and standard deviation of the actin intensity within the window.

4. Extracting hundreds of single-cell time series features using the "tsfresh" python

package (Christ et al., 2018). These features encoded properties of the temporal

segments, such as temporal peaks, derivatives, and statistics. The tsfresh feature selection

was based on the Benjamini-Yekutieli multiple test procedure (Benjamini & Yekutieli,

2001) to identify the most relevant features for characterizing the time series.

5. Training classifiers to distinguish between differentiated and undifferentiated cells. We
trained random forest classifiers, which are considered effective with high dimensional
and relatively small datasets (Breiman, 2001), as validated empirically on our data (Fig.
S4). Hyperparameter tuning was performed using a grid search with a 5-fold cross-
validation (motility classifier: {''max_depth': 12, 'min_samples_leaf': 1, 'n_estimators':
100}, actin intensity classifier: {'max_depth': 20, 'min_samples_leaf': 1, 'n_estimators':
200}).

6. Evaluating the trained classifies' performance. We assessed the discrimination
performance of our motility/actin classifiers on an independent experiment that was not
used for training. We partitioned time series to overlapping temporal segments (102,929
ERKi-treated cells segments, 7,214 DMSO-treated cells segments), selected temporal
segments for evaluation as described above for 577 differentiated and 180
undifferentiated temporal segments, extracted motility and actin intensity time series,
performed feature extraction using "tsfresh", and evaluated the performance of the
corresponding trained models. The AUC of the motility and the actin intensity classifiers
were 0.8 and 0.81, correspondingly (Fig. 3F); accuracy was 0.76 and 0.81; precision was
0.97 and 0.93; recall was 0.71 and 0.81.

**Inference of single cells differentiation trajectories**

Each single cell trajectory was partitioned into overlapping temporal segments of 2.5 hours, with
an overlapping lag of 5 minutes (one frame). We calculated motility & actin intensity time series,
applied "tsfresh", selected features according to training, and applied the corresponding trained
models on these feature vectors to retrieve a differentiation score for each segment defining
single-cell differentiation trajectories.

**Correlation of differentiation score with time**

The correlation between the single-cell differentiation scores and time was computed through the
critical time interval where differentiation occurred (7.5-14.5). We used the Spearman
correlation coefficient as a measurement for the monotonic increase in differentiation along a
trajectory.

**Prediction of the onset of differentiation process**

The onset of the single cell differentiation process was determined as the last stable time point below a threshold in the differentiation scores. This time point was defined as the last time point of the longest sequence with differentiation scores that ranged between values 0.2-0.3.

**Quantification of single cell predicted duration of the differentiation process**

The *differentiation process duration* is a proxy for the time a single cell undergoes differentiation. The duration of the single-cell differentiation process was determined as the time passed from the predicted onset of differentiation to reaching a high stable threshold in the differentiation scores. The high stable threshold was defined as the first time point of the longest sequence with differentiation scores that ranged between 0.7-0.8. The differentiation process duration was calculated as the time passed between the low and high stable thresholds.

**Simple single-cell measurements and corresponding classifiers**

We calculated single-cell time series of the following measurements:

- Local density: the number of nuclei within a radius of 50 µm around the cell.
- Speed: $speed(t) = \sqrt{(x_t - x_{t-1})^2 + (y_t - y_{t-1})^2}$, where $x_t$, $y_t$ are the nuclei $(x, y)$ position at time $t$.
- Mean actin intensity: mean actin intensity in a quantification window of 32X32 µm around the nuclei.
- Persistence: The ratio between a single cell's displacement and its full path length. Persistence of 1 implies that the cell migrated in a straight line.

For each measurement, and for all four together, we trained random forest classifiers with the mean value in each temporal segment to discriminate between undifferentiated and differentiated cells. We evaluated the discrimination performance of each of the five classifiers as described above.

**Quantification of single-cell terminal differentiation time:**

The *terminal differentiation time* of a single cell is an estimation based on the first time of the longest sequence of differentiation scores that are higher than a threshold value of 0.78 (to avoid local peaks).

**Manual annotation of fusion events timing**

68 nuclei from 6 fibers were backtracked to the frame when they fused into the fiber syncytium (Fig. S20).

**Statistical analysis**

Pearson correlation (scipy.stats.pearsonr function) was used to assess the correlation between the terminal differentiation time and fusion since we assumed a linear correlation between them (Fig. 4D). Spearman correlation (using scipy.stats.pearsonr function) was used for correlating the monotonic increase in the differentiation trajectories with time. D'Agostino's K-squared test (using scipy.stats.normaltest) was used to determine the normality of distributions: duration of the differentiation process, terminal differentiation time, fusion time, and duration between differentiation and fusion.

**Software and data availability**

Source code and sample data are publicly available, https://github.com/amitshakarchy/muscle-formation-regeneration.

## Acknowledgments

This research was supported by the Israeli Council for Higher Education (CHE) via the Data Science Research Center, Ben-Gurion University of the Negev, Israel (to AZ), and by the Wellcome Leap Delta Tissue program (to AZ). This project also received funding from the European Research Council (ERC StG # 851080 to O.A.). O.A. also acknowledges funding from the David Barton Center for Research on the Chemistry of Life and the Ruth and Herman Albert Scholarship Program for New Scientists. O.A. is an incumbent of the Miriam Berman Presidential Development Chair. We thank Shahar Golan for critically reading the manuscript.

## Author Contribution

AZ and OA conceived the study. GZ and AH designed the experimental assay and performed all experiments. AS developed analytic tools, analyzed, and interpreted the data with the help of RM. AS, GZ, RM and KH curated and annotated data. AS, OA, and AZ drafted the manuscript. OA and AZ mentored the authors. All authors wrote and edited the manuscript and approved its content.

## Competing Financial Interests

OA is a co-founder of ProFuse technologies LTD.

 **Figures**

 **Figure 1**

**Figure 1. Myoblast differentiation ex vivo leads to dynamic changes in motility and actin intensity**
(**A**) Overlay images of primary myoblasts fixed at different time points after ERK inhibition and stained for MyoG (Red), MyHC (green), and the nuclei (Hoechst, Blue). Magnification 5x, Scale bar: 100 µm. (**B**) Percentage of MyoG positive cells in differentiating cultures (ERKi, Red) and proliferating controls (DMSO, Black) over time. (**C**) Overlay image of primary myoblasts expressing the nuclear marker tDTomatto-NLS (magenta) and the actin marker LifeAct-EGFP (cyan) 23h after ERKi treatment. Magnification 10x. Scale bars 100 µm. (see Video S1). (**D**) Mean (line) and standard deviation (shade) of single cell speed over time during differentiation (ERKi; ~3000 cells). (**D**) Mean (line) and standard deviation (shade) of actin intensity over time of an entire field of view of differentiating cells (ERKi; Methods).

676  **Figure 2**

**Figure 2. Inference of single cells differentiation trajectories by machine learning applied to actin/motility dynamics**

**(A)** Training Random Forest classifiers to predict single cells' differentiation state - cartoon. Left: single cell motility/actin time series are partitioned into temporal segments of 2.5 hours each. Positive labels were assigned to the ERKi-treated cells' segment (top, orange) starting 2.5 hours before the first fusion event (orange star on the dashed timeline). Negative labels were assigned to all segments of DMSO-treated cells (blue). Right: features extracted from the positive (orange) and negative (blue) time series (top) were used to train classification models. Two models, one based on motility and the other on actin intensity, were trained based on time series extracted from the single-cell trajectories. **(B-C)** Classification performance. Area under the receiver operating characteristic (ROC) curve (AUC) over time for classifiers trained with motility (B) and actin intensity (C) time-series. The AUC was calculated for 789 cells from an independent experiment. Classification performance of a random model (AUC = 0.5) is marked with a dashed horizontal line. **(D-E)** Mean (solid line) and standard deviation (shade) of the differentiation score over time of ERKi (orange) and DMSO (blue) treated cells using the motility (B) and the actin intensity (C) classifiers (ERK: 575 cells; DMSO: 103 cells). The analysis for the entire experiment is shown at (Fig. S21). **(F)** Distribution of single-cell Spearman correlation between the classifier's score and time, calculated for motility (orange) and actin (red) classifiers (N=575 cells). Median correlation coefficient values were 0.55 (motility) and 0.7 (actin). 66.1% (motility) and 73.4% (actin) of the cells showed significant correlation (p-value < 0.05). **(G)** Representative single cells differentiation trajectories inferred by the motility (top) and the actin (bottom) classifiers. Trajectories are colored according to the Spearman correlation between their corresponding differentiation score and time. **(H)** Distribution of the single cell predicted duration of the differentiation process, as measured by the motility (yellow) and actin intensity (red) classifiers' prediction: the time passed between a stable low threshold of 0.2-0.3 and a high stable threshold of 0.7-0.8 (full details in Methods). The median predicted duration of the differentiation process was 3.3 (motility) and 4.5 (actin intensity) hours.

703 **Figure 3**

**Figure 3. Simple single-cell measurements are insufficient for continuous cell state transition characterization**

(**A-E**) Mean (solid line) and standard deviation (shade) of single cell characteristics over time of ERKi- (orange) and DMSO- (blue) treated cells. N=575 cells. Single-cell properties included are local cell density (A), mean speed (B), mean actin intensity (C), mean actin intensity temporal derivative (D), and persistence in migration (E). (**F**) Single-cell state classification performance. Area under the receiver operating characteristic (ROC) curve (AUC) for classifiers trained using local density, speed, mean actin intensity, mean actin intensity derivative, persistent migration, and integration of these properties to discriminate ERKi- and DMSO-treated cells experimental conditions. The right-most bars show AUCs for classifiers trained with motility and actin dynamics. AUC scores were calculated for 757 temporal segments of differentiated/undifferentiated cells from an experiment that was not used for model training (full details in Methods). (**G**) The single-cell correlation distribution between the differentiation score and time for all the classifiers shown in panel E (median shown in white). Dashed horizontal line shows no correlation. The right-most distributions show correlations for classifiers trained with motility and actin dynamics.

 **Figure 4**

**Figure 4. Correlation between terminal differentiation and fusion time**

(A) Annotations of single-cell fusion into a representative myofiber over time. Cells marked in white are already fused, and cells marked in yellow are fusing into the fiber. Scale bar 100 µm. (B) Distribution of single cells' fusion times (green) and terminal differentiation times determined by motility (yellow) and actin intensity (red) classifiers. The dashed vertical gray rectangle highlights the differentiation time interval of 7.5-14.5 hours. All three distributions were normal-like as assessed by the D'Agostino's K-squared test not rejecting the null hypothesis that the terminal differentiation time was normally distributed (motility classifier: p-value = 0.36, actin classifier p-value = 0.64; fusion time p-value = 0.1). The "terminal differentiation" state was determined using a differentiation score threshold of 0.78 (the same threshold was also used in panels C-D). The models identified 56 (motility) and 52 (actin intensity) cells that reached a terminal differentiation state. 71% (motility) and 65% (actin intensity) of the identified cells reached a terminally differentiated state by 15 hours post-induction. The median time of terminal differentiation was 12.63 (motility) and 14.2 (actin intensity); the median fusion time was 16.8 hours. (C) Distribution of the duration between single cell terminal differentiation and fusion, for terminal differentiation determined by motility (yellow) and actin (red) classifiers. Both distributions were normal-like as assessed by the D'Agostino's K-squared test not rejecting the null hypothesis that the duration was normally distributed (motility classifier: p-value = 0.13, actin classifier: p-value = 0.13). Median differentiation-to-fusion duration was 3.1 (motility) and 3 (intensity) hours. (D) Associating single cell terminal differentiation time (x-axis) and fusion time (y-axis), determined by the motility (yellow) and the actin (red) classifiers. Pearson correlation coefficients were 0.52 (motility) and 0.73 (actin intensity), p < 0.001 for both actin and motility classifiers.

743    **Figure 5**

**Figure 5. Differentiation and fusion can be uncoupled**

**(A)** Overlay images of primary myoblasts treated with p38i (left) and ERKi+p38i (left), fixed at 24hr and stained for MyoG (Red) and nuclei (Hoechst, Blue). Magnification 5x, Scale bar: 100 μm. Overlay images of myoblasts undergo differentiation but do not fuse. Scale bar 100 µm. **(B)** Fusion Index: Percentage of fused nuclei in ERKi and ERKi-p38i treated cultures. **(C)** Percentage of MyoG positive cells over time under proliferation conditions (p38i; Black) and differentiation conditions (ERKi + p38i; Red). DMDO and ERKi from Fig. 1 B are shown for convenience (semitransparent). **(D-E)** Mean differentiation score over time of ERKi- (orange), DMSO- (blue), ERKi+p38i- (green), and p38i- alone (purple) treated cells using the motility (C) and actin intensity (D) classifiers (ERKi: 575 cells; ERKi+p38i: 208 cells; p38i: 202 cells; DMSO: 103 cells). ERKi- and DMSO- treated cells differentiation scores are the same as in Fig. 2. The analysis for the entire experiment is shown at (Fig. S19).

**Figure 6**

**Figure 6. Mass spectroscopy analysis reveals differential protein expression between ERK1/2**
**inhibition and p38-ERK1/2 co-inhibition**

(**A**) Heatmap of significantly changed proteins (fold change > 2 and p-value < 0.05) between ERK1/2
inhibition samples and p38-ERK1/2 co-inhibition samples, after unsupervised hierarchical clustering.
Each column represents the sample and each row represents the differentially expressed protein. High
expression marked in red; low expression marked in green. Sample clustering tree is shown on the top.
(**B**) Bar Plot of Gene Ontology (GO) Annotation Enrichment Analysis compering ERK1/2 inhibition
samples and p38-ERK1/2 co-inhibition samples. Colors denote different protein groups. (**C**) Volcano plot
of differentially expressed proteins between ERK1/2 inhibition samples and p38-ERK1/2 co-inhibition
samples, red represents up-regulated proteins and blue represents down-regulated proteins.

## References

Abmayr, S. M., & Pavlath, G. K. (2012). Myoblast fusion: lessons from flies and mice.

*Development* , *139*(4), 641–656. https://doi.org/10.1242/dev.068353

Bains, W., Ponte, P., Blau, H., & Kedes, L. (1984). Cardiac actin is the major actin gene product

in skeletal muscle cell differentiation in vitro. *Molecular and Cellular Biology*, *4*(8),

1449–1453. https://doi.org/10.1128/mcb.4.8.1449-1453.1984

Benjamini, Y., & Yekutieli, D. (2001). The Control of the False Discovery Rate in Multiple

Testing under Dependency. *Annals of Statistics*, *29*(4), 1165–1188.

http://www.jstor.org/stable/2674075

Bentzinger, C. F., Wang, Y. X., & Rudnicki, M. A. (2012). Building muscle: molecular

regulation of myogenesis. *Cold Spring Harbor Perspectives in Biology*, *4*(2).

https://doi.org/10.1101/cshperspect.a008342

Bi, P., Ramirez-Martinez, A., Li, H., Cannavino, J., McAnally, J. R., Shelton, J. M., Sánchez-

Ortiz, E., Bassel-Duby, R., & Olson, E. N. (2017). Control of muscle formation by the

fusogenic micropeptide myomixer. *Science (New York, N.Y.), 356*(6335), 323–327.

https://doi.org/10.1126/science.aam9361

Bischoff, R. (1986). Proliferation of muscle satellite cells on intact myofibers in culture.

*Developmental Biology*, *115*(1), 129–139. https://doi.org/10.1016/0012-1606(86)90234-4

Breiman, L. (2001). Random Forests. *Machine Learning*, *45*(1), 5–32.

https://doi.org/10.1023/A:1010933404324

Buggenthin, F., Buettner, F., Hoppe, P. S., Endele, M., Kroiss, M., Strasser, M., Schwarzfischer,

M., Loeffler, D., Kokkaliaris, K. D., Hilsenbeck, O., Schroeder, T., Theis, F. J., & Marr,

C. (2017). Prospective identification of hematopoietic lineage choice by deep learning.

*Nature Methods*, *14*(4), 403–406. https://doi.org/10.1038/nmeth.4182

Choi, H. J., Wang, C., Pan, X., Jang, J., Cao, M., Brazzo, J. A., 3rd, Bae, Y., & Lee, K. (2021).

Emerging machine learning approaches to phenotyping cellular motility and

morphodynamics. *Physical Biology*, *18*(4). https://doi.org/10.1088/1478-3975/abffbe

Christ, M., Braun, N., Neuffer, J., & Kempa-Liehr, A. W. (2018). Time Series FeatuRe

Extraction on basis of Scalable Hypothesis tests (tsfresh – A Python package).

*Neurocomputing*, *307*, 72–77. https://doi.org/10.1016/j.neucom.2018.03.067

Copperman, J., Gross, S. M., Chang, Y. H., Heiser, L. M., & Zuckerman, D. M. (2021).

Morphodynamical cell-state description via live-cell imaging trajectory embedding. In

*bioRxiv* (p. 2021.10.07.463498). https://doi.org/10.1101/2021.10.07.463498

Deng, S., Bothe, I., & Baylies, M. K. (2015). The formin Diaphanous regulates myoblast fusion

through actin polymerization and Arp2/3 regulation. *PloS Genetics, 11*(8), e1005381.

https://doi.org/10.1371/journal.pgen.1005381

Eigler, T., Zarfati, G., Amzallag, E., Sinha, S., Segev, N., Zabary, Y., Zaritsky, A., Shakked, A.,

Umansky, K.-B., Schejter, E. D., Millay, D. P., Tzahor, E., & Avinoam, O. (2021).

ERK1/2 inhibition promotes robust myotube growth via CaMKII activation resulting in

myoblast-to-myotube fusion. *Developmental Cell*, *56*(24), 3349-3363.e6.

https://doi.org/10.1016/j.devcel.2021.11.022

Eulenberg, P., Köhler, N., Blasi, T., Filby, A., Carpenter, A. E., Rees, P., Theis, F. J., & Wolf, F.

A. (2017). Reconstructing cell cycle and disease progression using deep learning. *Nature*

*Communications*, *8*(1), 463. https://doi.org/10.1038/s41467-017-00623-3

Farnebäck, G. (2003). Two-Frame Motion Estimation Based on Polynomial Expansion. *Image*

*Analysis*, 363–370. https://doi.org/10.1007/3-540-45103-X_50

Fischer, U., Ludwig, N., Raslan, A., Meier, C., & Meese, E. (2016). Gene amplification during

myogenic differentiation. *Oncotarget*, *7*(6), 6864–6877.

https://doi.org/10.18632/oncotarget.6845

Gardner, S., Gross, S. M., David, L. L., Klimek, J. E., & Rotwein, P. (2015). Separating

myoblast differentiation from muscle cell fusion using IGF-I and the p38 MAP kinase

inhibitor SB202190. *American Journal of Physiology. Cell Physiology*, *309*(7), C491-

500. https://doi.org/10.1152/ajpcell.00184.2015

Goglia, A. G., Wilson, M. Z., Jena, S. G., Silbert, J., Basta, L. P., Devenport, D., & Toettcher, J.

E. (2020). A Live-Cell Screen for Altered Erk Dynamics Reveals Principles of

Proliferative Control. *Cell Systems*, *10*(3), 240-253.e6.

https://doi.org/10.1016/j.cels.2020.02.005

Griffin, C. A., Apponi, L. H., Long, K. K., & Pavlath, G. K. (2010). Chemokine expression and

control of muscle cell migration during myogenesis. *Journal of Cell Science, 123*(18),

3052–3060. https://doi.org/10.1242/jcs.066241

Gruenbaum-Cohen, Y., Harel, I., Umansky, K.-B., Tzahor, E., Snapper, S. B., Shilo, B.-Z., &

Schejter, E. D. (2012). The actin regulator N-WASp is required for muscle-cell fusion in

mice. *Proceedings of the National Academy of Sciences of the United States of America,*

*109*(28), 11211–11216. https://doi.org/10.1073/pnas.1116065109

Gut, G., Tadmor, M. D., Pe'er, D., Pelkmans, L., & Liberali, P. (2015). Trajectories of cell-cycle

progression from fixed cell populations. *Nature Methods*, *12*(10), 951–954.

https://doi.org/10.1038/nmeth.3545

Hernández-Hernández, J. M., García-González, E. G., Brun, C. E., & Rudnicki, M. A. (2017).

The myogenic regulatory factors, determinants of muscle development, cell identity and

regeneration. *Seminars in Cell & Developmental Biology*, *72*, 10–18.

https://doi.org/10.1016/j.semcdb.2017.11.010

Hurme, T., & Kalimo, H. (1992). Activation of myogenic precursor cells after muscle injury.

*Medicine and Science in Sports and Exercise*, *24*(2), 197–205.

https://www.ncbi.nlm.nih.gov/pubmed/1549008

Jacques, M.-A., Dobrzyński, M., Gagliardi, P. A., Sznitman, R., & Pertz, O. (2021). CODEX, a

neural network approach to explore signaling dynamics landscapes. *Molecular Systems*

*Biology*, *17*(4), e10026. https://doi.org/10.15252/msb.202010026

Jena, S. G., Goglia, A. G., & Engelhardt, B. E. (2022). Towards "end-to-end" analysis and

understanding of biological timecourse data. *Biochemical Journal*, *479*(11), 1257–1263.

https://doi.org/10.1042/BCJ20220053

Kimmel, J. C., Chang, A. Y., Brack, A. S., & Marshall, W. F. (2018). Inferring cell state by

quantitative motility analysis reveals a dynamic state system and broken detailed balance.

*PloS Computational Biology*, *14*(1), e1005927.

https://doi.org/10.1371/journal.pcbi.1005927

Krauss, R. S., Cole, F., Gaio, U., Takaesu, G., Zhang, W., & Kang, J.-S. (2005). Close

encounters: regulation of vertebrate skeletal myogenesis by cell-cell contact. *Journal of*

*Cell Science, 118*(11), 2355–2362. https://doi.org/10.1242/jcs.02397

Lepper, C., Partridge, T. A., & Fan, C.-M. (2011). An absolute requirement for Pax7-positive

satellite cells in acute injury-induced skeletal muscle regeneration. *Development* ,

*138*(17), 3639–3646. https://doi.org/10.1242/dev.067595

Luo, Z., Shi, J., Pandey, P., Ruan, Z.-R., Sevdali, M., Bu, Y., Lu, Y., Du, S., & Chen, E. H.

(2022). The cellular architecture and molecular determinants of the zebrafish fusogenic

synapse. *Developmental Cell, 57*(13), 1582-1597.e6.

https://doi.org/10.1016/j.devcel.2022.05.016

Lundberg, S., & Lee, S.-I. (2017). A unified approach to interpreting model predictions. In *arXiv*

*[cs.AI]*. arXiv.

https://proceedings.neurips.cc/paper/2017/hash/8a20a8621978632d76c43dfd28b67767-

Abstract.html

*mastodon: Mastodon – a large-scale tracking and track-editing framework for large, multi-view*

*images*. (n.d.). Github. Retrieved December 15, 2022, from https://github.com/mastodon-

sc/mastodon

Mayr, C. H., Simon, L. M., Leuschner, G., Ansari, M., Schniering, J., Geyer, P. E., Angelidis, I.,

Strunz, M., Singh, P., Kneidinger, N., Reichenberger, F., Silbernagel, E., Böhm, S.,

Adler, H., Lindner, M., Maurer, B., Hilgendorff, A., Prasse, A., Behr, J., … Schiller, H.

B. (2021). Integrative analysis of cell state changes in lung fibrosis with peripheral

protein biomarkers. *EMBO Molecular Medicine*, *13*(4), e12871.

https://doi.org/10.15252/emmm.202012871

Morris, E. J., Jha, S., Restaino, C. R., Dayananth, P., Zhu, H., Cooper, A., Carr, D., Deng, Y.,

Jin, W., Black, S., Long, B., Liu, J., Dinunzio, E., Windsor, W., Zhang, R., Zhao, S.,

Angagaw, M. H., Pinheiro, E. M., Desai, J., … Samatar, A. A. (2013). Discovery of a

novel ERK inhibitor with activity in models of acquired resistance to BRAF and MEK

inhibitors. *Cancer Discovery*, *3*(7), 742–750. https://doi.org/10.1158/2159-8290.CD-13-

0070

Nowak, S. J., Nahirney, P. C., Hadjantonakis, A.-K., & Baylies, M. K. (2009). Nap1-mediated

actin remodeling is essential for mammalian myoblast fusion. *Journal of Cell Science,*

*122*(Pt 18), 3282–3293. https://doi.org/10.1242/jcs.047597

Onel, S.-F., & Renkawitz-Pohl, R. (2009). FuRMAS: triggering myoblast fusion in Drosophila.

*Developmental Dynamics: An Official Publication of the American Association of*

*Anatomists, 238*(6), 1513–1525. https://doi.org/10.1002/dvdy.21961

Otey, C. A., Kalnoski, M. H., & Bulinski, J. C. (1988). Immunolocalization of muscle and

nonmuscle isoforms of actin in myogenic cells and adult skeletal muscle. *Cell Motility*

*and the Cytoskeleton, 9*(4), 337–348. https://doi.org/10.1002/cm.970090406

Pietzsch, T., Saalfeld, S., Preibisch, S., & Tomancak, P. (2015). BigDataViewer: visualization

and processing for large image data sets. *Nature Methods*, *12*(6), 481–483.

https://doi.org/10.1038/nmeth.3392

Powell, J. A. (1973). Development of normal and genetically dystrophic mouse muscle in tissue

culture. *Experimental Cell Research, 80*(2), 251–264. https://doi.org/10.1016/0014-

4827(73)90295-4

Prigge, J. R., Wiley, J. A., Talago, E. A., Young, E. M., Johns, L. L., Kundert, J. A., Sonsteng,

K. M., Halford, W. P., Capecchi, M. R., & Schmidt, E. E. (2013). Nuclear double-

fluorescent reporter for in vivo and ex vivo analyses of biological transitions in mouse

nuclei. *Mammalian Genome: Official Journal of the International Mammalian Genome*

*Society*. https://doi.org/10.1007/s00335-013-9469-8

Quinn, M. E., Goh, Q., Kurosaka, M., Gamage, D. G., Petrany, M. J., Prasad, V., & Millay, D. P.

(2017). Myomerger induces fusion of non-fusogenic cells and is required for skeletal

muscle development. *Nature Communications, 8*(1), 15665.

https://doi.org/10.1038/ncomms15665

Rappez, L., Rakhlin, A., Rigopoulos, A., Nikolenko, S., & Alexandrov, T. (2020). DeepCycle

reconstructs a cyclic cell cycle trajectory from unsegmented cell images using convolutional neural networks. *Molecular Systems Biology*, *16*(10), e9474. https://doi.org/10.15252/msb.20209474

Richardson, B. E., Nowak, S. J., & Baylies, M. K. (2008). Myoblast fusion in fly and vertebrates: new genes, new processes and new perspectives. *Traffic (Copenhagen, Denmark), 9*(7), 1050–1059. https://doi.org/10.1111/j.1600-0854.2008.00756.x

Riedl, J., Crevenna, A. H., Kessenbrock, K., Yu, J. H., Neukirchen, D., Bista, M., Bradke, F., Jenne, D., Holak, T. A., Werb, Z., Sixt, M., & Wedlich-Soldner, R. (2008). Lifeact: a versatile marker to visualize F-actin. *Nature Methods*, *5*(7), 605–607. https://doi.org/10.1038/nmeth.1220

Riedl, J., Flynn, K. C., Raducanu, A., Gärtner, F., Beck, G., Bösl, M., Bradke, F., Massberg, S., Aszodi, A., Sixt, M., & Wedlich-Söldner, R. (2010). Lifeact mice for studying F-actin dynamics. *Nature Methods, 7*(3), 168–169. https://doi.org/10.1038/nmeth0310-168

Rubinstein, N., Chi, J., & Holtzer, H. (1976). Coordinated synthesis and degradation of actin and myosin in a variety of myogenic and non-myogenic cells. *Experimental Cell Research, 97*(2), 387–393. https://doi.org/10.1016/0014-4827(76)90630-3

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schmidt, M., Schüler, S. C., Hüttner, S. S., von Eyss, B., & von Maltzahn, J. (2019). Adult stem cells at work: regenerating skeletal muscle. *Cellular and Molecular Life Sciences: CMLS*,

*76*(13), 2559–2570. https://doi.org/10.1007/s00018-019-03093-6

Schroeder, T. (2011). Long-term single-cell imaging of mammalian stem cells. *Nature Methods*,

*8*(4 Suppl), S30-5. https://doi.org/10.1038/nmeth.1577

Singh, K., & Dilworth, F. J. (2013). Differential modulation of cell cycle progression

distinguishes members of the myogenic regulatory factor family of transcription factors.

*The FEBS Journal*, *280*(17), 3991–4003. https://doi.org/10.1111/febs.12188

Stallaert, W., Kedziora, K. M., Taylor, C. D., Zikry, T. M., Ranek, J. S., Sobon, H. K., Taylor, S.

R., Young, C. L., Cook, J. G., & Purvis, J. E. (2022). The structure of the human cell

cycle. *Cell Systems*, *13*(1), 103. https://doi.org/10.1016/j.cels.2021.12.006

Stringer, C., Wang, T., Michaelos, M., & Pachitariu, M. (2021). Cellpose: a generalist algorithm

for cellular segmentation. *Nature Methods*, *18*(1), 100–106.

https://doi.org/10.1038/s41592-020-01018-x

Stumpf, P. S., Smith, R. C. G., Lenz, M., Schuppert, A., Müller, F.-J., Babtie, A., Chan, T. E.,

Stumpf, M. P. H., Please, C. P., Howison, S. D., Arai, F., & MacArthur, B. D. (2017).

Stem Cell Differentiation as a Non-Markov Stochastic Process. *Cell Systems*, *5*(3), 268-

282.e7. https://doi.org/10.1016/j.cels.2017.08.009

Szkalisity, A., Piccinini, F., Beleon, A., Balassa, T., Varga, I. G., Migh, E., Molnar, C.,

Paavolainen, L., Timonen, S., Banerjee, I., Ikonen, E., Yamauchi, Y., Ando, I., Peltonen,

J., Pietiäinen, V., Honti, V., & Horvath, P. (2021). Regression plane concept for

analysing continuous cellular processes with machine learning. *Nature Communications*,

*12*(1), 2532. https://doi.org/10.1038/s41467-021-22866-x

Valls, P. O., & Esposito, A. (2022). Signalling dynamics, cell decisions, and homeostatic control

in health and disease. *Current Opinion in Cell Biology*, *75*, 102066.

https://doi.org/10.1016/j.ceb.2022.01.011

Wang, W., Douglas, D., Zhang, J., Kumari, S., Enuameh, M. S., Dai, Y., Wallace, C. T.,

Watkins, S. C., Shu, W., & Xing, J. (2020). Live-cell imaging and analysis reveal cell

phenotypic transition dynamics inherently missing in snapshot data. *Science Advances*,

*6*(36). https://doi.org/10.1126/sciadv.aba9319

Wang, W., Poe, D., Yang, Y., Hyatt, T., & Xing, J. (2022). Epithelial-to-mesenchymal transition

proceeds through directional destabilization of multidimensional attractor. *Elife*, *11*.

https://doi.org/10.7554/eLife.74866

Wu, Z., Chhun, B. B., Popova, G., Guo, S.-M., Kim, C. N., Yeh, L.-H., Nowakowski, T., Zou, J.,

& Mehta, S. B. (2022). DynaMorph: self-supervised learning of morphodynamic states of

live cells. *Molecular Biology of the Cell, 33*(6), ar59. https://doi.org/10.1091/mbc.E21-

11-0561

Yang, K. D., Damodaran, K., Venkatachalapathy, S., Soylemezoglu, A. C., Shivashankar, G. V.,

& Uhler, C. (2020). Predicting cell lineages using autoencoders and optimal transport.

*PloS Computational Biology*, *16*(4), e1007828.

https://doi.org/10.1371/journal.pcbi.1007828

Yin, H., Price, F., & Rudnicki, M. A. (2013). Satellite cells and the muscle stem cell niche.

*Physiological Reviews*, *93*(1), 23–67. https://doi.org/10.1152/physrev.00043.2011

Zaritsky, A., Jamieson, A. R., Welf, E. S., Nevarez, A., Cillay, J., Eskiocak, U., Cantarel, B. L.,

& Danuser, G. (2021). Interpretable deep learning uncovers cellular properties in label-

free live cell images that are predictive of highly metastatic melanoma. *Cell Systems*,

*12*(7), 733-747.e6. https://doi.org/10.1016/j.cels.2021.05.003

Zetser, A., Gredinger, E., & Bengal, E. (1999). P38 mitogen-activated protein kinase pathway

promotes skeletal muscle differentiation. Participation of the Mef2c transcription factor.

*The Journal of Biological Chemistry*, *274*(8), 5193–5200.

https://doi.org/10.1074/jbc.274.8.5193

Zhang, Q., Vashisht, A. A., O'Rourke, J., Corbel, S. Y., Moran, R., Romero, A., Miraglia, L.,

Zhang, J., Durrant, E., Schmedt, C., Sampath, S. C., & Sampath, S. C. (2017). The

microprotein Minion controls cell fusion and muscle formation. *Nature Communications,*

*8*(1), 15664. https://doi.org/10.1038/ncomms15664

**Supplementary figures**

984
985 **Figure S1. The number of differentiating myoblasts MyoG-expressing increase over time**
986 Images of primary myoblasts fixed at different time points after ERK inhibition and stained for nuclei
987 (Hoechst, blue) and MyoG (red), along with Brightfield (gray) and Merge for reference. Magnification
988 5x, Scale bar 100 µm.
989

990
**Figure S2. Immunofluorescence staining of MyoG and MyHC**
Representative immunofluorescence (IF) images of myoblasts at 0,6,8,10,12,14,16, 24 hours after
treatment with DMSO, p38i 5 μM or ERKi 1 μM or the combination of ERKi-p38i. Cells were stained
using anti-MyoG (red), anti-MyHC (cyan), and the nuclear dye Hoechst 33342 (gray). Magnification 5x,
Scale bar: 100 μm.
996

997
998 **Figure S3. Dynamic changes in motility and actin intensity measured in semi-manually tracked**
999 **single cells**
1000 (**A-B**) Mean (line) and standard deviation (shade) of single cell speed (A) and mean actin intensity (B)
1001 over time during differentiation (ERKi; N = 575 cells). (**C-D**) Representative single cell's speed (C) and
1002 mean actin intensity (D) trajectories.
1003

1004

**Figure S4. Comparison of classification algorithms**
(**A-B**) Area under the receiver operating characteristic (ROC) curve (AUC) for classifiers trained with
actin intensity (A) and motility (B) time series, using random forest (RF), gradient boosting (GB), logistic
regression (LR), k-nearest neighbors (KNN), and support vector machines (SVM). Blue/green – flipped
train/test experiments. Average AUCs for actin intensity classifiers were 0.78 (RF), 0.77 (GB), 0.76 (LR),
0.59 (KNN), 0.54 (SVM). Average AUCs for motility classifiers were 0.8 (RF), 0.8 (GB), 0.64 (LR), 0.59
(KNN), 0.59 (SVM).
1012

1013

**Figure S5. Performance analysis of a classifier trained on both motility and actin dynamics**
(A) Area under the receiver operating characteristic (ROC) curve (AUC) over time for a combined model (N=678 cells). (B) AUC over time for a combined model- flipped experiments for train/test (848 cells). (C) Accuracy, precision, recall and area under the receiver operating characteristic (ROC) curve (AUC) for classifiers trained with motility (yellow), actin intensity (red) and a combination of motility and actin intensity (green) time series. Average accuracy rates were 0.74, 0.77 and 0.8; average precision rates were 0.77, 0.76, and 0.8; Average recall rates were 0.84, 0.84 and 0.87; average AUC rates were 0.8, 0.78 and 0.84 correspondingly. All metrics were calculated for 678 cells from an independent experiment. Overall, the combined classifier exhibits better classification performance. (D) Mean (solid line) and standard deviation (shade) of the differentiation score over time of ERKi- (orange) and DMSO- (blue) treated cells using the combined classifier. Dashed vertical gray rectangle highlights the time interval of 7.5-14.5 hours, where the model predicted the differentiation occurs (ERK: 575 cells; DMSO: 103 cells). (E) Mean (solid line) and standard deviation (shade) of the differentiation score over time of ERKi- (orange) and DMSO- (blue) treated cells using the combined classifier- flipped experiments for train/test (ERK: 538 cells; DMSO: 310 cells).

1030

**Figure S6. Random forests feature importance**

Top ten features most relevant for differentiation classification using random forest classifiers'
importance (A-B) and SHapley Additive exPlanation (SHAP) (C-D). Both approaches find very similar
important features: variance of consecutive change in displacement, permutation entropy of displacement,
variance of consecutive change in actin intensity, complexity-invariant distance of actin intensity and sum
of absolute consecutive changes in actin intensity. **(A-B)** Feature importance of random forest classifiers
trained to discriminate undifferentiated/differentiated cell states, on features extracted from motility
(yellow) and actin intensity (red), time series using "tsfresh" package. The 10 most important features are
shown. **(C-D)** SHAP summary plots of motility (C) and actin intensity (D) classifiers, produced by the
SHAP python package (Lundberg & Lee, 2017). The plot illustrates the feature relevance and combines
feature attributions to the model's predictive performance. Color is dependent on the feature values. The
10 most important features are shown.

1043
**Figure S7. Inference of differentiation trajectories - flipped experiments for train/test**
**(A-B)** Classification performance over time for the entire experiment. Area under the receiver operating
characteristic (ROC) curve (AUC) over time for classifiers trained with motility (B) and actin intensity
(C) time-series. The AUC was calculated for 736 cells from an independent experiment. Classification
performance of a random model (AUC = 0.5) is marked with a dashed horizontal line. Compare with Fig.

2B-C. **(C-D)** Mean (solid line) and standard deviation (shade) of the differentiation score over time of
ERKi- (orange) and DMSO- (blue) treated cells using the motility (B) and the actin intensity (C)
classifiers over time for the entire experiment. Dashed vertical gray rectangle highlights the time interval
of 7.5-14.5 hours, where both models predicted the differentiation occurs. Compare with Fig. S21.

1053

**Figure S8. Classification sensitivity analysis: temporal segment size**

**(A-B)** Mean (solid line) and standard deviation (shade) of the differentiation score over time of ERKi-
(orange) and DMSO- (blue) treated cells using the motility (B) and the actin intensity (C) classifiers,
trained with different sizes of temporal segment. Temporal segment's size (in hours) is shown above each
graph. Corresponding AUCs are reported in the figure. The temporal segment size for both classifiers was
2.5 hours.
1060

1061

1062  **Figure S9. Classification sensitivity analysis: actin intensity quantification window size**

1063 Top: an illustration of the actin quantification window size. Bottom: Mean (solid line) and standard
1064 deviation (shade) of the differentiation score over time of ERKi- (orange) and DMSO- (blue) treated cells
1065 using the actin intensity classifier, trained using these window sizes. Corresponding AUCs are reported in
1066 the figure.

1067

1068 **Figure S10. Prediction interval of single cells differentiation trajectories shows robustness across**
1069 **multiple model training.**

1070    Mean differentiation score (blue line) and differentiation trajectories (gray lines) predicted by 100
1071    independent models for both motility (left) and actin intensity (right) models. The figure shows the
1072    prediction interval of representative single cells differentiation trajectories.
1073

1074

**Figure S11. Differentiation scores' temporal derivative and integration over time**
**(A-B)** Mean (solid line) and standard deviation (shade) of temporal change in differentiation score for
motility (A) and actin intensity (B) models (ERK: 575 cells; DMSO: 103 cells). **(C-D)** Mean (solid line)
and standard deviation (shade) of the cumulative change in differentiation score for motility. Dashed

vertical gray rectangle highlights the time interval of 7.5-14.5 hours, where both models predicted the
differentiation occurs. (C) and actin intensity (D) models. Dashed vertical gray rectangle highlights the
time interval of 7.5-14.5 hours, where both models predicted the differentiation occurs (ERK: 575 cells;
DMSO: 103 cells). **(E-F)** Average cumulative change in differentiation score over time of Erki treated
cells, for motility **(E)** and actin intensity models **(F)**. P-values resulting from a t-test between the
cumulative change in differentiation score of cells treated with Erki and DMSO are demonstrating
significant (green, p-value <= 0.05) and non-significant (red, p-value > 0.05) difference (ERK: 575 cells;
DMSO: 103 cells).

1087

1088

1089     **Figure S12. Predicted onset of differentiation is heterogeneous**
1090     Distribution of single cells predicted onset of differentiation state transition as predicted by the motility
1091     (yellow) and the actin intensity (red) models. N = 343 (motility) and N = 390 (actin intensity) cells were
1092     included in the calculation according to our definition of the onset (see methods). Median start times were
1093     9.42 hours (motility) and 8.67 hours (actin intensity).
1094

1095

**Figure S13. Motility and actin intensity models agree on monotonically increasing differentiation**
**trajectories**

**(A)** Distribution of single cells agreement between the predictions of motility and actin intensity
classifiers, determined by the Pearson correlation coefficient of the correlation between the inferred
trajectories in the differentiation time interval of 7.5-14.5 hours. The agreement was assessed for 575
ERKi-treated cells and 103 DMSO-treated cells. The Median Pearson correlation coefficient (green
dashed line) was 0.48. Colored rectangles mark cells with a correlation coefficient range of [-1, -0.5]
(blue, 10 cells), [-0.5, 0] (yellow, 20 cells) [0, 0.5] (green, 198 cells), and [0.5, 1] (red, 374 cells). **(B)** Pie
chart representing single cell population fractions in different correlation coefficients ranges,
corresponding to the colored rectangles described above. **(C)** Cellular properties analysis of cells sub-
groups, partitioned by the agreement rate between the predictions of the motility and actin intensity
models. Top: mean (solid line) and standard deviation (shade) of cellular properties measured over time:
local density, mean actin intensity, persistence, and speed. Bottom: monotonicity rate, as defined by the
Spearman correlation coefficient between differentiation score and time at the time interval of 7.5-14.5.
Median monotonicity rates were 0.43 (motility) and 0.482 (actin) for range [-1, -0.5] (N=10 cells), 0.06
(motility) and 0.5 (actin) for range [-0.5, 0] (N=20 cells), 0.11 (motility) and 0.53 (actin) for range [0, 0.5]
(N=198 cells), and 0.66 (motility) and 0.72 (actin) for range [-1, -0.5] (N=374 cells).
1113

1114

**Figure S14: Exploration of disagreement between the motility and actin models.**
Left: representative single cells differentiation scores predicted using the motility (yellow) and actin (red)
models. Right: representative snapshots of these cells at time points where the models highly disagree.

**(A)** The examined single cell is barely visible. Thus, actin intensity measurements consist of other cells
rather than the examined one. The Pearson correlation coefficient was -0.91. **(B)** The examined cell
crawled below another cell at several time points, causing bias in the actin intensity measurements. The
Pearson correlation coefficient was -0.23. **(C)** The examined cell entered a crowded region, thus the
intensity measurements represent other cells. The Pearson correlation coefficient was -0.91.
1123

1124

**Figure S15: Persistence in migration is associated with differentiation**
**(A)** Representative single cell's differentiation score (yellow) as predicted by the motility-based model,
and persistence in migration rate (purple) through time. **(B)** Distribution of single cells Pearson
correlation between the difference in differentiation scores and the difference in persistence rates. Values
were calculated within time intervals of 50 minutes. Mean Pearson correlation coefficient was 0.51.
88.07% of cells showed significant correlation (p-value < 0.05; N=575 cells). **(C)** Single cells difference
in differentiation scores (y-axis) over difference in the persistence rate (x-axis) between the beginning and
end of the critical time window where we identified differentiation occurs (7.5 – 14.5 hours). Red
diagonal line indicates the regression line. Pearson correlation coefficient was 0.55 (p-value < 0.0001).
1134

1135

**Figure S16. Local density does not improve quantification of the continuous differentiation state**
Distribution of single cell correlation between the differentiation score and time for classifiers trained
using features that include or exclude local density. Dashed horizontal line shows no correlation. Median
values (shown in white) were 0.09 (local density), 0.62 (local density + actin intensity), 0.67 (actin
intensity), 0.53 (local density + motility), 0.53 (motility). The correlations of the actin intensity classifier
were higher without the local density feature (Wilcoxon rank sign test p-value = $5.7 * 10^{-10}$), the
correlations of the motility classifier were not improved by including the local density feature. N = 575
cells. *** - p-value < 0.0001, n.s – not significant.

1145

**Figure S17. Sensitivity analysis: threshold for terminal differentiation**
Pearson correlation coefficient of the correlation between the (predicted) terminal differentiation time and
the manually annotated fusion time, for different terminal differentiation thresholds. The number of cells
that are identified as terminally differentiated depends on the threshold, thus as the threshold increases-
the number of identified cells decreases.

1151

Before Washing  After Washing

1152
1153
**Figure S18: Co-inhibition of p38 and ERK1/2 leads to properly differentiated ready to fuse cells**
Immunofluorescence images of primary myoblasts treated with p38i+ERKi for 24 hours, before and 24
hours after the inhibitors were washed. The cells were fixed and stained for MyoG (Red), MyHC (green),
and the nuclei (Hoechst, Blue), scale bar 100um. 24 hours after treatment with p38i+ERKi, most of the
cells expressed MyoG, were MyHC negative, and did not undergo fusion. Thus, the cells were
differentiated but un-fused. 24 hours after the inhibitors were washed, there was a decrease in MyoG
positive cells, and most cells express MyHC, indicating that cells completed the differentiation process
and fused.
1162

1163
**Figure S19: Differentiation scores for p38i perturbation**
**(A-B)** Mean differentiation score over time of ERKi- (orange), DMSO- (blue), ERKi+p38i- (green) and
p38i- (purple) treated cells using the motility (A) and actin intensity (B) classifiers. Dashed vertical gray

rectangle highlights the differentiation time interval of 7.5-14.5 hours. ERKi- and DMSO- treated cells
differentiation scores are the same as shown in Fig. 5A-B. Since co-treated cells undergo differentiation
but not fusion, the differentiation score did not decrease after 14.5 hours, unlike ERKi-treated cells that
begin to massively fuse at these times and thus change their motility and actin dynamics (ERKi: 575 cells;
ERKi+p38i: 208 cells; p38i: 202 cells; DMSO: 103 cells). **(C-D)** Mean differentiation score over time
(solid line) and standard deviation (shade) of the cell populations described above (A, B), predicted by the
motility (C) and actin intensity (D) classifiers (ERKi: 575 cells; ERKi+p38i: 208 cells; p38i: 202 cells;
DMSO: 103 cells). **(E-F)** Average cumulative change in differentiation score over time of ERKi treated
cells using the motility (E) and actin intensity models (F). P-values resulting from a t-test between the
cumulative change in differentiation score of cells treated with ERKi and ERKi +p38i are color coded.
(ERKi: 575 cells; ERKi+p38i: 208 cells). **(G-H)** Average cumulative change in differentiation score over
time of Erki+p38i treated cells using the motility (G) and actin intensity models (H). P-values resulting
from a t-test between the cumulative change in differentiation score of cells treated with Erki+p38i and
ERKi are color coded. (ERKi: 575 cells; ERKi+p38i: 208 cells).
1181

1182

1183  **Figure S20: Nuclei fusion manual annotation**
1184  Overlay image showing nuclei in fibers, which were selected for backtracking. All nuclei (magenta) and
1185  actin (cyan) are shown.

1186

**Figure S21. Differentiation scores over time for the entire experiment**
**(A-B)** Mean (solid line) and standard deviation (shade) of the differentiation score over time of ERKi-
(orange) and DMSO- (blue) treated cells using the motility (A) and the actin intensity (B) classifiers.
Dashed vertical gray rectangle highlights the time interval of 7.5-14.5 hours, where both models predicted
the differentiation occurs. The increase of the untreated cells' differentiation score in concurrence with
unchanged (actin) or reduced (motility) scores for ERKi-treated cells around 14.5 hours could be due to
altered motility/actin dynamics of untreated cells in denser microenvironments and due to differentiated
ERKi-treated cells undergoing fusion.

## Supplementary video legends

**Video S1: Proliferating (DMSO) versus Differentiating (ERKi) myoblasts.**
Actin and nuclear reporters (LifeAct-GFP/ nTnG+/+ ) lineage traced primary myoblasts showing
proliferating myoblasts (DMSO-treated myoblasts; left) and differentiating myoblasts (ERKi-treated
myoblasts; right). Time lapse images were acquired using a 10× objective with a 5 minute interval
between frames. Imaging started 1 hour and 30 minutes after treatment. (Time scale: hh:mm). Scale bar
100 µm. Linear adjustments to brightness and contrast were made using ImageJ.

**Video S2: A video of representative single cell trajectories fusing into a single myofiber.**
Cells were tracked and manually edited using Mastodon FIJI plugin (mastodon: Mastodon – a large-scale
tracking and track-editing framework for large, multi-view images, n.d.).

**Video S3: Proliferating (p38i) versus Differentiating without fusion (ERKi-p38i) myoblasts.**
Actin and nuclear reporters (LifeAct-GFP/ nTnG+/+ ) lineage traced primary myoblasts showing
proliferating myoblasts (p38i-treated myoblasts; left) and differentiating myoblasts without fusion (ERKi-
p38i-treated myoblasts; right). Time lapse images were acquired using a 10× objective with a 5 minute
interval between frames. Imaging started 1 hour and 30 minute after treatment. (Time scale: hh:mm).
Scale bar 100 µm. Linear adjustments to brightness and contrast were made using ImageJ.