Harnessing DNA replication stress to target RBM10 deficiency in lung adenocarcinoma

Feras E. Machour[1], Enas Abu-Zhayia[1], Joyce Kamar[2], Alma Sophia Barisaac[1]

Itamar Simon[2], Nabieh Ayoub*[1]

[1]Department of Biology, Technion - Israel Institute of Technology, Haifa 3200003, Israel.

[2]Department of Microbiology and Molecular Genetics, Institute of Medical Research Israel-

Canada, Faculty of Medicine, The Hebrew University, Jerusalem, Israel

*Corresponding author: Prof. Nabieh Ayoub, Faculty of Biology

Technion - Israel Institute of Technology

Haifa 3200003, Israel.

Tel: +972-4-8294232 Fax: +972-4-8225153

Email: ayoubn@technion.ac.il

Abstract (150 words)

The splicing factor RBM10 is frequently mutated in lung adenocarcinoma (LUAD) (9-25%).

Most RBM10 cancer mutations are loss-of-function, correlating with increased tumorigenesis

and limiting targeted therapy efficacy in EGFR-mutated lung cancer. Notably, therapeutic

strategies leveraging RBM10 deficiency remain unexplored. Hence, we conducted RBM10

CRISPR-Cas9 synthetic lethality (SL) screen and identified ~250 RBM10 SL genes, including

WEE1 kinase. WEE1 inhibition sensitized RBM10-deficient LUAD cells *in-vitro* and *in-vivo*.

Mechanistically, we identified a splicing-independent role of RBM10 in promoting replication

fork progression that underpins RBM10-WEE1 SL. Also, we revealed that RBM10 is

associated with active replication forks, which is reliant on PRIM1, an enzyme synthesizing

RNA primers for Okazaki fragments. Functionally, we demonstrated that RBM10 serves as an

anchor for recruiting HDAC1 and facilitates H4K16 deacetylation to maintain replication fork

stability. Collectively, our data revealed a hitherto unrecognized function of RBM10 in fine-

tuning DNA replication, and provide therapeutic arsenal for targeting RBM10-deficient

tumors.

Introduction

Lung cancer has the highest mortality rate among all types of cancer worldwide[1,2]. Non-small cell lung cancer (NSCLC) comprises 85% of all lung cancer cases and is subdivided into different subtypes including lung adenocarcinoma (LUAD) (40%)[3,4]. Most LUAD patients are diagnosed at advanced or metastatic stage, when treatment options are limited to surgery, chemotherapy, and few targeted therapies[4]. EGFR and KRAS are the most mutated oncogenes in LUAD amongst Asian (60%) and Caucasian (33%) cohorts, respectively[5,6]. Accordingly, several targeted therapies, including EGFR and KRAS inhibition, are currently used in the clinic for the treatment of advanced stages of LUAD harboring KRAS and EGFR mutations[7,8]. However, effective treatment of LUAD remains elusive due to the genetic diversity of the disease and the development of therapeutic resistance. Thus, there is a critical need for identifying new therapeutic targets for personalized treatment of LUAD. Interestingly, RNA-binding proteins are broadly dysregulated in human cancers including LUAD and play a key role in carcinogenesis and metastatic progression[9–17]. Therefore, alterations in RNA-binding proteins might provide excellent therapeutic targets for treating lung cancer. One very attractive and yet unexplored therapeutic target is the RNA-binding protein, RBM10.

RBM10 is mapped to the X chromosome[18], and its mutations cause TARP syndrome, an X-linked disorder that leads to pre- and postnatal lethality in affected males[19–25]. RBM10 is a key regulator of alternative splicing that mainly promotes exon skipping of its target genes[26–32]. Also, RBM10 impacts chromatin structure and chromosome segregation independently of its splicing activity[33–36]. Mounting evidence show that RBM10 acts as a tumor suppressor, as its depletion increases cell proliferation and enhances mouse xenograft tumor formation[6,29,32,37–42]. Concordantly, RBM10 overexpression suppresses LUAD tumor growth *in vivo*[6,43]. RBM10 exerts its tumor suppressive activity through various mechanisms[29,40,44]. For example, it was shown that RBM10 inhibits cell proliferation and tumor growth through alternative splicing regulation of NUMB and EIF4H genes[6,32,39]

RBM10 is the most mutated splicing factor in LUAD and is mutated in 9% of all LUAD cases[5]. Strikingly, this percentage rises to 21% in invasive subtypes of LUAD, and 25% in multiple primary LUAD[45,46]. Moreover, RBM10 mutation rate is higher than KRAS and TP53 mutations in early stages of LUAD in patients from Asian origin[47,48]. Most RBM10 mutations in LUAD correspond to loss-of-function and are associated with poor survival[49,50]. Notably, RBM10 mutations mostly co-occur with EGFR and KRAS mutations and were shown to reduce the efficiency of tyrosine kinase inhibitors (TKIs) in LUAD patients harboring EGFR

mutations[6,51]. Altogether, these findings highlight the urgency of therapeutic targeting of

RBM10 deficiency in LUAD.

Herein, we performed a genome-wide CRISPR-Cas9 synthetic lethality (SL) screen in isogenic

LUAD cell line harboring RBM10 cancer mutation and identified ~250 high-scoring RBM10

SL genes, including WEE1 and Aurora A kinases. We show that pharmacological inhibition of

WEE1 selectively sensitizes RBM10-deficient LUAD cells, including patient-derived cells

harboring RBM10 cancer mutations, *in vitro* and in mouse xenograft model, and this effect was

further exacerbated when combined with Aurora A inhibition. Mechanistically, we identified

a splicing-independent role of RBM10 in promoting DNA replication fork progression that

underpins RBM10-WEE1 SL. We also show that RBM10 is associated with active replication

forks, which is contingent upon PRIM1, an enzyme responsible for synthesizing RNA primers

of Okazaki fragments. Functionally, we demonstrated that RBM10 loss disrupts the

localization of HDAC1 at replication forks, leading to elevated levels of H4K16ac, which

destabilizes the replication fork and induces replication stress. Collectively, our data revealed

a hitherto unrecognized function of the RNA-binding protein RBM10 in fine-tuning DNA

replication, and identify DNA replication stress as an SL pathway with RBM10 loss. Moreover,

we provide a repertoire of RBM10 SL targets that can be harnessed therapeutically to eradicate

RBM10-deficient tumors with immediate clinical applicability.

Results

Genome-wide CRISPR-Cas9 screen reveals synthetic lethal partners of RBM10 in LUAD

cells

Given that RBM10 is one of the most mutated genes in LUAD, we sought to identify genetic

vulnerabilities to RBM10 loss. To achieve this, we conducted an unbiased genome-scale

CRISPR-Cas9 SL screen in LUAD HCC827 cell line. HCC827 cells harbor a common

oncogenic deletion in EGFR (E746-A750), and are a pertinent model to screen for genetic

vulnerabilities to RBM10 loss since RBM10 and EGFR mutations often co-occur in LUAD,

and RBM10 loss was shown to limit the response to common EGFR-targeted therapy[6,51].

Therefore, we sought to generate isogenic HCC827 cells stably expressing comparable levels

of flag-Cas9 endonuclease and differ only in the status of RBM10. Toward this end, we

designed a sgRNA targeting RBM10 exon 2 to introduce a frameshift mutation V176Rfs*, that

exhibits a complete absence of RBM10 protein, while constitutively expressing Cas9 protein

(HCC827-Cas9[RBM10-KO]) (Fig. 1a and Supplementary Fig. 1a,b). Interestingly, the introduced

mutation closely resembles a previously characterized LUAD cancer mutation E177* that

abrogates RBM10 splicing activity and leads to RBM10 protein loss[50]. Notably, HCC827-

Cas9[RBM10-KO] cells exhibit an increase in exon inclusion of RBM10 splicing targets NUMB and

EIF4H, phenocopying the defective splicing phenotype of RBM10-deficient LUAD[6,32] (Fig.

1b and Supplementary Fig. 1c).

Next, we conducted a genome-scale CRISPR-Cas9 SL screen in the isogenic HCC827-Cas9[WT]

and HCC827-Cas9[RBM10-KO] using the TKOv1 sgRNA library, which contains 91,320 sgRNAs

targeting 17,232 protein-coding genes[52]. Cells were collected at 0 and 15 days following

TKOv1 library transduction and subjected to next-generation sequencing to measure sgRNA

abundance (Supplementary Fig. 1d). To identify bona fide RBM10 SL genes that are

selectively essential in RBM10-KO cells, we analyzed the screen results using three distinct

scoring methods: Bayesian Analysis of Gene Essentiality (BAGEL2)[53], Model-Based Analysis

of Genome-wide CRISPR-Cas9 Knockout (MAGeCK)[54], and CRISPR Count Analysis

(CCA)[55]. These methods unveiled 210, 280, and 357 RBM10 SL genes, respectively (Fig. 1c,

Supplementary Fig. 1e, and Supplementary Table 1). Of note, a comparative analysis of

RBM10 SL genes among the three different assays, revealed 60 RBM10 SL genes that are

common to all three analysis methods (Supplementary Fig. 1f). Gene-set enrichment analysis

revealed significant enrichment of several pathways including mRNA splicing, cell cycle

checkpoints, S-phase, and DNA replication (Fig. 1d).

To functionally validate the screen results, we conditionally knocked-down 5 candidate

RBM10 SL genes annotated to various enriched pathways (SF3B1, WAPAL, WEE1, AURKA,

PEX10), and 2 common essential (XRN2, LSM7), followed by clonogenic survival assay.

While depletion of common essential genes had a similar effect in inhibiting the proliferation

of both parental and RBM10-KO cells, depletion of candidate RBM10 SL genes selectively

inhibited the proliferation of HCC827-Cas9[RBM10-KO] cells, confirming successful identification

of bona-fide RBM10 SL genes (Fig. 1e and Supplementary Fig. 1g).

Interestingly, several RBM10 SL genes, such as WAPAL, WEE1, CHEK1, ATRIP, and

MRE11, were identified using the three aforementioned scoring methods. These genes have

known functions in DNA replication stress response and replication fork protection, suggesting

DNA replication stress as a synthetic lethal pathway with RBM10 deficiency in LUAD (Fig.

1f). In support of this, RBM10 was previously identified as a candidate replication stress

response gene by a genome-wide siRNA screen[56], and a phosphorylation target of ATR[57].

Additionally, RBM10 was shown to interact with several proteins involved in DNA replication

and repair in fission yeast[36]. Fittingly, analysis of "The Cancer Genome Atlas" (TCGA[58])

mutation data from LUAD patients revealed that RBM10 deficiency is associated with increased tumor mutational burden and replication stress gene signature (Supplementary Fig. 1h,i). Collectively, these observations suggest a role of RBM10 in DNA replication and genomic stability.

Figure 1: Genome-wide CRISPR-Cas9 screen reveals synthetic lethal partners of RBM10 in LUAD cells. (a) Immunoblot analysis for RBM10 and flag-Cas9 protein expression in isogenic RBM10-deficient HCC827 cells constitutively expressing flag-Cas9. β-actin is used as a loading control. The positions of molecular weight markers are indicated to the right. **(b)** RT-PCR analysis of NUMB exon 9 alternative splicing in parental (WT) and HCC827-Cas9$^{RBM10-KO}$ cells. RNA was isolated from the indicated cell lines and analyzed by RT-PCR using primers flanking NUMB exons 8-10. Left: Representative agarose gel image showing amplification of two NUMB variants that differ in exon 9 inclusion. Right: Precent-spliced-in

(PSI) quantification of NUMB exon 9 inclusion. Data are presented as mean ± s.d. (n=3).
Statistical significance was determined by unpaired t-test. ***P<0.001*. **(c)** Results of CRISPR-
Cas9 SL screen in WT and HCC827-Cas9[RBM10-KO] cells performed in triplicates. CRISPR
Counts Analysis (CCA) score is plotted against the difference in gene essentiality score,
expressed as Bayes Factor (BF), between RBM10-KO cells and WT cells. RBM10 SL genes
are shown in blue. **(d)** Gene ontology analysis of RBM10 SL genes identified in the CRISPR-
Cas9 SL screen. **(e)** Clonogenic survival of WT and HCC827-Cas9[RBM10-KO] cells transduced
with inducible vector expressing either scramble shRNA or shRNA targeting the indicated
genes upon the addition of doxycycline (DOX). Top: Representative images of crystal violet
staining of the indicated cell lines treated with DOX. Bottom: Quantification of clonogenic
survival in DOX-treated cells normalized to untreated cells. Data are presented as mean ± s.d.
(n=3). Statistical significance was determined by unpaired t-test.
P<0.01;P<0.001;****P<0.0001*. **(f)** STRING interaction network of RBM10 SL genes
involved in replication stress response and cell cycle. The thickness of the connecting line
indicates interaction confidence.
163

RBM10 promotes DNA replication fork progression and replication stress response

Herein, we sought to determine the impact of RBM10 loss on replication fork progression.

DNA fiber analysis showed that HCC827-Cas9[RBM10-KO] cells display a significant replication

fork slowdown, suggesting that RBM10 is required for proper fork progression (Fig. 2a).

Similar results were also observed in a second RBM10-KO clone (KO2) isolated from

HCC827-Cas9 cells, arguing against a clonal effect (Supplementary Fig. 2a). To confirm that

the effect of RBM10 loss on replication fork progression is not cell-line specific, and since

TP53 mutations often co-occur with RBM10 mutations in LUAD[58], we tested the effect of

RBM10 knockout on replication fork progression in p53-deficient LUAD cells NCI-H1299

(H1299[RBM10-KO]) (Supplementary Fig. 2b,c). Results show that RBM10 loss in H1299 cells also

led to significant reduction in replication fork progression, suggesting that the effect of RBM10

on DNA replication is independent of common co-occurring mutations such as EGFR and

TP53 (Fig 2b). Immunostaining analysis revealed that replication fork slowdown upon RBM10

loss is accompanied by an increase in the levels of pRPA32-S33 and γH2AX, primarily

observed in S phase cells marked by EdU staining (Fig. 2c-e). Similarly, western blot analysis

demonstrated a significant increase in pRPA32-S33, pCHK1, and γH2AX in RBM10 knockout

(KO) cells synchronized at the S phase using a double-thymidine block (Supplementary Fig.

2d). Next, we investigated the effect of RBM10 loss on the recovery from hydroxyurea (HU)

induced replication stress. Results showed that RBM10-deficient cells display a significant

increase in replication stress and DNA damage markers after release from HU, with no

significant effect on replication fork restart (Fig. 2f and Supplementary Fig. 2e). Corollary to

this, RBM10-deficient cells are markedly more sensitive than RBM10-proficient cells to

increasing HU concentrations (Fig. 2g). Collectively, our results demonstrate that RBM10 fosters efficient DNA replication fork progression and fork recovery during replication stress.

Figure 2: RBM10 promotes DNA replication fork progression and replication stress response. (a,b) Replication fork speed measurement using DNA combing assay in WT and RRBM10-KO HCC827 (a) or H1299 (b) cells. Horizontal bars represent mean value of replication fork speed (n>200). Statistical significance was determined by Mann-Whitney test. *****P<0.0001*. **(c)** Representative immunofluorescence microscopy images of pRPA32-S33 and γH2AX foci in H1299^{WT} and H1299$^{RBM10-KO}$ cells. EdU is used to mark S-phase cells. DAPI is used to stain nuclei. Scale bar, 20μm. **(d,e)** Quantification of γH2AX (d) and pRPA32-S33 (e) foci in S-phase (EdU-positive) H1299^{WT} and H1299$^{RBM10-KO}$ cells. Data are presented as mean foci number per nucleus ± SEM (n>200 cells) and representative of 3 independent experiments. Statistical significance was determined by Mann-Whitney test. *****P<0.0001*. **(f)** H1299^{WT} and H1299$^{RBM10-KO}$ cells were treated with 2mM hydroxyurea (HU) for 2 hr and subjected to immunoblot analysis at the indicated times after release from HU. Histone H3 is used as a loading control. UT=untreated. The positions of molecular weight markers are indicated to the right. **(g)** Short-term cell viability assay in parental (WT) and RBM10-KO HCC827-Cas9 cells treated with increasing concentrations of HU. Data are presented as mean ± s.d. (n=3).

RBM10 is associated with active DNA replication fork in a PRIM1-dependent manner

To gain molecular insights into the involvement of RBM10 in replication fork progression, we first determined its sub-cellular localization. Biochemical fractionation revealed that RBM10 is enriched at the chromatin-bound fraction (Supplementary Fig. 3a), supporting the notion that RBM10 has a direct role in regulating replication fork progression. To further substantiate this, cells expressing RBM10 fused to EGFP were subjected to GFP-trap followed by mass spectrometry (MS) analysis to map RBM10 interactome. Results revealed 410 RBM10-interacting proteins annotated to several pathways, most notably the spliceosome, consistent with RBM10 function in alternative splicing, as well as DNA replication. (Fig. 3a and Supplementary Table 2). Intriguingly, we revealed that RBM10 interacts with replication fork components and proteins involved in replication stress response (Supplementary Fig. 3b). Next, we validated the authenticity of RBM10 interactome and showed that RBM10 interacts with MCM5, PCNA, HDAC1, PARP1, and RAD51 (Fig. 3b).

To further substantiate RBM10 interaction with replication forks, first we confirmed the interaction between endogenous RBM10 and PCNA in LUAD cells (Supplementary Fig. 3c). Second, we tested whether RBM10 is associated with active replication forks in cells using proximity ligation assay (PLA). Results showed that RBM10 is in close proximity to PCNA (Fig. 3c). Moreover, PLA combined with EdU click chemistry revealed that RBM10 is associated with EdU-labelled nascent DNA, marking sites of active DNA replication (Fig. 3d). Collectively, our data provide firm evidence that RBM10 interacts with active DNA replication forks under native conditions.

Next, we aimed to gain molecular insights into the mechanism governing the association of RBM10 with DNA replication forks. Given that RBM10 is a well-established RNA-binding protein, we hypothesized that its binding to RNA molecules facilitates its association with chromatin. To explore this hypothesis, we treated cells with RNase A followed by chromatin fractionation. Our results revealed that RNase A treatment profoundly disrupts RBM10 association with chromatin (Supplementary Fig. 3d). Consistent with this observation, an RBM10 deletion mutant lacking the first ZnF domain (RBM10$^{\Delta ZNF1}$), known to bind RNA molecules[59,60], lost its association with active DNA replication forks, as determined by PLA (Fig. 3e). Remarkably, a recent work has shown that PRIM1, an enzyme involved in synthesizing RNA primers for Okazaki fragments, facilitates the recruitment of 53BP1 to DNA replication forks[61]. Hence, we investigated whether a similar scenario exists with RBM10. Our results reveal that siRNA knockdown of PRIM1 pronouncedly diminishes the association of

240 RBM10 with active DNA replication forks similar to 53BP1 (Fig. 3f and Supplementary Fig.

241 4a,b). Collectively, these findings robustly support the notion that Okazaki RNA primers play

242 a pivotal role in mediating the recruitment of RBM10 to replication forks.

Figure 3: PRIM1-dependent RBM10 association with DNA replication forks promotes HDAC1 recruitment to limit replication stress. (a) Gene ontology analysis of RBM10 interactome in unchallenged cells as identified by GFP-trap followed by mass spectrometry analysis. **(b)** HEK293T cells expressing EGFP-RBM10 or EGFP only were subjected to GFP-trap followed by immunoblot analysis using the indicated antibodies. Bottom: GFP-trap was performed on HEK293T cells additionally expressing Flag-MCM5. The positions of molecular weight markers are indicated to the right. **(c,d)** Representative immunofluorescence microscopy images of RBM10:PCNA (c) and RBM10:EdU-biotin (d) proximity ligation assay (PLA) foci in H1299[WT] and H1299[RBM10-KO] cells. Scale bar, 20μm. **(e)** Left: Representative immunofluorescence microscopy images of RBM10:EdU-biotin PLA in H1299[RBM10-KO] expressing Flag-RBM10[WT], Flag-RBM10[ΔZNF1], or vector only (VO). DAPI is used to stain nuclei. Scale bar, 20μm. Right: Quantification of RBM10-EdU-biotin PLA intensity per nucleus. Data are presented as mean PLA intensity per nucleus ± SEM (n>300 cells). Statistical significance was determined by Mann-Whitney test. *****P<0.0001*. **(f)** As in (e), except H1299[WT] were transfected with siRNA against PRIM1 (siPRIM1) or control siRNA (siCtrl) 72

hr prior to PLA assay. Scale bar, 50µm. **(g)** Immunoblot analysis of H1299[RBM10-KO] cells expressing Flag-RBM10[WT], Flag-RBM10[ΔRRM2], Flag-RBM10[ΔZNF1], or vector only (VO). Top: Representative immunoblot image using the indicated antibodies. α-Tubulin is used as a loading control. Bottom: Quantification of γH2AX band intensity (relative to H3) and pRPA32-S33 band intensity (relative to RPA32). Data are presented as mean ± s.d. (n=3). Statistical significance was determined by unpaired t-test. **P<0.01;****P<0.0001;ns, not significant.** **(h,i)** Left: Representative immunofluorescence microscopy images of HDAC1:EdU-biotin (h) and H4K16ac:EdU-biotin (i) PLA in H1299[WT] and H1299[RBM10-KO] cells. Cells were labelled with EdU for 20 min and were either left untreated (UT), treated with 1 mM HU for 1 h (HU). DAPI is used to stain nuclei. Scale bar, 20µm. Right: Quantification of PLA intensity per nucleus. Data are presented as mean PLA intensity per nucleus ± SEM (n>50). Statistical significance was determined by Mann-Whitney test. *P<0.05;**P<0.01;***P<0.001.* **(j)** Top: Representative immunofluorescence microscopy images of R-loops detected by S9.6 antibody in H1299[WT] and H1299[RBM10-KO] cells. Cells overexpressing RNaseH1 are used as a negative control. DAPI is used to stain nuclei. Scale bar, 10µm. Bottom: Quantification of S9.6 signal in H1299[WT] and H1299[RBM10-KO] cells. Data are presented as mean intensity per nucleus ± SEM (n>50 cells) and representative of 3 independent experiments. Statistical significance was determined by Mann-Whitney test. ****P<0.0001.*

The role of RBM10 in replication stress response is distinct from its splicing activity

Since RBM10 modulates the splicing of hundreds of genes, we sought to test whether the splicing activity of RBM10 underlies its role in the DNA replication stress response. To address this, we reintroduced RBM10-deficient cells with a vector expressing either flag-RBM10-WT or RBM10[ΔZNF1] or RBM10 mutant that lack RRM2 domain (RBM10[ΔRRM2]), known to modulate RBM10 splicing activity[60]. Subsequently, we assessed the ability of these RBM10 mutants to restore both splicing efficiency and the levels of replication stress markers to normal levels.

Our results demonstrate that while RBM10[ΔZNF1] successfully restores the splicing efficiency of RBM10 target genes, NUMB and EIF4H, it still exhibits increased levels of replication stress markers compared to cells expressing RBM10-WT (Fig. 3g and Supplementary Fig. 4c,d). Conversely, RBM10-deficient cells expressing RBM10[ΔRRM2] mutant show defects in the splicing of RBM10 target genes but display normal levels of replication stress markers compared to control cells. These findings collectively support the notion that the role of RBM10 in DNA replication is independent of its splicing activity. To further validate this observation, we conducted transcriptome analysis of RBM10 in isogenic HCC827 and NCI-H1299 cells. Results revealed no discernible changes in the expression of core DNA replication and repair genes (Supplementary Fig. 4e and Supplementary Table 2). Our findings align with previously reported transcriptome analyses of RBM10 in various cell lines, demonstrating that RBM10 does not regulate the expression or splicing of core DNA replication and repair genes[6,32].

RBM10 targets HDAC1 to active replication fork to fine-tune histone acetylation

To elucidate the mechanistic role of RBM10 in DNA replication progression and replication stress response, we focused on RBM10-HDAC1 interaction that has been identified in the RBM10 interactome and validated through immunoprecipitation (Fig. 3b and Supplementary Fig. 3b). Also, fission yeast Rbm10 was previously shown to interact with Clr6, a homolog of human HDAC1/2[36]. In addition, previous reports have shown that HDAC1 is localized at active DNA replication forks, where it deacetylates histones to prevent replication fork collapse[62–65]. Prompted by this, we sought to investigate whether RBM10 is essential for recruiting HDAC1 to DNA replication forks. Our results demonstrated a pronounced disruption in the association of HDAC1 with DNA replication in RBM10-deficient cells under normal and stress conditions (Fig. 3h). Consequently, the levels of H4K16 acetylation were increased in RBM10-deficient cells compared to control cells (Fig. 3i). In summary, these data suggest that RBM10 serves as an anchor for recruiting HDAC1 to active replication forks to facilitate the deacetylation of H4K16, a modification recently recognized as crucial for replication fork stability[65,66]. Since a recent work demonstrated that histone deacetylation counteracts the levels of R-loops at active replication forks to mitigate transcription-induced replication stress[66], we postulated that the increased levels of H4K16ac in RBM10-deficient cells is also accompanied by increased R-loops levels. Indeed, we observed that RBM10-deficient cells exhibit elevated levels of R-loops in S-phase cells, indicating that replication stress observed upon RBM10 loss is mediated, at least in part, by transcription-replication conflicts and R-loop accumulation (Fig. 3j and Supplementary Fig. 4f,g).

Additionally, since RBM10 interacts with RAD51 protein (Figure 3b) which has a central role in homology repair (HR) of double-strand breaks (DSB) and its depletion impairs HR repair leading to replication stress[67,68], we sought to investigate the functional relevance of their interaction[67,68]. We aimed to determine whether the replication stress in RBM10-deficient cells is due to defective HR repair. Toward this, we assessed the efficacy of HR repair of DSBs using the Cas9-mClover-LMNA assay[69]. Results indicate that RBM10 deficiency has no discernible effect on the percentage of mClover-positive cells (representing cells that repaired DSBs through HR) compared to control cells (Supplementary Fig. 4h). This observation suggests that the replication stress observed in RBM10-deficient cells is not attributed to defective HR repair. This data raises an interesting possibility that the RBM10-RAD51 interaction might be required for replication fork reversal rather than HR repair.

WEE1 kinase inhibition selectively sensitizes RBM10-deficient LUAD cells

Herein, we sought to exploit the role of RBM10 in replication stress to selectively target

RBM10-deficient LUAD cells. Towards this, we focused on WEE1 kinase, since it is a high

scoring RBM10 SL gene that was identified in the three scoring methods (BAGEL, MAGeCK

and CCA) (Fig. 1b and Supplementary Fig. 1e,f), and is involved in several RBM10 SL

pathways, such as S-phase and G2/M cell cycle checkpoints, and was shown to protect the

stability of stalled replication forks[70–72]. Moreover, WEE1 has a small molecule inhibitor,

MK1775 (AZD1775), which is currently in several phase II clinical trials for treatment of

advanced solid tumors (ClinicalTrials.gov), highlighting the clinical potential of targeting

RBM10-deficient LUAD[73–75]. To test whether the observed RBM10-WEE1 synthetic lethality

(Fig. 1e) is dependent on WEE1 kinase activity, we pharmacologically inhibited WEE1 using

MK1775 in parental HCC827-Cas9^{WT} and two HCC827-Cas9$^{RBM10-KO}$ cell lines. Short and

long-term survival assays revealed that HCC827-Cas9$^{RBM10-KO}$ cells exhibit pronounced

sensitivity to MK1775 treatment compared to parental cells (Fig. 4a,b, Supplementary Fig. 5a).

The hypersensitivity of RBM10-deficient cells to MK1775 was also observed in RBM10-KO

(KO3) cells derived from naïve HCC827 cells that do not stably express Cas9, suggesting that

RBM10-WEE1 synthetic lethality is not due to a clonal effect or cytotoxicity caused by stable

expression of Cas9 endonuclease (Supplementary Fig. 5b,c). In addition, complementing

RBM10-deficient cells with a vector expressing flag-RBM10-WT suppressed the sensitivity to

MK1775 compared to cells expressing empty vector (Fig. 4c).

Similar to p53-proficient HCC827, our results showed that H1299$^{RBM10-KO}$ cells are

hypersensitive to MK1775 compared to control cells (Fig. 4d,e). To increase the therapeutic

potential of our observations, we tested RBM10-WEE1 synthetic lethality in two RBM10-

deficient patient-derived cell lines, NCI-H1944 (containing KRAS^{G13D} mutation) and NCI-

H1975 (containing EGFR$^{L858R/T790M}$ mutation) harboring p.A683Rfs*10 and p.G905Afs*7

truncating RBM10 mutations, respectively. Towards this, NCI-H1944 and NCI-H1975 cells

were transduced with flag-RBM10 expression vectors (Supplementary Fig. 5d,e) and treated

with MK1775 followed by cell viability assays. Results showed that RBM10 expression

rescues the sensitivity of RBM10-deficient cell lines to MK1775, suggesting that RBM10 loss

occurring during LUAD tumorigenesis potentiates the sensitivity to WEE1 inhibition (Fig.

4f,g). Altogether, our results recapitulate RBM10-WEE1 synthetic lethality observed in our

CRISPR-Cas9 SL screen and demonstrate that WEE1 inhibition selectively sensitizes RBM10-

deficient LUAD tumor cells from different origins and independently of common co-occurring

mutations such as EGFR, KRAS and TP53, highlighting the clinical relevance of targeting

RBM10 loss in LUAD using MK1775.

Figure 4: WEE1 kinase inhibition selectively sensitizes RBM10-deficient LUAD cells. (a)
Short-term cell viability assay and EC_{50} determination in WT and two RBM10-KO clones (KO
and KO2) of HCC827-Cas9 cells treated with increasing concentrations of MK1775. Data are
presented as mean ± s.d. (n=3). **(b)** Clonogenic survival of WT and two RBM10-KO clones
treated with the indicated concentrations of MK1775. Left, representative images of plates
stained with crystal violet. Right, quantification of clonogenic survival. Data are presented as
mean ± s.d. (n=3). **(c)** As in (a), except of using HCC827$^{RBM10-KO3}$ cells expressing flag-
RBM10-WT or vector only. **(d)** As in (a), except of using H1299^{WT} and H1299$^{RBM10-KO}$ cells.
(e) Clonogenic survival of H1299^{WT} and H1299$^{RBM10-KO}$ cells treated with the indicated
concentrations of MK1775. Top: Representative images of plates stained with crystal violet.
Bottom: Quantification of clonogenic survival. Data are presented as mean ± s.d. (n=3). **(f,g)**
As in (a), except of using NCI-H1944 (f) or NCI-H1975 (g) cells expressing either flag-
RBM10-WT or vector only.

DNA damage and premature mitotic entry underpin RBM10-WEE1 synthetic lethality

We sought to decipher the molecular basis underlying the hypersensitivity of RBM10-deficient cells to WEE1 inhibition. We showed that H1299[RBM10-KO] cells treated with increasing concentrations of MK1775 exhibit elevated levels of phosphorylated histone H2A.X (γH2AX), suggesting that DNA damage accumulation upon WEE1 inhibition is exacerbated in RBM10-deficient cells (Fig. 5a). Likewise, higher levels of stalled forks and replication stress markers including phosphorylated RPA32 on positions 4 and 8 (pRPA32-S4/S8), RPA32 on position 33 (pRPA32-S33), and CHK1 on position 345 (pCHK1-S345) were observed in MK1775-treated H1299[RBM10-KO] cells compared to control cells (Fig. 5b and Supplementary Fig 6a,b). Since WEE1 inhibition in RBM10-deficient cells leads to excessive replication stress, we investigated the effect of MK1775 treatment on replication fork progression. Results showed that WEE1 inhibition exacerbates the decrease in replication fork rate in RBM10-deficient cells compared to parental cells (Supplementary Fig. 6c). To determine whether the increase in replication stress following WEE1 inhibition was due to an increase in DNA breakage, we used alkaline and neutral comet assays to measure the single-stranded DNA (ssDNA) breaks and double-stranded DNA (dsDNA) breaks, respectively. Results show that MK1775 treatment in RBM10-KO cells led to increased accumulation of both ssDNA and dsDNA breaks, with the increase in ssDNA breaks being more prominent (Fig. 5c,d and Supplementary Fig. 6d). Next, we utilized native BrdU staining to further confirm the increase in ssDNA in H1299[RBM10-KO] cells upon MK1775 treatment (Fig. 5e,f). To address whether the accumulation of ssDNA upon WEE1 inhibition is due to ssDNA gaps arising during DNA replication, we used a modified DNA combing assay incorporating a digestion step with S1 nuclease[76]. Results revealed that labelled DNA tracks in MK1775-treated RBM10-KO cells showed high sensitivity to S1 nuclease digestion compared to MK1775-treated parental cells, suggesting WEE1 inhibition in RBM10-KO cells leads to accumulated ssDNA gaps during DNA replication (Fig 5g). Altogether, our results suggest that WEE1 inhibition exacerbates replication stress upon RBM10 loss and that replication stress underpins, at least partly, RBM10-WEE1 synthetic lethality.

Next, we sought to determine the effect of WEE1 inhibition on cell cycle progression in RBM10-deficient cells. Towards this, parental and H1299[RBM10-KO] cells were co-stained for pH3(Ser10), a marker of early mitosis, and EdU as a marker for S-phase following MK1775 treatment. Remarkably, unlike parental cells, a significant fraction (~45%) of MK1775-treated

H1299[RBM10-KO] cells were co-stained with EdU and pH3(Ser10), suggesting that WEE1 inhibition leads to premature mitotic entry in RBM10-deficient cells (Fig. 5h,i). Similar results were also obtained in parental and HCC827[RBM10-KO] cells (Fig. 5j). Consequently, we observed that MK1775-treated H1299[RBM10-KO] cells exhibit high levels of micronucleation, extensive nuclear γH2AX staining and apoptosis (Fig. 5k,l and Supplementary Fig. 6e). Altogether, we concluded that the sensitivity of RBM10-deficient cells to MK1775 is mediated by increased replication stress and premature mitotic entry leading to mitotic catastrophe and cell death.

Figure 5: DNA damage and premature mitotic entry underpin RBM10-WEE1 synthetic lethality. (a) Quantification of γH2AX staining in H1299[WT] and H1299[RBM10-KO] cells treated with 300nM MK1775 for 24 hr. Data are presented as mean nuclear intensity ± SEM (n>590 cells) and representative of 3 independent experiments. Statistical significance was determined by Mann-Whitney test. ****$P<0.0001$. **(b)** Immunoblot analysis of DNA damage and replication stress markers in H1299[WT] and H1299[RBM10-KO] cells after MK1775 treatment. Cells were either left untreated (UT) or treated with the indicated concentrations of MK1775 for 24 hr followed by immunoblot analysis using the indicated antibodies. Histone H3 is used as a loading control. The positions of molecular weight markers are indicated to the right. **(c,d)** Alkaline comet assay in H1299[WT] and H1299[RBM10-KO] cells treated either with DMSO or the

indicated concentrations of MK1775 for 24 hr. (c) Representative fluorescence microscopy images of alkaline comet assay. (d) Quantification of DNA damage represented by comet tail moment. Data are presented as mean tail moment ± SEM (n>150) and representative of 3 independent experiments. Statistical significance was determined by Mann-Whitney test. ***P<0.001;****P<0.0001*. **(e,f)** Analysis of ssDNA levels using native BrdU staining in H1299[WT] and H1299[RBM10-KO] cells treated with 300nM MK1775 for 24 hr. (e) Representative immunofluorescence microscopy image for native BrdU staining. DAPI is used to stain nuclei. Scale bar, 50μm. (f) Quantification of native BrdU staining. Data are presented as mean nuclear intensity ± SEM (n>100 cells for untreated cells, n>300 for MK1775-treated cells) and representative of 3 independent experiments. Statistical significance was determined by Mann-Whitney test. *****P<0.0001*. **(g)** H1299[WT] and H1299[RBM10-KO] cells were either untreated or treated with 300nM MK1775 for 4 hr followed by DNA combing analysis as indicated. The length of CldU+ region of IdU+ CldU+ replication tracks was measured following digestion with S1 nuclease for 1 hr. Data are presented as mean track length ± SEM (n>100). Statistical significance was determined by Mann-Whitney test. *P<0.05;****P<0.0001*. **(h)** Representative immunofluorescence microscopy images showing premature mitotic entry in H1299[WT] and H1299[RBM10-KO] cells treated with MK1775. EdU incorporation and pH3(Ser10) were used to determine DNA synthesis and mitotic entry, respectively. DAPI is used to stain nuclei. Scale bar, 20μm. **(i,j)** Quantification of premature mitotic entry events in parental (WT) and H1299[RBM10-KO] (i) and HCC827[RBM10-KO] (j) cells. Premature mitosis events were calculated as the percentage of EdU-positive cells from the pH3(Ser10)-positive (mitotic) cells and presented as mean ± s.d. (n=6). Statistical significance was determined by unpaired t-test. *****P< 0.0001*. **(k)** Representative immunofluorescence microscopy images of γH2AX and DAPI staining in H1299[WT] and H1299[RBM10-KO] cells treated with 300nM MK1775 for 24 hr. Scale bar, 10μm. **(l)** Quantification of micronuclei in H1299[WT] and H1299[RBM10-KO] cells treated with 300nM MK1775 for 24 hr. Data are presented as the average percentage of cells with micronuclei ± SEM (n=3). Statistical significance was determined by unpaired t-test. *****P<0.0001*.

MK1775 inhibits the growth of RBM10-deficient LUAD tumors *in vivo*

Given that the WEE1 inhibitor MK1775 is currently undergoing phase II clinical trials for treating several tumor types[77], we sought to test the therapeutic potential of WEE1 inhibition against RBM10-deficient LUAD cells *in vivo*. Our results showed that MK1775 treatment prominently reduced the size of xenograft tumors derived from HCC827[RBM10-KO] and H1299[RBM10-KO] cells, but not their parental counterparts (Fig. 6a,b and Supplementary Fig. 7a,b). Consistent with the observed γH2AX phenotype *in vitro*, RBM10-deficient tumors exhibit increased levels of DNA damage in MK1775-treated mice (Fig. 6c,d and Supplementary Fig. 7c).

Collectively, these findings strongly suggest that RBM10 loss leads to increased sensitivity to WEE1 inhibition *in vivo*, independently of EGFR or TP53 status. To further substantiate the clinical relevance of targeting RBM10-deficient LUAD, we tested the anti-tumor effect of MK1775 against tumor xenografts derived from patient-derived LUAD cells, NCI-H1944 and

NCI-H1975, harboring naturally occurring RBM10 cancer mutations. Results showed that WEE1 inhibition leads to remarkable tumor growth inhibition in RBM10-deficient NCI-H1975 and NCI-H1944 compared to RBM10-proficient cells (Fig. 6e). Altogether, our results

demonstrate the therapeutic efficacy of MK1775 as a single-agent in eradicating LUAD tumors harboring RBM10 deleterious cancer mutations.

Figure 6: MK1775 inhibits the growth of RBM10-deficient LUAD cells *in vivo*. (a,b) Tumor weight of parental (WT) and HCC827-Cas9$^{RBM10-KO}$ (a) and H1299$^{RBM10-KO}$ (b) xenografts treated with either MK1775 or vehicle. MK1775 was administered once daily at 40mg/kg for 15 days. Results are shown as mean tumor weight ± SEM (n=6). Statistical significance was determined by unpaired t-test. *P<0.05;**P<0.01*. **(c,d)** Quantification of γH2AX immunostaining in xenograft tumor sections from parental and HCC827-Cas9$^{RBM10-KO}$ (c) and H1299$^{RBM10-KO}$ (d) tumors treated with either MK1775 or vehicle. Data are presented as the average percentage γH2AX positive cells ± SEM of 3 different tumor sections for each group. Statistical significance was determined by unpaired t-test. **P<0.01;***P<0.001*. **(e)** Tumor weight of NCI-H1944 and NCI-H1975 xenografts treated with either MK1775 or vehicle. MK1775 was administered once daily at 40mg/kg for 15 days. Results are shown as mean tumor weight ± SEM (n=5 for NCI-H1944, n=6 for NCI-H1975). Statistical significance was determined by unpaired t-test. **P<0.01;****P<0.00001*. **(f)** Clonogenic survival of H1299^{WT} and H1299$^{RBM10-KO}$ cells treated with the indicated concentrations of alisertib. Left, representative images of plates stained with crystal violet. Right, quantification of clonogenic

survival. Data are presented as mean ± s.d. (n=3). **(g)** Short-term cell viability assay and EC_{50}
determination in H1299^{WT} and H1299$^{RBM10-KO}$ cells treated with increasing concentrations of
alisertib with or without treatment with 300nM MK1775. Data are presented as mean ± s.d.
(n=3). **(h)** Tumor growth inhibition in H1299^{WT} and H1299$^{RBM10-KO}$ xenografts treated with
MK1775 alone, MK1775 and alisertib combination, or vehicle. MK1775 alone or MK1775 and
alisertib combination were administered once daily at 30mg/kg for 15 days. Results are shown
as percentage tumor growth inhibition relative to vehicle ± SEM (n=6 for MK1775 alone and
vehicle, n=5 for MK1775 + alisertib combination). Statistical significance was determined by
unpaired t-test. ***P<0.001*.

Aurora kinase A inhibition exacerbates the anti-tumor activity of WEE1 inhibition in
RBM10-deficient LUAD
Given that both WEE1 and Aurora A (AURKA) scored highly in RBM10 SL screen and both
kinases are targeted by small molecule inhibitors undergoing clinical trials, we tested the
efficacy of the combined Aurora A and WEE1 inhibition on the growth of RBM10-deficient
LUAD. First, we observed that the Aurora A inhibitor, alisertib, had a profound effect on the
clonogenic survival of RBM10 KO cells, concurrent with the CRISPR SL screen results (Fig.
6f and Supplementary Fig. 7d). Moreover, while alisertib alone led to a significant reduction
in RBM10-deficient cell viability, combined MK1775 and alisertib treatment further
exacerbated cell toxicity (Fig. 6g). Prompted by this, we tested the effect of the combined
MK1775 and alisertib treatment on the growth of RBM10-deficient xenograft tumors. Results
showed that the combined treatment leads to a profound and significant reduction in RBM10-
deficient tumor size that is greater than the reduction upon MK1775 treatment alone (Fig. 6h).
Therefore, we concluded that Aurora kinase A inhibition might be a viable therapeutic strategy
to enhance the anti-tumor activity of WEE1 inhibition in RBM10-deficient LUAD tumors.

Discussion
We present a model that underscores a new function of RBM10 in both DNA replication
progression and replication stress response (Fig. 7). RBM10 interactome analysis revealed that
it interacts with various replication fork components that are required for proper replication
fork progression, as well as recovery from replication stress. Also, we show that RBM10
depletion leads to replication fork slowdown, which is accompanied by an increase in DNA
damage. Our data are in line with a genome-wide siRNA screen that identified RNA processing
factors, including RBM10, as replication stress response genes[56]. Moreover, previous studies
showed that RBM10 is associated with human pre-replication complex[78] and interacts with
several proteins involved in DNA replication and repair in fission yeast[36]. In this regard, it
would be of interest to determine whether other RNA splicing and processing factors are also

implied in DNA replication. In this study, we show that the RNA-binding protein RBM10 is recruited to active DNA replication forks in a PRIM1-dependent manner to fine-tune the levels of histone acetylation at active DNA replication fork. Given that PRIM1 knockdown alleviates RBM10 association with active replication forks, we propose a model that this association requires RNA molecules, presumably the RNA primers of Okazaki fragments. In this regard, we revealed that the RBM10 ZnF1 domain, known for its RNA-binding capability, mediates its association with replication fork. Interestingly, this ZnF1 domain is found in various proteins including MDM2, NEIL3, ZRANB3, and FUS that have been previously implicated in regulating DNA replication progression and stress response[79–82]. Future investigations will be necessary to determine whether the association of these proteins with DNA replication forks is mediated by the RNA primers of Okazaki fragments. Functionally, we demonstrate that RBM10 counteracts the formation of R-loops in S-phase cells, concurrently facilitating the association of HDAC1 with active replication forks to deacetylate H4K16. These findings align with recent studies that described a role of HDAC1/2 in chromatin remodeling at active replication forks to maintain normal replication fork progression and replication fork stability[64–66]. Interestingly, it was shown that HDAC1/2 decrease the levels of the transcriptionally active H4K16 acetylation at replication forks, thereby reducing the levels of R-loops and mitigating transcription-induced replication stress[66]. Further work will be required to determine the impact of RBM10 depletion on the acetylation levels of other histone and non-histone residues that are implicated in DNA replication regulation. While our data suggest that RBM10 function in DNA replication is distinct from its splicing activity (Fig. 3g), we cannot exclude the possibility that RBM10 may regulate the splicing or expression of genes involved in regulating replication fork progression.

In this study, we performed a genome-wide CRISPR-Cas9 knockout screen employing isogenic LUAD cells carrying RBM10 cancer mutation, a type of nonsense mutation occurring with frequencies ranging from 9% to 25% in LUAD. We identified ~250 RBM10 SL genes, such as WEE1 and Aurora A kinases, which can be therapeutically exploited for eradicating RBM10-deficient LUAD tumors. Indeed, pharmacological inhibition of WEE1 using MK1775 markedly reduces RBM10-deficient cell proliferation and tumor growth. Given that MK1775 is currently in phase II clinical trials for the treatment of several types of tumors, we propose that it can be also used against RBM10-deficient LUAD with immediate clinical applicability. Interestingly, the sensitivity of RBM10-deficient cells to WEE1 inhibition is further exacerbated by the combined treatment with alisertib, an Aurora Kinase A inhibitor. This

synergistic effect between these two inhibitors is in agreement with a previous study showing

that simultaneous inhibition of Aurora A and WEE1 prompts cells to traverse the G2–M

checkpoint in the presence of DNA damage and impaired spindle assembly, culminating in

mitotic catastrophe. Consequently, it leads to a synergic anti-tumor effect in p53-deficient

squamous cell carcinoma of the head and neck[83]. Together, these observations provide a basis

for testing this combination for clinical treatment of RBM10-deficient LUAD patients, thus

increasing therapeutic benefits and minimizing the risk for developing resistance.

Mechanistically, RBM10-WEE1 synthetic lethality is mediated, at least in part, by increased

replication stress leading to premature mitotic entry and cell death. This finding supports the

notion that the sensitivity of RBM10-deficient cells to WEE1 inhibition is dependent on the

emerging role of RBM10 in DNA replication. Our data are therefore in agreement with

previous studies showing synergistic interaction between WEE1 inhibition and replication

stress-inducing agents[84–87].

Intriguingly, a previous report showed that RBM10 deficiency in LUAD is associated with

enriched immune pathways, elevated tumor mutational burden, and increased human leukocyte

antigens (HLA) expression, suggesting that RBM10 deficiency may enhance anti-tumor

immunity in LUAD[88]. In agreement with this, our RBM10 transcriptome in HCC827 and

H1299 cell lines showed that RBM10 depletion leads to increased expression of

proinflammatory cytokines such as IL1R2, TNFRSF1B, BMP2, IL7R and CCL2. Further work

will be vital to determine whether WEE1 inhibition in RBM10-deficient tumors further

amplifies the upregulation of genes associated with immune pathways, which might enhance

the response to immune therapies.

Unbiased genome-wide CRISPR-Cas9 SL screens revolutionized the discovery of novel SL

interactions that can be exploited for developing new cancer therapies[89,90]. Our CRISPR-Cas9

knockout screen revealed ~250 RBM10 SL genes involved in various cellular pathways, such

as RNA splicing and cell cycle. These genetic interactions advance our understanding of the

physiological functions of RBM10 and provide a repertoire of novel targets that can be

harnessed to formulate new strategies for targeting RBM10-deficiency. Interestingly, RBM10

is frequently mutated in various human cancers such as endometrial (8%) and urothelial (5%)

carcinomas[58]. Therefore, our findings might be of clinical interest to other types of human

cancer harboring RBM10 mutations.

Figure 7: Model depicting the novel role of RBM10 in DNA replication and replication stress response. (Top) RBM10 association with active DNA replication forks is dependent on PRIM1, which synthetizes the RNA primer of Okazaki fragments. RBM10 promotes the recruitment of HDAC1 to ongoing and stressed replication forks and ensures the deacetylation of H4K16, thereby limiting R-loop formation and maintaining fork stability. (Bottom) In RBM10-deficient cells, defective HDAC1 recruitment and H4K16 deacetylation at stressed replication forks contributes to R-loop accumulation, fork destabilization, and ssDNA gap formation leading to replication stress. High levels of replication stress render RBM10-deficient tumor cells sensitive to WEE1 kinase inhibition, leading to the accumulation of DNA damage and mitotic catastrophe resulting in tumor cell death. Created with BioRender.com.

Acknowledgments

We thank Avigail Yampolsky and Yarin Attar for helping in validating the RBM10 CRISPR-

Cas9 screen results, Dvir Aran for his help in analyzing RBM10 mutations in cancer, Yossi

Shiloh and Yehuda Assaraf for their insightful comments on the project. We also thank Nadav

Sharon, Assaf Bester, and Juan Valcárcel for generously sharing reagents and plasmids. We

thank Janette Zavin for histological processing. We thank our lab members for critical reading

of the manuscript. We thank bioRENDER (biorender.com) for assisting in drawing the model

and the schematics presented in the figures. Research in the Ayoub lab is supported by grants

from the Israel Science Foundation (2511/19) and Israel Cancer Research Fund (22-110-PG).

Research in the Simon lab is supported by grants from the Israel Science Foundation (1283/21)

and the Binational Science Foundation (2019688 and 2021085). F.E.M. and E.R.A. are

supported by Clore fellowship. A.B. is supported by Joan and Irwin Jacobs fellowship. N.A. is

supported by the Neubauer Family foundation.

Author Contributions

F.E.M. wrote the original draft and performed all the experiments and analysis described in

this manuscript except the ones indicated below. E.R.A. cloned flag-MCM5 expression

plasmid and helped in setting up the CRISPR-Cas9 screen and proofreading the manuscript.

J.K. performed the experiments described in Figure 2a, 2b, 5g and Supplementary Figure 2e,

5c. A.B. performed the experiments described in Supplementary Figure 2d, 3d and helped in

characterizing RBM10 splicing activity. I.S. helped in analyzing the data related to RBM10

role in DNA replication progression and editing the manuscript. N.A. conceived and supervised

the study.

Corresponding Author

Correspondence to Nabieh Ayoub (ayoubn@technion.ac.il).

Competing Interests

The authors declare that they have no competing interest.

Methods

Cell lines and cell culture

HCC827, NCI-H1299, NCI-H1944, and NCI-H1975 cell lines were obtained from ATCC and grown at 37°C and 5% CO2 and cultured in RPMI-1640 medium (Gibco) supplemented with 10% heat-inactivated fetal bovine serum (Gibco), 2mM L-glutamine (Gibco), and 100unit/mL penicillin and 100μg/mL streptomycin (Gibco). HEK293T cell line was obtained from ATCC and grown at 37°C and 5% CO2 and cultured in Dulbecco's Modified Eagle Medium (DMEM) (Gibco) supplemented with 10% heat-inactivated fetal bovine serum (Gibco), 2mM L-glutamine (Gibco), and 100unit/mL penicillin and 100μg/mL streptomycin (Gibco).

Lentiviral transduction

Lentiviral particles were generated by co-transfecting HEK293T cells plated in 10cm plates with 1.64pmol target vector, together with 1.3pmol psPAX2 (Addgene #12260) and 0.72pmol pMD2.G (Addgene #12259) using 3:1 μg PEI to μg DNA ratio. Media containing the viral particles were collected 48 hr post-transfection and filtered with 0.45 μm filters. The indicated cell lines were transduced with the lentiviral particles in the presence of 10μg/ml polybrene (Sigma-Aldrich H9268). At 48 hr post-infection, transduced cells were selected using the appropriate antibiotics.

Generation of Flag-Cas9 expressing cells and RBM10 knockout cell lines

To generate HCC827 cell line constitutively expressing Cas9, cells were transduced with Lenti-Cas9-2A-Blast (Addgene #73310) lentiviral particles followed by selection with 10μg/ml blasticidin (Invivogen #ant-bl) for 72 hr. A single clone stably expressing Flag-Cas9 fusion was selected and validated by western blot and immunofluorescence. To generate isogenic RBM10 knockout in HCC827 cells expressing flag-Cas9 (HCC827-Cas9), guide RNAs (gRNA) were designed and cloned into pSPgRNA plasmid (Addgene #47108) using BbsI restriction enzyme. HCC827-Cas9 cells were transfected with RBM10 gRNA1 (HCC827-Cas9[RBM10-KO] and naïve HCC827[RBM10-KO3]) or gRNA2 (HCC827-Cas9[RBM10-KO2]) followed by clonal isolation. Single clones were screened for RBM10 knockout using western blot and validated by immunofluorescence and sequencing. The sequences of RBM10 gRNAs are included in Supplementary Table 3. To generate RBM10 knockout in naïve HCC827 and NCI-H1299 cells, RBM10 gRNA1 was cloned into pSpCas9(BB)-2A-GFP (PX458) (Addgene #48138) and transfected to cells. At 24 hr after transfection, GFP-positive cells were sorted using BD LSRFortessa™ cell analyzer (BD Biosciences) and plated in 96-well plates at a

dilution of one cell per well. Single clones were screened for RBM10 knockout using western

blot and validated by immunofluorescence and sequencing.

Genome-wide CRISPR-Cas9 synthetic lethal screen

240 million cells of HCC827-Cas9^{WT} and HCC827-Cas9$^{RBM10-KO}$ were transduced in triplicates

with TKOv1 sgRNA library, which contains 91,320 sgRNAs targeting 17,232 protein-coding

genes[52], at MOI=0.3 and an average of 500-fold coverage of the library. 24 hr post-

transduction, cells were selected with 2 μg/mL puromycin for 2 days. Following selection, 80

$\times 10^7$ cells were harvested as day 0 samples (T0) and the remaining cells were plated and

passaged every 5 days. Cells were then collected at day 15 (T15) for isolation of genomic DNA

(gDNA) together with the T0 samples. Genomic DNA extraction was performed with Blood

& Cell Culture DNA Maxi Kit (Qiagen) according to the manufacturer's instructions. gRNA

inserts were amplified using two rounds of PCR reactions. First, a 600bp region of the lentiviral

library vectors containing the sgRNA region was amplified using ordinary PCR with Q5 high-

fidelity DNA polymerase for 25 cycles. A total of 120μg gDNA was used as template for each

sample (60 reactions: 2μg gDNA/50μl reaction). Second, the amplified PCR products were

subjected to another PCR reaction with Illumina next-generation sequencing adaptors for 10

cycles and 200bp product was amplified. The resulting amplicons were pooled and gel purified

using the NucleoSpin Gel and PCR Clean-up (Macherey-Nagel). Samples were sequenced on

a MiSeq platform with ~10 million reads per sample (Illumina). Demultiplexing and mapping

of sequencing reads to the gRNA library was performed using MAGeCK[54]. Fold-change for

each gRNA was calculated in each replicate versus the T0 sample. 3 analysis methods were

used to predict RBM10 synthetic lethal genes: (1) Bioinformatic and Bayesian Analysis of

Gene Essentiality (BAGEL) algorithm[53] was devised to calculate the essentiality score (Bayes

factor – BF) of each gene in parental (WT) and RBM10-KO cells. RBM10 SL genes were

defined as genes with BF>5 (i.e. essential) in RBM10-KO cells and BF<2 (i.e. non-essential)

in WT cells. (2) CRISPR Counts Analysis (CCA)[55] was used to calculate synthetic lethality

score for each gene and genes with score > 0.8 were considered candidate RBM10 SL genes.

(3) MAGeCK was used to calculate log fold-change (LFC) for each gene and genes with LFC<-

1 in RBM10-KO cells with LFC difference > 1 between WT and RBM10-KO were considered

RBM10 SL genes. Pathway enrichment analysis and gene ontology were conducted using

ShinyGO[91]. STRING interaction network was performed using string-db.org.

Conditional short hairpin RNA (shRNA) knockdown

shRNA oligonucleotides directed against the indicated genes were annealed and cloned into

Tet-pLKO-puro lentiviral vector (Addgene #21915) digested with *Eco*RI and *Age*I. Lentiviral

particles were generated as described above and used to transduce parental and RBM10-KO

HCC827-Cas9 cells, followed by selection with 1μg/ml puromycin (Invivogen #ant-pr) for 3

days. Cells were maintained in complete medium in the presence of 0.6μg/ml puromycin. To

induce shRNA-mediated gene expression knockdown, cells were cultured in the presence of

0.2μg/ml doxycycline (DOX) for 48 hr prior to clonogenic survival assays. Gene expression

knockdown was validated by RT-qPCR. The shRNA and qPCR primer sequences used in this

study are included in Supplementary Table 4.

Generation of expression vectors

To generate EGFP-RBM10 expression vectors, RBM10 was amplified from pDest26-RBM10-

wt plasmid[32] (gift from Prof. Juan Valcárcel) and cloned into pEGFP-C1 expression plasmid.

To generate RBM10 deletion mutants, all-round PCR was used to delete the relevant regions.

To constitutively restore RBM10 expression in RBM10-deficient cells, full-length RBM10 and

RBM10 deletion mutants were subcloned to Lenti-Cas9-2A-Blast plasmid followed by

lentivirus generation and transduction. To generate Flag-MCM5 expression vector, MCM5 was

cloned from cDNA into p3X-Flag-CMV10 expression vector. Primer sequences used for

cloning and mutagenesis are included in Supplementary Table 4.

Transfections

Cell transfections with plasmid DNA or siRNA were performed using Polyethylenimine (PEI)

and Lipofectamine RNAiMax, respectively, following the manufacturer's instructions. siRNAs

used in this study are: RBM10 siRNA #1 (ThermoFisher Scientific HSS112075, 5′-

CAAACGCCGAGAGAAGUGCU-3′); RBM10 siRNA #2 (ThermoFisher Scientific

HSS112075, 5'-TTTGCCAAGGGTTCTAAGAG-3'); PRIM1 siRNA (Euphera Biotech

EHU105881); Stealth RNAi negative control (Invitrogen).

RNA isolation, RT-PCR and qPCR

Total RNA was isolated from cells using the TRIzol reagent according to the manufacturer's

instructions (Ambion) and treated with RQ1 DNase (M6101, Promega). 1μg RNA was used

for cDNA synthesis using the qScript cDNA Synthesis Kit (Quanta) with random primers. RT-

PCR for alternative splicing assay of NUMB and EIF4H genes was performed by standard

PCR using Red Load Taq Master (Jena Bioscience) followed by gel electrophoresis in agarose

gel stained with ethidium bromide. Band intensities were quantified using GelDoc software.

Real-time qPCR for measuring mRNA levels was performed using Step-One-Plus real-time

PCR System (Applied Biosystems) and the Fast SYBR Green Master mix (Applied Biosystems) with three technical repeats for each PCR. Data analysis and quantification were performed using StepOne software V2.2 supplied by Applied Biosystems. Primer sequences used for RT-PCR and qPCR are included in Supplementary Table 4.

Small molecule inhibitors and drugs

MK1775 (A5755), Alisertib (MLN8237) (A4110), and hydroxyurea (B2102) were purchased from APExBio. Caffeine was purchased from Sigma-Aldrich (C8961).

Immunoblotting

Whole-cell protein extracts were prepared using hot lysis buffer (1%SDS, 5mM EDTA, 50mM Tris, pH7.5) supplemented with protease inhibitor mixture (Calbiochem). Protein samples were diluted and boiled in 5X protein loading buffer (10% SDS, 500mM DTT, 50%Glycerol, 250mM Tris-HCL and 0.5%bromophenol blue dye) and separated on an SDS-polyacrylamide gel. Proteins were transferred to a PVDF membrane (Millipore) and immunoblotted with the indicated antibodies. The following primary antibodies were used for immunoblotting: RBM10 (Sigma-Aldrich cat.no. HPA034972, 1:10,000), FLAG-tag (Sigma-Aldrich cat.no. F7425, 1:4,000), β-actin (Sigma-Aldrich cat.no. A5441, 1:10,000), α-Tubulin (Santa Cruz Bio cat.no. sc-5286, 1:1,000), pRPA32-S4/S8 (Bethyl cat.no. A300-245A, 1:1,000), pRPA32-S33 (Bethyl cat.no. A300-246A, 1:1,000), RPA32 (Abcam cat.no. ab2175, 1:500), γH2AX (Cell Signaling Technology cat.no. 2577, 1:1000), pChk1-S345 (Cell Signaling Technology cat.no. 2348, 1:1,000), Histone H3 (Abcam cat.no. ab1791, 1:30,000), Rad51 (Santa Cruz Bio cat.no. sc-8349, 1:1,000), PARP1 (Enzo Life Sciences cat. no. ALX-210-895, 1:2,000), HDAC1 (GeneTex cat. no. GTX100513, 1:2,000), PCNA (Santa Cruz Bio cat. no. sc-56, 1:1,000), GAPDH (Abcam cat.no. ab8245, 1:1,000), GFP (Santa Cruz Bio cat.no. sc-9996, 1:1,000). The following secondary antibodies were used: anti-rabbit IgG-HRP (Jackson ImmunoResearch cat.no. 111–035-003, 1:20,000), anti-mouse IgG-HRP (Amersham cat.no. NXA931, 1:10,000). Immunoblot intensity quantification was performed using ImageJ.

Biochemical fractionation

Cells were lysed with Buffer A for 5 min at 4°C. Cell lysates were centrifuged at 1500× g for 5 min at 4°C and the supernatant (cytoplasmic fraction) was removed. Then, the pellet was incubated with Buffer B (3mM EDTA, 0.2mM EGTA, 1mM DTT, PMSF, and protease inhibitor mixture) for 10 min on ice followed by centrifugation at 1700× g for 5 min at 4°C to extract nuclear-soluble fraction. To prepare chromatin-bound fraction, pellet was resuspended with hot lysis buffer (1%SDS, 5mM EDTA, 50mM Tris, pH7.5), boiled for 15 min, and

sonicated with two 15 sec pulses of 35% amplitude, centrifuged at maximum speed for 20 min

at 12°C, and the supernatant was recovered. Fractions were subjected to immunoblot analysis

with the indicated antiabodies. For chromatin-bound fractionation with RNaseA, pellet was

first resuspended in Buffer A containing 5 µg RNase A and incubated for 2 hr at room

temperature followed by extraction with hot lysis buffer.

GFP-trap

HEK293T cells expressing either EGFP only or EGFP-RBM10 plasmids were subjected to

GFP-TRAP assay as previously described[92]. Where indicated, cells were also co-transfected

with Flag-MCM5 plasmid. Briefly, EGFP-RBM10 or EGFP-only expressing cells were lysed

in RIPA buffer (150mM NaCl, 50mM Tris pH 7.4, 1% NP40, 0.5% deoxycholate, 0.1% SDS)

supplemented with protease inhibitor cocktail and then diluted 1:4 with binding buffer (150mM

NaCl, 50mM Tris pH 7.4). 2 milligram of cell lysate was incubated for 4 hr at 4°C with 15µl

GFP-TRAP beads followed by western blot analysis. For GFP-trap followed by mass

spectrometry (MS), HEK293T cells were transfected with EGFP-RBM10, lysed and subjected

to GFP-trap as described above. Beads were washed 5 times in PBSx1 followed by elution and

trypsin digestion. Digested peptides were then analyzed by mass spectrometry (MS) at the

Smoler Proteomics Center in the department of Biology in the Technion. The samples were

analyzed by LC-MS/MS using Q-Exactive plus mass spectrometer (Thermo Scientific),

coupled to Easy nano LC-1000 capillary UHPLC (Thermo Scientific). Protein identification

and intensity analysis was performed using maxquant[93], searching against the human section

of the Uniprot database. RBM10-interacting proteins enrichment analysis and gene ontology

was conducted using ShinyGO[91].

Immunoprecipitation

For endogenous RBM10 immunoprecipitation, H1299 cells were harvested and lysed in RIPA

buffer (150mM NaCl, 50mM Tris pH 7.4, 1% NP40, 0.5% deoxycholate, 0.1% SDS)

supplemented with protease inhibitor cocktail. Immunoprecipitations were performed using

0.5µg IgG or RBM10 antibody and protein A magnetic beads (GenScript #L00273) incubated

with 2mg lysate overnight at 4°C. Immunoprecipitated samples were subjected to immunoblot

analysis using the indicated antibodies.

Immunofluorescence

Cells were seeded onto glass coverslips and grown for 24 hr followed by drug treatments as

indicated in the figure legends. Cells were fixed with 4% (wt/vol) paraformaldehyde for 10

min, permeabilized with 0.5% Triton X-100 in PBS for 10 min, blocked with blocking buffer

(4% (wt/vol) BSA, 0.15% Tween 20 and 0.15% Triton X-100 in PBS) for 1 hr at room

temperature and incubated with the indicated antibodies for 1 hr at 37°C. Excess antibody was

washed three times with wash buffer (0.15% Tween 20 and 0.15% Triton X-100 in PBS × 1),

and cells were stained with secondary antibodies for 1 hr at room temperature in the dark, and

then washed as above. Coverslips were mounted onto glass slides using VECTASHIELD®

Antifade Mounting Medium with DAPI (Vectorlabs). For immunofluorescence staining of

γH2AX, pRPA32-S33, and BrdU, cells were pre-extracted with 0.25% Triton X-100 in PBS

for 15 min on ice prior to fixation. For EdU incorporation, cells were pulsed with 10μM EdU

(5-ethynyl-2-deoxyuridine, Invitrogen A10044) for 30 min at 37°C, 5% CO_2. EdU click

reactions were performed after cell permeabilization by incubating cells with EdU staining

buffer containing 2mM $CuSO_4$, 100mM ascorbic acid and 1μM AlexaFluor 647 azide

(Invitrogen A10277) in PBS for 30 min at room temperature. Images were acquired using

inverted Zeiss LSM-700 or LSM-710 confocal microscopes. Image analysis was performed

using ImageJ software. The following primary antibodies were used for immunofluorescence:

BrdU FITC (BD Biosciences, cat.no. 51-23614 1:1,000), RBM10 (Sigma-Aldrich cat.no.

HPA034972, 1:2,000), pRPA32-S33 (Bethyl cat.no. A300-246A, 1:500), γH2AX (Millipore

cat.no. 05-636, 1:2,500), FLAG-tag (Sigma-Aldrich cat.no. F7425, 1:1,000), pH3(Ser10) (Cell

Signaling Technology cat.no. 9706, 1:1,000), Anti-DNA-RNA Hybrid Antibody, clone S9.6

(Sigma-Aldritch cat.no. MABE1095, 1:1,000). The following secondary antibodies were used:

anti-rabbit IgG Alexa Fluor®488 (Invitrogen cat.no. A-21206, 1:500), anti-mouse IgG Alexa

Fluor®488 (Invitrogen cat.no. A-21202, 1:500), anti-rabbit Alexa Fluor®568 (Invitrogen cat.no.

A10042, 1:500), anti-mouse Alexa Fluor®568 (Invitrogen cat.no. A10037, 1:500), anti-rabbit

Alexa Fluor®647 (Invitrogen cat.no. A-31573, 1:500), anti-mouse IgG Alexa Fluor®647

(Invitrogen cat.no. A32787, 1:500).

Proximity ligation assay (PLA)

Proximity ligation assay (PLA) was performed using the Duolink in situ red starter kit (Sigma-

Aldrich) following the manufacturer's instructions. Rabbit anti-RBM10 (Sigma-Aldrich

HPA034972, 1:2,000) and mouse anti-PCNA (Santa Cruz Bio cat. no. sc-56, 1:250) antibodies

were incubated for 1 hr at 37°C prior to PLA analysis. PLA assay between RBM10 and EdU-

labelled nascent DNA was performed as previously described[94]. Briefly, cells were pulse

labelled with 100μM EdU for 10 min at 37°C, followed by pre-extraction in 0.25% Triton X-

100 in PBS for 10 min on ice and fixation in 4% (wt/vol) paraformaldehyde. Cells were

incubated in EdU click reaction buffer containing 2mM $CuSO_4$, 100mM ascorbic acid and

10μM biotin-azide (Invitrogen B10184) in PBS for 1 hr at room temperature. Cells were

incubated in blocking buffer for 1 hr at room temperature followed by incubation with rabbit

anti-RBM10 (Sigma-Aldrich HPA034972, 1:2000), rabbit anti-HDAC1 (GeneTex

GTX100513, 1:2,000), Rabbit anti-H4K16ac (Abcam ab109463, 1:2,000), rabbit anti-53BP1

(Novus biological NB100-305, 1:1,000) and mouse anti-biotin (Jackson ImmunoResearch

cat.no. 200-002-211, 1:2000) antibodies for 1 hr at 37°C. Next, PLA assay was performed using

the Duolink in situ red starter kit for experiments described in Fig. 3c,d (Sigma-Aldrich) or

NaveniFlex Cell MR Atto647N kit (Navinci) following the manufacturer's instructions.

Images were acquired using inverted Zeiss LSM-700 confocal microscope. Image analysis was

performed using ImageJ software. PLA intensity was calculated as the product of number of

spots and the mean intensity of the spots per nucleus.

Cell synchronization

857   To synchronize cells at G1/S border using double-thymidine block, cells were treated with

858   2 mM thymidine overnight, released for 10 h and treated again with thymidine overnight. Cells

859   were released into growth medium containing nocodazole (120 ng/ml) to enrich mitotic cells.

860   Cell cycle profiles were analyzed by Cytek Aurora© after staining with propidium iodide

861   (Sigma-Aldritch).

Alkaline and neutral comet assays

Alkaline and neutral comet assay was performed using the CometAssay Kit (Trevigen, 4250-

050-K) according to the manufacturer's instructions. Briefly, 1×10^5 cells/mL were mixed with

molten LMAgarose (at 37°C) at a ratio of 1:10 (v/v), and 70 μL of this mixture was transferred

onto a CometSlide. Slides were placed flat at 4°C in the dark for 30 minutes. Cells were then

lysed and subjected to alkaline or neutral electrophoresis with voltage of 1V/cm. DNA was

stained with SYBR green and slides were imaged using ZEISS LSM700 confocal microscope.

Comet tail moment was scored using CometScore 2.0 software.

Short-term growth delay assay

For determining drug sensitivity, cells were seeded in 96-well plates in duplicates at a density

of 2,000-5,000 cells per well. 24 hr post-seeding, drugs were added at the indicated

concentrations. Cell viability was measured 72 hr after drug treatment using the CellTiter 96®

AQueous One Solution Cell Proliferation Assay (Promega) following the manufacturer's

protocol, and absorbance was measured using Epoch Microplate Spectrophotometer (BioTek).

Cell viability was normalized to the viability of untreated cells.

Clonogenic survival assays

For colony survival assay, cells seeded onto 6-well plates (500 cells per well) and left untreated
or treated with the indicated doses of drugs. After 14 days, cells were fixed in methanol and
stained with 0.25% crystal violet. Individual colonies were counted, and relative survival was
calculated by normalizing to the number of colonies in control wells.

Annexin-V apoptosis assay
Apoptosis was assessed by annexin V-FITC kit (Invitrogen) according to the manufacturer's
instructions. Parental (WT) and RBM10-KO cells were treated with the indicated
concentrations of MK1775 for 24 hr prior to analysis. Samples (10,000 cells) were analyzed
using Cytek Aurora[©]. Results were calculated as the percentage of positive annexin V-FITC
cells out of total cells counted.

Endogenous homologous recombination assay
Homologous recombination repair assay was performed using Cas9-mediated knock-in of
green fluorescent mClover into the first exon of the LMNA gene, as previously described[69].
Briefly, cells were plated in 6-well plates and co-transfected with 1.6 μg pX330-LMNA-
gRNA1 plasmid containing Cas9 and gRNA against LMNA exon 1 and 0.4 μg pCR2.1-
CloverLamin plasmid containing HR donor sequence. In addition, 0.4 μg pDsRed-Monomer-
C1 was included per transfection as a transfection control unless otherwise indicated in the
figure legend. 12−16 h post transfection, culture medium was renewed and where indicated,
Caffeine (4 mM) was added. Seventy-two hours post-transfection, cells were collected and
analyzed by Cytek Aurora[©] . HR efficiency is percentage mClover-expressing cells from
DsRed-Monomer positive cells.

DNA combing assays
DNA combing analysis was performed as described[95]. Briefly, unsynchronized cells were pulse
labelled with 25uM Idu for 30 min, followed by 600μM CldU for another 30 min, followed by
1.5 hr growth without any label. For combing analysis with hydroxyurea (HU), cells were pulse
labelled with 25uM Idu for 30 min, followed by incubation with 4mM HU for 2 hr, followed
by pulse labelling 600μM CldU for another 30 min. Cells were harvested into 2% agarose
plugs, incubated for 48 hr in a digestion buffer (1% N Laurouylsarcosine, 0.2% Na
Deoxycholate, 10mM Tris-HCl pH 7.5, 100mM EDTA, and 1mg/ml proteinase K). Next, TE
50X was added overnight, and the plugs were dissolved into 50mM MES pH6 with 100mM
NaCl. The DNA was combed on Genomic vision combing slides, denatured and stained with
the following primary antibodies: Rat anti-BrdU (Abcam #ab6326, 1:200), mouse anti-BrdU
(BD Biosciences #347580, 1:20), and anti-ssDNA antibody (DSHB #AB10805144, 1:20). The

following secondary antibodies were used: Goat anti-mouse Alexa Fluor®647 (Jackson Immunoresearch #115-605-166, 1:400), goat anti-rat Alexa Fluor®594 (ThermoFisher Scientific #A11007, 1:400), and anti-mouse IgG2a Alexa Fluor®488 (ThermoFisher Scientific #A21131, 1:500). The slides were photographed with a fluorescent microscope and analyzed using Olympus CellSens program. For DNA combing analysis with S1 nuclease digestion, cells were treated as indicated and then pulse labelled with 25uM Idu for 30 min, followed by 600μM CldU for 60 min. Subsequently, cells were permeabilized (10 mM PIPES, pH 6.8, 0.1 M NaCl, 0.3 M sucrose, 3 mM MgCl2, EDTA-free Protease Inhibitor Cocktail (Roche)) at room temperature for 10 min, followed by incubation with S1 nuclease (20 U/ml) (Promega M5761) in S1 buffer for 60 min at 37 °C

RNA sequencing

Three biological replicates of RNA samples were purified from parental and RBM10-KO HCC827 and H1299 cells, or HCC827 cells transfected with control or RBM10 siRNA#1. RNA sequencing libraries were prepared using TruSeq mRNA library preparation kit. Sequencing was performed at The Crown Genomics institute of the Nancy and Stephen Grand Israel National Center for Personalized Medicine,Weizmann Institute of Science using a NovaSeq 6000 system with S1 flow cell to obtain 150 bp paired-end reads. Average read depth was 60 million reads per sample. The quality of the raw FASTQ files was assessed using FastQC software (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). For differential gene expression analysis, raw sequencing reads were aligned to GENCODE GRCh38 genome assembly using Salmon package[96] and differential gene analysis was performed in R using the DESeq2 package[97]. Pathway enrichment analysis and gene ontology was conducted using ShinyGO[91].

Mouse xenograft model

Animal studies and protocols were approved by the Committee on the Ethics of Animal Experiments of the Technion, Israel Institute of Technology (IL-069-07-22). Immunocompromised NOD.CB17-Prkdcscid/NcrCrl (NOD SCID) mice were purchased from Envigo. Mouse model of LUAD cell lines subcutaneous xenografts were established by transplanting cells into 6-week-old female NOD SCID mice. Mice were randomly assigned into control and treatment groups. Prior to initiating any experiments, mice were allowed one week acclimation to housing conditions at the Technion Animal Facility. All mice were housed in a strict pathogen-free environment. 2.5×10^6 of the indicated cells were resuspended in 100μl PBS and mixed 1:1 with Matrigel (High Concentration – Corning) and injected

subcutaneously into the right flank of 6 weeks-old female NOD SCID mice. Tumors were

measured using digital calipers. Tumor volumes were calculated using the formula: $0.5 \times length$

$\times width^2$. When tumors reached ~100mm^3, mice were randomly assigned to treatment groups

(n = 6 mice for each group, unless otherwise indicated). Each group received either control

(vehicle), MK1775 (40mg/kg, p.o.), or MK1775 + alisertib (30mg/kg each, p.o.) once daily for

15 days. Mice were euthanized when the tumors of control mice reached ~1500mm^3 and

tumors were excised, weighed, and photographed.

Immunohistochemistry

Tumors excised from mice were fixed in 4% formalin at 4^{o}C overnight, washed three times

with PBS, and immersed in 30% (wt/vol) sucrose solution at 4^{o}C overnight. Then, tumors were

embedded in optimal cutting temperature compound (TissueTek) and frozen at -20^{o}C and.

10μm cryosections were prepared and permeabilized using 0.5% Triton X-100 in PBS followed

by immunofluorescence staining as described above, except primary antibody incubation with

γH2AX antibody (Millipore cat.no. 05-636,1:500) were performed at 4^{o}C overnight.

Data availability

Raw next-generation sequencing data for RBM10 SL CRISPR-Cas9 screen have been

deposited in ArrayExpress and are available through accession number E-MTAB-13626.

RNA-seq sequencing data generated in this study have been deposited in ArrayExpress

and are available through accession number E-MTAB-13617. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE

repository with the dataset identifier PXD040109. All data supporting the findings of this

study are available from the corresponding author upon request.

References

1.   Sung, H. *et al.* Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence
and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* **71**, 209–
249 (2021).
2.   Siegel Mph, R. L. *et al.* Cancer statistics, 2023. *CA Cancer J Clin* **73**, 17–48 (2023).
3.   Zappa, C. & Mousa, S. A. Non-small cell lung cancer: current treatment and future
advances. *Transl Lung Cancer Res* **5**, 288 (2016).
4.   Herbst, R. S., Morgensztern, D. & Boshoff, C. The biology and management of non-
small cell lung cancer. *Nature 2018 553:7689* **553**, 446–454 (2018).
5.   Collisson, E. A. *et al.* Comprehensive molecular profiling of lung adenocarcinoma.
Nature 2014 511:7511 **511**, 543–550 (2014).
6.   Zhang, S. *et al.* RNA binding motif protein 10 suppresses lung cancer progression by
controlling alternative splicing of eukaryotic translation initiation factor 4H.
EBioMedicine **61**, 103067 (2020).

7.   Xie, M., Xu, X. & Fan, Y. KRAS-Mutant Non-Small Cell Lung Cancer: An Emerging
Promisingly Treatable Subgroup. *Front Oncol* **11**, 1535 (2021).
8.   Cooper, A. J., Sequist, L. v. & Lin, J. J. Third-generation EGFR and ALK inhibitors:
mechanisms of resistance and management. *Nature Reviews Clinical Oncology 2022*
19:8 **19**, 499–514 (2022).
9.   Pio, R. & Montuenga, L. M. Alternative splicing in lung cancer. *J Thorac Oncol* **4**,
674–678 (2009).
10.   Bonnal, S. C., López-Oreja, I. & Valcárcel, J. Roles and mechanisms of alternative
splicing in cancer — implications for care. *Nature Reviews Clinical Oncology 2020*
17:8 **17**, 457–474 (2020).
11.   Zhao, J. *et al.* Systematic profiling of alternative splicing signature reveals prognostic
predictor for prostate cancer. *Cancer Sci* **111**, 3020–3031 (2020).
12.   Coomer, A. O., Black, F., Greystoke, A., Munkley, J. & Elliott, D. J. Alternative
splicing in lung cancer. *Biochim Biophys Acta Gene Regul Mech* **1862**, (2019).
13.   de Fraipont, F., Gazzeri, S., Cho, W. C. & Eymin, B. Circular RNAs and RNA splice
variants as biomarkers for prognosis and therapeutic response in the liquid biopsies of
lung cancer patients. *Front Genet* **10**, (2019).
14.   Cherry, S. & Lynch, K. W. Alternative splicing and cancer: insights, opportunities, and
challenges from an expanding view of the transcriptome. *Genes Dev* **34**, 1005–1016
(2020).
15.   Li, Z. *et al.* The RNA-Binding Motif Protein Family in Cancer: Friend or Foe? *Front*
Oncol **11**, (2021).
16.   Stanley, R. F. & Abdel-Wahab, O. Dysregulation and therapeutic targeting of RNA
splicing in cancer. *Nat Cancer* **3**, 536–546 (2022).
17.   Machour, F. E. *et al.* RBM6 splicing factor promotes homologous recombination
repair of double-strand breaks and modulates sensitivity to chemotherapeutic drugs.
Nucleic Acids Res **49**, 11708–11727 (2021).
18.   Inoue, A., Takahashi, K. P., Kimura, M., Watanabe, T. & Morisawa, S. Molecular
cloning of a RNA binding protein, S1-1. *Nucleic Acids Res* **24**, 2990–2997 (1996).
19.   Owczarek-Lipska, M., Markus, F., Bültmann, E., Korenke, G. C. & Neidhardt, J. A
TARP Syndrome Phenotype Is Associated with a Novel Splicing Variant in RBM10.
Genes (Basel) **13**, (2022).
20.   Daicheng, H., Shiwen, X., Jingxuan, Z., Junbo, H. & Bo, W. A Frameshift RBM10
Variant Associated With TARP Syndrome. *Front Genet* **13**, (2022).
21.   Gebauer, F., Schwarzl, T., Valcárcel, J. & Hentze, M. W. RNA-binding proteins in
human genetic disease. *Nat Rev Genet* **22**, 185–198 (2021).
22.   Højland, A. T. *et al.* First reported adult patient with TARP syndrome: A case report.
Am J Med Genet A **176**, 2915 (2018).
23.   Imagawa, E., Konuma, T., Cork, E. E., Diaz, G. A. & Oishi, K. A novel missense
variant in RBM10 can cause a mild form of TARP syndrome with developmental
delay and dysmorphic features. *Clin Genet* **98**, 606–612 (2020).
24.   Gripp, K. W. *et al.* Long-term survival in TARP syndrome and confirmation of
RBM10 as the disease-causing gene. *Am J Med Genet A* **155**, 2516–2520 (2011).
25.   Johnston, J. J. *et al.* Massively parallel sequencing of exons on the X chromosome
identifies RBM10 as the gene that causes a syndromic form of cleft palate. *Am J Hum*
Genet **86**, 743–748 (2010).
26.   Wang, Y. *et al.* Integrative analysis revealed the molecular mechanism underlying
RBM10-mediated splicing regulation. *EMBO Mol Med* **5**, 1431–1442 (2013).
27.   Loiselle, J. J. & Sutherland, L. C. RBM10: Harmful or helpful-many factors to
consider. *J Cell Biochem* **119**, 3809–3818 (2018).

28. Rodor, J., FitzPatrick, D. R., Eyras, E. & Cáceres, J. F. The RNA-binding landscape of
RBM10 and its role in alternative splicing regulation in models of mouse early
development. *RNA Biol* **14**, 45–57 (2017).
29. Dou, X. Q. *et al.* Alternative splicing of VEGFA is regulated by RBM10 in
endometrial cancer. *Kaohsiung J Med Sci* **36**, 13–19 (2020).
30. Inoue, A. *et al.* RBM10 regulates alternative splicing. *FEBS Lett* **588**, 942–947 (2014).
31. Sutherland, L. C. *et al.* Splicing arrays reveal novel RBM10 targets, including SMN2
pre-mRNA. *BMC Mol Biol* **18**, 1–14 (2017).
32. Bechara, E. G., Sebestyén, E., Bernardis, I., Eyras, E. & Valcárcel, J. RBM5, 6, and 10
differentially regulate NUMB alternative splicing to control cancer cell proliferation.
Mol Cell **52**, 720–733 (2013).
33. Yamada, H. *et al.* Src Family Tyrosine Kinase Signaling Regulates FilGAP through
Association with RBM10. *PLoS One* **11**, (2016).
34. Mohan, N., Kumar, V., Kandala, D. T., Kartha, C. C. & Laishram, R. S. A Splicing-
Independent Function of RBM10 Controls Specific 3' UTR Processing to Regulate
Cardiac Hypertrophy. *Cell Rep* **24**, 3539–3553 (2018).
35. Kunimoto, H. *et al.* RBM10 regulates centriole duplication in HepG2 cells by
ectopically assembling PLK4-STIL complexes in the nucleus. *Genes Cells* **25**, 100–
110 (2020).
36. Weigt, M. *et al.* Rbm10 facilitates heterochromatin assembly via the Clr6 HDAC
complex. *Epigenetics Chromatin* **14**, (2021).
37. Cao, Y. *et al.* RNA-binding motif protein 10 represses tumor progression through the
Wnt/β- catenin pathway in lung adenocarcinoma. *Int J Biol Sci* **18**, 124–139 (2022).
38. Inoue, A. RBM10: Structure, functions, and associated diseases. *Gene* **783**, (2021).
39. Hernández, J. *et al.* Tumor suppressor properties of the splicing regulatory factor
RBM10. *RNA Biol* **13**, 466–472 (2016).
40. Atsumi, T. *et al.* Rbm10 regulates inflammation development via alternative splicing
of Dnmt3b. *Int Immunol* **29**, 581–591 (2017).
41. Xiao, W. *et al.* RBM10 regulates human TERT gene splicing and inhibits pancreatic
cancer progression. *Am J Cancer Res* **11**, 157 (2021).
42. Cao, Y. *et al.* RBM10 recruits METTL3 to induce N6-methyladenosine-MALAT1-
dependent modification, inhibiting the invasion and migration of NSCLC. *Life Sci* **315**,
(2023).
43. Jin, X. *et al.* RBM10 inhibits cell proliferation of lung adenocarcinoma via
RAP1/AKT/CREB signalling pathway. *J Cell Mol Med* **23**, 3897–3904 (2019).
44. Jung, J. H. *et al.* RNA-binding motif protein 10 induces apoptosis and suppresses
proliferation by activating p53. *Oncogene* **39**, 1031–1040 (2020).
45. Wang, Y. *et al.* Distinct gene mutation profiles among multiple and single primary
lung adenocarcinoma. *Front Oncol* **12**, 6746 (2022).
46. Vinayanuwattikun, C. *et al.* Elucidating Genomic Characteristics of Lung Cancer
Progression from In Situ to Invasive Adenocarcinoma. *Scientific Reports 2016 6:1* **6**,
1–10 (2016).
47. Chen, H. *et al.* Genomic and immune profiling of pre-invasive lung adenocarcinoma.
Nat Commun (2019) doi:10.1038/s41467-019-13460-3.
48. Seiler, M. *et al.* Somatic Mutational Landscape of Splicing Factor Genes and Their
Functional Consequences across 33 Cancer Types. *Cell Rep* (2018)
doi:10.1016/j.celrep.2018.01.088.
49. Rao, W. *et al.* Frequently mutated genes in predicting the relapse of stage I lung
adenocarcinoma. *Clin Transl Oncol* (2023) doi:10.1007/S12094-023-03074-Z.

50. Zhao, J. *et al.* Functional analysis reveals that RBM10 mutations contribute to lung
adenocarcinoma pathogenesis by deregulating splicing. *Sci Rep* **7**, (2017).
51. Nanjo, S. *et al.* Deficiency of the splicing factor RBM10 limits EGFR inhibitor
response in EGFR-mutant lung cancer. *J Clin Invest* **132**, (2022).
52. Hart, T. *et al.* High-Resolution CRISPR Screens Reveal Fitness Genes and Genotype-
Specific Cancer Liabilities. *Cell* (2015) doi:10.1016/j.cell.2015.11.015.
53. Hart, T. & Moffat, J. BAGEL: a computational framework for identifying essential
genes from pooled library screens. *BMC Bioinformatics* **17**, (2016).
54. Li, W. *et al.* MAGeCK enables robust identification of essential genes from genome-
scale CRISPR/Cas9 knockout screens. *Genome Biol* **15**, 554 (2014).
55. Adam, S. *et al.* The CIP2A–TOPBP1 axis safeguards chromosome stability and is a
synthetic lethal target for BRCA-mutated cancer. *Nature Cancer 2021 2:12* **2**, 1357–
1371 (2021).
56. Kavanaugh, G. *et al.* A whole genome RNAi screen identifies replication stress
response genes. *DNA Repair (Amst)* **35**, 55–62 (2015).
57. Matsuoka, S. *et al.* ATM and ATR substrate analysis reveals extensive protein
networks responsive to DNA damage. *Science (1979)* (2007)
doi:10.1126/science.1140321.
58. Weinstein, J. N. *et al.* The Cancer Genome Atlas Pan-Cancer analysis project. *Nat
Genet* **45**, 1113–1120 (2013).
59. Nguyen, C. D. *et al.* Characterization of a Family of RanBP2-Type Zinc Fingers that
Can Recognize Single-Stranded RNA. *J Mol Biol* **407**, 273–283 (2011).
60. Collins, K. M. *et al.* An RRM–ZnF RNA recognition module targets RBM10 to exonic
sequences to promote exon exclusion. *Nucleic Acids Res* **45**, 6761–6774 (2017).
61. Leriche, M. *et al.* 53BP1 interacts with the RNA primer from Okazaki fragments to
support their processing during unperturbed DNA replication. *Cell Rep* **42**, 113412
(2023).
62. Agudelo Garcia, P. A. *et al.* Histone acetyltransferase 1 is required for DNA
replication fork function and stability. *Journal of Biological Chemistry* **295**, 8363–
8373 (2020).
63. Kehrli, K. *et al.* Class i histone deacetylase HDAC1 and WRN RECQ helicase
contribute additively to protect replication forks upon hydroxyurea-induced arrest.
Journal of Biological Chemistry **291**, 24487–24503 (2016).
64. Bhaskara, S. *et al.* Histone deacetylases 1 and 2 maintain S-phase chromatin and DNA
replication fork progression. *Epigenetics Chromatin* **6**, 1–22 (2013).
65. Gaggioli, V. *et al.* Dynamic de novo heterochromatin assembly and disassembly at
replication forks ensures fork stability. *Nature Cell Biology 2023 25:7* **25**, 1017–1032
(2023).
66. de Vivo, A. *et al.* OTUD5 limits replication fork instability by organizing chromatin
remodelers. *Nucleic Acids Res* **51**, 10467–10483 (2023).
67. Ait Saada, A., Lambert, S. A. E. & Carr, A. M. Preserving replication fork integrity
and competence via the homologous recombination pathway. *DNA Repair (Amst)* **71**,
135–147 (2018).
68. Chakraborty, S., Schirmeisen, K. & Lambert, S. A. The multifaceted functions of
homologous recombination in dealing with replication-associated DNA damages. *DNA
Repair (Amst)* **129**, 103548 (2023).
69. Pinder, J., Salsman, J. & Dellaire, G. Nuclear domain 'knock-in' screen for the
evaluation and identification of small molecule enhancers of CRISPR-based genome
editing. *Nucleic Acids Res* **43**, 9379–9392 (2015).

70. Ghelli Luserna Di Rorà, A., Cerchione, C., Martinelli, G. & Simonetti, G. A WEE1
family business: regulation of mitosis, cancer progression, and therapeutic target. *J*
Hematol Oncol **13**, (2020).
71. Elbæk, C. R., Petrosius, V. & Sørensen, C. S. WEE1 kinase limits CDK activities to
safeguard DNA replication and mitotic entry. *Mutation Research/Fundamental and*
Molecular Mechanisms of Mutagenesis **819–820**, 111694 (2020).
72. Elbæk, C. R. *et al.* WEE1 kinase protects the stability of stalled DNA replication forks
by limiting CDK2 activity. *Cell Rep* **38**, (2022).
73. Bukhari, A. B., Chan, G. K. & Gamper, A. M. Targeting the DNA Damage Response
for Cancer Therapy by Inhibiting the Kinase Wee1. *Front Oncol* **12**, 347 (2022).
74. da Costa, A. A. B. A., Chowdhury, D., Shapiro, G. I., D'Andrea, A. D. &
Konstantinopoulos, P. A. Targeting replication stress in cancer therapy. *Nat Rev Drug*
Discov **22**, 38–58 (2023).
75. Hirai, H. *et al.* Small-molecule inhibition of Wee1 kinase by MK-1775 selectively
sensitizes p53-deficient tumor cells to DNA-damaging agents. *Mol Cancer Ther* **8**,
2992–3000 (2009).
76. Cong, K. *et al.* Replication gaps are a key determinant of PARP inhibitor synthetic
lethality with BRCA deficiency. *Mol Cell* **81**, 3128-3144.e7 (2021).
77. Kong, A. & Mehanna, H. WEE1 Inhibitor: Clinical Development. *Curr Oncol Rep* **23**,
(2021).
78. Coulombe, P. *et al.* The ORC ubiquitin ligase OBI1 promotes DNA replication origin
firing. *Nature Communications 2019 10:1* **10**, 1–14 (2019).
79. Giansanti, C. *et al.* MDM2 binds and ubiquitinates PARP1 to enhance DNA
replication fork progression. *Cell Rep* **39**, 110879 (2022).
80. Aliyaskarova, U. *et al.* NEIL3-mediated proteasomal degradation facilitates the repair
of cisplatin-induced DNA damage in human cells. *Sci Rep* **13**, (2023).
81. Vujanovic, M. *et al.* Replication Fork Slowing and Reversal upon DNA Damage
Require PCNA Polyubiquitination and ZRANB3 DNA Translocase Activity. *Mol Cell*
67, 882-890.e5 (2017).
82. Jia, W. *et al.* Fused in sarcoma regulates DNA replication timing and kinetics. *J Biol*
Chem **297**, (2021).
83. Lee, J. W. *et al.* Combined Aurora Kinase A (AURKA) and WEE1 Inhibition
Demonstrates Synergistic Antitumor Effect in Squamous Cell Carcinoma of the Head
and Neck. *Clin Cancer Res* **25**, 3430–3442 (2019).
84. Zheng, H., Shao, F., Martin, S., Xu, X. & Deng, C. X. WEE1 inhibition targets cell
cycle checkpoints for triple negative breast cancers to overcome cisplatin resistance.
Sci Rep **7**, (2017).
85. Chen, D. *et al.* Wee1 Inhibitor AZD1775 Combined with Cisplatin Potentiates
Anticancer Activity against Gastric Cancer by Increasing DNA Damage and Cell
Apoptosis. *Biomed Res Int* **2018**, (2018).
86. Aarts, M. *et al.* Forced mitotic entry of S-phase cells as a therapeutic strategy induced
by inhibition of WEE1. *Cancer Discov* **2**, 524–539 (2012).
87. Rajeshkumar, N. V. *et al.* MK-1775, a potent Wee1 inhibitor, synergizes with
gemcitabine to achieve tumor regressions, selectively in p53-deficient pancreatic
cancer xenografts. *Clin Cancer Res* **17**, 2799–2806 (2011).
88. Liu, B. *et al.* RBM10 Deficiency Is Associated With Increased Immune Activity in
Lung Adenocarcinoma. *Front Oncol* **11**, 677826 (2021).
89. Awwad, S. W., Serrano-Benitez, A., Thomas, J. C., Gupta, V. & Jackson, S. P.
Revolutionizing DNA repair research and cancer therapy with CRISPR-Cas screens.
Nat Rev Mol Cell Biol (2023) doi:10.1038/S41580-022-00571-X.

1179 90. Gallo, D. *et al.* CCNE1 amplification is synthetic lethal with PKMYT1 kinase
1180   inhibition. *Nature 2022 604:7907* **604**, 749–756 (2022).
1181 91. Ge, S. X., Jung, D., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool
1182   for animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).
1183 92. Khoury-Haddad, H. *et al.* PARP1-dependent recruitment of KDM4D histone
1184   demethylase to DNA damage sites promotes double-strand break repair. *Proc Natl*
1185   *Acad Sci U S A* **111**, (2014).
1186 93. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates,
1187   individualized p.p.b.-range mass accuracies and proteome-wide protein quantification.
1188   *Nature Biotechnology 2008 26:12* **26**, 1367–1372 (2008).
1189 94. Roy, S., Luzwick, J. W. & Schlacher, K. SIRF: Quantitative in situ analysis of protein
1190   interactions at DNA replication forks. *Journal of Cell Biology* **217**, 1521–1536 (2018).
1191 95. Kaykov, A., Taillefumier, T., Bensimon, A. & Nurse, P. Molecular Combing of Single
1192   DNA Molecules on the 10 Megabase Scale. *Scientific Reports 2016 6:1* **6**, 1–9 (2016).
1193 96. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides
1194   fast and bias-aware quantification of transcript expression. *Nature Methods 2017 14:4*
1195   **14**, 417–419 (2017).
1196 97. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
1197   dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 1–21 (2014).
1198
1199