Geographic variation of the *for* gene reveals signatures of local adaptation in *Drosophila melanogaster*

Dylan J. Padilla Perez

School of Life Sciences, Arizona State University, Tempe, Arizona 85287, USA.

Corresponding author: dpadil10@asu.edu; **Phone:** +1 (480) 646-7769.

Running title: Allelic variation at the *for* gene.

This manuscript includes:

Abstract: 220 words

Main text: $\approx$ 4,000 words

Figures: 6

References: 61

Abstract

In the early 1980s, the observation that *Drosophila melanogaster* larvae differed in their foraging behavior laid the foundation for the work that would later lead to the discovery of the foraging gene (*for*) and its associated foraging phenotypes, rover and sitter. Since then, the molecular characterization of the *for* gene and our understanding of the mechanisms that maintain its allelic variants in the laboratory have progressed enormously. However, the significance and dynamics of such variation are yet to be investigated in nature. With the advent of next-generation sequencing, it is now possible to identify loci underlying adaptation of populations in response to environmental variation. Here, I present results of the first genotype-environment association analysis that quantifies variation at the *for* gene among samples of *D. melanogaster* structured across space and time. These samples consist of published genomes of adult flies collected worldwide, and at least twice per site of collection (during spring and fall). An analysis of genetic differentiation based on F_{st} and individual admixture estimates revealed an east-west gradient in population structure. The results indicate that different patterns of gene flow as expected under models of isolation by environment and isolation by distance are likely driving the genetic differentiation among populations. Overall, this study is essential for understanding the mechanisms underlying the evolution of foraging behavior in *D. melanogaster*.

Keywords: foraging mode, landscape genetics, polymorphism, heterogeneous environments, path length.

Introduction

Species distributed over broad geographic ranges face drastic variation in environmental conditions. Such species provide an opportunity to understand the mechanisms underlying evolutionary responses to environmental variation. *Drosophila melanogaster* represents a good example of a species with a widespread distribution range, being found on every continent and most islands (Markow and O'Grady, 2005). Since the early 1900s until the present days, this species has been a central model to understand the genetic basis of adaptation (Flatt, 2020). Identifying genes underlying adaptation of populations is a key issue in evolutionary biology (Tenaillon and Tiffin, 2008). For instance, the foraging gene (*for*) is best known for underlying different foraging strategies adopted by *D. melanogaster* in response to environmental variation (Anreiter and Sokolowski, 2019). Detailed characterization of this gene in laboratory experiments revealed the existence of two allelic variants: rover and sitter (Sokolowski, 1980). In the presence of nutritive yeast, larval rovers travel longer distances and eat less than sitters do. This difference is not evident in the presence of a non-nutritive agar medium (Kaun et al., 2007), suggesting a gene-by-environment interaction (Burns et al., 2012). Similarly, Pereira and Sokolowski (1993) showed that adult rovers walked significantly farther from the food source after eating than did sitters, but there were no differences in walking behavior in the absence of food. This behavioral variation should also be observed in the wild, because food availability varies temporally and spatially across the range of *D. melanogaster*. Because these behavioral differences between rovers and

sitters arise from genetic variation at the *for* gene, one should expect the frequency of alleles of this gene to vary over space and time.

Extensive geographic and temporal variation in various genetic markers has been identified in *D. melanogaster* (Hoffmann and Weeks, 2007; Kapun et al., 2020; Sezgin et al., 2004). In some cases, clinal genetic variants have been directly linked to phenotypic variants (Lee et al., 2013; Paaby et al., 2014, 2010; Schmidt et al., 2008). For example, associations between the *Adh* allozyme locus and development, as well as the *hsp70* gene and heat knockdown resistance have been discovered from a mid-latitude population along the east coast of Australia. There is also good evidence indicating that *D. melanogaster* evolved Bergmann's clines on more than one continent. When raised in a common environment, both South American and Australian flies from high latitudes developed faster and matured at larger sizes than their counterparts from low latitudes (James et al., 1997). Similarly, seasonal stresses at high latitudes, such as overwinter survival and ephemeral food resources, impose strong selection for increased body size and somatic maintenance, whereas warm climates with ample feeding opportunities favor reduced size and decreased maintenance (James et al., 1997; Paaby and Schmidt, 2009). Because the foraging behavior of organisms influences their ability to gather and assimilate food to fuel their somatic maintenance, one may expect the allele frequency of the *for* gene to differ in genotypes living at high latitudes and over stressful time (e.g., winter), relative to those living at lower latitudes and less stressful conditions.

Although recent studies have provided good insights on the maintenance of the *for*

gene polymorphism in the laboratory, the significance and dynamics of this variation are yet to be investigated in nature. Studies based on experimental evolution suggest that density-dependent selection during the larval stage of Drosophila, rather than the adult stage, can maintain genetic variation in foraging behavior. For instance, a high larval density favors the rover variant and low larval density favors the sitter variant (Sokolowski et al., 1997). Similarly, negative frequency-dependent selection promotes the increase in frequency of one variant when it is rare relative to the other variant (Fitzpatrick et al., 2007). In addition, Anreiter et al. (2017) showed evidence of an epigenetic regulator of the *for* gene (the *G9a* methyltransferase) responsible for rover-sitter differences in adult foraging behavior. Specifically, they showed that rovers bearing *G9a* alleles, had significantly greater foraging success than sitters, but this rover–sitter difference disappeared when they carried *G9a* null alleles. In sum, researchers have now detailed advances on the mechanisms that produce variation in the Drosophila *for* gene under controlled conditions. Yet, our understanding of the environmental drivers of the genetic variation that influences the foraging behavior of flies in nature remains unknown. A good approach to fill this gap is referred to as genotype-environment association analysis (Joost et al., 2007; Rellstab et al., 2015), an emerging field that integrates tools from landscape genetics and population genomics to identify the environmental factors that have shaped present-day genetic variation, and the gene variants that drive local adaptation.

Here, I present results of the first genotype-environment association analysis that quantifies variation at the *for* gene among samples of *D. melanogaster* structured across space and time. These samples consist of published genomes of adult flies col-

lected worldwide, and at least twice per site of collection (during spring and fall). Thus, not only could I test whether the allele frequency of the *for* gene change geographically, but also in response to seasonality. I found a distinct genetic structure among the samples collected in America and Europe. An analysis of genetic differentiation based on F_{st} estimates revealed two major genetic clusters; one with all the American samples grouped together, and the other one grouping the European samples. Similarly, there is some degree of genetic differentiation among samples within each continent, suggesting a total of 9 ancestral populations (k) with unique patterns of allelic variation at the *for* gene. Although seasonality contributed to the genetic differentiation of populations, its effect depends on the site of collection. The results also indicated that different patterns of gene flow as expected under models of isolation by environment (IBE) and isolation by distance (IBD) are likely driving the genetic differentiation of the *for* gene among populations of *D. melanogaster*. Overall, this study is essential for understanding the mechanisms underlying the evolution of foraging behavior in *D. melanogaster*.

Materials and Methods

Data source

The data used in this study represent a subset of a dataset assembled by the Drosophila Genome Nexus project (DGN), the European Drosophila Population Genomics Consortium (DrosEU), and the Real Time Evolution Consortium (DrosRTEC). This dataset is known as the Drosophila Evolution over Space and Time (DEST), and it is coupled

with environmental metadata (see `https://github.com/DEST-bio/DEST_freeze1/`
`blob/main/populationInfo/samps_10Nov2020.csv`). It constitutes the most com-
prehensive spatiotemporal sampling of *D. melanogaster* populations to date, including
estimates of genome-wide allele frequencies for more than 270 populations. Collec-
tively, the data are based on sequencing of more than 13,000 flies collected around
the world over multiple years (Kapun et al., 2021). Importantly, the analyses of
this study relied on pooled samples stored in the Genomic Data Structure (GDS)
file named "dest.all.PoolSNP.001.50.10Nov2020.ann.gds", available at `https://dest.`
`bio/data-files/SNP-tables`. Pooled samples from this file consist of 33-40 wild-
caught males, which are more easily distinguishable morphologically from similar
species than females are. Detailed information on quality control, mapping, SNP
calling and dataset merging can be obtained from the DEST dataset mapping pipeline
(see `https://dest.bio/sourcecode`).

I filtered the GDS object to restrict my focus on Single Nucleotide Polymorphism
variants (SNPs) at the *for* gene. As annotated at `https://flybase.org/` (genome
assembly version 6.32), the *for* gene is found on the left arm of chromosome 2 from
base pairs 3,622,074-3,656,953. I also included 5,000 bp to both sides of this region
as potential regulatory elements. I further filtered the reads to get a subset of pools
collected at least once during spring and winter across multiple years, ranging from
2014 to 2019 (see Table S1 for the exact number of season per locality). Although the
data enabled me to assess the effect of seasonality on allelic variation at the *for* gene,
I failed to implement a temporal analysis (i.e., testing for an effect of year), because
both seasons could not always be sampled in the same year. Because the filtering pro-

cess drastically reduced the number of pools in a few populations, I grouped together

pools from the nearest neighboring localities such that the smallest populations had 5

pools. The resulting number of pools are distributed among 15 populations collected

in Austria EU ($n = 6$), Charlottesville USA ($n = 6$), Denmark EU ($n = 5$), Esparto

USA ($n = 5$), Finland EU ($n = 6$), France EU ($n = 10$), Germany EU ($n = 12$),

Michigan-Wisconsin USA ($n = 9$), New York-Massachusetts USA ($n = 6$), Pennsylva-

nia USA ($n = 18$), Russia EU ($n = 6$), Spain EU ($n = 7$), Tuolumne USA ($n = 5$),

Turkey EU ($n = 13$), and Ukraine EU ($n = 38$). The DEST dataset provides the

coordinates of each collection site, enabling one to obtain data about the seasonal-

ity of temperature and precipitation. These climatic variables were derived from the

monthly values of temperature and rainfall for global land areas at a resolution of 30

s of a longitude/latitude degree spatial resolution (this is about 900 m at the equa-

tor) available at WorldClim v.2 (Fick and Hijmans, 2017). In addition, I extracted

log-transformed values of net primary production (NPP)—the net amount of solar

energy converted to plant organic matter through photosynthesis—measured in units

of elemental carbon. This variable represents the primary source of trophic energy for

the world's ecosystems and can be obtained from the NASA Socioeconomic Data and

Applications Center (sedac) (Imhoff et al., 2004). These environmental variables were

used in a genotype-environment association analysis as described below.

Quality control

To avoid the influence of atypical observations driven by samples with low coverage,

I applied a minimum effective coverage N_{eff} filter of 28 (Nunez et al., 2022). I also

removed pools and SNP loci with more than 20% missing genotypes. To do so, I used the *missingno* function from the "poppr" package of R v2.9.3 Kamvar et al. (2014). Subsequently, I tested for deviations from Hardy–Weinberg equilibrium (HWE) using the function *hw.test* from the "pegas" package of R v1.1 Paradis (2010). This function calculated p-values of an exact test with a Monte Carlo permutation of alleles (1,000 replicates). I considered loci to be out of HWE if they deviated significantly in more than 50% of populations. This decision was based on a "false discovery rate" correction of the p-values.

Genetic differentiation and population structure

To test for genetic differentiation among populations at the level of the *for* gene, I estimated pairwise values of F_{st} adopting the method suggested by Weir and Cockerham (1984). The F_{st} values serve as a tool for describing the partitioning of genetic diversity within and among populations (Wright, 1931). For example, large populations among which there is much migration tend to show little differentiation with values of F_{st} close to 0, whereas small populations among which there is little migration tend to be highly differentiated with values of F_{st} close to 1. I used the function *gl.fst.pop* from the package "dartR" v2.7.2 (Gruber et al., 2018), running 1,000 bootstraps across all loci to generate 95% confidence intervals of the values of F_{st}. When the confidence intervals did not include zero, the F_{st} value in question was considered significantly different from zero.

I estimated population structure based on admixture proportions. Specifically, the admixture analysis uses a sparse negative matrix factorization method (SNMF), avail-

able in the R package "LEA" v3.2.0 (Frichot and François, 2015). Similar to Bayesian clustering programs such as CLUSTER, the function *snmf* in LEA estimates individual admixture coefficients from a matrix of allele frequencies. Assuming k ancestral populations, the function provides least-squares estimates of ancestry proportions and estimates an entropy criterion to evaluate the quality of fit of the statistical model through cross-validation. The entropy criterion enabled one to choose the number of ancestral populations that best explained the data. Specifically, I used the function *snmf* in LEA to compute individual ancestry coefficients for a number of ancestral populations (k), ranging between $k = 1$ and $k = 10$. For each value of k, I ran the algorithm 50 times and calculated the cross-entropy for each run.

Seasonality analysis

To assess the effect of seasonality on allele frequencies at the *for* gene among populations, I used a constrained ordination analysis, which enables one to determine the genetic variation that can be explained by certain environmental variables or constraints. Accordingly, I performed a Constrained Correspondence Analysis (cca) through the function *cca* in the R package "vegan" v2.6-4. This function requires an independent variable (season), and a dependent variable (allele frequencies matrix). To determine whether seasonality had a significant effect on the allele frequencies across populations, I used a permutation test using the mock *anova* function. The output of this function decomposes the total variation into constrained and unconstrained components. When the constrained variation is lower than the observed environmental variation, one says that the constraint is significant.

Detecting adaptation

To detect signature of local adaptation, I performed a genotype-environment association analysis based on a Latent Factor Mixed Model (LFMM) with a ridge penalty (Jumentier et al., 2022). The LFMM method is a univariate test that requires a matrix of allele frequencies as a dependent variable and an independent variable. To define the independent variable, I performed a Principal Component Analysis (PCA) on the environmental variables described earlier in the "Data source" section, using the function *rda* from the R package "vegan" v2.6-4. This function computes the scores of a PCA and returns a scree plot for the eigenvalues of the sample covariance matrix. I identified the number of significant components for the environmental matrix based on a Broken Stick model. The Broken Stick model is one proposed method for estimating the number of statistically significant principal components. The idea is to model N variances by taking a stick of unit length and breaking it into N pieces by randomly selecting break points from a uniform distribution (Jackson, 1993). According to the broken-stick stopping rule, only PC1 was retained as the observed eigenvalue was higher than the corresponding random broken stick component. Thus, I used PC1 as a synthetic predictor for the model. The LFMM method also requires an estimate of the number of populations in the data (k), which relied on the admixture analysis discussed earlier. I ran the model using the default genomic inflation factor correction (GIF) and a False Discovery Rate (FDR) of 0.05 by converting the p-values to q-values. The FDR is the expected proportion of false positives among the list of positive tests (Storey and Tibshirani, 2003). For example, an FDR threshold of 0.05 applied to a dataset with an ideal p-value distribution would produce 5% false positives (false

discoveries) among the set of positive tests. However, if the distribution of p-values deviates from the ideal distribution, this same FDR threshold of 0.05 would produce more or fewer false discoveries among the set of positive tests, depending on the skew of the distribution.

Finally, I explored patterns of isolation by environment (IBE) and isolation by distance (IBD) based on the environmental variables that contributed to most of the variation explained by PC1. Accordingly, I first computed distance matrices such as a genetic distance matrix according to (Nei, 1987), and both a primary production distance matrix and a geographic distance (coordinates) matrix based on Euclidean distance. Subsequently, I performed a Partial Mantel test on genetic distance and primary production distance, holding the geographic distance constant. I also ran the same test holding the primary production distance matrix constant to account for the effect of geographic distance. To do this, I used the *partial.mantel* function from the R package "vegan" v2.6-4, which ran a Monte-Carlo test based on 1,000 permutations and assessed the significance of the correlation between the distance matrices. I also evaluated the relationship between these distances with a Mantel correlogram, a method that does not make the assumption of linearity. For this test, I ran the function *mgram* from the R package "ecodist" v2.0.7 (Goslee and Urban, 2007), setting the number of permutations to 1,000 and using the Sturge's rule to determine an appropriate number of distance classes. The function returns a matrix with one row for each distance class and 6 columns; two of which represent estimates of the Mantel *r*, and its associated p-value (see supporting code).

Results

In general, none of the loci nor populations were consistently out of Hardy-Weinberg equilibrium. I filtered out of the analyses only one sample that had more than 20% of missing genotypes, and 62 loci that had more than 20% of missing data. The quality control procedure generated a new dataset of 1,550 loci (SNPs), and 3,100 alleles from 151 pools distributed among 15 populations.

I found an overall F_{st} value of 0.032, but the pairwise differences in F_{st} ranged from 0 to 0.069. Interestingly, the results revealed two major clusters separating the pools collected in America from those collected in Europe (Figure 1). However, significant differentiation was also observed within each continent.

Figure 1.

The genetic admixture analysis suggested a pattern of genetic clusters that approximates that observed from the F_{st} values. The *snmf* run with $k = 9$ best accounted for population structure as it produced the lowest cross-entropy value (Figure 2a). Overall, I observed a distinct east-west gradient in population structure according to the admixture proportions (Figure 2b). This gradient seems to be stronger among the populations collected in America than those collected in Europe, although the structure of pools from Ukraine, Turkey, and Russia stands out (Figure 3).

Figure 2.

Figure 3.

According to the constrained correspondence analysis, there is a significant interaction between season and locality ($df = 33 - 117; Chi - square = 0.067, p = 0.001$). Despite the substantial divergence among populations in response to seasonality, some of the American and European populations seem to be overlapping and cannot be completely separated by the effect of seasonality (Figure 4).

Figure 4.

The principal component analysis of the environmental variables showed that only PC1 explained more than the broken stick components (42.850%), with longitude ($score = -0.592$), latitude ($score = -0.598$), and NPP ($score = 0.525$) contributing to most of the variation. Accordingly, I adopted the PC1 as a synthetic predictor variable to be used in the genotype-environment association analysis. From the latent factor mixed model with $k = 9$, I extracted locus-specific z-scores and p-values (Figure 5a). Using $k = 9$, the default GIF correction, and an FDR threshold of 0.05, I detected 2 candidate SNPs under selection in response to the PC1 environmental predictor (Figure 5b).

Figure 5.

The IBE and IBD analyses based on the partial mantel test showed a clear increase of genetic distance with NPP distance (Mantel statistic $r = 0.361, p - value < 0.001$) and geographic distance (Mantel statistic $r = 0.926, p - value < 0.001$). However, a

smoothed local mean (read line), indicated that the relationship between the genetic

distance and the NPP distance may not be linear (Figure 6b). Importantly, the Mantel

correlograms supported the same pattern; populations are relatively similar at short

distances, while this similarity decreases with distance (Figure 6c-d).

Figure 6.

Discussion

This study shows that populations of *D. melanogaster* are geographically differenti-

ated based on SNP variants at the foraging gene (*for*). This finding was supported by

estimates of the F_{st} index and individual admixture proportions among populations.

Patterns of genetic differentiation among populations might have been shaped by both

neutral processes such as demographic history, and adaptive forces such as environ-

mental heterogeneity. Some evidence indicate that North American populations of

D. melanogaster have been affected by African and European migrations associated

with human cargo transportation and fruit trade around the world, contributing to

genetic admixture worldwide in recent years (Bergland et al., 2016; David and Capy,

1988; Kao et al., 2015; Lachaise et al., 1988). Numerous studies have also revealed

clinal variation in phenotypes, chromosomal arrangements, and genotypes across en-

vironmental gradients for this species, implicating spatially varying selection (Flatt,

2016; Hoffmann and Weeks, 2007). This study suggests that genetic variation at the

for gene seems to be influenced by seasonality, but this effect varies according to the

site of collection. Although an effect of seasonality seems evident among populations,

some of the American and European populations cannot be separated by the effect of seasons. To understand such variation, the effect of natural selection should be disentangled from demographic events (Miller et al., 2020; Sexton et al., 2014; Smith et al., 2020), and other unobserved factors. Those unobserved factors include uneven sampling designs, genome-sequencing biases, relatedness among individuals, gene interactions that affect phenotypic variation, and linkage disequilibrium within haplotypes (François et al., 2016). To deal with these factors, I applied a stringent correction to the p-values across the statistical test performed in genotype-association analysis (Balding, 2006; Pearson and Manolio, 2008; Korte and Farlow, 2013). Displaying the empirical distribution of the significance values (see Figure 5a) is a good way to show that confounding effects were removed from the analysis and that false discoveries could be controlled (Storey and Tibshirani, 2003; McCarthy et al., 2008). Once confounding errors are removed, adjusting for multiple comparisons could then be achieved through the application of FDR control algorithms (Benjamini and Hochberg, 1995; Storey and Tibshirani, 2003).

According to the genotype–environment association analysis, I found two potential loci under selection; one was identified as an intronic variant and the other one corresponds to an upstream gene variant. The genotype–environment association analysis identifies loci that show strong correlations with one or more environmental variables, and corrects for confounding effects when the environment is correlated with population structure. When modeling the effect of a synthetic predictor (PC1) on genetic variation at the *for* gene, I found that spatially varying selection may be one of the drivers of genetic variation among populations of *D. melanogaster*. The observation

that NPP was one the variables that contributed the most to the variation captured by the predictor of the model supports the idea that NPP may be a good measure of food availability in nature, as it constitutes the primary energy source for Earth's ecosystems and complex food webs by supplying food energy to the planet's heterotrophic organisms, such as *D. melanogaster* (Imhoff and Bounoua, 2006). If NPP varies spatially and temporally across the range of this species, genotypes better able to find food might prevail in environments with low NPP. Consistent with this hypothesis, Williams et al. (2004) found that food shortages in nature may often favor increased mobility in individuals of *D. melanogaster* to find food sources. Specifically, the authors showed evidence that flies become highly active when deprived of food. Yet, some organisms can adjust their foraging behavior to meet environmental changes in food availability (Schoener, 1971). For example, northern pikes and white-tailed deers may reduce food intake during winter, when food availability is low (Johnson, 1966; Silver et al., 1969). In the context of the rover-sitter variants, a mechanism known as balancing selection could explain the maintenance of these variants if they are exposed to NPP heterogeneity in space and time. Although this mechanism seems plausible in Drosophila, a recent study suggested that strong balancing selection at the *for* gene beginning recently (i.e., following migration out of Africa) is unlikely (Turner et al., 2015). The effect of NPP as a spatially selective force could not be ruled out though, as the balancing selection test performed by the authors was based on a lineage-specific McDonald-Kreitman model, which does not consider the effect of the environment.

The significant correlations between genetic distance with both NPP and geographic distances indicate that genetic differentiation between populations is consistent

with signatures of IBE and IBD. Accordingly, gene flow may be strong among populations that occur in close proximity and exploit similar environments. Different patterns of gene flow are expected under IBE and IDB models (Sexton et al., 2014). One scenario predicts that with limited dispersal and no selection, genetic drift would cause populations to become more differentiated at greater distances, and that this process should be more pronounced as mean population size decreases (Wright, 1943). Thus, under strict IBD, distance predicts differentiation as a result of dispersal limitation and drift, irrespective of environmental differences. A second scenario predicts gene flow to be strong among similar environments to a greater extent than predicted under IBD. However, adaptation to local environments can disrupt patterns of IBD, and subsequently IBE could arise if maladapted immigrants from different environments are selected against, causing strong barriers to gene flow. A good example of this scenario comes from plants growing on and near mine tailings, where adaptation to different soil types has occurred (Antonovics, 1968). McNeilly and Antonovics (1968) showed that reproductive barriers arose through differences in flower bud development as a result of local adaptation to different soil types. Interestingly, a pattern of isolation by geographical and environmental factors has been reported recently in *D. melanogaster* (Yue et al., 2021). Using microsatellite genotypes, the authors showed an effect of precipitation on patterns of genetic variation across a latitudinal gradient. Similarly, a significant correlation between pairwise F_{st} and geographic distance was discussed in the context of IBD in a study that examined putatively neutral SNPs, mitochondrial haplotypes, as well as inversion and transposable element insertion polymorphisms (Kapun et al., 2020).

Currently, a number of studies have demonstrated that the foraging behavior of *D. melanogaster* can be strongly influenced by the *for* gene, but none of them have explored the environmental drivers of variation at this gene based on a genotype-environment association study. To my knowledge, this study provides the first evidence that naturally-occurring populations of *D. melanogaster* are genetically differentiated at the *for* gene as a result of environmental selective forces. Because the samples used in this study consisted of consensus sequences derived from pooled samples (33-40 flies), identifying foraging strains whiting populations could not be possible (sitters vs rovers). Regardless of this limitation, I showed evidence of an important genetic variation at the *for* gene in this study, but additional evidence is required to support a hypothesis of a differential frequency of these strains in nature that can be attributed to balancing selection. Lastly, field studies could disentangle the contributions of selection and environmental structuring to patterns of gene flow. For example, one can conduct an experimental transplant study to understand how dispersal and habitat-specific selection interact to influence populations occupying heterogeneous environments. Although IBE and strong selection should promote local genetic adaptation, it is still unclear how often this is the case in Drosophila.

Acknowledgements

I thank Drs Michael Angilletta, Jay Taylor, and Joaquin Nunez for providing helpful comments on earlier versions of this manuscript. I also thank the Drosophila Genome Nexus project, the European Drosophila Population Genomics Consortium, and the Real Time Evolution Consortium for providing easy access to the data used in this

405 study.

Data Accessibility Statement

The analyses of this study relied on pooled samples stored in the Genomic Data Struc-

ture file named "dest.all.PoolSNP.001.50.10Nov2020.ann.gds", available at `https://`

`dest.bio/data-files/SNP-tables`. In addition, the environmental metadata can be

found at `https://github.com/DEST-bio/DEST_freeze1/blob/main/populationInfo/`

`samps_10Nov2020.csv`. Lastly, a fully reproducible workflow of the data analyses, in-

cluding R scripts and additional supporting material, is available in the following

repositories: Github `https://dylan-padilla.github.io/for-evo/`, a Dryad link

will be available upon acceptance .

Conflict of interest

416 The author has declared no competing interests.

Author Contributions

418 DP performed the data analyses, wrote, and edited the original draft of this manuscript.

419 The author agreed to be held accountable for the work performed herein.

References

421 Anreiter, I., Kramer, J. M., and Sokolowski, M. B. (2017). Epigenetic mechanisms
422 modulate differences in drosophila foraging behavior. *Proceedings of the National*
423 *Academy of Sciences*, 114(47):12518–12523.

424 Anreiter, I. and Sokolowski, M. B. (2019). The foraging gene and its behavioral effects:
425 Pleiotropy and plasticity. *Annual review of genetics*, 53:373–392.

Antonovics, J. (1968). Evolution in closely adjacent plant populations vi. manifold effects of gene flow. *Heredity*, 23(4):508–524.

Balding, D. J. (2006). A tutorial on statistical methods for population association studies. *Nature reviews genetics*, 7(10):781–791.

Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300.

Bergland, A. O., Tobler, R., González, J., Schmidt, P., and Petrov, D. (2016). Secondary contact and local adaptation contribute to genome-wide patterns of clinal variation in drosophila melanogaster. *Molecular ecology*, 25(5):1157–1174.

Burns, J. G., Svetec, N., Rowe, L., Mery, F., Dolan, M. J., Boyce, W. T., and Sokolowski, M. B. (2012). Gene–environment interplay in *Drosophila melanogaster*: Chronic food deprivation in early life affects adult exploratory and fitness traits. *Proceedings of the National Academy of Sciences*, 109(supplement_2):17239–17244.

David, J. R. and Capy, P. (1988). Genetic variation of drosophila melanogaster natural populations. *Trends in Genetics*, 4(4):106–111.

Fick, S. E. and Hijmans, R. J. (2017). Worldclim 2: new 1-km spatial resolution climate surfaces for global land areas. *International journal of climatology*, 37(12):4302–4315.

Fitzpatrick, M. J., Feder, E., Rowe, L., and Sokolowski, M. B. (2007). Maintaining a behaviour polymorphism by frequency-dependent selection on a single gene. *Nature*, 447(7141):210–212.

Flatt, T. (2016). Genomics of clinal variation in drosophila: disentangling the interactions of selection and demography. *Wiley Online Library*, pages 1023–1026.

Flatt, T. (2020). Life-history evolution and the genetics of fitness components in *Drosophila melanogaster*. *Genetics*, 214(1):3–48.

François, O., Martins, H., Caye, K., and Schoville, S. D. (2016). Controlling false discoveries in genome scans for selection. *Molecular ecology*, 25(2):454–469.

Frichot, E. and François, O. (2015). Lea: An r package for landscape and ecological association studies. *Methods in Ecology and Evolution*, 6(8):925–929.

Goslee, S. C. and Urban, D. L. (2007). The ecodist package for dissimilarity-based analysis of ecological data. *Journal of Statistical Software*, 22:1–19.

Gruber, B., Unmack, P. J., Berry, O. F., and Georges, A. (2018). dartr: An r package to facilitate analysis of snp data generated from reduced representation genome sequencing. *Molecular ecology resources*, 18(3):691–699.

Hoffmann, A. A. and Weeks, A. R. (2007). Climatic selection on genes and traits after a 100 year-old invasion: a critical look at the temperate-tropical clines in *Drosophila melanogaster* from eastern australia. *Genetica*, 129:133–147.

Imhoff, M. L. and Bounoua, L. (2006). Exploring global patterns of net primary production carbon supply and demand using satellite observations and statistical data. *Journal of Geophysical Research: Atmospheres*, 111(D22).

Imhoff, M. L., Bounoua, L., Ricketts, T., Loucks, C., Harriss, R., and Lawrence, W. T. (2004). Global patterns in human consumption of net primary production. *Nature*, 429(6994):870–873.

Jackson, D. A. (1993). Stopping rules in principal components analysis: a comparison of heuristical and statistical approaches. *Ecology*, 74(8):2204–2214.

James, A. C., Azevedo, R. B., and Partridge, L. (1997). Genetic and environmental responses to temperature of *Drosophila melanogaster* from a latitudinal cline. *Genetics*, 146(3):881–890.

Johnson, L. (1966). Experimental determination of food consumption of pike, *Esox lucius*, for growth and maintenance. *Journal of the Fisheries Board of Canada*, 23(10):1495–1505.

Joost, S., Bonin, A., Bruford, M. W., Després, L., Conord, C., Erhardt, G., and Taberlet, P. (2007). A spatial analysis method (sam) to detect candidate loci for selection: towards a landscape genomics approach to adaptation. *Molecular ecology*, 16(18):3955–3969.

Jumentier, B., Caye, K., Heude, B., Lepeule, J., and François, O. (2022). Sparse latent factor regression models for genome-wide and epigenome-wide association studies. *Statistical Applications in Genetics and Molecular Biology*, 21(1).

Kamvar, Z. N., Tabima, J. F., and Grünwald, N. J. (2014). Poppr: an r package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, 2:e281.

Kao, J. Y., Zubair, A., Salomon, M. P., Nuzhdin, S. V., and Campo, D. (2015). Population genomic analysis uncovers african and european admixture in drosophila melanogaster populations from the south-eastern united states and caribbean islands. *Molecular ecology*, 24(7):1499–1509.

Kapun, M., Barrón, M. G., Staubach, F., Obbard, D. J., Wiberg, R. A. W., Vieira, J., Goubert, C., Rota-Stabelli, O., Kankare, M., Bogaerts-Márquez, M., et al. (2020). Genomic analysis of european *Drosophila melanogaster* populations reveals longitudinal structure, continent-wide selection, and previously unknown dna viruses. *Molecular Biology and Evolution*, 37(9):2661–2678.

Kapun, M., Nunez, J. C., Bogaerts-Márquez, M., Murga-Moreno, J., Paris, M., Outten, J., Coronado-Zamora, M., Tern, C., Rota-Stabelli, O., Guerreiro, M. P. G., et al. (2021). Drosophila evolution over space and time (dest): a new population genomics resource. *Molecular biology and evolution*, 38(12):5782–5805.

Kaun, K. R., Riedl, C. A., Chakaborty-Chatterjee, M., Belay, A. T., Douglas, S. J., Gibbs, A. G., and Sokolowski, M. B. (2007). Natural variation in food acquisition mediated via a drosophila cgmp-dependent protein kinase. *Journal of Experimental Biology*, 210(20):3547–3558.

Korte, A. and Farlow, A. (2013). The advantages and limitations of trait analysis with gwas: a review. *Plant methods*, 9(1):1–9.

Lachaise, D., Cariou, M.-L., David, J. R., Lemeunier, F., Tsacas, L., and Ashburner, M. (1988). Historical biogeography of the drosophila melanogaster species subgroup. *Evolutionary biology*, pages 159–225.

Lee, S. F., Eyre-Walker, Y. C., Rane, R. V., Reuter, C., Vinti, G., Rako, L., Partridge, L., and Hoffmann, A. A. (2013). Polymorphism in the neurofibromin gene, nf1, is associated with antagonistic selection on wing size and development time in *Drosophila melanogaster*. *Molecular ecology*, 22(10):2716–2725.

Markow, T. A. and O'Grady, P. (2005). *Drosophila: a guide to species identification and use*. Elsevier.

McCarthy, M. I., Abecasis, G. R., Cardon, L. R., Goldstein, D. B., Little, J., Ioannidis, J. P., and Hirschhorn, J. N. (2008). Genome-wide association studies for complex traits: consensus, uncertainty and challenges. *Nature reviews genetics*, 9(5):356–369.

McNeilly, T. and Antonovics, J. (1968). Evolution in closely adjacent plant populations iv. barriers to gene flow. *Heredity*, 23(2):205–218.

Miller, A. D., Coleman, M. A., Clark, J., Cook, R., Naga, Z., Doblin, M. A., Hoffmann, A. A., Sherman, C. D., and Bellgrove, A. (2020). Local thermal adaptation and limited gene flow constrain future climate responses of a marine ecosystem engineer. *Evolutionary Applications*, 13(5):918–934.

Nei, M. (1987). *Molecular evolutionary genetics*. Columbia university press.

Nunez, J. C., Lenhart, B. A., Bangerter, A., Murray, C. S., Yu, Y., Nystrom, T. L., Tern, C., Erickson, P. A., and Bergland, A. O. (2022). A cosmopolitan inversion drives seasonal adaptation in overwintering drosophila. *bioRxiv*, pages 2022–12.

Paaby, A. B., Bergland, A. O., Behrman, E. L., and Schmidt, P. S. (2014). A highly pleiotropic amino acid polymorphism in the *Drosophila insulin* receptor contributes to life-history adaptation. *Evolution*, 68(12):3395–3409.

Paaby, A. B., Blacket, M. J., Hoffmann, A. A., and Schmidt, P. S. (2010). Identification of a candidate adaptive polymorphism for drosophila life history by parallel independent clines on two continents. *Molecular ecology*, 19(4):760–774.

Paaby, A. B. and Schmidt, P. S. (2009). Dissecting the genetics of longevity in *Drosophila melanogaster*. *Fly*, 3(1):29–38.

Paradis, E. (2010). pegas: an r package for population genetics with an integrated–modular approach. *Bioinformatics*, 26(3):419–420.

Pearson, T. A. and Manolio, T. A. (2008). How to interpret a genome-wide association study. *Jama*, 299(11):1335–1344.

Pereira, H. S. and Sokolowski, M. B. (1993). Mutations in the larval foraging gene affect adult locomotory behavior after feeding in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences*, 90(11):5044–5046.

Rellstab, C., Gugerli, F., Eckert, A. J., Hancock, A. M., and Holderegger, R. (2015). A practical guide to environmental association analysis in landscape genomics. *Molecular Ecology*, 24(17):4348–4370.

Schmidt, P. S., Zhu, C.-T., Das, J., Batavia, M., Yang, L., and Eanes, W. F. (2008). An amino acid polymorphism in the couch potato gene forms the basis for climatic adaptation in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences*, 105(42):16207–16211.

Schoener, T. W. (1971). Theory of feeding strategies. *Annual review of ecology and systematics*, 2(1):369–404.

Sexton, J. P., Hangartner, S. B., and Hoffmann, A. A. (2014). Genetic isolation by environment or distance: which pattern of gene flow is most common? *Evolution*, 68(1):1–15.

Sezgin, E., Duvernell, D. D., Matzkin, L. M., Duan, Y., Zhu, C.-T., Verrelli, B. C., and Eanes, W. F. (2004). Single-locus latitudinal clines and their relationship to temperate adaptation in metabolic genes and derived alleles in *Drosophila melanogaster*. *Genetics*, 168(2):923–931.

Silver, H., Colovos, N., Holter, J., and Hayes, H. (1969). Fasting metabolism of white-tailed deer. *The Journal of Wildlife Management*, pages 490–498.

Smith, A. L., Hodkinson, T. R., Villellas, J., Catford, J. A., Csergő, A. M., Blomberg, S. P., Crone, E. E., Ehrlén, J., Garcia, M. B., Laine, A.-L., et al. (2020). Global gene flow releases invasive plants from environmental constraints on genetic diversity. *Proceedings of the National Academy of Sciences*, 117(8):4218–4227.

Sokolowski, M. B. (1980). Foraging strategies of *Drosophila melanogaster*: a chromosomal analysis. *Behavior genetics*, 10:291–302.

Sokolowski, M. B., Pereira, H. S., and Hughes, K. (1997). Evolution of foraging behavior in drosophila by density-dependent selection. *Proceedings of the national academy of sciences*, 94(14):7373–7377.

Storey, J. D. and Tibshirani, R. (2003). Statistical significance for genomewide studies. *Proceedings of the National Academy of Sciences*, 100(16):9440–9445.

Tenaillon, M. I. and Tiffin, P. L. (2008). The quest for adaptive evolution: a theoretical challenge in a maze of data. *Current opinion in plant biology*, 11(2):110–115.

Turner, T., Giauque, C. C., Schrider, D. R., and Kern, A. D. (2015). Behavioral
variation in *Drosophila melanogaster*: no evidence for common alleles of large-effect
at the foraging gene in a population from north carolina, usa. *USA. bioRxiv*, 4325.

Weir, B. S. and Cockerham, C. C. (1984). Estimating f-statistics for the analysis of
population structure. *evolution*, pages 1358–1370.

Williams, A. E., Rose, M. R., and Bradley, T. J. (2004). The respiratory pattern in
Drosophila melanogaster selected for desiccation resistance is not associated with the
observed evolution of decreased locomotory activity. *Physiological and Biochemical*
Zoology, 77(1):10–17.

Wright, S. (1931). Evolution in mendelian populations. *Genetics*, 16(2):97.

Wright, S. (1943). Isolation by distance. *Genetics*, 28(2):114.

Yue, L., Cao, L.-J., Chen, J.-C., Gong, Y.-J., Lin, Y.-H., Hoffmann, A. A., and Wei,
S.-J. (2021). Low levels of genetic differentiation with isolation by geography and
environment in populations of *Drosophila melanogaster* from across china. *Heredity*,
126(6):942–954.

Figure 1: Heatmap depicting genetic differentiation among populations of *D. melanogaster* based on pairwise F_{st} values. In some cases, samples from adjacent localities were combined to avoid low sample sizes. Abbreviations are as follows: PA = Pennsylvania; MI-WI = Michigan and Wisconsin; NY-MA = New York and Massachusetts.

Figure 2: Population structure analysis based on individual ancestry coefficients for a number of ancestral populations. **(a)** Cross-entropy values for each *snmf* run with *k* ranging between $k = 1$ and $k = 10$. The red arrow indicates the most likely value of *k*. **(b)** Admixture proportion across populations of *D. melanogaster*. Colors indicate genetic clusters.

Figure 3: Mean admixture proportions across populations of *D. melanogaster* surveyed in America and Europe. Colors in the pie charts represent genetic clusters.

Figure 4: Constrained correspondence analysis depicting the spatial distribution of populations based on the effect of season and locality. "Spider" diagrams connect the pools to the population centroid.

Figure 5: Genotype-environment association test based on a latent factor mixed model with ridge penalty. **(a)** Distribution of adjusted p-values using the default genomic inflation factor. **(b)** Manhattan plot of loci (SNPs) potentially affected by the PC1 predictor variable. Loci highlighted in orange were considered to be under selection according to a False Discovery Rate of 0.05.

Figure 6: Patterns of isolation by distance and isolation by environment according to Partial Mantel tests and Correlograms. **(a-b)** Geographic and net primary production distances as a function of genetic distance. Colors represent estimated probability densities and the red line a smoothed local mean. **(c-d)** Mantel correlation at different distance classes as a function of geographic and net primary production distances. Filled dots indicate significant correlations.

 # Supplementary material

Table S1: Number of seasons surveyed across localities. Abbreviations are as follows: Aus: Austria, Charl = Charlottesville, Denm = Denmark, Esp = Esparto, Fin = Finland, Fran = France, Germ = Germay, PA = Pennsylvania; MI-WI = Michigan and Wisconsin; NY-MA = New York and Massachusetts, Tuol = Tuolume, Ukr = Ukraine.

	Aus	Charl	Denm	Esp	Fin	Fran	Germ	MI-WI	NY-MA	PA	Russia	Spain	Tuol	Turkey	Ukr
fall	3	3	4	3	2	4	6	5	3	9	3	4	4	4	14
spring	3	3	1	2	4	6	6	4	3	8	3	3	1	9	24