Dual-wield NTPases: a novel protein family mined from AlphaFold DB

Koya Sakuma[1], Ryotaro Koike[1] & Motonori Ota[1,2]

1. Graduate School of Informatics, Nagoya University
2. Institute for Glyco-core Research, Nagoya University

Email: K.S. (ksakuma@i.nagoya-u.ac.jp), M.O. (mota@i.nagoya-u.ac.jp)

AlphaFold protein structure database (AlphaFold DB) archives a vast number of predicted models. We conducted systematic data mining against AlphaFold DB and discovered an uncharacterized P-loop NTPase family. The structure of the protein family was surprisingly novel, showing an atypical topology for P-loop NTPases, noticeable two-fold symmetry and two pairs of independent putative active sites. Our findings show that structural data mining is a powerful approach to identifying undiscovered protein families.

Introduction

Characterizing protein structures is essential for understanding their function, and structures are typically solved by experimental approaches and deposited in the Protein Data Bank (PDB) (Burley et al. 2022). When the solved protein adopts a novel structure that appeared at the first time, the finding is usually reported by the researchers who determined it. However, more recently, public databases produced by state-of-the-art structure prediction, such as the AlphaFold protein structure database (AlphaFold DB) and ESM metagenomic Atlas (ESM Atlas), are changing this situation (Varadi et al. 2022; Lin et al. 2023). These databases are approximately three orders of magnitude larger than the PDB and contain numerous experimentally unsolved protein structures. Structural models never seen by human-being must be hiddenly archived there since the models were generated automatically by artificial intelligences and deposited without any human curations, providing opportunities for finding novel proteins based only on the structural information *in-silico*.

Dedicated data mining demands a clearly stated working hypothesis. While several groups have pursued intensive model classifications against AlphaFold DB (Bordin et al. 2023; Inigo Barrio Hernandez et al. 2023; Durairaj et al. 2023), this bird's-eye approach could miss unique and intriguing proteins. To find these hidden gems, we defined a very specific database-search question: are there monomeric proteins that contain multiple phosphate-binding loops (P-loops) on a single continuous β-sheet? The P-loop or Walker-A motif is a local functional motif that recognizes phosphate groups and shared among P-loop NTPases, such as ATPases, GTPases and nucleotide kinases (NKs) (Walker et al. 1982; Saraste, Sibbald, and Wittinghofer 1990; Leipe, Koonin, and Aravind 2003; Leipe et al. 2002; Leipe, Koonin, and Aravind 2003). In general, one P-loop resides on a single continuous β-sheet of a three-layered α/β/α sandwich architecture. Our preliminary search against the PDB supported this observation because no structure has multiple P-loops in a β-sheet. However, the possibility that a single β-sheet possesses multiple P-loops should not be excluded. We hypothesized that such experimentally unobserved multiple-P-loop structures exist in AlphaFold DB and can be discovered via systematic data mining.

Results

By computationally scanning more than 0.2 billion entries in AlphaFold DB version 4 (Varadi et al. 2022; Kim, Mirdita, and Steinegger 2023), we extracted 15,977 single-chained structures possessing multiple P-loops. We then analyzed the hydrogen-bond network and extracted 839 structures with multiple P-loops on a single continuous β-sheet (Frishman and Argos 1995). The structures were grouped into 11 clusters based on structural similarity (Van Kempen et al. 2023). As a result, we found an uncharacterized family of P-loop proteins, dual-wield P-loop NTPase (dwNTPase), as the largest cluster with 711 members. All structural models in this cluster were predicted with high confidence scores, i.e., the average predicted Local Distance Difference Test (pLDDT) was 94.27, indicating that the predictions were reliable (Figure supplement 1) (Jumper et al. 2021).

The overall architecture of dwNTPases was highly novel and showed noticeable two-fold symmetry. Figure 1a shows the structure of a representative dwNTPase from *Bacillus thuringiensis* (Bt. UniProt accession No.: A0A1Y0TWD8). Two P-loop domains are tightly packed and surrounded by two long bridging α-helices and two framing α-helices. Each of the domains comprises six β-strands, and twelve strands form a continuous single β-sheet, which reveals a previously unobserved single-sheeted dual-P-loop architecture (Figure 1b). Two β-hairpins from each domain form a pier-like four-stranded β-sheet to connect two domains. Two canonical P-loops independently form two putative ligand binding sites that penetrate through the molecule and resemble tunnels rather than pockets (Figure 1c). A search against the PDB clarified that no similar structures have been reported (Van Kempen et al. 2023; Minami et al. 2018). Similarly, a SwissProt subset of AlphaFold DB contained no similar structures (Van Kempen et al. 2023; Minami et al. 2018), indicating that the dwNTPase family has no reliable annotations manually verified by UniProt curators.

We found that the P-loop domain of dwNTPases was structurally atypical for a P-loop NTPase by searching against the PDB (Figure 2a) (Van Kempen et al. 2023; Minami et al. 2018). A crystal structure of mutual gliding-motility protein MglAa from *Myxococcus xanthus* (PDB ID: 6h35), a bacterial small and monomeric GTPase, was the only known P-loop NTPase that showed relevant structural similarity to the dwNTPase P-loop domain (Galicia et al. 2019). The P-loop domain of dwNTPase has an additional β-strand at the N-terminus (strand 0) compared to the MglAa structure (Figure 2b). Two strands constituting the pier sheet and a successive α-helix are also appended. In contrast, the domain lacks two C-terminal β-strands (strands 6 and 7) and some other surrounding secondary structural elements (SSEs). These unique arrangements of SSEs give rise to the atypical topology that does not resemble other P-loop NTPases (Figure supplement 2 and 3)(Minami et al. 2018; Chandonia et al. 2022). Furthermore, the P-loop domain has a long loop rather than a helix conserved in other P-loop NTPases (Figure supplement 4), which we named the switch loop (Figure 2a). These atypical features of the P-loop domain make it difficult to assign dwNTPase to known classes of P-loop NTPases.

Despite these novel features of dwNTPase, an iterative structure search by Foldseek against the entire AlphaFold DB revealed that 2,219 similar structures were deposited, most of which originated from bacteria in various Firmicutes (Table 1 and Supplement Data Table 1) (Varadi et al. 2022; Van Kempen et al. 2023). Similar searches against ESM Atlas culled by 30% sequence identity found 748 similar structures (Supplement Data Table 2) (Lin et al. 2023). We classified dwNTPase structures into six sub-classes based on the conservation of motifs and domains (Figure supplement 5). The bona fide dwNTPase structure with two P-

loops intact (class 1) was the most abundant, suggesting functional constraints exist to conserve the two active P-loops. A BLAST search against the non-redundant database revealed that dwNTPase had been classified as the PRK06851 family protein in the NCBI conserved domain database (McGinnis and Madden 2004; J. Wang et al. 2023). Thus, we concluded that dwNTPases constitute a conserved protein family among bacteria.

Discussion

   The molecular functions of dwNTPases were investigated by analyzing conserved residues. The most symmetric class of dwNTPases (class 1) possesses two clusters of conserved residues shared between both halves (Figure 3a and Figure supplement 6). We found Cys66/Cys248 (residue numbers follow Bt. dwNTPase), Asp74/Asp256, Asp87/Asp269 and His92/His274 formed putative metal binding sites. Molecular dynamics simulations of the Bt. dwNTPase structure complexed with two ATPs, two Mg^{2+} ions and two Zn^{2+} ions showed that the Zn^{2+} ions were stably coordinated by two aspartates and the γ-phosphate group of ATPs (Figure 3b) (Abraham et al. 2015; Huang et al. 2017), which resembles the active site structure of metal-dependent nucleotidyl-transfer enzymes (Figure 3c) (Yang 2008). The side chains of Cys66/Cys248 and His92/His274 remained unoccupied (Figure 3d), suggesting that they may have roles other than metal-binding. As the pair of cysteine and histidine residues are reminiscent of the catalytic triad/dyad in cysteine proteases (Figure 3e), we hypothesize that dwNTPases have additional hydrolase/ligase activity (Dodson and Wlodawer 1998).

   In addition to these conserved residues, we identified other regions characteristic of dwNTPases. First, each P-loop domain has conserved lysine residues (Lys36/Lys218) that precede the P-loops and interact with two switch loops. Because the switch loop partially conceals the ligand binding tunnels (Figure supplement 7a) and is highly flexible in MD simulations (Figure supplement 7b), the conserved lysine residues may play sensor-like roles to trigger NTPase activity, depending on the binding of other ligands to the tunnels. Additionally, the P-loops are surrounded by several charged or polar residues that support the recognition of NTPs and Mg^{2+} ions (Figure supplement 7c) and are not conserved in known P-loop NTPases (Leipe et al. 2002; Leipe, Koonin, and Aravind 2003).

   The biological functions of dwNTPases remain elusive because their structures show limited homology to NTPases with known functions, indicating that dwNTPases are responsible for unique biological functions. The two-fold symmetry implies that the substrates of dwNTPases also possess two-fold symmetry, such as double-stranded DNA, or that the cleft between two P-loop domains recognizes ligand molecules in a similar manner to periplasmic heme-binding proteins (Figure supplement 8) (Mattle et al. 2010). The left half (residues 1–139 and 321–369) of the structure in Figure 1a is more positively charged than the right half (140–320), indicating that each half plays different functional roles (Figure 1c and Figure supplement 9). Because dwNTPases are distributed among various Firmicutes, especially among Bacillus (Table 1), their biological functions may be related to spore-formation, which is characteristic of the species. The evolutionary origin of dwNTPases is unknown. Although it is plausible that dwNTPases gained two-fold symmetry via gene duplication, the origin of the P-loop domains remains unclear. Detailed phylogenetic analysis may explain the evolution of P-loop NTPases, including dwNTPases (Leipe et al. 2002; Leipe, Koonin, and Aravind 2003). Structural and biochemical studies are

required and should provide greater insight into the biological significance of the dwNTPase
family.
In summary, we demonstrated that structural data mining based on specific working
hypothesis can discover uncharacterized protein families and is a powerful approach to
exploring dark proteomes (Taylor et al. 2009; Perdigão et al. 2015), the unwatched region of
the protein universe, which will help and encourage the design of experimental studies.
144

145
146
Figure 1 | AlphaFold2 predicted model of a dual-wield NTPase structure (AF-
A0A1Y0TWD8-F1-model_v4). a, Overall dwNTPase structure colored according to a
purple-white-orange gradient from the N- to C-terminus. **b,** Topology diagram of
dwNTPase. Blue and red arrows represent β-strands pointing up and down that form the
large β-sheets in the P-loop domains. Green arrows represent the pier sheet. White
rectangles are α-helices. Grey and black lines indicate junctions projecting behind and out
of the β-sheets, respectively. Blue dotted lines represent hydrogen bonds connecting the two
halves of the large β-sheet. **c,** The location and shape of the ligand binding tunnels. Color
bar is at the bottom.

156
Figure 2 | P-loop domain. a, Front and top views of the dwNTPase P-loop domain colored
according to a purple-white-orange gradient from the N- to C-terminus. P-loop, switch loop
and pier-sheet are indicated by labels. **b**, Topology diagrams and cartoon representations of
dwNTPase P-loop domains and MglAa structure. Arrows and rectangles represent β-
strands and α-helices. Secondary structures that align between two structures are colored
blue.
163

Table 1 | Phylogenetic classification of dwNTPases

Domain	Phylum	Class	Count
Bacteria	Firmicutes	Bacilli	877
		Clostridia	705
		Desulfuribacillia	1
		Erysipelotrichia	5
		Negativicutes	13
		Tissierellia	32
		Unclassified Firmicutes	66
	Proteobacteria	Deltaproteobacteria	8
	Chloroflexi	Anaerolineae	1
	Tenericutes	Mollicutes	1
Archaea	Euryarchaeota	Methanomada group	1
Others			17

We performed structural alignment of all 2219 structures against the representative dwNTPase structure. To ensure fragmented structures were excluded, 1843 structures showing TM-scores > 0.85 were selected. Entries with no phylogenetic information available in UniProt were ignored. The structures (1727 in total) were classified by their species. Others include environmental samples, metagenomes, unclassified bacteria and Firmicutes from environmental samples.

173
Figure 3 | Putative functionally relevant residues. a, Conserved residues in the putative
ligand binding tunnels. His, Cys and Asp are colored blue, orange and red, respectively. P-
loops and their conserved residues are colored cyan and grey. **b**, Coordination of metal ions
by two aspartate side chains observed in MD simulations. ATP is shown in orange stick
representation. Side chains of relevant residues are shown as sticks and CPK coloring. Green
and grey spheres represent Mg^{2+} and Zn^{2+} ions, respectively. The P-loop is colored cyan. **c**,
The active site structure of RNase H (PDB ID: 1zbl). The side chain of metal coordinating
amino acid residues Asp and Asn are shown as sticks and CPK coloring, where Asn is a
mutation from Asp. Mg^{2+} ions are shown as spheres. The Mg^{2+} ion coordinating with the
side chain of Asp and Asn is colored green. Nucleic acid residues that contact the Mg^{2+} ion
are shown in orange. **d**, The catalytic triad-like side chain configuration observed during the
MD simulations. The triad-like side chain cluster is circled. The black dotted line indicates
the hydrogen bond between the side chains of His92 and Asp74, which HBPLUS detected.
e, The active site structure of TEV protease (PDB ID: 1lvm). Side chains of the Cys-His-Asp
catalytic triad are shown as sticks, CPK coloring and circled.
189

Materials and methods

Identification of structures containing P-loop-like fragments

AlphaFold DB (v4 UniProt) was downloaded from the Foldcomp database (Varadi et al. 2022; Kim, Mirdita, and Steinegger 2023). P-loop NTPase protein structures were extracted by converting the models into the sequences of ABEGO, where A, B, E and G, respectively, denote backbone dihedral angles (phi, psi) for α, β, left-handed β and left-handed α on the Ramachandran plot (Wintjens, Rooman, and Wodak, n.d.). O denotes other conformations unassignable on the Ramachandran plot, typically a cis-peptide conformation. Typical P-loop (Walker-A) motifs have conformations represented by EBBGAG or BBBGAG, both of which can be seen in the crystal structure of α and β subunits of bovine mitochondrial F1-ATPase (chain A and chain D of PDB ID: 1BMF) (Abrahams et al. 1994). Because the P-loop is a junction between a β-strand and an α-helix, we extended the ABEGO motifs to "BBBEBBGAGAAAAA" or "BBBBBBGAGAAAAA" and extracted all the structures containing any of them by sequence pattern matching. We then calculated the Cα root-mean-square deviations (RMSDs) of the matched substructures against the reference P-loop fragment (residues 166–179 of 1BMF, chain A) and filtered out substructures with RMSDs larger than 2.0 Å. We obtained 15,977 proteins containing multiple P-loop-like fragments and built a custom Foldcomp database for subsequent procedures (Kim, Mirdita, and Steinegger 2023; Steinegger and Söding 2017).

Identification of dual-wield NTPases

Visual inspection revealed that most structures with multiple P-loop-like fragments within a single chain were tandem repeats of known P-loop NTPase domains connected by flexible linkers. Such proteins were excluded by analyzing structures using an in-house program that detects the β-sheet hydrogen-bond network and extracting structures possessing two P-loop-like fragments on a single continuous β-sheet (Frishman and Argos 1995). We obtained 839 structures possessing two P-loop-like fragments on a single β-sheet. These structures were clustered by TM-score calculations with Foldseek into 11 clusters (Van Kempen et al. 2023). The largest cluster contained 711 members, which corresponded to dual-wield NTPases. For these structures, we performed all-against-all structure alignment using MICAN and defined the structure with the largest average TM-score as the representative (AF-A0A1Y0TWD8-F1-model_v4) (Minami et al. 2018).

Extraction of structures similar to dwNTPase from AlphaFold DB and ESM Atlas

We performed iterative structure searches using FoldSeek to enumerate as many structures that resemble dwNTPase as possible (Van Kempen et al. 2023). In the first stage, we performed a structure search against AlphaFold DB using all 711 structures initially mined from AlphaFold DB as queries. After removing overlapping structures, we obtained 1377 structures. Using these structures as seeds, we again performed a Foldseek search and obtained 135 new non-overlapping structures. The third iteration of Foldseek search yielded some non-specific hits. Therefore, we stopped this iteration, manually selected similar structures and discarded the remaining structures. Consequently, we obtained 2219 dwNTPase structures from AlphaFold DB. Similarly, we performed structural searches against the highquality_clust30 subset of ESM-atlas using 711 structures found in AlphaFold

234 DB as queries and obtained 748 structures with a TM-score larger than 0.5 (Lin et al. 2023;
235 Van Kempen et al. 2023; Xu and Zhang 2010).
236
Whole structure search against the PDB and Swiss-Prot subset of AlphaFold DB
To assess the novelty of the dwNTPase structure and gain insights into the function, we
performed structural searches against PDB100 and the Swiss-Prot subset of AlphaFold DB
(version 4) using the Foldseek server in the TM-align mode and the representative structure
as the query (Burley et al. 2022; Varadi et al. 2022; Van Kempen et al. 2023). No relevant
(TM-score $\geqq$ 0.5) hit was found among these databases. We used MICAN to perform
rigorous one-against-all searches without pre-filtering; however, no similar (TM-score $\geqq$
0.5) structures were found among the PDB (2023-09-Jan) and the Swiss-Prot subset of
AlphaFold DB (version 2) (Minami et al. 2018).
246
Domain structure search against the PDB, Swiss-Prot subset of AlphaFold DB and SCOPe
We searched structures similar to the P-loop domain of the representative structure
(residues 1–110) against the PDB100 and the Swiss-Prot subset of AlphaFold DB by using
the Foldseek server (Van Kempen et al. 2023). No relevant hit was found. We used MICAN
to perform a rigorous structure search without pre-filtering against the PDB (2023-09-Jan)
and the Swiss-Prot subset of AlphaFold DB (version 2) (Minami et al. 2018). We obtained
358 and 2931 relevant hits (TM-score $\geqq$ 0.5) from the PDB and Swiss-Prot. We performed
clustering by mmseq2 with sequence identity set at 35% and obtained 15 and 137 clusters
(Steinegger and Söding 2017). The alignments were checked by visual inspection of all
cluster representatives. We found that some structures showed similar topology to the P-
loop domain of dwNTPase: 6h35, Q1DB04 and Q9UBK7 from the PDB and Swiss-Prot,
which are annotated as GTPase or GTP-binding proteins (Galicia et al. 2019). The
remaining hits showed RecA-like topology and were not topologically identical to
dwNTPase because the RecA-like topology has an all-parallel β-sheet, whereas dwNTPases
have anti-parallel-containing β-sheets. Similarly, we performed structural comparisons
against domain structures classified as G-proteins (SCOP concise classification string
(SCCS): c.37.8), NKs (c.37.1) and RecA-like proteins (c.37.11) in the SCOPe version 2.08
using MICAN (Minami et al. 2018; Chandonia et al. 2022). The groups of G-proteins, NKs
and RecA-like proteins contained 255, 212 and 118 parsed domain structures, respectively,
and we selected the structures that showed the highest TM-score in the group for
visualization (Figure supplement 2).
268
Calculation of sequence identities between two halves of dwNTPase structures
270 We selected 1,903 structures with more than 340 residues from the set of dwNTPases
271 extracted from AlphaFold DB. A structure was self-aligned by MICAN in the rewiring
272 mode, which ignores the sequential order of SSEs (Minami et al. 2018). The sequence
273 identity was calculated based on the second-best alignment by MICAN.
274
Identification of putative catalytic residues and a side chain pattern search against the PDB
276 The potential function of dwNTPases was examined by performing a sequence search and
277 alignment to identify conserved residues by HHblits against UniRef30_2022_02 (Remmert
278 et al. 2012; Suzek et al. 2015). After three iterations, 2687 sequences were extracted from the
279 database. To exclude fragmented sequences most likely originating from partial matches to

the P-loop consensus motif, we removed aligned sequences with more than ten gaps against the representative sequence and obtained a Multiple Sequence Alignment (MSA) with 138 sequences. From this MSA, the site-wise entropy of the alignments was calculated to identify conserved residues, and the top 10 residues around the two tunnels were listed. We defined tunnel 1 as residues 61–100 and tunnel 2 as residues 243–282. From tunnel 1, residues 62, 66, 67, 69, 73, 74, 75, 87, 88 and 100 were identified. From tunnel 2, residues 244, 246, 247, 248, 252, 255, 256, 261, 263 and 274 were identified. According to the orientation of side chains toward the tunnel, we selected Cys66, Ser67, Asp74 and Asp87 as candidates for probable functional residues in tunnel 1. Similarly, Cys248, Asp256 and His274 were selected for tunnel 2. Considering the symmetry of the dwNTPase structure, Cys66/Cys248, Asp74/Asp256, Asp87/Asp269 and His92/His274 were considered clusters of functional residues in tunnels 1 and 2. We performed a side chain pattern search against the PDB using the strucmotif-search program to determine whether protein structures possessed similar side chain configurations (Bittrich, Burley, and Rose 2020). The set of residues Cys66, Asp74, Asp87 and His92 in the representative structure were selected as queries, and a search was performed against all structures in the PDB (2022-28-12), with the threshold for the structure similarity set to 1.0 Å. The side chain pattern search gave no hits and indicated that the putative catalytic residues have a novel configuration of conserved residues.

Docking of ATP, Mg and Zn

We transplanted ligand structures from existing PDB structures to model the complex structures. The P-loop region of an ATPase crystal structure (PDB ID: 6j18) was superposed to the P-loop of the representative structure by MICAN in PyMOL, and the ATP and Mg^{2+} models were extracted (Minami et al. 2018; S. Wang et al. 2020; Schrodinger 2015). Similarly, His125 from a zinc finger motif (PDB ID: 2hgh) was superposed to His92 and His274, and the coordinating Zn^{2+} ions were extracted (Lee et al. 2006). The extracted ligand molecules were merged with the representative structure.

MD simulations

MD simulations were performed by Gromacs version 2022.04 with the charmm36 force field (Abraham et al. 2015; Huang et al. 2017). The size of simulation boxes was determined by the molecule size with margins of 13 Å. After *in-vacuo* energy minimization to remove steric clashes, the protein-ligand complex was solvated by the TIP3P water model with 0.1 M NaCl, and the system was neutralized by adding additional Na^+ or Cl^- ions, depending on the total charge of the protein and ligands. The energy was minimized by the steepest descent and equilibrated by 100 ps NVT and NPT simulations with harmonic restraints on the non-hydrogen atoms. The temperature and pressure of the system were controlled to 300 K and 1 bar by the V-rescale thermostat and Parrinello-Rahman barostat. Electrostatic interactions were computed by the particle mesh Ewald method, and bonds involving hydrogen atoms were constrained by the LINCS algorithm. For each docked model, we performed 20 trajectories of 100 ns simulations with a 2 fs time step.

Figure preparation

The images of molecular structures were created by PyMOL and Mol* viewer (Schrodinger 2015; Sehnal et al. 2021). The surface electrostatic potential was calculated by the PyMOL APBS plugin (Jurrus et al. 2018). Hydrogen bonds were detected by HBPLUS and visualized

326 by PyMOL (McDonald and Thornton 1994). Secondary structure elements were assigned
327 by DSSP and illustrated by ESPript (Kabsch and Sander 1983; Robert and Gouet 2014).
328

329 **Author contributions**
330 K.S. designed and initiated the study, performed structure mining, structure searches,
331 structure comparisons, complex modeling, MD simulations, interpreted data and prepared
332 and wrote the manuscript. R.K. and M.O. interpreted data and wrote the manuscript.
333

Acknowledgements
K.S. would like to thank George Chikenji for providing the program to analyze hydrogen-
bonding patterns in β-sheets, Milot Mirdita for providing instructions on building the
custom Foldcomp databases, Naoya Kobayashi and Shintaro Minami for their suggestions
on extracting functionally relevant residues to infer functions from structures, Nobu C.
Shirai for his constructive comments on the manuscript, and Shigeo S. Sugano and Ryo
Ozuka for discussing the feasibility and significance of structure-based protein mining,
which inspired K.S. to design this study. Structure searches by Foldseek and MD simulations
were carried out on the supercomputer "Flow" at the Information Technology Center,
Nagoya University. This study was supported by KAKENHI grant numbers JP21H00394 to
R.K. and 20H05932 to M.O. We thank Edanz (https://jp.edanz.com/ac) for editing a draft of
this manuscript.

References

Abraham, Mark James, Teemu Murtola, Roland Schulz, Szilárd Páll, Jeremy C. Smith, Berk Hess, and Erik Lindahl. 2015. "GROMACS: High Performance Molecular Simulations through Multi-Level Parallelism from Laptops to Supercomputers." *SoftwareX* 1–2 (September): 19–25. https://doi.org/10.1016/j.softx.2015.06.001.

Abrahams, Jan Pieter, Andrew G W Leslie, Rene Lutter, and John E Walker. 1994. "Structure at 2.8 A Resolution of F1·ATPase from Bovine Heart Mitochondria" 370.

Bittrich, Sebastian, Stephen K. Burley, and Alexander S. Rose. 2020. "Real-Time Structural Motif Searching in Proteins Using an Inverted Index Strategy." *PLOS Computational Biology* 16 (12): e1008502. https://doi.org/10.1371/journal.pcbi.1008502.

Bordin, Nicola, Ian Sillitoe, Vamsi Nallapareddy, Clemens Rauer, Su Datt Lam, Vaishali P. Waman, Neeladri Sen, et al. 2023. "AlphaFold2 Reveals Commonalities and Novelties in Protein Structure Space for 21 Model Organisms." *Communications Biology* 6 (1): 160. https://doi.org/10.1038/s42003-023-04488-9.

Burley, Stephen K., Helen M. Berman, Jose M. Duarte, Zukang Feng, Justin W. Flatt, Brian P. Hudson, Robert Lowe, et al. 2022. "Protein Data Bank: A Comprehensive Review of 3D Structure Holdings and Worldwide Utilization by Researchers, Educators, and Students." *Biomolecules* 12 (10): 1425. https://doi.org/10.3390/biom12101425.

Chandonia, John-Marc, Lindsey Guan, Shiangyi Lin, Changhua Yu, Naomi K Fox, and Steven E Brenner. 2022. "SCOPe: Improvements to the Structural Classification of Proteins – Extended Database to Facilitate Variant Interpretation and Machine Learning." *Nucleic Acids Research* 50 (D1): D553–59. https://doi.org/10.1093/nar/gkab1054.

Dodson, Guy, and Alexander Wlodawer. 1998. "Catalytic Triads and Their Relatives." *Trends in Biochemical Sciences* 23 (9): 347–52. https://doi.org/10.1016/S0968-0004(98)01254-7.

Durairaj, Janani, Andrew M. Waterhouse, Toomas Mets, Tetiana Brodiazhenko, Minhal Abdullah, Gabriel Studer, Mehmet Akdel, et al. 2023. "What Is Hidden in the Darkness? Deep-Learning Assisted Large-Scale Protein Family Curation Uncovers Novel Protein Families and Folds." Preprint. Bioinformatics. https://doi.org/10.1101/2023.03.14.532539.

Frishman, Dmitrij, and Patrick Argos. 1995. "Knowledge-Based Protein Secondary Structure Assignment." *Proteins: Structure, Function, and Bioinformatics* 23 (4): 566–79. https://doi.org/10.1002/prot.340230412.

Galicia, Christian, Sébastien Lhospice, Paloma Fernández Varela, Stefano Trapani, Wenhua Zhang, Jorge Navaza, Julien Herrou, Tâm Mignot, and Jacqueline Cherfils. 2019. "MglA Functions as a Three-State GTPase to Control Movement Reversals of Myxococcus Xanthus." *Nature Communications* 10 (1): 5300. https://doi.org/10.1038/s41467-019-13274-3.

Huang, Jing, Sarah Rauscher, Grzegorz Nawrocki, Ting Ran, Michael Feig, Bert L de Groot, Helmut Grubmüller, and Alexander D MacKerell. 2017. "CHARMM36m: An Improved Force Field for Folded and Intrinsically Disordered Proteins." *Nature Methods* 14 (1): 71–73. https://doi.org/10.1038/nmeth.4067.

Inigo Barrio Hernandez, Jingi Yeo, Jürgen Jänes, Tanita Wein, Mihaly Varadi, Sameer
        Velankar, Pedro Beltrao, and Martin Steinegger. 2023. "Clustering Predicted
        Structures at the Scale of the Known Protein Universe." *BioRxiv*, January,
        2023.03.09.531927. https://doi.org/10.1101/2023.03.09.531927.

Jumper, John, Richard Evans, Alexander Pritzel, Tim Green, Michael Figurnov, Olaf
        Ronneberger, Kathryn Tunyasuvunakool, et al. 2021. "Highly Accurate Protein
        Structure Prediction with AlphaFold." *Nature* 596 (7873): 583–89.
        https://doi.org/10.1038/s41586-021-03819-2.

Jurrus, Elizabeth, Dave Engel, Keith Star, Kyle Monson, Juan Brandi, Lisa E. Felberg,
        David H. Brookes, et al. 2018. "Improvements to the APBS Biomolecular Solvation
        Software Suite." *Protein Science* 27 (1): 112–28. https://doi.org/10.1002/pro.3280.

Kabsch, Wolfgang, and Christian Sander. 1983. "Dictionary of Protein Secondary
        Structure: Pattern Recognition of Hydrogen-Bonded and Geometrical Features."
        Biopolymers 22 (12): 2577–2637. https://doi.org/10.1002/bip.360221211.

Kim, Hyunbin, Milot Mirdita, and Martin Steinegger. 2023. "Foldcomp: A Library and
        Format for Compressing and Indexing Large Protein Structure Sets." Edited by
        Lenore Cowen. *Bioinformatics* 39 (4): btad153.
        https://doi.org/10.1093/bioinformatics/btad153.

Lee, Brian M., Jing Xu, Bryan K. Clarkson, Maria A. Martinez-Yamout, H. Jane Dyson,
        David A. Case, Joel M. Gottesfeld, and Peter E. Wright. 2006. "Induced Fit and
        'Lock and Key' Recognition of 5S RNA by Zinc Fingers of Transcription Factor
        IIIA." *Journal of Molecular Biology* 357 (1): 275–91.
        https://doi.org/10.1016/j.jmb.2005.12.010.

Leipe, Detlef D., Eugene V. Koonin, and L. Aravind. 2003. "Evolution and Classification of
        P-Loop Kinases and Related Proteins." *Journal of Molecular Biology* 333 (4): 781–
        815. https://doi.org/10.1016/j.jmb.2003.08.040.

Leipe, Detlef D, Yuri I Wolf, Eugene V Koonin, and L Aravind. 2002. "Classification and
        Evolution of P-Loop GTPases and Related ATPases." *Journal of Molecular Biology*
        317 (1): 41–72. https://doi.org/10.1006/jmbi.2001.5378.

Lin, Zeming, Halil Akin, Roshan Rao, Brian Hie, Zhongkai Zhu, Wenting Lu, Nikita
        Smetanin, et al. 2023. "Evolutionary-Scale Prediction of Atomic-Level Protein
        Structure with a Language Model."

Mattle, Daniel, Antra Zeltina, Jae-Sung Woo, Birke A. Goetz, and Kaspar P. Locher. 2010.
        "Two Stacked Heme Molecules in the Binding Pocket of the Periplasmic Heme-
        Binding Protein HmuT from Yersinia Pestis." *Journal of Molecular Biology* 404 (2):
        220–31. https://doi.org/10.1016/j.jmb.2010.09.005.

McDonald, Ian K., and Janet M. Thornton. 1994. "Satisfying Hydrogen Bonding Potential
        in Proteins." *Journal of Molecular Biology* 238 (5): 777–93.
        https://doi.org/10.1006/jmbi.1994.1334.

McGinnis, S., and T. L. Madden. 2004. "BLAST: At the Core of a Powerful and Diverse Set
        of Sequence Analysis Tools." *Nucleic Acids Research* 32 (Web Server): W20–25.
        https://doi.org/10.1093/nar/gkh435.

Minami, Shintaro, Kengo Sawada, Motonori Ota, and George Chikenji. 2018. "MICAN-
        SQ: A Sequential Protein Structure Alignment Program That Is Applicable to
        Monomers and All Types of Oligomers." Edited by Alfonso Valencia.
        Bioinformatics 34 (19): 3324–31. https://doi.org/10.1093/bioinformatics/bty369.

Perdigão, Nelson, Julian Heinrich, Christian Stolte, Kenneth S. Sabir, Michael J. Buckley, Bruce Tabor, Beth Signal, et al. 2015. "Unexpected Features of the Dark Proteome." *Proceedings of the National Academy of Sciences* 112 (52): 15898–903. https://doi.org/10.1073/pnas.1508380112.

Remmert, Michael, Andreas Biegert, Andreas Hauser, and Johannes Söding. 2012. "HHblits: Lightning-Fast Iterative Protein Sequence Searching by HMM-HMM Alignment." *Nature Methods* 9 (2): 173–75. https://doi.org/10.1038/nmeth.1818.

Robert, Xavier, and Patrice Gouet. 2014. "Deciphering Key Features in Protein Structures with the New ENDscript Server." *Nucleic Acids Research* 42 (W1): W320–24. https://doi.org/10.1093/nar/gku316.

Saraste, Matti, Peter R. Sibbald, and Alfred Wittinghofer. 1990. "The P-Loop — a Common Motif in ATP- and GTP-Binding Proteins." *Trends in Biochemical Sciences* 15 (11): 430–34. https://doi.org/10.1016/0968-0004(90)90281-F.

Schrodinger. 2015. "The PyMOL Molecular Graphics System, Version 1.8."

Sehnal, David, Sebastian Bittrich, Mandar Deshpande, Radka Svobodová, Karel Berka, Václav Bazgier, Sameer Velankar, Stephen K Burley, Jaroslav Koča, and Alexander S Rose. 2021. "Mol* Viewer: Modern Web App for 3D Visualization and Analysis of Large Biomolecular Structures." *Nucleic Acids Research* 49 (W1): W431–37. https://doi.org/10.1093/nar/gkab314.

Steinegger, Martin, and Johannes Söding. 2017. "MMseqs2 Enables Sensitive Protein Sequence Searching for the Analysis of Massive Data Sets." *Nature Biotechnology* 35 (11): 1026–28. https://doi.org/10.1038/nbt.3988.

Suzek, Baris E., Yuqi Wang, Hongzhan Huang, Peter B. McGarvey, Cathy H. Wu, and the UniProt Consortium. 2015. "UniRef Clusters: A Comprehensive and Scalable Alternative for Improving Sequence Similarity Searches." *Bioinformatics* 31 (6): 926–32. https://doi.org/10.1093/bioinformatics/btu739.

Taylor, William R., Vijayalakshmi Chelliah, Siv Midtun Hollup, James T. MacDonald, and Inge Jonassen. 2009. "Probing the 'Dark Matter' of Protein Fold Space." *Structure* 17 (9): 1244–52. https://doi.org/10.1016/j.str.2009.07.012.

Van Kempen, Michel, Stephanie S. Kim, Charlotte Tumescheit, Milot Mirdita, Jeongjae Lee, Cameron L. M. Gilchrist, Johannes Söding, and Martin Steinegger. 2023. "Fast and Accurate Protein Structure Search with Foldseek." *Nature Biotechnology*, May. https://doi.org/10.1038/s41587-023-01773-0.

Varadi, Mihaly, Stephen Anyango, Mandar Deshpande, Sreenath Nair, Cindy Natassia, Galabina Yordanova, David Yuan, et al. 2022. "AlphaFold Protein Structure Database: Massively Expanding the Structural Coverage of Protein-Sequence Space with High-Accuracy Models." *Nucleic Acids Research* 50 (D1): D439–44. https://doi.org/10.1093/nar/gkab1061.

Walker, J.E., M. Saraste, M.J. Runswick, and N.J. Gay. 1982. "Distantly Related Sequences in the Alpha- and Beta-Subunits of ATP Synthase, Myosin, Kinases and Other ATP-Requiring Enzymes and a Common Nucleotide Binding Fold." *The EMBO Journal* 1 (8): 945–51. https://doi.org/10.1002/j.1460-2075.1982.tb01276.x.

Wang, Jiyao, Farideh Chitsaz, Myra K Derbyshire, Noreen R Gonzales, Marc Gwadz, Shennan Lu, Gabriele H Marchler, et al. 2023. "The Conserved Domain Database in 2023." *Nucleic Acids Research* 51 (D1): D384–88. https://doi.org/10.1093/nar/gkac1096.

Wang, Shuhui, Kaixuan Zhou, Xiaolin Yang, Bing Zhang, Yao Zhao, Yu Xiao, Xiuna Yang,
et al. 2020. "Structural Insights into Substrate Recognition by the Type VII
Secretion System." *Protein & Cell* 11 (2): 124–37. https://doi.org/10.1007/s13238-
019-00671-z.
Wintjens, Rene T, Marianne J Rooman, and Shoshana J Wodak. n.d. "Automatic
Classification and Analysis of Aa-Turn Motifs in Proteins."
Xu, Jinrui, and Yang Zhang. 2010. "How Significant Is a Protein Structure Similarity with
TM-Score = 0.5?" *Bioinformatics* 26 (7): 889–95.
https://doi.org/10.1093/bioinformatics/btq066.
Yang, Wei. 2008. "An Equivalent Metal Ion in One- and Two-Metal-Ion Catalysis." *Nature*
Structural & Molecular Biology 15 (11): 1228–31.
https://doi.org/10.1038/nsmb.1502.
496

497
Figure supplement 1 | Model confidence of initially mined dwNTPase structures. a, The cartoon shows the representative dwNTPase structure (UniProt accession No.: A0A1Y0TWD8). The cartoon is colored according to the residue-wise values of pLDDT. The coloring scheme is shown on the right of the structure, which is the same as displayed in AlphaFold DB. **b**, Distribution of total pLDDT values among the 711 dwNTPase structures initially mined from AlphaFold DB. **c–e**, The most confident, second worst and worst predictions among the 711 dwNTPase structures (A0A7V6TLZ4, A0A7C6J2V5 and W4RL53) gave average pLDDTs of 95.67, 88.43 and 88.01, respectively. Structures were created by Mol* viewer.

6h35
TM-score: 0.5046
RMSD: 3.524 Å
Aligned strands: 1,2,3,4 and 5

d1f5na2
TM-score: 0.4453
RMSD: 3.762 Å
Aligned strands: 1,3,4, and 5

d4ivra_
TM-score: 0.4270
RMSD: 3.524 Å
Aligned strands: 1,2,4, and 5

d1g64a_
TM-score: 0.4753
RMSD: 3.827
Aligned strands: 1,2,4, and 5

Figure supplement 2 | Structure of the dwNTPase P-loop domain compared with other
representative P-loop NTPase protein structures. Aligned structures of the dwNTPase P-
loop domain and known NTPases. **a**, MglAa, (**b**) G-proteins, (**c**) NKs and (**d**) Rec-A like.
The P-loop domain of dwNTPase and known NTPase are shown on the top and bottom of
the panels, respectively. The latter three structures were selected from c.37.8, c.37.1 and
c.37.11 SCCSs in SCOPe that showed the highest similarity to the dwNTPase P-loop domain.
The aligned regions are colored blue. The PDB or SCOPe IDs are shown below the structures.
The RMSD, TM-score and list of aligned strands in the dwNTPase P-loop domain are
summarized at the bottom of the panels.

Figure supplement 3 | Topology diagram of the dwNTPase P-loop domain compared with other representative P-loop NTPases. a, Comparisons with MglAa, (**b**) G-proteins, (**c**) NKs and (**d**) RecA-like proteins. Arrows represent strands. Grey and black lines indicate junctions projecting behind and out of the β-sheet. The pair of aligned strands are colored in the same colors. Unaligned strands are colored white. Helices are omitted for clarity.

Figure supplement 4 | Comparison of the switch loop with the helical region conserved among P-loop NTPases. Typical P-loop NTPase structures are compared to the dwNTPase P-loop domain (**a**). MglAa (**b**), Ras (**c**), adenylate-kinase (**d**) and F1-ATPase (**e**) are selected from G-proteins (SCCS: c.37.8), NKs (c.37.1) and Rec-A like protein (c.37.11) from SCOPe. The switch loop or corresponding helical regions are orange, and the P-loops are cyan. C-terminal portions of the structures that follow strand 5 are omitted for clarity. Protein names and PDB IDs are shown above the panels.

Class 1: both P-loops intact (1259)   Class 2: P-loops 1 collapsed (294)

Class 3: P-loops 2 collapsed (315)   Class 4: both P-loops collapsed (23)

Class 5: domain 1 collapsed (15)   Class 6: domain 2 collapsed (48)

Figure supplement 5 | Variations in the dwNTPase structure. Depending on the conservation of P-loops and domains, the dwNTPase family can be classified into six classes. The class 1 structure (e.g., Bt. dwNTPase, Uniport accession No. A0A1Y0TWD8) preserves both P-loops intact. The class 2 structure (A0A1T4Y6S3) has P-loop 1 degenerate and P-loop 2 intact. The class 3 structure (A0A1Y4J6T6) has P-loop 1 intact and P-loop 2 degenerate. The class 4 structure (A0A1I3I7D1) has both P-loops degenerate. The class 5 structure (A0A1Y4EVI7) has domain 1 degenerate and domain 2 intact. The class 6 structure (A0A1O1VWI5) has domain 1 intact and domain 2 degenerate. The number of structures that belong to respective classes is shown in parentheses. The structures are colored according to a purple-white-orange gradient from the N- to C-terminus. Note that the structures were rotated to show the collapsed region in front.

Figure supplement 6 | Sequence logo of dwNTPase. Using HHblits, we performed a sequence search against the UniRef30 database and constructed multiple sequence alignments. Black, blue, green and red represent hydrophobic, positively charged, polar and negatively charged residues. The amino acid sequence of the representative structure (Bt. dwNTPase) and the secondary structures assigned by DSSP are shown. The vertical axis represents the bit score, and the horizontal axis represents the residue index. Putative functional residues discussed in the main text are indicated by purple stars.

Figure supplement 7 | Other characteristic residues and substructures found in dwNTPases. a, Location of the switch loop relative to the tunnel and the hydrogen bond formed between conserved lysine and glutamate residues. The switch loop and P-loop are shown in orange and cyan cartoon, and the rest of the structure is a grey surface. The side chain of Lys36 is shown. **b,** Root-mean-square fluctuation (RMSF) of the Cα atoms during 20 trajectories of 100 ns MD simulations. The positions of switch loops are indicated by orange bars and labels reading sl1/sl2. The numbers above the plot show the residue index of the start/end of the switch loops. **c,** Conserved residues around P-loops supporting the recognition of ATP molecules. The side chain atoms of conserved residue recognizing ATPs or Mg^{2+} ions are shown as sticks. P-loops, ATPs and Mg^{2+} ions are colored cyan, orange and green, respectively. Additional loops unobserved in other P-loop proteins are colored grey.

Figure supplement 8 | Comparison with periplasmic heme-binding proteins. (Left) A
crystal structure of periplasmic heme-binding protein HmuT (PDB ID: 3nu1) liganded with
two heme molecules represented in orange CPK. (Right) The structure of dwNTPase is
shown at the same scale. The cleft between two P-loop domains found in dwNTPase is
indicated by an arrow. The chain is shown in cartoon representation colored according to a
blue-white-red gradient from the N- to C-terminus.

Figure supplement 9 | Distribution of the net charge in the left and right halves of the dwNTPase structure. a, The representative structure was divided into left and right halves and used as a template to find corresponding halves in other structures. The structures are colored according to a purple-white-orange gradient from the N- to C-terminus. **b**, From the initially mined 711 dwNTPase structures, 707 structures with all secondary structures intact were selected. The structures were aligned to the left half of the representative structure, and the aligned regions were assigned as left halves. The remaining regions of the structures were assigned as the right halves. In calculating net charges, each Arg, His and Lys residue was counted to have a +1 charge, and each Asp and Glu residue was counted to have a –1 charge.