FLAgellum Member 8 modulates extravasation and extravascular distribution of

African trypanosomes.

Estefanía Calvo Alvarez[1]*, Jean Marc Tsagmo Ngoune[1], Anneli Cooper[2], Christelle

Travaillé[1,3], Aline Crouzols[1], Annette MacLeod[2] and Brice Rotureau*[1,4]

[1] Trypanosome Transmission Group, Trypanosome Cell Biology Unit, INSERM U1201,

Department of Parasites and Insect Vectors, Institut Pasteur, Université Paris Cité,

Paris, France.

[2] Wellcome Centre for Integrative Parasitology, College of Medical, Veterinary, and Life

Sciences, Henry Wellcome Building for Comparative Medical Sciences, Glasgow,

Scotland, United Kingdom.

[3] Photonic BioImaging (UTechS PBI), Institut Pasteur, Université Paris Cité, Paris,

France.

[4] Parasitology Unit, Institut Pasteur of Guinea, Conakry, Guinea.

*Corresponding authors:

estefaniacalvoalvarez@gmail.com & brice.rotureau@pasteur.fr

Keywords: *Trypanosoma brucei*, FLAM8, flagellum, extravasation, dissemination,

sensing

Short title: FLAM8 modulates extravascular trypanosome dissemination

6 Figures, 3 Supplementary Figures and 1 Supplementary Table

Abstract (231/300 words)

In the mammalian host, the biology of tissue-dwelling *Trypanosoma brucei* parasites is not completely understood, especially the mechanisms involved in their extravascular colonization. The trypanosome flagellum is an essential organelle in multiple aspects of the parasites' development. The flagellar protein termed FLAgellar Member 8 (FLAM8) acts as a docking platform for a pool of Cyclic AMP response protein 3 (CARP3) that is involved in signaling. FLAM8 exhibits a stage-specific distribution suggesting specific functions in the mammalian and vector stages of the parasite. Analyses of knockdown and knockout trypanosomes in their mammalian forms demonstrated that FLAM8 is not essential *in vitro* for survival, growth, motility and stumpy differentiation. Functional investigations in experimental infections showed that *FLAM8*-deprived trypanosomes can establish and maintain an infection in the blood circulation and differentiate into insect transmissible forms. However, quantitative bioluminescence imaging and gene expression analysis revealed that *FLAM8*-null parasites exhibit a significantly impaired dissemination in the extravascular compartment, that is restored by the addition of a single rescue copy of *FLAM8*. *In vitro* trans-endothelial migration assays revealed significant defects in trypanosomes lacking *FLAM8*, possibly due to cAMP signaling impairments when a pool of CARP3 is not stabilized within FLAM8-related scaffold. FLAM8 is the first flagellar component shown to modulate *T. brucei* distribution in the host tissues, possibly through sensing functions, contributing to the maintenance of extravascular parasite populations in mammalian anatomical niches, especially in the skin.

Author Summary (192/200 words)

Trypanosoma brucei parasites cause neglected tropical diseases termed human and

animal African trypanosomiases. Transmitted by the bite of an infected tsetse fly, upon

deposition in the skin of a mammalian host, these parasites occupy both the

vasculature and extravascular tissues. Currently, the biology of tissue-resident

parasites is not well understood, and the parasite factors that mediate extravascular

colonization are not known. Using quantitative *in vivo* bioluminescence imaging and

ex vivo gene expression quantification in host infected tissues and blood, we reveal

that the flagellar parasite protein FLAM8 modulates the extravascular dissemination of

trypanosomes in the mammalian host. FLAM8 is known to act as a docking platform

for signaling complexes in the flagellum, but we observe that it does not influence

parasite differentiation into transmissible stages. However, we show that the absence

of FLAM8 results in the loss of a key component of the flagellar adenylate cyclase

signaling complexes, and reduces parasite migration through endothelial cell

monolayers, suggesting that FLAM8 is important for parasite exchanges between the

intravascular and the extravascular compartments. This work identifies a key

trypanosome flagellar component involved in host-parasite interactions, including the

modulation of parasite tropism and extravascular dissemination.

Introduction

Trypanosoma brucei is an extracellular parasite responsible for African trypanosomiases, including sleeping sickness in humans and *nagana* in cattle. African trypanosomes are blood and tissue-dwelling protists transmitted by the bite of the blood-feeding tsetse fly (*Glossina* genus). In the mammalian host, parasites face different micro-environments, including deadly challenges by multiple types of host immune responses and variable availability of carbon sources. This requires major morphological and metabolic adaptations, driven by the activation of specific gene expression programs, that are critical for life-cycle progression [1-3]. Recently, the importance of extravascular tropism for *T. brucei* has been re-discovered in animal models: in addition to the brain, parasites occupy most mammalian tissues, especially the skin and the adipose tissues [4-7]. However, the role of extravasation and sequestration of trypanosomes in specific anatomical niches and the underlying molecular processes are not understood yet.

The trypanosome flagellum is an essential organelle anchored along the surface of the cell body and present in all stages of its development [8]. It is essential for parasite viability [9], cell division and morphogenesis [10], attachment to the tsetse salivary glands [11] and motility [12]. In the insect host, the flagellum remains at the forefront of the cell and is likely to be involved in sensory and signaling functions required for host-parasite interactions [13, 14]. In the mammalian host, flagellar motility was shown to be critical for establishment and maintenance of bloodstream infection [15]. Nevertheless, the contributions of the trypanosome flagellum to parasite tropism and spatiotemporal dissemination outside vessels in the mammalian host remained to be explored.

A proteomic analysis of intact flagella purified from the insect stage of the parasite

identified a group of flagellar membrane and matrix proteins with unique patterns and

dynamics [16]. Amongst them, one large protein (3,075 amino acids) termed FLAgellar

Member 8 (FLAM8) is present only at the distal tip of the flagellum in the insect

procyclic form [16-18]. Interestingly, FLAM8 is redistributed along the entire length of

the flagellum in mammalian forms, including in stumpy transmissible stages [19], which

may imply a stage-specific function for this protein. Therefore, we hypothesized that

FLAM8 could be involved in host-parasite interactions or in developmental

morphogenesis. Here, we investigated the roles of FLAM8 in mammalian form

parasites *in vitro* in terms of survival, proliferation, motility, and differentiation, as well

as the *in vivo* dynamics of the intravascular and extravascular parasite burden over

the course of murine infection. Intriguingly, experimental infections in mice monitored

and quantified by bioluminescence imaging and gene expression analyses

demonstrated the involvement of FLAM8 in parasite dissemination in the extravascular

compartments. In addition, *in vitro* transmigration studies detected impaired

extravasation ability of *FLAM8*-deprived parasites, possibly resulting from the loss of

components of flagellar adenylate cyclase signaling complexes.

Results

FLAM8 RNAi silencing does not affect parasite survival

The differential distribution of FLAM8 in the flagellum of the different trypanosome stages [19] raises the question of its specific functions during the parasite cycle. To investigate the potential role(s) of FLAM8 in the mammalian host, bloodstream form parasites were first engineered for inducible RNAi knockdown of *FLAM8* in a monomorphic strain expressing a mNeonGreen-tagged version of FLAM8. In order to monitor their behavior in mouse by whole-body imaging approaches, these *FLAM8::mNG FLAM8*[RNAi] mutants were subsequently transformed to overexpress a chimeric triple reporter protein [20]. Upon RNAi induction with tetracycline for 72 h *in vitro*, *FLAM8* expression was reduced by 60% at the mRNA level (Fig. 1A) and became undetectable at the protein level by immunofluorescence (Fig. 1B). Parasite growth was monitored over 6 days upon induction of RNAi and no impact on proliferation was observed (Fig. 1C), which suggests that FLAM8 is possibly not essential for survival of mammalian forms of the parasite in cell culture conditions.

Then, the linear correlation between the emitted bioluminescence and the number of parasites was assessed in an IVIS Spectrum imager prior to *in vivo* challenge (Fig. S1). To get insights into the function of FLAM8 in the mammalian host, groups of male BALB/c mice were infected by the intraperitoneal route with 10^5 parasites of the parental, non-induced and induced cell lines (Fig. 1D). *In vivo* RNAi silencing of *FLAM8* was maintained in mice by the addition of doxycycline in sugared drinking water 48 h prior infection and until the end of the experiment. The course of the infection was monitored daily by i) quantifying the parasitemia, and ii) acquiring the bioluminescent signal emitted by the parasites in the entire organism with an IVIS Spectrum imager. The number of parasites in the extravascular compartment at a given timepoint can be

extrapolated by subtracting the known number of trypanosomes in the vascular system

(parasitemia x blood volume, according to body weight) from the total number of

parasites in the organism (total bioluminescence). No differences were detected,

neither in the establishment of the infection and the subsequent variations in the

number of intravascular parasites (Fig. 1E, IV), nor in the number of parasites

occupying extravascular tissues (Fig. 1F, EV). Similar population profiles were

observed over the course of the infection for both intra- and extravascular parasites in

each of the three independent groups of infected mice (Fig. 1D-F). In the extravascular

compartment, no significant differences were detected in terms of parasite

dissemination over the entire animal body in any group (Fig. 1G).

FLAM8 knockout does not affect trypanosome growth *in vitro*

Considering that i) *FLAM8* RNAi silencing efficiency was only partial (40% *FLAM8*

mRNA left after 72 h of induction), ii) the efficacy of doxycycline-induced *FLAM8*

repression could have been even lower *in vivo*, and that iii) the parental strain used for

this first strategy was monomorphic (i.e. unable to differentiate into tsetse adapted

stumpy stages), we reasoned that a gene knockout approach in a pleomorphic strain

would be more appropriate to evaluate the potential role(s) of FLAM8 during the

mammalian host infection. Therefore, a ∆*FLAM8* knockout cell line was generated in

pleomorphic trypanosomes by homologous recombination. The full replacement of one

FLAM8 allele and the partial replacement of the second one (to allow the use of a

shorter *in situ* rescue sequence than the long *FLAM8* coding sequence) by distinct

resistance cassettes was verified by whole-genome sequencing and PCR (Fig. S2).

The absence of full or truncated FLAM8 protein expression was further assessed by

immunofluorescence analysis with an anti-FLAM8 antibody [19] targeting the region of

FLAM8 that was not replaced in the second allele (Fig. 2A). This was also confirmed

at the mRNA level by RT-qPCR (Table S1). The trypanosome cell lines generated

were further transformed to express the chimeric triple reporter construct [20], and the

linear correlation between the bioluminescence signal and the total number of

parasites was analyzed for all strains both *in vitro* and *in vivo* (Fig. S3). In Δ*FLAM8*

knockout parasites where a rescue copy of one *FLAM8* allele was added back into its

endogenous locus, the FLAM8 distribution was restored along most of the flagellum

length, as assessed by immunofluorescence analysis (Fig. 2A). No impact on parasite

growth in culture conditions was observed (Fig. 2B). Next, we investigated whether the

loss of FLAM8 could have impacted the total length of the flagellum based on the

measurement of the signal obtained with the axonemal marker mAb25: no difference

was observed among BSF lines (Fig. 2C). In addition, the absence of *FLAM8* did not

affect parasite speed or linearity *in vitro*, neither in terms of speed nor in linearity (Fig.

2D-E). Compatible with our previous observations after RNAi silencing, these results

show that pleomorphic trypanosomes tolerated the loss of *FLAM8 in vitro*.

FLAM8 knockout affects trypanosome distribution in the mammalian host

Then, functional investigations in the mammalian host were performed by infecting

BALB/c mice either with the pleomorphic parental strain, three distinct Δ*FLAM8*

knockout subclones (resulting from independent recombination events) or one

Δ*FLAM8* strain bearing a rescue copy of *FLAM8* (Fig. 3). The infections were

monitored daily for 4 weeks by quantifying the parasitemia and the bioluminescence

signals emitted from whole animals (Fig. 3A). Null mutant parasites were able to

establish an infection in the bloodstream as well as in the extravascular compartment.

Within the vasculature, the overall amounts of parasites throughout the whole

experimental infection were comparable in all strains (Fig. 3B), yet with a slightly lower

parasitemia observed at the first peak of parasitemia in mice infected with the three

Δ*FLAM8* subclones as compared to mice infected with the parental strain (Fig. 3B). On

the other hand, quantitative analyses of extravascular parasites showed a different

scenario. Unlike in the intravascular compartment, significantly lower numbers of

extravascular trypanosomes were observed between day 5 to 12, and from day 19

post-infection until the end of the experiment at day 27, evidencing an impaired

extravascular colonization for Δ*FLAM8* parasites as compared to parental controls

(Fig. 3C). To note, distribution profiles of all pleomorphic strains were different from

those observed in mice infected with monomorphic parasites: larger amounts of

trypanosomes were seen occupying the extravascular compartment reaching up to 5-

8×10^9 parasites, while the maximum amount found in the bloodstream never exceeded

7×10^7 trypanosomes (Fig. 3B and C, respectively).

Furthermore, quantification of the parasite spreading over the whole animal bodies

showed that the depletion of *FLAM8* resulted in significantly impaired dissemination of

Δ*FLAM8* null mutant parasites in the extravascular compartment (days 5 to 13, 19 and

27 post-infection). This was mostly restored in trypanosomes bearing a rescue copy of

FLAM8 (Fig. 3D).

198 *FLAM8* knockout impairs extravascular trypanosome dissemination

We reasoned that the accuracy of the intra- Vs. extravascular parasite population

estimation from bioluminescence detection on small regions of interest could be

limited. Therefore, the same experiment was repeated, with comparable trends in

terms of extravascular parasite populations and parasite dissemination according to

the strain (Fig. 4A-C). At the end of this second experimental challenge (Day 24),

trypanosomes were removed from the vascular system by saline perfusion prior to

organ dissection. Individual organs were then collected for quantifying the expression

of different parasite genes by RT-qPCR. First, the absence of *FLAM8* transcripts was

confirmed in animals infected with Δ*FLAM8* parasites, whereas FLAM8 mRNAs were

detected in mice infected with both the parental and rescue strains (Table S1).

Normalized *Tubulin* expression was then used to compare the parasite densities

between organs and strains (Fig. 4D-E). In all mice, the highest number of

extravascular parasites was observed in the skin. In all organs, a marked decrease in

the number extravascular parasites was detected from mice infected with Δ*FLAM8*

parasites as compared to mice infected by parental and rescue parasites (Fig. 4D).

When considering the average difference between the number of parasites in organs

versus blood by strains, the statistically significant decrease in extravascular FLAM8-

deprived parasite densities appeared even more clearly (Fig. 4E). The rescue cell line

recapitulated EV parasite profiles in all organs examined with an average delta that

was not statistically different from the parental line.

We reasoned that this reduced extravascular dissemination of FLAM8-deprived

parasites could possibly be due to: (1) a lower proliferation rate, (2) a defect in motility,

(3) a higher rate of differentiation into non-proliferative stumpy forms, and / or (4) an

extravasation defect. The two first hypotheses could immediately be discarded as no

difference was observed between strains, neither in cell growth *in vitro* (Fig. 2B) and

in vivo (Fig. 3B and 4A), nor in cell motility *in vitro* (Fig. 2D-E). The two last hypotheses

were then successfully tested.

The absence of FLAM8 does not affect parasite differentiation

The systemic reduction in the extravascular Δ*FLAM8* parasite population could result

from a disequilibrium in the parasite differentiation into non-proliferative transmissible

forms. Therefore, we assessed whether the absence of FLAM8 could have impacted

the differentiation of proliferative slender into transmissible stumpy forms in the blood

and organs by immunofluorescence or RT-qPCR. In cultured parasites, the absence

of *FLAM8* did not significantly alter differentiation (Fig. 5A-B), and freshly differentiated

FLAM8-deprived stumpy parasites were able to further differentiate and maintain *in*

vitro as procyclic trypanosomes, as were the parental and rescue strains (Fig. 5C-D).

In vivo, the natural differentiation of proliferative slender into non-proliferative stumpy

parasites was confirmed by anti-PAD1 immunofluorescence staining of blood sampled

at the first peak of parasitemia (Fig. 5E, Day 5 of the first *in vivo* challenge), as well as

by RT-qPCR on blood and dissected organs at the end of the experiment (Fig. 5F, Day

24 of the second *in vivo* challenge). For each sample, *Tubulin* expression was used to

normalize the *PAD1* mRNA levels to compare the average levels of parasites

expressing *PAD1* mRNAs over the total parasite populations between organs and

strains (Fig. 5F), with a higher Delta CqPAD1-CqTub correlating with a lower amount

of *PAD1* transcripts in the organ. *PAD1* transcripts were detected in each organ at

least in one individual, showing that all tested organs represent a suitable environment

for the parasites to differentiate into transmissible forms (Fig. 5F). In mice infected with

the parental and rescue strains, the highest levels of *PAD1* transcripts were detected

in blood and skin, suggesting an accumulation or an increased rate of differentiation in

organs directly involved in parasite transmission. No *PAD1* transcripts were detected

in the gut, liver and spleen in mice infected with the parental and rescue strains,

whereas *PAD1* transcripts were detected in these three organs from mice infected with

the Δ*FLAM8* strains. However, when considering the average Delta CqPAD1-CqTub

in the entire organisms by strains, the relative proportions of parasites expressing

PAD1 mRNAs were not significantly different among groups, confirming that FLAM8 is

not involved in parasite differentiation (Fig. 5F).

Parasite extravasation is impaired in *FLAM8*-deprived trypanosomes

To test the last hypothesis, we asked whether the absence of FLAM8 might influence the way parasites traverse the vessel walls to access extravascular tissues. We first established a transmigration assay using human umbilical vein endothelial cells (HUVECs) grown to confluence on polyester transwell inserts with 3 μM pores, separating two chambers to mimic the vascular endothelium *in vitro* (Fig. 6A). Once HUVECs have reached confluence, 10^6 parental, *FLAM8*-deprived or rescue trypanosomes were added to the upper chamber and incubated for 24h. The number of trypanosomes that migrated through the endothelial monolayer into the lower chamber and the number of non-migrating trypanosomes remaining in the upper chamber of the transwell system were counted by flow cytometry to determine the transmigration percentage. Interestingly, all *FLAM8*-knockout subclones showed a significant reduction in transmigration relative to parental controls (25,6%, 70,7% and 33,8% of parasites crossing for KO1, 2 and 3, respectively) (Fig. 6B). Rescue trypanosomes expressing only one *FLAM8* allele exhibited an intermediate transmigration phenotype in which 59,2% of the parasites migrated through the endothelial monolayer (Fig. 6B), which was lower than for parental trypanosomes but significantly higher than for KO1 and KO3 null mutants (Fig. 6B). This demonstrates a strong impairment of parasites lacking *FLAM8* to cross through a confluent layer of endothelial cells, which was partially restored in rescue parasites.

Knowing that, in mammalian forms, FLAM8 was recently identified to be part of a flagellar complex including the cyclic AMP response protein 3 (CARP3) [21], we reasoned that this trans-endothelial crossing impairment could be caused by a defect in sensing and / or signaling. To assess how the absence of FLAM8 was impacting CARP3 localization and/or abundance, CARP3 was imaged by IFA in all strains (Fig.

6C). As expected, CARP3 was detected in a punctiform pattern along the entire

flagellum length in parental mammalian forms, but not in ∆*FLAM8* parasites, and the

presence of a *FLAM8* rescue copy in ∆*FLAM8* parasites restored the detection of

CARP3, yet in lower amounts than in parental cells (Fig. 6C).

In total, these data show that FLAM8 is likely involved in a cellular pathway modulating

trypanosome extravasation and consequently trypanosome dissemination in the

extravascular compartment of the mammalian host.

Discussion

The differential localization of FLAM8 from the very distal tip in tsetse midgut procyclic parasites to the entire flagellum length in the mammalian-infectious stages [19] prompted us to speculate that FLAM8 could play a distinct and specific role in each host. Here, we present for the first time the connection of a flagellar protein with the efficiency of trypanosomes to disseminate outside the mammalian host vasculature, especially in the skin. Quantitative analyses of experimental animal infections monitored by bioluminescence imaging and gene expression analysis showed that the absence of FLAM8 impairs parasite extravasation and dissemination in the host extravascular compartment over the time of the infection, which was mostly recovered by the integration of a single rescue copy of the *FLAM8* gene in the endogenous locus.

1. FLAM8 and trypanosome transmission

In the bloodstream, the balance between proliferative slender parasites and tsetse-adapted stumpy forms responds to a quorum sensing mechanism involving the production of oligopeptides and their reception through a specific transporter [22]. The absence of FLAM8 did not alter the ability of the parasites to differentiate into transmissible stumpy forms at the two observed time points (first peak of parasitemia and 3 weeks after infection). Although stumpy proportions were not evaluated over the entire course of the infection, this evidence suggests that FLAM8 does not play an integral role in this process. In addition, the retained ability of *FLAM8*-deprived stumpy cells to differentiate into procyclic forms *in vitro* suggests that they could further pursue their cyclical development upon ingestion by a tsetse fly.

Extravascular trypanosomes occupy the interstitial space of several organs, including the central nervous system, testes, adipose tissues, and skin [4-7, 23-25]. The relevance of skin-dwelling trypanosomes in parasite transmission has been

demonstrated by xenodiagnosis experiments, early after the infective bite [26], or later

in the infection [4], even in the absence of detectable parasitemia. More recently, the

presence of extravascular trypanosomes was confirmed in the skin of confirmed and

suspected cases of sleeping sickness [27]. Here, we show that all tested organs

represent a suitable environment for the parasites to differentiate into transmissible

forms, yet the highest amounts of *PAD1* transcripts were detected in blood and skin,

suggesting an increased rate of differentiation, or an accumulation of differentiated

stumpy forms in organs directly involved in parasite transmission. In total, the impaired

spreading of *FLAM8*-null parasites over the extravascular compartment would

mathematically reduce the probability for parasites to be ingested by tsetse flies. This

could be especially significant for the skin, that has the highest parasite density, and

the highest overall parasite load due to its size, at the direct interface with insect

vectors.

2. On the possible cellular function(s) of FLAM8

Proliferative slender trypanosomes are highly mobile [28] and this motility was proved

to influence virulence *in vivo*. For instance, the lack of propulsive motility in flagellar

dynein light chain 1 (*LC1*) knockout mutants resulted in the inability of trypanosomes

to establish an infection in the bloodstream [15]. Here, *FLAM8* depletion in pleomorphic

mammalian forms did not alter parasite growth and cell motility in matrix-containing

medium. However, quantitative analyses showed that *FLAM8*-null trypanosomes were

less numerous in extravascular tissues as compared to parental controls. Assuming

that parasite motility could be different in tissues and interstitial spaces with biophysical

properties distinct from those in the blood [28, 29], one cannot exclude that the motility

of Δ*FLAM8* knockout parasites might be somehow altered in the extravascular

compartment. Intravital imaging for motility analyses at the cell level would be needed to confirm this hypothesis.

Historically, most studies on *T. brucei* virulence in experimental infections have considered the blood circulation almost as the sole host compartment parasitized by trypanosomes, whereas extravascular parasite niches and the underlying exchanges between both compartments have been underestimated for long. De Niz and colleagues recently identified adhesion molecules as key players for tissue tropism [30]. They showed that reservoir establishment happens before vascular permeability is compromised, suggesting that extravasation is an active mechanism, and depends on trypanosome interactions with endothelial surface adhesion molecules, such as E-selectin, P-selectins, or ICAM2 [30]. Here, we showed that the absence of FLAM8 led to a strong impairment in parasite extravasation. The fact that *FLAM8*-null trypanosomes were not able to disseminate properly over the extravascular compartment could somehow imply defects in the way parasites sense their microenvironment or interact with the host endothelial cells, resulting in an alteration of their extravascular tropism.

The insect forms´ coordinated social motility *in vitro* is linked to cAMP signaling at the flagellar tip [31, 32], i. e. where FLAM8 localizes [19]. The architecture of an adenylate cyclase complex in the tip nanodomain was recently shown to be essential for social motility and salivary gland colonization [21]. In this complex, CARP3 interacts with the catalytic domain of adenylate cyclases and regulates abundance of multiple adenylate cyclase isoforms. We recently demonstrated that the CARP3 tip localization depends on the presence of FLAM8 acting as a scaffold protein [21]. Thus, trypanosome migration and transmission in the tsetse vector specifically depend on adenylate cyclase complex-mediated signaling in the tip nanodomain including FLAM8 [21].

Interestingly, we have recently shown that CARP3 and FLAM8 are both redistributed

along the entire length of the flagellum during their differentiation to the mammalian

stage, and that they further remain associated in flagellar complexes in mammalian

forms [21]. Here, this was confirmed by the absence of CARP3 from the flagellum of

FLAM8-deprived mammalian forms. Altogether, these data may imply a stage-specific

function of CARP3-containing signaling complexes depending on FLAM8 for their sub-

flagellar localization. Environmental sensing and / or signaling, possibly through direct

contact with host cell receptors may play a role in extravasation. We propose that the

absence of FLAM8 would destabilize or delocalize these signaling complexes,

impairing parasite sensing, signaling and / or adhesion functions, hence preventing the

parasite to efficiently cross vascular endothelia.

To our knowledge, FLAM8 is the first flagellar component affecting parasite

extravasation and their subsequent dissemination in the extravascular host tissues *in*

vivo. Further investigations on the FLAM8 interactions with other possible partners in

the flagellum would help to unravel the roles of this fascinating and essential organelle,

especially regarding the modulation of trypanosome tropism, extravasation and

spreading in their mammalian hosts, and its implications in parasite virulence and

transmission.

Materials and Methods

Strains, culture and *in vitro* differentiation

The AnTat 1.1E Paris pleomorphic clone of *Trypanosoma brucei brucei* was derived from a strain originally isolated from a bushbuck in Uganda in 1966 [33]. The monomorphic *T. brucei* strain Lister 427 [34] was also used. All bloodstream forms (BSF) were cultivated in HMI-11 medium supplemented with 10% (v/v) fetal bovine serum (FBS) [35] at 37°C in 5% CO_2. For *in vitro* slender to stumpy BSF differentiation, we used 8-pCPT-2′-O-Me-5′-AMP, a nucleotide analog of 5′-AMP from BIOLOG Life Science Institute (Germany). Briefly, $2x10^6$ cultured pleomorphic AnTat 1.1E slender forms were incubated with 8-pCPT-2′-O-Me-5′-AMP (5 µM) for 48 h [36]. For specific experiments, *in vitro* differentiation of BSF into procyclic forms was performed by transferring freshly differentiated short stumpy forms into SDM-79 medium supplemented with 10% (v/v) FBS, 6 mM cis-aconitate and 20 mM glycerol at 27°C [37].

Monomorphic BSF "Single Marker" (SM) trypanosomes are derivatives of the Lister 427 strain, antigenic type MITat 1.2, clone 221a [38], and express the T7 RNA polymerase and tetracycline repressor. *FLAM8^RNAi* cells express complementary single-stranded RNA corresponding to a fragment of the *FLAM8* gene from two tetracycline-inducible T7 promoters facing each other in the pZJM vector [39] integrated in the rDNA locus [40]. Addition of tetracycline (1 µg/mL) to the medium induces expression of sense and anti-sense RNA strands that can anneal to form double-stranded RNA (dsRNA) and trigger RNAi. For *in vivo* RNAi studies in mice, doxycycline hyclate (Sigma Aldrich) was added in sugared drinking water (0.2 g/L doxycycline hyclate combined with 50 g/L sucrose).

Generation of FLAM8 RNAi mutants

For the generation of the *FLAM8^RNAi* cell lines, a 380 bp (nucleotides 6665-7044)

For the generation of the *FLAM8^RNAi* cell lines, a 380 bp (nucleotides 6665-7044)

fragment of *FLAM8* (Tb927.2.5760), flanked by 5' HindIII and 3' XhoI restriction sites

to facilitate subsequent cloning, was selected using the RNAit algorithm

(http://trypanofan.bioc.cam.ac.uk/software/RNAit.html) to ensure that the targeted

sequence was distinct from any other genes to avoid any cross-RNAi effects [41]. This

FLAM8 DNA fragment was synthesized by GeneCust Europe (Dudelange,

Luxembourg) and inserted into the HindIII-XhoI digested pZJM vector [39].

The pZJM-FLAM8 plasmid was linearized with NotI prior to transfection using

nucleofector technology (Lonza, Italy) as described previously [42]. The cell line was

further engineered for endogenous tagging of *FLAM8* with an mNeonGreen (mNG) at

its C-terminal end by using the p3329 plasmid [43], carrying a *FLAM8* gene fragment

corresponding to *FLAM8* ORF nucleotides 8892-9391. Prior to nucleofection, NruI

linearization of p3329-FLAM8-mNG plasmid was performed.

For *in vivo* experiments in mice, *FLAM8^RNAi* parasites were finally modified by

integrating a plasmid encoding for the chimeric triple reporter which combines the red-

shifted firefly luciferase PpyREH9 and the tdTomato red fluorescent protein fused with

a TY1 tag [20]. Transformants were selected with the appropriate antibiotic

concentrations: phleomycin (1 µg/mL), blasticidin (5 µg/mL), G418 (2 µg/mL), and

puromycin (0.1 µg/mL). Clonal populations were obtained by limiting dilution. Cell

culture growth was monitored with an automatic Muse cell analyzer (Merck Millipore,

Paris).

Generation of *FLAM8* KO mutants

For generating the *FLAM8* knockout and rescue cell lines, all insert templates were

synthesized by GeneCust Europe (Dudelange, Luxembourg). For breaking the first

allele, the 300 first nucleotides of the *FLAM8* gene flanking sequences were added on

each side of a HYG resistance cassette (Fig. S2). For a complete disruption of the

FLAM8 locus, a second selectable marker (PAC) was flanked with the *FLAM8*-flanking

sequence at 5' and by 300 nucleotides of the *FLAM8* ORF (nucleotides 501-800) at 3'.

For generating an add-back rescue cell line, due to the large size of the *FLAM8* ORF

(9,228 nucleotides), the PAC selection marker was replaced by a BLE marker cassette

flanked by the 300 first nucleotides of the *FLAM8* 5' untranslated region (UTR) and by

the nucleotides 1 to 500 of the *FLAM8* ORF for reinsertion into the endogenous locus

of the knockout cell line. PCR amplifications of the DNA fragments bearing the *FLAM8*

flanking sequences and the appropriate resistance markers were used for

nucleofection and generation of all cell lines. The primers used are listed below: 5'-

CATGACTTTACGTGTTTGGGCAC-3' (FW, located 82 bp upstream the flanking

5'UTR sequence); 5'-CTTGCTTGTTTCTGTTTCGCAAC-3' (RV, 130 bp downstream

the flanking 3'UTR sequence, used to replace one WT allele by HYG resistance

cassette); 5'- GCACACTAAAACTCATTGAAAGCC-3' (RV, 926 bp downstream the

ATG codon of *FLAM8*, used for second WT allele replacement by PAC cassette and

rescue line generation). All knockout and rescue cell lines were further transfected to

express the chimeric triple reporter protein PpyRE9H/TY1/tdTomato for multimodal *in*

vivo imaging approaches as described elsewhere [20]. Selection-marker recovery was

confirmed by screening individual clones in multi-well plates. Transformants were

selected with the appropriate antibiotic concentrations: phleomycin (1 µg/mL),

blasticidin (5 µg/mL), puromycin (0.1 µg/mL) and hygromycin (2.5 µg/mL). Clonal

populations were obtained by limiting dilution and cell culture growth was monitored

with an automatic Muse cell analyzer (Merck Millipore, Paris).

Knockout and rescue cell lines were validated by whole-genome sequencing (BGI,

Hong Kong). Briefly, genomic DNA from parental and mutant cell lines were sequenced

by the HiSeq4000 sequence system (Illumina), generating about 10 million 100-bp

reads and compared to that of the *T. brucei brucei* AnTat 1.1E Paris reference strain.

In addition, some validation of the construct integrations in mutants were performed by

PCR analysis according to standard protocols (Fig. S2 and Table 1).

Table 1. Oligonucleotides used for PCR validation of the ∆*FLAM8* knockout and rescue

cell lines.

Primer	Sequence	Target	
1	CATGACTTTACGTGTTTGGGCAC	*FLAM8* WT allele (82 nt upstream 5'UTR)	F
2	GCACACTAAAACTCATTGAAAGCC	*FLAM8* WT allele (926 nt downstream ATG)	R
3	CGTCCGAGGGCAAAGGAATAG	Hygromycin cassette	R
4	GACCGCGCACCTGGTGCATG	Puromycin cassette	R
5	GTGGCCGAGGAGCAGGACTGA	Phleomycin cassette	R

Orientation of primers: F, forward; R, reverse.

Motility analyses

In silico 2D tracking was performed as previously described [44]. For each BSF strain,

10 to 20 movies were recorded for 20 seconds (50 ms of exposure). Trypanosomes at

1×10^6 parasites/mL were maintained in matrix-dependent HMI-11 medium containing

0,5% methylcellulose at 37°C and were observed under the 10x objective of an

inverted DMI-4000B microscope (Leica) coupled to an ORCA-03G (Hamamatsu) or a

PRIME 95B (Photometrics) camera. Movies were converted with the MPEG Streamclip

V.1.9b3 software (Squared 5) and analyzed with the medeaLAB CASA Tracking V.5.5

software (medea AV GmbH). Results were analyzed as mean ± SD of three

independent experiments.

In vitro bioluminescence quantification and analysis

To perform the parasite density / bioluminescence intensity assay, BSF parasites were counted, centrifuged, and resuspended in fresh HMI-11 medium. Then, 100 µL of this suspension containing 10^6 parasites were transferred into black clear-bottom 96-well plates and serial 2-fold dilutions were performed in triplicate adjusting the final volume to 200 µL with 300 µg/mL of beetle luciferin (Promega, France). Luciferase activity was quantified after 10 min of incubation with an IVIS Spectrum imager (PerkinElmer). Imaging data analysis was performed with the Living Image 4.3.1 software (PerkinElmer) by drawing regions of interest with constant size for well plate quantification. Total photon flux was calculated after removal of intensity values from WT parasites and / or parasite-free medium corresponding to the background noise. Results were analyzed as mean ± SD of three independent experiments (Fig. S1 and S3 A-B).

Mouse infection and ethical statements

Seven-week-old male BALB/c mice were purchased from Janvier Laboratory (sub-strain BALB/cAnNRj) and used as models for experimental infection and monitoring of the bioluminescence signal with the IVIS Spectrum imager (PerkinElmer). This study was conducted in strict accordance with the recommendations from the Guide for the Care and Use of Laboratory Animals of the European Union (European Directive 2010/63/UE) and the French Government. The protocol was approved by the "Comité d'éthique en expérimentation animale de l'Institut Pasteur" CETEA 89 (Permit number: 2012-0043 and 2016-0017) and undertaken in compliance with the Institut Pasteur Biosafety Committee (protocol CHSCT 12.131). BR is authorized to perform experiments on vertebrate animals (license #A-75-2035) and is responsible for all the experiments conducted personally or under his supervision. For *in vivo* infections,

groups of four and three animals (*FLAM8* knockdown and knockout-infected mice, respectively) were injected intraperitoneally (IP) with 10^5 slender BSF parasites, washed in TDB (Trypanosome Dilution Buffer: 5 mM KCl, 80 mM NaCl, 1 mM $MgSO_4$*$7H_2O$, 20 mM Na_2HPO_4, 2 mM NaH_2PO_4, 20 mM glucose) and resuspended in 100 µl of PBS prior animal inoculation.

In vivo bioluminescence imaging (BLI) analyses

Infection with bioluminescent parasites was monitored daily by detecting the bioluminescence signal in whole animals with the IVIS Spectrum imager (PerkinElmer). The equipment consists of a cooled charge-coupled camera mounted on a light-tight chamber with a nose cone delivery device to keep the mice anaesthetized during image acquisition with 1.5-2% isoflurane. A heated stage is comprised within the IVIS Spectrum imager to maintain optimum body temperature. D-luciferin potassium salt (Promega) stock solution was prepared in sterile PBS at 33.33 mg/mL and stored in a -20°C freezer. To produce bioluminescence, mice were inoculated by the intraperitoneal route (IP) with 150 µL of D-luciferin stock solution (250 mg/Kg body weight). After 10 minutes of incubation to allow substrate dissemination, all mice were anaesthetized in an oxygen-rich induction chamber with 1.5-2% isoflurane, and dorsal and ventral BLI images were acquired by using automatic exposure (0.5 seconds to 5 minutes) depending on signal intensity.

Images were analyzed with the Living Image software version 4.3.1 (PerkinElmer). Data were expressed in total photons/second (p/s) corresponding to the total flux of bioluminescent signal according to the selected area (ventral and dorsal regions of interest with constant size covering the total body of the mouse). The background noise was removed by subtracting the bioluminescent signal of the control mouse from the infected ones for each acquisition. For parasite dissemination analyses, a minimum

value of photons/second (p/s) was set for all animals in every time point to quantify the exact dissemination area (in cm^2) over the whole animal body. Parasitemia was determined daily following tail bleeds and assayed by automated fluorescent cell counting with a Muse cytometer (Merck-Millipore, detection limit at 10^2 parasites/mL) according to the manufacturer's recommendations. The quantification of the total intravascular parasite population was assessed by calculating the blood volume of all animals according to their body weight and referring to daily parasitemia.

To quantify the total number of parasites by BLI (intravascular plus extravascular trypanosomes), an *in vivo* standard curve was performed (Fig. S3 C-D). Since the bioluminescent emission of cultured parental, KO subclones and rescue parasites was not significantly different (Fig. S3 A-B), the *in vivo* standard curve was obtained by injecting IP increasing numbers (10^3, 10^4, 10^5, 10^6 and 10^7 parasites/animal) of parental trypanosomes only (Fig. S3 C-D). After 2.5h, animals received 150 µL of D-luciferin stock solution IP (250 mg/kg body weight), 10 minutes prior image acquisition. During this time, mice were anaesthetized with 1.5-2% isoflurane, and images were acquired in the IVIS Spectrum imager by using automatic exposure settings. A region of interest with a constant size was used to correlate the number of injected parasites with the whole-body BLI signal. Non-infected controls were imaged and the total BLI values used to subtract the background signal or noise. The signals in photons/second were used to construct a standard curve to further interpolate the total number of trypanosomes present in each animal during the entire experimental infection period (Fig. S3 D). Subsequently, to obtain the number of extravascular parasites, the calculated total number of parasites present in the vascular system was subtracted from the total number of trypanosomes per animal body, resulting in estimating the

547 total parasite population colonizing the extravascular compartments at a given time

548 point.

Endothelial transmigration assay

Single donor cryopreserved Primary Human Umbilical Vein Endothelial Cells

(HUVECs) were obtained from Promocell and maintained as per the manufacturer's

instructions in 75 cm^2 flasks at 37°C with 5% CO_2 in endothelial cell growth medium

(Promocell) with 100 U/mL penicillin, and 100 µg/mL streptomycin (Gibco). Cells were

passaged at 80-90% confluence by dissociation with 0.04% Trypsin-0.03% EDTA

(Promocell), split at 1:3-1:5 ratios, and maintained for up to six passages. Polyester

transwell inserts for 24 well plates with 3 µm pore size (Corning) were coated with 10

µg/mL bovine fibronectin (Promocell) for one hour, the excess removed, and 600 µL

pre-warmed endothelial cell growth medium added to the lower chamber (Fig 6A). The

upper chamber was seeded with 2x10^4 HUVECs per insert in a volume of 100 µL

endothelial cell growth medium. Media was exchanged in the upper and lower chamber

every two days until monolayer confluence reached (approximately 6 days).

Confluence was confirmed by FITC-dextran permeability assay and crystal violet

staining of a sacrificed transwell. Briefly, 1 mg/mL FITC-70kDa dextran (Sigma-Aldrich)

in endothelial cell growth medium was added to the upper chamber, incubated for 20

minutes and leakage to the lower chamber measured on a Qubit 4 Fluorometer

(Invitrogen). Leakage of <1% of the FITC-dextran was observed at confluence.

Monolayer integrity was confirmed by brightfield imaging of a transwell insert stained

with crystal violet as per the manufacturer's instructions (Millipore). To perform the

trypanosome transmigration assays, the confluent transwells were exchanged into a

new 24 well plate containing pre-warmed assay media (endothelial cell growth medium

supplemented with 20% trypanosome growth media) for two hours prior to performing

the assay. Cultured trypanosomes were collected by centrifugation at 900×g for 5 min and resuspended in assay media at $2x10^6$/mL. The media in the upper chamber was replaced with 100 μL of assay media containing $2x10^5$ trypanosomes and incubated overnight at 37°C, 5% CO_2. Each cell line was tested in triplicate transwells. Unbiased quantification of trypanosome transmigration was determined after 24 hours by collecting the media from the upper and lower chambers and transferring 100 μL to 96 well plates for automated counting using a Guava Easycyte HT system with a green laser and a custom counting protocol for tdTomato fluorescent trypanosomes. The cell counts per ml were used to calculate the number of trypanosomes in each compartment and proportion of transmigration into the lower chamber. Trypanosome transmigration was compared between the FLAM8 mutant cell lines and the parental reference cell line using the Generalised Linear Model function in R with a Gaussian family function and proportion transmigration as the dependant variable. A probability value of $p < 0.05$ was considered significant.

Ex vivo RT-qPCR of mouse tissues

Biological samples. After four weeks of *in vivo* monitoring, parental-, KO- and rescue-infected mice were euthanized by an overdose of an anesthetic/analgesic mixture of ketamine and xylazine (100 mg/kg and 10 mg/kg, respectively). Terminal bleeding was then performed through the inferior *vena cava*, and the collected blood was directly transferred in 1 ml RNAlater (ThermoFisher Scientific), snap frozen in liquid nitrogen for long-term preservation. Next, perfusion was initiated by injecting 0.9% NaCl (pre-heated at 37°C), to completely remove the blood from the animal's body (both organs and vasculature), thus ensuring that subsequent analyses would only contain tissue-dwelling trypanosomes. Finally, spleen, lungs, kidney, gut, testis, liver and skin tissue samples were preserved in 1 ml RNAlater, snap frozen and stored in liquid nitrogen.

RNA isolation. Total RNA from samples was isolated using the QIAzol Reagent

(Qiagen) according to the manufacturer's instructions. Frozen tissue samples were

weighed and processed on ice to prevent thawing. Briefly, 30 mg of tissue were added

to 700 µl of QIAzol lysis reagent, or 250 µl of blood in 550 µl of QIAzol lysis reagent,

and samples were homogenized using the Precellys® Evolution Homogenizer (Bertin,

USA) with 2.8 mm stainless steel beads for 2 cycles of 1 min at 5000 rpm each,

followed by 15 sec of resting between them. Homogenates were then incubated for 5

min at room temperature before the addition of 140 µl chloroform (0.2 volume of

starting QIAzol lysis reagent), thoroughly mixed by vortexing for 15 seconds, incubated

5 min at room temperature and centrifuged at $12,000 \times g$, 4°C, 15 min. The aqueous

phase containing the RNA was subsequently mixed with 1.5 volumes of absolute

ethanol and transferred into a RNeasy Mini spin column (QIAGEN®) following the

manufacturer's recommendations. The concentration of each RNA sample was

measured by spectrophotometric analysis in a NanoDrop 2000c (ThermoFisher

Scientific). Finally, RNA quality was determined by capillary electrophoresis in a 2100

Bioanalyzer (Agilent). Extracted RNA was stored at -80 °C prior to RT-qPCR analyses.

DNAse treatment and validation. Extracted RNA samples were subjected to a second

DNase treatment using Invitrogen's DNA-*free* kit (Life Technologies) according to

manufacturer's protocol. DNase treatment confirmation was performed by running a

qPCR targeting the *Tubulin* locus as housekeeping gene control. The primer pair used

is as follows: forward (FW), 5'-ACTGGGCAAAGGGCCACTAC-3'; reverse (RV), 5'-

CTCCTTGCAGCACACATCGA-3', with an amplicon size of 105 bp. Reactions were

done in a volume of 20 µl containing: 1 µl of RNA template (17 ng), 10 µl of 2X GoTaq

qPCR Master Mix Buffer, 2 µl of both FW and RV primers (at 10X) and 5 µl of Nuclease-

Free Water. Amplification was accomplished in a QuantStudio™ 3 thermal cycler

(Applied Biosystems) using the following program: 2 min at 95°C; 40 cycles at 94°C 15 sec; 55°C 1 min; and final 30 sec at 60°C. The absence of residual DNA in the DNAse-treated RNA samples was thus confirmed when no amplification was observed. RNA samples were further employed for gene expression quantification.

Primer design of target genes. For *FLAM8* amplification, primers were designed to recognize the 324-bp region suppressed during *FLAM8* knockout generation: FW, 5'-GCATCGTTCGTGAGGTTGGA-3'; RV, 5'-GTTCCTCTTCGTCATCTGGTTCA-3'. The amplicon size was 88 bp. For Protein Associated to Differentiation 1 (PAD1) quantification in infected samples, primer sequences were described elsewhere (Saldivia et al. 2016) and are listed below: FW, 5'-TCATGGTTTCGCCATTCTCGTAACC-3'; RV, 5'-CTCAGCCACTTCTCTCCTACAACAC-3'. Amplicon size is 156 bp.

Real time RT-PCR assay. One step RT-PCR kit (Promega) was used. Reactions were prepared in a volume of 20 µl, containing: 1 µl of RNA template (17 ng), 10 µl of 2X GoTaq qPCR Master Mix Buffer, 0.4 µl of 50X GoScript™ RT Mix for 1-Step RT-qPCR, 2 µl of both FW and RV primers (10X), and 4,6 µl of Nuclease-Free Water. Reverse transcription and amplification were accomplished in one step in a QuantStudio™ 3 thermal cycler (Applied Biosystems) using the following incubation program: 15 min at 42°C; 10 min at 95°C; 40 cycles of 95°C during 30 sec; 55°C for *Tubulin*, 58°C for *FLAM8* or 60°C for *PAD1* during 1 min; final 72°C during 30 sec. A melt curve program was included: 15 sec at 95°C; 55°C for *Tubulin*, 58°C for *FLAM8* or 60°C for *PAD1* during 1 min; 95°C for 10 sec. Amplicons were then analyzed by gel electrophoresis. Negative and positive controls consisted of RNA extracted from uninfected mice and cultured trypanosomes, respectively.

Table 2. Oligonucleotides used for *ex vivo* RT-qPCR of infected tissues.

Primer		Sequence	Purpose
Tubulin	F	ACTGGGCAAAGGGCCACTAC	Housekeeping control. Total trypanosome quantification (both SL and ST parasites) through standard curve generation.
	R	CTCCTTGCAGCACACATCGA	
FLAM8	F	GCATCGTTCGTGAGGTTGGA	Detection of parasites expressing *FLAM8*. The amplified sequence is absent in *FLAM8* KO mutants.
	R	GTTCCTCTTCGTCATCTGGTTCA	
PAD-1	F	TCATGGTTTCGCCATTCTCGTAACC	Quantification of transmissible ST parasites within *ex vivo* samples.
	R	CTCAGCCACTTCTCTCCTACAACAC	

Orientation of primers: F, forward; R, reverse.

RT-qPCR data analysis. All samples were amplified in triplicates and Cq mean values

were calculated. Considering that the same initial amounts of total mRNAs extracted

from each organ were used as RT-qPCR templates, the total number of parasites in

each sample was calculated for comparisons by using a *Tubulin* RT-qPCR standard

curve. Nine pools of cultured parasites (p) increasing by 10-folds from 10^1 to 10^8 were

extracted and tested in triplicates by *Tubulin* RT-qPCR to generate a standard curve.

The resulting standard curve's equation Cq = -2,87 x Log10(p) + 35,412 allowed us to

calculate the total number of parasites per sample according to the Cq values obtained

by *Tubulin* RT-qPCR on each sample. For normalization purposes, the difference

between the calculated number of parasites in each sample of a given mouse and the

calculated number of parasites in the blood sample from the same mouse (Delta

number of parasites) was calculated and plotted. *Tubulin* expression (CqTub) was also

used to normalize the *PAD1* mRNA levels (CqPAD1): the difference between the

CqPAD1 and the CqTub values was calculated for each organ of each mouse and

plotted as the Delta CqPAD1-CqTub. It allowed us to compare the relative proportions

of parasites expressing *PAD1* mRNAs between organs and strains, a higher Delta

CqPAD1-CqTub correlating with a lower amount of *PAD1* transcripts in the organ.

Immunofluorescence analysis (IFA)

Cultured parasites were washed twice in TDB and spread directly onto poly-L-lysine

coated slides. For methanol fixation, slides were air-dried for 10 min, fixed in methanol

at -20°C for 5 min and rehydrated for 20 min in PBS. For immunodetection, slides were

incubated for 1 h at 37°C with the appropriate dilution of the first antibody in 0.1% BSA

in PBS. After 3 consecutive 5 min washes in PBS, species and subclass-specific

secondary antibodies coupled to the appropriate fluorochrome (Alexa 488, Cy3, Cy5

Jackson ImmunoResearch) were diluted 1/400 in PBS containing 0.1% BSA and were

applied for 1 h at 37°C. After washing in PBS as indicated above, slides were finally

stained with 4',6-diamidino-2-phenylindole (DAPI, 1 µg/mL) for visualization of

kinetoplast and nuclear DNA content and mounted under cover slips with ProLong

antifade reagent (Invitrogen), as previously described [8]. Slides were observed under

an epifluorescence DMI4000 microscope (Leica) with a 100x objective (NA 1.4), an

EL6000 (Leica) as light excitation source and controlled by the Micro-Manager V1.4.22

software (NIH), and images were acquired using an ORCA-03G (Hamamatsu) or a

PRIME 95B (Photometrics) camera. Images were analyzed with ImageJ V1.8.0 (NIH).

The monoclonal antibody mAb25 (anti-mouse IgG2a, 1:10) was used as a flagellum

marker as it specifically recognizes the axoneme protein *Tb*SAXO1 [45]. FLAM8 was

detected using: i) a specific rabbit serum (1:500) kindly provided by Paul McKean

(University of Lancaster, UK), or ii) a monoclonal anti-mNeonGreen antibody (anti-

mouse IgG2c, 1:100, ChromoTek). CARP3 was detected using a polyclonal CARP3

antiserum (1:150) [21]. Stumpy BSF were identified at the molecular level with a rabbit

polyclonal anti-PAD1 antibody (kindly provided by Keith Matthews, University of

Edinburgh; dilution 1:300) [46]. In the case of RNAi knockdown experiments, IFA

signals were normalized using the signal obtained in non-induced controls as a

reference.

Measurements, normalization, and statistical analyses

Standardization of fluorescent signals was carried out by parallel setting of raw

integrated density signals in all the images to be compared in ImageJ V1.8.0 (NIH).

For clarity purposes, the brightness and contrast of several pictures were adjusted

after their analysis in accordance with editorial policies. Statistical analyses and plots

were performed with XLSTAT 2019.2.01 (Addinsoft) in Excel 2016 (Microsoft) or Prism

V9.3.1 (GraphPad). Statistical analyses include: (1) linear regression for

bioluminescence / fluorescence intensity vs. parasite density and RT-qPCR standard

curve, (2) two-sided ANOVA tests with Tukey or Dunnett's ad-hoc post-tests for inter-

group comparisons for growth curves and ΔCq comparisons of RT-qPCR data, all at

95% confidence.

Acknowledgements

We thank M. Bonhivers, D. Robinson, P. McKean, K. Matthews, and K. Gull for providing various plasmids and antibodies. We gratefully acknowledge the UTechS Photonic BioImaging (Imagopole), C2RT, Institut Pasteur, supported by the French National Research Agency (France BioImaging; ANR-10–INSB–04; Investments for the Future). We are grateful to P. Bastin for his strong scientific and human support. We warmly thank M. Boshart, P. Bastin, and S. Bonnefoy for their critical reading of the manuscript.

Funding

This work was supported by the Institut Pasteur, the French Government Investissement d'Avenir programme - Laboratoire d'Excellence "Integrative Biology of Emerging Infectious Diseases" (ANR-10-LABX-62-IBEID) and the French National Agency for Scientific Research (projects ANR-14-CE14-0019-01 EnTrypa and ANR-18-CE15-0012 TrypaDerm). AML and AnC are funded by a Wellcome Senior Fellowship to AML (209511/Z/17/Z). None of these funding sources has a direct scientific or editorial role in the present study.

Author contributions

ECA and BR designed the study. ECA, JMTN, CT, AnC and AlC performed the experiments. ECA, JMTN, AnC and BR analyzed the data. ECA and BR wrote the manuscript. ECA, AML and BR discussed the manuscript and acquired the funding.

Competing interest

All authors declare no financial relationships with any organizations that might have an interest in the submitted work in the previous three years, no other relationships or

725   activities that could appear to have influenced the submitted work, and no other

726   relationships or activities that could appear to have influenced the submitted work.

 Figure captions

 Figure 1. Characterization of the *FLAM8::mNG FLAM8^{RNAi}* strain *in vitro* and *in*

 ***vivo* functional investigation in the mammalian host. A)** Expression of *FLAM8*

 mRNA assessed by RT-PCR by the comparative $\Delta\Delta C_T$ method in control, non-induced

 and induced *FLAM8::mNG FLAM8^{RNAi}* parasites (72 h). **B)** Immunofluorescence

 pictures of non-induced (upper panels) and induced (bottom panels) *FLAM8::mNG*

 FLAM8^{RNAi} BSF during 72 h. Methanol-fixed trypanosomes were stained with an anti-

 mNG antibody (green) and DAPI for DNA content (blue). The scale bars represent 5

 μm. **C)** Growth curves of control, non-induced and induced *FLAM8::mNG FLAM8^{RNAi}*

 BSF parasites. All cell lines received 1 μg tetracycline for 6 days. Control parasites are

 Lister 427 "Single Marker" BSF parasites that do not bear the pZJM-FLAM8 plasmid

 for RNAi silencing. Results represent the mean (± standard deviation, SD) of three

 independent experiments. **D)** Groups of 4 BALB/c mice were injected IP with either

 control, non-induced or induced *FLAM8^{RNAi}* BSF trypanosomes. One PBS-injected

 BALB/c animal was used as negative control. Representative normalized *in vivo*

 images of the bioluminescence radiance signal (in photons / second / cm^2 / steradian)

 emitted from BALB/c mice infected with control, non-induced and induced

 FLAM8::mNG FLAM8^{RNAi} parasites 4 days post-infection (non-infected technical

 control mice were negative for bioluminescence, not shown). RNAi silencing of *FLAM8*

 was maintained *in vivo* by the addition of doxycycline in sugared drinking water 48 h

 prior infection and until the end of the experiment. **E)** Number of parasites in the blood

 (intravascular, IV) of infected BALB/c mice during the course of the infection (5 days).

 F) Number of parasites in the extravascular compartment (extravascular, EV) of the

 same infected mice as in E). **G)** Dissemination of control, non-induced and induced

FLAM8::mNG FLAM8^RNAi parasites, measured over the entire animal body (in cm^2)

through the total bioluminescent surface, during the entire infection course. Results

represent means ± standard deviation (SD).

Figure 2. Characterization of Δ*FLAM8* null mutants *in vitro*. A)

Immunofluorescence pictures of parental, Δ*FLAM8* knockout and rescue pleomorphic

BSF parasites labelled with the anti-FLAM8 (green) and mAb25 (axoneme in magenta)

antibodies, DAPI staining for DNA content (blue). Scale bars show 5 μm. **B)** Growth

curve of one parental, three Δ*FLAM8* subclones and one rescue pleomorphic BSF

trypanosome cell lines over 6 consecutive days. **C)** Measurements of the flagellum

length based on the axonemal marker mAb25 profiles in parental, three Δ*FLAM8*

subclones and rescue parasites. No statistical differences were found. **D, E)** Motility

tracking analysis showing the average speeds (D) and linearity (E) of BSF cell lines in

matrix-dependent culture medium. No statistical differences were observed. The

number of parasites considered for quantifications (N) is indicated above graphs (C),

(D) and (E). Results represent the mean ± standard deviation (SD) of three

independent experiments. Statistical tests included one-way ANOVA and Tukey's ad-

hoc post-tests for multiple comparisons.

Figure 3. Functional investigations on the Δ*FLAM8* null mutants *in vivo* in the

mammalian host. Groups of 3 BALB/c mice were injected IP with either one parental,

three ΔF*LAM8* null subclones or one rescue strains. One PBS-injected BALB/c animal

was used as negative control. **A)** Normalized *in vivo* images of the bioluminescence

radiance intensity (in photons / second / cm^2 / steradian) emitted 8 days post-infection

in BALB/c mice infected with parental, three Δ*FLAM8* subclones or rescue parasites

(non-infected control mice C- were negative for bioluminescence). **B)** Total number of

parasites in the blood of infected mice (intravascular, IV) during the infection (4 weeks).

Statistically significant differences (p<0.01) are indicated with one, two or three

asterisks (*, **, ***) representing differences between the parental strain and one, two

or three ΔFLAM8 subclones, respectively. **C)** Total number of extravascular (EV)

trypanosomes in the same mice. Statistically significant differences between the

parental strain and ΔFLAM8 subclones are indicated as in B). **D)** Dissemination of the

parental, three ΔFLAM8 subclones and rescue parasite strains, measured over the

entire animal body (in cm^2) through the total surface of bioluminescent signal, during

the entire infection course. Statistically significant differences between the parental

strain and ΔFLAM8 subclones are indicated as described above. Results represent

means ± standard deviation (SD). Statistical tests included two-way ANOVA and

Tukey's ad-hoc post-tests for multiple comparisons.

Figure 4. The absence of *FLAM8* reduces extravascular trypanosome

dissemination. In a second experimental infection, groups of 3 BALB/c animals were

injected IP with either parental, three ΔF*LAM8* null subclones or rescue strains. One

PBS-injected BALB/c animal was used as negative control. **A)** Total number of

intravascular parasites (IV) during the infection period. **B)** Total number of

extravascular (EV) trypanosomes in the same mice. **C)** Dissemination of the parental,

three ΔFLAM8 subclones and rescue parasite strains, measured over the entire animal

body (in cm^2) through the total surface of bioluminescent signal, during the entire

infection course. Statistically significant differences (p<0.01) are indicated with one or

three asterisks (*, ***) representing differences between the parental strain and one or

three ΔFLAM8 subclones, respectively. **D)** Number of EV parasites per dissected

organ and strain obtained by RT-qPCR quantification according to normalized *Tubulin*

expression. **E)** Number of EV trypanosomes calculated as the average difference

between the number of parental, Δ*FLAM8* and rescue parasites in all organs vs.

intravascular trypanosomes by the RT-qPCR method. Statistical differences

(p<0.0001) according to one-way ANOVA and Dunnett's comparison tests.

Figure 5. Parasite differentiation is not impacted in *FLAM8*-depleted

trypanosomes. A) Representative immunofluorescence pictures of stumpy parasites

after *in vitro* differentiation from proliferative slenders of parental (left panel), three

Δ*FLAM8* subclones (middle panels) and rescue (right panel) parasite strains upon *in*

vitro treatment with a nucleotide 5'-AMP analog. Methanol-fixed parasites were

labelled with the anti-PAD-1 antibody (red), and DAPI staining for DNA content (blue).

Scale bars show 5 μm. **B)** Quantification of the proportion of stumpy trypanosomes in

all pleomorphic cell lines after *in vitro* differentiation. The number of parasites

considered for quantification (n) is indicated above the graph. No significant differences

were found (one-way ANOVA and Tukey's comparisons test). Results represent the

mean ± standard deviation (SD) of three independent experiments. **C)** Selected

immunofluorescence images of freshly *in vitro* differentiated procyclic cells.

Trypanosomes were doubly labelled with anti-FLAM8 (green) and mAb25 (axoneme

in magenta), DAPI staining showing DNA content in blue. Scale bars show 5 μm. **D)**

Upon differentiation, early procyclic forms (ePCF) of parental, three Δ*FLAM8*

subclones and rescue parasite strains were equally diluted, and their *in vitro* growth

assessed. Statistical differences (p<0,01) were observed only when comparing

parental and KO 2 ePCF trypanosomes (one-way ANOVA and Tukey's comparison

test). Results represent the mean ± standard deviation (SD) of four independent

experiments. **E)** Representative immunofluorescence images of naturally differentiated

stumpy trypanosomes from parental, one selected ∆*FLAM8* subclone and rescue

strains isolated from mouse blood during the first peak of parasitemia (first

experimental *in vivo* infection). Parasites were labelled with anti-PAD-1 antibody (red),

and DAPI staining for DNA content (blue). Scale bars show 5 µm. **F)** Relative

proportions of *PAD-1*-expressing parental, three ∆*FLAM8* subclones and rescue

strains by RT-qPCR on blood and dissected organs at day 23 of the second *in vivo*

challenge. *Tubulin* expression was used to normalize the *PAD-1* mRNA levels. **G)**

Average of whole-organism *PAD-1* expression referred to *Tubulin* normalization in

parental-, ∆*FLAM8*- and rescue-infected mice.

Figure 6. Parasites lacking *FLAM8* are impaired in transmigrating through

endothelial cells *in vitro*. A) Schematic representation of the trans-endothelial

migration assay, showing the transwell system (Boyden chamber) containing the upper

and the lower compartments, the monolayer of endothelial HUVEC cells and slender

trypanosomes seeded on the top of the chamber for 24 hours. **B)** After this period,

parasites within the upper and lower chambers were counted and values further used

to calculate the proportion of parental, three ∆*FLAM8* subclones and rescue parasites

that migrated through the endothelial monolayer into the lower compartment.

Trypanosome transmigration was compared between the FLAM8 mutant cell lines and

the parental reference line (100% of trans-endothelial migration) using the Generalised

Linear Model function in R with a Gaussian family function and proportion of

transmigration as the dependent variable. A probability value of $p < 0.001$ was

considered significant (*). Error bars show SD. **C)** Representative immunofluorescence

images of cultured slender trypanosomes from parental, ∆*FLAM8* subclone and rescue

strains. Parasites were labelled with anti-CARP3 antibody (green intensity

normalized), and DAPI staining for DNA content (blue). Scale bars show 5 µm.

Supporting information captions

Figure S1. Validation of the triple-reporter efficiency in monomorphic *FLAM8^RNAi* **parasite lines.** Linear correlation between the number of parasites and the bioluminescence (in p/s) emitted by monomorphic 427 *FLAM8::mNG FLAM8^RNAi* BSF overexpressing the triple reporter chimeric protein [20], acquired by the IVIS Spectrum imager. Parasites without the RNAi plasmid (control, C), non-treated with tetracycline (non-induced, NI) and treated with tetracycline (induced, I) are shown. Representative bioluminescent image of serial 1/2 dilutions performed in a 96-well plate (in photons / second / cm² / steradian). RNAi induction was triggered by the addition of 1 µg tetracycline and / or doxycycline for 72 h. Results represent the mean ± standard deviation (SD) of three independent experiments.

Figure S2. Molecular validation of the Δ*FLAM8* null mutant cell lines. A) Whole-genome sequencing results showing *FLAM8* wild-type allele (WT, upper panel), the partial loss of the 5' *FLAM8* ORF in Δ*FLAM8* knockout trypanosomes (middle panel; the black arrow is showing the absence of reads at the *FLAM8* 5' ORF in the knockout line); and the restoration of the full *FLAM8* gene in rescue parasites (bottom panel; the black arrow is showing the absence of reads within the *pac* cassette due to the insertion of the new construct bearing the *ble* resistance marker), relative to the number of reads per 100-nt read length. The presence of the correct antibiotic cassettes is shown for knockout and rescue parasites (right middle and bottom panels). Δ*FLAM8* knockout and rescue parasites also bear a construct for expression of a triple reporter (TR) as assessed by the detection of *bsd* reads. **B)** Schemes showing the structure of the *FLAM8* locus in wild-type parasites (upper scheme) and the integration

plan of the different cassettes for ∆*FLAM8* knockout (middle panel, *HYG*- and *PAC*-containing schemes) and Rescue parasites (lower panel *HYG*- and *BLE*-containing schemes). **C)** PCR confirmation of the successful integrations of all reporter cassettes. Primer pairs used for PCRs are indicated at the bottom of each line and correspond to those drawn on the schemes in B), along with the expected band sizes of the corresponding diagnostic PCR. BSD: blasticidin; HYG: hygromycin; PAC: puromycin; BLE: phleomycin.

Figure S3. *In vitro* and *in vivo* validation of the triple-reporter efficiency in pleomorphic ∆*FLAM8* mutant cell lines. A) Representative bioluminescent image (in photons / second) of serial 1/2 dilutions performed in a 96-well plate of pleomorphic parental, ∆*FLAM8* knockout subclones and rescue parasites overexpressing the triple reporter chimeric protein [20]. **B)** Linear correlation between the number of parasites and the emitted bioluminescence (in photons / second) acquired by the IVIS Spectrum imager. Results represent the mean ± standard deviation (SD) of three independent experiments. **C)** Representative ventral view images of mice infected with increasing amounts of parental trypanosomes (10^3, 10^4, 10^5, 10^6 and 10^7 parasites/animal) acquired with the IVIS Spectrum imager 2.5 hours after IP injection. **D)** *In vivo* standard curve showing the correlation between the number of injected parasites and the bioluminescent signal (in photons/second, R^2 = 1). The standard curve was further employed to calculate the total number of parasites present in infected animals through whole-body BLI signal.

References

1.   MacGregor P, Szoor B, Savill NJ, Matthews KR. Trypanosomal immune

evasion, chronicity and transmission: an elegant balancing act. Nat Rev Microbiol.

2012;10(6):431-8. Epub 2012/05/01. doi: 10.1038/nrmicro2779. PubMed PMID:

22543519.

2.   Smith TK, Bringaud F, Nolan DP, Figueiredo LM. Metabolic reprogramming

during the *Trypanosoma brucei* life cycle. F1000Res. 2017;6. Epub 2017/06/20. doi:

10.12688/f1000research.10342.2. PubMed PMID: 28620452; PubMed Central

PMCID: PMCPMC5461901.

3.   Rotureau B, Van Den Abbeele J. Through the dark continent: African

trypanosome development in the tsetse fly. Frontiers in cellular and infection

microbiology. 2013;3:53. Epub 2013/09/26. doi: 10.3389/fcimb.2013.00053. PubMed

PMID: 24066283; PubMed Central PMCID: PMC3776139.

4.   Capewell P, Cren-Travaille C, Marchesi F, Johnston P, Clucas C, Benson RA,

et al. The skin is a significant but overlooked anatomical reservoir for vector-borne

African trypanosomes. eLife. 2016;5. doi: 10.7554/eLife.17716. PubMed PMID:

27653219; PubMed Central PMCID: PMCPMC5065312.

5.   Caljon G, Van Reet N, De Trez C, Vermeersch M, Perez-Morga D, Van Den

Abbeele J. The Dermis as a Delivery Site of *Trypanosoma brucei* for Tsetse Flies.

PLoS Pathog. 2016;12(7):e1005744. doi: 10.1371/journal.ppat.1005744. PubMed

PMID: 27441553; PubMed Central PMCID: PMCPMC4956260.

6.   Trindade S, Rijo-Ferreira F, Carvalho T, Pinto-Neves D, Guegan F, Aresta-

Branco F, et al. *Trypanosoma brucei* Parasites Occupy and Functionally Adapt to the

Adipose Tissue in Mice. Cell Host Microbe. 2016. doi: 10.1016/j.chom.2016.05.002.

PubMed PMID: 27237364.

7.   Crilly NP, Mugnier MR. Thinking outside the blood: Perspectives on tissue-

resident *Trypanosoma brucei*. PLoS Pathog. 2021;17(9):e1009866. Epub 2021/09/17.

doi: 10.1371/journal.ppat.1009866. PubMed PMID: 34529724; PubMed Central

PMCID: PMCPMC8445408.

8.   Rotureau B, Subota I, Bastin P. Molecular bases of cytoskeleton plasticity during

the *Trypanosoma brucei* parasite cycle. Cell Microbiol. 2011;13(5):705-16. Epub

2010/12/17. doi: 10.1111/j.1462-5822.2010.01566.x. PubMed PMID: 21159115.

9.   Broadhead R, Dawe HR, Farr H, Griffiths S, Hart SR, Portman N, et al. Flagellar

motility is required for the viability of the bloodstream trypanosome. Nature.

2006;440(7081):224-7. Epub 2006/03/10. doi: nature04541 [pii]

10.1038/nature04541. PubMed PMID: 16525475.

10.   Kohl L, Robinson D, Bastin P. Novel roles for the flagellum in cell

morphogenesis and cytokinesis of trypanosomes. EMBO J. 2003;22(20):5336-46.

Epub 2003/10/09. doi: 10.1093/emboj/cdg518. PubMed PMID: 14532107.

11.   Tetley L, Vickerman K. Differentiation in *Trypanosoma brucei*: host-parasite cell

junctions and their persistence during acquisition of the variable antigen coat. J Cell

Sci. 1985;74:1-19. Epub 1985/03/01. PubMed PMID: 4030903.

12.   Langousis G, Hill KL. Motility and more: the flagellum of *Trypanosoma brucei*.

Nat Rev Microbiol. 2014;12(7):505-18. Epub 2014/06/17. doi: 10.1038/nrmicro3274.

PubMed PMID: 24931043.

13.   Rotureau B, Morales MA, Bastin P, Spath GF. The flagellum-MAP kinase

connection in Trypanosomatids: a key sensory role in parasite signaling and

development? Cell Microbiol. 2009;11(5):710-18. Epub 2009/02/12. doi: CMI1295 [pii]

10.1111/j.1462-5822.2009.01295.x. PubMed PMID: 19207727.

14.   Roditi I, Schumann G, Naguleswaran A. Environmental sensing by African

trypanosomes. Curr Opin Microbiol. 2016;32:26-30. Epub 2016/05/01. doi:

10.1016/j.mib.2016.04.011. PubMed PMID: 27131101.

15.   Shimogawa MM, Ray SS, Kisalu N, Zhang Y, Geng Q, Ozcan A, et al. Parasite

motility is critical for virulence of African trypanosomes. Sci Rep. 2018;8(1):9122. Epub

2018/06/16. doi: 10.1038/s41598-018-27228-0. PubMed PMID: 29904094; PubMed

Central PMCID: PMCPMC6002391.

16.   Subota I, Julkowska D, Vincensini L, Reeg N, Buisson J, Blisnick T, et al.

Proteomic analysis of intact flagella of procyclic *Trypanosoma brucei* cells identifies

novel flagellar proteins with unique sub-localisation and dynamics. Molecular & cellular

proteomics : MCP. 2014. Epub 2014/04/18. doi: 10.1074/mcp.M113.033357. PubMed

PMID: 24741115.

17.   Fort C, Bonnefoy S, Kohl L, Bastin P. Intraflagellar transport is required for the

maintenance of the trypanosome flagellum composition but not its length. J Cell Sci.

2016;129(15):3026-41. Epub 2016/06/28. doi: 10.1242/jcs.188227. PubMed PMID:

27343245.

18.   Bertiaux E, Morga B, Blisnick T, Rotureau B, Bastin P. A Grow-and-Lock Model

for the Control of Flagellum Length in Trypanosomes. Curr Biol. 2018. Epub

2018/11/20. doi: 10.1016/j.cub.2018.10.031. PubMed PMID: 30449671.

19.   Calvo-Alvarez E, Bonnefoy S, Salles A, Benson FE, McKean PG, Bastin P, et

al. Redistribution of FLAgellar Member 8 during the trypanosome life cycle:

Consequences for cell fate prediction. Cell Microbiol. 2021;23(9):e13347. Epub

2021/04/26. doi: 10.1111/cmi.13347. PubMed PMID: 33896083; PubMed Central

PMCID: PMCPMC8459223.

20. Calvo-Alvarez E, Cren-Travaille C, Crouzols A, Rotureau B. A new chimeric

triple reporter fusion protein as a tool for *in vitro* and *in vivo* multimodal imaging to

monitor the development of African trypanosomes and *Leishmania* parasites. Infect

Genet Evol. 2018. Epub 2018/01/18. doi: 10.1016/j.meegid.2018.01.011. PubMed

PMID: 29339220.

21. Bachmaier S, Giacomelli G, Calvo-Alvarez E, Vieira LR, Van Den Abbeele J,

Aristodemou A, et al. A multi-adenylate cyclase regulator at the flagellar tip controls

African trypanosome transmission. Nat Commun. 2022;13(1):5445. Epub 2022/09/17.

doi: 10.1038/s41467-022-33108-z. PubMed PMID: 36114198; PubMed Central

PMCID: PMCPMC9481589.

22. Rojas F, Silvester E, Young J, Milne R, Tettey M, Houston DR, et al.

Oligopeptide Signaling through TbGPR89 Drives Trypanosome Quorum Sensing. Cell.

2018. Epub 2018/12/07. doi: 10.1016/j.cell.2018.10.041. PubMed PMID: 30503212.

23. Goodwin LG. The pathology of African trypanosomiasis. Trans R Soc Trop Med

Hyg. 1970;64(6):797-817. Epub 1970/01/01. PubMed PMID: 5495630.

24. Biteau N, Asencio C, Izotte J, Rousseau B, Fevre M, Pillay D, et al.

Trypanosoma brucei gambiense Infections in Mice Lead to Tropism to the

Reproductive Organs, and Horizontal and Vertical Transmission. PLoS Negl Trop Dis.

2016;10(1):e0004350. Epub 2016/01/07. doi: 10.1371/journal.pntd.0004350. PubMed

PMID: 26735855; PubMed Central PMCID: PMCPMC4703293.

25. Carvalho T, Trindade S, Pimenta S, Santos AB, Rijo-Ferreira F, Figueiredo LM.

Trypanosoma brucei triggers a marked immune response in male reproductive organs.

PLoS Negl Trop Dis. 2018;12(8):e0006690. Epub 2018/08/16. doi:

10.1371/journal.pntd.0006690. PubMed PMID: 30110342; PubMed Central PMCID:

PMCPMC6093638.

26. Caljon G, Van Den Abbeele J, Sternberg JM, Coosemans M, De Baetselier P, Magez S. Tsetse fly saliva biases the immune response to Th2 and induces anti-vector antibodies that are a useful tool for exposure assessment. Int J Parasitol. 2006;36(9):1025-35. Epub 2006/06/17. doi: 10.1016/j.ijpara.2006.05.002. PubMed PMID: 16777113.

27. Camara M, Soumah AM, Ilboudo H, Travaille C, Clucas C, Cooper A, et al. Extravascular Dermal Trypanosomes in Suspected and Confirmed Cases of *gambiense* Human African Trypanosomiasis. Clin Infect Dis. 2021;73(1):12-20. Epub 2020/07/09. doi: 10.1093/cid/ciaa897. PubMed PMID: 32638003; PubMed Central PMCID: PMCPMC8246823.

28. Bargul JL, Jung J, McOdimba FA, Omogo CO, Adung'a VO, Kruger T, et al. Species-Specific Adaptations of Trypanosome Morphology and Motility to the Mammalian Host. PLoS Pathog. 2016;12(2):e1005448. Epub 2016/02/13. doi: 10.1371/journal.ppat.1005448. PubMed PMID: 26871910; PubMed Central PMCID: PMCPMC4752354.

29. Sun SY, Kaelber JT, Chen M, Dong X, Nematbakhsh Y, Shi J, et al. Flagellum couples cell shape to motility in *Trypanosoma brucei*. Proc Natl Acad Sci U S A. 2018;115(26):E5916-E25. Epub 2018/06/13. doi: 10.1073/pnas.1722618115. PubMed PMID: 29891682; PubMed Central PMCID: PMCPMC6042131.

30. De Niz M, Bras D, Ouarne M, Pedro M, Nascimento AM, Henao Misikova L, et al. Organotypic endothelial adhesion molecules are key for *Trypanosoma brucei* tropism and virulence. Cell Rep. 2021;36(12):109741. Epub 2021/09/23. doi: 10.1016/j.celrep.2021.109741. PubMed PMID: 34551286; PubMed Central PMCID: PMCPMC8480282.

31.   Shaw S, DeMarco SF, Rehmann R, Wenzler T, Florini F, Roditi I, et al. Flagellar

cAMP signaling controls trypanosome progression through host tissues. Nat Commun.

2019;10(1):803. Epub 2019/02/20. doi: 10.1038/s41467-019-08696-y. PubMed PMID:

30778051; PubMed Central PMCID: PMCPMC6379439.

32.   Shaw S, Knusel S, Abbuhl D, Naguleswaran A, Etzensperger R, Benninger M,

et al. Cyclic AMP signalling and glucose metabolism mediate pH taxis by African

trypanosomes. Nat Commun. 2022;13(1):603. Epub 2022/02/03. doi: 10.1038/s41467-

022-28293-w.   PubMed   PMID: 35105902;   PubMed   Central   PMCID:

PMCPMC8807625.

33.   Le Ray D, Barry JD, Easton C, Vickerman K. First tsetse fly transmission of the

"AnTat" serodeme of *Trypanosoma brucei*. Ann Soc Belg Med Trop. 1977;57(4-5):369-

81. Epub 1977/01/01. PubMed PMID: 610616.

34.   Bohringer S, Hecker H. Quantitative ultrastructural differences between strains

of the *Tryponasoma brucei* subgroup during transformation in blood. J Protozool.

1974;21(5):694-8.   Epub   1974/11/01.   doi: 10.1111/j.1550-7408.1974.tb03731.x.

PubMed PMID: 4449091.

35.   Hirumi H, Hirumi K. Continuous cultivation of *Trypanosoma brucei* blood stream

forms in a medium containing a low concentration of serum protein without feeder cell

layers. J Parasitol. 1989;75(6):985-9. Epub 1989/12/01. PubMed PMID: 2614608.

36.   Barquilla A, Saldivia M, Diaz R, Bart JM, Vidal I, Calvo E, et al. Third target of

rapamycin complex negatively regulates development of quiescence in *Trypanosoma*

brucei.   Proc   Natl   Acad   Sci   U   S   A.   2012;109(36):14399-404.   doi:

10.1073/pnas.1210465109.   PubMed   PMID: 22908264;   PubMed   Central   PMCID:

PMCPMC3437835.

37.   Czichos J, Nonnengaesser C, Overath P. *Trypanosoma brucei*: cis-aconitate

and temperature reduction as triggers of synchronous transformation of bloodstream

to procyclic trypomastigotes in vitro. Exp Parasitol. 1986;62(2):283-91. Epub

1986/10/01. doi: 10.1016/0014-4894(86)90033-0. PubMed PMID: 3743718.

38.   Doyle JJ, Hirumi H, Hirumi K, Lupton EN, Cross GA. Antigenic variation in

clones of animal-infective *Trypanosoma brucei* derived and maintained in vitro.

Parasitology.   1980;80(2):359-69.   Epub   1980/04/01.   doi:

10.1017/s0031182000000810. PubMed PMID: 6102753.

39.   Wang Z, Morris JC, Drew ME, Englund PT. Inhibition of *Trypanosoma brucei*

gene expression by RNA interference using an integratable vector with opposing T7

promoters.   J   Biol   Chem.   2000;275(51):40174-9.   Epub   2000/10/03.   doi:

10.1074/jbc.M008405200. PubMed PMID: 11013266.

40.   Wirtz E, Leal S, Ochatt C, Cross GA. A tightly regulated inducible expression

system for conditional gene knock-outs and dominant-negative genetics in

Trypanosoma brucei. Mol Biochem Parasitol. 1999;99(1):89-101. Epub 1999/04/24.

PubMed PMID: 10215027.

41.   Redmond S, Vadivelu J, Field MC. RNAit: an automated web-based tool for the

selection of RNAi targets in *Trypanosoma brucei*. Mol Biochem Parasitol.

2003;128(1):115-8. Epub 2003/04/23. doi: 10.1016/s0166-6851(03)00045-8. PubMed

PMID: 12706807.

42.   Burkard G, Fragoso CM, Roditi I. Highly efficient stable transformation of

bloodstream forms of *Trypanosoma brucei*. Mol Biochem Parasitol. 2007;153(2):220-

3.   Epub   2007/04/06.   doi: 10.1016/j.molbiopara.2007.02.008.   PubMed   PMID:

17408766.

43.   Kelly S, Reed J, Kramer S, Ellis L, Webb H, Sunter J, et al. Functional genomics

in *Trypanosoma brucei*: a collection of vectors for the expression of tagged proteins

from endogenous and ectopic gene loci. Mol Biochem Parasitol. 2007;154(1):103-9.

Epub 2007/05/22. doi: S0166-6851(07)00099-0 [pii]

10.1016/j.molbiopara.2007.03.012. PubMed PMID: 17512617.

44.   Rotureau B, Ooi CP, Huet D, Perrot S, Bastin P. Forward motility is essential for

trypanosome infection in the tsetse fly. Cell Microbiol. 2014;16(3):425-33. Epub

2013/10/19. doi: 10.1111/cmi.12230. PubMed PMID: 24134537.

45.   Dacheux D, Landrein N, Thonnus M, Gilbert G, Sahin A, Wodrich H, et al. A

MAP6-related protein is present in protozoa and is involved in flagellum motility. PLoS

One. 2012;7(2):e31344. Epub 2012/02/23. doi: 10.1371/journal.pone.0031344.

PubMed PMID: 22355359; PubMed Central PMCID: PMCPMC3280300.

46.   Dean S, Marchetti R, Kirk K, Matthews KR. A surface transporter family conveys

the trypanosome differentiation signal. Nature. 2009;459(7244):213-7. Epub

2009/05/16. doi: nature07997 [pii]

10.1038/nature07997. PubMed PMID: 19444208.

1089

Fig1

Fig2

Fig3

Fig4

Fig5

Fig6