Gene expression plasticity followed by genetic change during

colonization a high-elevation environment

Huishang She[1], Yan Hao[1], Gang Song[1], Xu Luo[3], Fumin Lei[1,2,4], Weiwei Zhai[1,2,4], Yanhua Qu[1,2*]

[1] Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy

of Sciences, Beijing, China

[2] College of Life Sciences, University of Chinese Academy of Sciences, Beijing, China

[3] Faculty of Biodiversity and Conservation, Southwest Forestry University, Kunming, China

[4] Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming

650223, China

*, corresponding author

Abstract

Phenotypic plasticity facilitates organismal invasion of novel environments, and the resultant phenotypic change may later be modified by genetic change, so called "plasticity first". Herein we quantify gene expression plasticity and regulatory adaptation in a wild bird (Eurasian Tree Sparrow) from its original lowland (ancestral stage), experimentally implemented hypoxia acclimation (plastic stage) and colonized highland (colonized stage). Using a group of co-expressed genes from the cardiac and flight muscle, respectively, we demonstrate that gene expression plasticity to hypoxia tolerance is more often reversed than reinforced at the colonized stage. By correlating gene expression change with muscle phenotypes, we show that colonized tree sparrows reduce maladaptive plasticity that largely associates with decrease hypoxia tolerance. Conversely, adaptive plasticity that is congruent with increase hypoxia tolerance is often reinforced in the colonized tree sparrows. Genes displaying large levels of reinforcement or reversion plasticity (*i.e.*, 200% of original level) show greater genetic divergence between ancestral and colonized populations. Overall, our work demonstrates that gene expression plasticity at the initial stage of high-elevation colonization can be reversed or reinforced through selection-driven adaptive modification.

Key words: Gene expression, plasticity, acclimation, genetic variation, plasticity reinforcement, plasticity reversion

Introduction

Understanding the interactions between organisms and their environments is becoming increasingly important as environmental change and human activity change the original distribution of many species (Visser 2008; Wingfield et al. 2011; McCairns et al. 2016; Oostra et al. 2018). While some organisms can easily cope with environmental change, others run the risk of local or global extinction. An organismal response to environmental change partially depends on its ability of phenotypic change (Pfennig et al. 2010; Scoville & Pfrender 2010; Murren et al. 2015; Fox et al. 2019). For organisms that have recently colonized a new environment, phenotypic change may involve two stages. At the early stage, species may rapidly change their phenotype through plasticity without involving genetic changes. Under a persisting strong selective pressure, selection may bring genetic change to canalize phenotypic changes. Consequently, phenotypic plasticity has been considered to play an important stepping stone for genetically based evolutionary change at a late stage, so called "plasticity first" hypothesis (*i.e.*, West-Eberhard 2003; Schwander & Leimar 2011; Levis & Pfennig 2016; Corl et al. 2018).

Under this hypothesis, plasticity may serve adaptive evolution in two alternative ways (Ho & Zhang 2018; Ho et al. 2020). On the one hand, if plasticity changes the phenotype in the same direction as adaptive evolution does, genetic variation can strengthen the change towards an optimal phenotypic value in the new environment (reinforcement plasticity). On the other hand, if the plastic change works in the opposite direction as that driven by adaptive evolution, the subsequent genetic change would have to revert the initial plastic response (reversion/reduction plasticity). As previous studies on morphological and physiological traits have revealed mixed evidence for the reinforcement and the reversion of plasticity (Pigliucci et al. 2006; Lande 2009; Pfennig et al. 2010; Moczek et al. 2011; Levis & Pfennig 2016; Corl et al. 2018), it remains unclear if and how genetic change acts to reinforce or reverse plasticity in the wild species (*i.e.*, Campbell-Staton et al. 2021). The potential challenges are how to quantify phenotypic change relative to plasticity and adaptive evolution, and how to define what genes are involved in the phenotypic change, especially

as many phenotypes are likely polygenically mediated (Novembre & Barton 2018; Pritchard

et al. 2010).

Comparison of gene expression provides a potential approach to quantify the relative

contributions of phenotypic plasticity and genetic variation since gene expression can bridge

an organisms' genotype to its cellular biology and, by extension, higher order physiological

processes (Wray et al. 2003; Carrol 2008; Morris et al. 2014). While entire transcriptional

program is orchestrated by the gene regulatory network, it is possible to trace the genes that

are co-expressed or underpin the phenotypic change (Fukao et al. 2011). In addition, gene

expression evolves in a stabilizing manner, where regulatory elements accumulate mutations

that keep gene expression at an optimal level for physiological functions (Coolon et al. 2014;

Gilad et al. 2006; Hodgins-Davis et al. 2015). Herein we used gene expression data to

quantify regulatory plasticity and adaptation, as well as the relevant genetic changes in a

group of wild birds that have recently colonized a highland environment, the Eurasian Tree

Sparrows, *Passer montanus* (hereafter referred as tree sparrows).

The tree sparrow is a human commensal that has spread over a wide variety of habitats in

the Eurasian continent, and that has been successfully introduced to Australia and North

America (Widmann 1889; Graham et al. 2011). This species colonized the Qinghai-Tibet

plateau approximately 2600 years ago (Qu et al. 2020), possibly concurrently with the

introduction of barley agriculture in the highland. The Qinghai-Tibet plateau is a harsh

environment with an average elevation of 4500 m above sea level (m a.s.l.) where only a few

endemic animals can survive. For mammals and birds living at high elevations, hypoxia is one

of the strongest selective pressures that drive physiological changes (Barve et al. 2016;

McClelland & Scott, 2019). Compared to the lowland ancestral population, highland tree

sparrows have evolved physiological and gene expression change for hypoxia tolerance (Sun

et al. 2016; Qu et al. 2020), but how these changes relate to plasticity and adaptive evolution

are largely unknown.

In order to explore this, we acclimated a group of lowland birds to an experimentally

implemented hypoxic condition similar to that where we collected highland birds (*i.e.*, 3200

m a.s.l). We could assess gene expression plasticity and regulatory adaptation in hypoxia

tolerance by studying tree sparrows collected a) in their original lowland environment

(representing the physiologically ancestral stage), b) shortly after their exposure to the new

environmental stimulus (representing the ancestral plasticity at the initially plastic stage),

and c) after having adapted to the highland environment (the colonized stage). Using the

framework described in Figure 1, we quantified gene expression change stemming from

hypoxia acclimation and highland colonization and defined genes with the expression

plasticity being reinforced or reversed at the colonization stage. We test whether these

genes increase genetic variation between ancestral and colonized populations using a

permutation test. We use this framework in four datasets including groups of co-expressed

genes associated to hypoxia tolerance and those correlated to muscle phenotypes for the

cardiac and flight muscle, respectively. These independent comparisons congruently show

that gene expression at the plastic stage is more often reversed than reinforced at the

colonized stage, and the selection-driven genetic change depends on the magnitude of

reinforcement and/or reversion plasticity.

Results

Co-expressed genes involved in hypoxia tolerance generally reverse gene expression

plasticity

We sequenced transcriptomic data of the cardiac and flight muscles from the lowland tree

sparrows that had been kept under the hypoxia acclimation in this study (n=7), and

re-analyzed transcriptomic data in the lowland and highland tree sparrows collected by Qu et

al. (2020)(n=14, Supplementary Table 1). We focused on the flight and cardiac muscles

because they affect thermogenic capacity and oxygen transport and delivery and are thus

critical for the highland adaptation (Scott et al. 2015; Storz et al. 2010). Considering that

response to environmental change often involves complex coordinated biological response,

such as the co-regulation of genes underpinning physiological functions, we analyzed the

resultant expression data for all individuals using weighted gene correlation network analysis

(WGCNA, Langfelder & Horvath 2008). Weighted correlation network is a system biology

method for identifying groups of highly co-expressed genes (modules), summarizing

module-level expression and identifying important genes within modules (hub genes).

Consequently, these co-expressed genes can capture expression-stage correlations due to

the hypoxia-dependent expression. We identified six and nine modules of co-expressed

genes with their expression changes that are significantly associated with the three stages

(*i.e.*, ancestral, plastic and colonized stages) for the flight and cardiac muscle, respectively

(Figure 2a and Supplementary Figure S1). Across these modules, we identified 2413 and 508

hub genes based on the gene significance function (GS>1st quartile and *P*<0.05) for the flight

and cardiac muscles, respectively (Figure 2b-c).

Using these candidate genes we examined how gene expression plasticity changed across

the three stages. We compared the gene expression levels between the birds at the ancestral

stage and those at the plastic stage, as well as between the birds at the ancestral stage and

those at the colonized stage. We attributed genes showing the same direction of change to

the reinforcement group and those showing opposite direction of change to the reversion

group as described in Ho and Zhang (2018). Using a threshold of >50% increase/reduction of

gene expression between two comparisons, we attributed 36 and 15 genes to the

reinforcement group and 209 and 60 genes to the reversion group for flight and cardiac

muscle, respectively. Interestingly, we found fractions of genes with reversion plasticity are

larger than those of genes with reinforcement plasticity (Figure 2d). To confirm the

robustness of this observation with respect to random sampling errors, we carried out a

parametric bootstrap procedure as described in Ho and Zhang (2019), which aimed to

identify genes resulting from genuine differences rather than random sampling errors.

Bootstrap results also confirmed that genes exhibiting reversing plasticity significantly

outnumber those exhibiting reinforcing plasticity (Figure 2d).

We then explore whether this result is robust with different magnitudes of plasticity by using

a range of thresholds of <150% and >100%, <200% and >150%, and >200% to attribute gene

to reinforcement and reversion groups. These different categories consistently showed an

excess of genes with reversion plasticity (six out of eight comparisons, two-tailed binomial

test, $P<0.05$, Fig. 2e-f). To further confirm the robustness of the discovered pattern against

the arbitrariness of threshold selection, we also adopted different categorizations according

the magnitude of plasticity (*i.e.*, 20%, 40% and 60% bin settings along the spectrum of the

reinforcement/reversion plasticity), and these analyses showed similar results

(Supplementary Figure S2). Altogether, these results suggest that high-elevation colonization

of tree sparrows generally reverses gene expression plasticity.

If the reinforcement and/or reversion gene expression plasticity in hypoxic tolerance were

the target of selection in the high-elevation environment, we would expect to see an

increase in the genetic divergence between the ancestral and colonized populations. To

evaluate this we searched for genomic signatures of selection using resequencing data from

12 lowland and 11 highland tree sparrows generated from a previous study (Qu et al. 2020).

Significantly elevated genetic divergence (*i.e.*, F_{ST}) within the candidate genes beyond the

genic background would provide evidence for selection acting on this group of co-expressed

genes (*i.e.*, polygenic adaptation). We thus calculated average F_{ST} for the genes with

reinforcement or reversion plasticity using all SNPs included in the genic region, 2kb

up-stream and down-stream regions, respectively. We considered the up-stream and

down-stream regions because gene regulation may refer to genetic variation in nearby

non-coding regions. By comparing to the F_{ST} distributions generated from 100 random

samplings of the same number of SNPs from the genic background (see Methods), we found

that the empirical F_{ST} values significantly increased in the genes with the levels of

reinforcement/reversion plasticity reaching above 200% ($P < 0.05$, Figure 2e-f). In all cases,

the empirical F_{ST} values increased significantly in the 2kb up-stream and/or down-stream

regions of the candidate genes, but not in the genic regions. This observation thus suggests

that selection-driven genetic change is more targeted on the non-coding regions and

depends on the magnitude of reinforcement/reversion plasticity.

Gene expression and muscle phenotype analyses show adaptive plasticity and maladaptive

plasticity

The evolutionary outcome of the phenotypic plasticity for a population colonizing a novel

environment depends on the direction of plastic change with respect to the local optimum

of novel environment. If the plastic change is close to the local optimum, *i.e.*, adaptive

plasticity, natural selection likely reinforces adaptive plasticity. Conversely, if the plasticity

moves the phenotypic change away from the local optimum, *i.e.*, maladaptive plasticity,

natural selection should reduce or reverse the reaction norm and restore the phenotypic

change back to the original ancestral values (Campbell-Staton et al. 2021). In order to

investigate phenotypic plasticity and its evolutionary consequence, we analyzed two

datasets that included gene expression and muscle phenotypes collected from the lowland

ancestral and highland colonized tree sparrows (Qu et al. 2020).

Based on the correlation analysis between gene expression and muscle phenotypic values of

the ancestral and colonized tree sparrows, we categorized the direction of

expression-phenotype correlation as positive or negative regulators for the muscle

phenotype-associated genes ($P < 0.05$, Figure 3a). We identified 2037 and 1866 muscle

phenotype-associated genes in the flight and cardiac muscle, respectively (Figure 3b-c). Out

of these genes we found 328 positive regulators and 1709 negative regulators for the flight

muscle, and 843 positive regulators and 1023 negative regulators for cardiac muscle

(Supplementary Figure S3). We then characterized the differences in gene expression

observed for the ancestral-plastic groups and the ancestral-colonized groups as congruent or

incongruent with increased hypoxia tolerance (see Methods). Because the muscle

phenotypic changes observed in the colonized tree sparrows have been shown to enhance

oxygen delivery and improve metabolic capacity in the highland animals (Giordano 2005;

Storz et al. 2010; Scott 2011; Scott et al. 2015), we deemed that the gene expressions in

which the direction of plastic change matched the expectation for increase hypoxia

tolerance as putatively adaptive, while those that opposed this expectation were considered

to be putatively maladaptive (Figure 3a). Using a threshold of >50% of plasticity change, we

found that the tree sparrows displayed more genes with maladaptive plasticity than genes

with adaptive plasticity (two-tailed binomial test, flight muscle, 198 vs. 48, $P<2.2e^{-16}$, cardiac

muscle 186 vs. 166, $P<0.05$). These results are robust with random sampling errors as our

parametric bootstrap analyses also revealed similar results (Figure 3d).

We subsequently examined expression-stage relationship using the thresholds (<150%

and >100%, <200% and >150%, and >200%) and bin settings (20%, 40% and 60% bin settings)

as above-mentioned to explore the magnitudes of adaptive and maladaptive plasticity. We

found that the fractions of genes with maladaptive plasticity were larger than those of genes

with adaptive plasticity (two tailed binomial test, $P<0.01$, Figure 3d). This observation is

robust with different thresholds and bin settings for the flight muscle, but slightly sensitive to

bin setting for the cardiac muscle, indicating by more than 50% of comparisons supporting

an excess of genes with maladaptive plasticity (Supplementary Figure S4). These results

suggest that gene expression plasticity induced by hypoxia exposure at the early stage may

be mainly maladaptive for muscle physiological performance in highland environment.

The observed adaptive and maladaptive gene expressions in the tree sparrows suggest that

selection may have acted to reinforce and/or reverse gene expression plasticity. We then

quantified the genetic divergence of these genes between ancestral and colonized tree

sparrows by testing for statistically significant increase of F_{ST} as compared to the F_{ST}

distributions generated from 100 random samplings (see Methods). We found that the

empirical F_{ST} values were significantly larger in the genes within the <200% and >150%

category (cardiac muscle, $P<0.05$) and >200% category (flight and cardiac muscle, $P<0.05$),

but not in those within other categories with low levels of reinforcement and reversion

plasticity. Additionally, we found that genetic divergence between the ancestral and

colonized tree sparrows tends to increase in the 2kb up-stream and down-stream regions of

these genes ($P < 0.05$, Figure 3e-f). These results suggest that selection on regulatory

adaptation (*i.e.*, genetic divergence) may only be expected if large strength of selection is

needed to bring initial plasticity toward adaptive optimum of the novel environment.

Discussion

An understanding of phenotypic plasticity and its consequence for adaptive evolution is

important in evolutionary biology but remains challenging (Ghalambor et al. 2007; 2015;

Campbell-Staton et al. 2021; Kenkel & Matz 2017; Rivera et al. 2021). Using an integrative

approach of field-based studies, experimentally implemented acclimation experiment,

multiomic and muscle phenotypic data, our study demonstrates the importance of gene

expression plasticity in facilitating initial survival and population persistence at early stage of

colonization. In particular, our results provide novel insight into an issue of active debate

within evolutionary biology (*i.e.*, reinforcement vs. reversion plasticity) by showing that the

evolutionary consequence of plasticity depends on the adaptive plasticity (reinforcement) or

maladaptive plasticity (reversion). We also demonstrate that natural selection drives genetic

divergence when the magnitude of reinforcement and/or reversion plasticity becomes

intense, providing novel insights into the mechanism of plasticity first and gene as followers

(Schwander & Leimar 2011).

Our gene expression and sequence analyses provide several lines of evidence in support for

gene expression plasticity contribution to the high-elevation colonization of tree sparrows.

First, we show that co-expressed genes and muscle phenotype associated genes change

either the intensity or the direction of their expression profiles, and that gene expressions at

the plastic stage are more often reversed than reinforced at the colonization stage. Second,

by combining gene expression and muscle phenotype, we show that selection operates

intensely on maladaptive plasticity during the initial stage of responding novel environment

(Ghalambor et al. 2007; 2015; Ho and Zhang 2018; Campbell-Staton et al. 2021). Third, the

genes with intensified levels of reinforcement and reversion expression plasticity show large

genetic divergence (F_{ST}) in their up-stream and down-stream regions between the ancestral

and colonized populations, suggesting that selection has driven genetic changes in the

noncoding regions after the tree sparrows had colonized the novel environment.

These observations thus support that plasticity serves as a stepping stone in a successful

colonization of a new environment and that genetic changes that evolve at a late stage will

modify the plastic change to an optimal phenotype in the new environment (Corl et al. 2018;

Schwander & Leimar, 2011). It is interesting to think about why phenotypic change at the

plastic stage sometimes differs from those at the colonized stage. A likely reason is that

initially plastic change allows organisms to survive upon a sudden environmental shift but

the fitness is much reduced compared to that after a long-term colonization to the novel

environment (Fischer et al. 2016; Huang & Agrawal 2016; Leonard & Lancaster 2020; Kuo et

al. 2023). Thus, the overall physiological states of the organisms right after an environmental

shift may result in low fitness. To coordinate the phenotype values to close the organismal

fitness, a genetic change may be required modifying plastic change of phenotype to

approach the optimum to the novel environment (Storz 2021). Thus, the consequence of

genetic change on plasticity is dependent on the cost of plastic change (Ho and Zhang 2018;

Corl et al. 2018).

A limitation of our study is that we could only consider hypoxic condition in the acclimation

experiment, while also other factors associated with high-elevation environments, *e.g.*, low

temperature and UV radiation, may be of importance. Although we assume that hypoxia is

the dominant selective force in the Qinghai-Tibet Plateau (see also Sun et al. 2016; Qu et al.

2020), the use of a single environmental stimulus makes our conclusions more conservative.

Also, it is reasonable to assume that plastic change in organisms is time-dependent and that

our results cannot catch gene expression occurring at later stage of plastic response (*i.e.*,

after one month). Since we could only run the acclimation experiment for one month it is

likely we have not identified all genes related to hypoxia tolerance. Despite this, we could

observe a clear pattern of reinforcement and reversion plasticity and the associated genetic

changes. Our sample sizes are rather small (n=12 for the flight muscle and n=9 for the cardiac

muscle), mostly because of the logistical challenge of keeping tree sparrows in the necessary

common garden experiments. In order to compensate for this, we utilized comparative

transcriptomics to generate the four independent datasets (a group of co-expressed genes

and a group of phenotype-associated genes for the flight and cardiac muscles, respectively).

As the analyses of these four independent datasets show similar results, we believe the

conclusions drawn from our study are robust.

Despite some limitations, our study demonstrates an easy implemented framework for

quantifying phenotype plasticity and how to relate this to genetic change. Short-term

acclimation response of lowland populations has been found to be different from the

populations adapted to the high-elevation environments. For example, people newly

exposed to high-elevation environment increase their red blood cell as compared to people

at sea level, which is likely mediated by *HIF* and *EPO* genes in hypoxia response cascade

(Erzurum et al. 2007; Peng et al. 2017). This physiological change, however, is less activated

in native people at high-elevation when compared to people at sea level, demonstrating an

example of reversion of phenotypic plasticity (Storz 2021). Together with previous studies,

our work shows that short-term acclimation response and long-term adaptive evolution in

hypoxia response may affect different evolutionary pathways. Our work provides

understanding how plasticity facilitates species to invade new habitats and survive

environmental change. Such an understanding is important, in particular when numerous

species are being introduced to new regions of the globe through human intervention and

spread as invasive species (Visser 2008).

Materials and Methods

Sampling

We generated transcriptomic data from hypoxia-acclimated lowland tree sparrows by

exposing five lowland tree sparrows to hypoxic condition for 30 days using a hypoxic

chamber (Qu et al. 2020). The oxygen content in the chamber was set to 14% of the oxygen

content (simulating the oxygen concentration at 3200 m a.s.l., 70% of ~20%, latter of which is

the content of oxygen content at sea level), which is similar to condition where we collected

the highland tree sparrows (Qinghai Lake, 3200 m a.s.l.). We collected flight muscle from the

four individuals and cardiac muscle from three individuals. In addition, we extended the

analyses of the transcriptomic and muscle histological data from tree sparrows that were

collected from lowland (Beijing and Hebei, 100 m a.s.l.) and highland (Qinghai Lake, 3200 m

a.s.l., Figure 1) collected in previous study (Qu et al. 2020). In total, we used twelve samples

for flight muscle and nine samples for cardiac muscle (Supplementary Table 1). The ancestral

lowland and hypoxia-exposed lowland tree sparrows were collected from the same locality

(Beijing) and the same season (pre-breeding). Only adult birds with similar body weight

(approximately 18g) were used.

Transcriptome sequencing

RNA libraries were constructed for the flight and cardiac muscles and sequenced on an

Illumina HiSeq4000 platform. After filtering low-quality, adapter-contaminated, and N-rich

reads (>10%), a total of 134, 128 and 120 million reads of transcriptional data were

generated for the lowland, hypoxia-exposed lowland tree sparrows and highland tree

sparrows, respectively (Supplementary Table S1). We mapped cleaned reads against the tree

sparrow genome using STAR (Dobin et al. 2013). We compared log-transformed transcript

per million (TPM) across all genes and determined the most conserved genes (*i.e.*, coefficient

of variance ≤ 0.3 and average TPM ≥ 1 for each sample) for the flight and cardiac muscles,

respectively (Hao et al. 2023). We then compared the median expression levels of these

conserved genes and found no difference among the lowland, hypoxia-exposed lowland, and

highland tree sparrows (Wilcoxon signed-rank test, *P*<0.05, Supplementary Figure S5),

suggesting that batch effect had little influence on the transcriptomic data. We then used

TPM values to calculate gene expression level and intensity using RSEM (Li & Dewey 2011).

WGCNA analyses

We used WGCNA v. 1.41-1 (Langfelder and Horvath, 2008) to identify regulatory architecture

for the flight muscle and cardiac muscle transcriptomes. Specifically, we used a principal

component analysis (PCA) to summarize modules of genes expression with

blockwiseModules function, and then used module eigengene values of the first principal

component (PC) to test correlation between module expression and stages. We used Pearson

linear correlation (cor function) to identify genes with expression reinforcement and

nonlinear regression (polynomial regression model: Model = lm(y ~ poly(x, 5, raw = TRUE),

data=data) to identify genes with expression reversion. Regulatory modules were identified

as branches of the resulting cluster tree via the dynamic tree-cutting method and highly

correlated modules (*P*<0.1) were merged. We used "GS > 1st quartile of GS" and "*P*<0.05" to

identify important genes within modules (hub genes).

Identify gene regulation associated with muscle phenotype

To identify the genes with expression levels that correlated to evolved differences in the

muscle phenotypes in the tree sparrows, we used the lowland and highland individuals from

Qu et al. (2020) for which RNAseq and muscle phenotypic data were available. We correlated

the levels of gene expression with muscle phenotypes in the lowland and highland

individuals. We used three phenotypes for the flight muscle (capillary per fiber, fiber area

and perimeter) and two for the cardiac muscle (capillary density and fiber diameter). We

used Pearson correlation to test the association between each of the phenotypic traits and

expression profile of each gene (TPM). A threshold of *P*<0.05 was used to detect

muscle-associated genes. We then followed Campbell-Staton et al. (2021) to categorize

genes as positive (positively correlated with muscle phenotypes) or negative regulators

(negatively correlated with muscle phenotypes).

Subsequently, we compared the direction of gene expression plasticity at the plastic stage

with those of the positive and negative regulators. Briefly, the direction of gene expression

plasticity at the plastic stage matched the expectation for positive regulator (increased

expression) or negative regulator (decreased expression) in the colonized tree sparrows was

considered to indicate adaptive plasticity. Conversely, those that opposed this expectation

were considered to be maladaptive, *i.e.*, the direction of gene expression plasticity at the

plastic stage showed decreased expression for the positive regulators and increased

expression for negative regulators (Figure 3a).

Reinforcement and reversion plasticity analyses

To investigate the magnitude of reinforcement and reversion plasticity, we followed the

method described in Ho and Zhang (2018). Specifically, expression levels of each gene in the

three stages were treated as E_lowland (expression level at ancestral stage), E_hypoxia

(expression level at plastic stage) and E_highland (expression level at colonized stage). We

used a threshold of 50% of the ancestral gene expression level, *i.e.*, 50% * E_lowland, to

detect an excess of gene expression change. We identified genes with an excess of plastic

change if it satisfied the condition of |(E_hypoxia-E_lowland)|> 50% * E_lowland. Likewise,

we identified genes with an excess of evolutionary change if it met the condition of

|(E_highland-E_hypoxia)|> 50%* E_lowland. If the genes showed same direction in the

expression change at the plastic and colonized stages, we regarded these genes as

reinforcement expression. If the genes showed opposite direction in the expression change

at the plastic and colonized stages, we regarded these genes as reversion expression. We

categorized these genes to reinforcement and reversion groups and then tested if the

proportions of the genes in the two groups were significantly different from the expected

proportions using two-tailed binomial tests.

To explore whether an excess of genes with reversion plasticity was subject to random

sampling errors (Mallard et al. 2018), we used a parametric bootstrap method as described

in Ho and Zhang (2019). Specifically, we simulated the mean expression level of a gene at

each stage following a Gaussian distribution within the mean equal to the observed mean

expression of this gene at this stage and the standard deviation equal to the estimated

standard error. We then draw a random variable from the above Gaussian distribution to

represent an observation of the mean expression level of this gene at this stage. We drew

random variables representing the mean expression level of gene at each of the three stages

(*i.e.*, ancestral, plastic and colonized stages), and then computed gene expression changes at

plastic stage and colonized stage and determined the reinforcement or reversion plasticity.

This process is repeated 1000 times for each gene. If this gene exhibited reinforcement or

reversion plasticity at least 950 repeats (*P*<0.05), we considered this gene showing

reinforcement or reversion plasticity. The Gaussian distribution and random sampling were

generated using rnorm function of R.

To test the impact of the choice of thresholds on the results, we analyzed the data using a

range of thresholds, *i.e.*, <100% and >50%, <150% and >100%, <200% and >150%,

and >200%. This setting aimed to obtain independent genesets for subsequently genetic

divergence analyses, because a geneset selected by relax threshold (*i.e.*, 50%) also included

genesets obtained by the stringent threshold (*i.e.*, 100%, 150% and 200%). In addition, we

also implemented other categorization scheme to check the robustness of the results.

Specifically, we used three different bin settings (*i.e.*, 20%, 40% and 60% bin settings along

the spectrum of the reinforcement/reversion plasticity), to group genes according their

magnitude of plasticity. For each category in the three bin settings, we compared whether

the proportion of the genes with reversion plasticity differed from that with reinforcement

plasticity using two-tailed binomial tests.

Comparing genetic divergence for genes with reinforcement and reversion expression

plasticity

To calculate the genetic divergence between the lowland ancestral and highland colonized

tree sparrows, we used 12 lowland and 11 highland individuals for which re-sequencing data

were available (Qu et al. 2020). After mapping the raw reads to the tree sparrow genome

using BWA v0.7.17 (Li & Durbin 2009), we obtained mean sequence coverage of 17× for each

individual (Table S2). We called variants with GATK v 3.7 (McKenna et al. 2010) and Samtools

v1.2 (http://www.htskib.org/) and filtered single nucleotide polymorphism (SNP) using VCF

tools and GATK with minimum coverage = 138, root mean square mapping quality >= 20,

distance of adjacent SNPs >=5, distance to a gap >=5bp and read quality value >=30.

We calculated average F_{ST} using Vcftools (Danecek et al. 2011) with SNPs included in the

genic region, 2kb up-stream and down-stream regions of the genes within each of the four

categories, *i.e.*, <100% and >50%, <150% and >100%, <200% and >150%, and >200%. The

up-stream and down-stream regions were considered because gene regulation may refer to

genetic variation in the nearby genic regions. We permuted F_{ST} distributions for the genes

within each category by random sampling the same number of SNPs, allowing for a

fluctuation of 5% of total SNPs. We compared the empirical F_{ST} values to permuted F_{ST}

distributions generated from 100 samplings and considered a threshold of *P*<0.05 to be

statistically significant (empirical F_{ST}>95% percentile of permuted F_{ST} distribution). We

calculated empirical F_{ST} values and permuted F_{ST} distributions for the genic region, 2kb

up-stream and down-stream regions separately. To avoid the sampling bias (*i.e.*, sampling

SNPs from neutral regions), our SNP sampling was constrained to genic region, 2k up-stream

and down-stream regions of all genes (*i.e.*, 16925 genes). All analyses were conducted with

the R statistical software package (R Foundation for Statistical Computing 2017).

Data accessibility

DNA sequencing reads used in this study have been deposited in Short Read Archive under

the project number PRJNA417520.

Authors' contributions

Q.Y. designed research; S.H., Y.H., X.L., S.G. and Q.Y. performed research; F.L. provided

highland samples; Q.Y. and S.H. wrote the paper.

Acknowledgement

We acknowledge Ying Xiong for logistic work in the hypoxia-exposed experiment. This

research was funded by the Third Xinjiang Scientific Expedition and Research (XIKK)

(2022xjkk0205), Second Tibetan Plateau Scientific Expedition and Research (2019QZKK0501)

and the National Natural Science Foundation of China (NSFC32020103005).

Reference:

Barve S, Dhondt AA, Mathur VB. and Cheviron ZA. 2016. Life-history characteristics influence

physiological strategies to cope with hypoxia in Himalayan birds. *Proc. R. Soc. B.* 283:

20162201.

Campbell-Staton SC, Velotta JP, Winchell KM. 2021. Selection on adaptive and maladaptive

genes expression plasticity during thermal adaptation to urban heat islands. *Nat.*

Commun. 12: 6195.

Corl A, Bi K, Luke C, Challa SA, Stern JA, Sinervo B, Nielsen R. 2018. The genetic basis of

adaptation following plastic changes in coloration in a novel environment. *Cur. Biol.* 28:

2970–2977.

Coolon JD, McManus CJ, Stevenson KR, Graveley BR, Wittkopp PJ. 2014. Tempo and mode of

regulatory evolution in Drosophila. *Genome Res*. 24: 797–808.

Danecek P. et al. 2011. The variant call format and VCFtools. Bioinformatics 27, 2156–2158.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.

2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15–21.

Erzurum SC, et al. 2007. Higher blood flow and circulating NO products offset high-altitude

hypoxia among Tibetans. *Proc. Natl. Acad. Sci. U.S.A.* 104: 17593–17598.

Fischer EK, Ghalambor CK, Hoke KL. 2016. Can a network approach resolve how adaptive vs

nonadaptive plasticity impacts evolutionary trajectories? *Integr. Comp. Biol.* 56: 877–

888.

Fox RJ, Donelson JM, Schunter C, Ravasi T and Gaitán-Espitia JD. 2019. Beyond buying time:

the role of plasticity in phenotypic adaptation to rapid environmental change. *Phil. Trans.*

R. Soc. B. 374: 20180174.

Fukao T, Yeung E, Bailey-Serres J. 2011. The submergence tolerance regulator sub1a

mediates crosstalk between submergence and drought tolerance in rice. *The Plant Cell*

23:412–427.

Ghalambor CK, Hoke KL, Ruell EW, Fischer EK, Reznick DN, Hughes KA. 2015. Non-adaptive

plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature*

525:372–375.

Ghalambor CK, McKay JK, Carroll SP, Reznick, DN. 2007. Adaptive versus non-adaptive

phenotypic plasticity and the potential for contemporary adaptation in new

environments. *Funct. Ecol.* 21: 394–407.

Gilad Y, Oshlack A, Rifkin SA. 2006. Natural selection on gene expression. *Trends in Genetics*

22: 456–461.

Giordano FJ. 2005. Oxygen, oxidative stress, hypoxia, and heart failure. *J. Clin. Invest.* 115:

550–508.

Graham J, Harnevich C, Young N, Newman G, Stohlgren T. 2011. How will climate change

affect the potential distribution of Eurasian tree sparrows *Passer montanus* in North

America? *Curr. Zool.* 57:648–654.

Hao et al. 2023. Divergent contributions of coding and noncoding sequences to initial

high-altitude adaptation in passerine birds endemic to the Qinghai–Tibet Plateau. *Mol. Ecol.* Doi: 10.1111/mec.16942.

Ho WC, Zhang J. 2018. Evolutionary adaptations to new environments generally reverse plastic phenotypic changes. *Nat. Commun.* 9: 350.

Ho WC, Zhang J. 2019. Genetic gene expression changes during environmental adaptations tend to reverse plastic changes even after correction for statistical nonindependence. *Mol. Biol. Evol.* 36: 604–612.

Ho WC, Li D, Zhu Q, Zhang J. 2020. Phenotypic plasticity as a long-term memory easing readaptations to ancestral environments. *Sci. Adv.* 6: eaba3388.

Hodgins-Davis A, Rice DP, Townsend JP. 2015. Gene Expression Evolves under a House-of-Cards Model of Stabilizing Selection. *Mol. Biol. Evol.* 32:2130–2140.

Huang Y, Agrawal AF. 2016. Experimental evolution of gene expression and plasticity in alternative selective regimes. *PLoS Genet.* 12: 1–23.

Kenkel C, Matz M. 2017. Gene expression plasticity as a mechanism of coral adaptation to a variable environment. *Nat. Ecol. Evol.* 1: 0014.

Kuo KC, Yao CT, Liao BY, Weng MP, Dong F, Hsu YC, Hung CM. 2023. Weak gene-gene interaction facilitates the evolution of gene expression plasticity. *BMC Biol.* 21: 57.

Lande R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. *J. Evol. Biol.* 22: 1435143514

Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinform.* 9: 559.

Leonard AM, Lancaster LT. 2020. Maladaptive plasticity facilitates evolution of thermal tolerance during an experimental range shift. *BMC Evol. Biol.* 20: 1–11.

Levis NA, Pfennig DW. 2016. Evaluating 'plasticity-first' evolution in nature: key criteria and empirical approaches. *Trends Ecol. Evol.* 31: 563–574.

Li B, Dewey CN. 2011. RSEM: Accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform.* 12: 323.

Li H, Durbin R. 2009. Fast and accurate short read alignment with burrows-wheeler transform. *Bioinformatics* 25: 1754–60.

Mallard F, Jaksic AM, Schlotterer C. 2018. Contesting the evidence for nonadaptive plasticity.

Nature 555: E21–22.

McCairns RJS, Smith S, Sasaki M, Bernatchez L, Beheregaray LB. 2016. The adaptive potential

of subtropical rainbowfish in the face of climate change: heritability and heritable

plasticity for the expression of candidate genes. *Evol. Appl.* 9: 531–545.

McClelland GB, Scott R. 2019. Evolved mechanisms of aerobic performance and hypoxia

resistance in high-altitude natives. *Annu. Rev. Physiol*. 81: 561–583.

McKenna A. et al. 2010. The genome analysis toolkit: a map reduce framework for analyzing

next-generation DNA sequencing data. *Genome Res.* 20: 1297–303.

Moczek AP, Sultan S, Foster S, Ledultan SC, Dworkin I, Nijhout HF, Abouheif E, Pfennig DW.

2011. The role of developmental plasticity in evolutionary innovation. *Proc. Biol. Sci.* 278,

2705–2713.

Morris MRJ, Richard R, Leder EH, Barrett RDH, Aubin-Horth N, Rogers SM. 2014. Gene

expression plasticity evolves in response to colonization of freshwater lakes in

threespine stickleback. *Mol. Ecol.* 23, 3226–3240.

Murren CJ, et al. 2015. Constraints on the evolution of phenotypic plasticity: limits and costs

of phenotype and plasticity. *Heredity* 115: 293–301.

Novembre J, Barton NH. 2018. Tread lightly interpreting polygenic tests of selection. *Genetics*

208: 1351–1355.

Oostra V, Saastamoinen M, Zwaan B, Wheat C, 2018. Strong phenotypic plasticity limits

potential for evolutionary responses to climate change. *Nat. Commun.* 9:1005.

Peng Y, et al. 2017. Down-regulation of EPAS1 transcription and genetic adaptation of

Tibetans to high-altitude hypoxia. *Mol. Biol. Evol.* 34:818–830.

Pfennig DW, Wund MA, Snell-Rood EC, Cruickshank T, Schlichting CD, Moczek AP. 2010.

Phenotypic plasticity's impacts on diversification and speciation. *Trends Ecol. Evol.* 25:

459–467.

Pigliucci M, Murren CJ, Schlichting CD. 2006. Phenotypic plasticity and evolution by genetic

assimilation. J. Exp. Biol. 209: 2362–2367.

Pritchard JK, Pickrell JK, Coop G. 2010. The genetics of human adaptation: hard sweeps, soft

sweeps, and polygenic adaptation. *Curr. Biol.* 20: R208–215.

Qu Y, et al. 2020. Rapid phenotypic evolution with shallow genomic differentiation during
early stages of high elevation adaptation in Eurasian Tree Sparrows. *Nat. Sci. Rev.*, 7:
113.

R Foundation for Statistical Computing. 2017. R: A Language and Environment for Statistical
Computing, https://www.r-project.org.

Rivera HE, Aichelman HE, Fifer JE, Kriefall NG, Wuitchik DM, Wuitchik SJS, Davies SW. 2021. A
framework for understanding gene expression plasticity and its influence on stress
tolerance. *Mol. Ecol.* 30: 1381-1397.

Schwander T, Leimar O. 2011. Genes as leaders and followers in evolution. *Trends Ecol. Evol.*
26: 143–151.

Scott GR. 2011. Elevated performance: the unique physiology of birds that fly at high
altitudes. *J. Exp Biol.* 214:2455–2462.

Scott GR, Elogio TS, Lui MA, Storz JF, Cheviron ZA. 2015. Adaptive modifications of muscle
phenotype in high-altitude deer mice are associated with evolved changes in gene
regulation. *Mo. Biol. Evol.* 32: 1962–1976.

Scoville AG, Pfrender ME. 2010. Phenotypic plasticity facilitates recurrent rapid adaptation to
introduced predators. *Proc. Natl. Acad. Sci. U.S.A.* 107: 4260–4263

Storz JF, Scott GR, Cheviron AZ. 2010. Phenotypic plasticity and genetic adaptation to
high-altitude hypoxia in vertebrates. *J. Exp. Biol.* 213:4125–4136.

Storz JF. 2021. High-altitude adaptation: mechanistic insights from integrated genomics and
physiology. *Mol. Biol. Evol.* 38: 2677–2691.

Sun YF, Ren ZP, Wu YF, Lei FM, Dudley R, Li DM. 2016. Flying high: limits to flight performance
by sparrows on the Qinghai-Tibetan Plateau. *J. Exp. Biol.* 219: 3642–3648.

Visser ME. 2008. Keeping up with a warming world; assessing the rate of adaptation to
climate change. *Proc. R. Soc. Lond. B.* 275: 649–649.

West-Eberhard MJ. 2003. Developmental Plasticity and Evolution, Oxford University Press.

Widmann O. 1889. History of the House Sparrow Passer domesticus and the European Tree
Sparrow Passer montanus at Saint Louis, Mo. Washington DC: U.S. Department of
Agriculture, Division of Economic Ornithology and Mammology.

Wingfield JC, Kelley JP, Angelier F, Chastel O, Lei F, Lynn SE, Miner B, Davis JE, Li DM, Wang G.

2011. Organism-environment interactions in a changing world: a mechanistic approach.

J. Ornithol. 152: S279–S288.

Wray GA, Hahn MW, Abouheif E, Balhoff JP, Pizer M, Rockman MV, Romano LA. 2003. The

evolution of transcriptional regulation in eukaryotes. *Mol. Biol. Evol.* 20:1377–1419.

Figure 1. A schematic representation of quantifying gene expression plasticity and genetic change in relevant genes. **Experimental design**, transcriptomic data of the flight and cardiac muscle were collected from tree sparrows obtained in their original lowland habitat (ancestral stage), experimental exposure to the hypoxic condition (plastic stage), and colonization to a high-elevation environment (colonized stage). **Identify candidate genes**, co-expressed genes that display correlation with hypoxia tolerance were identified by a WGCNA analysis and muscle phenotype associated genes were identified by correlating gene expression levels and each of the muscle phenotypes. **Reinforcement/reversion plasticity**, the genes with reinforcement or reversion plasticity were defined by comparing expression changes between the ancestral and plastic stages (plastic change), and between plastic and colonized stages (evolved change). The strength of reinforcement/reversion plasticity is measured by a range of thresholds, *i.e.*, <100% and >50%, <150% and >100%, <200% and >150%, and >200%. **Quantify genetic divergence**, genetic divergence (*i.e.*, F_{ST}) between lowland and highland populations was estimated for the genic region, 2kb up-stream and down-stream regions of the candidate genes. The empirical F_{ST} values were compared with permuted F_{ST} distributions generated from 100 random samplings from the genic background. A threshold of $P<0.05$ was used to determine statistical significance.

Figure 2. The WGCNA analysis identified co-expressed genes and reinforcement and reversion of expression plasticity. (a) Pearson linear correlation is used to identify genes with expression reinforcement and nonlinear regression to identify genes with expression reversion. Regulatory modules were identified as branches of the resulting cluster tree via the dynamic tree-cutting method and highly correlated modules ($P<0.1$) were merged. We used "GS > 1st quartile of GS" and "$P<0.05$" to identify important genes within modules (hub genes). (b) and (c) Six and nine modules respectively were identified to be associated with different stages for the flight and cardiac muscles ($P<0.01$). Within each module, colored dots show the genes with the expression levels significantly associated with the stages (gene significance function>1st quartile and $P<0.05$). (d) Frequencies of genes with reinforcement and reversion plasticity (>50%) and their subsets that acquire strong support in the parametric bootstrap analyses ($\geq$ 950/1000). (e) and (f) Left, genes with expression plasticity being reinforced (red) or reversed (blue) at the colonized stage identified for the flight (e) and cardiac muscles (f), respectively. There are more genes showing reversion plasticity than those showing reinforcement plasticity. Two-tailed binomial test, NS, nonsignificant; ***, $P < 0.001$. Right, the F_{ST} values (vertical lines) significantly increase in the 2kb up-stream and/or down-stream regions of the genes having the magnitude of reinforcement/reversion plasticity > 200%. Vertical lines, the empirical F_{ST} values; shades, permutated F_{ST} distributions generated from 100 random samplings. NS, non-significant, *, $P < 0.05$; **, $P < 0.01$.

Figure 3. Genes with expression levels associated with muscle phenotypes and adaptive and maladaptive plasticity. (a) Gene expression profiles were correlated with muscle phenotypes. The direction of gene expression plasticity at the plastic stage matched the expectation for positive regulator (increased expression) and negative regulators (decreased expression) in the colonized tree sparrows was considered to indicate adaptive plasticity. Conversely, those that opposed this expectation were considered to be maladaptive, *i.e.*, the direction of gene expression plasticity at the plastic stage showed decreased expression for the positive regulators and increased expression for negative regulators. A, ancestral stage; P, plastic stage; C, colonized stage. (b) and (c) Gene expression profiles were correlated with three flight muscle phenotypes (b, capillary per fiber, fiber area and perimeter) and two cardiac muscle phenotype (c, capillary density and fiber diameter). Red dots show genes with adaptive gene expression plasticity and blue dots show genes with maladaptive gene expression plasticity. (d) Frequencies of genes with adaptive and maladaptive plasticity (>50%) and their subsets that acquire strong support in the parametric bootstrap analyses (≥ 950/1000). (e) and (f) Left, candidate genes were classified to four categories with different magnitudes of adaptive or maladaptive plasticity. More genes show maladaptive plasticity than those show adaptive plasticity. Two-tailed binomial test, **, $P < 0.01$; ***, $P < 0.001$. Right, the empirical F_{ST} values significantly increased in the 2k up-stream and/or down-stream regions of the genes having adaptive/maladaptive plasticity <200% and >150% (cardiac muscle) and >200% (flight and cardiac muscles). (e) Flight muscle, (f) Cardiac muscle. Vertical lines, the empirical F_{ST} values; shades, permutated F_{ST} distributions generated from 100 times sampling. *, $P < 0.05$; **, $P < 0.01$.