Host immunosenescence compromises *Mycobacterium tuberculosis* clearance

Running Title: Older TB hosts have compromised T$_H$1 responses

Falak Pahwa[1], Shweta Chaudhary[1], Ashish Gupta[1], Shivam Chaturvedi[2], Ranjan Kumar Nanda[1,*]

[1]Translational Health Group, International Centre for Genetic Engineering and Biotechnology, New Delhi-110067, India

[2]Bio-experimentation Facility, International Centre for Genetic Engineering and Biotechnology, New Delhi-110067, India

*Corresponding author:

Ranjan Kumar Nanda (Ph.D.)

Group Leader

Translational Health Group

International Centre for Genetic Engineering and Biotechnology, New Delhi, India

Email: ranjan@icgeb.res.in

Abstract

Immunosenescence increases susceptibility to infectious diseases like tuberculosis (TB) in older subjects (≥60 years) and may impact containment of *Mycobacterium tuberculosis* (Mtb) during therapeutic intervention. A deeper understanding of cellular and molecular changes with age could inform new strategies to improve therapeutic outcomes. Here, we monitored the immunopathology, frequency and functionality of immune cells across extreme age groups of C57BL/6 mice following low aerosol dose infection (100-120 cfu) with Mtb H37Rv and treatment with rifampicin and isoniazid (RIF-INH). Up to 6 weeks of infection, tissue (lung, spleen and liver) mycobacterial load in old (17-19 months; M) and aged (31M) C57BL/6 mice was similar compared to young (2-4M) mice. However, at two weeks post-treatment, older mice showed a slower rate of Mtb clearance in the lungs. Old Mtb-infected mice had higher splenic T-follicular cytotoxic (T_{FC})-like cells and proteomic analysis of flow-sorted $CD4^+CD44^+$ T cells revealed deregulated mitochondrial proteins (4-hydroxy-2-oxoglutarate aldolase, aspartate aminotransferase and prostaglandin E synthase), pointing to impaired mitochondrial function. Collectively, these findings suggest that age-associated immune alterations may impair immunometabolic processes contributing to delayed Mtb clearance. These results highlight the potential importance of targeting immunometabolic dysfunction to improve TB treatment outcomes in older populations and reduce morbidity.

Keywords: Aging, Tuberculosis, Immunosenescence, T_{FH} cells, Proteomics

Summary

The elderly population is particularly susceptible to tuberculosis (TB), making it crucial to understand the cellular and molecular mechanisms contributing to the decline in immune responses with age. Evaluating the immunopathology, frequency and functionality of immune cells and Mtb-specific antibody responses across different age groups is essential for developing adjunct therapies for geriatric TB patients.

Introduction

A recent WHO report predicts that, by 2030, one in six people globally will be over 60 years of age[1]. The elderly population is particularly vulnerable to infectious diseases such as tuberculosis (TB) due to age-associated decline in immune function, often referred to as immunosenescence[2]. *Mycobacterium tuberculosis* (Mtb), the causative pathogen of TB, also triggers hypermethylation and transcriptional alterations in host immune cells, affecting the immune system[3]. Older adults not only exhibit reduced vaccine effectiveness but also show higher susceptibility to TB, with epidemiological studies indicating 2–3 times higher TB incidence and up to four times greater mortality than younger TB patients. In several East Asian regions, including Japan, Korea, and Hong Kong, 45–69% of TB cases were reported in 2020-21 from older age groups[4]. Similarly, projections from Taiwan calculated that 78% of new TB cases by 2035 will be from the geriatric population, which currently contributes to 82.1% of TB-associated mortality[5,6]. These findings underscore the growing clinical relevance of TB in an aging world.

Despite this growing burden, current TB research relies largely on young adult mouse models (6–8 weeks old, equivalent to ~18-year-old humans), which do not adequately capture the immune landscape of older hosts[7]. Evidence from studies suggests that old (>18 months; M) mice exhibit higher bacterial burden, delayed CD4$^+$ T cell responses, and altered immune activation compared to younger counterparts[8,9,10]. However, focused studies addressing how immunosenescence shapes bacterial clearance, immune cell dynamics, and host responses during TB treatment remain scarce. To address this gap, we examined the impact of aging on TB infection and treatment outcomes using a murine model. Young (2–4M), mid-aged (9-12M; 38-44 human years), old (17–19M; ~54–60 human years), and aged (31M; >80 human years) C57BL/6 mice were infected with a low aerosol dose of Mtb H37Rv. Following treatment with rifampicin and isoniazid (RIF-INH), we monitored

bacterial clearance kinetics across tissues and profiled systemic and tissue-specific immune

responses, including circulating cytokines, Mtb-specific antibodies, liver micronutrients, and the

proteome of splenic CD4$^+$CD44$^+$ T cells. Through this approach, we aim to identify age-associated

processes that compromise treatment efficacy, which can inform strategies to reduce relapse risk in

geriatric TB patients.

Results

Lung Mtb burden in C57BL/6 mice is unaffected by age or sex till 6 weeks post-infection: The

influence of host age and sex on susceptibility to Mtb infection remains an area of interest,

particularly at early time points. Thus, female and male C57BL/6 mice of wider age groups were

aerosol infected with low-dose of H37Rv (Fig. 1A). Mice were categorized as young (2-4M),

middle-aged (9M females/12M males) and old (17-19M males). Prior to Mtb infection, body

weights were higher in the middle-aged and old mice than in younger controls (Supplementary Fig.

S1A). During the study period, old mice lost ~3% weight, whereas young mice gained ~8% weight

(Fig. 1B). Young female mice gained more body weight (~20%) than males (~10%, Fig. 1B). At 6

weeks post infection (w.p.i.), middle-aged and old male mice showed significant body weight loss

(Fig. 1B). Our findings reveal that, despite age-related variations in body weight, lung Mtb burden

progressively increased up to 4 w.p.i. before plateauing, with no significant differences attributable

to age or sex by 6 w.p.i. (Fig. 1C).

Age-dependent delays in TB treatment responses are associated with altered liver function,

immune dysregulation and inflammaging: To evaluate the impact of age on treatment response,

a subset of mice received RIF-INH following Mtb infection. Remarkably, old mice retained

significantly higher (~3 log$_{10}$ cfu) lung Mtb load at 2 w.p.t. compared to the young group (~1.5 log$_{10}$

cfu). And by 4 weeks, both mice age groups achieved complete bacterial clearance, i.e. resulting

below the limit of detection (LOD; Fig. 2A and 2B). Multiple foci of inflammatory cells and

granulomatous lung lesions were observed with prominent foamy cells later (at 6 w.p.i.) in old mice

but earlier (by 4 w.p.i.) in young controls (Supplementary Fig. S1). Lung inflammation was resolved

in young mice by 4 w.p.t. but persisted in old mice (Supplementary Fig. S1). Also, middle-aged

mice were able to resolve the inflammation by 4 w.p.t. irrespective of sex (Supplementary Fig. S2).

In extrapulmonary sites, like spleen and liver, mycobacterial clearance was observed by 2 w.p.t. in
both the age groups (Fig. 2B).
Liver enzymes play a critical role in drug metabolism, and metal ions play a significant role in their
function. Age-related changes may influence this variability in liver function and micronutrient
status[11]. Thus, we monitored the liver micronutrient levels of these mice groups using inductively
coupled plasma mass spectrometry (ICP-MS, Supplementary Fig. S3). Old mice had higher hepatic
Cu (4.28×), low Zn (0.14×) levels with similar Fe and Se levels compared to controls
(Supplementary Fig. S3). Also, old mice receiving treatment showed higher hepatic Cu levels
(Supplementary Fig. S3). Liver micronutrient status impacts activities of hepatic enzymes like
alanine aminotransferase (ALT) and aspartate aminotransferase (AST), commonly used markers of
liver function and injury. Old mice exhibited higher serum ALT and AST activities than young mice
(Fig. 2C). Mtb infection did not impact serum AST or ALT activity in old mice, but ALT activity
was higher in younger mice post-infection (Fig. 2C). Strikingly, old mice receiving treatment had
higher serum AST activity (Fig. 2C).
To assess age-related differences in humoral immunity against Mtb, serum anti-H37Rv specific IgG
titers were measured. The overall IgG response was significantly higher in old mice compared to
young mice (Fig. 2D). Importantly, old mice showed higher IgG responses at a later time point i.e.
4 w.p.t., while young mice showed it at 2 w.p.t. (Fig. 2D and Supplementary Fig. S4). While robust
humoral responses are beneficial for pathogen control, the delayed yet heightened antibody response
in old mice could indicate persistent antigenic stimulation and likely reflects altered immune
regulation with aging. Circulatory IL-6 levels were higher in old mice, consistent with a state of
chronic low-grade inflammation or inflammaging (Fig. 2E). Higher circulatory IFN-**γ** levels were
observed in Mtb-infected young and old mice, which resolved partially in treated groups (Fig. 2E).
We observed higher lung IFN-γ levels in Mtb-infected young mice compared to old mice (Fig. 2F).
Old mice receiving treatment had declined lung IFN-γ levels (Fig. 2F). Thus, the interplay between
micronutrient imbalance, delayed humoral immunity and inflammaging, likely underpins the
compromised mycobacterial clearance observed in older hosts.
108
Age-related T cell dysregulation impairs immune control of Mtb infection: $CD4^+$ T cells are
critical for Mtb control; however, in older hosts, accumulation of dysfunctional T cells leads to a

compromised immune response. T cells were profiled using activation markers (CD44, PD1),

homing receptor (CXCR5), senescence (KLRG1, CD153) and lineage-defining transcription factors

(Tbet for T_H1 and FoxP3 for T_{reg} cells) and a significant change in their distribution was observed

in old mice (Supplementary Fig. S5). Notably, high expression of CD44, CXCR5, PD1 and CD153

in CD4$^+$ and CD8$^+$ T cells of old mice was observed (Supplementary Figs. S5 and S6). Increased

circulatory CD4$^+$KLRG1$^+$PD1$^+$ cells were observed in old mice, indicative of exhausted T cell

phenotypes (Supplementary Fig. S5).

Next, we monitored the immune cell distribution in lungs and spleen of Mtb-infected and RIF-INH

treated mice groups (Fig. 3A, 3B and Supplementary Fig. S7). Old mice showed a lower proportion

of splenic CD4$^+$, higher CD8$^+$ T cells, along with higher CXCR5 and PD1 expression compared to

young mice (Fig. 3C and Supplementary Fig. S7). Interestingly, lungs of old mice showed

accumulation of CD4$^+$CD44$^+$CXCR5$^+$ (T_{FH})-like cells, similar to young mice post Mtb infection

(Supplementary Fig. S8). Also, CD8$^+$CD44$^+$CXCR5$^+$ (T_{FC})-like cells were observed to be higher in

the lungs and spleen of old Mtb-infected mice at 4 w.p.i. (Fig. 3C and Supplementary Figs. S7 and

S8).

In a separate experimental set, at 4 w.p.i., lung Mtb load was found to be similar between young

(2M) and aged (31M) male C57BL/6 mice groups (Supplementary Fig. S9). Old and aged mice had

overall higher splenic CD4$^+$CD44$^+$CXCR5$^+$PD1$^+$, CD4$^+$CD44$^+$CXCR5$^+$PD1$^+$ICOS$^+$ and

CD8$^+$CD44$^+$CXCR5$^+$PD1$^+$ cells compared to young mice, irrespective of infection and treatment

status (Fig. 3C and Supplementary Fig. S9). The detailed distribution of the immune cells observed

in multiple tissues of 2, 4, 17, 19 and 31M old mice groups is tabulated in Supplementary Table S1.

The increased abundance of T_{FC}-like cells in Mtb-infected old mice suggests dysfunctional T_{FH} cell

activity, which may have contributed to compromised pathogen clearance.

Splenic T cell proteome analysis of old mice upon Mtb infection and anti-TB treatment shows

perturbed mitochondrial pathways: To further elucidate the mechanisms underlying the delayed

Mtb clearance, we conducted global proteomic profiling of flow-sorted CD4$^+$CD44$^+$ T cells (Fig.

3D and Supplementary Table S2). In set-I, 109 proteins were identified, and Mtb-infected young

and old mice showed 23 and 30 deregulated ($p \leq 0.1$, log$_2$fold change (log$_2$fc): I/H > |0.58|) proteins,

respectively (Fig. 3E, 3F and Supplementary Table S3). In set-II, out of 41 identified proteins, ~20

were deregulated in mice groups receiving treatment (Fig. 3G, 3H and Supplementary Table S4).
Proteins related to immune response (e.g., complement, serotransferrin) were suppressed in old
mice, potentially reflecting age-related immune decline. Young mice showed few changes, with
upregulation of stress response proteins like galectin-1 and superoxide dismutase, possibly
reflecting a more robust acute response (Supplementary Table S3). A mitochondrial protein, 4-
hydroxy-2-oxoglutarate aldolase (HOGA), was downregulated in the cells of Mtb-infected mice
groups irrespective of their age (Supplementary Table S3). Upon Mtb infection, old mice showed
an upregulation of Prostaglandin E synthase 3 (log_2fc: I/H = 1.15), the inhibition of which increases
mitochondrial function (Supplementary Table S3). Also, old mice showed elevated levels of
mitochondrial AST after treatment (log_2fc: T/I = 3.15), indicative of greater metabolic adaptation
or mitochondrial stress, consistent with increased serum AST and altered hepatic micronutrient
status observed earlier (Supplementary Table S4). Old mice showed lower HOGA abundance after
treatment, implying that mitochondrial stress persists longer in cells of old mice (Supplementary
Table S4). Together, these proteomic insights provide a molecular basis for the observed immune
dysregulation, inflammaging, and impaired mycobacterial clearance in older hosts, linking cellular
metabolic stress in T cells during Mtb infection and therapy.

Discussion
As the global demography is shifting towards the older ($\geq$60 years) age groups, it is expected that
immunosenescence and lack of an effective TB vaccine for adults will be key drivers for the elevated
numbers of geriatric TB patients. Importantly, in countries where TB was contained successfully,
the number of new TB cases is increasing, for example, in Kansas City and North Carolina in the
United States. The majority of the published reports focused on understanding host-pathogen
interactions in Mtb-infected mice of 6-8 weeks (~2M), representing human age equivalence of 18-
20 years[7]. Thus, understanding how the older host handles Mtb infection is crucial for developing
appropriate interventions.
TB demonstrates a marked gender disparity, with males showing increased susceptibility
(male/female ratio of 1.7 in TB prevalence), which is associated with reduced B cell follicle
formation in the lung[12,13,14]. On the contrary, females exhibit increased T cell activity, production
of T_H1-type cytokines, particularly IFN-γ and robust antibody responses[15]. Above 65 years of age,

genomic disparities between the sexes become more pronounced, with males having higher innate

and pro-inflammatory activity[16]. Thus, we focused on using old male mice to study the outcomes

upon Mtb infection and post-treatment.

A low aerosol dose of Mtb infection to young (2-4M), middle-aged (9/12M), old (17-19M) and aged

(31M) C57BL/6 mice yielded similar lung mycobacterial burden up to 6 w.p.i.. Whereas earlier

reports showed better Mtb control in older mice at earlier time points[8,9,17,18]. For example, Turner et

al reported that 18M C57BL/6 mice aerosol infected with Erdman strain had lower lung load at 14

d.p.i., compared to 3M controls[9]. 24M B6D2F1 hybrid mice infected with H37Rv showed lower

load at 20 d.p.i. than young 2M controls[8]. Variations in reported tissue Mtb loads in earlier reports

and this study may stem from differences in the mouse strain and their sex, Mtb strain and its dose,

infection method, and observation timings. Important to highlight that, to the best of our knowledge,

this report demonstrates for the first time that old mice showed a compromised Mtb clearance after

2 weeks of RIF-INH treatment compared to younger controls. These results corroborate the reported

delayed viral clearance in 17M BALB/c mice infected with PR8 influenza virus[19].

Aging also impacts hepatic function, affecting metabolism of exogenous molecules, thereby leading

to hepatoxicity[20]. Despite their Mtb infection status, old mice showed a higher hepatic Cu/Zn ratio.

Accumulation of Cu and Zn in the phagosomes increases bactericidal activity; however, Cu overload

is detrimental to the host[21]. Low Zn levels in older mice may impair Cu/Zn superoxide-dismutase

function, leading to pro-inflammation and weakened hepatic function[22]. Mtb pathogenesis lowers

the available Fe pool and is a typical host response in counteracting Mtb replication in TB patients[23].

Liver Fe level was reduced in young mice post treatment; however, higher hepatic Fe pool in the

treated old mice may help Mtb for its growth and multiplication. Impaired liver function, indicated

by higher AST levels, can lead to reduced drug metabolism and elimination. Elevated serum AST

and ALT support the concept that aging is linked to increased liver stress, which may contribute to

altered responses to liver insults such as drug treatment post-infection.

Immune cells like $CD4^+$ T cells play a critical role in controlling Mtb, and their function might be

partially impaired in older hosts[24]. Higher lung mycobacterial load was predicted in 18M mice at 6

w.p.i. compared to 3M mice, due to a delayed $CD4^+$ T cell response[24,25]. We observed that Mtb-

infected old C57BL/6 mice had similar T_{FH}-like cells in the lung and a higher frequency in the spleen

at 4 w.p.i. compared to younger controls. However, a higher proportion of T_{FC}-like cells in the lung

and spleen was observed in Mtb-infected old mice. The contribution of T_H1 cells in TB is well
established. However, our understanding of T follicular cells, which link humoral and adaptive
immunity, during Mtb infection is limited[26]. T_{FC} cells suppress T_{FH} cell helper function and antibody
responses through multiple mechanisms, inhibiting antibody production to maintain self-tolerance
and regulate immune responses[27,28]. Age is reported to influence T_{FH} cell function and
differentiation negatively, compromising the host responses[29]. Localization of T_{FH}-like cells has
been demonstrated within protective granuloma-associated lymphoid tissue in lungs of 2M
C57BL/6 mice which mediates Mtb control[30].

Understanding tissue level T cell distribution in Mtb-infected old mice has been attempted earlier;
however, limited reports captured the changes during TB treatment[8,9,17,24,26]. At 2 w.p.t., young mice
showed increased splenic T_{FH}-like cells, which remained similar in old mice, suggesting the inability
of the older host to elicit a proper T_{FH} response. Compromised Mtb clearance in old mice might be
due to the impaired T_{FH} cell functionality, which was reported earlier in >20M old C57BL/6
mice[29,31].

The splenic cell proteome showed significant variation in the abundance of key mitochondrial
enzymes, e.g., HOGA and AAT, involved in the catabolism of hydroxyproline (Hyp), a product of
protein degradation (Fig. 4)[32]. AAT catalyzes 4-hydroxy glutamate (4-OH-Glu) to 4-hydroxy-2-
oxo-glutarate (HOG), which is converted into pyruvate and glyoxylate using HOGA[32]. A higher
abundance of AAT and lower HOGA levels in the splenic cells of Mtb-infected old mice might lead
to HOG accumulation. HOG inhibits glyoxylate reductase, which contributes to excessive oxalate
formation independent of the lactate dehydrogenase pathway and is reported to cause mitochondrial
dysfunction[33,34]. We found higher levels of Prostaglandin E synthase-3 in Mtb-infected old mice,
which is reported to negatively affect the proliferation of alveolar macrophages (AMs)[35]. Also,
inhibition of prostaglandin E2 receptor enhanced mitochondrial function in AMs of 18-22M mice[35].
Reports demonstrated the potential of mitochondrial-based interventions to enhance aged CD4+ T
cell functionality with implications for improving immune responses in geriatric TB patients[36,37].

This study has a few limitations. Due to a limited number of lung T_{FH} cells, we could not profile
their proteome level changes and focused on the splenic CD4+CD44+ T cells. Mouse models in TB
research have limitations for extrapolating findings to humans due to differences in disease
pathology, lacking key features like granuloma liquefaction, cavitation, and fibrosis. Also, mice

exhibit different metabolic and aging patterns compared to humans, with distinct changes in organ

metabolism, immune responses, and biomarkers of aging.

In summary, this study revealed an impaired lung Mtb clearance in old C57BL/6 mice receiving

RIF-INH, mainly due to inflammaging, delayed T_H1 responses, higher T_{FC} cell frequency and

mitochondrial dysfunction in T cells. The findings from this study help us to better understand the

effect of immunosenescence on poorer treatment outcomes in geriatric TB patients.

Materials and Methods

Experimental groups, Mtb infection and treatment: Procedures adopted in this study were

performed following the recommended Guidelines for Care and Use of Laboratory Animals and

approved by the Animal Ethics Committee of ICGEB- New Delhi (Ref No.

ICGEB/IAEC/07032020/TH-10). C57BL/6 mice of different age groups (2,4,9,12,17,19 and 31

months) were procured from the ICGEB bio-experimentation facility. These mice were transferred

to the BSL-III Tuberculosis Aerosol Challenge Facility (TACF) for Mtb infection experiments.

Mice were fed a chow diet, water ad libitum and maintained at 20-22ºC; 45-60% humidity with a

12h light/dark cycle. Mice were weight-adjusted, randomly grouped and after 1 week of

acclimatization in TACF, infected with a low dose (100-120 cfu) of Mtb H37Rv strain in the

Madison aerosol exposure chamber. After 2 w.p.i., a subset of mice received RIF and INH (100

mg/L each) in drinking water and fresh drug solutions were replaced every 48 hours.

Cfu assay: Tissues (lung, spleen and liver) were homogenized in PBS (1 mL) and plated on 7H11

agar supplemented with Middlebrook 10% OADC. These plates were incubated at 37ºC and

bacterial colonies were enumerated 3 weeks after.

Histopathology: Formalin-fixed lungs were paraffin-embedded, sectioned and stained with

hematoxylin and eosin, followed by capturing images using Nikon Eclipse TS2. The slides were

scored based on the severity of inflammation, granuloma formation and necrosis.

Liver micronutrient profiling: Liver (~50 mg) was subjected to microwave digestion post addition

of H_2O_2 and HNO_3 (1:4). These were diluted and introduced to iCAP™-TQ ICP-MS using argon as

carrier. Data acquisition was done using Qtegra (Thermo Scientific, version: 2.10.4345.39) along

with running different concentrations of multielement standard mix (Supelco, #92091).

ALT and AST assay: Serum ALT and AST activities were measured using kits (Cayman Chemical,

#700260 and #701640) and steps provided by the manufacturer.

Immune cell isolation and flow cytometry: Harvested tissues were incubated with collagenase D

(1 µg/mL) and DNase I (0.5 mg/mL) and stained with: CD3-BV510, CD4-BV785, CD8-AF700,

CD44-BV605, CXCR5-PE/Dazzle, PD1-PE-Cy7, KLRG1-BV421, CD153-PE, CD25-AF488 and

CD127-BV650 and live dead-APC-Cy7 on ice for 45 min. After washing with buffer (0.2% FBS in

$1\times$ PBS), cells were treated with fixation and permeabilization buffer and kept on ice for 30 min.

Intracellular staining was done with FoxP3-AF647 and Tbet-BV711 and incubated at RT for 45

min, followed by washing. Data (~1 million events/sample) was acquired using BD LSRFortessa

X-20 and analyzed using FCS Express (DeNovo software, version 6).

Another set of immune cells were stained with: CD3-FITC, CD4-SB600, CD8a-BV421, CD44-

BV510, CXCR5-PE, ICOS-PerCP/Cy5.5, PD1-PE-Cy7 and live dead-APC-Cy7 on ice for 45 min.

After washing with buffer (3% FBS in $1\times$ PBS), splenic $CD4^+CD44^+$ T cells were sorted using

BD FACS Aria III Fusion. Data was analyzed using FlowJo (BD Biosciences, version 10.8.1).

Serum cytokine estimation: Two times diluted serum samples were used for estimating cytokines

using capture bead-based assay LEGENDplex (BioLegend) along with the standard (BioLegend,

#740371, lot no. B330381). The samples were acquired on BD LSRFortressa X-20 using PE and

APC as the reporter and bead classification channel, respectively. Data was analyzed using Qognit

software.

Serum IgG antibody quantification: ELISA plate was coated with Mtb H37Rv whole cell lysate

(10 µg/mL) and incubated overnight at 4°C. Then, wash buffer (0.05% Tween-20 in $1\times$ PBS) was

added, followed by blocking in 3% BSA in $1\times$ PBS containing 0.05% Tween-20 and incubated at

37°C for 2h at 180 rpm. After blotting the plate dry, sera were serially diluted four-fold in 0.1%

BSA in $1\times$ PBS containing 0.05% Tween-20, and 100 µL was added per well and incubated

overnight at 4°C. The plate was washed thrice followed by incubation with anti-mouse IgG

(Southern Biotech, #1031-05) at 37°C for 2h at 180 rpm. The plate was washed six times and

developed by adding o-phenylenediamine dihydrochloride, incubated at RT for 5 min. The reaction

was terminated by adding 1N HCl and the absorbance was measured at 490 nm.

Splenic $CD4^+CD44^+$ cell proteome analysis: Proteins were isolated using TRIzol reagent from

sorted splenic $CD4^+CD44^+$ cells and quantified using BCA Protein Assay. An equal amount (~10

μg) of proteins from a set of samples (n=10) was taken for TMT10plex (Thermo Scientific, #90113).

TMT-labelled peptides from all samples were pooled, dried and subjected to strong cation exchange

chromatography and eluted using ammonium formate. Later, these fractions (n = 5) were cleaned

up using C18 spin columns and dried, resuspended in formic acid (0.1%, 20 μL) and injected into

an Orbitrap Fusion Lumos Tribrid Mass Spectrometer connected to a nano-LC 1200. Data was

analysed with Proteome Discoverer version 2.3 using *Mus musculus* (ID: UP000000589; 55,087

protein count). Proteins qualifying the parameters [$\log_2$ fold change $\geq \pm 0.58$; $\log_{10}$p-value ≥ 1; false

discovery rate<0.05] were selected as important deregulated molecules.

Statistical analysis: Data are presented as mean ± standard deviation. Details regarding the sample

size and statistical analysis have been provided in the figure legends.

Acknowledgements: FP was supported with a Junior and Senior Research fellowship from the

Department of Biotechnology (DBT), Government of India. SC was a Shyama Prasad Mukherjee

Fellow supported by the Council of Scientific and Industrial Research, Government of India. A

Junior Research Fellowship from the DBT supports AG. The core support from the International

Centre for Genetic Engineering and Biotechnology (ICGEB), New Delhi, to RKN is highly

acknowledged. Haripriya Priyadarsini, Adyasha Sarangi, Nidhi Yadav and Mothe Sravya are

acknowledged for their help and support during the mouse experiments. Help from the staff of the

bio-experimentation facility, BSL-III Tuberculosis Aerosol Challenge Facility (TACF) at ICGEB-

New Delhi, Shilpa Sharma and Jaya Baranwal, is acknowledged. DBT supports the TACF. We

thank Dr. Anmol Chandele for her insightful comments and suggestions for revising the manuscript.

Part of these research findings were presented in Keystone Symposia on Molecular Basis of Healthy

Aging in 2023; World Conference on Lung Health 2022, organized by The Union, and the 2nd

biennial meeting on Sex Differences in the Immune Systems Conference in 2022, organized by

Trinity College Dublin & Whitehead Institute. Some figures have been created with BioRender. The

authors declare no competing financial interests.

Author Contributions: FP and RKN conceptualized the project. FP and SC carried out the mouse

experiments and cytokine profiling. FP performed the flow cytometry data acquisition and liver

micronutrient and proteomics sample preparation. AG carried out antibody titer, ELISA

experiments and helped in proteomic data analyses. ShiC calculated histopathological scores. FP and RKN analyzed the data, wrote the original draft and revised it with the suggestions received from all the co-authors. RKN acquired funding, shared resources and administered the project. All co-authors reviewed and approved the manuscript.

Declaration of interests: The authors declare no competing interests.

Supporting Information: The article contains supplemental information.

References

1. Website: https://www.who.int/news-room/fact-sheets/detail/ageing-and-health (Accessed 10/02/2024)

2. Yew, W.W., Yoshiyama, T., Leung, C.C., & Chan, D.P. (2018). Epidemiological, clinical and mechanistic perspectives of tuberculosis in older people. *Respirology*, **23**, 567–575.

3. Bobak, C.A., Abhimanyu, Natarajan, H., Gandhi, T., Grimm, S.L., Nishiguchi, T., ... & DiNardo, A.R. (2022). Increased DNA methylation, cellular senescence and premature epigenetic aging in guinea pigs and humans with tuberculosis. *Aging (Albany NY)*, **14**, 2174–2193.

4. Teo, A.K.J., Morishita, F., Islam, T., Viney, K., Ong, C.W., Kato, S., ... & Marais, B.J. (2023). Tuberculosis in older adults: Challenges and best practices in the Western Pacific Region. *The Lancet Regional Health–Western Pacific*, **36**, 100770.

5. Ku, C.C., & Dodd, P.J. (2019). Forecasting the impact of population ageing on tuberculosis incidence. *PLoS One*, **14**, e0222937.

6. Yen, Y.F., Feng, J.Y., Pan, S.W., Chuang, P.H., Su, V.F., & Su, W.J. (2017). Determinants of mortality in elderly patients with tuberculosis: A population-based follow-up study. *Epidemiology & Infection*, **145**, 1374–1381.

7. Flurkey, K., Currer, J.M., Harrison, D.E., & Fox, J.G. (2007). The mouse in biomedical research. *American College of Laboratory Animal Medicine series. Elsevier, AP: Amsterdam*, 637–672.

8. Cooper, A.M., Callahan, J.E., Griffin, J.P., Roberts, A.D., & Orme, I.M. (1995). Old mice are able to control low-dose aerogenic infections with *Mycobacterium tuberculosis*. *Infection and Immunity*, **63**, 3259–3265.

9. Turner, J., Frank, A.A., & Orme, I.M. (2002). Old mice express a transient early resistance to pulmonary tuberculosis that is mediated by CD8 T cells. *Infection and Immunity*, **70**, 4628–4637.

10. Vesosky, B., Rottinghaus, E.K., Davis, C., & Turner, J. (2009). CD8 T Cells in old mice contribute to the innate immune response to *Mycobacterium tuberculosis* via interleukin-12p70-dependent and antigen-independent production of gamma interferon. *Infection and Immunity*, **77**, 3355–3363.

11. Murugaiha, J.S. (2021). Micronutrient deficiency in pulmonary tuberculosis-perspective on
hepatic drug metabolism and pharmacokinetic variability of first-line anti-tuberculosis drugs:
special reference to fat-soluble vitamins A, D, & E and nutri-epigenetics. *Drug Metabolism*
Letters, **14**, 166–176.

12. Dibbern, J., Eggers, L., & Schneider, B.E. (2017). Sex differences in the C57BL/6 model of
Mycobacterium tuberculosis infection. *Scientific Reports*, **7**, 10957.

13. Gupta, M., Srikrishna, G., Klein, S. L., & Bishai, W.R. (2022). Genetic and hormonal
mechanisms underlying sex-specific immune responses in tuberculosis. *Trends in*
Immunology, **43**, 640–656.

14. Hertz, D., Dibbern, J., Eggers, L., von Borstel, L., & Schneider, B.E. (2020). Increased male
susceptibility to *Mycobacterium tuberculosis* infection is associated with smaller B cell
follicles in the lungs. *Scientific Reports*, **10**, 5142.

15. Klein, S.L., & Flanagan, K.L. (2016). Sex differences in immune responses. *Nature Reviews*
Immunology, **16**, 626–638.

16. Márquez, E.J., Chung, C.H., Marches, R., Rossi, R.J., Nehar-Belaid, D., Eroglu, A., ... &
Ucar,D. (2020). Sexual-dimorphism in human immune system aging. *Nature*
Communications, **11**, 751.

17. Orme, I.M. (1987). Aging and immunity to tuberculosis: Increased susceptibility of old mice
reflects a decreased capacity to generate mediator T lymphocytes. *The Journal of Immunology*,
138, 4414–4418.

18. Lafuse, W.P., Wu, Q., Kumar, N., Saljoughian, N., Sunkum, S., Ahumada, O.S., ... & Rajaram,
M.V. (2022). Psychological stress creates an immune suppressive environment in the lung that
increases susceptibility of aged mice to *Mycobacterium tuberculosis* infection. *Frontiers in*
Cellular and Infection Microbiology, **12**, 990402.

19. Toapanta, F.R., & Ross, T.M. (2009). Impaired immune responses in the lungs of aged mice
following influenza infection. *Respiratory Research*, **10**, 112.

20. Lossow, K., Kopp, J.F., Schwarz, M., Finke, H., Winkelbeiner, N., Renko, K., ... & Kipp, A.P.
(2020). Aging affects sex-and organ-specific trace element profiles in mice. *Aging (Albany*
NY), **12**, 13762–13790.

21. Rowland, J.L., & Niederweis, M. (2012). Resistance mechanisms of *Mycobacterium*
tuberculosis against phagosomal copper overload. *Tuberculosis*, **92**, 202–210.

22. Sakiyama, H., Fujiwara, N., Yoneoka, Y., Yoshihara, D., Eguchi, H., & Suzuki, K. (2016). Cu,
Zn-SOD deficiency induces the accumulation of hepatic collagen. *Free Radical Research*, **50**,
666–677.

23. Cercamondi, C.I., Stoffel, N.U., Moretti, D., Zoller, T., Swinkels, D.W., Zeder, C., ... &
Zimmermann, M.B. (2021). Iron homeostasis during anemia of inflammation: A prospective
study of patients with tuberculosis. *Blood,* **138**, 1293–1303.

24. Orme, I.M., Griffin, J.P., Roberts, A.D., & Ernst, D.N. (1993). Evidence for a defective
accumulation of protective T cells in old mice infected with *Mycobacterium*
tuberculosis. *Cellular Immunology*, **147**, 222–229.

25. Friedman, A., Turner, J., & Szomolay, B. (2008). A model on the influence of age on immunity
to infection with *Mycobacterium tuberculosis*. *Experimental Gerontology*, **43**, 275–285.

26. Crotty, S. (2019). T follicular helper cell biology: A decade of discovery and
diseases. *Immunity*, **50**, 1132–1148.

27. Kim, H.J., Verbinnen, B., Tang, X., Lu, L., & Cantor, H. (2010). Inhibition of follicular T-
helper cells by CD8+ regulatory T cells is essential for self tolerance. *Nature*, **467**, 328–332.

28. Yu, D., & Ye, L. (2018). A portrait of CXCR5+ follicular cytotoxic CD8+ T cells. *Trends in*
Immunology, **39**, 965–979.

29. Sage, P.T., Tan, C.L., Freeman, G.J., Haigis, M., & Sharpe, A.H. (2015). Defective TFH cell
function and increased TFR cells contribute to defective antibody production in aging. *Cell*
Reports, **12**, 163–171.

30. Swanson, R.V., Gupta, A., Foreman, T.W., Lu, L., Choreno-Parra, J.A., Mbandi, S.K., ... &
Khader, S.A. (2023). Antigen-specific B cells direct T follicular-like helper cells into lymphoid
follicles to mediate *Mycobacterium tuberculosis* control. *Nature Immunology*, **24**, 855–868.

31. Silva-Cayetano, A., Fra-Bido, S., Robert, P.A., Innocentin, S., Burton, A.R., Watson, E.M., ...
& Linterman, M.A. (2023). Spatial dysregulation of T follicular helper cells impairs vaccine
responses in aging. *Nature Immunology*, **24**, 1124–1137.

32. Buchalski, B., Wood, K.D., Challa, A., Fargue, S., Holmes, R.P., Lowther, W.T., & Knight, J.
(2020). The effects of the inactivation of Hydroxyproline dehydrogenase on urinary oxalate
and glycolate excretion in mouse models of primary hyperoxaluria. *Biochimica et Biophysica*
Acta (BBA)-Molecular Basis of Disease, **1866**, 165633.

33. Riedel, T.J., Knight, J., Murray, M.S., Milliner, D.S., Holmes, R.P., & Lowther, W.T. (2012).
4-Hydroxy-2-oxoglutarate aldolase inactivity in primary hyperoxaluria type 3 and glyoxylate
reductase inhibition. *Biochimica et Biophysica Acta*, **1822**, 1544–1552.

34. Patel, M., Yarlagadda, V., Adedoyin, O., Saini, V., Assimos, D.G., Holmes, R.P., & Mitchell,
T. (2018). Oxalate induces mitochondrial dysfunction and disrupts redox homeostasis in a
human monocyte derived cell line. *Redox Biology*, **15**, 207–215.

35. Chen, J., Deng, J.C., Zemans, R.L., Bahmed, K., Kosmider, B., Zhang, M., ... & Goldstein, D.
R. (2022). Age-induced prostaglandin E2 impairs mitochondrial fitness and increases mortality
to influenza infection. *Nature Communications*, **13**, 6759.

36. Headley, C.A., Gautam, S., Olmo-Fontanez, A., Garcia-Vilanova, A., Dwivedi, V., Akhter, A.,
... & Turner, J. (2024). Extracellular delivery of functional mitochondria rescues the
dysfunction of CD4+ T cells in aging. *Advanced Science*, **11**, 2303664.

37. Headley, C.A., Gautam, S., Olmo-Fontanez, A., Garcia-Vilanova, A., Dwivedi, V., Schami, A.,
... & Turner, J. (2024). Mitochondrial Transplantation promotes protective effector and

memory CD4+ T cell response during *Mycobacterium tuberculosis* infection and diminishes

exhaustion and senescence in elderly CD4+ T cells. *Advanced Science*, **11**, 2401077.

Figure 1. Lung Mtb burden remained similar till 6 weeks post-infection across age groups and sexes in C57BL/6 mice. **A**. Schematic of the experimental design used for Mtb H37Rv infection of C57BL/6 mice [female (♀) and male (♂)] of different age groups. **B**. Changes in body weight of mice groups (healthy: H, ···· ; infected: I, —) are presented, n=5-10/age group/condition; dashed vertical line represents day of infection. **C**. Lung bacterial burden (in $\log_{10}$cfu/mL) in mice post Mtb infection; n=5-10/timepoint/age group/condition; dashed horizontal line represents limit of detection (LOD). Young (2-4 months) in blue, middle-aged (9-12 months) in green and old (17-19 months) mice in brown; cfu= colony forming unit; w.p.i.= weeks post infection; p-value: ** <0.005 at 95% confidence interval by Mann-Whitney test. Data is represented as mean ± SD. *Additional details available in Supplementary figures S1 and S2* .

Figure 2. Old C57BL/6 mice receiving two weeks of RIF-INH treatment showed delayed lung Mtb clearance. **A**. Lung bacterial burden (in $\log_{10}$cfu/mL) in male C57BL/6 mice belonging to infected (I, at 2, 4 and 6 w.p.i.) and treated (T, at 2 and 4 w.p.t.; treatment started at 2 w.p.i.) groups; n=10/timepoint/age group/condition; dashed horizontal line represents limit of detection (LOD). A set of two independent experiments is depicted. **B**. Bacterial load (in $\log_{10}$cfu/mL) in the lung, spleen and liver; individual data points at 4 w.p.i./2 w.p.t. presented in histograms (n=10/age group/condition). **C**. Serum Alanine transaminase (ALT) and Aspartate aminotransferase (AST) activity in mice of different groups (n=5/age group/condition)) at 4 w.p.i./2 w.p.t.. **D**. Area under the curve (AUC) showing Mtb-specific (H37Rv lysate) IgG quantification in serum of healthy: H (n=15/age group), I (n=27-28/age group) and T (n= 16-17/age group) at 4 w.p.i./2 w.p.t.. **E**. Serum cytokines (in picogram per milliliter): IL-6, IFN-γ in the different mice groups (n=6-7/age group/condition) at 4 w.p.i./2 w.p.t.. **F**. Lung IFN-γ in the different mice groups (n=4-5/age group/condition) at 4 w.p.i./2 w.p.t.. Young (2-4 months) in blue and old (17-19 months) mice in brown; cfu= colony forming unit; w.p.i.= weeks post infection; w.p.t.= weeks post treatment; p-values: * ≤0.05, ** <0.005, *** <0.0005 and **** <0.0001 at 95% confidence interval by Mann-Whitney test. Data is represented as mean ± SD. *Additional details available in Supplementary figures S1 and S4.*

Figure 3. Deregulated splenic CD4$^+$CD44$^+$ T cell proteome profile observed in old C57BL/6 mice upon Mtb infection and RIF-INH treatment. **A**. Schematic showing Mtb H37Rv infection of male C57BL/6 mice of two age groups (young: 2-4 and old: 17-19 in months) followed by treatment with RIF-INH starting at 2 w.p.i.. **B**. Method adopted for immune cell isolation from the spleen, flow cytometry data acquisition and data analysis. **C**. Frequency of CD4$^+$CD44$^+$CXCR5$^+$, CD4$^+$CD44$^+$CXCR5$^+$PD1$^+$ and CD4$^+$CD44$^+$CXCR5$^+$PD1$^+$ICOS$^+$ and CD8$^+$CD44$^+$CXCR5$^+$ cells in the spleen of C57BL/6 mice at 4 w.p.i./2 w.p.t. are presented. Healthy (H, n=3/age group), Mtb H37Rv infected (I, n= 6-10/age group) and RIF-INH treated (T, n=3/age group). **D**. Schematic representation of method adopted for TMT10plex-T cell proteome analysis (for TMT set-1) by liquid chromatography- mass spectrometry (LC-MS/MS). Volcano plot of splenic CD4$^+$CD44$^+$ T cell proteome showing significantly deregulated proteins (in -log$_{10}$p-value ≥ 1; log$_2$fold change > |0.58|) in (**E**) Mtb infected versus healthy (young), (**F**) Mtb infected versus

healthy (old), (**G**) RIF-INH treated versus Mtb infected (young) and (**H**) RIF-INH treated versus Mtb infected (old); ▲/▲: up-/down-regulation; HOGA: 4-hydroxy-2-oxoglutarate aldolase; AST: aspartate aminotransferase. Young mice in blue and old mice in brown; cfu= colony forming unit; w.p.i.= weeks post infection; p-values: * ≤0.05, ** <0.005, *** <0.0005 and **** <0.0001 at 95% confidence interval by Mann-Whitney test. Data is represented as mean ± SD. *Additional details available in Supplementary figures S7, S8, S10 and Supplementary tables S1, S2, S3 and S4.*

Figure 4. Old C57BL/6 mice showed a compromised lung *Mycobacterium tuberculosis* (Mtb) H37Rv clearance in the early time points of rifampicin and isoniazid treatment. Illustrative summary of the results observed in the study. (**1**) Following Mtb infection via aerosol challenge, (**2**) antigen presentation takes place in the spleen. (**3a**) T cells take longer to reach the inflamed lung of old mice. (**3b**) Increased levels of proinflammatory cytokines: interferon-gamma (IFN-γ) and decreased Mtb-specific IgG levels observed in Mtb-infected old mice. (**4**) Mtb disseminates to the liver via hepatic artery where iron (Fe) was observed to be accumulated along with higher Aspartate aminotransferase (AST) activity, suggestive of higher reactive oxygen species (ROS) generation (**5**) which also in turn could increase circulatory proinflammatory mediators. Old mice had low splenic T follicular helper (T_{FH}: CD4$^+$CD44$^+$CXCR5$^+$PD1$^+$) cells, and presented a delayed lung Mtb clearance at 2 weeks post treatment. Also, proteome analysis of splenic CD4$^+$CD44$^+$ T cells of old mice showed alterations in mitochondrial proteins (HOGA: 4-hydroxy-2-oxoglutarate aldolase and AST) which are related to hydroxyproline (Hyp) degradation pathway. Increased AST and decreased HOGA might lead to buildup of 4-hydroxy-2-oxo-glutarate (4-OH-2-oxo-Glr) from glutamate (Glu), directing the reaction to oxalate (Oxa) via pathway independent of lactate dehydrogenase (LDH), pyruvate (Pyr) and glyoxylate (Gly-ox). Figure created with *BioRender*.com