A Mechanistic Insight into Sources of Error of Visual Working Memory in Multiple Sclerosis

Ali Motahharynia,[1,2†] Ahmad Pourmohammadi,[1,2,4†] Armin Adibi,[1,2] Vahid Shaygannejad,[1,2,3] Fereshteh Ashtari,[1,2,3] Iman Adibi,[1,2,3*] and Mehdi Sanayei[1,2,4*]

Author affiliations:

1. Center for Translational Neuroscience, Isfahan University of Medical Sciences, Isfahan, Iran

2. Isfahan Neuroscience Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

3. Department of Neurology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

4. School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

* Corresponding authors:

Mehdi Sanayei
School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran, 1956836484, Iran,
Tel: +98 212 611 3274
Email: mehdi.sanayei@gmail.com

Iman Adibi
Department of Neurology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, 81746 75731, Iran,
Tel: +98 313 620 1286
Email: i.adibi@gmail.com
ORCID: 0000-0003-4409-9404

[†]These authors contributed equally to this work.

Abstract

Working memory (WM) is one of the most affected cognitive domains in multiple sclerosis (MS), which is mainly studied by the previously established binary model for information storage (slot model). However, recent observations based on the continuous reproduction paradigms have shown that assuming dynamic allocation of WM resources (resource model) instead of the binary hypothesis will give more accurate predictions in WM assessment. Moreover, continuous reproduction paradigms allow for assessing the distribution of error in recalling information, providing new insights into the organization of the WM system. Hence, by utilizing two continuous reproduction paradigms, memory-guided localization (MGL) and analog recall task with sequential presentation, we investigated WM dysfunction in MS. Our results demonstrated an overall increase in recall error and decreased recall precision in MS. Furthermore, computational modeling of the results from the sequential paradigm determined that imprecision in decoding information and swap error (mistakenly reporting the feature of other presented items) were responsible for WM dysfunction in MS. Overall, this study offered a sensitive measure for assessing WM deficit and provided new insight into the organization of the WM system in MS population.

Keywords: Binding; Multiple sclerosis; Resource model; Swap error; Working memory

Introduction

Multiple sclerosis (MS) is a debilitating inflammatory disorder characterized by demyelinating central nervous system (CNS) plaques creating a progressive neurodegenerative state with heterogeneous clinical characteristics (Benedict et al., 2020; Dobson & Giovannoni, 2019). Impairment in cognitive function is a common clinical manifestation of MS, which detrimentally affects different aspects of patients' daily life, from decreased physical performance and productivity to unemployment (Benedict et al., 2020; Clemens & Langdon, 2018). One of the frequently affected domains of cognition in MS is working memory (WM) (Vacchi et al., 2017), which due to its essential role in several cognitive processes (Miller et al., 2018; Nee & D'Esposito, 2016), is one of the main areas of MS research (Costers et al., 2021; Covey et al., 2011; Hulst et al., 2017; Rocca et al., 2014; Vacchi et al., 2017).

Multiple neuropsychological cognitive paradigms, such as paced auditory serial addition test (PASAT), n-back, and delayed-match to sample, were developed to investigate different aspects of WM deficit in MS (Costers et al., 2021; PARMENTER et al., 2006; Pourmohammadi et al., 2022; Rocca et al., 2014; Stojanovic-Radic et al., 2015; Vacchi et al., 2017). The basis of these change detection paradigms is the slot model of WM (Ma et al., 2014). In this quantized model, WM is considered a short-term storage for a limited number of items (Cowan, 2001; Ma et al., 2014; MILLER, 1956), storing the information in a binary format. This assumption creates an all-or-none condition in which only the stored items in these limited slots will be remembered (Ma et al., 2014). Nonetheless, recent observations from analog recall paradigms assessing the precision of WM determined dynamic allocation of WM resources (Bays et al., 2009; Bays & Husain, 2008; Gorgoraptis et al., 2011; Schneegans & Bays, 2016). Each stored item in this framework possesses a fraction of WM storage in which the allocated space changes dynamically between them (Bays & Husain, 2008; Ma et al., 2014; Schneegans & Bays, 2016). This concept is the foundation for the resource-based model of WM.

The analog nature of inputs in resource-based model paradigms makes it possible to investigate the resolution and variability of stored memory (Peich et al., 2013; Zokaei et al., 2015). Also, assessing the distribution of error in these paradigms further helped in uncovering the underlying structure of the visual WM system (Liang et al., 2016; Lugtmeijer et al., 2021; McMaster et al., 2022; Peich et al., 2013; Zokaei et al., 2020). An analog recall task is a paradigm in which

subjects need to simultaneously recall multiple features of items in a continuous space, hence
requiring encoding the information of connected features in addition to their distinct value (e.g.,
object, location, and object-location binding information) (McMaster et al., 2022; Peich et al.,
2013; Zokaei et al., 2020). According to the study of Bays et al., there are three different sources
of error for recalling information in visual WM tasks with connected features (Bays et al., 2009).
They are identified as i: the Gaussian variability in response around the target value (target
response proportion), ii: Gaussian variability in response around the non-target value, i.e.,
mistakenly reporting feature of other presented items (swap error), and iii: random responses
(uniform response proportion) (Bays et al., 2009; Zokaei et al., 2020).

Studies based on analog recall paradigms unraveled new insights into the sources of recall error
in neurodegenerative disorders. It was determined that random response and swap error
contribute to the impairment of visual WM in Parkinson's and Alzheimer's diseases, respectively
(Liang et al., 2016; Zokaei et al., 2020). However, regardless of the influential impact of these
study designs on the discovery of novel mechanistic insights into the organization of visual WM
system, it has not gotten enough attention in the field of MS research.

The current study followed our previous study in which the quantity of MS-related visual WM
was assessed based on the slot model (Pourmohammadi et al., 2022). Here, we aimed to evaluate
recall precision (quality) based on the resource-based model paradigms. In this regard, we
developed two analog recall paradigms, a memory-guided localization (MGL) and two analog
recall tasks with sequential bar presentation (3-bar and 1-bar). Primarily using the simplistic
design of MGL, recall error (absolute error) and precision of visual WM were assessed.
Similarly, recall error and precision were evaluated using the two designed analog recall
paradigms with sequential presentation, i.e., the low memory load, 1-bar, and high memory load,
3-bar conditions, respectively. Furthermore, the classifying performance of these paradigms in
distinguishing different groups was assessed. Finally, the distribution of recall error in analog
paradigms with sequential presentation was further investigated using the probabilistic model of
Bays, Catalao, and Husain (Bays et al., 2009). In both MGL and sequential paradigms (low and
high memory load conditions), recall error and precision were impaired in MS. The dissociable
function of these paradigms in classifying MS subtypes (relapsing-remitting and secondary
progressive) gave some clues about the underlying structure of WM deficit in progressive states
of the disease. Investigation into sources of error in the high memory load condition revealed that

target response proportion and swap error (non-target response proportion) contribute to visual
WM dysfunction in MS.

Results

In the memory-guided localization (MGL) paradigm (Figure 1A), 45 patients (19 relapsing-
remitting MS, RRMS, and 26 secondary progressive MS, SPMS) and 24 healthy controls
participated. As mentioned in the materials and methods section, the mean data of five
participants (one healthy control, three RRMS, and one SPMS) were excluded from further
analysis. In the sequential paradigm with 3 bar (Figure 1B), from a total of 76 patients (42
RRMS and 34 SPMS) and 49 healthy controls who participated, three healthy controls, three
RRMS, and two SPMS were excluded. The demographic and clinical data of participants were
summarized in Table 1 and Table 2.

Recall Error in Multiple Sclerosis

In the MGL paradigm (Figure 1A), recall error was evaluated as a function of distance using a
mixed-model analysis of variance (ANOVA). Recall error was significantly different between
groups [$F(2,61) = 14.57$, $P < 10^{-5}$] and distances ([$F(2,61) = 85.03$, $P < 10^{-23}$], Figure 2A). A
significant interaction was also observed between group and distance [$F(4,61) = 7.24$, $P < 10^{-4}$].
Tukey post hoc test determined that recall error was significantly higher in SPMS [1.86º ± 0.92º
visual degree] compared to healthy control [0.97 ± 0.26, $P < 10^{-4}$] and RRMS [1.09 ± 0.27, $P <$
10^{-3}]. No significant difference was detected between RRMS and healthy control [$P = 0.83$].
Similarly, recall error as a function of delay interval was also evaluated. Recall error was
significantly different between delay intervals ([$F(4,61) = 18.89$, $P < 10^{-12}$], Figure 2D). No
significant interaction was observed between group and delay interval [$F(8,61) = 0.69$, $P = 0.70$].
After adjusting for cognitive ability, assessed by the Montreal cognitive assessment (MoCA)
screening tool (cognitive performance was significantly different between groups), the effect of
group on recall error remained significant (Appendix 1—table 1).
While reaction time (RT) was significantly different between groups [$F(2,61) = 26.44$, $P < 10^{-8}$]
and distances ([$F(2,61) = 25.94$, $P < 10^{-9}$], Figure 2C), it was not significantly different between
delay intervals ([$F(4,61) = 0.97$, $P = 0.43$], Figure 2F). No significant interaction was observed

between group and distance [$F(4,61) = 2.06$, $P = 0.09$] or group and delay interval [$F(8,61) =$

0.86, $P = 0.55$]. The statistical results of RT were summarized in Appendix 2—table 1.

Recall error was evaluated for sequential tasks using the same method. In the 'high memory load'

condition (i.e., sequential paradigm with 3 bar, Figure 1B), recall error was significantly different

between groups [$F(2,114) = 28.18$, $P < 10^{-9}$] and bar orders ([$F(2,114) = 48.74$, $P < 10^{-17}$],

Figure 2G). No significant interaction was observed between them [$F(4,114) = 1.21$, $P = 0.31$].

Tukey post hoc test showed that recall error was significantly higher in RRMS [0.68 ± 0.20

radian] and SPMS [0.76 ± 0.17] compared to healthy control [0.48 ± 0.16, $P < 10^{-5}$, $P < 10^{-8}$,

respectively]. However, no significant difference was detected between RRMS and SPMS

groups [$P = 0.14$]. After adjusting for gender, age, education, and cognitive ability (they were

significantly different between groups), the group's effect on recall error remained significant

(Appendix 1—table 2). Moreover, while RT was also significantly different between groups

[$F(2,114) = 12.95$, $P < 10^{-5}$] and bar orders ([$F(2,114) = 5.92$, $P < 0.004$], Figure 2I), no

significant interaction was observed between group and bar order [$F(4,114) = 0.16$, $P = 0.96$].

The statistical results of RT were summarized in Appendix 2—table 2.

In the 'low memory load' condition (i.e., 1-bar paradigm, Figure 1C), recall error was

significantly different between groups ([$F(2,114) = 36.85$, $P < 10^{-12}$], Figure 2G). Tukey post hoc

test showed that recall error was significantly higher in RRMS [0.33 ± 0.12 radian] and SPMS

[0.43 ± 0.12] compared to healthy control [0.22 ± 0.08, $P < 10^{-4}$, $P < 10^{-8}$, respectively]. We also

observed a significant difference between RRMS and SPMS [$P < 10^{-3}$]. After adjusting for

gender, age, education, and cognitive ability, the group's effect on recall error remained

significant (Appendix 1—table 3). Correspondingly, RT differed significantly between groups

([$F(2,114) = 12.59$, $P < 10^{-4}$], Figure 2I). The statistical results of RT were summarized in

Appendix 2—table 2.

Recall Precision in Multiple Sclerosis

In the MGL paradigm, recall precision was also significantly different between groups [$F(2,61)$

$= 13.74$, $P < 10^{-4}$] and distances ([$F(2,61) = 23.39$, $P < 10^{-8}$], Figure 2B). No significant

interaction was observed between group and distance [$F(4,61) = 0.91$, $P = 0.46$]. Post hoc

analysis determined recall precision was significantly lower in SPMS [0.87 ± 0.52 /º] than in

both RRMS [1.32 ± 0.51, $P < 0.039$] and healthy control [1.70 ± 0.60, $P < 10^{-5}$]. Recall precision

was not significantly different between RRMS and healthy control [$P = 0.08$]. Moreover, this

effect remained significant after adjusting for cognitive ability (Appendix 1—table 1). We also

determined the effects of delay interval on recall precision. In this analysis, groups [$F(2,61) =$

$14.57, P < 10^{-5}$], delay intervals [$F(4,61) = 7.30, P < 10^{-4}$], and their interaction [$F(8,61) = 2.44$.

$P < 0.02$] had significant effects on recall precision (Figure 2E). Post hoc analysis showed the

same pattern of result [SPMS = 0.84 ± 0.47; RRMS = 1.39 ± 0.46; healthy = 1.63 ± 0.61; SPMS

vs. RRMS: $P < 0.005$; SPMS vs. healthy: $P < 10^{-5}$, RRMS vs. healthy: $P = 0.34$]. This effect

remained significant after adjusting for cognitive ability (Appendix 1—table 1).

Similarly, in the high memory load condition, recall precision was significantly different

between groups ([$F(2,114) = 25.23, P < 10^{-9}$], Figure 2H) and bar orders [$F(2,114) = 20.70, P <$

10^{-8}]. However, no significant interaction was observed between them [$F(4,114) = 1.84, P =$

0.12]. Besides, while in post hoc analysis, recall precision was significantly higher in healthy

control [2.39 ± 0.78 /radian] than in both RRMS [$1.68 \pm 0.50, P < 10^{-6}$] and SPMS patients [1.52

$\pm 0.31, P < 10^{-7}$], no difference was observed between RRMS and SPMS participants [$P = 0.49$].

After adjusting for gender, age, education, and cognitive ability, the effect of group on recall

precision remained significant (Appendix 1—table 2).

Accordingly, the low memory load condition showed the same pattern. Recall precision

significantly differed between groups ([$F(2,114) = 25.48, P < 10^{-9}$], Figure 2H). Post hoc

analysis determined that recall precision was significantly higher in healthy control [6.10 ± 2.41

/radian] than in RRMS [$4.16 \pm 1.98, P < 10^{-4}$] and SPMS [$2.95 \pm 1.05, P < 10^{-8}$]. Moreover, there

was a significant difference between RRMS and SPMS [$P < 0.031$], which after adjusting for

gender, age, education, and cognitive ability, remained significant (Appendix 1—table 3).

Distribution of Error in Recalling Information in Multiple Sclerosis

The distribution of error in recalling information was assessed further to investigate the

underlying mechanisms of WM impairment in MS. In this regard, the recorded data from the

sequential paradigms was fitted to a probabilistic model developed by Bays et al. (Bays et al.,

2009). In line with the results of recall error and precision, for the high memory load condition,

circular standard deviation (SD) of von Mises distribution of recall error was significantly

different between groups [$F(2,114) = 26.79, P < 10^{-9}$] and bar orders ([$F(2,114) = 14.95, P < 10^{-9}$

^{6}], Figure 3A). At the same time, they had no significant interaction [$F(4,114) = 1.19, P = 0.31$].

von Mises SD of recall error was lower in healthy control [0.51 ± 0.12] than both RRMS [0.69 ±

0.20, $P < 10^{-5}$] and SPMS [0.76 ± 0.15, $P < 10^{-8}$]. There was no difference between RRMS and

SPMS in the von Mises SD parameter [$P = 0.16$]. In the low memory load condition, von Mises

SD was affected by groups ([$F(2,114) = 33.07$, $P < 10^{-11}$], Figure 3A). Again, von Mises SD was

lower in healthy control [0.25 ± 0.10] than both RRMS [0.38 ± 0.14, $P < 10^{-5}$] and SPMS [0.47 ±

0.12, $P < 10^{-8}$]. There was a significant difference between RRMS and SPMS in von Mises SD

[$P < 0.02$]. After adjusting for gender, age, education, and cognitive ability, the group's effect on

von Mises SD remained significant in both high and low memory load conditions (Appendix 1—

table 2 and Appendix 1—table 3).

According to the study by Bays et al. (Bays et al., 2009), there are three sources of error for

recalling information. The sources of these errors were defined as Gaussian variability in

response around i: target (target response proportion, Figure 3B) and ii: non-target values (swap

error, Figure 3C) and iii: random responses (uniform response proportion, Figure 3D). In the

high memory load condition, target response proportion was significantly different between

groups [$F(2,114) = 11.04$, $P < 10^{-4}$] and bar orders ([$F(2,114) = 10$, $P < 10^{-4}$], Figure 3B). No

significant interaction was observed between group and bar order [$F(4,114) = 0.43$, $P = 0.78$].

Target proportion was higher in healthy control [0.88 ± 0.11] than both RRMS [0.79 ± 0.12, $P <$

0.003] and SPMS [0.76 ± 0.14, $P < 10^{-4}$]. There was no significant difference in target proportion

between RRMS and SPMS groups [$P = 0.54$]. After adjusting for gender, age, years of

education, and cognitive ability, the effect of group on target proportion remained significant

(Appendix 1—table 2). Moreover, the nearest neighbor analysis was performed to further

evaluate the effect of target proportion in the absence of swap error. Removing the effect of swap

error allowed us to assess the isolated effect of orientation recall. The findings from the nearest

neighbor analysis showed a similar pattern of results (Figure 3—figure supplement 1). The

isolated effect of orientation was significantly different between groups [$F(2,114) = 29.26$, $P <$

10^{-10}] among different bar orders [$F(2,114) = 7.07$, $P < 0.002$]. No significant interaction was

observed between group and bar order [$F(4,114) = 0.58$, $P = 0.67$]. After adjusting for gender,

age, education, and cognitive ability, the results of nearest neighbor analysis remained significant

(Appendix 1—table 2).

In the low memory load condition, group significantly affected target response proportion

([$F(2,114) = 3.11$, $P < 0.049$], Figure 3B). While target proportion in the healthy group [0.98 ±

0.03] was significantly higher than SPMS patients [0.94 ± 0.09, $P < 0.04$], after adjusting for

gender, age, years of education, and cognitive ability, this effect became insignificant (Appendix

1—table 3). In addition, the target proportion did not significantly differ between healthy vs.

RRMS [0.96 ± 0.08, $P = 0.48$] and RRMS vs. SPMS population [$P = 0.37$].

In line with the above findings, swap error was higher in the MS population. In the high memory

load condition, swap error was significantly different between groups [$F(2,114) = 7.11$, $P <$

0.002] and bar orders ([$F(2,114) = 31.05$, $P < 10^{-11}$], Figure 3C). No significant interaction was

observed between group and bar order [$F(4,114) = 1.45$, $P = 0.22$]. Swap error was lower in

healthy control [0.07 ± 0.06] than in both RRMS [0.11 ± 0.09, $P < 0.05$] and SPMS patients

[0.14 ± 0.09, $P < 0.002$]. There was no significant difference in swap error between RRMS and

SPMS [$P = 0.41$]. After adjusting for gender, age, years of education, and cognitive ability, the

group's effect on swap error remained significant (Appendix 1—table 2). Moreover, while in the

high memory load condition, the uniform response proportion was different between groups

([$F(2,114) = 5.50$, $P < 0.006$], Figure 3D), no such differences were observed between bar orders

[$F(2,114) = 0.81$, $P = 0.45$) or the interaction between them [$F(4,114) = 0.18$, $P = 0.95$]. Post hoc

analysis revealed that uniform proportion was lower in healthy control [0.05 ± 0.08] than both

RRMS [0.09 ± 0.08, $P < 0.03$] and SPMS [0.10 ± 0.08, $P < 0.02$]. Moreover, there was no

significant difference in uniform response proportion between RRMS and SPMS groups [$P =$

0.95]. Additionally, although after adjusting for gender, the effect of group on uniform

proportion remained significant, adding age, years of education, or cognitive ability made this

effect insignificant (Appendix 1—table 2). The result of uniform response proportion in the low

memory load condition is mathematically same as the target proportion (uniform proportion = 1

– target proportion, since there was no swap error in the 1-bar condition).

Dissociable Function of MGL and Sequential Paradigms

The classifying ability of MGL and sequential paradigms in differentiating healthy control from

MS patients was assessed based on recall error parameters using the receiver operating

characteristic (ROC) analysis (Figure 4A-C). The accuracy of MGL and sequential paradigms

with 3 bar and 1 bar in differentiating MS patients from healthy participants was 80% (Figure

4A), 83.4% (Figure 4B), and 86.2% (Figure 4C), respectively. A closer look at Figures 2A, 2D,

and 2G suggested that these paradigms differentiate MS and healthy control with distinct

patterns. Hence, we separately applied ROC analysis to healthy control vs. RRMS, healthy
control vs. SPMS, and RRMS vs. SPMS for MGL (Figure 4D) and sequential paradigms with 3
bar (Figure 4E) and 1 bar (Figure 4F). While the MGL paradigm had good performance in
differentiating SPMS from healthy control [90.1%] and SPMS from RRMS [84.7%], it had poor
ability in distinguishing healthy control from RRMS ([64.1%], Figure 4D). Accordingly,
although the 3-bar paradigm also had good accuracy in differentiating healthy control from
SPMS [88.4%] and better results (compared to MGL) in discriminating healthy control from
RRMS [79.3%], it did a poor job in discriminating MS subtypes ([62%], Figure 4E).
Complementary to the above findings, the 1-bar paradigm showed an in-between pattern of
results. The 1-bar paradigm accurately discriminates healthy control from SPMS [94.4%], while
it also performed well in differentiating healthy control from RRMS [79.6%]. However,
compared to MGL, it had a weaker ability to discriminate MS subtypes ([72.3%], Figure 4F).

Discussion

In the present study, we investigated the visual working memory (WM) deficits in MS using two
continuous reproduction paradigms: memory-guided localization (MGL) and analog recall
paradigms with sequential presentation. Our results align with the previous reports regarding
WM deficit in MS (Costers et al., 2021; PARMENTER et al., 2006; Pourmohammadi et al.,
2022; Stojanovic-Radic et al., 2015; Vacchi et al., 2017). Complementary to these reports, which
envisaged a binary model for storing information (slot-based model), we assessed recall
precision in WM using analog reproduction paradigms (resource-based model). Further, by
utilizing the unique design of our sequential paradigm, which allows us to assess the distribution
of error in recalling information, we introduced a new mechanistic insight into the visual WM
dysfunction in MS.
Although the results from both MGL and sequential paradigms showed an overall decrease in
recall precision and increased recall error in MS population, the post hoc analysis demonstrated
inconsistent results. In the MGL paradigm, while SPMS patients performed worse than other
groups, no significant difference was observed between healthy control and RRMS. This result
contrasted with the sequential paradigm with 3 bar (high memory load condition), in which MS
subtypes (RRMS vs. SPMS) were not significantly different, and healthy control performed

better than RRMS and SPMS. The situation also varied for the 1-bar paradigm (low memory load condition), in which all three groups performed with different levels of precision. ROC analysis further confirmed these results, which determined a dissociation between the classifying performance of MGL and sequential paradigms. Additionally, although the low memory load condition was better than the high memory load condition in distinguishing MS subtypes, its classifying performance was not as good as the MGL paradigm. It seems that these paradigms evaluated distinct aspects of WM dysfunction in MS.

The dissociable function of our paradigms could arise from using different types of stimuli (location in MGL vs. orientation in sequential paradigms), in which spatial WM was assessed in the MGL paradigm. As the spatial WM process was associated with the function of the hippocampus (Rolls, 2018), the observed difference could indicate more hippocampal disruption in SPMS patients. This finding is in line with previous studies that showed more hippocampal regional loss (Sicotte et al., 2008) and increased hippocampus neuroinflammatory activity in SPMS compared to RRMS (Cree et al., 2021), suggesting that assessment of the spatial WM could be a specific marker for disorganization of the WM system in SPMS.

Another explanation is the long delay interval in MGL, which assessed the maintenance of information. Therefore, the observed difference could be due to additional impairment of SPMS patients in keeping that information. This finding is in line with our previous study, which showed that change detection paradigms with long delay intervals were promising in differentiating MS subtypes (Pourmohammadi et al., 2022). At the same time, one may debate that this difference was related to the longer stimulus presentation time in the MGL paradigm (1000ms vs. 500ms in the sequential paradigm). However, since the stimulus presentation time was adequate in sequential paradigms and the stimuli were presented consecutively, it did not seem that the inadequate time for encoding information was responsible for this difference (Peich et al., 2013; Zokaei et al., 2015).

Additionally, one may argue that the observed dissociation could be due to the extra binding process needed in the sequential paradigm with 3 bar. However, the results from the low memory load condition and our findings from the sequential paradigm nearest-neighbor analysis, which provided a proxy to assess the isolated effect of orientation, demonstrated a similar pattern of dissociation in the absence of a binding effect. Based on these findings, we concluded that the binding process was not responsible for the observed dissociation. Nevertheless, since the

evaluated binding process was an intra-term association (i.e., conjunctive binding), we could not
be assured that the same results would be reached for an inter-term association (i.e., relational
binding) (Parra et al., 2015). This issue becomes more interesting when we realize that the
relational binding function is mainly centered on the hippocampus (Liang et al., 2016; Parra et
al., 2015), the structure we presumed was responsible for the observed dissociation in the MGL
paradigm.

Finally, due to the diffuse pattern of involved brain areas in MS and evidence demonstrating that
brain networks accounted for different WM processes, it is reasonable to assume that distinct
WM-related networks, instead of a single region, were responsible for the observed dissociable
patterns (Bastin et al., 2019; Dobson & Giovannoni, 2019; Figueroa-Vargas et al., 2020;
Lugtmeijer et al., 2021; Vacchi et al., 2017).

We applied a probabilistic model to disentangle the error distribution in recalling information in
the sequential paradigms to further investigate the underlying mechanism behind the visual WM
dysfunction in MS. Our finding revealed that in addition to imprecision in decoding information,
swap error (mistakenly reporting a non-target feature) contributed to WM dysfunction in
individuals with MS. Swap error has been associated with various mechanisms, including the
variability in cue feature dimension, cue-independent sources, and strategic guessing (McMaster
et al., 2022). The recent study by McMaster et al. comprehensively investigated these hypotheses
and determined that the variability in cue feature dimension could solely account for the swap
error mechanism (McMaster et al., 2022). The neural correlates of swap error could be explained
by the imprecision in decoding information in the cue feature dimension (i.e., color in this study)
due to the noise in neural activity. Hence, a non-cued item may mistakenly be considered as the
actual cue, leading to the reporting of the corresponding non-target feature as a response
(Schneegans & Bays, 2017). Therefore, the swap error observed in this study can be attributed to
the variability in the color dimension. The neural underpinning of this observation in MS
population could be related to their impaired neural activity. Recent studies have demonstrated
disturbed neural activity during WM tasks in MS (Costers et al., 2021; Figueroa-Vargas et al.,
2020). For instance, Costers et al. showed a disturbed theta band oscillation, which plays a
crucial role in WM encoding information, in the hippocampal area of MS population (Costers et
al., 2021). While this evidence provides a primary insight into the possible underlying

mechanisms of WM dysfunction in MS, future electrophysiological studies using the cued recall

paradigms could further elucidate this matter.

Swap errors (misbinding errors) have been observed in various neurological disorders, including

different types of Alzheimer's disease (Cecchini et al., 2022; Della Sala et al., 2012; Liang et al.,

2016; Zokaei et al., 2020), epileptic patients with temporal lobe lobectomy (Zokaei et al., 2019),

and voltage-gated potassium channel complex antibody (VGKC-Ab) limbic encephalitis

(Pertzov et al., 2013). From a neuroanatomical perspective, previous studies proposed that

impairment in hippocampus and medial temporal lobe regions contribute to problems in

relational binding (Della Sala et al., 2012; Liang et al., 2016; Pertzov et al., 2013; Zokaei et al.,

2019). Furthermore, whereas regions related to conjunctive binding have yet to be better

understood, occipital and parietal regions seem to be mainly related to conjunctive processing

(Cecchini et al., 2022). Additionally, a recent study by Valdés Hernández and colleagues also

demonstrated the possible role of globus pallidus in conjunctive binding (Valdés Hernández et

al., 2020). On the other hand, a study by Vacchi et al. demonstrated less activation of the

superior and inferior parietal lobule during an N-back fMRI task in individuals with MS (Vacchi

et al., 2017). Moreover, globus pallidus was shown to be involved in MS (Fujiwara et al., 2017).

Hence, considering the convergence of brain regions involved in conjunctive processing

(assessed through the current sequential paradigm) with those shown to be affected in MS, the

parietal regions and globus pallidus could be the candidate regions responsible for the observed

impairment. Additionally, based on the evidence showing the involvement of hippocampal

regions in MS (Rocca et al., 2018; Sicotte et al., 2008), we also expect to see relational

processing impairment; however, the current study design did not allow us to evaluate this

condition. Further neuroimaging studies are needed to validate these findings. Eventually, the

insignificant results from the low memory load condition suggested more impairment in visual

WM under high memory load situations, which was not unexpected.

Despite these findings, our study had some limitations that should be addressed in future

research. The current study only assessed WM dysfunction using behavioral paradigms. Further

structural and functional evaluations should be performed to confirm our suggested brain areas

associated with conjunctive binding and spatial WM deficit in MS. Concurrent assessment of

brain networks using fMRI and EEG or volumetric studies alongside behavioral paradigms could

address this issue. Furthermore, although we hypothesized that relational binding could be a

specific marker for the progressive state of the disease based on the more disruption of
hippocampal-related areas in SPMS, the current study design did not allow us to evaluate this
assumption. Future studies assessing the source of WM deficit regarding relational binding could
elucidate this statement. Additionally, although clinical and cognitive assessments were
performed to mitigate the possible confounding effects of cognitive, visual, and motor
impairments on the outcomes of the study, it cannot be concluded that no confounding effects
occurred. In future studies, including a control condition matched to the experimental paradigm
where only the memory components are removed could better clarify this issue. Finally,
considering the aim of this study, which was to develop a practical apparatus for WM assessment
in clinical settings, we did not use an eye tracker. Although we attempted to minimize the effect
of eye movement in both paradigms by instructing the participants to fixate on a central fixation
point, using an eye tracker is necessary to validate these findings further.

In summary, using the resource-based model paradigms, we demonstrated that recall error and
precision were impaired in MS. We provided new insight regarding the progressive state of the
disease by assessing the plausible mechanisms related to the dissociable behavior of MGL and
sequential paradigms in classifying MS subtypes. Furthermore, applying a computational model
capable of disentangling error distribution in recalling information allowed us to uncover new
insight regarding WM deficit in the MS population. Our results determined that decreased signal-
to-noise ratio and swap error were responsible for WM deficit in MS. Overall, this study
provided a sensitive measure for assessing WM impairment and gave new insight into the
organization of WM dysfunction in MS.

Materials and Methods

1. Participants

409　A total of 121 patients with confirmed MS (61 RRMS and 60 SPMS) and 73 healthy control
410　volunteers participated in this study. Patients were recruited in a full-census manner from the
411　Comprehensive Multiple Sclerosis Clinic at Kashani Hospital in Isfahan between February 2021
412　and January 2022. The following criteria were used for participant selection: diagnosis based on
413　the 2017 McDonald criteria (Thompson et al., 2018), age between 18 to 55 years, diagnosis

within 1-18 years prior to entering the study, an expanded disability status scale (EDSS) score of 0-6.5, no history of clinical relapse or corticosteroid therapy within two months before entering the study, normal or corrected-to-normal visual acuity and color vision (based on the recorded bedside neurologic examination in their profile), and no impairment in factors that could interfere with the study including visual acuity, visual field, extraocular movement, scotoma, nystagmus, or tremor in the upper extremity, assessed as part of the EDSS examination. Additionally, normal performance in the Nine-Hole Peg Test (9-HPT < 45 seconds), the absence of a history of brain surgeries, major neurological disorders (stroke, epilepsy, brain tumor, or CNS infection), psychiatric disorders (major depressive disorder, bipolar disorder, or schizophrenia), drug or alcohol abuse, and chronic systemic disorders (diabetes, renal failure, liver failure, chronic obstructive pulmonary disorder (COPD), hypothyroidism, or hyperthyroidism) were required for participation in this study. Moreover, the control group should not have a family history of MS in their first-degree relatives.

2. Procedure

The study began with a clinical interview and neurologic examination to collect clinical and demographic information from the participants. Participants also completed 9-HPT and the Persian version of the MoCA test. MoCA is a standard cognitive screening tool (Badrkhahan et al., 2020; Nasreddine et al., 2005) with a scoring system ranging from 0 to 30, where distinct ranges correspond to different levels of cognitive function. Scores of $\geq$ 26 indicate normal cognitive performance, 18-25 denotes mild cognitive impairment, 10-17 determine moderate cognitive impairment, and $\leq$ 10 is considered severe impairment.

Participants received verbal and written instructions before beginning the assessment. A 10-trial training block was then performed to ensure their understanding of the tasks. After this, the actual assessment started.

Written informed consent was taken from all participants before the start of the study. This study followed the latest update of the Declaration of Helsinki ("World Medical Association Declaration of Helsinki," 2013) and was approved by the Iranian National Committee of Ethics in Biomedical Research (Approval ID: IR.MUI.MED.REC.1400.441).

3. Visual working memory paradigms

Visual WM was assessed using two analog recall tasks, memory-guided localization (MGL) and analog recall paradigms with sequential bar presentation. Stimuli were presented on a 15□ cathode ray tube (CRT, 75Hz refresh rate) monitor at a distancing view of 48cm. The paradigms were run in a dimly lit room on a computer with a Linux operating system and MATLAB software (MATLAB 2019a, The MathWorks, Inc., Natick, Massachusetts, USA) with Psychtoolbox 3 extension (Brainard, 1997; Pelli, 1997).

3.1. Memory-guided localization

Each trial was initiated by presenting a central fixation point (diameter of 0.51°) for 2 seconds (s), followed by the presentation of a target (a filled green circle with a diameter of 1.29°) for 1s. The target randomly appeared at different eccentricities (3.22°, 6.44°, or 9.66°) on each trial. In each block, targets were presented with equal probability at each eccentricity in random order (pseudo-random selection). While encouraging participants to maintain fixation on the central fixation point, participants were asked to memorize the location of the target circle for a delay period of 0.5, 1, 2, 4, or 8s (chosen pseudo-randomly). After the delay period, the fixation point changed from a circle to a cross, indicating the end of the delay period. Participants were asked to locate the target's position using the computer mouse and confirm their response by pressing the left button on the mouse. Subsequently, visual feedback was presented, showing them the correct position of the target and their response (Figure 1A). Participants completed six blocks of 30 trials. They also completed a 10-trial training block before the start of the study. Recall error (absolute error), Euclidian distance between the target's location and subject response in visual degree, and reaction time (RT) were recorded for further assessment.

3.2. Sequential paradigm with bar stimuli

Two designs of analog report tasks with sequential bar presentation, i.e., the low memory load condition (1-bar) and high memory load condition (3-bar), were developed to evaluate the visual WM deficit in MS. In the high memory load condition, each trial started with a small central fixation point (0.26°) for 2s, followed by a sequence of three distinguishable colored bars (red, green, and blue) at the center of the screen in a pseudo-random order. Each bar (2.57° by 0.19°) was presented for 500ms, followed by a 500ms blank interval. The minimum angular difference between the consecutively presented bars was 10 degrees. Participants were asked to memorize

both the orientation and color of the presented bars. After the bars were presented, a single bar, the 'probe bar' cued with the color of one of the presented bars, was displayed. Participants were asked to adjust the orientation of the probe bar, presented vertically, to match the orientation of the bar with the same color (target bar). To do that, they used a computer mouse and confirmed their response by clicking the right button. They received visual feedback, which showed the correct orientation of the target bar, their response, and the difference between them in angular degree (Figure 1B). The high memory load condition consisted of six blocks, each with 30 trials. The low memory load condition had the same structure as the high memory load condition except for presenting one bar instead of three (Figure 1C). After the high memory load condition, subjects participated in 30 trials of the low memory load condition. Due to the 1-bar design of low memory load condition, the swap error was absent, so fewer trials were needed (Peich et al., 2013). Before starting the paradigm, they also participated in a 10-trial training block with the same structure as the low memory load condition. The orientation of presented bars, subject response, recall error (absolute error), angular difference between the target value and subject response, and RT were recorded.

4. Statistical analysis

Statistical analyses were conducted using IBM SPSS Statistics for Mac, version 26 (IBM Corp., Armonk, N.Y., USA). The values were reported as mean ± standard deviation (SD). Data with extreme outliers (values greater than 3rd quartile + 3×interquartile range or less than 1st quartile – 3×interquartile range) in MGL and sequential paradigm with 3 bar were excluded from further analysis. The level of significance was set at p-value < 0.05.

Clinical and demographic profiles of the participants, except for the gender, cognitive ability, and treatment history, were compared using one-way ANOVA or Kruskal-Wallis H test (three groups comparison) and independent sample t-test or Mann-Whitney U test (two groups comparison). The post hoc Tukey's and Dunn's multiple comparison tests were performed following the significant results of ANOVA and Kruskal-Wallis H test. Also, Bonferroni correction for multiple tests was performed following Dunn's post hoc analysis, and the adjusted p-value was reported.

The cognitive ability of participants was classified according to the mentioned ranges of MoCA score. Patients' disease-modifying therapy (DMT) was classified into platform and non-platform

treatments. In our study, platform treatments include interferon beta-1a and glatiramer acetate,
and non-platform treatments contain rituximab, ocrelizumab, fingolimod, dimethyl fumarate, and
natalizumab. These categorical variables, along with gender, were compared using the Chi-
squared test.

For the MGL paradigm, recall error, recall precision (defined as the reciprocal of the standard
deviation of recall error), and RT were compared between groups (Healthy, RRMS, and SPMS)
among different conditions (distance or delay, mixed model ANOVA, between- and within-
subjects comparisons). For the sequential paradigm, since the data was in a circular space, based
on the method proposed by Fisher (Fisher, 1993), we used the analog report MATLAB toolbox
of Bays Lab (https://www.paulbays.com/toolbox/) to calculate the circular mean of recall error
for each subject. Similarly, recall precision, defined as the reciprocal of the circular standard
deviation of recall error, was calculated for each subject. For both high and low memory load
conditions (3-bar and 1-bar), recall error, recall precision, and RT were compared between
groups with respect to the order of the presented bars (mixed model and one-way ANOVA). To
further investigate the sources of error in recalling information and uncover the involved
mechanisms in visual WM impairment, the Mixture Model
(https://www.paulbays.com/toolbox/), a probabilistic model developed before (Bays et al., 2009;
Schneegans & Bays, 2016), was utilized. The Mixture Model considers three possible sources for
information recall. They are defined as the Gaussian variability in reporting the target and non-
target values and random responses (Bays et al., 2009; Schneegans & Bays, 2016). In our study,
they referred to reporting the orientation of the target bar (target proportion), misreporting the
orientation of the other two non-target bars instead of the target bar (swap error), and random
response (uniform proportion). Using the Mixture Model, the probabilities of target, non-target,
and uniform responses and the von Mises distribution concentration score were calculated for
each subject. The sources of error were evaluated by comparing the probability of target, non-
target, and uniform responses between groups among different bar orders (mixed model
ANOVA). Also, von Mises SD, defined as the circular standard deviation of the concentration
score of von Mises distribution, was assessed using the same method. Moreover, nearest
neighbor analysis was performed to further evaluate the isolated effect of orientation (Pertzov et
al., 2013). In this set of analyses, we removed the effect of swap error by defining recall error as
the difference between the subject response and the nearest presented bar in each trial. The

isolated effect of orientation was assessed between groups among different bar orders (mixed

model ANOVA). Finally, due to the 1-bar design of the low memory load condition, swap error

was not present; hence, only the target proportion, uniform proportion, and von Mises SD were

compared (one-way ANOVA).

For each comparison, hierarchical regression analyses were performed to evaluate the possible

confounding effect of demographic and clinical variables (significantly different between

groups) on results. Finally, the dissociable function of MGL and sequential paradigms in

distinguishing healthy control from MS patients, healthy control from MS subtypes, and MS

subtypes from each other were assessed by performing ROC analysis. Recall error was used for

classification purposes, and the area under the curve (AUC) was calculated as a measure of the

paradigms' accuracy in group classification.

Acknowledgment

We sincerely thank Farinaz Tabibian, Leila Sadat Razian, Maryam Mokhtari, and Zahra Mohamad Hoseiny for their cooperation in coordinating patients, Shahrzad Mohammadpour for advice about model fitting, and Mohammadamin Motaharinia for help regarding programming issues and his comments on the first draft of this manuscript.

Additional information

Funding

This study was supported by Isfahan University of Medical Sciences (grant number: 2400104). The funder had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

Ali Motahharynia, Conceptualization, Methodology, Software, Investigation, Formal analysis, Data curation, Visualization, Writing—original draft, Writing—review & editing; Ahmad Pourmohammadi, Conceptualization, Methodology, Software, Investigation, Data curation, Writing—review & editing; Armin Adibi, Investigation, Writing—review & editing; Vahid Shaygannejad, Resources, Writing—review & editing, Funding acquisition; Fereshteh Ashtari, Resources, Writing—review & editing, Funding acquisition; Iman Adibi, Conceptualization, Methodology, Resources, Formal analysis, Writing—review & editing, Supervision, Funding acquisition; Mehdi Sanayei, Conceptualization, Methodology, Software, Formal analysis, Writing—review & editing, Supervision

Competing interests

The authors declare that no competing interests exist.

Ethics

This study followed the latest update of the Declaration of Helsinki and was approved by the Iranian National Committee of Ethics in Biomedical Research (Approval ID: IR.MUI.MED.REC.1400.441).

Data availability

Anonymized data will be available upon request from corresponding authors. The corresponding authors will consider the request against the data-sharing policy in the protocol and ethical approval of the study.

Code availability

The source codes for the analog report MATLAB toolbox are available at https://www.paulbays.com/toolbox/.

References

Badrkhahan, S. Z., Sikaroodi, H., Sharifi, F., Kouti, L., & Noroozian, M. (2020). Validity and reliability of the Persian version of the Montreal Cognitive Assessment (MoCA-P) scale among subjects with Parkinson's disease. *Applied Neuropsychology: Adult*, *27*(5), 431–439. https://doi.org/10.1080/23279095.2019.1565762

Bastin, C., Besson, G., Simon, J., Delhaye, E., Geurten, M., Willems, S., & Salmon, E. (2019). An integrative memory model of recollection and familiarity to understand memory deficits. *Behavioral and Brain Sciences*, *42*, e281. https://doi.org/10.1017/S0140525X19000621

Bays, P. M., Catalao, R. F. G., & Husain, M. (2009). The precision of visual working memory is set by allocation of a shared resource. *Journal of Vision*, *9*(10), 7–7. https://doi.org/10.1167/9.10.7

Bays, P. M., & Husain, M. (2008). Dynamic Shifts of Limited Working Memory Resources in Human Vision. *Science*, *321*(5890), 851–854. https://doi.org/10.1126/science.1158023

Benedict, R. H. B., Amato, M. P., DeLuca, J., & Geurts, J. J. G. (2020). Cognitive impairment in multiple sclerosis: clinical management, MRI, and therapeutic avenues. *The Lancet Neurology*, *19*(10), 860–871. https://doi.org/10.1016/S1474-4422(20)30277-5

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436. http://www.ncbi.nlm.nih.gov/pubmed/9176952

Cecchini, M. A., Parra, M. A., Brazzelli, M., Logie, R. H., & Della Sala, S. (2022). Short-term memory conjunctive binding in Alzheimer's disease: A systematic review and meta-analysis. *Neuropsychology*. https://doi.org/10.1037/neu0000825

Clemens, L., & Langdon, D. (2018). How does cognition relate to employment in multiple sclerosis? A systematic review. *Multiple Sclerosis and Related Disorders*, *26*, 183–191. https://doi.org/10.1016/j.msard.2018.09.018

Costers, L., Van Schependom, J., Laton, J., Baijot, J., Sjøgård, M., Wens, V., De Tiège, X., Goldman, S., D'Haeseleer, M., D'hooghe, M. B., Woolrich, M., & Nagels, G. (2021). The role of hippocampal theta oscillations in working memory impairment in multiple sclerosis. *Human Brain Mapping*, *42*(5), 1376–1390. https://doi.org/10.1002/hbm.25299

Covey, T. J., Zivadinov, R., Shucard, J. L., & Shucard, D. W. (2011). Information processing speed, neural efficiency, and working memory performance in multiple sclerosis: Differential relationships with structural magnetic resonance imaging. *Journal of Clinical and Experimental Neuropsychology*, *33*(10), 1129–1145. https://doi.org/10.1080/13803395.2011.614597

Cowan, N. (2001). The magical number 4 in short-term memory: A reconsideration of mental storage capacity. *Behavioral and Brain Sciences*, *24*(1), 87–114. https://doi.org/10.1017/S0140525X01003922

Cree, B. A. C., Arnold, D. L., Chataway, J., Chitnis, T., Fox, R. J., Pozo Ramajo, A., Murphy, N., & Lassmann, H. (2021). Secondary Progressive Multiple Sclerosis. *Neurology*, *97*(8), 378–388. https://doi.org/10.1212/WNL.0000000000012323

Della Sala, S., Parra, M. A., Fabi, K., Luzzi, S., & Abrahams, S. (2012). Short-term memory binding is impaired in AD but not in non-AD dementias. *Neuropsychologia*, *50*(5), 833–840.

https://doi.org/10.1016/j.neuropsychologia.2012.01.018

Dobson, R., & Giovannoni, G. (2019). Multiple sclerosis – a review. *European Journal of Neurology*, *26*(1), 27–40. https://doi.org/10.1111/ene.13819

Figueroa-Vargas, A., Cárcamo, C., Henríquez-Ch, R., Zamorano, F., Ciampi, E., Uribe-San-Martin, R., Vásquez, M., Aboitiz, F., & Billeke, P. (2020). Frontoparietal connectivity correlates with working memory performance in multiple sclerosis. *Scientific Reports*, *10*(1), 9310. https://doi.org/10.1038/s41598-020-66279-0

Fisher, N. I. (1993). *Statistical Analysis of Circular Data*. Cambridge University Press. https://doi.org/10.1017/CBO9780511564345

Fujiwara, E., Kmech, J. A., Cobzas, D., Sun, H., Seres, P., Blevins, G., & Wilman, A. H. (2017). Cognitive Implications of Deep Gray Matter Iron in Multiple Sclerosis. *American Journal of Neuroradiology*, *38*(5), 942–948. https://doi.org/10.3174/ajnr.A5109

Gorgoraptis, N., Catalao, R. F. G., Bays, P. M., & Husain, M. (2011). Dynamic Updating of Working Memory Resources for Visual Objects. *Journal of Neuroscience*, *31*(23), 8502–8511. https://doi.org/10.1523/JNEUROSCI.0208-11.2011

Hulst, H. E., Goldschmidt, T., Nitsche, M. A., de Wit, S. J., van den Heuvel, O. A., Barkhof, F., Paulus, W., van der Werf, Y. D., & Geurts, J. J. G. (2017). rTMS affects working memory performance, brain activation and functional connectivity in patients with multiple sclerosis. *Journal of Neurology, Neurosurgery & Psychiatry*, *88*(5), 386–394. https://doi.org/10.1136/jnnp-2016-314224

Liang, Y., Pertzov, Y., Nicholas, J. M., Henley, S. M. D., Crutch, S., Woodward, F., Leung, K., Fox, N. C., & Husain, M. (2016). Visual short-term memory binding deficit in familial Alzheimer's disease. *Cortex*, *78*, 150–164. https://doi.org/10.1016/j.cortex.2016.01.015

Lugtmeijer, S., Schneegans, S., Lammers, N. A., Geerligs, L., de Leeuw, F. E., de Haan, E. H. F., Bays, P. M., & Kessels, R. P. C. (2021). Consequence of stroke for feature recall and binding in visual working memory. *Neurobiology of Learning and Memory*, *179*, 107387. https://doi.org/10.1016/j.nlm.2021.107387

Ma, W. J., Husain, M., & Bays, P. M. (2014). Changing concepts of working memory. *Nature Neuroscience*, *17*(3), 347–356. https://doi.org/10.1038/nn.3655

McMaster, J. M. V., Tomić, I., Schneegans, S., & Bays, P. M. (2022). Swap errors in visual working memory are fully explained by cue-feature variability. *Cognitive Psychology*, *137*, 101493. https://doi.org/10.1016/j.cogpsych.2022.101493

Miller, E. K., Lundqvist, M., & Bastos, A. M. (2018). Working Memory 2.0. *Neuron*, *100*(2), 463–475. https://doi.org/10.1016/j.neuron.2018.09.023

MILLER, G. A. (1956). The magical number seven plus or minus two: some limits on our capacity for processing information. *Psychological Review*, *63*(2), 81–97. http://www.ncbi.nlm.nih.gov/pubmed/13310704

Nasreddine, Z. S., Phillips, N. A., Bã©dirian, V., Charbonneau, S., Whitehead, V., Collin, I., Cummings, J. L., & Chertkow, H. (2005). The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment. *Journal of the American Geriatrics Society*, *53*(4), 695–699.

https://doi.org/10.1111/j.1532-5415.2005.53221.x

Nee, D. E., & D'Esposito, M. (2016). *The Representational Basis of Working Memory* (pp. 213–230). https://doi.org/10.1007/7854_2016_456

PARMENTER, B. A., SHUCARD, J. L., BENEDICT, R. H. B., & SHUCARD, D. W. (2006). Working memory deficits in multiple sclerosis: Comparison between the n -back task and the Paced Auditory Serial Addition Test. *Journal of the International Neuropsychological Society*, *12*(5), 677–687. https://doi.org/10.1017/S1355617706060826

Parra, M. A., Fabi, K., Luzzi, S., Cubelli, R., Hernandez Valdez, M., & Della Sala, S. (2015). Relational and conjunctive binding functions dissociate in short-term memory. *Neurocase*, *21*(1), 56–66. https://doi.org/10.1080/13554794.2013.860177

Peich, M.-C., Husain, M., & Bays, P. M. (2013). Age-related decline of precision and binding in visual working memory. *Psychology and Aging*, *28*(3), 729–743. https://doi.org/10.1037/a0033236

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: transforming numbers into movies. *Spatial Vision*, *10*(4), 437–442. http://www.ncbi.nlm.nih.gov/pubmed/9176953

Pertzov, Y., Miller, T. D., Gorgoraptis, N., Caine, D., Schott, J. M., Butler, C., & Husain, M. (2013). Binding deficits in memory following medial temporal lobe damage in patients with voltage-gated potassium channel complex antibody-associated limbic encephalitis. *Brain*, *136*(8), 2474–2485. https://doi.org/10.1093/brain/awt129

Pourmohammadi, A., Motahharynia, A., Shaygannejad, V., Ashtari, F., Adibi, I., & Sanayei, M. (2022). Working memory dysfunction differs between secondary progressive and relapsing multiple sclerosis: effects of clinical phenotype, age, disease duration, and disability. *Multiple Sclerosis and Related Disorders*, 104411. https://doi.org/10.1016/j.msard.2022.104411

Rocca, M. A., Barkhof, F., De Luca, J., Frisén, J., Geurts, J. J. G., Hulst, H. E., Sastre-Garriga, J., Filippi, M., Barkhof, F., Ciccarelli, O., De Stefano, N., Enzinger, C., Filippi, M., Frederiksen, J. L., Gasperini, C., Kappos, L., Palace, J., Rocca, M. A., Rovira, A., … Yousry, T. A. (2018). The hippocampus in multiple sclerosis. *The Lancet Neurology*, *17*(10), 918–926. https://doi.org/10.1016/S1474-4422(18)30309-0

Rocca, M. A., Valsasina, P., Hulst, H. E., Abdel-Aziz, K., Enzinger, C., Gallo, A., Pareto, D., Riccitelli, G., Muhlert, N., Ciccarelli, O., Barkhof, F., Fazekas, F., Tedeschi, G., Arévalo, M. J., & Filippi, M. (2014). Functional correlates of cognitive dysfunction in multiple sclerosis: A multicenter fMRI Study. *Human Brain Mapping*, *35*(12), 5799–5814. https://doi.org/10.1002/hbm.22586

Rolls, E. T. (2018). The storage and recall of memories in the hippocampo-cortical system. *Cell and Tissue Research*, *373*(3), 577–604. https://doi.org/10.1007/s00441-017-2744-3

Schneegans, S., & Bays, P. M. (2016). No fixed item limit in visuospatial working memory. *Cortex*, *83*, 181–193. https://doi.org/10.1016/j.cortex.2016.07.021

Schneegans, S., & Bays, P. M. (2017). Neural Architecture for Feature Binding in Visual Working Memory. *The Journal of Neuroscience*, *37*(14), 3913–3925. https://doi.org/10.1523/JNEUROSCI.3493-16.2017

Sicotte, N. L., Kern, K. C., Giesser, B. S., Arshanapalli, A., Schultz, A., Montag, M., Wang, H., & Bookheimer, S.

Y. (2008). Regional hippocampal atrophy in multiple sclerosis. *Brain*, *131*(4), 1134–1141.

https://doi.org/10.1093/brain/awn030

Stojanovic-Radic, J., Wylie, G., Voelbel, G., Chiaravalloti, N., & DeLuca, J. (2015). Neuroimaging and cognition

using functional near infrared spectroscopy (fNIRS) in multiple sclerosis. *Brain Imaging and Behavior*, *9*(2),

302–311. https://doi.org/10.1007/s11682-014-9307-y

Thompson, A. J., Banwell, B. L., Barkhof, F., Carroll, W. M., Coetzee, T., Comi, G., Correale, J., Fazekas, F.,

Filippi, M., Freedman, M. S., Fujihara, K., Galetta, S. L., Hartung, H. P., Kappos, L., Lublin, F. D., Marrie, R.

A., Miller, A. E., Miller, D. H., Montalban, X., … Cohen, J. A. (2018). Diagnosis of multiple sclerosis: 2017

revisions of the McDonald criteria. *The Lancet Neurology*, *17*(2), 162–173. https://doi.org/10.1016/S1474-

4422(17)30470-2

Vacchi, L., Rocca, M. A., Meani, A., Rodegher, M., Martinelli, V., Comi, G., Falini, A., & Filippi, M. (2017).

Working memory network dysfunction in relapse-onset multiple sclerosis phenotypes: A clinical-imaging

evaluation. *Multiple Sclerosis Journal*, *23*(4), 577–587. https://doi.org/10.1177/1352458516656809

Valdés Hernández, M. C., Clark, R., Wang, S.-H., Guazzo, F., Calia, C., Pattan, V., Starr, J., Della Sala, S., & Parra,

M. A. (2020). The striatum, the hippocampus, and short-term memory binding: Volumetric analysis of the

subcortical grey matter's role in mild cognitive impairment. *NeuroImage: Clinical*, *25*, 102158.

https://doi.org/10.1016/j.nicl.2019.102158

World Medical Association Declaration of Helsinki. (2013). *JAMA*, *310*(20), 2191.

https://doi.org/10.1001/jama.2013.281053

Zokaei, N., Burnett Heyes, S., Gorgoraptis, N., Budhdeo, S., & Husain, M. (2015). Working memory recall

precision is a more sensitive index than span. *Journal of Neuropsychology*, *9*(2), 319–329.

https://doi.org/10.1111/jnp.12052

Zokaei, N., Nour, M. M., Sillence, A., Drew, D., Adcock, J., Stacey, R., Voets, N., Sen, A., & Husain, M. (2019).

Binding deficits in visual short-term memory in patients with temporal lobe lobectomy. *Hippocampus*, *29*(2),

63–67. https://doi.org/10.1002/hipo.22998

Zokaei, N., Sillence, A., Kienast, A., Drew, D., Plant, O., Slavkova, E., Manohar, S. G., & Husain, M. (2020).

Different patterns of short-term memory deficit in Alzheimer's disease, Parkinson's disease and subjective

cognitive impairment. *Cortex*, *132*, 41–50. https://doi.org/10.1016/j.cortex.2020.06.016

Figure 1. Schematic design of visual working memory (WM) paradigms. (A) In the memory-

guided localization (MGL) paradigm, participants were asked to memorize and then localize the

position of the target circle following a random delay interval of 0.5, 1, 2, 4, or 8 seconds.

Following their response, visual feedback was presented. **(B)** In the sequential paradigm with 3

bar (high memory load condition), a sequence of three colored bars was presented consecutively.

Participants were asked to match the orientation of the probe bar to the previously presented bar

with the same color. Visual feedback was displayed following their response. **(C)** The 1-bar

paradigm (low memory load condition) has the same structure as the 3-bar paradigm except for

presenting one bar instead of three.

Figure 2. Recall error and precision of healthy control and multiple sclerosis (MS) subtypes (relapsing-remitting, RRMS and secondary progressive, SPMS) in visual WM paradigms. (**A**) Recall error, (**B**) recall precision, and (**C**) reaction time as a function of distance for the MGL paradigm. (**D, E, and F**) The same as a function of delay interval. (**G**) Recall error, (**H**) recall precision, and (**I**) reaction time as a function of bar order in the sequential paradigms with 3 bar (left of each subplot) and 1 bar (right of each subplot). Data are represented as mean ± SEM.

Figure 2—Source Data 1. Statistical reports corresponding to Figure 2.

Figure 3. The sources of recall error in high and low memory load conditions (3-bar and 1-bar, respectively). (A) von Mises SD (circular standard deviation of von Mises distribution), **(B)** Target response (probability of response around the target value), **(C)** swap error (probability of response around the non-target values), and **(D)** uniform response (probability of random response) for healthy control and MS subtypes in the sequential paradigms with 3 bar (left of each subplot) and 1 bar (right of each subplot). Data are represented as mean ± SEM.

Figure 3—figure supplement 1. Isolated effect of orientation in the high and low memory load conditions. The nearest neighbor analysis determined the isolated effect of orientation for healthy control, RRMS, and SPMS in the high memory load condition (left of each subplot). The effect of orientation for the same groups in the low memory load condition (right of each subplot).

810   **Figure 3—Source Data 1.** Statistical reports corresponding to Figure 3 and Figure 3—figure

811   supplement 1.

835   **Figure 4. Classifying performance of visual WM paradigms in differentiating healthy**

836   **control from MS and MS subtypes, and MS subtypes from each other.** Receiver operating

837   characteristic (ROC) curve demonstrated the accuracy of **(A)** MGL and sequential paradigms

838   with **(B)** 3 bar and **(C)** 1 bar in distinguishing healthy control from MS patients. The precision of

839   these paradigms in dissociating healthy control from MS subtypes (RRMS and SPMS) and MS

840 subtypes from each other is represented as the area under the curve (AUC) for **(D)** MGL and

841 sequential paradigms with **(E)** 3 bar and **(F)** 1 bar.

842 **Figure 4—Source Data 1.** Statistical reports corresponding to Figure 4.

843 **Table 1. Demographic and clinical profiles of participants in the MGL paradigm.**

	HC (n = 23)	RRMS (n = 16)	SPMS (n = 25)	*P*
Gender (F:M)	13:10	14:2	17:8	0.12
Age (year)	35.9 ± 8.34	37.25 ± 6.63	39.28 ± 5.56	0.25
Education (years)	13.30 ± 2.74	13.69 ± 3.34	13.56 ± 3.22	0.86
Cognitive ability[a] (NL:MCI)	23:0	14:2	19:6	< 0.05 *
Disease duration (years)	N/A	8.562 ± 3.20	11.56 ± 3.28	< 0.02 *
EDSS	N/A	1.28 ± 0.79	2.740 ± 1.23	< 0.0002 *
DMT (Platform: Non-platform)	N/A	2:14	0:25	0.07

845 All data, except for the categorical information, are presented as mean ± standard deviation.

846 [a] assessed based on the Montreal cognitive assessment (MoCA) test classification.

847 Gender, cognitive ability, and DMT were compared using the chi-squared test. Age and education were compared

848 using one-way ANOVA and Kruskal-Wallis H test, respectively. Disease duration and EDSS score were compared

849 using Mann-Whitney U test and independent sample t-test, respectively.

850 HC = Healthy control, RRMS = Relapsing-remitting multiple sclerosis, SPMS = Secondary progressive multiple

851 sclerosis, NL = Normal (MoCA score ≥ 26), MCI = Mild cognitive impairment (MoCA score = 18-25), EDSS =

852 Expanded disability status scale, DMT = Disease-modifying therapy, Platform treatment = interferon beta-1a and

853 glatiramer acetate, Non-platform treatment = rituximab, ocrelizumab, fingolimod, dimethyl fumarate, and

854 natalizumab, N/A = Not applicable

855 ***P* < 0.05**

866 **Table 2. Demographic and clinical profiles of participants in the sequential paradigms.**

	HC (n = 46)	RRMS (n = 39)	SPMS (n = 32)	*P*
Gender (F:M)	16:30	23:16	22:10	< 0.008 *
Age (year)	30.5 ± 10.37	32.03 ± 6.72	39.00 ± 6.43	< 10^{-6} *
Education (years)	16.95 ± 2.23	13.87 ± 3.41	13.67 ± 2.73	< 10^{-7} *
Cognitive ability[a] (NL:MCI)	42:4	28:11	18:13	< 0.003 *
Disease duration (years)	N/A	6.60 ± 3.84	9.37 ± 4.43	< 0.007 *
EDSS	N/A	1.49 ± 1.01	3.86 ± 1.74	< 10^{-7} *
DMT (Platform: Non-platform)	N/A	5:34	1:31	0.14

All data, except for the categorical information, are presented as mean ± standard deviation.

[a] assessed based on the Montreal cognitive assessment (MoCA) test classification. One value for SPMS is missing.

Gender, cognitive ability, and DMT were compared using the chi-squared test. Age and education were compared

using the Kruskal-Wallis H test. Disease duration and EDSS score were compared using an independent sample t-

test and Mann-Whitney U test, respectively.

Dunn's post hoc test was performed following significant results of age and education, and the adjusted p-value

following Bonferroni correction for multiple tests are reported:

Age: Healthy vs. RRMS: $P = 0.27$, Healthy vs. SPMS: $P < 10^{-6}$*, and RRMS vs. SPMS: $P < 0.005$*.

Education: Healthy vs. RRMS: $P < 10^{-5}$*, Healthy vs. SPMS: $P < 10^{-5}$*, and RRMS vs. SPMS: $P = 1$.

HC = Healthy control, RRMS = Relapsing-remitting multiple sclerosis, SPMS = Secondary progressive multiple

sclerosis, NL = Normal (MoCA score ≥ 26), MCI = Mild cognitive impairment (MoCA score = 18-25), EDSS =

Expanded disability status scale, DMT = Disease-modifying therapy, Platform treatment = interferon beta-1a and

glatiramer acetate, Non-platform treatment = rituximab, ocrelizumab, fingolimod, dimethyl fumarate, and

natalizumab, N/A = Not applicable

*P < 0.05

Appendix 1—table 1. Hierarchical regression analysis for MGL paradigm.

	Statistical reports
Independent variables	**Recall parameters** **Error/ Precision (distance)/ Precision (delay) (F, P)**
Group	$(14.57, < 10^{-5}*)$/ $(13.74, <10^{-4}*)$/ $(14.57, < 10^{-5}*)$
Group Cognitive ability	$(10.91, < 10^{-4}*)$/ $(9.7, < 10^{-3}*)$/ $(10.62, < 10^{-3}*)$ $(24.01, < 10^{-5}*)$/ $(6.29, <0.02*)$/ $(5.59, < 0.03*)$

886 Cognitive ability: assessed based on the Montreal cognitive assessment (MoCA) test classification.

887 MGL = Memory-guided localization

888 *$P < 0.05$

895 **Appendix 1—table 2. Hierarchical regression analysis for sequential paradigm with 3 bar, high memory load**
896 **condition.**

	Statistical reports
Independent variables	**Recall parameters** **Error/ Precision/ von Mises SD/ Target proportion/ Swap error/ Uniform proportion / Nearest neighbor (F, P)**
Group	$(28.18, < 10^{-9}*)$ / $(25.23, < 10^{-9}*)$/ $(26.79, < 10^{-9}*)$/ $(11.04, < 10^{-4}*)$/ $(7.11, < 0.002*)$/ $(5.50, < 0.006*)$/ $(29.26, < 10^{-10}*)$
Group Gender	$(22.47, < 10^{-8}*)$/ $(19.90, < 10^{-7}*)$/ $(20.37, < 10^{-7}*)$/ $(9.98, < 0.0002*)$/ $(7.22, < 0.002*)$/ $(4.38, < 0.02*)$/ $(22.64, < 10^{-8}*)$ $(3.89, = 0.051)$/ $(4.16, < 0.05*)$/ $(8.71, < 0.004*)$/ $(0.002, = 0.97)$/ $(0.47, = 0.49)$/ $(0.57, = 0.45)$/ $(7.82, < 0.007*)$
Group Age	$(18.84, < 10^{-7}*)$/ $(17.68, < 10^{-6}*)$/ $(18.53, < 10^{-6}*)$/ $(6.76, < 0.002*)$/ $(6.32, < 0.003*)$/ $(2.85, = 0.06)$/ $(19.60, < 10^{-7}*)$ $(9.39, < 0.003*)$/ $(7.80, < 0.007*)$/ $(5.51, < 0.03*)$/ $(5.24, < 0.03*)$/ $(0.07, = 0.79)$/ $(15.04, < 0.0002*)$/ $(12.04, < 10^{-3}*)$
Group Education	$(14.03, < 10^{-5}*)$/ $(12.79, < 10^{-4}*)$/ $(15.22, < 10^{-5}*)$/ $(4.17, < 0.02*)$/ $(4.33, < 0.02*)$/ $(0.90, = 0.41)$/ $(14.2, < 10^{-5}*)$ $(8.21, < 0.005*)$/ $(5.46, < 0.03*)$/ $(3.21, = 0.08)$/ $(6.64, < 0.02*)$/ $(0.52, = 0.47)$/ $(10.08, < 0.002*)$/ $(9.57, 0.003*)$
Group Cognitive ability[a]	$(19.26, < 10^{-7}*)$/ $(18.23, < 10^{-6}*)$/ $(19.69, < 10^{-7}*)$/ $(5.71, < 0.005*)$/ $(3.41, < 0.04*)$/ $(2.95, = 0.056)$/ $(20.93, < 10^{-7}*)$ $(14.91, < 10^{-3}*)$/ $(6.81, < 0.02*)$/ $(5.68, 0.02*)$/ $(16.10 < 10^{-3}*)$/ $(11.82, < 10^{-3}*)$/ $(5.73, < 0.02*)$/ $(8.94, < 0.004*)$

898 Cognitive ability: assessed based on the Montreal cognitive assessment (MoCA) test classification.

899 [a] One MoCA value in SPMS group is missing.

900 *$P < 0.05$

Appendix 1—table 3. Hierarchical regression analysis for sequential paradigm with 1 bar, low memory load condition.

	Statistical reports
Independent variables	**Recall parameters** **Error/ Precision/ von Mises SD/ Target or Uniform proportion (F, P)**
Type	$(36.85, < 10^{-12}*)$/ $(25.48, < 10^{-9}*)$/ $(33.07, < 10^{-11}*)$/ $(3.11, < 0.05*)$
Group Gender	$(29.46, < 10^{-10}*)$/ $(20.05, < 10^{-7}*)$/ $(26.92, < 10^{-9}*)$/ $(1.97, = 0.14)$ $(8.09, < 0.006*)$/ $(4.41, < 0.04*)$/ $(3.45, = 0.07)$/ $(2.44, = 0.12)$
Group Age	$(23.85, < 10^{-8}*)$/ $(18.48, < 10^{-6}*)$/ $(22.03, < 10^{-8}*)$/ $(1.87, = 0.16)$ $(8.27, < 0.005*)$/ $(1.57, = 0.21)$/ $(6.89, < 0.001*)$/ $(0.57, = 0.45)$
Group Education	$(23.04, < 10^{-8}*)$/ $(15.41, < 10^{-5}*)$/ $(18.65, < 10^{-6}*)$/ $(2.59, = 0.08)$ $(4.03, < 0.05*)$/ $(2.42, = 0.12)$/ $(6.36, < 0.02*)$/ $(0.01, = 0.91)$
Group Cognitive ability[a]	$(27.35, 10^{-9}*)$/ $(20.6, < 10^{-7}*)$/ $(25.54, < 10^{-9}*)$/ $(1.87, = 0.16)$ $(4.46, < 0.04*)$/ $(0.35, = 0.56)$/ $(5.25, < 0.03*)$/ $(0.08, = 0.77)$

Cognitive ability: assessed based on the Montreal cognitive assessment (MoCA) test classification.

[a] One MoCA value in SPMS group is missing.

*$P < 0.05$

926   **Appendix 2—table 1. Statistical results of reaction time (in second) for MGL paradigm.**

	Statistical results
Healthy	1.59 ± 0.41
RRMS	2.33 ± 0.65
SPMS	3.02 ± 0.87
Between groups (mixed-model ANOVA)	$F(2,61) = 26.44, P < 10^{-8}*$
Healthy vs. RRMS (Tukey)	$P < 0.004*$
Healthy vs. SPMS (Tukey)	$P < 10^{-8}*$
RRMS vs. SPMS (Tukey)	$P < 0.008*$
Within groups (Distance, mixed-model ANOVA)	$F(2,61) = 25.94, P < 10^{-9}*$
Interaction (Groups $\times$ Distance, mixed-model ANOVA)	$F(4,61) = 2.06, P = 0.09$
Within groups (Delay, mixed-model ANOVA)	$F(4,61) = 0.97, P = 0.43$
Interaction (Groups $\times$ Delay, mixed-model ANOVA)	$F(8,61) = 0.86, P = 0.55$

927
928   MGL = Memory-guided localization, RRMS = Relapsing-remitting multiple sclerosis, SPMS = Secondary

929   progressive multiple sclerosis.

930   Data are represented as mean $\pm$ standard deviation.

931   *$P < 0.05$
932

936   **Appendix 2—table 2. Statistical results of reaction time (in second) for sequential paradigms (3 bar and 1**
937   **bar)**

	Statistical results (3-bar)	Statistical results (1-bar)
Healthy	2.62 ± 1.01	2.25 ± 0.92
RRMS	3.14 ± 0.96	2.78 ± 1.04
SPMS	4.18 ± 2.00	3.66 ± 1.72
Between groups (mixed-model ANOVA)	$F(2,114) = 12.95, P < 10^{-5}*$	$F(2,114) = 12.59, P < 10^{-4}*$
Healthy vs. RRMS (Tukey)	$P = 0.17$	$P = 0.11$
Healthy vs. SPMS (Tukey)	$P < 10^{-5}*$	$P < 10^{-5}*$
RRMS vs. SPMS (Tukey)	$P < 0.005*$	$P < 0.009*$
Within groups (Bar order, mixed-model ANOVA)	$F(2,114) = 5.92, P < 0.004*$	N/A
Interaction (Groups $\times$ Bar order, mixed-model ANOVA)	$F(4,114) = 0.16, P = 0.96$	N/A

939   RRMS = Relapsing-remitting multiple sclerosis, SPMS = Secondary progressive multiple sclerosis.

940   Data are represented as mean $\pm$ standard deviation.

941   *$P < 0.05$

A

MGL paradigm
6 blocks of 30 trials

Fixation point
2 sec

Target circle
1 sec

Varied delay interval
0.5-8 sec

Feedback
2 sec

B

Sequential paradigm with 3 bar
6 blocks of 30 trials

Fixation point
2 sec

Presentation time
0.5 sec

Blank interval
0.5 sec

Presentation time
0.5 sec

Blank interval
0.5 sec

Presentation time
0.5 sec

Blank interval
0.5 sec

Delay interval
1 sec

Probe bar

Feedback
2 sec

C

Sequential paradigm with 1 bar
30 trials