Human-aided dispersal and population bottlenecks facilitate parasitism escape in the most invasive mosquito species

Maxime Girard*[1], Edwige Martin[1], Laurent Vallon[1], Van Tran Van[1], Camille Da Silva Carvalho[1], Justine Sack[1], Zélia Bontemps[1,2], Julie BaltenNeck[1,3] , Florence Colin[1], Pénélope Duval[1], Simon Malassigné[1], Ian Hennessee[5], Lucrecia Vizcaino[6], Yamila Romer[4], Nsa Dada[7,8], Khan Ly Huynh Kim[9], Trang Huynh Thi Thuy[9], Christophe Bellet[10], Gregory Lambert[11], Fara Nantenaina Raharimalala[12], Natapong Jupatanakul[13], Clement Goubert[14], Matthieu Boulesteix[15], Patrick Mavingui[16], Emmanuel Desouhant[15], Patricia Luis[1], Rémy Cazabet[17], Anne-Emmanuelle Hay[1], Claire Valiente Moro[1], Guillaume Minard[1]

[1]Univ. Lyon, Université Claude Bernard Lyon 1, CNRS, INRAe, VetAgro Sup, UMR Ecologie Microbienne, F-69622 Villeurbanne, France.
[2]Department of Medical Biochemistry and Microbiology, Science for Life Laboratories, Uppsala University, Uppsala, Sweden.
[3]Université de Lyon, INSA Lyon, Université Claude Bernard Lyon 1, CNRS, UMR 5240 MAP, Microbiologie, Adaptation, Pathogénie, F-69622 Villeurbanne, France.
[4]Emory College of Arts and Science, Emory University, Atlanta, USA.
[5]Rollins School of Public Health, Emory University, Atlanta, USA.
[7]Centre de Recherche pour la lutte contre les Maladies Infectieuses Tropicales/Tropicales Infectious Diseases Research Centre, Université d'Abomey-Calavi, Cotonou, Benin.
[8]School of Life Sciences, Arizona State University, Tempe, AZ, USA
[9]Department of Medical Entomology and Zoonotics, Pasteur Institute in Ho Chi Minh City, Vietnam.
[10]Entente Interdépartementale Rhône-Alpes pour la Démoustication, Chindrieux, France.
[11]Entente Interdépartementale de Démoustication du Littoral Méditerranéen, Montpellier, France.
[12]Pasteur Institute, Antananarivo, Madagascar.
[13]National Center for Genetic Engineering and Biotechnology (BIOTEC), Pathum Thani, Thailand.
[14]Canadian Centre for Computational Genomics, McGill Genome Centre, Human Genetics, McGill University, Montreal, QC, Canada.
[15]Université de Lyon, Université Claude Bernard Lyon 1, Laboratoire de Biométrie et de Biologie Evolutive, UMR CNRS 5558, VetAgro Sup, Villeurbanne, France.
[16]UMR PIMIT, Processus Infectieux en Milieu Insulaire Tropical, CNRS 9192, INSERM U1187, IRD 249, Université de La Réunion, Sainte-Clotilde, La Réunion, France.
[17]Univ de Lyon, CNRS, Université Lyon 1, LIRIS, UMR5205, F-69622 Villeurbanne, France.

Running title: Human-aided parasitism escape in mosquitoes

Contact information: Maxime Girard, 8 Rue Raphael Dubois 69100 VILLEURBANNE, France, phone: +33472431143, email: maxime.girard@univ-lyon1.fr

Keywords: Parasitism escape, Invasion, Human transportation, *Aedes albopictus*, *Ascogregarina taiwanensis*, Biogeography, Microbial communities, Mosquito

Abstract.

During biological invasion process, species encounter new environments and partially escape some ecological constraints they faced in their native range, while they face new ones. The Asian tiger mosquito *Aedes albopictus* is one of the most iconic invasive species introduced in every inhabited continent due to international trade. It has also been shown to be infected by a prevalent yet disregarded microbial entomoparasite *Ascogregarina taiwanensis*. In this study, we aimed at deciphering the factors that shape the global dynamics of *As. taiwanensis* infection in natural *Ae. albopictus* populations. We showed that *Ae. albopictus* populations are highly colonized by several parasite genotypes but recently introduced ones are escaping it. We further performed experiments based on the invasion process to explain such pattern. To that end, we hypothesized that (i) mosquito passive dispersal (*i.e.* human-aided egg transportation) may affect the parasite infectiveness, (ii) founder effects (*i.e.* population establishment by a small number of mosquitoes) may influence the parasite dynamics and (iii) unparasitized mosquitoes are more prompt to found new populations through active flight dispersal. The two first hypotheses were supported as we showed that parasite infection decreases over time when dry eggs are stored and that experimental increase in mosquitoes' density improves the parasite horizontal transmission to larvae. Surprisingly, parasitized mosquitoes tend to be more active than their unparasitized relatives. Finally, this study highlights the importance of global trade as a driver of biological invasion of the most invasive arthropod vector species.

Significance.

Global trade expansion has facilitated the introduction of invasive species such as the Asian tiger mosquito *Aedes albopictus*. Eventually, invasive species might escape their natural enemies and this phenomenon exemplifies their invasion success. In this study, we combined field observations and laboratory experiments to decipher the ecological consequences of the invasion process on the interaction dynamics between *Ae. albopictus* and its most prevalent natural parasite *As. taiwanensis*. We observed a decrease in parasitism in recently introduced populations and provide experimental evidence to explain how human-aided mosquito transportation and mosquito population bottlenecks were a burden for the parasite.

Introduction

Despite repeated calls on actions to prevent drastic consequences of global change, our society is now facing many ecological challenges that could profoundly affect its future (1). The cumulative effects of biological invasion and human activities are a threat to biodiversity due to change in global species distributions but also species loss and drift in community composition related to parasite transmission or competitive exclusion (2–7). Introduced species tend to raise health, agricultural, goods and services costs that weaken the global economy (8,9). Finally, some invasive species are considered as a threat to human health due to their pathogenicity or their ability to transmit pathogens (10). Currently, 37% of the 66 most-studied invasive species are considered as a threat while uncertainties remain on others (11). During biological invasion, invaders may be accompanied by parasites that may originate from their native range or an intermediate location (12). Co-introduction of invasive species with their parasites can lead to native species infection also referred as host shifting when the infection concerns different host species or spillover when the infection concerns different host populations (13). Invasive species may also acquire the parasite from native species. Finally, following introduction, invasive species may also escape the parasite colonizing them in their native or intermediate areas (14). Those events could contribute to exacerbate the performance and invasion success of recently introduced host species.

Both stochastic and selective pressures may lead to parasite release (15). Subsampling of individuals in the native or intermediate area during the invasion process may lead to uninfected newly formed populations. This is particularly true when the parasite prevalence is low within the source population (16,17). Furthermore, unsuitable conditions during the invasion path or colonization process may also lead to differential loss of the parasite due to *e.g.* unsuitable transportation conditions, higher host resistance at the invasion front and non-permissive environmental conditions (18). Other studies suggest that demographic changes may also alter the parasite transmission among individuals at the invasion front (19,20). Indeed, new populations are often founded by a small number of individuals that may be less prompt to exchange parasite due to lower horizontal transmissions and parasite virulence (21). In some cases, lower inter-individual transmission due to low population densities is counter balanced by a plastic change in host permissiveness toward parasites named the Density Dependent Prophylaxis (DDP) (22). In such cases, the immune system is relaxed in conditions of low host densities and reciprocally upregulated in conditions of high host densities. It is therefore expected that parasitism release is more likely to be exemplified if reduction in population densities occurs in absence of DDP. Finally, parasitism release may be the result of a parasite-driven counter selection on the host propensity to disperse. As an

example, the parasite *Ophryocystis elektroscirrha* decreases flight performance of the
Monarch butterfly *Danaus plexippus* reducing the chance of infected individuals to migrate
(23).

The Asian tiger mosquito *Aedes albopictus* is one of the most invasive species and
poses threats for human health (24) due to its ability to replicate and transmit at least 19 viruses
such as Chikungunya, Dengue or Zika, as well as human and animal filarial nematodes
(25,26). Originating from South and East Asia, its distribution has rapidly increased all over the
world since 1980's (27). Many studies pointed out the strong influence of human-aided
transportation on intermediate and long distance dispersal of desiccated mosquito eggs and,
in a lesser extent, adults (27–33). Actually, *Ae. albopictus* mosquitoes are often transported as
desiccated eggs on attractive containers (*e.g.* used tires and lucky bamboo) (28,34–39). Eggs
hatch whenever the containers get flooded after being introduced in a new area. Under optimal
conditions, previous studies have shown that *Ae. albopictus* eggs can survive a month-long
desiccation period (40). However, due to harsh transports conditions and population
subsampling, the demography of recently introduced mosquito populations is often small
(41,42). Such founder effects involve genetic drift, genetic bottlenecks and reductions in
population size at the invasion front; however, these populations were shown to experience
rapid growth in subsequent years (43,44,42,45,46). The final step of the invasion process is
the local spread of mosquitoes and is both mediated either *via* flight or *via* passive
transportations again (42,47).

Ae. albopictus populations are highly infected by an Apicomplexan gregarine parasite
belonging to the *Ascogregarina* genus (48). Various species of *Ascogregarina* are colonizing
mosquitoes with host specificity (49). *As. taiwanensis* is specifically associated with *Ae.*
albopictus even though rare spillover events toward non-host mosquito species were reported
(50–54). Its biological cycle is synchronized with the mosquito development (55). Oocysts
released outside the eggshell are ingested by first instar larvae in water containers and
replicate into developing individuals until adults release them. This parasite can be horizontally
transmitted either when adults die, defecate or emerge within larval habitats. *As. taiwanensis*
may have strong ecological consequences for its mosquito host although it is considered as a
weak parasite (56). It was shown to have detrimental effects on mosquito traits such as
development and survivorship at high density or when food is limited (57–59). A previous study
conducted in Florida (USA) suggested that *Ae. albopictus* may escape *As. taiwanensis* at the
invasion front (60). The authors demonstrated a drastically reduced parasite prevalence in
recently introduced mosquito populations (<3 years). To reinforce those preliminary
observations, we have conducted a field study in mainland France showing a similar trend of
parasitism release at the invasion front. To estimate whether stochastic selection of uninfected

individuals was likely to occur, we have also tested older populations in the native range and

intermediate locations. We showed that the parasite is one of the most prevalent and dominant

members of the mosquito microbiota in settled populations among contrasted locations across

the world. Considering that stochastic selection of uninfected individuals was unlikely, we have

thus conducted hypothesis driven experiments to test factors that may explain the observed

release from the parasite pressure during biological invasion.

Our experiments have been designed to answer three questions based on the mosquito

invasion process: (i) Does long distance dissemination through human-aided egg

transportation limit the parasite maintenance? (ii) Does the decrease in host density following

introduction affect the parasite dissemination probability in absence of DDP? and (iii) Does the

parasite limit mosquito local active dissemination through flight? To answer those questions,

we first tested the effect of desiccation on the parasite maintenance to reflect conditions

occurring during long distance dissemination. Then, we tried to reproduce the decrease in

mosquito populations reflecting founder effect through two different experiments to assess the

impact of both adult and larvae densities on the parasite infection success. Finally, the impact

of the parasitism on mosquito flight has been assessed through mosquito movement counting.

Combined with previously published data, our results support that population density decrease

following founding events and human-aided transportation contribute both to parasitism

escape. Furthermore, our results suggest that the parasite *de novo* colonizes the mosquito

with high densities two years after introduction.

Material and methods

Rearing parasitized and unparasitized mosquito lines

A total of ~100 larvae were collected in 2017 in Villeurbanne (N: 45°46'18990'' E:

4°53'24615'') and Pierre-Bénite (N: 45°42'11534'' E: 4°49'28743'') in France. This population

named AealbVB was grown into larger populations (~5000 individuals) for 15 generations in a

Biosafety level 2 insectary at 28°C, with a relative humidity of 80% and a day/night cycle of

18h/6h. Once the eggs were laid, they were immediately placed into water and ~100 adults

individuals were crushed and spread to make sure that the parasite was loaded. The water

was never changed until the individuals molt into the 2nd instar since individuals get infected

during the 1st instar. This protocol pursued for each generation. Uninfected individuals originate

from the same initial population. To ensure that the parasite did not infect them, several actions

were performed to shunt the infection path. First, the eggs were not immersed immediately but

water was removed and eggs were kept dry at least 7 days before being flooded again to avoid

direct transmission of this parasite when it is released by adults in water. Then, no contact was

permitted between infected individuals and the water in which larvae were reared to avoid any

direct transmission from feces or adult bodies. Finally, as the infection occurs during the first
stage of larval development, the water was changed during the first instar to reduce the
transmission probability. The absence of infection in the unparasitized population and
presence of infection in the parasitized population was regularly checked by crushing adult
individuals in water and observing the mixture at a magnification of x400 on an optical
microscope (Leica).

Sample collection and prevalence estimation

Live mosquito samples were collected with aspirators, net and BG-traps in 7 countries
and 17 sites. They were identified using morphological characteristics (61), stored in 70%
ethanol during transportation and frozen until use. Information related to site, country, year of
collection, sex, climate, GPS coordinates and the population age (*i.e.* time between population
introduction and sample collections) are reported in **Table S1**. This last information was
obtained from local entomological surveillance services in France (EID and EIRAD), as well as
from published reports in Italy (62), Spain (63) and USA (64). The date of introduction in
Madagascar is relatively uncertain and was estimated to be older than 1904 (65). Populations
from Thailand and Vietnam were considered as natives. DNA was extracted from surface
sterilized mosquito whole bodies and PCR diagnostic were performed to determine the
parasite prevalence (see **Supplementary material section**). To complete those dataset,
supplemental data of *As. taiwanensis* presence-absence were extracted from previously
published studies corresponding to mosquitoes collected in the USA (60,66,57,67) either from
the manuscript or from graph by using the ImageJ software v.1.54f (68). Data are available
on **Table S2**.

Metabarcoding of eukaryotic and prokaryotic microbial communities

Using DNA extracted from surface sterilized mosquitoes, prokaryotic and eukaryotic
communities within entire individuals were characterized through high throughput sequencing
of hypervariable regions from the 16S rDNA and 18S rDNA genes respectively. The V5-V6
region of 16S rDNA gene was amplified using the primers 784F (5'-AGG ATT AGA TAC CCT
GGT A-3') and 1061R (5'-CRR CAC GAG CTG ACG AC-3'). The V1-V2 region of the 18S
rDNA was amplified using the primers Euk82F (5′ GAA ACT GCG AAT GGC TC 3') and
Euk516R (5′ ACC AGA CTT GCC CTC C 3'). The details concerning PCR cycles are presented
in supplementary materials. The PCR products were then sent to Biofidal sequencing company
for purification and 2x300bp Miseq sequencing (Illumina). A total of 12,836,921 and
14,248,382 reads were obtained from 16S and 18S rDNA respectively and were demultiplexed
using the Mothur pipeline (69) that is described in supplementary materials. After removing
singletons and subsampling, 5,886 and 5,264 OTUs were retrieved respectively for 16S rDNA

and 18S rDNA. Contaminant OTUs were removed if their proportion in samples were not at
least 10-fold in comparison with controls. Finally, after quality control, data were homogenized
by sampling 1,000 and 500 reads for 16S rDNA and 18S rDNA, respectively. The data have
finally been converted into relative abundance, hereafter define as the proportion of reads
associated to each OTU in comparison to the total number of reads per sample. Since relative
abundances are of compositional nature, we have randomly selected 7 samples from each sex
and group of microbial similarities for which we conducted 16S and 18S rDNA qPCR to convert
relative abundances into absolute abundances *i.e.* number of copies associated with the given
OTU per mosquito sample (see more details in **Supplementary materials**). Miseq sequences
have been deposited on Zenodo under the project number 8252320
(https://zenodo.org/record/8252320).
239
Testing the ability of parasite maintenance during passive long-distance transportation
on desiccated eggs
As previously described, desiccated mosquito eggs can be transported by human
transportations during weeks. However, we do not know if such transportation could affect the
parasite. Since *Ae. albopictus* tends to prefer dark container (70), we let infected mosquitoes
lay eggs on water containers coated with a dark green blotting paper and preserved them in
the dark under optimal conditions (*i.e.* 28°C with a relative humidity of 80%). After respectively
35h, 1 week and 1 month, 100 eggs were harvested from the blotting paper and placed in a
water container to hatch. Larvae were reared at 28°C and fed 1/3 yeast (Biovers) and 2/3 fish
food and was provided *ad libitum* to the larvae. The adults that emerged from those containers
were collected. The number of parasites colonizing the adult mosquitoes was estimated with
qPCR on 10 males and females per time point and replicate. Primers and protocol for *As.*
taiwanensis specific qPCR were developed for this study to determine the absolute abundance
of the parasite (*i.e.* number of the parasite 18S copy per mosquito) and are detailed in the
supplementary material. The experiment was reproduced five times. This experiment was
repeated five times on five different blotting papers.
256
Testing the influence of mosquito densities on inter-individual transmission
Water containers were maintained with 50ml of sterile water in cages containing either
100 (low density) or 3000 (high density) infected adult individuals during a week to allow them
to release oocysts in water. The experiment was replicated 5 times for each condition (*i.e.* 5
donor mosquito cages were used for each tested mosquito density). Eggs from an
unparasitized population were hatched in sterile water and 50 larvae were added to the
container. Larvae were reared in similar conditions to the previously described experiment.

Adult males and females were collected at emergence and their parasite density was
measured with qPCR using 4 to 10 individuals per sex and replicate.

Testing Density Dependent Prophylaxis (DDP)

To test whether the inter-individual transmission could be related to the density of larvae
into the breeding sites, we hatched > 1000 larvae in containers to let them get infected with
the parasite released from their parents. After 2 days, we then harvested them and grew them
in new containers filled with distilled water at different densities (either 20, 50 or 100 larvae per
100ml of water). This protocol was replicated 5 times for each condition. Larvae were reared
on a shelve providing shadow. When adults emerged, they were collected and parasites
quantification was performed with qPCR (5 males and 5 females per replicate and larval
density). For technical reasons, highest density modality could not be performed for the 5[th]
replicate (*i.e.* poor hatching rate for this replicate). Larval rearing conditions were similar to
those in the previous experiment.

Testing the influence of the parasite on active local mosquito dispersal through flight

Absence of the parasite in recently introduced mosquito populations might be the result
of a poor propensity of infected mosquitoes to be active and fly. To test such hypothesis,
parasitized and unparasitized adults were collected 2 weeks after their emergence and each
individual was placed in a 25mm Ø activity tube containing a cotton soaked with a 10% sucrose
solution and sealed with cotton. The tube was placed in a LAM25 Locomotor Activity Monitor
system (Trikinetics). The system was crossed by 9 infrared beams that register an activity
signal whenever the mosquito crosses the beams. Mosquitoes were left for 15min to rest
before their activity was measured. The system was placed on a shelve that provides shadow
to the mosquitoes since they tend to prefer such habitats in nature (71). The total activity
measured with LAM systems can be mostly attributed to flight since walk distance do not
exceed 20% of mosquitoes activity (72). Two experiments each involving 16 parasitized males,
16 parasitized females, 16 unparasitized males and 16 unparasitized females were conducted
for 24h in a BSL2 insectary with 18h/6h day/night and 28°C. Whenever individuals died before
the end of the registration, they were removed from the final dataset.

Statistical and data analysis

Variations of As. taiwanensis prevalence in field populations. Statistical analyses
performed with the R software v.4.3.1 (73). The proportion of parasitized mosquitoes (defined
hereafter as parasite prevalence) has been modeled for field populations as a response
variable using a binomial distribution and a probit link function. The explanatory variables

tested were the sex and/or the mosquitoes collection site (defined hereafter as the mosquito
origin), using the full dataset or a subsampled dataset for each sex to study the impact of
factors without taking into account statistical interactions. The influence of each factor was
tested with a Type II Wald χ^2 test (*lme4* and *car* packages). We also correlated average
parasite prevalence in collection sites with the number of years after the mosquito introduction
as well as difference in prevalence between sites with the geographical distances separating
them using the Spearman rank index with a correlation test and the Moran's I autocorrelation
coefficient respectively (*stat* and *ape* packages). Geographical distances between mosquito
populations were calculated based on ellipsoid distance (*enmSdm,* package). Correlation
analyses were repeated with data corresponding to previously published datasets from
samples collected in the USA at different locations and time points after introduction.

Microbial communities associated with field populations. Microbiota diversity was
estimated using the number of OTUs per individual sample after rarefaction for richness,
Shannon index for alpha diversity and Bray-Curtis dissimilarity index for beta diversity (*vegan,*
package). Richness and alpha diversities were modeled using a Poisson distribution and a log
link function or a Gaussian distribution respectively and the impact of the sex and population
origin were used as fixed factors. The fixed factors were tested with a type II Anova or Wald χ^2
test (*lme4* and *car,* packages). The influence of sex and population origin on microbial Bray-
Curtis dissimilarity were tested using permutational multivariate analysis (*vegan,* package). A
similar analysis was used to compare the influence of the years after introduction and climates
for each sex on the microbiota. Moreover, a mantel correlation test was used to compare the
Beta diversity distances to geographical distances. Relative abundances of the 20 most
dominant prokaryotic and eukaryotic OTUs were represented as a heatmap
(*Complexheatmap,* package). The phylogenetic analysis of *As. taiwanensis* OTUs (based on
an alignment of the most abundant sequences in each OTU) was performed with Seaview and
the Muscle algorithm using other members of the Apicomplexa phylum. Phylogenetic trees
were then generated with a Maximum Likelihood method, an HKY evolutionary model and 100
bootstraps (performed with Seaview and represented with Figtree). To identify the
microorganisms which relative abundances covaried with the population age (years after
introduction of *Ae. albopictus* on each sampling sites), a Constrained Analysis of Principal
Coordinates (CAP) was performed (*vegan,* packages). The microbiota abundance correlation
networks were performed with a homemade *python* code and visualized with the *pyvis* library.
Briefly, the 200 most frequently retrieved OTUs for each community (prokaryotic and
eukaryotic) were selected and their paired non-parametric correlations were tested. If the
Spearman correlation (Rs) value was >0.25 (positive correlation) or <-0.25 (negative
correlation), a linked was added between the two pairs. An automatic positioning algorithm

was used based on mass-spring models. In this model, the links play a spring role which tend to bring the OTUs together while the OTUs play a role of opposite mass charges that are constantly repelled from each other. The algorithm used iterations to minimize the energy of the system. Individual *Ascogregarina taiwanensis* OTUs that vary according to mosquito population age were correlated with this last variable using non-linear Spearman correlations. Since relative abundances may induce bias due to their compositional nature, we have repeated CAP and correlation analyses with the selected samples for which we have performed qPCR.

Parasite prevalence, abundance and flight activity in controlled experiments. For the founder effect and desiccation experiments, the abundance of *As. taiwanensis* per mosquito were modeled with mixed models with a Gaussian and a Gamma inverse distribution respectively using mosquito sex, density or desiccation as fixed factors and the experimental replicates as random variables (*lme4*, package). Fixed factors were tested with an ANOVA or a type II Wald χ^2 test. Post-hoc Tukey-HSD tests were used to estimate pairwise differences (*emmeans*, package). Prevalence was also tested for the experiment in which enough parasitized and unparasitized mosquitoes were detected (at least 5 events of each). It was analyzed with mixed model using a binomial distribution with a logistic link function with the mosquito sex, desiccation as fixed factors and tested with a type II Wald χ^2 test. Flight activities (*i.e.*number of movements for each mosquitoes) was modeled with mixed model using a negative binomial, the sex of mosquitoes and infection status toward *As. taiwanensis* as fixed factors and the experimental replicates as random variables. The impact of the fixed factors was tested with a type II Wald χ^2 test. The datasets and scripts used for statistical analysis have been deposited on zenodo (https://zenodo.org/record/8252320).

Results

Recently introduced mosquitoes in mainland France harbor lower prevalence of As. taiwanensis

The prevalence of *As. taiwanensis* was estimated in 598 mosquitoes sampled among 17 populations using diagnostic PCR (**Table S1**; **Fig. S1**). Samples collected in mainland France were a combination of recently introduced populations and intermediate ones. For all other countries, native or intermediate populations were collected. Among the studied variables, the population origin significantly impacted the prevalence of the parasite and differences were observed between sexes along the sampling sites (**Table 1**). For those reasons, the two sexes were then analyzed separately. To explain features of the mosquito

origin that can influence the prevalence of the parasite, we further tested the impact of climate,

population age and geographical distance between mosquito populations. Variations in

climatic regions has no significant impact on parasite prevalence (**Table 1, Figure S2**).

However, it was positively correlated with the mosquito population age for both males and

females (**Table 1**, **Figure 1A, 1B**). In recent mosquito populations (< 2 years), the observed

prevalence of *As. taiwanensis* ranged from 0 to 25%, while it ranged from 37% to 100% in

populations that were native or introduced for more than 6 years. This effect was specifically

strong on French sites that are geographically closed from each other but harbor high

differences in parasite abundance correlating with differences in population ages. We could

therefore assume that spatial autocorrelation might influence this observed pattern. The

Moran's index revealed a very weak but significant spatial autocorrelation for Mosquitoes

populations at the global scale for both female and male mosquitoes (I = 0.27, p=0.038; I =

0.28, p = 0.013 for both sexes respectively). When considering the French populations

separately no significant autocorrelation was evidenced for females (I = 0.05, p=0.064) and a

very weak and yet significant autocorrelation was evidenced for males (I = 0.16, p = 0.02). To

separate the impact of population age from that of geographical distances, we included

prevalence data from previously published studies concerning recently introduced and older

populations from the USA (**Table S2**). Those data referred to prevalence of *As. taiwanensis* in

mosquito larvae and mixed adults and were included separately but in comparison with our

study (**Figure 1B**). The spatial autocorrelation was not significant in this new dataset (I = -0.52,

p = 0.22) while correlation between parasite prevalence and population age was comparable

with the one reported for French populations (R_s=0.46, *p*<0.001).

Ascogregarina taiwanensis *abundances correlate with the microbiota composition, the*

mosquito origin and the population age

To compare dynamics of *As. taiwanensis* within the microbiota of the different mosquito

populations, a metabarcoding analysis was conducted on a subset of 248 male and female

mosquitoes (**Table S1**). Mosquitoes were randomly selected for this analysis within the

previously described populations. The eukaryotic and prokaryotic microbiota of *Ae. albopictus*

were respectively dominated by two OTUs classified as *Wolbachia* (**Figure 2A**) and three

OTUs classified as *Ascogregarina* (**Figure 2A**). In total 8 different OTUs of *Ascogregarina* were

identified among the 200 most frequently retrieved OTUs but 5 showed low abundances.

Those three dominant *Ascogregarina* OTUs showed a prevalence of 74.4% and an average

abundance of 42.1±41.2% (mean±s.d.) among the eukaryotic microbiota while the two

dominant *Wolbachia* OTUs showed a prevalence of 93.4% and an average abundance of

39.4±34.4% among the prokaryotic microbiota. Other OTUs identified as *Aureobasidium* and
Penicillium genera for eukaryotes or as *Cutibacterium*, *Staphylococcus* and *Acinetobacter*
genera for prokaryotes were also prevalent. Study of factors involved in variation of microbiota
composition (*i.e.* β-diversity) between individuals revealed that the origin of mosquitoes was
the most important (R^2=0.32, *p*=0.001) structuring factor in interaction with the sex (R^2=0.04,
p=0.001) at a lower magnitude (see **Table S3**). Both eukaryotic and prokaryotic microbiota
dissimilarities were also correlated with the age of the population (**Figure S2**), climate (**Figure S3**)
S3) and geographical distance between individuals (**Figure S4**).

Based on their similarity, the eukaryotic community could be separated in three groups
(**Figure 2A**). The first group (60 individuals) was dominated by the most prevalent OTUs of
Ascogregarina (Otu000001), the second (43 individuals) by two different OTUs of
Ascogregarina (Otu000002 and Otu000003), and the last one (84 individuals) by neither of
those three OTUs. This pattern was confirmed with correlations networks that showed two
independent networks of positive correlations between relative abundances of OTUs identified
as *Ascogregarina* (**Figure 2B**). The first network involved the dominant Ascogregarina
(Otu000001) and fifth other underrepresented *Ascogregarina* OTUs (Otu000070, Otu000071,
Otu000085, Otu000092, Otu000096) while the second one involved solely the two others
dominant *Ascogregarina* OTUs (Otu000002 and Otu000003). The three dominant
Ascogregarina OTUs could be confidently classified within the *As. taiwanensis* species based
on a maximum likelihood phylogeny conducted on this same 18S rDNA region (**Figure 2C**).
The three most abundant *Ascogregarina taiwanensis* OTUs were explaining most part of the
eukaryotic microbiota that covaries with the age of the population based on a CAP analysis
(**Figure 2D**). Correlations of relative abundances of those three OTUs showed that Otu000002
and Otu000003 increased in abundance in old populations while Otu000001 significant vary
with the age of the population only for females with a low correlation index (**Figure 3A**). For
each of the three group of microbiota similarity, absolute abundances were calculated for a
subset of 7 females and 7 males. When converted in absolute abundances, the three most
abundant *Ascogregarina taiwanensis* OTUs were still the most discriminant ones covarying
with the age of the population in the CAP axis (**Figure 2E**). Otu000002 and Otu000003 showed
a significant positive correlation between their absolute abundance and the age of the
mosquitoes' population while this was not significant for Otu000001 (**Figure 3B**).

Parasitic success decreases over time when desiccated eggs are stored during human-
aided transportation

To evaluate whether egg desiccation may restrain the parasite infectivity through long
distance transportation, we collected eggs from five containers. Desiccated eggs were
separated into three groups (100 eggs/group) that we respectively stored 35h, 1 week and 1
month before being flooded (**Figure 4A**). Parasite quantifications in adults were performed on
10 individuals per time point and replicate. Results showed significant decreases of the
parasite absolute abundance, hereafter defined as the number of copies for specific genes of
As. taiwanensis, over time after desiccation for both males and females (**Figure 4B, Table 1**).
Based on 95% confidence interval of the model predictions, between 4421.4 and 86.6 copy of
oocysts per ng of DNA were quantified per mosquito at the beginning of the experiment while
this number was comprised between 134.3 and 0.2 after a month period of desiccation. Such
treatment also affected the prevalence of the parasite within mosquitoes (**Figure 4C, Table 1**).

Founder effects can reduce the amplitude of parasite infections due to low mosquito
population densities and an absence of Density-Dependent Prophylaxis (DDP)

Two experiments were designed to determine whether the density of adults releasing
oocysts and DDP (*i.e.* number of larvae receiving oocysts) could influence the parasite success
to colonize new adult mosquitoes (**Figure 5A**). When the number of adults releasing oocysts
in water was increased, we observed that higher densities of parasites (emmeans estimate ±
se: 207±45.7 and 191±49.8 for males and females respectively) colonized individuals
emerging from the water habitat (**Figure 5B, Table 1**). We also noticed that emerging females
harbored more parasite oocysts than males (emmeans estimate ± se: 100±33.5). However,
the parasite success was not significantly modified by larval densities either in males or in
females (**Figure 5C, Table 1**).

The parasite is not likely to impair mosquito dispersion through flight since parasitized
individuals are more active

The flight behavior of uninfected and infected mosquitoes was assessed using a
Locomotor Activity Monitor. We recorded individually the number of movements of parasitized
and unparasitized mosquitoes during 24h (**Figure 6A**). Contrarily to our expectations,
parasitized mosquitoes were significantly more active than unparasitized ones (emmeans
estimates: 1.34±0.45 and 0.89±0.45 for females and males respectively; **Figure 6B, Table 1**).
Although it was not statistically significant, the same pattern was observed for parasitized
males (49.4 moves) in comparison to unparasitized ones (18.9 moves). Therefore, those
results suggest that the low infection rate of mosquitoes in young populations is not related

with a higher propensity of uninfected individuals to disperse by flight but the parasite could,
conversely, promote the mosquito flight activity.

Discussion

In this study, we reported a positive correlation between *As. taiwanensis* prevalence
and *Ae. albopictus* population age for different mosquito populations. Such parasitism escape
has been evidenced in particular within recently introduced populations in France and USA.
The same pattern was observed regarding the abundance of 2 out of the 3 main *As.*
taiwanensis genotypes identified in the metabarcoding analysis. All in all, those convergent
tendencies suggest that *Ae. albopictus* might have escaped its natural parasite multiple time
along its invasion history. Such decrease might result from subsampling of non- or poorly-
infected individuals in the source population that formed the introduced ones. However, this is
more likely to occur if the parasite is rare in the source population (16,17) while our field study
suggest that the native or old populations sampled along potential source areas are highly
colonized by the parasite. Indeed, most recent mosquito records and population genetic
studies suggest that mosquito introduction in mainland France is the combination of south
European, north-east American introduced populations as well as old invasive populations
from the Indian Ocean and native populations from Asia (74,75). Based on PCR diagnostics
and microbial community analysis on populations from those areas, we can argue that the
parasite is likely to be prevalent and dominant in source population. Three dominant genotypes
of *As. taiwanensis* were observed in those area and two of them (Otu0002 and Otu0003)
correlated with the laps time separating the mosquito introduction to its sampling. Indeed,
metabarcoding and qPCR analysis suggest that those two OTUs dominate the microbiota of
intermediate and native populations but not that of recently introduced ones. A recent study on
parasite infection in invasive species has shown that parasite characteristics is reliable to
predict parasite retention (76). The authors even demonstrated that specialist parasites (like
As. taiwanensis is*)* are usually more persistent to co-invade with their native host. All in all, we
cannot exclude that stochasticity due to subsampling of non-infected individuals may have
played a role in the absence of infection in recently introduced populations but since it is
unlikely, we argue that other mechanisms have probably led them to escape *As. taiwanensis.*
504

The Asian tiger mosquito invasive process is mainly induced by human-mediated large
scale dispersal of desiccated eggs as previously discussed (77) and is therefore facilitated by
desiccated egg ability to resist without hatching during transportation (78,24). The desiccation
and storage laps time that we selected (*i.e.* up to 1month) were included within the range the

eggs can survive without showing consistent decrease in survival (40). We demonstrated that
the abundance and prevalence of parasite infection in mosquitoes hatched from desiccated
eggs decreases as a function of the time the eggs were kept before being flooded. Similarly,
the gregarine *Apolocystis elongata* showed a reduced infectivity toward its earthworm host
Eisenia foetida three weeks after oocysts have been released on a dry soil (79). Sensitivity to
desiccation has also been reported as a major oocyst weakness in related Apicomplexan
species such as *Cryptosporidium* or *Eimeria* species (80,81). The molecular mechanism
involved in desiccation sensitivity has poorly been studied until now in Apicomplexa but is likely
to be due to the peculiar evolution of those microorganisms that recently diverged from marine
algae (82). At the opposite, resistance of cysts to desiccation have been acquired in some
Apicomplexan taxa such as *Toxoplasma* spp. and was associated to the production of Late
Embryogenesis Abundant (LEA)-related proteins (83). The particular biochemical properties of
those protein enable them to harbor chaperone-like activities and contribute to protein
stabilization that may reinforce the cyst wall integrity and to limit water loss. The lack of
acquisition of such protein in *As. taiwanensis* still needs to be demonstrated but may explain
why this parasite that is often released in water container poorly resist to desiccation.

Since transportation of desiccated eggs is a major path driving the mosquito
introduction (29–32,74,84,85), human-aided transportation could be a consistent factor
explaining *Ae. albopictus* parasitism escape after mosquito population introduction. However,
part of the mosquitoes are also transported as adults for a shorter period of time in car cabin
and this path that can hardly be related to parasitism escape (29). Habitat and host
permissiveness to the parasite after introduction may also be a key factor that could have
decreased the infection probability for a short period of time after mosquito introduction (86).
Those processes are complex and may involve biotic and abiotic components. We did not fully
address this question and argue that further studies must be conducted to better understand
the importance of genotype x genotype x environment (GxGxE) interactions in *Ae. albopictus*
- *As. taiwanensis* interactions. However, our results provide new insights for further
investigations. Firstly, the metabarcoding of eukaryotic and prokaryotic communities in *Ae.*
albopictus did not evidence that the parasite was replaced by another competitive parasite
from the invasion range. This part suggest that the biotic part associated to microorganism
does not seem very relevant to explain the host-parasite dynamics. Secondly, the habitats in
mainland France and in USA could be partially refractory to the parasite since we evidenced
for both localizations' correlation between geographical distances and parasite prevalence.
Further studies on biotic and abiotics are necessary to conclude on the environment
importance on this host-parasite relationship.

In other host-parasite systems such as snail-trematode, the host density has been
shown to be a good predictor of parasite colonization success (87). Such phenomena are often
predicted to occur for horizontally transmitted parasites due to inter-individual contacts leading
to higher transmission probability that increase with population density (88). Since inter-
individual horizontal transmission of parasites is mediated through oocysts released from
adults in water habitats, one can expect that a reduced number of mosquitoes invading a new
area would result in a limited number of parasites in the breeding sites to colonize new hosts.
Previous studies on *As. taiwanensis* evidenced that the parasite oocyst load released in water
influence the success of host infection (59). In this study, we have complemented those results
by showing that the number of adult mosquitoes releasing parasites positively increases the
number of parasites efficiently colonizing unparasitized conspecifics emerging from water
habitats. Interestingly an inter-seasonal mosquito and *As. taiwanensis* dynamics have been
shown to be correlated over the year (51). Those results converge toward the importance of
high mosquito population densities for an efficient parasite transmission, independently from
the probability of parasite acquisition. Previous studies on gregarious insects showed that an
adaptive response to host high density leads to a reduction of the parasite susceptibility
through the enhancement of their immune system (22). This phenomenon is also named
Density-Dependent Prophylaxis (DDP). For instance, in response to density raise, the bumble
bee *Bombus terrestris* (Hymenoptera) and the velvet bean caterpillar *Anticarsia gemmatalis*,
increase their phenoloxidase activity (immune response leading to a melanization process)
(89,90). In this study, we have shown an absence of larval density-mediated modulation of the
parasite infection cycle. Although the immune response has not been tested, we can then
predict that the parasite transmission drastically increases with the population size without
being slowed down by DDP.

We have highlighted the impact of the different invasion processes in the host-parasite
relationship. Contrarily to our hypothesis, local mosquito dispersal through flight has been
shown to be improved by parasitism. Local flight distance of field populations is generally
limited to a few hundred meters (47) and remains a secondary factor to explain *Ae. albopictus*
invasiveness. The hypothesis of gregarine dispersal through local flight is debatable due to
opposite examples. Indeed, the infectivity of the gregarine *Ophryocystis elektroscirrha* to
monarch butterfly (*Danaus plexippus*) has been proved to reduce host wing mass and
resistance (91). In the dragonfly (*Libellula pulchella*), gregarines are also negatively affecting
the flight performance by decreasing the concentration of muscle myosin and impairing muscle
contractile performance (92,93). The results reported here are preliminary. Indeed, laboratory
experiments on mosquitoes behavior using locomotor activity monitor do not take into account
various stimuli (e.g. breeding sites, hosts…) from the environment that might differentially

enhance the local dispersal of mosquitoes (94,95). Other factors such as the mosquito age
and nutrition would also be considered in the future while previous studies showed that the
activity varies according to those factors (96). If our result is confirmed under more complex
conditions, this would suggest that mosquitoes that have been colonized by the parasite after
introduction might be more prompt to locally disperse.

Recently, a mathematical model predicted *Ae. albopictus* competitiveness against the
tree hole mosquito *Ae. triseriatus* in presence of gregarine parasites (97). In most scenarii, *Ae.*
albopictus dominates *Ae. triseriatus*, but *As. taiwanensis* infection may decrease the
competitiveness of its host by increasing the larval development time and inducing a higher
mortality, according to other studies (98). The authors concluded that parasitism reduces *Ae.*
albopictus competitiveness. Selective pressure induced by competition might explain the loss
of parasites in recently introduced *Ae. albopictus* populations since its presence might restrain
the population settlement probability. Another hypothesis to explain the parasitism escape is a
dilution effect when *Ae. albopictus* colonizes new areas. Both parasite prevalence and intensity
could be reduced due to a wider range of potential host (induced by native mosquito species).
Indeed, they can colonize first instar larvae of other mosquito species but rarely achieve to
complete their life cycle and replicate inside them (49). To the best of our knowledge, no study
evidenced dilution of *As. taiwanensis* out of *Ae. albopictus*. However, dilution effect of
Ascogregarina barreti has been experimentally observed during co-inoculation of its native
host *Ae. triseriatus* with other mosquito species such as *Ae. japonicus* (99) and *Ae. albopictus*
(100). It was also suggested by field surveys reporting that the prevalence of *As. taiwanensis*
colonizing *Ae. albopictus* was higher when the mosquitoes did not share breeding sites with
Ae. aegypti (101). *Ae albopictus* is often reported as a strong competitor toward other
mosquitoes (102–107). Therefore, the parasite prevalence dynamics in young populations
could be due to an evolution of the dilution effect resulting from a higher diversity of mosquitoes
in breeding sites at the early stage of the colonization process that decrease over the time due
to interspecific competition. Field studies on the arthropod's community sharing breeding sites
with recent and ancient populations may help to find relevant alternative hosts to test this
hypothesis.

Conclusion

In this study, we provide new insights into the dynamics of *Ae. albopictus-As.*
taiwanensis relationship by combining field observations and laboratory experiments. In
French populations, the mosquito host has escaped its parasite after introduction and this
process is unlikely to be stochastic. The maintenance of non-flooded eggs in containers during

human-aided transportation and the decrease in mosquito founding population after introduction have likely altered the parasite maintenance and transmission. Local dispersal through flight is probably not affected by the parasite since the flight activity of unparasitized individuals was lower than that of their parasitized conspecifics. We need further investigations on the consequences of the rupture of this natural interaction to understand how it may affect the mosquito ecology, population dynamics and pathogen transmission.

Acknowledgement. Part of the mosquito collection was enabled by the international network of the 'Mosquito Microbiome Consortium' (https://mosquito-microbiome.org/). This project was funded by the French National Research Program for Environmental and Occupational Health of Anses, grant number 19-95. We would also like to thank the EC2CO program for its support to this project (project name 'INTERASCO') as well as the IDEX Breakthrough (project name 'MICRO-BEHAVE'). Sampling collections were also supported by the Thailand Program Management Unit for Human Resources & Institutional Development, Research and Innovation (PMU-B), NXPO, grant number B17F640002. This study used resources from the iBio platform and the FR BioEEnViS. Finally, the authors would like to thank James Swanson and Stephanie Jiang from Emory University for their help during sampling.

Data availability. All the reads, datasets and scripts used for this analysis have been deposited on zenodo (https://zenodo.org/record/8252320).

References

1. Atwoli L, Baqui AH, Benfield T, Bosurgi R, Godlee F, Hancocks S, et al. Call for Emergency Action to Limit Global Temperature Increases, Restore Biodiversity, and Protect Health: Wealthy Nations Must do Much More, Much Faster. Glob Heart. 2021;16(1):60.

2. Bellard C, Genovesi P, Jeschke JM. Global patterns in threats to vertebrates by biological invasions. Proceedings of the Royal Society B: Biological Sciences. 27 janv 2016;283(1823):20152454.

3. Bradshaw CJA, Leroy B, Bellard C, Roiz D, Albert C, Fournier A, et al. Massive yet grossly underestimated global costs of invasive insects. Nature Communications. 4 oct 2016;7:12986.

4. Didham RK, Tylianakis JM, Hutchison MA, Ewers RM, Gemmell NJ. Are invasive species the drivers of ecological change? Trends in Ecology & Evolution. 1 sept 2005;20(9):470-4.

5. Doherty TS, Glen AS, Nimmo DG, Ritchie EG, Dickman CR. Invasive predators and global biodiversity loss. Proc Natl Acad Sci U S A. 4 oct 2016;113(40):11261-5.

6. Mainka SA, Howard GW. Climate change and invasive species: double jeopardy. Integrative Zoology. 2010;5(2):102-11.

654   7.   Hulme PE. Unwelcome exchange: International trade as a direct and indirect driver of biological
655   invasions worldwide. One Earth. 21 mai 2021;4(5):666-79.

656   8.   Andersen MC, Adams H, Hope B, Powell M. Risk Assessment for Invasive Species. Risk Analysis.
657   2004;24(4):787-93.

658   9.   Diagne C, Leroy B, Vaissière AC, Gozlan RE, Roiz D, Jarić I, et al. High and rising economic costs of
659   biological invasions worldwide. Nature. avr 2021;592(7855):571-6.

660   10.   Hulme PE. Invasive species challenge the global response to emerging diseases. Trends in
661   Parasitology. 1 juin 2014;30(6):267-70.

662   11.   Shackleton RT, Shackleton CM, Kull CA. The role of invasive alien species in shaping local
663   livelihoods and human well-being: A review. J Environ Manage. 1 janv 2019;229:145-57.

664   12.   Chalkowski K, Lepczyk CA, Zohdy S. Parasite Ecology of Invasive Species: Conceptual Framework
665   and New Hypotheses. Trends Parasitol. août 2018;34(8):655-63.

666   13.   Roberts M, Dobson A, Restif O, Wells K. Challenges in modelling the dynamics of infectious
667   diseases at the wildlife–human interface. Epidemics. 1 déc 2021;37:100523.

668   14.   Dunn AM. Chapter 7 Parasites and Biological Invasions. In: Advances in Parasitology [Internet].
669   Academic Press; 2009 [cité 26 juin 2023]. p. 161-84. (Natural History of Host-Parasite
670   Interactions; vol. 68).

671   15.   Torchin ME, Lafferty KD, Kuris AM. Parasites and marine invasions. Parasitology. oct
672   2002;124(7):137-51.

673   16.   Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ. Is invasion success explained by the enemy
674   release hypothesis? Ecology Letters. 2004;7(8):721-33.

675   17.   Drake JM. The paradox of the parasites: implications for biological invasion. Proceedings of the
676   Royal Society of London Series B: Biological Sciences. 7 nov 2003;270(suppl_2):S133-5.

677   18.   Blakeslee AMH, Fowler AE, Keogh CL. Chapter Two - Marine Invasions and Parasite Escape:
678   Updates and New Perspectives. In: Lesser M, éditeur. Advances in Marine Biology [Internet].
679   Academic Press; 2013 [cité 26 juin 2023]. p. 87-169.

680   19.   Llopis-Belenguer C, Blasco-Costa I, Balbuena JA, Sarabeev V, Stouffer DB. Native and invasive
681   hosts play different roles in host–parasite networks. Ecography. 2020;43(4):559-68.

682   20.   Llaberia-Robledillo M, Balbuena JA, Sarabeev V, Llopis-Belenguer C. Changes in native and
683   introduced host–parasite networks. Biol Invasions. 1 févr 2022;24(2):543-55.

684   21.   Nørgaard LS, Zilio G, Saade C, Gougat-Barbera C, Hall MD, Fronhofer EA, et al. An evolutionary
685   trade-off between parasite virulence and dispersal at experimental invasion fronts. Ecol Lett. avr
686   2021;24(4):739-50.

687   22.   Wilson K, Cotter S (2008) Density-dependent prophylaxis in insects. In: Ananthakrishnan TN,
688   Whitman TW (eds) Phenotypic plasticity of insects: mechanisms and consequences, vol 1.
689   Science Publishers, Enfeld, UK. 381–420.

23. Bradley CA, Altizer S. Parasites hinder monarch butterfly flight: implications for disease spread in
migratory hosts. Ecology Letters. 2005;8(3):290-300.

24. Bonizzoni M, Gasperi G, Chen X, James AA. The invasive mosquito species Aedes albopictus:
current knowledge and future perspectives. Trends Parasitol. sept 2013;29(9):460-8.

25. Lequime S, Dehecq JS, Matheus S, de Laval F, Almeras L, Briolant S, et al. Modeling intra-
mosquito dynamics of Zika virus and its dose-dependence confirms the low epidemic potential of
Aedes albopictus. PLoS Pathog. 31 déc 2020;16(12):e1009068.

26. Paupy C, Delatte H, Bagny L, Corbel V, Fontenille D. Aedes albopictus, an arbovirus vector: From
the darkness to the light. Microbes and Infection. 1 déc 2009;11(14):1177-85.

27. Goubert C, Minard G, Vieira C, Boulesteix M. Population genetics of the Asian tiger mosquito
Aedes albopictus, an invasive vector of human diseases. Heredity (Edinb). sept
2016;117(3):125-34.

28. Bennett K, Gómez Martínez C, Almanza A, Rovira JR, McMillan WO, Enriquez V, et al. High
infestation of invasive Aedes mosquitoes in used tires along the local transport network of
Panama. Parasites Vectors. déc 2019;12(1):1-10.

29. Eritja R, Palmer JRB, Roiz D, Sanpera-Calbet I, Bartumeus F. Direct Evidence of Adult Aedes
albopictus Dispersal by Car. Sci Rep. 24 oct 2017;7(1):14399.

30. Madon MB, Mulla MS, Shaw MW, Kluh S, Hazelrigg JE. Introduction of Aedes albopictus (Skuse)
in southern California and potential for its establishment. J Vector Ecol. juin 2002;27(1):149-54.

31. Medley KA, Jenkins DG, Hoffman EA. Human-aided and natural dispersal drive gene flow across
the range of an invasive mosquito. Mol Ecol. 1 janv 2015;24(2):284-95.

32. Miller MJ, Loaiza JR. Geographic Expansion of the Invasive Mosquito Aedes albopictus across
Panama—Implications for Control of Dengue and Chikungunya Viruses. PLOS Neglected Tropical
Diseases. 8 janv 2015;9(1):e0003383.

33. Schmidt TL, Swan T, Chung J, Karl S, Demok S, Yang Q, et al. Spatial population genomics of a
recent mosquito invasion. Molecular Ecology. 2021;30(5):1174-89.

34. Cornel AJ, Hunt RH. Aedes albopictus in Africa? First records of live specimens in imported tires
in Cape Town. Journal of the American Mosquito Control Association. 1991;7(1):107-8.

35. Dalla Pozza G, Majori G. First record of Aedes albopictus establishment in Italy. J Am Mosq
Control Assoc. sept 1992;8(3):318-20.

36. Laird M, Calder L, Thornton RC, Syme R, Holder PW, Mogi M. Japanese Aedes albopictus among
four mosquito species reaching New Zealand in used tires. J Am Mosq Control Assoc. mars
1994;10(1):14-23.

37. Moore CG, Mitchell CJ. Aedes albopictus in the United States: ten-year presence and public
health implications. Emerging Infect Dis. sept 1997;3(3):329-34.

38. Pluskota B, Storch V, Braunbeck T, Beck M, Becker N. First record of Stegomyia albopicta (Skuse)
(Diptera: Culicidae) in Germany. European Mosquito Bulletin. 2008;26:1-5.

39. Schaffner F, Van Bortel W, Coosemans M. First record of Aedes (Stegomyia) albopictus in
Belgium. J Am Mosq Control Assoc. juin 2004;20(2):201-3.

40. Juliano SA, O'Meara GF, Morrill JR, Cutwa MM. Desiccation and thermal tolerance of eggs and
the coexistence of competing mosquitoes. Oecologia. 1 févr 2002;130(3):458-69.

41. Toma L, Severini F, Di Luca M, Bella A, Romi R. Seasonal patterns of oviposition and egg hatching
rate of Aedes albopictus in Rome. J Am Mosq Control Assoc. mars 2003;19(1):19-22.

42. Sherpa S, Renaud J, Guéguen M, Besnard G, Mouyon L, Rey D, et al. Landscape does matter:
Disentangling founder effects from natural and human-aided post-introduction dispersal during
an ongoing biological invasion. Journal of Animal Ecology. 2020;89(9):2027-42.

43. Kambhampati S, Black WC 4th, Rai KS. Geographic origin of the US and Brazilian Aedes albopictus
inferred from allozyme analysis. Heredity (Edinb). août 1991;67 (Pt 1):85-93.

44. Minard G, Tran FH, Van VT, Goubert C, Bellet C, Lambert G, et al. French invasive Asian tiger
mosquito populations harbor reduced bacterial microbiota and genetic diversity compared to
Vietnamese autochthonous relatives. Front Microbiol [Internet]. 22 sept 2015.

45. Gojković N, Ludoški J, Krtinić B, Milankov V. The First Molecular and Phenotypic Characterization
of the Invasive Population of Aedes albopictus (Diptera: Culicidae) from the Central Balkans. J
Med Entomol. 3 sept 2019;56(5):1433-40.

46. Schmidt TL, Chung J, Honnen AC, Weeks AR, Hoffmann AA. Population genomics of two invasive
mosquitoes (Aedes aegypti and Aedes albopictus) from the Indo-Pacific. PLoS Negl Trop Dis. juill
2020;14(7):e0008463.

47. Vavassori L, Saddler A, Müller P. Active dispersal of Aedes albopictus: a mark-release-recapture
study using self-marking units. Parasites Vectors. déc 2019;12(1):1-14.

48. Thongsripong P, Chandler JA, Green AB, Kittayapong P, Wilcox BA, Kapan DD, et al. Mosquito
vector-associated microbiota: Metabarcoding bacteria and eukaryotic symbionts across habitat
types in Thailand endemic for dengue and other arthropod-borne diseases. Ecol Evol.
2018;8(2):1352-68.

49. Lantova L, Volf P. Mosquito and sand fly gregarines of the genus Ascogregarina and Psychodiella
(Apicomplexa: Eugregarinorida, Aseptatorina) - Overview of their taxonomy, life cycle, host
specificity and pathogenicity. Infect Genet Evol. 4 mai 2014;

50. Erthal JA, Soghigian JS, Livdahl T. Life Cycle Completion of Parasite Ascogregarina taiwanensis
(Apicomplexa: Lecudinidae) in Non-Native Host Ochlerotatus japonicus (Diptera: Culicidae).
Journal of Medical Entomology. 1 sept 2012;49(5):1109-17.

51. Garcia JJ, Fukuda T, Becnel JJ. Seasonality, prevalence and pathogenicity of the gregarine
Ascogregarina taiwanensis (Apicomplexa: Lecudinidae) in mosquitoes from Florida. J Am Mosq
Control Assoc. sept 1994;10(3):413-8.

52. Levine ND. Progress in taxonomy of the Apicomplexan protozoa. J Protozool. nov
1988;35(4):518-20.

53. Munstermann LE, Wesson DM. First record of Ascogregarina taiwanensis (Apicomplexa:
Lecudinidae) in North American Aedes albopictus. J Am Mosq Control Assoc. juin
1990;6(2):235-43.

54. Reeves WK, McCullough SD. Laboratory susceptibility of Wyeomyia smithii (Diptera: Culicidae) to
Ascogregarina taiwanensis (Apicomplexa: Lecudinidae). J Eukaryot Microbiol. oct
2002;49(5):391-2.

55. Tseng M. Ascogregarine parasites as possible biocontrol agents of mosquitoes. J Am Mosq
Control Assoc. 2007;23(2 Suppl):30-4.

56. Boisard J, Florent I. Why the –omic future of Apicomplexa should include gregarines. Biology of
the Cell. 2020;112(6):173-85.

57. Aliabadi BW, Juliano SA. Escape from gregarine parasites affects the competitive interactions of
an invasive mosquito. Biol Invasions. 1 sept 2002;4(3):283-97.

58. Comiskey NM, Lowrie RC, Wesson DM. Effect of nutrient levels and Ascogregarina taiwanensis
(Apicomplexa: Lecudinidae) infections on the vector competence of Aedes albopictus (Diptera:
Culicidae) for Dirofilaria immitis (Filarioidea: Onchocercidae). J Med Entomol. janv
1999;36(1):55-61.

59. Soghigian J, Livdahl T. Differential response to mosquito host sex and parasite dosage suggest
mixed dispersal strategies in the parasite Ascogregarina taiwanensis. PLOS ONE. 13 sept
2017;12(9):e0184573.

60. Blackmore MS, Scoles GA, Craig GB. Parasitism of Aedes aegypti and Ae. albopictus (Diptera:
Culicidae) by Ascogregarina spp. (Apicomplexa: Lecudinidae) in Florida. J Med Entomol. nov
1995;32(6):847-52.

61. Rueda LM. Pictorial keys for the identification of mosquitoes (Diptera: Culicidae) associated with
dengue virus transmission. 2004;(589):1-60.

62. Knudsen AB, Romi R, Majori G. Occurrence and spread in Italy of Aedes albopictus, with
implications for its introduction into other parts of Europe. J Am Mosq Control Assoc. juin
1996;12(2 Pt 1):177-83.

63. Collantes F, Delacour S, Alarcón-Elbal PM, Ruiz-Arrondo I, Delgado JA, Torrell-Sorio A, et al.
Review of ten-years presence of Aedes albopictus in Spain 2004–2014: known distribution and
public health concerns. Parasites Vectors. déc 2015;8(1):1-11.

64. Womack ML, Thuma TS, Evans BR. Distribution of Aedes albopictus in Georgia, USA. J Am Mosq
Control Assoc. juin 1995;11(2 Pt 1):237.

65. Edwards FW. Notes on the Mosquitos of Madagascar, Mauritius and Réunion. Bulletin of
Entomological Research. août 1920;11(2):133-8.

66. Comiskey NM, Lowrie RC Jr, Wesson DM. Role of Habitat Components on the Dynamics of Aedes
albopictus (Diptera: Culicidae) from New Orleans. Journal of Medical Entomology. 1 mai
1999;36(3):313-20.

67. Seabourn P, Spafford H, Yoneishi N, Medeiros M. The Aedes albopictus (Diptera: Culicidae) microbiome varies spatially and with Ascogregarine infection. PLoS Negl Trop Dis. août 2020;14(8):e0008615.

68. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nat Methods. juill 2012;9(7):671-5.

69. Schloss PD. Secondary structure improves OTU assignments of 16S rRNA gene sequences. ISME J. mars 2013;7(3):457-60.

70. Clements AN. The Biology of Mosquitoes: Sensory reception and behaviour. Chapman & Hall; 1999. 764 p.

71. Flacio E, Engeler L, Tonolla M, Lüthy P, Patocchi N. Strategies of a thirteen year surveillance programme on Aedes albopictus (Stegomyia albopicta) in southern Switzerland. Parasites Vectors. déc 2015;8(1):1-18.

72. Ziegler R, Blanckenhorn WU, Mathis A, Verhulst NO. Video analysis of the locomotory behaviour of Aedes aegypti and Ae. japonicus mosquitoes under different temperature regimes in a laboratory setting. Journal of Thermal Biology. 1 avr 2022;105:103205.

73. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. 2016.

74. Vega-Rúa A, Marconcini M, Madec Y, Manni M, Carraretto D, Gomulski LM, et al. Vector competence of Aedes albopictus populations for chikungunya virus is shaped by their demographic history. Commun Biol. 24 juin 2020;3(1):1-13.

75. Sherpa S, Blum MGB, Capblancq T, Cumer T, Rioux D, Després L. Unraveling the invasion history of the Asian tiger mosquito in Europe. Mol Ecol. 8 mars 2019;

76. Schatz AM, Park AW. Patterns of host–parasite coinvasion promote enemy release and specialist parasite spillover. Journal of Animal Ecology. 2023;92(5):1029-41.

77. Roche B, Léger L, L'Ambert G, Lacour G, Foussadier R, Besnard G, et al. The Spread of Aedes albopictus in Metropolitan France: Contribution of Environmental Drivers and Human Activities and Predictions for a Near Future. PLOS ONE. 11 mai 2015;10(5):e0125600.

78. Juliano SA, Lounibos LP. Ecology of invasive mosquitoes: effects on resident species and on human health. Ecol Lett. mai 2005;8(5):558-74.

79. Miles HB. The Mode of Transmission of the Acephaline Gregarine Parasites of Earthworms. The Journal of Protozoology. 1962;9(3):303-6.

80. King BJ, Monis PT. Critical processes affecting Cryptosporidium oocyst survival in the environment. Parasitology. mars 2007;134(3):309-23.

81. Jenkins MC, Parker C, O'Brien C, Miska K, Fetterer R. Differing Susceptibilities of Eimeria acervulina, Eimeria maxima, and Eimeria tenella Oocysts to Desiccation. Journal of Parasitology. 1 oct 2013;99(5):899-902.

82. Janouškovec J, Paskerova GG, Miroliubova TS, Mikhailov KV, Birley T, Aleoshin VV, et al. Apicomplexan-like parasites are polyphyletic and widely but selectively dependent on cryptic

plastid organelles. McCutcheon J, Weigel D, Howe C, McFadden G, éditeurs. eLife. 16 août
2019;8:e49662.

83. Arranz-Solís D, Warschkau D, Fabian BT, Seeber F, Saeij JPJ. Late Embryogenesis Abundant
Proteins Contribute to the Resistance of Toxoplasma gondii Oocysts against Environmental
Stresses. mBio. 21 févr 2023;14(2):e02868-22.

84. Bennett K, Gómez-Martínez C, Almanza A, Rovira J, McMillan W, Enriquez V, et al. High
infestation of invasive Aedes mosquitoes in used tires along the local transport network of
Panama. Parasites & Vectors. 27 mai 2019;12.

85. Schmidt TL, Rašić G, Zhang D, Zheng X, Xi Z, Hoffmann AA. Genome-wide SNPs reveal the drivers
of gene flow in an urban population of the Asian Tiger Mosquito, Aedes albopictus. PLOS
Neglected Tropical Diseases. 18 oct 2017;11(10):e0006009.

86. Dunn AM. Parasites and biological invasions. Adv Parasitol. 2009;68:161-84.

87. Buck JC, Hechinger RF, Wood AC, Stewart TE, Kuris AM, Lafferty KD. Host density increases
parasite recruitment but decreases host risk in a snail–trematode system. Ecology.
2017;98(8):2029-38.

88. Watve MG, Sukumar R. Parasite abundance and diversity in mammals: correlates with host
ecology. Proc Natl Acad Sci USA. 12 sept 1995;92(19):8945-9.

89. Ruiz-González MX, Moret Y, Brown MJF. Rapid induction of immune density-dependent
prophylaxis in adult social insects. Biology Letters. 23 déc 2009;5(6):781-3.

90. Silva FWS, Serrão JE, Elliot SL. Density-dependent prophylaxis in primary anti-parasite barriers in
the velvetbean caterpillar. Ecological Entomology. 2016;41(4):451-8.

91. Davis AK, Roode JC de. Effects of the parasite, Ophryocystis elektroscirrha, on wing
characteristics important for migration in the monarch butterfly. Animal Migration. 1 déc
2018;5(1):84-93.

92. Marden JH, Cobb JR. Territorial and mating success of dragonflies that vary in muscle power
output and presence of gregarine gut parasites. Animal Behaviour. oct 2004;68(4):857-65.

93. Schilder RJ, Marden JH. Parasites, proteomics and performance: effects of gregarine gut
parasites on dragonfly flight muscle composition and function. Journal of Experimental Biology.
15 déc 2007;210(24):4298-306.

94. Girard M, Martin E, Vallon L, Raquin V, Bellet C, Rozier Y, et al. Microorganisms Associated with
Mosquito Oviposition Sites: Implications for Habitat Selection and Insect Life Histories.
Microorganisms. août 2021;9(8):1589.

95. Wooding M, Naudé Y, Rohwer E, Bouwer M. Controlling mosquitoes with semiochemicals: a
review. Parasites & Vectors. 17 févr 2020;13(1):80.

96. Kaufmann C, Collins LF, Brown MR. Influence of Age and Nutritional Status on Flight Performance
of the Asian Tiger Mosquito Aedes albopictus (Diptera: Culicidae). Insects. 26 juill
2013;4(3):404-12.

97. Stump E, Childs LM, Walker M. Parasitism of Aedes albopictus by Ascogregarina taiwanensis
lowers its competitive ability against Aedes triseriatus. Parasites & Vectors. 25 janv
2021;14(1):79.

98. Comiskey NM, Lowrie RC, Wesson DM. Role of habitat components on the dynamics of Aedes
albopictus (Diptera: Culicidae) from New Orleans. J Med Entomol. mai 1999;36(3):313-20.

99. Westby KM, Sweetman BM, Van Horn TR, Biro EG, Medley KA. Invasive species reduces parasite
prevalence and neutralizes negative environmental effects on parasitism in a native mosquito.
Journal of Animal Ecology. 2019;88(8):1215-25.

100.   McIntire KM, Chappell KM, Juliano SA. How do noncompetent hosts cause dilution of
parasitism? Testing hypotheses for native and invasive mosquitoes. Ecology. oct
2021;102(10):e03452.

101.   Dos Passos RA, Tadei WP. Parasitism of Ascogregarina taiwanensis and Ascogregarina culicis
(Apicomplexa: Lecudinidae) in larvae of Aedes albopictus and Aedes aegypti (Diptera: Culicidae)
from Manaus, Amazon region, Brazil. J Invertebr Pathol. mars 2008;97(3):230-6.

102.   Bagny Beilhe L, Delatte H, Juliano SA, Fontenille D, Quilici S. Ecological interactions in Aedes
species on Reunion Island. Medical and Veterinary Entomology. 2013;27(4):387-97.

103.   Costanzo KS, Mormann K, Juliano SA. Asymmetrical Competition and Patterns of Abundance
of Aedes albopictus and Culex pipiens (Diptera: Culicidae). Journal of Medical Entomology. 1 juill
2005;42(4):559-70.

104.   de Oliveira S, Villela DAM, Dias FBS, Moreira LA, Maciel de Freitas R. How does competition
among wild type mosquitoes influence the performance of Aedes aegypti and dissemination of
Wolbachia pipientis? PLoS Negl Trop Dis. 9 oct 2017;11(10):e0005947.

105.   Juliano SA. Species Introduction and Replacement Among Mosquitoes: Interspecific Resource
Competition or Apparent Competition? Ecology. 1998;79(1):255-68.

106.   Kesavaraju B, Leisnham PT, Keane S, Delisi N, Pozatti R. Interspecific Competition between
Aedes albopictus and A. sierrensis: Potential for Competitive Displacement in the Western
United States. PLoS ONE. 28 févr 2014;9(2):e89698.

107.   O'Donnell DL, Armbruster P. Comparison of Larval Foraging Behavior of Aedes albopictus and
Aedes japonicus (Diptera: Culicidae). Journal of Medical Entomology. 1 nov 2007;44(6):984-9.

Competing Interests. The authors declare no competing interests.

Figure legends

Fig. 1. Correlation between the prevalence of *As. taiwanensis* and the time after
introduction of the mosquito populations in each sampling site. (A) Correlation between
parasite prevalence (*i.e.* proportion of infected individuals) and the population ages (number
of years separating the mosquito introduction from the collection date) for females (triangle)
and males (square) have been investigated in different countries (colors). (B) A comparative

analysis has been realized between previously published data and the data from this study regarding the correlation between parasite prevalence and population ages. The data from previously published studies includes larvae (circles) and mixed adults (diamond) and are colored in red meanwhile the data from this study are presented in grey. Each dot represents a sampling site. Native populations were arbitrarily set at 200 years to be ranked at the end of the x-axis.

Fig. 2. Heatmap of the microbiota associated with *Ae. albopictus*. (A) The 20 most abundant prokaryotic and eukaryotic OTUs were represented in columns. The number attached to *Wolbachia* and *Ascogregarina* OTUs corresponds to the OTUs number in the following analysis. The 'Abundance' color scale shows the OTU proportion in the community (pink). Rows are classified depending on the central scale that represent three different groups based on eukaryotic community dissimilarity (Bray-Curtis distance) and that correlates with variations in the three *As. taiwanensis* OTUs proportions. Individuals belonging to the G1 group are dominated by *Ascogregarina* Otu000001, those belonging to the G2 group are dominated by *Ascogregarina* Otu000002 and *Ascogregarina* Otu000003 and those belonging to the G3 group are not dominated by *Ascograrina* OTUs. Three additional scales show colors corresponding to the individuals' origin site, age (years after introduction), and sex. (B) A network analysis showing positive correlations between microbial OTUs (eukaryotes are represented in yellow while prokaryotes are in blue and each link represents a correlation with ρ>0.25). The left network in the zoom is dominated by the *As taiwanensis* Otu000001 and is correlated with minor OTUs assigned to *Ascogregarina*. The right network in the zoom shows correlations between the *As. taiwanensis* Otu000002 and Otu000003. (C) Phylogenetic analysis revealed the belonging of OTU identified as Gregarinasina to the species *Ascogregarina taiwanensis*. (D) The main eukaryotic OTUs according to the coordinate of the axis (CAP1) extracted from a constrained analysis of principal coordinates (Capscale) on the OTU relative abundance highlighting correlations between OTUs and the age of the population. The three *As. taiwanensis* main OTUs are circled in violet. (E) The main eukaryotic OTUs according to the coordinate of the axis (CAP1) extracted from a constrained analysis of principal coordinates (Capscale) on the OTU absolute abundance highlighting correlations between OTUs and the age of the population. The three *As. taiwanensis* main OTUs are circled in violet.

Fig. 3. Correlations between *Ascogregarina* OTUs abundance and the mosquito population age. (A) The age of the mosquito populations and the relative abundance of the three dominant *Ascogregarina* are represented for female and male mosquitoes. (B) The age of the mosquito populations and the absolute abundance of the three dominant *Ascogregarina*

are represented for female and male mosquitoes. The p-value represent the significance of the spearman correlation between OTUs absolute abundances and the age of the population (number of years between introduction and sampling). The native population was set to 200 years after introduction to be ranked at the end of the x-axis. The points are colored based on the mosquito population country.

Fig. 4. Impact of egg maintenance outside of water (desiccated) on the infectivity of *As. taiwanensis*. (A) Desiccated eggs have been stored for 35h, 1 week or 1 month in a container before being flooded to trigger egg hatching. Five containers were tested for each time point. The parasite abundance was estimated on emerging adults by qPCR. (B) The impact of desiccation time on parasite abundance was represented for each repeat (R1 to R5) and both sexes. (C) The impact of desiccation has also been tested on the parasite prevalence and was represented for each repeat (R1 to R5) and both sexes. The p-value and associated statistics have been obtained through spearman correlation analysis.

Fig. 5. Density of adults releasing oocysts positively impacts the gregarine colonization and it is not counter balanced by density dependent prophylaxis. (A)The impact of adult and larval mosquito density on the parasite success has been tested through two different experiments. In a first experiment, the density of adults releasing oocysts in water habitats was controlled. They were allowed to released oocysts through feces and body of dead individuals for a week before rearing larvae. In a second experiment, we tested density dependent prophylaxis. To that end, larvae were infected with the same water for 2 days before being separated in batches of 20, 50 and 100 individuals in 100ml of water. Each experiment was repeated five times. The parasite abundance was estimated on emerging adults by qPCR. (B) The impact of larval density (*i.e.* number of larvae per 100ml of water) and (C) the impact of adults density that released oocysts in the larval water habitat on the parasite abundance was represented for each repeat (R1 to R5) and both sexes. Each dot represents a single mosquito. The p-value obtained from a Wald χ^2 test conducted on a GLMM model represents the significance of the slope.

Fig. 6. *Ascogregarina taiwanensis* infection impacts on mosquito flight activity. (A) Parasitized or unparasitized mosquitoes activity was measured in a Locomotor Activity Monitor. Movements were recorded for 24h by measuring the number of times each individual pass through an infrared beam. The experiment was repeated on two generations. (B) The impact of parasitism on flight activity (*i.e.* number of time they passed in front of an infrared

987   beam during 24h) was measured for two generations (R1 and R2) and both sexes. The p-

988   value obtained from a Wald χ^2 test conducted on a GLMM model represents the significance

989   of the slope.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024 **Table 1. Summary of statistical analyses**

Category	Variable	Factor	Test	Details on sample	Subsample 2	χ²	*p-value*	Index
Field	Prevalence	Origin	GLM			280.65	**<0.0001*****	
		Sex				0.46	0.496	
		Origin x Sex				25.04	**0.008****	
		Origin			Female	198.09	**<0.0001*****	
					Male	107.61	**<0.0001*****	
		Climate				0	1	
		Climate x Sex				0	1	
		Age of the population	Spearman		Female		**<0.0001*****	0.50
					Male		**<0.0001*****	0.36
		Geographical distance	Moran		Female		**0.038***	0.27
					Male		**0.013***	0.28
					France x Female		0.064	0.05
					France x Male		**0.020***	0.16
					USA[1] x Mixed		0.688	-0.38
Experiment		Egg desiccation	GLMM ╪	Desiccation resistance	Female	20.124	**<0.0001*****	
					Male	17.456	**<0.0001*****	
		Adult mosquito density		Inter-individual transmission	Female	NA2	NA	
					Male	NA	NA	
		Larval mosquito density		Density Dependant Prophylaxis	Female	NA	NA	
					Male	NA	NA	
Field	Relative abundance	Years after introduction	Spearman	Ascogregarina|Otu000001	Female		**0.034***	0.18
					Male		0.465	-0.07
				Ascogregarina|Otu000002	Female		**<0.0001*****	0.51
					Male		**<0.0001*****	0.50
				Ascogregarina|Otu000003	Female		**0.0008*****	0.28
					Male		**<0.0001*****	0.40
		Geographical distance	Moran	Ascogregarina|Otu000001	Female		0.371	0.09
					Male		0.246	0.14
					France x Female		**0.034***	-0.43
					France x Male		0.107	0.00
				Ascogregarina|Otu000002	Female		**0.005****	0.48
					Male		0.316	0.03
					France x Female		0.576	-0.31
					France x Male		0.158	-0.03
				Ascogregarina|Otu000003	Female		**0.013***	0.48
					Male		0.426	0.03
					France x Female		0.127	-0.39
					France x Male		0.956	-0.24
	Absolute abundance	Years after introduction	Spearman	Ascogregarina|Otu000001	Female		0.87	-0.04
					Male		0.51	0.15
				Ascogregarina|Otu000002	Female		**0.007****	0.60
					Male		**0.007****	0.55
				Ascogregarina|Otu000003	Female		**0.001*****	0.63
					Male		**0.005****	0.62
Experiment	Parasite abundance	Egg desiccation	GLMM ╪	Desiccation resistance	Female	26.32	**<0.0001*****	
					Male	68.81	**<0.0001*****	

	Adult mosquito density		Inter-individual transmission	Female	15.07	**0.001*****
				Male	20.535	**<0.0001*****
	Larval mosquito density		Density Dependant Prophylaxis	Female	3.10	0.0784
				Male	1.14	0.2855
Flight Activity	Infection status	GLM	Locomotor Activity System	Female	11.58	**0.0151***
				Male	0.48	0.1988

*p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001, ǂ the experimental replicates were used as random variable, [1]refers to data extracted from the literature.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary material.

DNA extraction

Mosquito samples were rinsed three time with sterile water, surface disinfected 5min in ethanol and washed 5 times in sterile water. DNA was extracted from each mosquito individually crushed with three 3mm borosilicate beads and under agitation during 60s at 6m/s with a Fastprep homogenizer (MP Biomedicals). Homogenate was then suspended into a CTAB extraction buffer (2% hexadecyltrimethyl ammonium bromide, 1.4 M NaCl, 0.02 M EDTA, 0.1 M Tris pH 8.0, 0.2% 2-β-mercaptoethanol) and heated for 2h at 60°C. After incubation, 2mg/ml of RNase was added to the extraction buffer and incubated at 37°C for 5 min. DNA was separated from proteins and lipids with an equivalent volume of phenol-chloroform-isoamyl alcohol (25:24:1, v:v:v), vortexed for 15min and centrifuged for 30min at 16,100g. Aqueous phase was harvested and mixed with an equivalent volume of chloroform:isoamyl alcohol (24:1, v:v). The mixture was vortexed for 15min and centrifuged for 30min. The newly separated aqueous phase containing the nucleic acids was harvested. DNA was precipitated with isopropyl alcohol and centrifuged for 30min at 16,100g. DNA pellet were washed with 70% cold ethanol and centrifuged for 30min at 16,100g. Ethanol was removed and DNA was suspended in 20µl of sterile water. DNA extracts were quantified with a spectrophotometer (SAFAS) and diluted at 15ng/µl.

qPCR estimation of the eukaryotic and prokaryotic microbial density

Since variations in OTU abundances may be influenced by variations in absolute densities of microorganisms, qPCR essays were performed on samples for which variations in *As. taiwanensis* abundances were observed. Following the metabarcoding analysis and correlation network analysis, three different groups of *Ae. albopictus* microbiota were observed depending on their infection status with *As. taiwanensis* (see Results section). Prokaryotic and Eukaryotic qPCR essays were performed on 7 males and 7 females from each of those three groups using the same primers than in metabarcoding. Reaction mix contained 1X of SYBR Green iTaq (Bio-Rad), 3µM of each primer and 5ng of DNA. PCR cycles were performed on a CFX96 Touch Real-Time PCR detection system (Bio-Rad) following a 3min initial denaturation at 95 °C followed by 40cycles with 5s of denaturation at 95°C and 1min of hybridization and amplification at 60°C. A final denaturation of the PCR products was conducted by progressively increasing the temperature from 60°C to 90°C at 0.11°C/s. The Ct values were used to quantify the copies number of each gene with a standard range of 10^7 to 10^0 genomic DNA copies of *Escherichia coli* DH5-α for the 16S rDNA and a 10^7 to 10^0 genomic DNA copies of *Saccharomyces cerevisiae* CEN.PK2-2 for the 18SrDNA.

PCR-based estimation of Ascogregarina taiwanensis prevalence

Detection of *As. taiwanensis* in DNA from field collected individuals was conducted with diagnostic PCR using primers targeting the ITS1-5.8SrDNA-ITS2 intergenic region with the previously published AU (5'-ACCGCCCGTCCGTTCAATCG-3') and AT (5'-GAGAAGCCGTCGTCAATACAGC-3') primers (Morales et al., 2005). The mix contained 1X of Hoststart Master mix (Dutscher, 0.2ng/ml of BSA (Promega), 0.1µM of each primer, 0.1X of GC-Enhancer (Biofidal) and 30ng of mosquito genomic DNA. Reactions were performed with a 10min initial denaturation at 96°C, followed by 35 cycles of amplification including 20s at 96°C, 1min at 50°C, 2min at 72°C and a final elongation of 10min at 72°C. Amplification signals were identified under UV light on a Geldoc 2000 system (Biorad) after 20min of electrophoresis migration at 100V on a 1.5% agarose gel colored with Ethidium Bromide.

Design of qPCR primers for the quantification of Ascogregarina taiwanensis

Primers hybridizing *As. taiwanensis* were designed *in silico* based on an alignment of previously published 18S rDNA sequences of Apicomplexa (genebank accession numbers JX131299.1, DQ462455.1, DQ462454.1, JX131298.1, JX131300.1, JX131297.1, EF666482.1, JX131296.1, DQ462456.1, DQ462457.1, DQ462458.1, DQ462459.1, LC109003.1, AY179988.1, AY789075.1, AF129883.1, HQ224956.1, AB071177.1, DQ176427.1, EF024130.1, EF024388.1, EF024744.1, EF024808.1, EF024116.1, EF024963.1, EF024809.1, EF024338.1, EF024673.1, EF024471.1, EF024351.1, EF024347.1, EF024539.1, AY334569.1, KF130567.1, AY334568.1, AY490099.1, AY919720.1, AY919821.1, AY919809.1, KF130566.1, KF130601.1, EU910605.1, EF024102.1, AJ243064.2, KT072989.1, MF998086.1, FN546182.1, FJ459739.1) to select a conserved region in *As. taiwanensis* that diverged from other species. Two specific primers named AscoJSF (5'-CTMGGCTTGACTTCGGTCT-3') and AscoJSR (5'-TTCCATGCTGGAGTATTCAAGG-3') were designed and produced a fragment of 150bp. A standard concentration of the fragment cloned in a vector was prepared. It was first amplified with a PCR mix containing 1X of buffer, 1.5mM of $MgCl_2$, 200µM of dNTP, 200nM of each primer, 0.625 Units of Taq DNA Polymerase (New England Biolabs) and 30ng of DNA from a parasitized mosquito individual. The PCR program was ran with 95°C of initial denaturation for 4min, followed by 35 cycles including 30s at 95°C, 30s at 56°C, 15s at 68°C and a final elongation of 5min at 68°C. The PCR product was purified with a PCR purification kit MinElute (Qiagen), cloned into a PCR 2.1-TOPO TA vector (Invitrogen) and transferred into competent *Escherichia coli* Top10 cells (Invitrogen) following manufacturers' recommendations. Competent cells containing the vector and the insert were selected on LB agar supplemented with 25µg/ml of Kanamycin and 20µg/ml of Xgal and 23.8µg/ml of IPTG after 24h of incubation at 37°C. Presence of the insert was also verified with PCR on white clones that developed on

the medium. DNA from the competent cells was extracted with a DNeasy Blood and Tissue kit (Qiagen) following manufacturers' recommendations and linearized 1h at 37°C in a mix containing 0.2μg/μl of DNA, 1X of SurE/Cut Buffer B and 10U of BamH1 endonuclease (Roche). Reaction was inactivated at 65°C for 15min. This digested DNA was purified with a MinElute PCR purification kit (Qiagen), quantified with Quanti-it dsDNA BR Assay kit (Invitrogen) and used as a standard for qPCR with concentrations ranging from 10^8 to 10^0 copies. Reaction mixes for quantification included 0.5μM of each primer, 1X of iTaq Universal SYBR Green Supermix (Bio-Rad) and 5ng of DNA. The qPCR program included a 5min initial denaturation at 95°C, followed by 40 cycles with 15s at 95°C, 30s at 56°C and 10s at 60°C followed by a progressive denaturation of the qPCR products from 65°C to 97°C at 0.11°C/s.

Metabarcoding of eukaryotic and prokaryotic microbial communities

Both PCR amplifications were conducted under sterile conditions with 1X of Hotstart 5X Bioamp Master Mix (Biofidal), 0.2μM of each primers, 0.2mg/ml of Bovine Serum Albumin (New England Biolabs), 0.4X of GC rich Enhancer (Biofidal). For 16S rDNA, PCR program started with 10min of initial denaturation at 96°C followed by 35 cycles including 20s of denaturation at 96°C, 1min of annealing at 54°C, 30s of extension at 72°C and ended with a final extension of 10min at 72°C. For 18S rDNA, PCR program started with 10min of initial denaturation at 96°C followed by 35 cycles including 20s of denaturation at 96°C, 30s of annealing at 54°C, 30s of extension at 72°C and ended with a final extension of 10min at 72°C. Three negative controls corresponding to blank extractions and PCR were included for each gene.

Details concerning the mother pipelines used for bioinformatics treatment: A total of 12,836,921 and 14,248,382 reads were obtained from 16S and 18S rDNA respectively and were demultiplexed using the Mothur pipeline (69). Reads were trimmed based on size (270-310bp and 400-550bp for the 16S and 18S rDNA, respectively) and sequences were conserved if they contained no ambiguous position and if both strands of the paired-end reads aligned together. Chimeras were removed using CHIMERAVSEARCH. The alignment was performed using Silva v.132 database for bacteria and eukarya and reads that did not align were filtered out. However, for sequences that matched gregarine, a supplemental local alignment (BLAST) was performed against the Genbank database due to the lack of reference sequences in Silva. Clustering was performed using Opticlust method allowing a maximum dissimilarity rate of 3%.

Table S1. Aedes albopictus population used in the current study.

Country	Origin site	Months of sample	Years of sample	Years after introduction	Sex	n. samples prevalence	n. samples metabarcoding	Climate	GPS coordinates
France	Saint-Priest	July	2012	0	males	8	10	Semi-continental	N : 45°41'49''
					females	28	10		E : 4°58'50'
	Saint-Priest	August	2018	1	males	24	-	Semi-continental	N: 45°43'8'
					females	28	-		E : 4°55'9''
	Oullins	July	2018	1	males	20	10	Semi-continental	N: 45°42'42''
					females	18	10		E : 4°48'2'
	Feyzin	July	2018	1	males	18	10	Semi-continental	N: 45°40'25'
					females	22	10		E : 4°51'33''
	Villeurbanne	July	2018	5	males	26	10	Semi-continental	N : 45°46'18''
						30	-		E : 4°53'24''
	Nice	August	2012	7	males	16	-	Mediteraneen	N : 43°41'60''
					females	20	10		E : 7°17'33''
	Portes-Les-Valence	July	2013	2	males	7	10	Semi-continental Mediteraneen	N : 44°52'8''
					females	31	10		E : 4°52'9'
Italy	Turin	September	2011	15	males	17	10	Mediteraneen	N: 45°4'13'
					females	13	10		E : 7°41'12''
Spain	Barcelona	July	2013	9	females	22	10	Mediteraneen	N: 41°23'6.2'
									E : 2°10'24''
USA	Atlanta	October	2018	33	males	25	10	Humid Subtropical	N : 33°44'56''
					females	24	10		O : 84°23'16''
Madagascar	Diego Suarez	NA	2009	>105	males	18	-	Humid Tropical < 900m	S : 12°16'51.24''
					females	14	10		E : 49°17'27.092''
	Toamasina	December	2010	>105	males	22	10	Humid Tropical < 900m	S : 18°8'59.64''
					females	19	10		E : 49°24'8.312'

Country	Site	Month	Year	Population age	Sex	Sample size	Infected	Climate	Coordinates
	Tsimbaza-Park	December	2010	>105	males	13	10	Tropical > 900m	S : 18°55'40.395''
					females	25	10		E : 47°31'38.5'
Thailand	Chiang Mai	February	2019	Native	males	8	8	Tropical	N : 18° 42' 46.1" E : 98° 56' 17.1"
Vietnam	Hô-chi-minh	October	2012	Native	females	20	10	Tropical	N : 10°47'19'' E : 106°42'19''
	Binh Duong	October	2012	Native	females	19	10	Tropical	N : 10°47'19'' E : 106°42'19'
	Vüng Tau	October	2012	Native	males	25	10	Tropical	N : 10°22'26'' E : 107°4'13''

115 **Table S2. Number and details on mosquito prevalence from this study and the**
116 **previously published studies.**

Country	Site	Sex	Population age	Latitude (N)	Longitude (E)	Sample size	Uninfected	Infected	Prevalence	From
France	Feyzin	Female	1	45.6736111	4.85916666666666	22	22	0	0,00	This study
France	Feyzin	Male	1	45.6736111	4.85916666666666	18	18	0	0,00	This study
France	Nice	Female	7	43.7	7.29249999999999	25	3	22	0,88	This study
France	Nice	Male	7	43.7	7.29249999999999	16	2	14	0,88	This study
France	Oullins	Female	1	45.7116667	4.80055555555555	18	16	2	0,11	This study
France	Oullins	Male	1	45.7116667	4.80055555555555	20	19	1	0,05	This study
France	Portes-lès-Valence	Female	2	44.8688889	4.86916666666666	31	24	7	0,23	This study
France	Portes-lès-Valence	Male	2	44.8688889	4.86916666666666	7	2	5	0,71	This study
France	Saint Priest 2	Male	3	45.7188889	4.91916666666666	24	12	12	0,50	This study

Country	Location	Sex		Latitude	Longitude					Source
France	Saint Priest 1	Female	0	45.6858333	4.98055555555555	28	27	1	0,04	This study
France	Saint Priest 1	Male	0	45.6858333	4.98055555555555	8	6	2	0,25	This study
France	Saint Priest 2	Female	3	45.7188889	4.91916666666666	28	23	5	0,18	This study
France	Villeurbanne	Female	5	45.7711111	4.88888888888888	30	6	24	0,80	This study
France	Villeurbanne	Male	5	45.7188889	4.98055555555555	26	13	13	0,50	This study
Italy	Torino	Female	15	45.0702778	7.68666666666666	13	1	12	0,92	This study
Italy	Torino	Male	15	45.0702778	7.68666666666666	17	0	17	1,00	This study
Madagascar	Diego-Suarez	Female	>105	-12.2808333	49.2908333333333	18	1	17	0,94	This study
Madagascar	Diego-Suarez	Male	>105	-12.2808333	49.2908333333333	14	0	14	1,00	This study
Madagascar	Tamatave	Male	>105	-18.1497222	49.4022222222222	22	9	13	0,59	This study
Madagascar	Tamatave	Female	>105	-18.1497222	49.4022222222222	19	7	12	0,63	This study
Madagascar	Tsimbazaza	Male	>105	-18.9277778	47.5272222222222	13	2	11	0,85	This study
Madagascar	Tsimbazaza	Female	>105	-18.9277778	47.5272222222222	25	4	21	0,84	This study
Spain	Barcelona	Female	9	41.385	2.17333333333333	22	2	20	0,91	This study
USA	Atlanta	Female	33	33.7488889	-84.3877777777779	24	6	18	0,75	This study
USA	Atlanta	Male	33	33.7488889	-84.3877777777779	25	10	15	0,60	This study
Vietnam	Binh Duong	Female	Native	10.7886111	106.705277777777	19	7	12	0,63	This study
Vietnam	Ho Chi Minh City	Female	Native	10.7886111	106.705277777777	20	9	11	0,55	This study

Vietnam	Vung Tau	Female	Native	10.3738889	107.070277777777	25	4	21	0,84	This study
Thailand	Phuket	Male	Native	7.8788889	98.3983333333333	21	11	10	0,48	This study
France	Feyzin	Female	1	45.6736111	4.85916666666666	22	0	22	1,00	This study
France	Feyzin	Male	1	45.6736111	4.85916666666666	18	0	18	1,00	This study
France	Nice	Female	7	43.7	7.29249999999999	25	22	3	0,12	This study
France	Nice	Male	7	43.7	7.29249999999999	16	14	2	0,13	This study
France	Oullins	Female	1	45.7116667	4.80055555555555	18	2	16	0,89	This study
France	Oullins	Male	1	45.7116667	4.80055555555555	20	1	19	0,95	This study
France	Portes-lès-Valence	Female	2	44.8688889	4.86916666666666	31	7	24	0,77	This study
France	Portes-lès-Valence	Male	2	44.8688889	4.86916666666666	7	5	2	0,29	This study
France	Saint Priest 2	Male	3	45.7188889	4.91916666666666	24	12	12	0,50	This study
France	Saint Priest 1	Female	0	45.6969444	4.98055555555555	28	1	27	0,96	This study
France	Saint Priest 1	Male	0	45.6969444	4.98055555555555	8	2	6	0,75	This study
France	Saint Priest 2	Female	3	45.7188889	4.91916666666666	28	5	23	0,82	This study
France	Villeurbanne	Female	5	45.7711111	4.88888888888888	30	24	6	0,20	This study
France	Villeurbanne	Male	5	45.7188889	4.91916666666666	26	13	13	0,50	This study
Italy	Torino	Female	15	45.0702778	7.68666666666666	13	12	1	0,08	This study
Italy	Torino	Male	15	45.0702778	7.68666666666666	17	17	0	0,00	This study
Madagascar	Diego-Suarez	Female	105	-12.2808333	49.2908333333333	18	17	1	0,06	This study
Madagascar	Diego-Suarez	Male	105	-12.2808333	49.2908333333333	14	14	0	0,00	This study

Country	Location	Sex	N	Lat	Long					Reference
Madagascar	Tamatave	Male	105	-18.1497222	49.4022222222222	22	13	9	0,41	This study
Madagascar	Tamatave	Female	105	-18.1497222	49.4022222222222	19	12	7	0,37	This study
Madagascar	Tsimbazaza	Male	105	-18.9277778	47.5272222222222	13	11	2	0,15	This study
Madagascar	Tsimbazaza	Female	105	-18.9277778	47.5272222222222	25	21	4	0,16	This study
Spain	Barcelona	Female	9	41.385	2.17333333333333	22	20	2	0,09	This study
USA	Atlanta	Female	33	33.7488889	-84.38777777777779	24	18	6	0,25	This study
USA	Atlanta	Male	33	33.7488889	-84.38777777777779	25	15	10	0,40	This study
Vietnam	Binh Duong	Female	200	10.7886111	106.705277777777	19	12	7	0,37	This study
Vietnam	Ho Chi Minh City	Female	200	10.7886111	106.705277777777	20	11	9	0,45	This study
Vietnam	Vung Tau	Female	200	10.3738889	107.070277777777	25	21	4	0,16	This study
Thailand	Phuket	Male	200	7.8788889	98.3983333333333	21	10	11	0,52	This study
USA	East St Louis	Larvae	14	38.616405	-90.159821	55	0	55	1,00	Aliabadi et al. 2002
USA	Peoria_1	Larvae	2	40.703544	-89.579084	49	14	35	0,71	Aliabadi et al. 2002
USA	Peoria_2	Larvae	1	40.703545	-89.579086	52	43	9	0,17	Aliabadi et al. 2002
USA	Florida_1	Larvae	4	27.994401	-81.760253	883	300	583	0,66	Blackmore et al. 1995
USA	Florida_2	Larvae	1	27.994402	-81.760254	535	433	102	0,19	Blackmore et al. 1995
USA	Hawaï	Mixed	117	19.896766	-155.582782	118	19	99	0,84	Seabourn et al. 2020
USA	Orleans_Parish_1	Female	9	90,0715323	0,716049383	181	55	126	0,70	Comiskey et al. 1999
USA	Orleans_Parish_2	Male	9	90,0715323	0,716049383	62	14	48	0,77	Comiskey et al. 2000

120 **Figure S1. World map representing the collection sites of mosquitoes' populations used**

121 **in this study.**

Figure S2. Microbiota composition dissimilarity separation according to mosquito population ages. For both (A, B) female and (C, D) male individuals the dissimilarity (bray-curtis index) of (A, C) prokaryotic and (B, D) eukaryotic microbiota communities was represented. The points are colored with a gradient ranging from blue to red based on the age of each population (years after introduction). Native populations were set to 200 years.

Figure S3. Microbiota dissimilarity separation according to climate. For both (A, B) female
and (C, D) male individuals the dissimilarity (bray-curtis index) of (A, C) prokaryotic and (B, D)
eukaryotic microbiota communities was represented. The points are colored with a gradient
ranging from blue to red based on climatic regions. Native populations were set to 200 years.

Figure S4. Microbiota dissimilarity correlation with geographical distances. Spearman correlation analysis between the microbiota community dissimilarity estimated with the Bray-Curtis index and geographical ellipsoid distance. Correlations were represented for both (A, B) female and (C, D) male individuals and for (A, C) prokaryotic and (B, D) eukaryotic microbiota communities. The points are colored with a gradient ranging from blue to red based on geographical distance between both points being compared. The correlation coefficient (Rs) and p-value are represented on each graph.

Group: F = 2.79, *p* = 0.0754
Sexe: F = 0.19, *p* = 0.668
Group x Sex: F = 0.84, *p* = 0.442

Group: F = 210.94, ***p* < 0.0001*****
Sexe: F = 4.16 , ***p* = 0.041***
Group x Sex: F = 1.71, *p* = 0.42

Figure S5. Quantification of microbial densities. Quantifications were performed through

qPCR of the (A) 16S rDNA genes for the eukaryotic community or (B) 18S rDNA gene for the

prokaryotic community. Females and males are respectively colored in red and blue. G1

represent samples from various origins for which the *Ascogregarina taiwanensis* Otu000001

dominates the microbiota. G2 represents samples for which *As. taiwanensis* Otu000002 and

Otu000003 dominates the microbiota. Finally, G3 represents samples that are not dominated

by *As. taiwanensis.*