**Transposon accumulation at xenobiotic gene family loci: a**

**comparative genomic analysis in aphids**

Tobias Baril[1,2]*, Adam Pym[1,3], Chris Bass[1,4], Alex Hayward[1,5]*

[1] Centre for Ecology and Conservation, University of Exeter, Penryn Campus,

Cornwall, TR10 9FE, UK

[2] ORCID: 0000-0002-5936-7531

[3] ORCID: 0000-0001-8754-4016

[4] ORCID: 0000-0002-2590-1492

[5] ORCID: 0000-0001-7413-718X

* Corresponding Author information:

Tobias Baril: tb529@exeter.ac.uk

Alex Hayward: alex.hayward@exeter.ac.uk

## Abstract

The evolution of resistance is a major challenge for the sustainable control of pests and pathogens. Thus, a deeper understanding of the evolutionary and genomic mechanisms underpinning resistance evolution is required to safeguard health and food production. Several studies have implicated transposable elements (TEs) in xenobiotic resistance evolution in insects. However, analyses are generally restricted to one insect species and/or one or a few xenobiotic gene families (XGFs). We examine evidence for TE accumulation at XGFs by performing a comparative genomic analysis across 20 aphid genomes, considering major subsets of XGFs involved in metabolic resistance to insecticides: Cytochrome P450s, glutathione S-transferases, esterases, UDP-glucuronosyltransferases, and ABC transporters. We find that TEs are significantly enriched at XGFs compared to other genes. XGFs show similar levels of TE enrichment to housekeeping genes. But unlike housekeeping genes, XGFs are not constitutively expressed in germline cells, supporting the selective enrichment of TEs at XGFs, rather than enrichment due to chromatin availability. Hotspots of extreme TE enrichment occur around certain XGFs. In aphids of agricultural importance, we find particular enrichment of TEs around cytochrome P450 genes with known functions in the detoxification of synthetic insecticides. Our results provide evidence supporting a general role for TEs as a source of genomic variation at host XGFs, and highlight the existence of considerable variability in TE content across XGFs and host species. These findings demonstrate the need for detailed functional verification analyses to clarify the significance of individual TE insertions and elucidate underlying mechanisms at TE-XGF hotspots.

## Introduction

Xenobiotics are substances that are foreign to an organism, including naturally occurring plant allelochemicals and man-made insecticides (Li et al. 2007). Xenobiotic toxicity can

present a strong selective pressure, leading to the rapid evolution of resistance via efficient

xenobiotic removal, metabolism, and tolerance (Chung et al. 2007; Li et al. 2007).

Consequently, numerous insects have evolved resistance to both plant allelochemicals,

produced for plant defence, and synthetic insecticides, developed to protect economically

important crops from insect pests, or humans and livestock from insect vectored diseases

(Sparks et al. 2021; Sternberg and Thomas 2018). The intensive use of numerous synthetic

insecticides has led many insects to evolve resistance to multiple insecticide classes (Li et

al. 2007; Bradshaw et al. 2016), severely affecting our ability to control target insects (Singh

et al. 2021). The emergence of multiple insecticide resistance is a major global challenge,

threatening global food security in the case of agricultural pests, and human and animal

health in the case of disease vectors (Worner and Gevrey 2006; Deutsch et al. 2018).

Two major mechanisms are frequently implicated in conferring xenobiotic resistance

across a wide range of arthropods (Li et al. 2007; Kliot and Ghanim 2012): (i) 'target-site

resistance', involving structural changes (mutations) in the gene encoding the insecticide

target protein, that make it less sensitive to the toxic effect of the insecticide; (ii) 'metabolic

resistance', involving the increased production or activity of metabolic enzymes that break

down or sequester the insecticide. Regarding the latter, several major gene families are

associated with metabolic resistance, which act during three distinct phases of xenobiotic

metabolism (Kennedy and Tierney 2013): (i) Cytochrome P450 monooxygenases (CYP

genes); (ii) glutathione S-transferases (GSTs), esterases, and UDP-

glucuronosyltransferases (UGTs); (iii) ATP-binding cassette transporters (ABC transporters).

We briefly summarise these three phases below. During phase I, CYP genes catalyse

functionalisation reactions, where hydrophobic xenobiotics are typically converted to more

water-soluble metabolites through the addition of functional groups (Kennedy and Tierney

2013). CYP genes are diverse enzymes involved in several purposes from biosynthesis to

metabolism, and are sometimes referred to as 'nature's blowtorch', due to their high-valence

iron chemistry oxidation mechanism (Guengerich 2009). CYP genes can mediate resistance

to all classes of insecticides due to their broad substrate specificity and versatility, whilst they

are also involved in several other processes, such as juvenile hormone and ecdysteroid

metabolism, and fatty acid biosynthesis (Feyereisen 2005; Kennedy and Tierney 2013;

Feyereisen 1999). During phase II, GSTs, UGTs, and esterases catalyse conjugation

reactions between phase I substrates and endogenous molecules to form water-soluble

metabolites (Kennedy and Tierney 2013). During phase III, ABC transporters, which are cell

membrane transport proteins that efflux toxins and modified toxins from the cell, export

xenobiotics and products of phase I and II (Gott et al. 2017). In addition, ABC transporters

can also block the cellular import of xenobiotics to protect the host, which is termed phase 0.

(Gott et al. 2017). Insecticide resistance can arise via mutations acting on the genes

involved in the processes described above through various genetic mechanisms, including

up-regulation, changes to coding sequence, and gene amplification (Li et al. 2007). It is also

important to note that many xenobiotic gene family (XGF) members are involved in other

metabolic processes beyond xenobiotic resistance. For example, some GSTs have roles in

processes including eye pigmentation, intracellular transport, and cell signalling pathways

(Ketterman et al. 2011; Ranson and Hemingway 2005), whilst ABC transporters have

numerous roles in processes including heme biosynthesis, iron homeostasis, and protection

against oxidative stress (Dermauw and Van Leeuwen 2014). UGTs also have roles in the

modulation of endobiotics and olfactory processes (Ahn et al. 2012), and esterases have

roles in neurodevelopment and pheromone signalling (Gilbert and Gill 2014).

There is growing evidence that transposable elements (TEs) can play an important

role in the evolution of xenobiotic resistance in insects (Gilbert et al. 2021; Rostant et al.

2012), with examples from several major lineages: Diptera: *Drosophila* (Feyereisen 2005;

Bogwitz et al. 2005; Schmidt et al. 2010; Aminetzach et al. 2005; Remnant et al. 2013;

Mateo et al. 2014; Salces-Ortiz et al. 2020), *Culex* (Itokawa et al. 2011; Darboux et al. 2007)*,*

*Musca domestica* (Li et al. 2007; Kasai and Scott 2001), *Anopheles funestus* (Weedall et al.

2020) Lepidoptera: *Heliothis* (Gahan et al. 2001; Yang et al. 2007; Zhao et al. 2010; Chen

and Li 2007), *Pectinophora gossypiella* (Fabrick et al. 2011; Wang et al. 2019) Hemiptera:

*Myzus persicae* (Singh et al. 2020; Panini et al. 2021). TEs are DNA sequences capable of

moving from one genomic location to another within the genome. TEs occur in nearly all

eukaryotic genomes, and are implicated in the evolution of host genomic novelty through

diverse processes including the modification of regulatory networks, chromosomal

rearrangements, exon shuffling, and donation of coding sequence (Bourque et al. 2018;

Wells and Feschotte 2020; Cosby et al. 2019; Sundaram et al. 2014). In the case of

xenobiotic resistance, TEs are reported to have contributed to evolvability via myriad

mechanisms, including: gene amplification and duplication (Singh et al. 2020; Remnant et al.

2013), knockout of a susceptible allele in heterozygous individuals (Panini et al. 2021),

increases in detoxification gene expression (Bogwitz et al. 2005; Itokawa et al. 2011), allelic

succession leading to increases in resistance gene copy number (Schmidt et al. 2010), and

alternative splicing and production of truncated proteins that prevent interactions with

xenobiotics (Darboux et al. 2007; Gahan et al. 2001; Yang et al. 2007; Zhao et al. 2010).

Studies examining the contribution of TEs to resistance evolution suggest that there

can be an additive effect of successive TE insertions at focal host loci over time. For

example, in the well-known case of the cytochrome P450 (CYP) gene *CYP6G1* and DDT

resistance in *Drosophila*, successive TE insertions are linked with an increasing ability to

detoxify the insecticide (Catania et al. 2004; Daborn et al. 2002; Chung et al. 2007; Schmidt

et al. 2010). A similar process involving the accumulation of TEs, in this case at *CYP6CY3,*

is implicated in the evolution of resistance to nicotine, following a host plant shift to tobacco

in the aphid *Myzus persicae nicotianae* (Puinean et al. 2010; de Little et al. 2017). More

widely, there is evidence that the selective accumulation of TEs at rapidly evolving host loci

under strong selection may represent a general evolutionary process. For example, studies

have reported the enrichment of TEs at other gene classes, such as immune genes and

those involved in responses to external stimuli (van de Lagemaat et al. 2003; Song et al.

1998). Existing studies strongly suggest that TEs are involved in individual cases of

resistance to certain insecticides, but there is currently very limited understanding of how

associations with TEs vary among different classes of XGFs, or among whole clades of

insects. However, patterns in TE accumulation remain poorly described, and it is unclear to

what extent TEs may be enriched at XGFs compared to other host genes, or how much

variability exists across XGFs.

Here we consider patterns of TE accumulation at XGFs in the genomes of 20 species

sampled from across aphid phylogeny, for which high-quality genome assemblies are

available. The aphid family is a relevant clade within which to explore the genomic bases of

insecticide resistance, as it includes numerous globally important crop pests responsible for

causing billions of US dollars of crop losses annually (Van Emden and Harrington 2017), and

members that have been intensively targeted with insecticides and evolved resistance. For

example, the severely damaging global crop pest *Myzus persicae* (the green peach aphid) is

resistant to at least 82 insecticides, via at least eight different resistance mechanisms (Bass

et al. 2014; Mota-Sanchez and Wise 2023; Singh et al. 2021). We hypothesise similar

broadscale patterns for TE accumulation at XGFs among aphid genomes, given similar

challenges from xenobiotics that target conserved biological pathways. Meanwhile, given the

strong selection pressure that can arise from xenobiotic exposure, and repeated xenobiotic

challenges over time, we anticipate an ability to detect patterns at the species-level.

Since the TE landscapes of aphids are poorly described, we begin by characterising

patterns of TE content across our focal genomes. After this, we examine key fundamental

questions relating to the pattern of associations between TEs and XGFs in aphid genomes.

(i) Firstly, we test whether XGFs are enriched for TEs compared to other host genes, to

evaluate the overall signature of TE accumulation at XGFs. (ii) Secondly, we examine the

extent to which TE content varies among individual XGFs and among major XGF

classifications, to determine if contributions from TEs are more pronounced at certain types

of XGF. (iii) Thirdly, we consider if specific TE classifications are enriched at XGFs (e.g DNA

TEs, rolling circles, *Penelope*-like elements, LINEs, SINEs, and LTRs), to explore if

particular TEs are predisposed to contribute to the evolution of host resistance. (iv) Fourthly,

we assess whether patterns of TE accumulation at XGFs vary among aphid species to

examine if patterns vary across aphid phylogeny. (v) Fifthly, we investigate TE enrichment at

XGFs that have demonstrated associations with the detoxification of synthetic insecticides,

to test if these show particularly strong signatures of TE recruitment. (vi) Lastly, we explore

whether our results suggest a selective role for patterns in TE accumulation at XGFs versus

the alternative explanation of increased availability for TE insertion due to chromatin

availability.

## Results

### *Transposable Element Landscapes in Aphids*

TEs make considerable contributions to genome size in eukaryotes (Kidwell 2002; Gregory

2005). In aphids, we find that TE content varies from 8.21% (*Melanaphis sacchari*) to

35.52% (*Metopolophium dirhodum*) of total genome size (Figure 1B, Supplemental Table

S1). This is in accordance with levels of TE content identified among a large-scale survey of

196 insect species, where TE content varied from 0.08% to 53.93% of total genome size

(Peccoud et al. 2017) . Similarly, consistent with findings in other eukaryotic lineages, we

report a significant strong positive association between genome size and TE content in

aphids (Linear regression, $F_{1,19}$=46.35, p<0.01, $R^2$=0.69, Supplemental Figure S1). Aphids

show variation in sexual and asexual lifecycles (Van Emden and Harrington 2017). It is

hypothesised that asexual lineages would suffer from Muller's ratchet as deleterious

mutations accumulate (Muller 1964). Despite this, we find no significant difference in TE

content between holocyclic and anholocyclic lineages (Wilcoxon Rank Sum, $W_{14,6}$ = 35,

p>0.05) (Figure 1B).

TEs have a non-random distribution within host genomes (Bourque et al. 2018). As

expected, given the predominantly deleterious effects of TE insertion into host genes

(Bourque et al. 2018; Sultana et al. 2017), exonic regions exhibit the lowest levels of TE

sequence content in aphid genomes (mean TE content in exons = 1.90% of total genome

size). Across 19 of the 20 aphid genomes considered here, the greatest TE coverage

(expressed as proportion of total genome size) is found in 5' and 3' gene flanking regions,

where it varies between 3.49% (*M. sacchari*) and 17.94% (*M. dirhodum*) (Figure 2B).

However, this pattern is not universal across all aphid species. In *Sipha flava,* the majority of

TE coverage is present in intronic regions (10.60%), while in *M. cerasi*, the majority occurs in

intergenic space (11.28%) (Figure 2B).

The amount of TE sequence and the frequency of TE insertions in genic versus

intergenic regions in aphid genomes is not significantly correlated with genome

compactness, described as the ratio of genic (intron and exon) to intergenic (gene flanks and

intergenic space) base pairs (Spearman's Rank, S=1680, p>0.05, rho=-0.09, Supplemental

Figure S2, Supplemental Table S2). *S. flava* has the most compact genome, with a

genic:intergenic base pair ratio of 1.50:1 (250.0 Mb:166.8 Mb), and so presumably a higher

likelihood of TE insertion into genic regions compared to intergenic space, leading to the

observation that the majority of TE insertions are found in intronic regions. In contrast, *M.*

*cerasi* has the least compact genome, with a much lower genic:intergenic base pair ratio of

0.13:1 (76.1 Mb:559.2 Mb). In less compact genomes such as *M. cerasi*, TEs can

accumulate in expanded intergenic regions that act as 'genomic safe havens', where

insertion is less likely to exert deleterious effects (Arkhipova 2018). For context, the human

genome (GCF_009914755.1), has a TE content of 44%-69% (Nurk et al. 2022; de Koning et

al. 2011) and a genic:intergenic base pair ratio of 0.55:1 (1717.7 Mb:3117.3 Mb), making it

several times more compact than *M. cerasi*, but much less compact than *S. flava.*

Across all aphid species and in most individual genomes, the dominant TE

classification is DNA TEs, which comprise between 36.6% (*Schlechtendalia chinensis*) and

68.9% (*Aphis craccivora*) of total TE content ($\mu$ = 45.27%, SD = 18.00%, Figure 1B, Table 1,

Supplemental Table S1). SINEs are present only in the genomes of 7 of the 20 species

considered (Supplemental Table S1) and, where present, they comprise just 0.009% (*A.*

*pisum*) to 0.1% (*A. craccivora*) of total TE content.

There is little evidence of ancient TE activity in aphid genomes, as indicated by the

relatively low number of TEs showing high divergence from their respective TE consensus

models in repeat landscape plots (Figure 1C). On the other hand, there is considerable

evidence of relatively recent TE activity in most aphid species, with particular evidence of

very recent TE activity in *Myzus persicae* and *M. dirhodum*, indicated by large numbers of

TEs with low genetic distance from their respective consensus sequences (Figure 1C). The

relative absence of ancient TE activity is presumably a consequence of a relatively rapid

genomic turnover rate in aphids, and contrasts strongly with TE landscapes in mammals,

where a low rate of genomic turnover leads to considerable accumulations of ageing TEs

(Blass et al. 2012). Instead, the dynamics we observe for aphids are consistent with patterns

reported for other insects, such as lepidopterans (Lavoie et al. 2013; Baril and Hayward

2022).

The extent to which TE content differs among the genomes of closely related species

can vary greatly. We find no significant effect of phylogeny on TE abundance (No. of TEs

present in a genome) or TE diversity (No. distinct TE families present in a genome (Wicker

et al. 2007)) in aphids (Phylogenetic GLMM, TE abundance: posterior mean = 0.592, 95%

HPD = 0.005,0.999; TE diversity: posterior mean = 0.022, 95% HPD = 0.000,0.111). Thus,

more closely related aphid species do not possess similar numbers of TEs or a similar level

of TE diversity. We also find that most TE families present in aphid genomes are species-

specific, with almost half of the TE families present in a particular aphid genome being

unique to that species (μ = 46.97%) (Figure 2A). On average, 10.02% of the TE families

identified in an aphid genome are shared among all 20 species (Figure 2A). The remaining

43.01% of TE families are shared in similar proportions among 2 to 19 aphid species (i.e. μ =

2.53% per category, Figure 2A). The strong signature of species-specificity observed for

aphid TE families suggests that the independent gain of new TE families is a major factor

driving aphid TE landscapes. Collectively, these findings suggest a dynamic repeat

landscape in aphid genomes, characterised by relatively frequent gain and turnover of TEs.

## 231  *TEs are enriched at XGFs compared to other host genes*

Previous studies have implicated TEs in the evolution of xenobiotic resistance across a

range of insects and resistance loci (Li et al. 2007). However, the extent to which TE

association at XGFs represents a general mechanism for the evolution of new resistance

phenotypes remains unclear.

Across all aphid genomes considered here, we find significant enrichment of TEs,

both in TE coverage and TE count, at XGFs in comparison to all other genes excluding

XGFs (TE coverage: Wilcoxon Rank Sum, $W_{6488,401787}$ = 732,129,770, p<0.01; TE count:

Wilcoxon Rank Sum, $W_{6488,401787}$ = 770,552,556, p<0.01) (Figure 3, Supplemental Figure

S3). Specifically, XGFs in aphids have a mean of 2.4x TE coverage and 2.1x TE count

compared to all other genes.

## 243  *TE content varies among individual XGF loci*

We observe an enrichment of TEs around all major XGF categories considered compared to

other host genes: Cytochrome P450s (CYPs), 2.4x TE coverage, 2.1x TE count; glutathione

S-transferases (GSTs), 2.7x TE coverage, 2.3x TE count; esterases, 2.0x TE coverage, 1.8x

TE count; UDP-glucuronosyltransferases (UGTs), 2.0x TE coverage, 1.7x TE count; ABC

transporters (ABCs), 2.8x TE coverage, 2.4x TE count (Figure 3, Supplemental Figure S3,

Supplemental Table S3). Whilst all major XGF categories are enriched for TEs, there is no

significant difference in relative enrichment among XGF types, suggesting relatively equal

levels of enrichment across XGF types (Kruskal-Wallis, TE sequence: $\chi^2_4$ = 7.72, p = 0.10;

TE count: $\chi^2_4$ = 4.25, p = 0.37). However, considerable variability in TE content is present

within XGF categories, attributable to extremely large accumulations of TEs at certain

individual XGFs (Table 2, Figure 4). Some of these XGFs are unusually large in terms of

sequence length compared to the mean size for the XGF type in question, as indicated by

gene size *Z*-scores > 3 (Table 2), which is potentially a consequence of the increased

presence of TEs in their intronic regions (Figure 4). To investigate this, we consider gene

sizes with genic TEs removed. We find that 5/18 of the most TE-enriched XGFs have

significantly inflated gene sizes when TE insertions are removed, indicated by a *Z*-score of >

3. This suggests that these loci were significantly larger than average before accounting for

TE contributions. However, the remaining 13 highly TE-enriched XGFs have gene sizes

within the expected range in the absence of TEs, suggesting that TEs are responsible for the

significant increases in gene size at the majority of TE-enriched XGF loci. Individual XGFs

with the greatest enrichment of TE coverage and TE count are listed in Table 2 (i.e. >35kb

TE sequence and >80 TE insertions, compared to a non-XGF mean of 1.5kb TE sequence

and 1.48 TE insertions). Whilst all XGF types are represented in the TE hotspots listed in

Table 2, there are markedly more hotspots at UGTs and CYP genes, and fewer at ABCs,

GSTs, and Esterases.

## 270    *Specific TE types are enriched at XGFs*

While most TE types contribute to enrichment at XGFs, overall enrichment is primarily

driven by increases in DNA TE content, due to the high frequency of DNA TEs in aphid

genomes, where they comprise almost half of total TE content (Table 3).

The TE type with the greatest degree of enrichment in coverage around XGFs in

comparison to other genes is SINEs, although there is a slight non-significant reduction in

SINE count at XGFs (**TE coverage**, 3.2x enrichment: XGFs μ = 1,144bp, all other genes μ =

358bp; Wilcoxon Rank Sum, $W_{9,482} = 439$, p < 0.01; **TE count,** 0.82x enrichment: XGFs μ =

1.00, all other genes μ = 1.22; Wilcoxon Rank Sum, $W_{9,482} = 2592$, p = 0.14). This is notable,

since SINEs are generally very scarce in aphid genomes, and suggests a particular retention

of SINE sequence at XGFs (Figure 1B, Figure 3, Table 1, Supplemental Table S1).

However, given their scarcity, the relative contribution that SINEs make to TE enrichment at

XGFs compared to other TE types is extremely low. Conversely, DNA TEs are the most

abundant TE type in aphid genomes, and they are significantly enriched at XGFs both in TE

coverage and TE count (**TE coverage**, 2.0x enrichment: XGFs μ = 4,026bp, all other genes

μ = 1,974bp; Wilcoxon Rank Sum, $W_{6156,336473} = 616,739,338$, p < 0.01; **TE count**, 6.4x

enrichment: XGFs μ = 10.05, all other genes μ = 1.57; Wilcoxon Rank Sum, $W_{6156,336473} =$

$1,267,581,920$, p < 0.01) (Figure 3). Rolling circle TEs show significant, but lower levels of

enrichment (**TE coverage**, 1.1x enrichment: XGFs μ = 794bp, all other genes μ = 728bp;

Wilcoxon Rank Sum, $W_{1801,55720} = 46,195,509$, p<0.01; **TE count**, 1.8x enrichment: XGFs μ =

2.14, all other genes μ = 1.22; Wilcoxon Rank Sum, $W_{1801,55720} = 56,336,944$, p < 0.01). In

contrast, LINEs do not show significant levels of enrichment at XGFs in terms of TE

coverage, but there is a significant enrichment of LINE TE count (**TE coverage**, 1.03x

enrichment: XGFs μ = 992bp, all other genes μ = 959bp; Wilcoxon Rank Sum, $W_{2575,101686} =$

$128,128,748$, p = 0.06; **TE count**, 1.3x enrichment: XGFs μ = 1.79, all other genes μ = 1.41;

Wilcoxon Rank Sum, $W_{2575,101686} = 167,286,786$, p < 0.01). Similarly, we find only a slight,

non-significant, enrichment of LTR TE coverage at XGFs, but we do identify a significant

enrichment of LTR TE count (**TE coverage**, 1.4x enrichment: XGFs μ = 1,817bp, all other

genes μ = 1,238bp; Wilcoxon Rank Sum, $W_{1004,51486} = 25,656,118$, p = 0.69; **TE count**, 1.7x

enrichment: XGFs μ = 2.54, all other genes μ = 1.51; Wilcoxon Rank Sum, $W_{1004,51486} =$

$32,328,754$, p < 0.01). Only PLEs show a depletion, albeit insignificant, in TE coverage

around XGFs, but they also show significant enrichment in count around XGFs (**TE**

**coverage**, 0.7x enrichment: XGF μ = 456bp, all other genes μ = 608bp; $W_{47,6823} = 185,654$, p

= 0.06; **TE count**, 1.3x enrichment: XGF μ = 1.43, all other genes μ = 1.11; Wilcoxon Rank

Sum, $W_{47,6823}$ = 174,958, p < 0.05). However, like SINEs, PLEs account for a very small

proportion of total TE content (Supplemental Table S1).

## *XGF enrichment varies among aphid genomes*

We find no significant difference in TE enrichment among different XGF types (i.e CYP

genes, esterases, UGTs, GSTs, ABCs) (Kruskal-Wallis, TE coverage: $\chi^2_4$ = 7.72, p = 0.10;

TE count: $\chi^2_4$ = 4.25, p = 0.37, Figure 3). However, we do find significant variation in the

magnitude of TE enrichment at XGFs among aphid species (linear regression, TE coverage:

$F_{19,78}$ = 37.49, p < 0.01, adjusted $R^2$ = 0.88, TE count: $F_{19,78}$ = 28.28, p < 0.01, adjusted $R^2$ =

0.84). Specifically, in terms of TE coverage, *C. cedri* shows the highest levels of enrichment

across all XGFs, with a mean enrichment of 15.9x compared to non-XGFs (Figure 5,

Supplemental Table S3). This is considerably higher than the lowest level of mean TE

coverage enrichment, which was found in *M. sacchari* with 1.9x TE coverage enrichment at

XGFs compared to non-XGFs (Figure 5, Supplemental Table S3). In terms of TE count, the

highest level of enrichment at XGFs compared to non-XGFs was also found in *C. cedri*, with

a mean TE count enrichment of 28.5x the number of TE insertions, whilst the lowest level

was found in *A. craccivora*, with a mean enrichment of 4.6x the number of TE insertions at

XGFs compared to non-XGFs (Figure 5, Supplemental Table S3)

Overall, whilst we observe enrichment in both TE coverage and TE count at XGFs

compared to non-XGFs in aphid genomes, fold-change increases are larger for TE count.

High enrichment for TE count combined with lower increases in TE coverage suggest that

enrichment is driven by the presence of large numbers of TE fragments at XGFs, as

opposed to full-length TEs (Figure 6). Indeed, mean TE length at XGFs across aphids is just

536bp, which is much shorter than most full length TEs, excluding SINEs (Wells and

Feschotte 2020; Wicker et al. 2007). Further, 90.40% of TEs across XGFs are below

1,000bp in length (Figure 6). Therefore, it is likely that a general pattern in TE accumulation

at XGFs is TE truncation with retention of specific TE domains, as previously observed for

certain individual cases of TE co-option during the evolution of host insecticide resistance

(Singh et al. 2020; Remnant et al. 2013; Panini et al. 2021; Bogwitz et al. 2005; Itokawa et

al. 2011; Schmidt et al. 2010; Darboux et al. 2007; Gahan et al. 2001; Yang et al. 2007;

Zhao et al. 2010).

335
336

*TEs are enriched at XGFs associated with the detoxification of synthetic*

*insecticides in certain aphid species*

Cytochrome P450 genes (CYP genes) are a highly important gene family for the

detoxification of xenobiotics, including insecticides (Katsavou et al. 2022; Feyereisen 2005).

Indeed, increases in the expression of certain CYP genes can provide increases in

insecticide resistance (Nauen et al. 2022). Additionally, several cases of TE interactions with

CYP genes have been characterised (Chung et al. 2007; Chen and Li 2007; Daborn et al.

2002; Itokawa et al. 2011; Schmidt et al. 2010; Karunker et al. 2008; Nauen et al. 2022).

Therefore, we undertook a targeted analysis to examine evidence for the accumulation of

TEs at CYP genes with putative functions in xenobiotic resistance. For this, aphid CYP

genes were labelled based on the availability of evidence to support their xenobiotic

resistance functions from (Katsavou et al. 2022), as some CYP gene family members play

other roles, such as hormone breakdown. This resulted in the assignment of 2-29 CYP

genes per aphid genome to the xenobiotic detoxification category (Supplemental Table S5).

In *S. flava*, *C. cedri*, and *M. cerasi*, we find significant enrichment of TEs surrounding CYP

genes in CYP gene (sub)families frequently implicated in the breakdown of xenobiotics

compared to those with other functions (Figure 7A). However, we find the opposite pattern in

*S. graminum*, *M. dirhodum*, and *M. persicae*, where overall there is a significant depletion of

TEs surrounding CYP genes associated with xenobiotic resistance (Figure 7A).

Next, we considered patterns of TE accumulation at CYP genes specifically involved

in resistance to synthetic insecticides, rather than resistance to xenobiotics more generally.

For this, CYP genes associated with xenobiotic resistance functions were labelled to reflect

the compounds to which they confer resistance. CYP genes that confer resistance to

insecticides were then considered separately from CYP genes associated with the

detoxification of other xenobiotics, such as those offering resistance to naturally occurring

plant allelochemicals. We find a significant enrichment of TEs around CYP genes associated

with resistance to synthetic insecticides in *P. nigronervosa*, *A. pisum*, *M. cerasi*, and *M.*

*persicae* (Figure 7B). With the exception of *P. nigronervosa*, these are all species of "*most*

*agricultural importance*" (Van Emden and Harrington 2017), and are therefore likely to have

experienced particularly high exposure to synthetic insecticides, and extremely strong

selection to evolve resistance. The significant enrichment of TEs at XGFs associated with

resistance to synthetic insecticides in these species is thus consistent with their selective

recruitment for resistance evolution. However, no significant enrichment of TEs around CYP

genes associated with resistance to synthetic insecticides was found in 8 other species of

"*most agricultural importance*" included in our analysis: *A. craccivora, A. gossypii, D. noxia*,

*M. dirhodum, R. padi, S. graminum,* and *S. avenae* (Van Emden and Harrington 2017).

*Evidence for the selective enrichment of TEs at XGFs*

We hypothesised that TE enrichment at resistance loci could arise through two

distinct processes: (i) Chromatin availability, whereby XGFs are surrounded by open

chromatin due to frequent transcriptional activity or constitutive expression, and so represent

locations that are more accessible for transposition; (ii) Selective retention of beneficial TE

insertions. Given that XGFs are essential for the detoxification of harmful plant metabolites

and insecticides, and so are of key importance for survival and reproduction, TE insertions

that modify XGFs in beneficial ways may be selectively retained and so spread through host

populations. Consequently, the combined effects of XGFs as selective hotspots (especially

considering the extremely strong selection mediated by intensive insecticide regimes), and

the demonstrated evolutionary potential of TEs, could result in the selective retention of TE

insertions surrounding XGFs over time. Such effects may be magnified by the influence of

host stress under insecticide exposure, which has been implicated in increases in TE activity

(Chénais et al. 2012; Stapley et al. 2015; Horváth et al. 2017). Specifically, increased TE

activity may result in a higher likelihood of TE insertions occurring at any genomic location,

including XGFs. However, TE insertions at XGFs may be more likely to be selectively

retained, if they offer host benefits in the form of resistance mutations, due to their capacity

to contribute to host evolvability through diverse genetic mechanisms (Bourque et al. 2018;

Rebollo et al. 2012).

Only germline TE activity, as opposed to somatic TE activity, is relevant from the

perspective of TE accumulation at XGFs, since only TE insertions that occur in the germline

have the potential to be transmitted to the next generation (novel TE insertions in the

genomes of somatic cells are an evolutionary dead-end). TE density at a gene, defined as

the proportion of its sequence occupied by TEs, is positively correlated with germline gene

expression, suggesting that TEs preferentially insert into regions that are actively transcribed

in the germline (McVicker and Green 2010). Therefore, a key question to distinguish

between alternative hypotheses that explain TE enrichment at XGFs is, to what extent are

XGFs expressed in the germline? We addressed this question using RNA-seq data from a

recent study on *A. pisum* (Jaquiéry et al. 2022) to examine gene expression in testes and

ovaries for XGFs and a large panel of housekeeping genes. Housekeeping genes are

constitutively expressed in all cell types and so represent a 'maximally accessible' case with

which to examine TE enrichment (Butte et al. 2001) (Supplemental Table S4). We find that

germline housekeeping gene expression is significantly higher compared to XGFs

considering mean RPKM (reads per kilobase of transcript per million base pairs sequenced)

in *A. pisum* (Wilcoxon Rank Sum, $W_{612,112} = 52252$, p < 0.01, Figure 8C). The observed

pattern of expression of housekeeping genes and XGFs is also present in *Drosophila*

*melanogaster*, suggesting conservation across insect diversity (Supplemental Figure S4).

Consistent with chromatin availability as an explanation for TE enrichment, we find that TEs

are significantly enriched at housekeeping genes compared to all other host genes,

excluding XGFs (TE coverage: Wilcoxon Rank Sum, $W_{546,475492} = 218,813,232$, p < 0.01; TE

count: Wilcoxon Rank Sum, $W_{546,475492} = 170,683,160$, p < 0.01), to a similar level as that

observed for XGFs versus all other host genes (Figure 8A-B, Supplemental Figure S5).

However, given that germline expression of XGFs is consistently low across all XGF types

(Figure 8C), we conclude that germline expression cannot explain the observed enrichment

of TEs at XGFs. Instead, our findings support the alternative explanation for the enrichment

of TEs at XGFs, which is selective retention of TE sequences due to genetic contributions

towards the evolution of xenobiotic resistance mutations. There is no mechanistic basis to

suggest that TEs specifically target the regions surrounding XGFs as insertion sites,

compared to other genomic locations. Instead, under this model TE insertions at XGFs are

more likely to be retained and spread through the host population (compared to TE

insertions at other loci), due to the combination of their capacity to contribute to host

evolvability, and the extremely strong selection pressure imposed by insecticide treatment.

## 427 Discussion

Resistance evolution remains a major societal challenge that greatly affects efforts to control

harmful pests and pathogens (Van Emden and Harrington 2017). To develop new

approaches to overcome the challenge of resistance evolution, a deeper understanding of

the genomic mechanisms leading to the repeated emergence of resistant phenotypes is

required. Whilst isolated examples of TE involvement in the evolution of xenobiotic

resistance exist, the extent to which TEs act as a source of genomic novelty for resistance

evolution, and host evolution more generally, remains an open question.

We examine whether genes implicated in xenobiotic resistance are enriched for TE

insertions, potentially due to the generation of novel advantageous mutations for resistance

evolution. We find an enrichment of TEs, both in terms of TE sequence coverage and

number of TE insertions, at XGFs across all aphids and all XGF types. Further, we provide

evidence that this enrichment is the consequence of selection at XGFs, as opposed to

elevated insertion rates due to chromatin availability at XGFs in the germline.

Our results also uncover considerable variability in the enrichment of TEs around

XGFs both within and among aphid species and XGF types. This demonstrates that TEs are

not globally enriched at every XGF, but instead are enriched only at certain XGFs and

depleted at others. Consequently, the patterns we observe are driven by hotspots of TE

enrichment around certain XGF types, and in some cases individual XGFs, which show

extremely high levels of enrichment compared to other XGFs. For example, considering all

aphid species examined, we identify significant enrichment of TEs around GSTs. However,

this result is driven by a subset of aphids, with 11 of the 20 species showing no evidence of

significant enrichment around this XGF class (Supplemental Figure S4). This is consistent

with varying levels of selection acting on particular detoxification genes and particular aphid

species. Meanwhile, we find markedly more TE hotspots at UGTs and CYP genes, whereby

certain XGFs are massively enriched for TE fragments. These observations demonstrate the

nuances involved in patterns of TE accumulation at XGFs and underline a need for

functional validation analyses to test the impact of individual TE insertions on XGF

expression, and whether highly TE-enriched loci are indeed involved in xenobiotic

resistance.

Important crop pests are likely to experience much higher levels of insecticide stress

than non-target species, demonstrated by the increased number and diversity of insecticide

resistance mutations in target species, compared to non-target species (Mota-Sanchez and

Wise 2023). Consequently, variability in TE enrichment is expected among aphid species

and XGFs, depending on variation in the degree of insecticide exposure. This prediction is

confirmed when considering CYP genes, for which data clarifying the relationship to specific

insecticides is available (rather than just to xenobiotics in general). Specifically, we find that

TE enrichment is significantly higher around CYP genes that are implicated in the

detoxification of synthetic insecticides compared to those involved in the detoxification of

natural xenobiotics, but only in three out of ten aphids of "*major agricultural importance*" (Van

Emden and Harrington 2017) included in our study (*A. pisum, M. cerasi*, and *M. persicae*).

Consequently, either CYP genes are less important for survival in the other species (perhaps

due to differences in pest control strategies, such as a greater reliance on forms of non-

insecticide-based control, such as biocontrol strategies), or there are genomic mechanisms

that limit the co-option of TEs for resistance evolution in these species.

Demonstrated contributions of TEs to xenobiotic resistance often involve small

fragments of the original TE insertions, and contain motifs that act as host regulatory

elements (Singh et al. 2020; Remnant et al. 2013; Panini et al. 2021; Bogwitz et al. 2005;

Itokawa et al. 2011; Schmidt et al. 2010; Darboux et al. 2007; Gahan et al. 2001; Yang et al.

2007). Meanwhile, the emergence of a resistant phenotype frequently appears to result from

a single, or up to a few, TE insertions at each specific XGF locus, rather than large

accumulations of TE DNA (Schmidt et al. 2010; Daborn et al. 2002; Chung et al. 2007;

Panini et al. 2021). This presumably occurs due to genomic and selective processes that act

to retain the core TE regions that convey a selective benefit, while purging long repetitive

and potentially deleterious regions (e.g. that promote non-homologous recombination with

similar repetitive elements). Meanwhile, the impacts of these TE insertions can occur via

complex multistep processes including the retention of several truncated TE fragments (such

as for *CYP6G1*, Schmidt *et al.* (2010)). Thus, a key consideration is that TE-associated

resistance mechanisms go beyond simple enrichment at specific loci.

We find that DNA TEs are highly enriched at XGFs, despite the more widely

acknowledged role of TEs with more complex *cis*-regulatory elements, such as LTR

retrotransposons, in altering host gene expression (Chung et al. 2007). This may suggest

that other processes beyond TE-mediated increases in expression are involved in the

patterns of enrichment we uncover. Although, it is entirely possible that the DNA TEs

involved also contain sequences relevant for the recruitment of transcription factors that

increase expression. Meanwhile, at present we cannot currently explain the repeated cases

whereby certain XGFs are massively enriched in terms of TE coverage and TE count

compared to other XGFs. Thus, unravelling the influence of TEs at these loci represents a

key target for future research.

Gaining a deeper understanding of the mechanisms that drive patterns of association

between TEs and XGFs will require contributions from complementary approaches to the

broadscale comparative genomic analyses performed here. Of particular relevance are

functional validation studies to examine the influence of individual TE insertions on nearby

XGF loci, which our results provide considerable scope to address. Further analyses are

also required to elucidate the underlying mechanisms by which TE sequences may

contribute to resistance phenotypes, such as screening to investigate if implicated TE

sequences contain transcription factor binding sites, as well as information on chromatin

conformation and the availability of DNA within TE sequences at XGFs. Meanwhile, detailed

intra-species analyses can provide valuable evolutionary insights into individual TE

insertions of interest (e.g. Aminetzach et al. (2005)), and information on TE occupancy within

populations (e.g. Barrón et al. (2014)). Future work considering strains displaying different

resistance phenotypes using high-quality long-read datasets and individual TE annotations

(e.g Rech et al. (2022)) will enable the characterisation of TE variants likely contributing to

resistance evolution in aphids. Collectively, such approaches will ultimately allow evaluation

of the extent to which TEs are utilised in resistance mutations, versus the contributions

arising from other classes of genetic variation such as SNPs and CNVs.

# Materials and Methods

## *Estimating Host Phylogeny*

The genome assemblies of 20 aphid species, and 7 diverse hemipteran outgroup species

(*Ericerus pela*, *Maconellicoccus hirsutus*, *Phenacoccus solenopsis, Bemisa tabaci, Sogatella*

*furcifera, Diaphorina citri, Pachypsylla venusta*), were obtained along with gene annotations

where available (Supplemental Table S6). To estimate host phylogeny, a supermatrix

approach was used. Specifically, for each genome assembly, BUSCO (version 5.2.2) (Manni

et al. 2021a, 2021b) was used with the *hemiptera_odb10* geneset to identify conserved gene

orthologs in each genome assembly. The identifiers for all complete genes were extracted

and those present in less than three genomes were removed. The amino acid sequences for

each gene from all species were extracted and saved in individual FASTA files. For each

gene, sequences were aligned using MAFFT (version 7.453) (Katoh and Standley 2013) (--

auto). The subsequent alignments were concatenated using Phykit create_concat (Steenwyk

et al. 2021) to generate a supermatrix. To generate the species phylogeny, we performed

1,000 ultrafast bootstrap repetitions (-bb 1000) in IQ-TREE (version 1.6.12) (Nguyen et al.

2015), using the best fit amino acid model identified by Modelfinder (Kalyaanamoorthy et al.

2017) (Supplemental File S1).

## *Transposable Element Annotation*

To curate and annotate TEs, each genome assembly was annotated with the Earl Grey TE

annotation pipeline (version 1.3) (https://github.com/TobyBaril/EarlGrey) (Baril et al. 2022).

Briefly, known TEs from Arthropoda (-r arthropoda) were first annotated using both the Dfam

3.4 (Hubley et al. 2016) and Repbase RepeatMasker Edition (version 20181026) (Jurka et

al. 2005; Kapitonov and Jurka 2008) TE databases. Following this, Earl Grey identified *de*

*novo* TE families and refined these using an iterative 'BLAST, Extract, Extend' process (Platt

et al. 2016). Following final annotation with the combined library of known and *de novo* TE

families, annotations are processed to remove overlapping annotations and to defragment

annotations likely to originate from the same TE insertion. TE annotation GFF files are

provided in Supplemental File S2.

To characterise TEs as either shared or species-specific, *de novo* TE libraries from

each genome assembly were clustered to the TE family definition of Wicker et al. (2007),

implemented as described by Goubert et al. (2022) using cd-hit-est (-d 0 -aS 0.8 -c 0.8 -G 0 -

g 1 -b 500 -r 1) (Li and Godzik 2006; Fu et al. 2012). Sequences within each cluster were

designated a number between 0 and 20 to determine the number of other species that the

TE family was shared with, with 0 identifying TEs unique to a species, and 19 identifying TEs

shared across all 20 aphid species considered in this study. For known TE families from

Dfam and Repbase, each TE was labelled using the same system based on the number of

species the TE family was identified in.

Genetic distance from TE consensus sequence is used as a proxy for estimated

timing of TE activity (i.e TE age). Using genetic distance from TE consensus as a proxy for

insertion time should be taken with caution, as this can be influenced by the methodology

used to generate TE consensus sequences, and whether known TE sequences from other

organisms are used as a reference or purely *de novo* TE consensus sequences from the

species of interest. However, this metric can still be used to provide a rough relative estimate

of TE activity.

*Phylogenetic Heritability of TE Abundance and Diversity*

To determine the phylogenetic heritability of TE abundance, expressed as total TE count,

and TE diversity, expressed as total number of distinct TE families, we measured the

amount of variation explained by phylogenetic relationships using Bayesian phylogenetic

mixed models with Markov chain Monte Carlo estimations, with an intercept fitted as a fixed

effect and the phylogeny fitted as a random effect, in the MCMCglmm package (Hadfield

2010) in R (version 4.2.1) (R Core Team 2023) using the Rstudio IDE (RStudio Team 2015;

Racine 2013). A poisson error distribution and log link function were used, and inverse

gamma priors were specified for all R and G-side random effects (V = 1, v = 0.002). Models

were run for 11,000,000 iterations with a burn-in of 1,000,000 and a thinning interval of

1,000. This approach generated 10,000 posterior samples from which posterior mode and

95% CIs were calculated. The proportion of between-species variation explained by

phylogeny was calculated from the model using the equation $V_p / (V_p + V_s)$ where $V_p$ and $V_s$

represent the phylogenetic and species-specific components of between-species variance

(Freckleton et al. 2002), which is equivalent to phylogenetic heritability.

*Gene Predictions and Transposable Element Association with Genomic*

*Compartments*

For genome assemblies lacking a gene annotation GFF file at the time of analysis (*A.*

*gossypii*, *H. cornu, M. dirhodum, P. nigronervosa, R. padi, S. graminum, S. chinensis, S.*

*avenae, S. miscanthi*), genes were annotated using AUGUSTUS (version 3.3.3) (Stanke et

al. 2008; Keller et al. 2011) with the pea aphid reference gene set (--species=pea_aphid --

strand=both --genemodel=partial --introns=on --exonnames=on --gff3=on).

To prepare genome annotation files for intersection with TE loci, each gene

annotation GFF was processed to generate a GFF file containing coordinates of exons,

introns, 5' and 3' flanking regions, and intergenic space. Gene flanking regions are defined

here as 20kb directly upstream or downstream of the gene body. These flanking regions are

determined to identify TEs at a distance that could be described as the proximate promoter

region, rather than just accounting for the core promoter region, to include both promoter

and more distal enhancer regions for genes (Lenhard et al. 2012). Intergenic space

coordinates were generated using BEDTools complement (Quinlan and Hall 2010), to

generate inverse coordinates to those of genic compartments and flanking regions. To

quantify TE occupation in different genomic compartments (introns, exons, 5' and 3' flanking

regions, and intergenic space), BEDTools intersect was used to calculate overlap (-wao)

between all compartments and TEs.

*Identification of Xenobiotic Gene Family Loci and Housekeeping Genes*

The gene identities for all ABC transporters, cytochrome P450s, UDP-

glucuronosyltransferases, esterases, and glutathione S-transferases were manually curated

from the functional annotation of the *M. persicae* G006v2 genome (Singh et al. 2021). Using

these identifiers, exon coordinates were obtained from the *M. persicae* gene annotation

GFF3 file, and corresponding nucleotide sequences were extracted from the genome

assembly. The resulting exon sequences from *M. persicae* were used as queries to identify

XGFs in all other aphid assemblies using BLASTN (Camacho et al. 2009) with an e-value

threshold of $1\times10^{-10}$. BLAST hits for each species were manually inspected to extract those

that best represented each XGF, with the inclusion of gene duplications where appropriate.

Following this, a BED file of XGF coordinates for each species was generated including the

annotation of flanking regions as described above (Supplemental File S3).

Housekeeping genes (HKGs) are required for basic cellular functions and are

typically expressed in all cell types (Butte et al. 2001). Amino acid sequences of 28 HKGs

characterised for aphids (Bansal et al. 2012; Koramutla et al. 2016; Li et al. 2020; Kang et al.

2017; Yang et al. 2015, 2014) were obtained from GenBank (Benson et al. 2013) for *M.*

*persicae* (Supplemental File S4). Subsequently, a BLASTX search was performed, with

default parameters, against each aphid genome to identify HKG loci in the other 19 species.

For each species, HKGs were manually curated and a BED file of HKG coordinates,

including the annotation of flanking regions, was generated (Supplemental File S5). To

quantify TE occupation around XGFs and HKGs, BEDTools intersect was used to calculate

overlap (-wao) between annotated TEs and XGFs.

## Competing Interests

620    The authors declare no competing interests.

## Acknowledgements

TB was supported by a studentship from the Biotechnology and Biological Sciences

Research Council-funded South West Biosciences Doctoral Training Partnership

(grant number BB/M009122/1).

AP was supported by the European Research Council (ERC) under the European

Union's Horizon 2020 research and innovation program (grant agreement number

646625).

CB was supported by a Biotechnology and Biological Sciences Research Council

(BBSRC) grant (grant number BB/S006060/1) and the European Research Council

(ERC) under the European Union's Horizon 2020 research and innovation program

(grant agreement number 646625)

AH was supported by a Biotechnology and Biological Sciences Research Council

(BBSRC) David Phillips Fellowship (grant number BB/N020146/1).

*Myzus cerasi* DNA sequencing data were downloaded from AphidBase. Funding for

the sequencing was provided by ERC Starting Grant APHIDHOST-310190 awarded

to Jorunn Bos at the James Hutton Institute, United Kingdom.

For the purpose of open access, the author has applied a 'Creative Commons

Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

## 646  Author Contributions

647  TB performed TE annotation and downstream analyses, produced the figures, and

648  drafted the manuscript. AP manually curated xenobiotic resistance genes from *M.*

649  *persicae.* CB and AH contributed to the analytical design and data analysis, and

650  participated in writing the manuscript. AH conceived and coordinated the study. All

651  authors read and approved the final manuscript.

# REFERENCES

Ahn S-J, Vogel H, Heckel DG. 2012. Comparative analysis of the UDP-glycosyltransferase multigene family in insects. *Insect Biochem Mol Biol* **42**: 133–147.

Aminetzach YT, Macpherson JM, Petrov DA. 2005. Pesticide resistance via transposition-mediated adaptive gene truncation in Drosophila. *Science* **309**: 764–767.

Arkhipova IR. 2018. Neutral theory, transposable elements, and eukaryotic genome evolution. *Mol Biol Evol* **35**: 1332–1337.

Bansal R, Mamidala P, Mian MAR, Mittapalli O, Michel AP. 2012. Validation of reference genes for gene expression studies in Aphis glycines (Hemiptera: Aphididae). *J Econ Entomol* **105**: 1432–1438.

Baril T, Hayward A. 2022. Migrators within migrators: exploring transposable element dynamics in the monarch butterfly, Danaus plexippus. *Mob DNA* **13**: 5.

Baril T, Imrie RM, Hayward A. 2022. Earl Grey: a fully automated user-friendly transposable element annotation and analysis pipeline. https://www.researchsquare.com/article/rs-1812599/latest.pdf.

Barrón MG, Fiston-Lavier A-S, Petrov DA, González J. 2014. Population genomics of transposable elements in Drosophila. *Annu Rev Genet* **48**: 561–581.

Bass C, Puinean AM, Nauen R, Gutbrod O, Denholm I, Williamson MS, Slater R, Field LM, Zimmer CT, Foster SP. 2014. The evolution of insecticide resistance in the peach potato aphid, Myzus persicae. *Insect Biochem Mol Biol* **51**: 41–51.

Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2013. GenBank. *Nucleic Acids Res* **41**: D36-42.

Blass E, Bell M, Boissinot S. 2012. Accumulation and rapid decay of non-LTR retrotransposons in the genome of the three-spine stickleback. *Genome Biol Evol* **4**: 687–702.

Bogwitz MR, Chung H, Magoc L, Rigby S, Wong W, O'Keefe M, McKenzie JA, Batterham P, Daborn PJ. 2005. Cyp12a4 confers lufenuron resistance in a natural population of Drosophila melanogaster. *Proc Natl Acad Sci U S A* **102**: 12807–12812.

Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault M, Izsvák Z, Levin HL, Macfarlan TS, et al. 2018. Ten things you should know about transposable elements. *Genome Biol* **19**: 199.

Bradshaw CJA, Leroy B, Bellard C, Roiz D, Albert C, Fournier A, Barbet-Massin M, Salles J-M, Simard F, Courchamp F. 2016. Massive yet grossly underestimated global costs of invasive insects. *Nat Commun* **7**: 12986.

Butte AJ, Dzau VJ, Glueck SB. 2001. Further defining housekeeping, or "maintenance," genes Focus on "A compendium of gene expression in normal human tissues." *Physiol Genomics* **7**: 95–96.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: Architecture and applications. *BMC Bioinformatics* **10**: 1–9.

Catania F, Kauer MO, Daborn PJ, Yen JL, Ffrench-Constant RH, Schlötterer C. 2004.
World-wide survey of an Accord insertion and its association with DDT resistance in
Drosophila melanogaster. *Mol Ecol* **13**: 2491–2504.

Chen S, Li X. 2007. Transposable elements are enriched within or in close proximity to
xenobiotic-metabolizing cytochrome P450 genes. *BMC Evol Biol* **7**: 46.

Chénais B, Caruso A, Hiard S, Casse N. 2012. The impact of transposable elements on
eukaryotic genomes: From genome size increase to genetic adaptation to stressful
environments. *Gene* **509**: 7–15.

Chung H, Bogwitz MR, McCart C, Andrianopoulos A, Ffrench-Constant RH, Batterham P,
Daborn PJ. 2007. Cis-regulatory elements in the accord retrotransposon result in
tissue-specific expression of the Drosophila melanogaster insecticide resistance
gene Cyp6g1. *Genetics* **175**: 1071–1077.

Cosby RL, Chang N-C, Feschotte C. 2019. Host–transposon interactions: conflict,
cooperation, and cooption. *Genes Dev* **33**: 1098–1116.

Daborn PJ, Yen JL, Bogwitz MR, Le Goff G, Feil E, Jeffers S, Tijet N, Perry T, Heckel D,
Batterham P, et al. 2002. A single p450 allele associated with insecticide resistance
in Drosophila. *Science* **297**: 2253–2256.

Darboux I, Charles J-F, Pauchet Y, Warot S, Pauron D. 2007. Transposon-mediated
resistance to Bacillus sphaericus in a field-evolved population of Culex pipiens
(Diptera: Culicidae). *Cell Microbiol* **9**: 2022–2029.

de Koning APJ, Gu W, Castoe TA, Batzer MA, Pollock DD. 2011. Repetitive elements may
comprise over Two-Thirds of the human genome. *PLoS Genet* **7**.
http://dx.doi.org/10.1371/journal.pgen.1002384.

de Little SC, Edwards O, van Rooyen AR, Weeks A, Umina PA. 2017. Discovery of
metabolic resistance to neonicotinoids in green peach aphids (Myzus persicae) in
Australia. *Pest Manag Sci* **73**: 1611–1617.

Dermauw W, Van Leeuwen T. 2014. The ABC gene family in arthropods: Comparative
genomics and role in insecticide transport and resistance. *Insect Biochem Mol Biol*
**45**: 89–110.

Deutsch CA, Tewksbury JJ, Tigchelaar M, Battisti DS, Merrill SC, Huey RB, Naylor RL.
2018. Increase in crop losses to insect pests in a warming climate. *Science* **361**:
916–919.

Fabrick JA, Mathew LG, Tabashnik BE, Li X. 2011. Insertion of an intact CR1
retrotransposon in a cadherin gene linked with Bt resistance in the pink bollworm,
Pectinophora gossypiella. *Insect Molecular Biology* **20**: 651–665.
http://dx.doi.org/10.1111/j.1365-2583.2011.01095.x.

Feyereisen R. 2005. Insect cytochrome P450. *Comprehensive molecular insect science*.

Feyereisen R. 1999. Insect P450 enzymes. *Annu Rev Entomol* **44**: 507–533.

Freckleton RP, Harvey PH, Pagel M. 2002. Phylogenetic analysis and comparative data: a
test and review of evidence. *Am Nat* **160**: 712–726.

Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**: 3150–3152.

Gahan LJ, Gould F, Heckel DG. 2001. Identification of a gene associated with Bt resistance
in Heliothis virescens. *Science* **293**: 857–860.

Gilbert C, Peccoud J, Cordaux R. 2021. Transposable Elements and the Evolution of
Insects. *Annu Rev Entomol* **66**: 355–372.

Gilbert LI, Gill SS, eds. 2014. *Insect pharmacology*. Academic Press.

Gott RC, Kunkel GR, Zobel ES, Lovett BR, Hawthorne DJ. 2017. Implicating ABC
Transporters in Insecticide Resistance: Research Strategies and a Decision
Framework. *J Econ Entomol* **110**: 667–677.

Goubert C, Craig RJ, Bilat AF, Peona V, Vogan AA, Protasio AV. 2022. Correction: A
beginner's guide to manual curation of transposable elements. *Mob DNA* **13**: 15.

Gregory TR. 2005. Synergy between sequence and size in large-scale genomics. *Nat Rev
Genet* **6**: 699–708.

Guengerich FP. 2009. Cataloging the Repertoire of Nature's Blowtorch, P450. *Chem Biol* **16**:
1215–1216.

Hadfield JD. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The MCMCglmm R Package. *J Stat Softw* **33**: 1–22.

Horváth V, Merenciano M, González J. 2017. Revisiting the Relationship between
Transposable Elements and the Eukaryotic Stress Response. *Trends Genet* **33**:
832–841.

Hubley R, Finn RD, Clements J, Eddy SR, Jones TA, Bao W, Smit AFA, Wheeler TJ. 2016.
The Dfam database of repetitive DNA families. *Nucleic Acids Res* **44**: D81–D89.

Itokawa K, Komagata O, Kasai S, Masada M, Tomita T. 2011. Cis-acting mutation and
duplication: History of molecular evolution in a P450 haplotype responsible for
insecticide resistance in Culex quinquefasciatus. *Insect Biochem Mol Biol* **41**: 503–
512.

Jaquiéry J, Simon J-C, Robin S, Richard G, Peccoud J, Boulain H, Legeai F, Tanguy S,
Prunier-Leterme N, Letrionnaire G. 2022. Masculinization of the X-chromosome in
aphid soma and gonads. *Peer Community J* **2**.
http://dx.doi.org/10.24072/pcjournal.166.

Jurka J, Kapitonov VV, Pavlicek A, Klonowski P, Kohany O, Walichiewicz J. 2005. Repbase
Update, a database of eukaryotic repetitive elements. *Cytogenet Genome Res* **110**:
462–467.

Kalyaanamoorthy S, Minh BQ, Wong TKF, Haeseler A von, Jermiin LS. 2017. ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nature Methods 2017 14:6*
**14**: 587–589.

Kang Z-W, Liu F-H, Tian H-G, Zhang M, Guo S-S, Liu T-X. 2017. Evaluation of the reference
genes for expression analysis using quantitative real-time polymerase chain reaction
in the green peach aphid, Myzus persicae. *Insect Sci* **24**: 222–234.

Kapitonov VV, Jurka J. 2008. A universal classification of eukaryotic transposable elements
implemented in Repbase. *Nat Rev Genet* **9**: 411–412.

Karunker I, Benting J, Lueke B, Ponge T, Nauen R, Roditakis E, Vontas J, Gorman K,

Denholm I, Morin S. 2008. Over-expression of cytochrome P450 CYP6CM1 is
associated with high resistance to imidacloprid in the B and Q biotypes of Bemisia
tabaci (Hemiptera: Aleyrodidae). *Insect Biochem Mol Biol* **38**: 634–644.

Kasai S, Scott JG. 2001. Cytochrome P450s CYP6D3 and CYP6D1 are part of a P450 gene
cluster on autosome 1 in the house fly. *Insect Mol Biol* **10**: 191–196.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Mol Biol Evol* **30**: 772–780.

Katsavou E, Riga M, Ioannidis P, King R, Zimmer CT, Vontas J. 2022. Functionally
characterized arthropod pest and pollinator cytochrome P450s associated with
xenobiotic metabolism. *Pesticide Biochemistry and Physiology* **181**: 105005.
http://dx.doi.org/10.1016/j.pestbp.2021.105005.

Keller O, Kollmar M, Stanke M, Waack S. 2011. A novel hybrid gene prediction method
employing protein multiple sequence alignments. *Bioinformatics* **27**: 757–763.

Kennedy CJ, Tierney KB. 2013. Xenobiotic Protection/Resistance Mechanisms in
Organisms. In *Environmental Toxicology: Selected Entries from the Encyclopedia of
Sustainability Science and Technology* (ed. E.A. Laws), pp. 689–721, Springer New
York, New York, NY.

Ketterman AJ, Saisawang C, Wongsantichon J. 2011. Insect glutathione transferases. *Drug
Metab Rev* **43**: 253–265.

Kidwell MG. 2002. Transposable elements and the evolution of genome size in eukaryotes.
*Genetica* **115**: 49–63.

Kliot A, Ghanim M. 2012. Fitness costs associated with insecticide resistance. *Pest Manag
Sci* **68**: 1431–1437.

Koramutla MK, Aminedi R, Bhattacharya R. 2016. Comprehensive evaluation of candidate
reference genes for qRT-PCR studies of gene expression in mustard aphid, Lipaphis
erysimi (Kalt). *Sci Rep* **6**: 25883.

Lavoie CA, Ii RNP, Novick PA, Counterman BA, Ray DA. 2013. Transposable element
evolution in Heliconius suggests genome diversity within Lepidoptera. *Mob DNA* **21**:
1–10.

Lenhard B, Sandelin A, Carninci P. 2012. Metazoan promoters: emerging characteristics and
insights into transcriptional regulation. *Nat Rev Genet* **13**: 233–245.

Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**: 1658–1659.

Li X, Gong P, Wang B, Wang C, Li M, Zhang Y, Li X, Gao H, Ju J, Zhu X. 2020. Selection
and validation of experimental condition-specific reference genes for qRT-PCR in
Metopolophium dirhodum (Walker) (Hemiptera: Aphididae). *Scientific Reports* **10**.
http://dx.doi.org/10.1038/s41598-020-78974-z.

Li X, Schuler MA, Berenbaum MR. 2007. Molecular mechanisms of metabolic resistance to
synthetic and natural xenobiotics. *Annu Rev Entomol* **52**: 231–253.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021a. BUSCO Update: Novel
and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage
for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol Biol Evol* **38**: 4647–

817 4654.

818 Manni M, Berkeley MR, Seppey M, Zdobnov EM. 2021b. BUSCO: Assessing Genomic Data
819   Quality and Beyond. *Curr Protoc* **1**: e323.

820 Mateo L, Ullastres A, González J. 2014. A transposable element insertion confers xenobiotic
821   resistance in Drosophila. *PLoS Genet* **10**: e1004560.

822 McVicker G, Green P. 2010. Genomic signatures of germline gene expression. *Genome Res*
823   **20**: 1503–1511.

824 Mota-Sanchez D, Wise JC. 2023. The Arthropod pesticide resistance database.
825   http://www.pesticideresistance.org.

826 Muller HJ. 1964. The relation of recombination to mutational advance. *Mutat Res* **106**: 2–9.

827 Nauen R, Bass C, Feyereisen R, Vontas J. 2022. The Role of Cytochrome P450s in Insect
828   Toxicology and Resistance. *Annu Rev Entomol* **67**: 105–124.

829 Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective
830   Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol*
831   **32**: 268–274.

832 Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze AV, Mikheenko A, Vollger MR, Altemose
833   N, Uralsky L, Gershman A, et al. 2022. The complete sequence of a human genome.
834   *Science* **376**: 44–53.

835 Panini M, Chiesa O, Troczka BJ, Mallott M, Manicardi GC, Cassanelli S, Cominelli F,
836   Hayward A, Mazzoni E, Bass C. 2021. Transposon-mediated insertional mutagenesis
837   unmasks recessive insecticide resistance in the aphid Myzus persicae. *Proc Natl*
838   *Acad Sci U S A* **118**. http://dx.doi.org/10.1073/pnas.2100559118.

839 Peccoud J, Loiseau V, Cordaux R, Gilbert C. 2017. Massive horizontal transfer of
840   transposable elements in insects. *Proceedings of the National Academy of Sciences*
841   **114**: 4721–4726.

842 Platt RN, Blanco-Berdugo L, Ray DA. 2016. Accurate transposable element annotation is
843   vital when analyzing new genome assemblies. *Genome Biol Evol* **8**: 403–410.

844 Puinean AM, Foster SP, Oliphant L, Denholm I, Field LM, Millar NS, Williamson MS, Bass C.
845   2010. Amplification of a cytochrome P450 gene is associated with resistance to
846   neonicotinoid insecticides in the aphid Myzus persicae. *PLoS Genet* **6**: 1–11.

847 Quinlan AR, Hall IM. 2010. BEDTools: A flexible suite of utilities for comparing genomic
848   features. *Bioinformatics* **26**: 841–842.

849 R Core Team. 2023. R: A Language and Environment for Statistical Computing.
850   https://www.R-project.org.

851 Racine JS. 2013. RSTUDIO: A PLATFORM-INDEPENDENT IDE FOR R AND SWEAVE. *J*
852   *Appl Econometrics* **27**: 167–172.

853 Ranson H, Hemingway J. 2005. Mosquito glutathione transferases. *Methods Enzymol* **401**:
854   226–241.

855 Rebollo R, Romanish MT, Mager DL. 2012. Transposable Elements: An Abundant and
856   Natural Source of Regulatory Sequences for Host Genes. *Annu Rev Genet* **46**: 21–

857           42.

858 Rech GE, Radío S, Guirao-Rico S, Aguilera L, Horvath V, Green L, Lindstadt H, Jamilloux V,
859           Quesneville H, González J. 2022. Population-scale long-read sequencing uncovers
860           transposable elements associated with gene expression variation and adaptive
861           signatures in Drosophila. *Nat Commun* **13**: 1948.

862 Remnant EJ, Good RT, Schmidt JM, Lumb C, Robin C, Daborn PJ, Batterham P. 2013.
863           Gene duplication in the major insecticide target site, Rdl, in Drosophila melanogaster.
864           *Proc Natl Acad Sci U S A* **110**: 14705–14710.

865 Rostant WG, Wedell N, Hosken DJ. 2012. Transposable Elements and Insecticide
866           Resistance. *Advances in Genetics* 169–201. http://dx.doi.org/10.1016/b978-0-12-
867           394394-1.00002-x.

868 RStudio Team. 2015. RStudio: integrated development for R. *RStudio, Inc, Boston, MA URL*
869           *http://www rstudio com*.

870 Salces-Ortiz J, Vargas-Chavez C, Guio L, Rech GE, González J. 2020. Transposable
871           elements contribute to the genomic response to insecticides in Drosophila
872           melanogaster. *Philos Trans R Soc Lond B Biol Sci* **375**: 20190341.

873 Schmidt JM, Good RT, Appleton B, Sherrard J, Raymant GC, Bogwitz MR, Martin J, Daborn
874           PJ, Goddard ME, Batterham P, et al. 2010. Copy number variation and transposable
875           elements feature in recent, ongoing adaptation at the Cyp6g1 locus. *PLoS Genet* **6**:
876           e1000998.

877 Singh KS, Cordeiro EMG, Troczka BJ, Pym A, Mackisack J, Mathers TC, Duarte A, Legeai
878           F, Robin S, Bielza P, et al. 2021. Global patterns in genomic diversity underpinning
879           the evolution of insecticide resistance in the aphid crop pest Myzus persicae.
880           *Commun Biol* **4**: 847.

881 Singh KS, Troczka BJ, Duarte A, Balabanidou V, Trissi N, Carabajal Paladino LZ, Nguyen P,
882           Zimmer CT, Papapostolou KM, Randall E, et al. 2020. The genetic architecture of a
883           host shift: An adaptive walk protected an aphid and its endosymbiont from plant
884           chemical defenses. *Sci Adv* **6**: eaba1070.

885 Song WY, Pi LY, Bureau TE, Ronald PC. 1998. Identification and characterization of 14
886           transposon-like elements in the noncoding regions of members of the Xa21 family of
887           disease resistance genes in rice. *Mol Gen Genet* **258**: 449–456.

888 Sparks TC, Storer N, Porter A, Slater R, Nauen R. 2021. Insecticide resistance management
889           and industry: the origins and evolution of the Insecticide Resistance Action
890           Committee (IRAC) and the mode of action classification scheme. *Pest Manag Sci* **77**:
891           2609–2619.

892 Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntenically
893           mapped cDNA alignments to improve de novo gene finding. *Bioinformatics* **24**: 637–
894           644.

895 Stapley J, Santure AW, Dennis SR. 2015. Transposable elements as agents of rapid
896           adaptation may explain the genetic paradox of invasive species. *Mol Ecol* **24**: 2241–
897           2252.

898 Steenwyk JL, Buida TJ, Labella AL, Li Y, Shen X-X, Rokas A. 2021. PhyKIT: a broadly
899           applicable UNIX shell toolkit for processing and analyzing phylogenomic data.

*Bioinformatics.* http://dx.doi.org/10.1093/bioinformatics/btab096.

Sternberg ED, Thomas MB. 2018. Insights from agriculture for the management of
insecticide resistance in disease vectors. *Evol Appl* **11**: 404–414.

Sultana T, Zamborlini A, Cristofari G, Lesage P. 2017. Integration site selection by
retroviruses and transposable elements in eukaryotes. *Nat Rev Genet* **18**: 292–308.

Sundaram V, Cheng Y, Ma Z, Li D, Xing X, Edge P, Snyder MP, Wang T. 2014. Widespread
contribution of transposable elements to the innovation of gene regulatory networks.
*Genome Res* **24**: 1963–1976.

van de Lagemaat LN, Landry J-R, Mager DL, Medstrand P. 2003. Transposable elements in
mammals promote regulatory variation and diversification of genes with specialized
functions. *Trends Genet* **19**: 530–536.

Van Emden HF, Harrington R. 2017. *Aphids as crop pests.* Cabi.

Wang L, Wang J, Ma Y, Wan P, Liu K, Cong S, Xiao Y, Xu D, Wu K, Fabrick JA, et al. 2019.
Transposon insertion causes cadherin mis-splicing and confers resistance to Bt
cotton in pink bollworm from China. *Sci Rep* **9**: 7479.

Weedall GD, Riveron JM, Hearn J, Irving H, Kamdem C, Fouet C, White BJ, Wondji CS.
2020. An Africa-wide genomic evolution of insecticide resistance in the malaria vector
Anopheles funestus involves selective sweeps, copy number variations, gene
conversion and transposons. *PLoS Genet* **16**: e1008822.

Wells JN, Feschotte C. 2020. A Field Guide to Eukaryotic Transposable Elements. *Annu*
*Rev Genet* **54**: 539–561.

Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A, Leroy P,
Morgante M, Panaud O, et al. 2007. A unified classification system for eukaryotic
transposable elements. *Nat Rev Genet* **8**: 973–982.

Worner SP, Gevrey M. 2006. Modelling global insect pest species assemblages to
determine risk of invasion. *J Appl Ecol* **43**: 858–867.

Yang C, Pan H, Liu Y, Zhou X. 2014. Selection of Reference Genes for Expression Analysis
Using Quantitative Real-Time PCR in the Pea Aphid, Acyrthosiphon pisum (Harris)
(Hemiptera, Aphidiae). *PLoS ONE* **9**: e110454.
http://dx.doi.org/10.1371/journal.pone.0110454.

Yang C, Pan H, Liu Y, Zhou X. 2015. Temperature and Development Impacts on
Housekeeping Gene Expression in Cowpea Aphid, Aphis craccivora (Hemiptera:
Aphidiae). *PLoS One* **10**: e0130593.

Yang Y, Chen H, Wu Y, Yang Y, Wu S. 2007. Mutated cadherin alleles from a field
population of Helicoverpa armigera confer resistance to Bacillus thuringiensis toxin
Cry1Ac. *Appl Environ Microbiol* **73**: 6939–6944.

Zhao J, Jin L, Yang Y, Wu Y. 2010. Diverse cadherin mutations conferring resistance to
Bacillus thuringiensis toxin Cry1Ac in Helicoverpa armigera. *Insect Biochem Mol Biol*
**40**: 113–118.

940
941
942

# Figure Legends

Figure 1. Summary of TE content and activity in aphid genomes, plotted in phylogenetic

order. Nodes in the aphid phylogeny with bootstrap support <100 are labelled. Reproductive

mode (Van Emden and Harrington 2017) is labelled next to species names. Genome size is

represented by the black bars. TE content is expressed as percentage of total genome size

for each species, with major TE classifications represented by the colours indicated in the

key. Kimura 2-parameter distance (CpG adjusted) from each TE family consensus is used

as a proxy for relative TE activity, where a lower Kimura distance indicates more recent TE

activity. Due to challenges with accurately estimating TE age, divergence from consensus

better reflects relative TE activity, with lower Kimura distances signalling more recent TE

activity. Activity is organised such that recent TE activity is towards the RHS of the X axis.

Figure 2. Overview of the extent to which TE families are shared among aphid genomes,

and their insertion locations. A. Quantification of shared and unique TE families in aphids.

Main TE classifications are represented by the colours indicated in the key. B. TE genome

compartment occupancy, expressed by TEs as a percentage of total genome size. Gene

flanking regions are defined as 20kb directly upstream or downstream of the gene body.

Figure 3. Differences in TE coverage and TE count at XGFs compared to all other host

genes. Coverage and count of genic (exon and intron) XGF sequence plus flanking regions

(20kb directly upstream or downstream of the gene body) is shown. Each point represents a

single XGF locus. Black lines indicate the expected TE coverage difference whereby XGFs

and other genes are equally enriched for TEs (i.e. 0 enrichment). Coloured lines indicate

species mean coverage and count for each XGF locus and TE type. Major TE classifications

are indicated by the colours in the key.

Figure 4. Karyoplots illustrating the most TE-dense XGF loci, corresponding to the XGFs

highlighted in bold in Table 2. For comparison, the gene region represented in the shaded

box (gene10004) shows a non-XGF from M. persicae with representative TE coverage close

to the mean for non-XGFs (1,519bp). XGFs 125_CYP380C52 and 105_CYP380C19 are not

shown, as these are nested at the CYP cluster containing 109_CYP380C31. Sea green

shaded regions indicate gene flanking regions. Grey bars represent gene bodies, and black

bars show exonic regions. TEs are annotated above their respective gene tracks, with

colours indicating the main TE classifications, as depicted in the key at the bottom of the

figure.

Figure 5. TE enrichment at XGFs compared to genome-wide levels. Fold-change enrichment

for TE coverage and TE count at XGFs compared to within-genome levels at non-XGFs

across 20 aphid species.

Figure 6. Fragmented TEs at XGFs across all aphid species, presented as TE count by TE

length with a bin width of 250bp. TE count is presented on a log10 scale. The main TE

classifications are indicated by the colours indicated in the key. Mean TE length across all

TEs at XGFs is 536bp.

Figure 7. TE association with cytochrome P450 genes in aphids. A. TE coverage around

CYP genes linked to xenobiotic resistance versus those with other functions. Numbers on

the right indicate the number of genes in each group being compared. B.TE coverage

around CYP genes linked to resistance to synthetic insecticides versus those conferring

resistance to natural xenobiotics. Numbers on the right indicate the number of genes in each

group under comparison.

Figure 8. TE occupancy at XGFs, housekeeping genes, and all other genes, and expression

of XGFs and housekeeping genes in A. pisum (Jaquiéry et al. 2022). A. TE coverage around

XGFs, housekeeping genes, and all other genes. Box limits indicate upper and lower

quartiles, whilst central lines indicate mean TE coverage. B. TE count around XGFs,

housekeeping genes, and all other genes. Box limits indicate upper and lower quartiles,

whilst central lines indicate mean TE count. C. Expression of XGFs and housekeeping

genes in germline tissues (testis and ovary) in A. pisum, expressed as reads per kilobase of

transcript per million base pairs sequenced (RPKM), to determine expected expression of

XGFs in germline cell types. Gene types are indicated by colours in the key.

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

## Tables

| TE Classification | Mean Share of TE Content (% of total TE content) | Standard Deviation of Share of TE Content (% of total TE content) |
|---|---|---|
| DNA | 45.27 | 18.00 |
| Unclassified | 35.95 | 14.66 |
| LINE | 7.52 | 3.76 |
| LTR | 6.19 | 8.00 |
| Rolling Circle | 3.46 | 2.16 |
| SINE | 2.14 | 5.24 |
| Penelope | 0.40 | 0.75 |

1026 Table 1. Mean and standard deviation of the percentage of total TE content attributed to
1027 each of the main TE classifications for the 20 aphid genomes included in this study.

1028

| Aphid Species | Gene ID | XGF Type | Mean Gene Size (by gene type) | Gene Size in bp (Z-score) | XGF associated TE Coverage (bp) | Fold-increase above aphid-wide enrichment (XGF-type specific) | Fold-increase above aphid-wide enrichment (non-XGF genes) | XGF associated TE Count | Fold-increase above aphid-wide enrichment (XGF-type specific) | Fold-increase above aphid-wide enrichment (non-XGF genes) | Gene Size after removing genic TEs (Z-score) |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | **Greatest Enrichment of TE Coverage** | | | | | | |
| *Acyrthosiphon pisum* | 2399_UGT344E8_v2 | UDP-glucuronosyltransferase | 8,302 | 55,119 (4.63) | 55,257 | 9.05 | 36.38 | 118 | 9.08 | 79.73 | 24,991 (2.45) |
| *Acyrthosiphon pisum* | 2354_UGT344D11_v2 | UDP-glucuronosyltransferase | 8,302 | 57,143 (4.83) | 45,208 | 7.40 | 29.76 | 96 | 7.38 | 64.86 | 26,620 (2.68) |
| *Myzus persicae* | 125_CYP380C52 | Cytochrome P450 | 7,221 | 100,885 (9.70) | 44,205 | 6.28 | 29.10 | 102 | 6.00 | 68.92 | 56,825 (5.97) |
| *Myzus persicae* | 109_CYP380C31 | Cytochrome P450 | 7,221 | 99,455 (9.55) | 42,790 | 6.08 | 28.17 | 94 | 5.53 | 63.51 | 57,937 (6.10) |
| *Myzus persicae* | 352_gst_g525.t1 | Glutathione S-Transferase | 3,450 | 16,231 (2.65) | 40,940 | 5.02 | 26.95 | 45 | 2.50 | 30.41 | 10,854 (1.54) |
| *Myzus persicae* | 105_CYP380C19 | Cytochrome P450 | 7,221 | 61,287 (5.60) | 39,133 | 5.56 | 25.76 | 100 | 5.88 | 67.57 | 34,033 (3.17) |
| *Metopolophium dirhodum* | 2360_UGT344A13 | UDP-glucuronosyltransferase | 8,302 | 18,807 (1.04) | 38,003 | 6.22 | 25.02 | 45 | 3.46 | 30.41 | 8,327 (0.11) |
| *Metopolophium dirhodum* | 1793_UGT345A3 | UDP-glucuronosyltransferase | 8,302 | 6,504 (-0.18) | 37,746 | 6.18 | 24.85 | 29 | 2.23 | 19.59 | 4,684 (-0.40) |

| Species | Gene ID | Annotation | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| *Sitobion miscanthi* | 690_UGT341A5_v2 | UDP-glucuronosyltransferase | 8,302 | 4,437 (-0.38) | 36,666 | 6.00 | 24.14 | 42 | 3.23 | 28.38 | 3,950 (-0.50) |
| *Sitobion miscanthi* | 687_UGT341A5 | UDP-glucuronosyltransferase | 8,302 | 4,456 (-0.38) | 36,179 | 5.92 | 23.82 | 39 | 3.00 | 26.35 | 3,969 (-0.50) |
| *Metopolophium dirhodum* | 13_abc_g2222.t1 | ABC Transporter (ABC Subfamily G) | 11,211 | 14,244 (0.32) | 35,841 | 4.30 | 23.60 | 35 | 1.84 | 23.65 | 12,513 (0.06) |
| *Sitobion miscanthi* | 1630_UGT344E8_v2 | UDP-glucuronosyltransferase | 8,302 | 3,249 (-0.50) | 35,637 | 5.84 | 23.46 | 57 | 4.38 | 38.51 | 2,150 (-0.75) |
| *Sitobion miscanthi* | 72_est_g6279.t1 | Esterase | 10,888 | 35,579 (1.82) | 35,569 | 5.90 | 23.42 | 63 | 4.50 | 42.57 | 23,910 (1.03) |
| **Greatest Enrichment of TE Count** | | | | | | | | | | | |
| ***Acyrthosiphon pisum*** | **2399_UGT344E8_v2** | **UDP-glucuronosyltransferase** | **8,302** | **55,119 (4.63)** | **55,257** | **9.05** | **36.38** | **118** | **9.08** | **79.73** | **24,991 (2.45)** |
| ***Myzus persicae*** | **125_CYP380C52** | **Cytochrome P450** | **7,221** | **100,885 (9.70)** | **44,205** | **6.28** | **29.10** | **102** | **6.00** | **68.92** | **56,825 (5.97)** |
| ***Myzus persicae*** | **105_CYP380C19** | **Cytochrome P450** | **7,221** | **61,287 (5.60)** | **39,133** | **5.56** | **25.76** | **100** | **5.88** | **67.57** | **34,033 (3.17)** |
| ***Acyrthosiphon pisum*** | **2354_UGT344D11_v2** | **UDP-glucuronosyltransferase** | **8,302** | **57,143 (4.83)** | **45,208** | **7.40** | **29.76** | **96** | **7.38** | **64.86** | **26,620 (2.68)** |
| ***Myzus persicae*** | **109_CYP380C31** | **Cytochrome P450** | **7,221** | **99,455 (9.55)** | **42,790** | **6.08** | **28.17** | **94** | **5.53** | **63.51** | **57,937 (6.10)** |
| *Acyrthosiphon pisum* | 2423_abc_g11808.t1 | ABC Transporter (ABC Subfamily A) | 11,211 | 25,566 (1.53) | 25,867 | 3.11 | 17.03 | 84 | 4.42 | 56.76 | 19,631 (0.83) |
| *Sitobion avenae* | 1033_UGT344A27 | UDP-glucuronosyltransferase | 8,302 | 62,310 (5.34) | 33,066 | 5.41 | 21.77 | 82 | 6.31 | 55.41 | 45,645 (5.35) |
| *Acyrthosiphon pisum* | 1088_CYP380C8 | Cytochrome P450 | 7,221 | 10,900 (0.38) | 28,620 | 4.07 | 18.84 | 82 | 4.82 | 55.41 | 5,975 (-0.27) |
| *Sitobion miscanthi* | 911_UGT344A27 | UDP-glucuronosyltransferase | 8,302 | 64,596 (5.57) | 27,402 | 4.49 | 18.04 | 81 | 6.23 | 54.73 | 38,974 (4.41) |
| *Acyrthosiphon pisum* | 1_abc_g11808.t1 | ABC Transporter (ABC Subfamily A) | 11,211 | 39,350 (3.00) | 31,743 | 3.81 | 20.90 | 81 | 4.26 | 54.73 | 25,849 (1.50) |

1030  Table 2. XGF loci with the greatest enrichment for TE coverage and TE count in aphids. *Z*-scores indicate the number of standard deviations
1031  away from mean gene size for individual genes. *Z*-scores <-3 or >3 indicate statistically unusually small or large genes compared to mean XGF
1032  size, respectively. Entries in bold are XGFs that are highly enriched for both TE coverage and TE count. ABC transporter subfamilies have
1033  been labelled where appropriate (Dermauw and Van Leeuwen 2014).

1034

1035

| TE Type | Non-XGF | All XGFs | CYPs | GSTs | Esterases | UGTs | ABCs |
|---|---|---|---|---|---|---|---|
| **DNA** | 1,974 (2) | 4,026(10) | 4,445 (11) | 4,800 (11) | 3,513 (9) | 3,743 (8) | 4,609 (11) |
| **Rolling Circle** | 728 (1) | 794 (2) | 836 (2) | 849 (2) | 1,126 (3) | 706 (2) | 756 (2) |
| **Penelope** | 608 (1) | 456 (1) | 840 (2) | 175 (1) | 848 (2) | 185 (1) | 477 (1) |
| **LINE** | 959 (1) | 992 (2) | 965 (2) | 1,225 (2) | 1,492 (2) | 859 (2) | 1,126 (2) |
| **SINE** | 358 (1) | 1,144 (1) | - | - | - | 1,686 (1) | 602 (1) |
| **LTR** | 1,238 (2) | 1,817 (3) | 1,115 (1) | 2,528 (3) | 1,397 (2) | 1,124(2) | 3,826 (5) |
| **Unclassified** | 1,307 (1) | 2,255 (5) | 2,114 (5) | 2,290 (6) | 1,754 (4) | 2,508 (6) | 2,384 (6) |

1036    Table 3. Mean base pair coverage of each TE classification at XGFs among aphids.
1037    Numbers in brackets indicate mean TE copy number at each XGF for each TE classification.

1038

**A.** Genome Size **B.** TE Content **C.** Relative TE Activity

Outgroups

Coccidae — *Ericerus pela*
Pseudococcidae — *Maconellicoccus hirsutus*, *Phenacoccus solenopsis*
Aleyrodidae — *Bemisia tabaci*
Delphacidae — *Sogatella furcifera*
Liviidae — *Diaphorina citri*
Aphalaridae — *Pachypsylla venusta*

Aphididae

*Hormaphis cornu*
*Schlechtendalia chinensis*
*Cinara cedri*
*Eriosoma lanigerum*
*Sipha flava*
*Aphis craccivora*
*Aphis glycines*
*Aphis gossypii*
*Melanaphis sacchari*
*Schizaphis graminum*
*Rhopalosiphum maidis*
*Rhopalosiphum padi*
*Pentalonia nigronervosa*
*Acyrthosiphon pisum*
*Metopolophium dirhodum*
*Sitobion avenae*
*Sitobion miscanthi*
*Diuraphis noxia*
*Myzus cerasi*
*Myzus persicae*

Genome Size (Mb): 0 200 400 600

TE Coverage (% of Total Genome Size): 0 10 20 30 40 50

Percentage of Total Genome Size (%)

Kimura Distance (CpG Adjusted): 60 40 20 0

**Reproduction**
Holocyclic / Mostly Holocyclic
Anholocyclic / Mostly Anholocyclic

DNA   Rolling Circle   Penelope   LINE   SINE   LTR   Other (Simple repeat, microsatellite, RNA)   Unclassified

**A.** Shared TE Families

**B.** TE Localisation

**Outgroups**

*Hormaphis cornu*
*Schlechtendalia chinensis*
*Cinara cedri*
*Eriosoma lanigerum*
*Sipha flava*
*Aphis craccivora*
*Aphis glycines*
*Aphis gossypii*
*Melanaphis sacchari*
*Schizaphis graminum*
*Rhopalosiphum maidis*
*Rhopalosiphum padi*
*Pentalonia nigronervosa*
*Acyrthosiphon pisum*
*Metopolophium dirhodum*
*Sitobion avenae*
*Sitobion miscanthi*
*Diuraphis noxia*
*Myzus cerasi*
*Myzus persicae*

Intergenic Space    Gene Flanking Regions    Intron    Exon

Percentage of Total Families in Each Species

Number of Species Sharing TE Family

Unique    Pervasive

TE Coverage (% of Total Genome Size)

DNA    Rolling Circle    Penelope    LINE    SINE    LTR    Other (Simple repeat, microsatellite, RNA)    Unclassified

Difference in TE Coverage (kb): Xenobiotic Gene Families vs All Other Genes

Difference in TE Count: Xenobiotic Gene Families vs All Other Genes

DNA | Rolling Circle | Penelope | LINE | SINE | LTR | Unclassified

| | | | | |
|---|---|---|---|---|
| DNA | Rolling Circle | Penelope | LINE | SINE | LTR | Unclassified |

**A.** Resistance vs Other Functions

| Species | Value |
|---|---|
| *Hormaphis cornu* | 0/1 |
| *Schlechtendalia chinensis* | 2/3 |
| *Cinara cedri* | * 7/1 |
| *Eriosoma lanigerum* | 11/6 |
| *Sipha flava* | * 11/2 |
| *Aphis craccivora* | 25/50 |
| *Aphis glycines* | 24/51 |
| *Aphis gossypii* | 22/52 |
| *Melanaphis sacchari* | 19/39 |
| *Schizaphis graminum* | * 23/53 |
| *Rhopalosiphum maidis* | 20/51 |
| *Rhopalosiphum padi* | 21/60 |
| *Pentalonia nigronervosa* | 24/72 |
| *Acyrthosiphon pisum* | 29/128 |
| *Metopolophium dirhodum* | * 27/122 |
| *Sitobion avenae* | 25/118 |
| *Sitobion miscanthi* | 27/123 |
| *Diuraphis noxia* | 23/62 |
| *Myzus cerasi* | ** 23/156 |
| *Myzus persicae* | ** 27/124 |

TE Coverage (kb)

**B.** Resistance to
Synthetic vs Natural Compounds

| Species | Value |
|---|---|
| *Hormaphis cornu* | 0/0 |
| *Schlechtendalia chinensis* | 1/1 |
| *Cinara cedri* | 7/0 |
| *Eriosoma lanigerum* | 11/0 |
| *Sipha flava* | 11/0 |
| *Aphis craccivora* | 24/1 |
| *Aphis glycines* | 23/1 |
| *Aphis gossypii* | 22/0 |
| *Melanaphis sacchari* | 19/0 |
| *Schizaphis graminum* | 22/1 |
| *Rhopalosiphum maidis* | 19/1 |
| *Rhopalosiphum padi* | 20/1 |
| *Pentalonia nigronervosa* | * 22/2 |
| *Acyrthosiphon pisum* | * 24/5 |
| *Metopolophium dirhodum* | 23/4 |
| *Sitobion avenae* | 23/2 |
| *Sitobion miscanthi* | 24/3 |
| *Diuraphis noxia* | 21/2 |
| *Myzus cerasi* | * 21/2 |
| *Myzus persicae* | * 23/4 |

TE Coverage (kb)

Evidence for Role in Xenobiotic Resistance
No Evidence for Role in Xenobiotic Resistance

Resistance to Synthetic Compounds
Resistance to Natural Compounds

**A.** TE Coverage

Log$_{10}$(TE coverage (bp))

**B.** TE Count

TE Count

Cytochrome P450   ABC Transporter   Glutathione S-Transferase

UDP-Glucuronosyltransferase   Esterase   Housekeeping Genes

All Other Genes

**C.** Germline Gene Expression in *A. pisum*

Germline Gene Expression (RPKM)

Cytochrome P450   Esterase   Glutathione S-Transferase   UDP-Glucuronosyltransferase   ABC Transporter   Housekeeping Genes

Male Sample 1   Sexual Female Sample 1

Male Sample 2   Sexual Female Sample 2