Title: Accounting for unobserved population dynamics and aging error in close-kin mark-recapture assessments

Authors: John D. Swenson[1], Elizabeth N. Brooks[2], Dovi Kacev[3], Charlotte Boyd[4], Michael Kinney[5], Benjamin Marcy-Quay[6], Anthony Sévêque[7], Kevin Feldheim[8], Lisa M. Komoroske[1]

[1] Department of Environmental Conservation, The University of Massachusetts, Amherst, Massachusetts, USA

[2] Population Dynamics Branch, Northeast Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Woods Hole, Massachusetts, USA

[3] Marine Biology Research Division, Scripps Institute of Oceanography, San Diego, California, USA

[4] Conservation International, Africa Field Division, Nairobi, Kenya

[5] Life History Program, Pacific Island Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Honolulu, Hawaii, USA

[6] Rubenstein Ecosystem Science Laboratory, University of Vermont, Burlington, Vermont, USA

[7] Department of Wildlife, Fisheries and Aquaculture, Forest and Wildlife Research Center, Mississippi State University, Mississippi State, Mississippi, USA

[8] Pritzker Laboratory for Molecular Systematics and Evolution, The Field Museum, Chicago, Illinois, USA

Correspondence: John Swenson, Organismic and Evolutionary Biology Program, The University of Massachusetts, Amherst, Massachusetts, USA. E-mail: jswenson@umass.edu

Data availability: All code used in this work is annotated and freely available at https://github.com/JDSwenson/LemonSharkCKMR. The code will be deposited to Dryad and assigned a DOI upon acceptance. Kinship data for lemon sharks comes from: Feldheim, K. A., S. H. Gruber, J. D. DiBattista, E. A. Babcock, S. T. Kessel, A. P. Hendry, E. K. Pikitch, M. V. Ashley, and D. D. Chapman. 2014. Two decades of genetic profiling yields first evidence of natal philopatry and long-term fidelity to parturition sites in sharks. Molecular Ecology 23:110–117 and can be accessed at: https://doi.org/10.5061/dryad.1q9r8.

Keywords: abundance; CKMR; data-poor fisheries; demographic models; elasmobranch; population dynamics; shark

Abstract: Obtaining robust estimates of population abundance is a central challenge hindering the conservation and management of many threatened and exploited species. Close-kin mark-recapture (CKMR) is a genetics-based approach that has strong potential to improve monitoring of data-limited species by enabling estimates of abundance, survival, and other parameters for populations that are challenging to assess. However, CKMR models have received limited sensitivity testing under realistic population dynamics and sampling scenarios, impeding application of the method in population monitoring programs and stock assessments. Here, we use individual-based simulation to examine how unmodeled population dynamics and aging uncertainty affect the accuracy and precision of CKMR parameter estimates under different sampling strategies. We then present adapted models that correct the biases that arise from model misspecification. Our results demonstrate that a simple base-case CKMR model produces robust estimates of population abundance with stable populations that breed annually; however, if a population trend or non-annual breeding dynamics are present, or if year-specific estimates of abundance are desired, a more complex CKMR model must be constructed. In addition, we show that CKMR can generate reliable abundance estimates for adults from a variety of sampling strategies, including juvenile-focused sampling where adults are never directly observed (and aging error is minimal). Finally, we apply a CKMR model that has been adapted for population growth and intermittent breeding to two decades of genetic data from juvenile lemon sharks (*Negaprion brevirostris*) in Bimini, Bahamas, to demonstrate how application of CKMR to samples drawn solely from juveniles can contribute to monitoring efforts for highly mobile populations. Overall, this study expands our understanding of the biological factors and sampling decisions that cause bias in CKMR models, identifies key areas for future inquiry, and provides

55   recommendations that can aid biologists in planning and implementing an effective CKMR

56   study, particularly for long-lived data-limited species.

Introduction

Population abundance plays important roles in both fundamental and applied biological research and is associated with a wide range of ecological and evolutionary processes (Hassell 1975, Berryman 1989, Robertson 1996, Carbone et al. 2011, Ellegren and Galtier 2016). Abundance estimates and trends are also key metrics for conservation and management and are commonly used to assess conservation status (Wilson et al. 2011), quantify the impacts of threats and/or recovery efforts (Jennings 2000, Ward-Paige et al. 2012, Magera et al. 2013), and scale regulated harvest quantities (e.g., allowable biological catch, annual catch limits) for managed populations of target and non-target species. Consequently, a wide range of methods have been developed for estimating population abundance (Schwarz and Seber 1999, Wilson and Delahay 2001, McCauley et al. 2012).

Capture-mark-recapture (CMR) is one prominent and widely used method in which abundance is estimated by constructing capture histories for each sampled (or tagged) individual, estimating capture probabilities, and comparing the number of recaptured individuals to the total number of sampled individuals (Cormack 1964, Jolly 1965, Seber 1965). A number of variations of CMR methods have been developed over the years to account for varied population demographics and sampling schemes (Pollock 2000, Amstrup et al. 2010, Royle et al. 2013), but the approach remains largely intractable in situations where recapture rates are very low, as with many low density and highly mobile marine species (Kohler and Turner 2001, Webster et al. 2002, Boyd et al. 2018). In addition, estimating a capture probability for CMR requires an estimate of the rate at which tags are lost (Arnason and Mills 1981, Hyun et al. 2012) and reported (e.g., by fishermen or hunters; Green et al. 1983, Pollock et al. 2001, Sackett and Catalano 2017), and tag loss and reporting rates vary with the species and experimental design

(Oosthuizen et al. 2010). As such, their estimation is likely to require auxiliary studies that

demand more time and resources and may be reliant on cooperation from individuals that

encounter the tags. CMR provides direct information about the sampled demographic, but many

highly mobile marine species have spatially segregated life histories and are only available for

sampling in nearshore habitats as juveniles before transitioning to a less accessible pelagic

habitat as adults. In such cases, CMR results are restricted to providing direct information about

the juvenile portion of the population, while the population dynamics of adults can only be

modeled effectively if additional data are available and if key assumptions are met (Kendall

1999, Pollock 2000). As alternatives to CMR, surveys or transect-based methods can be helpful

tools to estimate regional abundance (Schwarz and Seber 1999). However, variability in survey

length, uncertainty surrounding the proportion of habitat sampled and shifts in habitat

availability, as well as changes in behavior arising from the presence of human observers and

observation error are common pitfalls that can make such methods unreliable or incomparable

across studies (McCauley et al. 2012, Boyd and Punt 2021, Davis et al. 2022).

While CMR, surveys, and transect-based methods can all be useful tools for abundance

estimation in certain contexts, applying them in an unbiased way can be prohibitively

challenging in many systems. When estimates of absolute abundance are infeasible, indices of

relative abundance are commonly used to assess populations of exploited species (Campbell

2015). In fisheries, abundance trends derived from catch and effort data (e.g., catch-per-unit-

effort, CPUE), in concert with biological reference points, can inform management by providing

critical information about whether a population is overfished or if overfishing is actively

occurring (Cortés and Brooks 2018). However, it is extremely challenging to account for all the

factors that could influence catchability (Maunder et al. 2006); hence, indices of relative

abundance derived from CPUE are rarely linearly proportional to actual abundance (Harley et al.

2001, Maunder and Punt 2004, Lynch et al. 2012). Fish or fisher behavior contributes to

hyperstability (Erisman et al. 2011, Ward et al. 2013) and biased inference about abundance

trends can result if CPUE data are interpreted in isolation, or if linearity between catch rate and

abundance is implicitly assumed (Maunder et al. 2006). Furthermore, estimating trends of

relative abundance for highly mobile species frequently requires the integration of multiple

independent surveys that suggest differing abundance trends, making it difficult to establish true

abundance patterns (Peterson et al. 2021). All of these issues are amplified in taxa such as

elasmobranchs (sharks, skates, and rays), where reported catch data are often unreliable (Cortés

and Brooks 2018). While CPUE can provide invaluable information regarding stock status and

harvest pressure when analyzed in the right context (e.g., via an integrated model that

incorporates additional data streams), there is an urgent need for methods that can provide robust

estimates of absolute population abundance in circumstances where catch data are unreliable or

strongly correlated with factors other than population trend (e.g., changes in fishing practices,

skill, or gear improvement, or environmental perturbations).

Close-kin mark-recapture (CKMR) is a genetics-based approach for estimating absolute

population abundance that overcomes many of the logistical challenges associated with CMR

and other abundance estimation methods (Skaug 2001, Bravington et al. 2016a). As such,

CKMR has great potential to expand monitoring efforts and improve or enable assessments of

species for which conventional methods are intractable. In contrast to conventional CMR, the

tags in CKMR are genotypes, and animals are considered "re-captured" when their kin are

identified (Bravington et al. 2016a). This removes the need for individual recapture and allows

for the estimation of adult abundance using samples collected solely from juveniles, as well as

samples obtained lethally through fishing or hunting (Bravington et al. 2016a, Hillary et al.

2018). While CKMR can theoretically leverage any relationship, the most common applications

so far have focused on parent-offspring pairs (POPs) (Bravington et al. 2016b, Ruzzante et al.

2019, Marcy☐Quay et al. 2020) and/or half-sibling pairs (HSPs) (Hillary et al. 2018, Patterson et

al. 2022a). Similar to conventional CMR, CKMR can estimate quantities beyond abundance,

including survival (Hillary et al. 2018), fecundity (Bravington et al. 2016b), dispersal (Feutry et

al. 2017, Conn et al. 2020, Patterson et al. 2022a), and, potentially, population growth rate,

though which parameters can be estimated depends on the form of the model and type of kin

pairs modeled. In cases where sampling is limited to juveniles, CKMR can provide added value

to conventional CMR by generating parameter estimates for the adult population while CMR

estimates parameters for the sampled (in this case juvenile) portion of the population. These

advantages and possibilities make CKMR an exciting tool to improve monitoring efforts and

population assessments for data-limited species of management and conservation concern, either

in conjunction with, or in place of, conventional CMR.

Despite CKMR's strong potential to provide key information for conservation and

management, its implementation has been slowed by a lack of clarity regarding the flexibility

and limitations of the method. Several studies have discussed factors that are likely to cause bias

if left unaccounted for in CKMR models (Bravington et al. 2016a, Conn et al. 2020, Waples and

Feutry 2021, Trenkel et al. 2022), but there have been few quantitative assessments of the biases

that arise from applying an overly simplistic CKMR model to a population with complex

dynamics (but see Conn et al. 2020, Waples and Feutry 2021). For example, a simple base-case

CKMR model (e.g., Equations 3.3 and 3.10 in Bravington et al. 2016a) produces an abundance

estimate that assumes abundance is constant over the modeled time period. However, real

populations experience interannual fluctuations in population size. If such changes are persistent

or severe (e.g., following an environmental disaster or introduction of heavy fishing pressure),

then it will be necessary to specify a more complex CKMR model that can accommodate a

changing population if year-specific abundance estimates are desired.

When modeling half-sibling relationships in particular, a simple base-case CKMR model

assumes that the probability of two individuals sharing a parent is a simple exponential function

of the year gap that separates their births. However, many long-lived species exhibit intermittent

breeding whereby one or more years elapse between reproductive events (Morbey and Shuter

2013, Shaw and Levin 2013, Desprez et al. 2018, Bauwens and Claus 2019, Skjæraasen et al.

2020), resulting in different probabilities of detecting half-siblings depending on the age gap

(Waples and Feutry 2021). Systematic intermittent breeding will cause bias in CKMR parameter

estimates if unaccounted for in the model (Waples and Feutry 2021). While it may be possible to

infer breeding periodicity based on the distribution of observed kin pairs in the data, instances of

off-cycle breeding, mixed breeding schedules (e.g., a population comprising both annual and

multiennial breeders), and aging uncertainty that leads to errors in cohort assignment may

obscure the signal (Rivalan et al. 2005, Cubaynes et al. 2011, Öst et al. 2018, Higgs et al. 2020).

Finally, a core component of CKMR is the use of age data, which is required to assign

individuals to the correct cohort (Bravington et al. 2016a). Direct aging is very challenging for

some taxa (Cailliet 2015), and length-based age assignment is prone to bias when growth curves

are based on size-selective sampling, as they often are (Gwinn et al. 2010). While more advanced

statistical methods can account for uncertainty in aging during the modeling process (Schwarz

and Runge 2009), it may also be possible to alleviate bias by targeting sampling to age classes

that can be reliably aged, such as young-of-the-year (YOY) which are often easily distinguished

from other age classes by their small size and/or the presence of umbilical scars (Feldheim et al.

2002). However, sampling constraints will not always permit long-term sampling of YOY and

the number of cohorts required to produce robust parameter estimates with CKMR is unclear. A

better understanding regarding the circumstances in which unobserved population dynamics or

sampling limitations are likely to bias CKMR model estimates, in combination with strategies to

mitigate that bias, will help ensure robust application of the method and facilitate its integration

into conservation and management frameworks.

Elasmobranchs (sharks, skates, and rays) are a group of highly vulnerable marine species

that play key ecological roles as apex- and meso-predators in ecosystems around the world

(Vaudo and Heithaus 2011, Ferretti et al. 2018) and are likely to benefit from future application

of CKMR. Around one-third of the 1200+ elasmobranch species are threatened with extinction,

due primarily to overfishing (Dulvy et al. 2021), while nearly half of elasmobranch species

(46%) are classified on the IUCN Red List of Threatened Species as Data Deficient and only a

small fraction of exploited populations are managed sustainably (Kindsvater et al. 2018).

Conventional methods for estimating abundance and mortality are intractable for many

elasmobranch populations because individual recapture rates for highly mobile elasmobranch

species can be very low (Kohler and Turner 2001), and it can be logistically challenging to

physically capture and mark larger species (Guttridge et al. 2017). In contrast to conventional

methods, CKMR requires only small tissue samples that can be obtained from adults via biopsy,

or from juveniles that are easier to handle than their adult counterparts. There is also no need for

individual recapture so each animal only needs to be captured and handled once, making this a

more feasible approach for many elasmobranch populations. In addition, when adults are

unavailable for sampling, the life histories of many elasmobranch species allow for the use of

juvenile-only CKMR models (e.g., half-sibling (HS) CKMR) that can estimate adult abundance

without sampling a single adult (Bravington et al. 2016a, Førland 2019). Considering that many

migratory elasmobranchs use nursery areas where juveniles are more readily available for

sampling than adults (Heupel et al. 2007), the potential for CKMR to provide novel insights into

difficult-to-study elasmobranch populations is vast.

CKMR has been applied to several elasmobranch populations to date (Bradford et al.

2018, Hillary et al. 2018, Bravington et al. 2019, Delaval et al. 2022, Trenkel et al. 2022,

Patterson et al. 2022a) and is likely to be an important tool to inform elasmobranch conservation

and management in the future. However, elasmobranch populations are susceptible to steep

population declines arising from overexploitation (Ferretti et al. 2018), commonly exhibit

multiennial breeding cycles (Nosal et al. 2021), and are exceptionally challenging to age (Cailliet

2015). As such, there is a risk that CKMR models that do not sufficiently account for these

factors will produce biased parameter estimates that will be unwittingly incorporated into

management frameworks, leading to incorrect management actions that ultimately threaten

elasmobranch populations.

To facilitate the robust application of CKMR to elasmobranchs and other long-lived taxa

facing similar challenges with abundance estimation, we investigated the sensitivity of CKMR

parameter estimates to unmodeled dynamics related to population growth and breeding schedule,

as well as uncertainty in age assignment. We used stochastic individual-based simulation to

generate distinct populations of lemon sharks (*Negaprion brevirostris*), a representative long-

lived species with promiscuous mating and multiennial breeding, under different population

dynamics scenarios and sampled each population using three sampling schemes that targeted

different age classes. Two different CKMR models were fit to each dataset: one that was naïve to

at least one component of the data-generating model (naïve model) and one that was adapted to account for all relevant population dynamics (adapted model). We compared the bias in parameter estimates from both models (naïve vs adapted) and across all three sampling schemes, including one in which age data were unreliable. Finally, we applied a model that was adapted for population growth and multiennial breeding to two decades of real genetic data from a small population of lemon sharks in Bimini, Bahamas, to generate a time-series of abundance estimates for the breeding population of females. Collectively, these results provide important insights into the ways in which unmodeled population dynamics, sampling selectivity, and aging error affect CKMR model performance, while also offering guidance regarding sampling design and model construction.

2. Methods

Our simulation framework comprised four primary components: 1) an individual-based population simulation that stochastically generated distinct populations with known parameters, 2) selective sampling of age classes from those populations, 3) construction of a pairwise comparison matrix from the samples, and 4) a CKMR model that was fit to the pairwise comparison matrix to estimate the known population parameters. The first three components comprised our data-generating model (DGM) while the latter formed our estimation model (Figure 1).

We then tested the interplay of population dynamics and model complexity by iteratively varying a subset of population parameters (Table 1) and fitting two CKMR models to the data: one that was naïve to the added dynamics, and one that was adapted to account for them. Each

scenario was repeated 500 times, with each iteration producing a population with a distinct

pedigree and parameter estimates.

2.1 Data generating model

Parameters governing our individual-based population simulations were designed to

replicate the life history traits and population dynamics of lemon sharks in Bimini, Bahamas,

(similar to White et al. 2014) (Appendix S1: Table S1). Females bred with one, two, or three

distinct males each breeding cycle and produced two or three pups with each male, resulting in a

range of 2-9 total pups produced per female per year. We set no limit on the number of females

an individual male could breed with. As a consequence, the variance in reproductive output for

males was much greater, ranging from 2 – 41 offspring per breeding male per year (median = 6).

After maturity, fecundity was age-invariant, so sex-specific lifetime reproductive output was

approximately equal across the population. Survival was assumed constant within each of three

life stages, which we designated as young-of-year (YOY; age 0), juvenile (age 1-11) and adult

(age 12-50). We assigned knife-edged maturity following White et al. (2014), so every individual

age 12 and over was available for breeding, while no individuals younger than age 12 were

allowed to breed.

2.1.1 Population growth

We varied population growth in our DGM by reducing or adding mortality on juveniles

and adults (Appendix S1: Table S1). For the slight increase and decline scenarios (Table 1:

scenarios 2.1 and 2.2), mortality was increased or reduced by ~1% from the beginning of the

simulation, resulting in a population growth rate of +/- 1% per year. To achieve more substantial

declines in population size (Table 1: scenario 2.3), we simulated a stable population for 80 years

and then stochastically imposed 4-7% added mortality for juvenile and adult age classes for years

81-90 (when imposed from the beginning of the simulation, the population invariably went

extinct). This produced a population that declined at a rate of ~7% per year for the last 10 years

of the simulation.

2.1.2 Intermittent breeding

Many elasmobranchs systematically breed on multiennial cycles (Feldheim et al. 2002,

2017, Nosal et al. 2021). To examine the bias that accrues when this trait is unaccounted for in a

CKMR model, we ran simulations where all females bred on a bi- or triennial cycle (Table 1:

scenario 3.1 – 3.3), including one scenario where we allowed 20% of breeding females to

stochastically breed off-cycle or fail to breed when they were on-cycle (scenario 3.2). Each

female in our simulation was assigned a breeding cycle at birth, which determined the first year

of reproduction for multiennial breeders. In scenarios with biennial breeders, this resulted in a

population where half of the females reproduced for the first time in the year they matured (age

12) and the other half reproduced for the first time the following year (age 13). For the scenario

with triennial breeding, a third group reproduced for the first time at age 14. Mature males were

assumed available to breed every year once they reached maturity at age 12.

2.2 Sampling

All simulated populations were sampled using three different schemes that selected for

different age classes: the first drew samples exclusively from young-of-year (age 0) individuals;

the second made juveniles of all ages except young-of-year (ages 1-11) available to sample; and

the third allowed sampling of all age classes (ages 0-50). These scenarios were chosen to

replicate potential sampling opportunities for elasmobranchs such as nursery areas (Feldheim et

al. 2002, Heupel et al. 2007), juvenile aggregation sites (Rowat et al. 2007, Jacoby et al. 2012),

and resident populations (Snelson and Williams 1981), respectively.

In each case, the population was initially sampled at four different intensities representing

0.5%, 1%, 1.5%, and 2% of the population. Samples were drawn annually and non-lethally for

four years at the end of the population simulation (i.e., years 87 – 90), following reproduction but

before mortality each year. With a stable population, sampling 1.5% of the population resulted in

an average of 616 total samples and 100-200 half-sibling pairs (HSPs), which is expected to

produce a reasonable CV for all three sampling schemes (Bravington et al. 2016a). Therefore,

following model validation, we focused on sampling 1.5% of the population for the remainder of

our simulations.

2.2.1 Aging uncertainty

A crucial component of CKMR is accurate aging, yet some taxa, including

elasmobranchs, are notoriously difficult to age, with most efforts relying on length-at-age growth

curves to assign age to sampled individuals (Cailliet 2015). To examine how imprecision in

growth curves affects CKMR parameter estimates, we first constructed an age-length key for

lemon sharks using data from a long-term study of the population in Bimini, Bahamas (Feldheim

et al. 2014), and calculated the standard deviation of lengths for individuals with known ages, the

majority of which (>95%) spanned ages 0-3. We then simulated lengths for each sampled

individual (which were assigned ages in our DGM) using a von Bertalanffy growth curve for the

species (Brown and Gruber 1988). Each individual was assigned a length by drawing a value

from a normal distribution with the mean length-at-age specified by the von Bertalanffy curve,

and the standard deviation derived empirically from our age-length key for individuals aged 0-2,

and arbitrarily from a CV of 5%, 10%, or 20% for individuals aged 3+. After assigning lengths to

each individual, we used a reverse von Bertalanffy growth curve with the same values for

theoretical age of zero size (t_0 = -2.302), asymptotic average length (L_{inf} = 317.65), and the

growth coefficient (K = 0.057) and then re-assigned ages to sampled individuals based on their

lengths, rounding to the nearest integer. This produced plausible, yet sometimes incorrect, ages

(similar to age-slicing; see Ailloud et al. 2015, e.g.). The re-assigned ages were then used to

construct the pairwise comparison matrix that was input to the CKMR model.

2.3 Pairwise comparison matrix

CKMR produces estimates of abundance and other population parameters by defining

kinship probabilities for every pair of sampled individuals given relevant covariates (e.g., age,

sex). We constructed two standard pairwise comparison matrices for each set of samples. The

first matrix contained positive and negative kinship assignments for half-siblings. To satisfy the

assumption of independent sampling, whenever full siblings or self-recaptures were present, all

but one individual/instance was removed prior to construction of the matrix. Once the matrix was

created, within-cohort comparisons were removed. Though CKMR models can be adapted to

incorporate within-cohort comparisons (Førland 2019), without considerable modifications to the

equations, within-cohort and cross-cohort comparisons will estimate different quantities (Waples

and Feutry 2021). As such, removing within-cohort comparisons is common practice for

application of HS CKMR (Bravington et al. 2016a). Kinship assignment in our simulations was

known without error, so each comparison was assigned as a positive if the two individuals being

compared were a half-sibling pair, and negative if not. Because half-siblings are genetically

indistinguishable from aunt/niece (uncle/nephew, etc.) pairs, we included one scenario (Table 1:

scenario 1.2) where we allowed such comparisons to contaminate the pool of half-siblings and

evaluated the degree to which these false positives affected parameter estimates.

The second matrix was composed of parent-offspring (PO) comparisons, which was only

relevant to the scenario that included sampling of adults. For each birth year represented in the

dataset, individuals that were alive in that year were split into potential offspring or parents based

on whether they were born in that year (potential offspring), reproductively mature at the time

(potential parent), or neither, in which case they were left out of the matrix corresponding to that

year. Each comparison was assigned as a positive if they were related as parent-offspring or

negative if not.

Once the appropriate half-sibling and parent-offspring comparisons were defined, all

matrices were collated and grouped by 1) type of relationship (half-sibling or parent-offspring),

2) birth year of younger individual in each comparison (a.k.a. y_j; see Section 2.4.1), 3) reference

year gap (a.k.a. $(y_j - t_0)$; see Section 2.4.2), and 4) birth year gap (a.k.a. δ; see Section 2.4.1), as

applicable. The number of observed kin pairs (Y) was then modeled as a random variable, with

the probability of success defined by Equations 1-9 below, and n equal to the total number of

comparisons in each group (see Appendix S1: S1.1 and Table S2 for more details on the pairwise

comparison matrix, and Appendix S1: S1.2 for more details on kinship types that can cause

issues for half-sibling CKMR e.g., aunt/niece pairs).

2.4 Estimation models

Kinship probabilities for each pairwise comparison in CKMR are derived from the

expected reproductive output of individual animals (defined by covariates such as age and sex)

relative to the total reproductive output of the population in the birth year of the younger

individual in each pairwise comparison (Bravington et al. 2016a). The specific equations we

used to define kinship probabilities in our CKMR models varied with the scenarios we tested,

with each scenario comparing a "naïve" model to an "adapted" model, where the naïve model

ignored one key dynamic of the simulated population and the adapted model accounted for it.

Our equations are based on the general equations defined in Bravington et al. (2016a).

2.4.1 Base-case CKMR model

Let $P\{K_{i,j} = MHSP\}$ be the probability that individuals i and j are a maternal half-sibling

pair (i.e. they share a mother but not a father). Probabilities for $K_{i,j}$ depend on the likelihood that

the same individual that birthed the older offspring (i) survived and gave birth to the younger

offspring (j). If we assume that all animals of reproductive age in the population during i and j's

birth years are equally likely to have birthed each of them, then the probability of kinship (K) can

be defined as

370
$$P\{K_{i,j} = MHSP\} \sim Binomial\left(\frac{\phi^\delta}{N_{(y_j)}},\ R_{(y_i,y_j)}\right) \tag{1,}$$

where,

ϕ is the annual survival probability for adults,

δ is the number of years between the birth years of individuals i and j (i.e. $y_j - y_i$) during

which any potential parent of i may have died a.k.a. the "birth year gap",

y_j is the birth year of individual j (the younger sibling),

$R_{(y_i, y_j)}$ reflects the total number of pairwise comparisons between individuals born in

years y_i and y_j, and

$N_{(yj)}$ is the total number of mature females in year y_j.

Now, let $P\{K_{i,j} = MPOP\}$ refer to the probability that individuals i and j are related as a

maternal parent-offspring pair (MPOP). In this case, survival only enters the equation for

MPOPs if sampling is non-lethal (as it was in our simulations) *and* if the potential parent was

sampled before the potential offspring was born. If, on the other hand, the potential mother i was

captured in or after offspring j's birth year and was reproductively mature, then we know that she

was alive in the year the offspring was born and, assuming constant fecundity across the

population, is equally likely to have birthed j as any other potential mother. Assuming the pool of

potential parents was filtered to only include individuals that were mature and were not known to

have died before j's birth year, the probability of kinship is

$$P\{K_{i,j} = MPOP\} \sim Binomial \left(\begin{array}{cc} \frac{\phi^{(y_j - c_i)}}{N_{(yj)}}, \ R_{(c_i, y_j)} & c_i < y_j \\[2ex] \frac{1}{N_{(yj)}}, \ R_{(c_i, y_j)} & c_i \geq y_j \end{array} \right) \tag{2},$$

where c_i refers to the year in which the potential parent was captured. For each offspring

birth year (y_j), it is crucial to ensure that individuals who were not reproductively mature are not

included as potential parents for that year. This restriction can be added directly to the model

(Bravington et al. 2016a, Conn et al. 2020) or implemented during construction of the pairwise

comparison matrix, as was done here.

Given that sampling was nonlethal for the parent, then mother i's survival to the year

of j's birth is conflated with detection probability (when $c_i < y_j$). In circumstances where the

individual recapture rate of adults is non-negligible, an additional parameter defining the adult

detection probability will be required to disentangle the state ($\Box$) and observation (detection

probability) processes. However, if sampling is sparse such that individual recaptures are

exceedingly rare, the cost of estimating an extra parameter for detection probability likely

outweighs the benefits.

Equations 1 and 2 define our base-case CKMR model. Though the probabilities presented

here focus on maternal kinship, the same probabilities apply to males and paternal kinship (see

Equations 8 and 9 below). While all parameter estimates presented here incorporate HSPs, POPs

were only included in the likelihood for the sampling scheme in which adults were sampled with

all other age classes; otherwise, the likelihood included HS kinship probabilities only.

2.4.2 Population growth model

To account for population growth/decline in our CKMR model, we defined a simple

exponential growth model to describe the population dynamics, where

$$N_t = N_{(t0)} \lambda^t \tag{3}.$$

As such, the kinship probabilities become:

$$414 \qquad P\{K_{i,j} = MHSP\} \sim Binomial\left(\frac{\phi^{\delta}}{N_{(t_0)}\lambda^{(y_j-t_0)}}, \; R_{(y_i,y_j)}\right) \qquad (4)$$

415               and

$$416 \qquad P\{K_{i,j} = MPOP\} \sim Binomial\left(\begin{array}{ll} \frac{\phi^{(y_j-c_i)}}{N_{(t_0)}\lambda^{(y_j-t_0)}}, \; R_{(c_i,y_j)} & c_i < y_j \\[2em] \frac{1}{N_{(t_0)}\lambda^{(y_j-t_0)}}, \; R_{(c_i,y_j)} & c_i \geq y_j \end{array}\right) \qquad (5).$$

418     Here, λ defines the annual population growth rate and t_0 refers to the initial model year,

419   also called the "reference year" (Bravington et al. 2016a). The reference year typically refers to

420   the earliest instance of y_j in the pairwise comparison matrix but could refer to any modeled year.

421   To assess the capacity of CKMR to generate year-specific abundance estimates, we fit the same

422   CKMR model to each dataset four times. In each instance, we set the reference year (t_0) to the

423   earliest instance of y_j. Then, we estimated N_t 10 years prior to the reference year (before data

424   were collected), in the reference year (t_0), 5 years prior to the last year of the simulation, and the

425   last year of the simulation (i.e., present). We also tested two different methods for generating

426   year-specific abundance estimates: one where t_0 was fixed to the first year of data (first instance

427   of y_j) and N_t was calculated as a derived quantity (Table 2 – our primary approach), and one

428   where t_0 was directly set to the year of interest (i.e., N was directly estimated in year t; see

429   Appendix S1: S1.3 for more discussion on CKMR with a changing population).

431   2.4.3 Intermittent breeding model

432     If a population – or subset of a population – systematically breeds on a non-annual

433   schedule, then CKMR estimates will be biased unless this behavior is accounted for in the model

(Waples and Feutry 2021). We accounted for intermittent breeding dynamics in our CKMR

model via the inclusion of parameters a and Ψ, where a refers to the number of years between

breeding (e.g., 2 for biennial breeders), and Ψ is the proportion of individuals that breed every a

years (similar to Patterson et al. 2022b). This implies that ($1 - \Psi$) individuals breed annually. We

assume that the proportion of on-cycle breeders that breed in a given year is $1/a$. Thus, the

effective number of female breeders in a given year ($\widetilde{N}_t$) is given by

$$(1 - \psi)N_t + \frac{\psi}{a}N_t \tag{6}$$

Accounting for interannual population dynamics (Eq. 3), the full probability of maternal

half-sibling kinship for a population that reproduces on a multiennial schedule becomes

$P\{K_{i,j} = MHSP\} \sim Binomial \left(\begin{array}{ll} \frac{a\phi^{\delta}(1-\psi)}{(a+\psi-a\psi)N_{(t_0)}\lambda^{(y_j-t_0)}}, \ R_{(y_i,y_j)} & \delta \text{ not evenly divisible by } a \\[2em] \frac{a\phi^{\delta}}{(a+\psi-a\psi)N_{(t_0)}\lambda^{(y_j-t_0)}}, \ R_{(y_i,y_j)} & \delta \text{ evenly divisible by } a \end{array} \right)$

(7).

If 100% of females breed on a biennial cycle (i.e. $a = 2$ and $\Psi = 1$), then the probability

of finding half-siblings that are separated by an odd number of birth years is 0. It is the presence

of δ intervals that are not evenly divisible by a that provide information on the parameter Ψ (see

Appendix S1: S1.4). Alternatively, when a consistent pattern is observed in the year gaps that

separate HSPs, one may be tempted to remove off-cycle comparisons from the pairwise

comparison matrix - since they have no chance of revealing a positive comparison - and fit a

model that is naïve to intermittent breeding (e.g., Eq. 4) to the on-cycle comparisons only. We

thus evaluated two different types of "naïve" model: one where we retained off-cycle

comparisons in the pairwise comparison matrix (the "naïve" model), and one where we removed

off-cycle comparisons (the "naïve - filtered" model).

In our simulations, intermittent breeding dynamics were only present for females, and all

males in the population were available for breeding each year; as such, Equation 7 only applied

to maternal comparisons, while kinship probabilities for paternal half sibling pairs (PHSPs)

continued to be defined as:

459
$$P\{K_{i,j} = PHSP\} \sim Binomial\left(\frac{\phi^{\delta}}{N_{(t_0)}\lambda^{(y_j - t_0)}}, \ R_{(y_i, y_j)}\right)$$
(8),

460   which is the same as the maternal kinship probability defined in Equation 4.

461   Because the parent is directly sampled in PO CKMR, there is no need to explicitly

462   account for breeding periodicity in the likelihood; therefore, we continued to use Equation 5 for

463   maternal PO comparisons when applicable. Similarly, kinship probabilities for paternal parent-

464   offspring pairs (PPOPs) mirrored those for MPOPs in Equation 5:

465
$$P\{K_{i,j} = PPOP\} \sim Binomial\left(\begin{array}{ll} \frac{\phi^{(y_j - c_i)}}{N_{(t_0)}\lambda^{(y_j - t_0)}}, \ R_{(c_i, y_j)} & c_i < y_j \\[2ex] \frac{1}{N_{(t_0)}\lambda^{(y_j - t_0)}}, \ R_{(c_i, y_j)} & c_i \geq y_j \end{array}\right)$$
(9).

For all multiennial simulations, we simulated a population with an approximately stable

growth rate, set the reference year (t_0) to the earliest instance of y_j, and estimated N_t in the

present (i.e., most recent year of sampling), as we expect this to be a common approach in real-

life applications of CKMR.

2.4.4 Estimation framework

We adopted a Bayesian approach to CKMR parameter estimation, which allows for the

incorporation of auxiliary data and/or expert knowledge as priors on model parameters (Kéry and

Schaub, 2012b). For the scenarios tested here, survival and other parameters were assigned

reasonably diffuse priors to reflect data-limited situations (Table 2). Though it is possible to

estimate sex-specific survival ($\square$) and population growth rates (λ), these parameters were shared

between males and females in our models. The posterior distributions for parameters were

approximated using Markov Chain Monte Carlo (MCMC) sampling, implemented using the

software JAGS (Plummer 2003) and applied in the R environment (Denwood 2016, R Core

Team 2021). We ran two Markov chains with a thinning rate of 20, drawing 40,000 samples

from the posterior distribution following a burn-in of 50,000 samples. These settings were

empirically derived by assessing autocorrelation among successive draws and convergence

among the chains. We assessed convergence of the final Markov chains with trace plots and the

Gelman-Rubin statistic (Gelman and Rubin 1992), and removed from further analysis any

simulation replicate with an Rhat value > 1.01, although these instances were rare (~1.5% of

simulations).

2.5 Application to lemon sharks

2.5.1 Bimini lemon shark dataset

A long-term genetic dataset from lemon sharks in Bimini, Bahamas, was used to illustrate

application of our multiennial CKMR model (Eq. 7) to a dataset derived entirely from juvenile

tissue samples (Feldheim et al. 2014). Lemon sharks are large viviparous (live-bearing)

elasmobranchs that reach sexual maturity at approximately 12 years of age (Brown and Gruber

1988) with a lifespan exceeding 30 years (Brooks et al. 2016). Female lemon sharks at Bimini

are regionally philopatric and return to Bimini to pup on a biennial schedule, while the males

with which they mate likely reproduce over a much larger area (Feldheim et al. 2002). Juveniles

use the shallow waters surrounding Bimini as a nursery and remain in the area until 2-3 years of

age or until they reach 90 cm in length (Morrissey and Gruber 1993) and generally do not move

between the North and South Islands (Gruber et al. 2001). The Bimini nursery contributes to a

larger Western Atlantic population that is classified as Vulnerable on the IUCN Red List

(Hansell et al. 2018, 2021, Carlson et al. 2021). The Bimini nursery has been intensively studied

since 1995, with an estimated 99% of newborn sharks sampled at the Bimini North Island each

year (Dibattista et al. 2007). The ability to heavily sample multiple litters has allowed for reliable

reconstruction of maternal genotypes, while paternal genotype reconstruction has often relied on

relatively few newborns, resulting in high confidence in maternal kinship assignment and lower

confidence in paternal kinship (Feldheim et al. 2002, 2004, 2014).

Given the disparities in kinship assignment and breeding range, we focused our CKMR

model on maternal comparisons to estimate abundance and survival of adult females. We used

samples collected from the North Island, a small isolated nursery for lemon sharks aged 0-3

years old (Chapman et al. 2009), from 1993-2015. Most individuals in our dataset were sampled

as YOY (92%) and easily identified by the presence of umbilical scars, so their ages were

known. We estimated abundance of total females in our CKMR model using Equation 7, and

derived the number of effective female breeders in each year using Equation 6. Thus, our scope

of inference for parameter estimation encompassed the adult females that visited the North Island

nursery to give birth during each modeled year, a number which is likely very small (White et al.

516 2014). We excluded sampled individuals without a known birth year from analysis as well as

517 same-cohort comparisons (Bravington et al. 2016a), and any individuals for which maternal

518 kinship assignment was uncertain.

519  There were many full sibling pairs in the dataset (1,515 individuals contributing to 1,129

520 pairs), but very few cross-cohort full siblings (only 4% of full sibling pairs). Including more than

521 one individual from each litter in a CKMR analysis can result in non-independence among

522 pairwise comparisons and unreliable estimates of variance, especially in small populations where

523 sampling effort may be high relative to the population size (i.e., non-sparse sampling; Bravington

524 et al. 2016a, 2019). Because Bimini lemon sharks were exhaustively sampled with relatively few

525 instances of cross-cohort full siblings, we hypothesized that the retention of littermates might

526 provide valuable data to the CKMR model, even if it reduced the reliability of variance

527 estimates. Therefore, we fit our multiennial CKMR model (Eq. 7) to two sets of data: one where

528 we included full littermates in the analysis (though we still removed all within-cohort

529 comparisons from the pairwise comparison matrix) and one where we only retained one

530 individual per mother/sire breeding pair, similar to our approach with the larger simulated

531 populations.

532  Finally, recognizing that the Bimini lemon shark dataset is unique in how thoroughly the

533 population was sampled, we also examined whether the model performed similarly with a

534 sparser dataset by randomly downsampling and reducing the number of samples from each year

535 to 30% of the full dataset. To account for random variation surrounding which samples were

536 retained, we iterated over the process 50 times, fit a CKMR model to each set of samples, and

537 report the average of the median and 95% highest posterior density intervals (HPDI) of the 50

538 posterior distributions.

2.5.2 Bimini lemon shark simulations

Preliminary application of our multiennial CKMR model (Eq. 7) to the real Bimini

dataset suggested the population likely experienced alternating periods of growth and decline

during the modeled period. Our demographic model (Eq. 3) assumes the population is growing

exponentially, and we suspected that this may result in an averaging effect and imprecise

parameter estimates over our multi-decadal time series of data, especially if an inconsistent trend

was present. To test this hypothesis and further examine the effects of applying our model to a

small population, we refined our DGM to produce a population of similar size and with similar

dynamics as Bimini lemon sharks (Feldheim et al. 2002, DiBattista et al. 2011, White et al.

2014) and then sampled 90% of the YOY from that population over 20 years to achieve a dataset

that resembled the real dataset.

We fit the first CKMR model after four years of sampling. Then, to replicate the type of

real-time estimates that could be produced by integrating CKMR into long-term monitoring

efforts, we iteratively added one year of samples to the dataset until reaching the end of the time

series, fitting three CKMR models each time a year of samples was added: one that included all

samples that had been collected up to the most recent year of sampling, one that subset for

samples within a five-year window of the most recent year of sampling, and one that subset for

samples within a three-year window. In each case, the reference year (t_0) was set to the birth year

of the second oldest individual in the dataset being used (i.e., the first instance of y_j), and

abundance ($N_{\female t}$) was estimated in the most recent year of sampling following Eq. 3. Abundance

trend was then tracked in two ways: via a time-series of female abundance, and estimation of λ

over the window of included samples. Finally, we applied the same approach to the real genetic

dataset to estimate the total number of adult females, then derived the number of year-specific

female breeders using Equation 6, and compared our estimates of yearly female breeders to

estimates that have been independently obtained for the population using a reconstructed

pedigree (DiBattista et al. 2011).

3. Results

3.1 Model validation

When the assumptions of the base-case model (Eqs. 1 and 2) were met, CKMR produced

unbiased estimates of abundance under all sampling schemes and intensities (Figure 2a), with

increasing precision as sampling intensity increased (Figure 2b). The model produced unbiased

estimates of abundance whether the likelihood included HSPs only (as in the "sample juveniles"

and "target YOY" scenarios) or jointly considered HSPs and POPs (as in the "sample all ages"

scenario), though we note improved precision for the latter. At very low sampling intensities

(0.5% of the target population sampled), fewer than 25 HSPs were identified for all sampling

schemes (Figure 2c) and fewer than 5 parent-offspring pairs (POPs) were identified for the

sampling scheme that included all ages (Figure 2d). In contrast, when 2% of the population was

sampled, over 200 HSPs were identified on average for all sampling schemes, while 10-40 POPs

were identified for the scenario in which all age classes were sampled. Including aunt/niece

(uncle/nephew, etc.) pairs as HSPs had a minimal effect on abundance estimation, as such

instances were rare in our dataset (Appendix S1: S1.2, Figure S1); similarly, while same-cohort

full siblings were common, instances of cross-cohort full siblings were rare (Appendix S1,

Figure S2). These results demonstrate that a simple base-case CKMR model can produce

unbiased abundance estimates across a range of potential sampling scenarios when population

dynamics align with the model's assumptions while suggesting that false positive HSPs arising

from misidentified aunt/niece (uncle/nephew, etc.) pairs are likely to be rare for randomly

sampled long-lived promiscuous species.

3.2 Population growth

When we simulated a population with a trend that was growing or declining in size and

compared year-specific truths to the abundance estimates generated by our base-case CKMR

model that was naïve to a population trend (Figure 3a-c, orange), the disparity between the

quantities grew the further the estimation year (year t) was projected into the past. These

disparities were rectified by adapting our CKMR model to include a population growth model

(Eq. 3) and deriving N_t from estimates of $N_{(t0)}$ and λ (Figure 3a-c, blue; Table 2). Uncertainty

accrued as N_t was projected further from the mode of the data (Figure 3d), even with a stable

population (Appendix S1, figure S3a). When we varied $N_{(t0)}$ to generate year-specific abundance

estimates rather than deriving N_t, the model showed similar, though not identical, trends

(Appendix S1: S1.3, figure S4), suggesting that the two approaches are functionally equivalent in

most cases.

Estimates of λ were highly correlated with abundance (Table 3; Appendix S1: S1.3) and

were mostly unbiased for scenarios that involved a population that was monotonically

increasing, declining, or stable (Figure 3e); however, the model tended to underestimate λ when

the population began a severe decline during the modeled time period ("severe decline"

scenario). Estimates of $\square$ were unbiased regardless of population trend but varied with the

number of age classes sampled (Appendix S1: Figure S3b) and were highly correlated with

abundance in the scenario that only included four cohorts (target YOY; Table 3). Combined,

these results suggest that a CKMR model that is adapted for population growth can give

unbiased year-specific abundance estimates across a range of scenarios, while estimates of

population trend should be interpreted with caution (see also section 3.5 and section 4.4 below

for considerations to improve trend estimation).

3.3 Intermittent breeding

When a CKMR model that is fully naïve to intermittent breeding (Figure 4, orange) was

applied to data from populations with females that bred on a consistent multiennial schedule,

estimates of female (Figure 4a) and male (Figure 4b) abundance were positively biased. Males

did not breed on a multiennial schedule, but they did share the survival parameter ($\square$) with

females, and this parameter was also overestimated with the naïve model (Figure 4c).

When the pairwise comparison matrix was filtered to remove off-cycle comparisons

before fitting a model that was otherwise naïve to intermittent breeding (Figure 4, white),

estimates of abundance and survival were unbiased for models that only included HSPs ("sample

juveniles" and "target YOY" scenarios), but only if 100% of females bred on the same schedule.

When off-cycle breeding was introduced (biennial* scenario), estimates of female abundance

were positively biased. When the model also included kinship probabilities for parent-offspring

pairs ("sample all ages" scenario), estimates of female abundance, male abundance, and survival

were all positively biased with the "naïve – filtered" model, reflecting the fact that the two

kinship probabilities refer to different quantities (see section 4.2 and Appendix S1:S1.5).

Parameter estimates were generally unbiased with the model that was adapted for

intermittent breeding (Equation 7), including the scenario with 100% annual breeders, in which

case the naïve and adapted models performed identically (the "naïve – filtered" approach was not

tested in this scenario because with annual breeding there were no off-cycle comparisons to

remove). When we compared estimates of ψ to the realized proportion of HSPs that came from

on-cycle females (the 'true ψ' in the simulated data), estimates of ψ were unbiased when all

females bred on a multiennial cycle, with or without stochastic off-cycle breeding (Figure S5),

though we note that when the multiennial model was applied to a population that bred annually,

ψ had a non-unique solution (but estimates of other parameters were unbiased; see Appendix S1:

S1.5 for further discussion about considerations surrounding intermittent breeding). Taken

together, these results demonstrate that our model that accounts for multiennial breeding can

accommodate variable or unknown breeding schedules and produce unbiased estimates of

abundance and survival for populations that breed annually, biennially, or triennially, with or

without instances of off-cycle breeding.

3.4 Aging uncertainty

When ages were misassigned to samples, older individuals were far more likely to be

assigned to the wrong cohort (Figure 5a). The probability of age misassignment roughly

corresponded to the slope of the von Bertalanffy growth curve, with the probability of age

misassignment being greatest as the curve approached its asymptote. Consequently, when

sampling was targeted to YOY, individuals were far more likely to be assigned to the correct

cohort (Figure 5a, right).

When multiple age classes were represented in the data, bias accrued in estimates of

female abundance (Figure 5b) and survival (Figure 5c) as the CV surrounding age assignment

widened, regardless of whether we simulated a population that bred annually or biennially.

Targeted sampling of YOY showed a different trend: the probability of misassigning an age-0

individual to the wrong cohort was very low, so increasing the CV on length-based age

assignment did not affect the bias of parameter estimates. These results confirm that reliable

aging is a key component of CKMR and that targeted sampling of age classes that can be reliably

aged can improve estimation when accurate aging for other age classes is challenging.

3.5 Application to lemon sharks

3.5.1 Bimini lemon shark simulations

When we simulated a small population that resembled the population of lemon sharks at

Bimini, Bahamas, and then heavily sampled that population and used all samples for

construction of the pairwise comparison matrix, results varied depending on which window of

data was used (Figure 6 a-c). When mortality was reduced in the 90-year simulation and the

population was growing (years 70-79), year-specific estimates of female abundance, abundance

trend (λ), and survival ($\square$) were mostly unbiased for the five-year window and the scenario that

included all available samples (Figure 6a-c). As mortality increased and the population stabilized

(years 80-84), bias began to accrue for the scenario that included all available samples. Bias

continued to rise for this scenario when mortality was increased to produce a declining

population (years 85-90). In contrast, results from the five-year window of samples were

unbiased whenever the sample window spanned a period that included a consistent population

trend: each time a shift in population trend occurred (years 80 and 85), the five-year window of

samples produced biased parameter estimates for the following year, but that bias was reduced as

the population trend stabilized. The three-year window of samples produced parameter estimates

that were both biased and imprecise, suggesting that more than three cohorts are needed to

produce reliable parameter estimates for a population that breeds biennially. All sample windows

estimated abundance at an absolute scale within an order of magnitude of the real population size

(see Appendix S1, Figure S6 for an illustrative example).

When we retained just one full sibling from each mother/father pairing, parameter

estimates showed similar tendencies as when the full dataset was used, with the exception that

estimates of λ were less biased for the three-year window (see Appendix S1: S1.6; Figure S7 a-

c). Combined, these results demonstrate that CKMR can reliably estimate abundance within an

order of magnitude for a small population that is heavily sampled, while inference of trends –

whether through a time-series of female abundance or estimation of λ – varies depending on the

window of samples included.

3.5.2 Bimini lemon shark data

Application of our multiennial CKMR model to real data from Bimini lemon sharks

showed a parabolic abundance trend and very low abundance regardless of whether we retained

full siblings (Figure 6d, Table 4) or filtered them (Appendix S1: Figure S7d, Table S3), and

whether the datasets were downsampled (Appendix S1: Table S4, Table S5, Figure S8) or not.

These trends were consistent regardless of which window of samples was used, though the

timing of peak abundance varied (see Appendix S1: S1.6 for more discussion). Estimates of

female breeders ($N_{\female b}$) were close to, but slightly higher than, estimates independently obtained

by DiBattista et al. (2011) using a pedigree-based approach.

Estimates of adult survival (ϕ) were generally high for the full and downsampled datasets,

but varied with the sample window (Appendix S1: Figure S9) and showed more variability when

the dataset was both filtered for full siblings and downsampled (Appendix S1: Figure S8b). The

three-year window of samples gave rise to survival estimates that were highly correlated with

estimates of abundance (Table 5) and did not vary across years (Appendix S1, Figure S9), again

suggesting that the three-year window is too short a time period for reliable estimation of

survival rates for populations that breed biennially. Overall, regardless of how the data were

subset, our results align with other studies that suggest low abundance (DiBattista et al. 2011)

and high survival rates (White et al. 2014) of adult females at Bimini; however, we also note that

estimation of abundance trend and survival were correlated with the number of cohorts included

in the analysis and the mortality regimes that the population experienced.

4. Discussion

Obtaining unbiased estimates of abundance is a central challenge for effective

conservation and management of many threatened and exploited populations and is especially

pertinent for populations of low density and highly mobile species where effective sampling of

adults is impractical. Our simulation results broadly concur with recent work supporting CKMR

as a promising approach to estimate abundance and survival in data-limited circumstances, but

emphasize the critical need to adapt CKMR models adequately to accommodate population

dynamics and life history traits that violate the assumptions of a simple base-case model. Further,

although we confirm sensitivity of CKMR to aging error, we also find that bias in parameter

estimates can be mitigated by sampling as few as four cohorts that can be reliably aged,

providing options for applying the method when accurate aging is difficult or when long-term

sampling is impractical. Our application to lemon sharks in Bimini, Bahamas demonstrates that

CKMR is a flexible framework that can be used to estimate abundance and survival of breeding

adults when only juveniles are available for sampling. Taken together, the results of our

application of CKMR to simulated and real populations with different population sizes, trends,

and breeding schedules support the recognition of CKMR's immense potential for monitoring

populations of low density and highly mobile species, while also highlighting several promising

avenues for future research.

4.1   Accounting for population growth/decline

A simple base-case CKMR model (e.g., Eqs. 1 and 2) estimates adult abundance over the

modeled period by assuming that population dynamics are stable and consistent over time. In

cases with sex-specific or transient population dynamics, or if estimates of underlying population

parameters are desired, population dynamics can be modeled with CKMR using latent variables.

Alternatively, year-specific abundance estimates can be obtained by modeling each year

independently, but this approach requires a particularly rich dataset (e.g., heavily sampled

salmonids; Ruzzante et al. 2019, Marcy☐Quay et al. 2020). As such, most data-limited situations

will likely benefit from leveraging all available data for a single abundance estimate. In such

cases, specifying an exponential growth model where population dynamics are broadly captured

in a parameter like λ allows data to be shared across cohorts to produce a single estimate of

abundance for a specified reference year (t_0). Then, abundance in any modeled year can be

derived from estimates of $N_{(t0)}$ and λ. In practical applications of CKMR, knowledge of a

species' life history in combination with Leslie matrix simulations can help inform a prior on λ to

improve precision of parameter estimates. If a fishery-independent index or fishery-dependent

CPUE index (from a fishery whose operations have been relatively constant) exists over the

modeled period, then trend data could also be integrated into the model via specification of the

prior on λ.

Including too many age classes in the data may hinder inference of abundance trends if a

population's trajectory shifts during the modeled time period, as estimates of λ will represent an

average of those trajectories. One way to mitigate this averaging effect is to subset the dataset for

smaller time windows over which averaging λ has a less pronounced effect. This approach

sacrifices precision and may not be possible in many data limited situations; even in a data-rich

scenario, the time window must be thoughtfully calibrated, as using too small a window provides

limited data to the exponent on $\square$, resulting in a high degree of correlation between $\square$ and N

(e.g., the target YOY scenario in Table 3 and the three-year window in Table 5). On the other

hand, our results suggest that using too large a window will result in an abundance trend that lags

behind the real trend if the real trend shifts during the modeled period. For our Bimini lemon

shark simulations, a five-year window produced a reasonable balance of precision and accuracy

for parameter estimates and also for reconstructing an abundance trend for a population that

experienced multiple mortality regimes during the modeled period. In application to real data

sets, one would not know the accuracy associated with a given window-size, but the pattern of

lag in estimated population trends as the window length is expanded may give some indication of

where shifts in the trajectory have occurred. Alternatively, one could test for a quadratic trend by

adding another parameter in the model for λ and then use model selection to determine the model

that best captures the true population trend. Expanding CKMR to estimate population trend in a

reliable and robust way is a ripe area for future research.

4.2   Intermittent breeding dynamics

When intermittent breeding coincides with a population that is most easily sampled

during the juvenile life stage (e.g., when adults are not directly observed), our results indicate

that abundance estimates derived from a naïve half-sibling CKMR model will be biased if all

pairwise comparisons are included in the model. In contrast, in circumstances where all

individuals breed on the same schedule such that there is no possibility of off-cycle breeding,

then filtering the pairwise comparison matrix to remove off-cycle comparisons and fitting a half-

sibling model that is otherwise naïve to intermittent breeding (e.g., Eq. 4) can give unbiased

parameter estimates (Figure 4, white). Importantly, when off-cycle comparisons are removed, the

non-breeding adults essentially become invisible to half-sibling CKMR (similar to infertile

adults; see section 3.2 of Bravington et al. 2016a). As such, a naïve HS model that is filtered to

remove off-cycle comparisons gives estimates of the number of effective female breeders ($\widetilde{N}$) in

year t (Appendix S1: S1.4), a number that may be substantially different than the total number of

adult females (N) in populations that breed intermittently. Without modification to the kinship

probabilities, this difference precludes the use of POPs in the same model because PO CKMR

gives estimates of N (note the bias for the "sample all ages" scenario in Figure 4). In addition,

our simulations suggest that if off-cycle breeding produces HSPs that are separated by a birth

year gap that does not align with the expected breeding cycle (e.g., when the birth year gap is

odd for a population that breeds on a biennial schedule), then parameter estimates will be biased

with a naïve model, whether off-cycle comparisons are retained in the pairwise comparison

matrix or not. Overall, applying a model that is naïve to intermittent breeding to a population that

breeds on a bi- or triennial schedule can produce unbiased parameter estimates if off-cycle

comparisons are removed, but only in limited situations (e.g., when there is no off-cycle

breeding).

The multiennial CKMR model presented here accommodates intermittent breeding via

inclusion of the parameters Ψ and a, which assigns a non-zero probability to off-cycle

comparisons without assuming the probability is the same as on-cycle comparisons, resulting in

estimates of N rather than $\widetilde{N}$ (see Appendix S1, S1.4 and S1.5 for further discussion). While the

parameter Ψ can be estimated, a must be fixed to the expected breeding cycle. If the breeding

cycle for a population is unknown, and if adults are not available for sampling, then it may be

possible to estimate a from the distribution of birth year gaps among identified half-siblings

(Waples and Feutry 2021). As a cursory example, if most HSPs were born in year gaps that are

divisible by 2, then fixing a to 2 would be logical. However, in real populations, reproductive

periodicity may be challenging to infer from the distribution of kin pairs, as environmental

conditions may cause individuals to fail to breed one year and then breed off-cycle the next

(Rivalan et al. 2005, Cubaynes et al. 2011, Morbey and Shuter 2013, Öst et al. 2018, Skjæraasen

et al. 2020). In our lemon shark dataset, for example, 30% of adult mothers bred off-cycle at

least once before returning to a biennial cycle. With elasmobranchs and other species that are

difficult to age, aging error will further obscure inference of breeding schedule based on

offspring birth years. Stochastic off-cycle breeding was not a problem for our multiennial model

as long as there were no systemic differences in lifetime fecundity (Bravington et al. 2016a; see

Appendix S1:S1.6 and Figure S10). Future work that adapts CKMR to estimate Ψ and a across a

range of scenarios, including populations with mixed mating schedules (Walker 2007, Driggers

and Hoffmayer 2009, Higgs et al. 2020), would further expand the potential of CKMR to

illuminate aspects of population breeding dynamics.

4.3   Aging error

CKMR depends heavily on accurate cohort assignment, which can be very challenging

for many species, including elasmobranchs. Our results confirm that age misassignment can

substantially bias CKMR parameter estimates. A hierarchical model that accounts for aging error

may help alleviate this issue, but such a model would require some estimate of the probability of

age misassignment (Hirst et al. 2004, Schwarz and Runge 2009) and selectivity (Henríquez et al.

2016, Francis 2016), and such data may not be available in data-limited situations. Estimating

the probability of age misassignment is not a trivial task, even for species with well-established

aging methods (e.g., teleosts, O'Sullivan 2007). For species that are challenging to age,

substantial upfront effort may be required to calibrate aging methods and estimate the degree of

error present. For example, patterns of DNA methylation can be used to estimate age (Jarman et

al. 2015) and these data can be obtained from the same tissue samples used for kinship

assignment in CKMR. However, epigenetic clocks are taxa-specific, and the discovery of

informative biomarkers requires calibration using representative samples of known ages, which

may be arduous to obtain in their own right (Polanowski et al. 2014, Beal et al. 2022). It is wise,

therefore, to consider how samples will be aged and how much error there is likely to be prior to

embarking on a large-scale CKMR study.

In cases where only YOY can be reliably aged, our results show that CKMR can generate

reliable abundance estimates from targeted sampling of as few as four cohorts of YOY, even for

a population that breeds bi- or triennially, though estimates of survival will improve as more

cohorts are added. If mature individuals are also available to sample – e.g., when visiting a

nursery site to breed – then sampling potential parents as well as YOY can enable the use of

POPs in the likelihood and improve precision of parameter estimates. Aging error in this case

would be less critical for adults as long as maturity can be confirmed in the year of sampling,

though care must be taken to ensure that potential parents and offspring are sampled

independently, as parameter estimates will be biased if the probability of sampling a parent is

correlated with the probability of sampling its offspring (Bravington et al. 2016a).

4.4   Population dynamics and abundance of lemon sharks in Bimini

Our application of CKMR to Bimini lemon sharks highlights the flexibility and potential of

CKMR for long-term monitoring of populations of low density highly mobile species with

geographically distinct life histories. Estimates of abundance from CKMR suggest that a very

small number of female lemon sharks give birth at the North Bimini Lagoon during each biennial

breeding cycle (Figure 6d, Table 4). These results align with a previous study that reconstructed

a pedigree for the population and identified the number of adults that successfully bred at the

North Island each year between 1995-2007 (DiBattista et al. 2011). In both cases, the number of

females that gave birth at the North Island during this time period was estimated to be very small

(<50 per year), with an increasing abundance trend through ~2006. At some point after or around

2006, results from CKMR suggest that the number of females using Bimini for breeding began to

decline. Intense dredging and mangrove deforestation took place around the North Bimini Island

in March 2001 in preparation for development of a mega-resort (Jennings et al. 2008). Although

the number of breeding females at the North Island counterintuitively increased immediately

after the disturbance (DiBattista et al. 2011), there was a transient drop in the survival rates of

age 0 and age 1 individuals, though the degree to which juvenile mortality was affected is

debated (Jennings et al. 2008, DiBattista et al. 2011). These cohorts would have reached maturity

and begun returning to Bimini for reproduction around 2011, which may explain the decreasing

trend around that time (Figure 6d). All sampling windows and methods of downsampling

showed a parabolic abundance trend over the time series, though the stationary point (where

population size was stable) varied depending on the window.

Although our results closely resemble those reported in DiBattista et al. (2011), we note that

our abundance estimates from CKMR were slightly higher. The degree to which our results

differed depended on whether we included full siblings in the analysis and whether we used the

full dataset or a downsampled dataset (see Appendix S1: S1.6 for more discussion on CKMR

with small populations). Abundance estimates were generally similar (<50 breeding females)

across the datasets we tested, except for a few instances when sampling was constrained to a

three-year window. When a population breeds biennially, sampling three years only includes one

year gap with possible positive comparisons (years one and three), which provides very limited

information to the exponent on $\square$ and impedes its estimation (see Appendix S1: Figure S9), as

well as the estimation of other parameters (note the high correlation with abundance in Table 5).

Though all three windows of samples we tested (three-year, five-year, all available) suggest

the population of breeding females at Bimini is small, the five-year window produced the least

biased parameter estimates in simulation, and application to the real data resulted in estimates

that aligned more closely with the estimates of Dibattista et al. (2011) than when all samples

were used. More complex models (e.g., that allow for quadratic abundance trends) would likely

improve performance of models that leverage long time-series of data; in the absence of such a

model, calibrating the time window of samples to correspond to likely abundance trends can help

alleviate the averaging effect of assuming exponential growth/decline over long time periods.

Adapting CKMR to produce more reliable parameter estimates for small populations will

require additional work (Bravington et al. 2016a); however, when dealing with abundance

estimates that are small enough to cause such issues, the practical implications of this known bias

are likely minimal. Discarding estimates from the three-year window of samples, our model that

retained full siblings in the dataset estimated a maximum of 47 females visited the North Island

across all 18 years of abundance estimation (Table 4, Figure 6d). Removing full siblings and

downsampling the dataset produced slightly higher values but still did not exceed 70 females

(Appendix S1: Figure S7, Table S3). These quantities are small enough that any additional

mortality would likely threaten the sustainability of this portion of the population.

4.5 Implications for sampling design

We have shown that application of HS CKMR to long-lived species can generate reliable

estimates of abundance – on its own, or in conjunction with PO CKMR - from a limited number

of cohorts when aging is reliable; however, estimates of survival (ϕ) were more reliable when

more cohorts were included in the dataset across our simulations. A dataset that spans enough

cohorts to reliably estimate parameters beyond abundance can be obtained by intensely sampling

multiple age classes over a small number of years, or by long-term sampling of nursery areas.

The former would require reliable aging of all sampled age classes to avoid biased parameter

estimates, especially for models that incorporate half-sibling kinship probabilities where

estimates of survival and abundance both depend on the birth year of sampled individuals. The

latter - long-term sampling of nursery areas - represents a promising method for monitoring low

density highly mobile populations, especially in circumstances where aging error is likely for

older age classes.

We are not the first to suggest that CKMR benefits from focusing sampling efforts on

individuals that can be reliably aged (Trenkel et al. 2022). Our results expand on this idea by

demonstrating that CKMR can produce robust abundance estimates from as few as four cohorts,

though estimates of survival will be less reliable as fewer age classes are included. In cases

where sampling of juveniles is focused on nursery areas, sufficient biological knowledge to

determine the scope of inference for CKMR will be required; if the target population uses

multiple nursery areas, then sampling multiple nurseries can allow the model to estimate

demographic connectivity (Patterson et al. 2022a). If sex-specific population dynamics are

present, as with Bimini lemon sharks, the associated CKMR model should account for this and

estimate parameters separately for each sex or focus solely on the sex for which the scope of

inference is well-understood, as we did with Bimini lemon sharks.

One of the more exciting aspects of CKMR is its potential to generate rapid estimates of

adult abundance without sampling a single adult (see Patterson et al. 2022a for an applied

example). Our results confirm that a sampling program that can procure as few as four or five

reliably aged cohorts can be used in combination with half-sibling CKMR to produce robust

estimates of present-day abundance as well as reasonable estimates of survival. In circumstances

where a genotyping panel, workflow for assigning kinship, and appropriate CKMR model are

already developed for a population, contemporary abundance estimates could conceivably be

obtained within weeks of sampling. As such, CKMR can offer a rapid and cost-effective method

for population monitoring in real time following an initial investment in the laboratory and

analytical workflows.

Conclusion and Future Directions

CKMR is a powerful tool for estimating population abundance of species that have been

historically difficult to assess. Reliable application of the method requires careful consideration

of the relevant population dynamics matched to an appropriate sampling scheme. Here, we have

identified a set of factors that must be considered for robust application of CKMR, proposed

methods for accounting for them, and highlighted areas in need of further research. Specifically,

we found that a half-sibling focused CKMR model can produce robust abundance estimates from

as few as four or five cohorts, while reliable estimates of survival will likely require more data.

Monotonic abundance trends can be dependably inferred by incorporating a simple exponential

growth model; however, more complex trends will require further model development or, at a

minimum, deployment of a sliding window of samples, which prevents long-term averaging of λ

and obfuscation of transient dynamics.

When ages are prone to misassignment, focusing sampling efforts on individuals with

known ages (e.g., YOY), or subsampling for these individuals if the dataset is sufficiently rich,

can alleviate bias in parameter estimates, particularly abundance. Long-term monitoring of

highly mobile species can be enhanced by CKMR via sampling of nursery areas when one or

both sexes are philopatric, and can provide estimates of present-day abundance and abundance

trends for adults that visit the nursery area without directly sampling a single adult. Overall, this

study highlights the sensitivity of simple base-case CKMR models to assumptions about

population dynamics and sampling, while also demonstrating that the CKMR framework is

easily adaptable to accommodate these factors, making it a promising tool for integration into

long-term monitoring programs.

Acknowledgements

The authors thank Paul Conn and Shane Baylis for early guidance with regards to CKMR model

construction and individual-based simulation during the initial phase of the project, and Andy

Lin and Patrick Sadil for assistance with code that was integrated into construction of the

pairwise comparison matrix. We are grateful to Chris Sutherland for invaluable support during

model validation, and to Mark Bravington, Mark Maunder, Kevin Piner, and Steve Teo for

conversations and suggestions that guided our sensitivity tests. The authors are exceedingly

grateful to Paul Conn and an anonymous reviewer for insightful comments on a prior version of

this manuscript, and members of the Molecular Ecology and Conservation Lab at UMass

Amherst provided helpful feedback on earlier drafts of this manuscript. Thanks to the hundreds

of volunteers and staff at the Bimini Biological Field Station that sampled and tagged lemon

sharks. This work was completed using resources provided by the University of Massachusetts'

Green High Performance Computing Cluster (GHPCC). Funding for this project was provided to

JDS by the American Fisheries Society via the Steven Berkeley Conservation Fellowship, the

Save Our Seas Foundation, and the UMass Amherst Organismic and Evolutionary Biology

program.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

References

Ailloud, L. E., M. W. Smith, A. Y. Then, K. L. Omori, G. M. Ralph, and J. M. Hoenig. 2015. Properties

of age compositions and mortality estimates derived from cohort slicing of length data. ICES

Journal of Marine Science 72:44–53.

Amstrup, S. C., T. L. McDonald, and B. F. Manly. 2010. Handbook of capture-recapture analysis.

Princeton University Press.

Arnason, A. N., and K. H. Mills. 1981. Bias and Loss of Precision Due to Tag Loss in Jolly–Seber

Estimates for Mark–Recapture Experiments. Canadian Journal of Fisheries and Aquatic Sciences

38:1077–1095.

Bauwens, D., and K. Claus. 2019. Intermittent reproduction, mortality patterns and lifetime breeding

frequency of females in a population of the adder (*Vipera beru* s). PeerJ 7:e6912.

Beal, A. P., S. Hackerott, K. Feldheim, S. H. Gruber, and J. M. Eirin-Lopez. 2022. Age group DNA

methylation differences in lemon sharks (Negaprion brevirostris): Implications for future age

estimation tools. Ecology and Evolution 12:e9226.

Berryman, A. A. 1989. The Conceptual Foundations of Ecological Dynamics. Bulletin of the Ecological

Society of America 70:230–236.

Boyd, C., J. Barlow, E. A. Becker, K. A. Forney, T. Gerrodette, J. E. Moore, and A. E. Punt. 2018.

Estimation of population size and trends for highly mobile species with dynamic spatial

distributions. Diversity and Distributions 24:1–12.

Boyd, C., and A. E. Punt. 2021. Shifting trends: Detecting changes in cetacean population dynamics in

shifting habitat. PLOS ONE 16:e0251522.

Bradford, R., R. Thomson, M. Bravington, D. Foote, R. M. Gunasekera, B. Bruce, D. Harasti, N. Otway,

and P. Feutry. 2018. A close-kin mark-recapture estimate of the population size and trend of east

coast grey nurse shark.

Bravington, M., P. Feutry, R. D. Pillans, R. Hillary, G. Johnson, T. M. Saunders, R. M. Gunasekera, N.

Bax, and P. M. Kyne. 2019. Close-Kin Mark-Recapture population size estimate of *Glyphis*

garricki in the Northern Territory.

Bravington, M. V., H. J. Skaug, and E. C. Anderson. 2016a. Close-Kin Mark-Recapture. Statistical

Science 31:259–274.

Bravington, M. V., P. M. Grewe, and C. R. Davies. 2016b. Absolute abundance of southern bluefin tuna

estimated by close-kin mark-recapture. Nature Communications 7:13162.

Brooks, J. L., T. L. Guttridge, B. R. Franks, R. D. Grubbs, D. D. Chapman, S. H. Gruber, J. D. Dibattista,

and K. A. Feldheim. 2016. Using genetic inference to re-evaluate the minimum longevity of the

lemon shark Negaprion brevirostris. Journal of Fish Biology 88:2067–2074.

Brown, C. A., and S. H. Gruber. 1988. Age Assessment of the Lemon Shark, Negaprion brevirostris,

Using Tetracycline Validated Vertebral Centra. Copeia 1988:747–753.

Cailliet, G. M. 2015. Perspectives on elasmobranch life-history studies: a focus on age validation and

relevance to fishery management. Journal of Fish Biology 87:1271–1292.

Campbell, R. A. 2015. Constructing stock abundance indices from catch and effort data: Some nuts and

bolts. Fisheries Research 161:109–130.

Carbone, C., N. Pettorelli, and P. A. Stephens. 2011. The bigger they come, the harder they fall: body size

and prey abundance influence predator–prey ratios. Biology Letters 7:312–315.

Carlson, J., P. Charvet, A. Ba, J. Bizzarro, D. Derrick, M. Espinoza, P. Doherty, E. Chartrain, M. Dia, and

G. H. L. Leurs. 2021. Negaprion Brevirostris. The IUCN Red List of Threatened Species:2021–1.

Chapman, D. D., E. A. Babcock, S. H. Gruber, J. D. Dibattista, B. R. Franks, S. A. Kessel, T. Guttridge,

E. K. Pikitch, and K. A. Feldheim. 2009. Long-term natal site-fidelity by immature lemon sharks

(Negaprion brevirostris) at a subtropical island. Molecular Ecology 18:3500–3507.

Conn, P. B., M. V. Bravington, S. Baylis, and J. M. Ver Hoef. 2020. Robustness of close‐kin mark–

recapture estimators to dispersal limitation and spatially varying sampling probabilities. Ecology

and Evolution 10:5558–5569.

Cormack, R. M. 1964. Estimates of Survival from the Sighting of Marked Animals. Biometrika 51:429–

438.

Cortés, E., and E. N. Brooks. 2018. Stock status and reference points for sharks using data-limited

methods and life history. Fish and Fisheries 19:1110–1129.

Cubaynes, S., P. F. Doherty, E. A. Schreiber, and O. Gimenez. 2011. To breed or not to breed: a seabird's

response to extreme climatic events. Biology Letters 7:303–306.

Davis, K. L., E. D. Silverman, A. L. Sussman, R. R. Wilson, and E. F. Zipkin. 2022. Errors in aerial

survey count data: Identifying pitfalls and solutions. Ecology and Evolution 12:e8733.

Delaval, A., V. Bendall, S. J. Hetherington, H. J. Skaug, M. Frost, C. S. Jones, and L. R. Noble. 2022.

Evaluating the suitability of close☐kin mark☐recapture as a demographic modelling tool for a

critically endangered elasmobranch population. Evolutionary Applications.

Denwood, M. J. 2016. runjags: An R Package Providing Interface Utilities, Model Templates, Parallel

Computing Methods and Additional Distributions for MCMC Models in JAGS. Journal of

Statistical Software 71:1–25.

Desprez, M., O. Gimenez, C. R. McMahon, M. A. Hindell, and R. G. Harcourt. 2018. Optimizing lifetime

reproductive output: Intermittent breeding as a tactic for females in a long-lived, multiparous

mammal. Journal of Animal Ecology 87:199–211.

DiBattista, J. D., K. A. Feldheim, D. Garant, S. H. Gruber, and A. P. Hendry. 2011. Anthropogenic

disturbance and evolutionary parameters: a lemon shark population experiencing habitat loss.

Evolutionary Applications 4:1–17.

Driggers, W. B., and E. R. Hoffmayer. 2009. Variability in the Reproductive Cycle of Finetooth Sharks,

Carcharhinus isodon, in the Northern Gulf of Mexico. Copeia 2009:390–393.

Dulvy, N. K., N. Pacoureau, C. L. Rigby, R. A. Pollom, R. W. Jabado, D. A. Ebert, B. Finucci, C. M.

Pollock, J. Cheok, D. H. Derrick, K. B. Herman, C. S. Sherman, W. J. VanderWright, J. M.

Lawson, R. H. L. Walls, J. K. Carlson, P. Charvet, K. K. Bineesh, D. Fernando, G. M. Ralph, J.

H. Matsushiba, C. Hilton-Taylor, S. V. Fordham, and C. A. Simpfendorfer. 2021. Overfishing

drives over one-third of all sharks and rays toward a global extinction crisis. Current Biology

31:4773-4787.e8.

Ellegren, H., and N. Galtier. 2016. Determinants of genetic diversity. Nature Reviews Genetics 17:422–

433.

Erisman, B. E., L. G. Allen, J. T. Claisse, D. J. Pondella, E. F. Miller, and J. H. Murray. 2011. The

illusion of plenty: hyperstability masks collapses in two recreational fisheries that target fish

spawning aggregations. Canadian Journal of Fisheries and Aquatic Sciences 68:1705–1716.

Feldheim, K. A., S. H. Gruber, and M. V. Ashley. 2002. The breeding biology of lemon sharks at a

tropical nursery lagoon. Proceedings of the Royal Society B: Biological Sciences 269:1655–1661.

Feldheim, K. A., S. H. Gruber, and M. V. Ashley. 2004. Reconstruction of Parental Microsa□lite

Genotypes Reveals Female Polyandry and Philopatry in the Lemon Shark, Negaprion

Brevirostris. Evolution 58:2332–2342.

Feldheim, K. A., S. H. Gruber, J. D. DiBattista, E. A. Babcock, S. T. Kessel, A. P. Hendry, E. K. Pikitch,

M. V. Ashley, and D. D. Chapman. 2014. Two decades of genetic profiling yields first evidence

of natal philopatry and long-term fidelity to parturition sites in sharks. Molecular Ecology

23:110–117.

Feldheim, K., A. Fields, D. Chapman, R. Scharer, and G. Poulakis. 2017. Insights into reproduction and

behavior of the smalltooth sawfish *Pristis pectinata*. Endangered Species Research 34:463–471.

Ferretti, F., B. Worm, G. L. Britten, M. R. Heithaus, and H. K. Lotze. 2018. Patterns and ecosystem

consequences of shark declines in the ocean. Ecology Letters:1055–1071.

Feutry, P., O. Berry, P. M. Kyne, R. D. Pillans, R. M. Hillary, P. M. Grewe, J. R. Marthick, G. Johnson,

R. M. Gunasekera, N. J. Bax, and M. Bravington. 2017. Inferring contemporary and historical

genetic connectivity from juveniles. Molecular Ecology 26:444–456.

Førland, B. 2019. Close-Kin Mark-Recapture Models:80.

Francis, R. I. C. C. 2016. Growth in age-structured stock assessment models. Fisheries Research 180:77–

86.

Green, A. W., G. C. Matlock, and J. E. Weaver. 1983. A Method for Directly Estimating the Tag-

Reporting Rate of Anglers. Transactions of the American Fisheries Society 112:412–415.

Guttridge, T. L., M. P. M. Van Zinnicq Bergmann, C. Bolte, L. A. Howey, J. S. Finger, S. T. Kessel, J. L.

Brooks, W. Winram, M. E. Bond, L. K. B. Jordan, R. C. Cashman, E. R. Tolentino, R. D. Grubbs,

and S. H. Gruber. 2017. Philopatry and Regional Connectivity of the Great Hammerhead Shark,

Sphyrna mokarran in the U.S. and Bahamas. Frontiers in Marine Science 4.

Gwinn, D. C., M. S. Allen, and M. W. Rogers. 2010. Evaluation of procedures to reduce bias in fish

growth parameter estimates resulting from size-selective sampling. Fisheries Research 105:75–

79.

Hansell, A. C., T. H. Curtis, J. Carlson, E. Cortés, G. Fay, and S. X. Cadrin. 2021. Stock Assessment of

the Lemon Shark off the Southeast United States. North American Journal of Fisheries

Management 41:35–48.

Hansell, A. C., S. T. Kessel, L. R. Brewster, S. X. Cadrin, S. H. Gruber, G. B. Skomal, and T. L.

Guttridge. 2018. Local indicators of abundance and demographics for the coastal shark

assemblage of Bimini, Bahamas. Fisheries Research 197:34–44.

Harley, S. J., R. A. Myers, and A. Dunn. 2001. Is catch-per-unit-effort proportional to abundance?

Canadian Journal of Fisheries and Aquatic Sciences 58:1760–1772.

Hassell, M. P. 1975. Density-Dependence in Single-Species Populations. Journal of Animal Ecology

44:283–295.

Henríquez, V., R. Licandeo, L. A. Cubillos, and S. P. Cox. 2016. Interactions between ageing error and

selectivity in statistical catch-at-age models: simulations and implications for assessment of the

Chilean Patagonian toothfish fishery. ICES Journal of Marine Science 73:1074–1090.

Heupel, M., J. Carlson, and C. Simpfendorfer. 2007. Shark nursery areas: concepts, definition,

characterization and assumptions. Marine Ecology Progress Series 337:287–297.

Higgs, J. M., E. R. Hoffmayer, J. A. Sulikowski, W. B. Driggers, D. A. Stiller, J. M. Hendon, J. M.

Higgs, E. R. Hoffmayer, J. A. Sulikowski, W. B. Driggers, D. A. Stiller, and J. M. Hendon. 2020.

Reproductive biology of the finetooth shark (Carcharhinus isodon) in the northern Gulf of

Mexico, with evidence of both annual and biennial reproduction. Marine and Freshwater

Research 72:693–708.

Hillary, R. M., M. V. Bravington, T. A. Patterson, P. Grewe, R. Bradford, P. Feutry, R. Gunasekera, V.

Peddemors, J. Werry, M. P. Francis, C. a. J. Duffy, and B. D. Bruce. 2018. Genetic relatedness

reveals total population size of white sharks in eastern Australia and New Zealand. Scientific

Reports 8:2661.

Hirst, D., S. Aanes, G. Storvik, R. B. Huseby, and I. F. Tvete. 2004. Estimating catch at age from market

sampling data by using a Bayesian hierarchical model. Journal of the Royal Statistical Society:

Series C (Applied Statistics) 53:1–14.

Hyun, S.-Y., J. H. Reynolds, and P. F. Galbreath. 2012. Accounting for Tag Loss and Its Uncertainty in a

Mark–Recapture Study with a Mixture of Single and Double Tags. Transactions of the American

Fisheries Society 141:11–25.

Jacoby, D. M. P., D. P. Croft, and D. W. Sims. 2012. Social behaviour in sharks and rays: analysis,

patterns and implications for conservation. Fish and Fisheries 13:399–417.

Jarman, S. N., A. M. Polanowski, C. E. Faux, J. Robbins, R. De Paoli-Iseppi, M. Bravington, and B. E.

Deagle. 2015. Molecular biomarkers for chronological age in animal ecology. Molecular Ecology

24:4826–4847.

Jennings, D. E., S. H. Gruber, B. R. Franks, S. T. Kessel, and A. L. Robertson. 2008. Effects of large-

scale anthropogenic development on juvenile lemon shark (Negaprion brevirostris) populations of

Bimini, Bahamas. Environmental Biology of Fishes 83:369–377.

Jennings, S. 2000. Patterns and prediction of population recovery in marine reserves. Reviews in Fish

Biology and Fisheries 10:209–231.

Jolly, G. M. 1965. Explicit Estimates from Capture-Recapture Data with Both Death and Immigration-

Stochastic Model. Biometrika 52:225–247.

Kendall, W. L. 1999. Robustness of Closed Capture–Recapture Methods to Violations of the Closure

Assumption. Ecology 80:2517–2525.

Kindsvater, H. K., N. K. Dulvy, C. Horswill, M.-J. Juan-Jordá, M. Mangel, and J. Matthiopoulos. 2018.

Overcoming the Data Crisis in Biodiversity Conservation. Trends in Ecology & Evolution

33:676–688.

Kohler, N. E., and P. A. Turner. 2001. Shark tagging: a review of conventional methods and studies. The

behavior and sensory biology of elasmobranch fishes: an anthology in memory of Donald

Richard Nelson:191–224.

Lynch, P. D., K. W. Shertzer, and R. J. Latour. 2012. Performance of methods used to estimate indices of

abundance for highly migratory species. Fisheries Research 125–126:27–39.

Magera, A. M., J. E. M. Flemming, K. Kaschner, L. B. Christensen, and H. K. Lotze. 2013. Recovery

Trends in Marine Mammal Populations. PLOS ONE 8:e77908.

Marcy☐Quay, B., S. A. Sethi, N. O. Therkildsen, and C. E. Kraft. 2020. Expanding the feasibility of fish

and wildlife assessments with close-kin mark–recapture. Ecosphere 11:e03259.

Maunder, M. N., and A. E. Punt. 2004. Standardizing catch and effort data: a review of recent approaches.

Fisheries Research 70:141–159.

Maunder, M., J. Sibert, A. Fonteneau, J. Hampton, P. Kleiber, and S. Harley. 2006. Interpreting catch per

unit effort data to assess the status of individual stocks and communities. ICES Journal of Marine

Science 63:1373–1385.

McCauley, D. J., K. A. McLean, J. Bauer, H. S. Young, and F. Micheli. 2012. Evaluating the

performance of methods for estimating the abundance of rapidly declining coastal shark

populations. Ecological Applications 22:385–392.

Morbey, Y. E., and B. J. Shuter. 2013. Intermittent breeding in the absence of a large cost of

reproduction: evidence for a non-migratory, iteroparous salmonid. Ecosphere 4:art150.

Morrissey, J. F., and S. H. Gruber. 1993. Home Range of Juvenile Lemon Sharks, Negaprion brevirostris.

Copeia 1993:425–434.

Nosal, A. P., D. P. Cartamil, A. J. Ammann, L. F. Bellquist, N. J. Ben-Aderet, K. M. Blincow, E. S.

Burns, E. D. Chapman, R. M. Freedman, A. P. Klimley, R. K. Logan, C. G. Lowe, B. X.

Semmens, C. F. White, and P. A. Hastings. 2021. Triennial migration and philopatry in the

critically endangered soupfin shark Galeorhinus galeus. Journal of Applied Ecology 58:1570–

1582.

Oosthuizen, W. C., P. J. N. De Bruyn, M. N. Bester, and M. Girondot. 2010. Cohort and tag-site-specific

tag-loss rates in mark–recapture studies: A southern elephant seal cautionary case. Marine

Mammal Science 26:350–369.

Öst, M., A. Lindén, P. Karell, S. Ramula, and M. Kilpi. 2018. To breed or not to breed: drivers of

intermittent breeding in a seabird under increasing predation risk and male bias. Oecologia

188:129–138.

O'Sullivan, S. 2007. Fisheries long term monitoring program: Fish Age Estimation Review. State of

Queensland. Department of Primary Industries and Fisheries, Brisbane, Australia.

Patterson, T. A., R. M. Hillary, P. M. Kyne, R. D. Pillans, R. M. Gunasekera, J. R. Marthick, G. J.

Johnson, and P. Feutry. 2022a. Rapid assessment of adult abundance and demographic

connectivity from juvenile kin pairs in a critically endangered species. Science Advances

8:eadd1679.

Patterson, T. A., R. Hillary, P. Feutry, R. Gunasakera, J. Marthick, and R. D. Pillans. 2022b, February 27.

Rapid estimation of cryptic adult abundance and breeding dynamics in a critically endangered

elasmobranch from close-kin mark recapture. *bioRxiv*.

Peterson, C. D., M. J. Wilberg, E. Cortés, and R. J. Latour. 2021. Dynamic factor analysis to reconcile

conflicting survey indices of abundance. ICES Journal of Marine Science 78:1711–1729.

Polanowski, A. M., J. Robbins, D. Chandler, and S. N. Jarman. 2014. Epigenetic estimation of age in

humpback whales. Molecular Ecology Resources 14:976–987.

Pollock, K. H. 2000. Capture-Recapture Models. Journal of the American Statistical Association 95:293–

296.

Pollock, K. H., J. M. Hoenig, W. S. Hearn, and B. Calingaert. 2001. Tag Reporting Rate Estimation: 1.

An Evaluation of the High-Reward Tagging Method. North American Journal of Fisheries

Management 21:521–532.

R Core Team 2021. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Rivalan, P., A.-C. Prévot-Julliard, R. Choquet, R. Pradel, B. Jacquemin, and M. Girondot. 2005. Trade-

off between current reproductive effort and delay to next reproduction in the leatherback sea

turtle. Oecologia 145:564–574.

Robertson, D. R. 1996. Interspecific Competition Controls Abundance and Habitat Use of Territorial

Caribbean Damselfishes. Ecology 77:885–899.

Rowat, D., M. G. Meekan, U. Engelhardt, B. Pardigon, and M. Vely. 2007. Aggregations of juvenile

whale sharks (Rhincodon typus) in the Gulf of Tadjoura, Djibouti. Environmental Biology of

Fishes 80:465–472.

Royle, J. A., R. B. Chandler, R. Sollmann, and B. Gardner. 2013. Spatial Capture-Recapture. Academic

Press.

Ruzzante, D. E., G. R. McCracken, B. Førland, J. MacMillan, D. Notte, C. Buhariwalla, J. Mills

Flemming, and H. Skaug. 2019. Validation of close‑kin mark–recapture (CKMR) methods for

estimating population abundance. Methods in Ecology and Evolution 10:1445–1453.

Sackett, D. K., and M. Catalano. 2017. Spatial Heterogeneity, Variable Rewards, Tag Loss, and Tagging

Mortality Affect the Performance of Mark–Recapture Designs to Estimate Exploitation: an

Example using Red Snapper in the Northern Gulf of Mexico. North American Journal of

Fisheries Management 37:558–573.

Schwarz, C. J., and G. A. Seber. 1999. A review of estimating animal abundance III. Statistical Science

14:427–456.

Schwarz, L. K., and M. C. Runge. 2009. Hierarchical Bayesian analysis to incorporate age uncertainty in

growth curve analysis and estimates of age from length: Florida manatee (Trichechus manatus)

carcasses. Canadian Journal of Fisheries and Aquatic Sciences 66:1775–1789.

Seber, G. A. F. 1965. A Note on the Multiple-Recapture Census. Biometrika 52:249–259.

Shaw, A. K., and S. A. Levin. 2013. The evolution of intermittent breeding. Journal of Mathematical

Biology 66:685–703.

Skaug, H. J. 2001. Allele-Sharing Methods for Estimation of Population Size. Biometrics 57:750–756.

Skjæraasen, J. E., K. Korsbrekke, G. E. Dingsør, Ø. Langangen, A. F. Opdal, and C. Jørgensen. 2020.

Large annual variation in the amount of skipped spawning for female Northeast Arctic haddock

Melanogrammus aeglefinus. Fisheries Research 230:105670.

Snelson, F. F., and S. E. Williams. 1981. Notes on the Occurrence, Distribution, and Biology of

Elasmobranch Fishes in the Indian River Lagoon System, Florida. Estuaries 4:110.

Trenkel, V. M., G. Charrier, P. Lorance, and M. V. Bravington. 2022. Close-kin mark–recapture

abundance estimation: practical insights and lessons learned. ICES Journal of Marine

Science:fsac002.

Vaudo, J., and M. Heithaus. 2011. Dietary niche overlap in a nearshore elasmobranch mesopredator

community. Marine Ecology Progress Series 425:247–260.

Walker, T. 2007. Spatial and temporal variation in the reproductive biology of gummy shark Mustelus

antarcticus (Chondrichthyes: Triakidae) harvested off southern Australia. Marine and Freshwater

Research – MAR FRESHWATER RES 58.

Waples, R. S., and P. Feutry. 2021. Close☐kin methods to estimate census size and effective population

size. Fish and Fisheries:faf.12615.

Ward, H. G. M., P. J. Askey, and J. R. Post. 2013. A mechanistic understanding of hyperstability in catch

per unit effort and density-dependent catchability in a multistock recreational fishery. Canadian

Journal of Fisheries and Aquatic Sciences 70:1542–1550.

Ward-Paige, C. A., D. M. Keith, B. Worm, and H. K. Lotze. 2012. Recovery potential and conservation

options for elasmobranchs. Journal of Fish Biology 80:1844–1869.

Webster, M. S., P. P. Marra, S. M. Haig, S. Bensch, and R. T. Holmes. 2002. Links between worlds:

unraveling migratory connectivity. Trends in Ecology & Evolution 17:76–83.

White, E. R., J. D. Nagy, and S. H. Gruber. 2014. Modeling the population dynamics of lemon sharks.

Biology Direct 9.

Wilson, G. J., and R. J. Delahay. 2001. A review of methods to estimate the abundance of terrestrial

carnivores using field signs and observation. Wildlife Research 28:151–164.

Wilson, H. B., B. E. Kendall, and H. P. Possingham. 2011. Variability in Population Abundance and the

Classification of Extinction Risk. Conservation Biology 25:747–757.

Table 1: Simulation scenarios for data-generating model and estimation model.

Scenario	Test	10 year mean λ	Model equation(s) used	Parameters estimated	Parameters derived	Parameters compared (naïve vs adapted)
1.1	Model validation	1	Eq. 2	$N_{♀}$, $N_{♂}$, □	NA	$N_{♀}$, $N_{♂}$, □
1.2	Misidentify aunt/niece pairs as HSPs	1	Eq. 2	$N_{♀}$, $N_{♂}$, □	NA	$N_{♀}$, $N_{♂}$, □
2.1	Slight population decline	0.99	Eq. 2, Eq. 3, Eq. 7, Eq. 8	$N_{♀(t0)}$, $N_{♂(t0)}$, □, λ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, □, λ
2.2	Slight population growth	1.01	Eq. 2, Eq. 3, Eq. 7, Eq. 8	$N_{♀(t0)}$, $N_{♂(t0)}$, □, λ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, □, λ
2.3	Severe population decline	0.93	Eq. 2, Eq. 3, Eq. 7, Eq. 8	$N_{♀(t0)}$, $N_{♂(t0)}$, □, λ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, □, λ
2.4	Stable population growth	1	Eq. 2, Eq. 3, Eq. 7, Eq. 8	$N_{♀(t0)}$, $N_{♂(t0)}$, □, λ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, □, λ
3.1	100% biennial breeders	1	Eq. 3, Eq. 4, Eq. 7, Eq. 8	$N_{♀(t0)}$, $N_{♂(t0)}$, □, λ, ψ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, $N_{♂(t)}$, □, ψ
3.2	100% biennial breeders w/ stochastic off-cycle breeding	1	Eq. 3, Eq. 4, Eq. 7, Eq. 8	$N_{♀(t0)}$, $N_{♂(t0)}$, □, λ, ψ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, $N_{♂(t)}$, □, ψ
3.3	100% triennial breeders	1	Eq. 3, Eq. 4, Eq. 7, Eq. 8	$N_{♀(t0)}$, $N_{♂(t0)}$, □, λ, ψ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, $N_{♂(t)}$, □, ψ
3.4	100% annual breeders	1	Eq. 3, Eq. 4, Eq. 7, Eq. 8	$N_{♀(t0)}$, $N_{♂(t0)}$, □, λ, ψ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, $N_{♂(t)}$, □, ψ
4.1	Minimal age-length uncertainty (5% CV)	1	Eq. 3, Eq. 4, Eq. 7, Eq. 8	$N_{♀(0)(t0)}$, $N_{♂(0)(t0)}$, □, λ, ψ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, □
4.2	Moderate age-length uncertainty (10% CV)	1	Eq. 3, Eq. 4, Eq. 7, Eq. 8	$N_{♀(0)(t0)}$, $N_{♂(0)(t0)}$, □, λ, ψ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, □
4.3	Substantial age-length uncertainty (20% CV)	1	Eq. 3, Eq. 4, Eq. 7, Eq. 8	$N_{♀(0)(t0)}$, $N_{♂(0)(t0)}$, □, λ, ψ	$N_{♀(t)}$, $N_{♂(t)}$	$N_{♀(t)}$, □

1232 *Note:* See Table 2 for parameter definitions.

1233

1234 **Table 2:** Model parameters and priors.

Parameter	Definition	Prior
$N_{s(t0)}$	Sex-specific abundance in year 0.	$N_{s(t0)} \sim \text{Normal}(\mu, \sigma)$ $\mu \sim \text{Uniform}(1, 10000)$ $\sigma \sim \text{Uniform}(1, 10000)$
Φ	Annual survival	Uniform(0.5, 0.95): default Uniform(0.5, 0.99): for small population simulations and application to Bimini data
λ	Annual finite population growth rate	Uniform(0.95, 1.05): default Uniform(0.80, 1.20): for severe decline scenario Uniform(0.70, 1.30): for small population simulations and application to Bimini data
ψ	Proportion of individuals that breed every a years	Uniform(0, 1)
a	Years between breeding	Fixed
Derived quantities	**Definition**	**Equation**
$N_{s(t)}$	Sex-specific abundance in year t	$N_{s(t0)} * \lambda^{(yj - t0)}$
$N_{\female b(t)}$	Abundance of breeding females in year t (in multiennial simulations)	$\dfrac{a + \psi - a\psi}{a} N_{(t)}$

1235 *Note: N_s is a general term that encompasses both $N_\female$, $N_\male$ when the sex-specific parameters were*
1236 treated the same.

1237 **Table 3:** Mean cross-correlation values between female abundance at t_0 ($N_{♀(t0)}$) and survival ($\square$)
1238 or population growth rate (λ) from population growth simulations (Table 1, scenario 2.1 – 2.4).

Parameter	$\square$	λ	t_0	Sampling scheme
$N_{♀(t0)}$	0.77	-0.13	88	target.YOY
$N_{♀(t0)}$	0.34	-0.76	77	sample all juveniles
$N_{♀(t0)}$	0.28	-0.80	77	sample all ages

1239

1240

Table 4: Year-specific abundance estimates for $N_{\female b(t)}$ from the real dataset for Bimini lemon sharks when all individuals were kept from each mother/father pairing.

Estimation year	All available samples				Five-year window				Three-year window			
	Median	95% HPDI	Total samples	MHSPs	Median	95% HPDI	Total samples	MHSPs	Median	95% HPDI	Total samples	MHSPs
1997	17	(15-19)	285	807	17	(15-19)	285	807	**14**	**(6-19)**	**257**	**570**
1998	15	(14-16)	367	1322	14	(13-15)	354	1148	12	(5-16)	245	324
1999	23	(22-24)	489	2279	24	(22-26)	461	1848	19	(8-27)	312	501
2000	17	(16-19)	569	3204	14	(12-16)	447	1623	8	(3-11)	284	566
2001	32	(29-35)	651	3623	15	(11-19)	474	1347	45	(18-65)	284	175
2002	34	(32-36)	722	4380	31	(29-34)	437	1213	26	(10-38)	233	166
2003	37	(34-39)	796	5298	35	(32-39)	429	933	21	(9-31)	227	227
2004	39	(37-41)	886	6219	29	(26-32)	397	915	18	(8-26)	235	249
2005	43	(39-46)	1038	7489	35	(32-39)	469	1228	**25**	**(10-36)**	**316**	**309**
2006	47	(45-51)	1148	8825	33	(30-37)	497	1213	**20**	**(8-28)**	**352**	**358**
2007	38	(36-41)	1268	10745	25	(23-26)	546	1743	**20**	**(8-27)**	**382**	**681**
2008	39	(36-41)	1328	11616	26	(23-28)	532	1536	25	(9-34)	290	212
2009	34	(32-35)	1451	13461	21	(18-25)	565	1931	20	(9-27)	303	555
2010	34	(32-35)	1501	14339	19	(15-23)	463	1144	18	(7-27)	233	114
2011	28	(27-29)	1626	17072	27	(25-29)	478	1941	19	(7-26)	298	582
2012	29	(28-30)	1637	17144	21	(18-25)	369	813	168	(26-442)	186	15
2013	27	(26-28)	1716	19006	19	(14-23)	388	1281	18	(7-25)	215	410
2014	26	(25-27)	1749	19519	16	(14-17)	298	603	10	(3-17)	123	27
2015	25	(24-26)	1847	21700	31	(28-34)	346	1066	26	(11-37)	210	217

Note: MHSPs refers to the number of maternal half-sibling pairs identified. Bold reflects instances that included at least one parameter where $1.01 < \text{Rhat} < 1.02$.

 Table 5: Cross-correlation among parameter estimates following application of CKMR to
 Bimini lemon sharks.

Parameter	$N_{♀(t0)}$	φ	$λ$	$ψ$	Time window
$N_{♀(t0)}$	1.00	0.13	-0.84	-0.08	all available samples
φ	0.13	1.00	0.32	-0.10	all available samples
$λ$	-0.84	0.32	1.00	0.02	all available samples
$ψ$	-0.08	-0.10	0.02	1.00	all available samples
$N_{♀(t0)}$	1.00	0.29	-0.73	-0.18	five-year window
φ	0.29	1.00	0.19	-0.11	five-year window
$λ$	-0.73	0.19	1.00	0.09	five-year window
$ψ$	-0.18	-0.11	0.09	1.00	five-year window
$N_{♀(t0)}$	1.00	0.83	-0.43	-0.54	three-year window
φ	0.83	1.00	-0.01	-0.46	three-year window
$λ$	-0.43	-0.01	1.00	0.21	three-year window
$ψ$	-0.54	-0.46	0.21	1.00	three-year window

1247 *Note:* Reported values represent the average cross-correlation values over all years (1997-2015).

1248

Appendix S1

S1.1. Constructing the pairwise comparison matrix

Close-kin mark-recapture models use a "pseudo-likelihood" framework to estimate

population parameters. Whereas a full likelihood would derive probabilities of kinship from the

population's full genealogy (which is infeasible in most cases), a pseudo-likelihood works by

specifying a probability of kinship for each pair of sampled individuals. Assuming each pairwise

comparison is independent, the pseudo-likelihood approximates the full likelihood and can be

used in its stead. This equivalence is central to CKMR and requires thoughtful construction of

the full pairwise comparison matrix that is used as the input to the CKMR model.

The full pairwise comparison matrix contains kinship probabilities for each pair of sampled

individuals. This matrix – which is likely to be tens of thousands or millions of rows depending

on sample size – can be condensed into fewer rows by grouping comparisons based on the

relevant covariates for the CKMR model (e.g. by reference year, birth year gap, and reference

year gap; Table S2). The relevant covariates vary based on the parameterization of the CKMR

model, which means that each model will allow for different degrees of grouping; the more the

data can be grouped, the faster the model will run at the expense of accounting for individual

variation.

Besides grouping the data – which will depend on how the model is constructed – there are

various considerations to keep in mind when constructing the final pairwise comparison matrix

for CKMR. First, comparisons with a zero percent chance of unveiling kin should be excluded.

Alternatively, careful construction of the model to ensure that such comparisons are assigned

zero probability of kinship in the model will suffice (but including both filters is optimal). For

parent-offspring relationships, this means removing comparisons between offspring and other

individuals that could not have birthed/sired them (e.g., because they themselves were not

sexually mature or alive during the offspring's birth year). For half-siblings, comparisons should

be excluded if the individuals were born too far apart for a single individual to have birthed both

of them. For instance, if a species is expected to live to a maximum age of 10 years old and is

reproductively mature at age 3 (assuming knife-edge maturity for this example), then

comparisons between individuals born more than 7 years apart should be excluded from the final

pairwise comparison matrix. That said, in most applications of CKMR, the boundaries of life

expectancy and reproductive maturity for a population will not be so clear-cut; as such, a

probabilistic approach that is directly integrated into the CKMR model may serve better when

the year gap is near the expected maximum.

Besides including impossible comparisons, an additional consideration for half-sibling

CKMR is whether or not to include within-cohort comparisons. The rationale and consequences

of this decision have been outlined elsewhere (Bravington et al. 2016a, Waples and Feutry

2021); for a CKMR model that incorporates half-sibling relationships and does not explicitly

account for same-year comparisons, the most straightforward approach to this issue is to only

retain cross-cohort comparisons in the final pairwise comparison dataframe.

Finally, it is worth mentioning that it can be easy to accidentally double-count comparisons;

as such, the pairwise comparison matrix should be double-checked to ensure that each

comparison is only included once. We have provided annotated R code that can serve as a

starting point (see Data Availability statement).

1292

S1.2. Instances of aunt/niece pairs

Including aunt/niece (uncle/nephew, etc.) pairs as HSPs had a minimal effect on our results

(Appendix S1, Figure S1a), and the small amount of bias that was introduced to the "sample all

ages" scenario was rectified by including a filter that removed all half-sibling comparisons that

spanned a year gap greater than the age of maturity (11 years). Aunt/niece pairs were rare in our

simulated dataset (Appendix S1, Figure S1b), as were full siblings (Appendix S1, Figure S2).

Conflating aunt/niece pairs with HSPs is likely to be a greater issue for circumstances in which

full siblings are more prevalent and/or when the first age at reproduction is early enough that

constraining year gap comparisons to the age at reproduction is impractical. For long-lived

promiscuous species like lemon sharks, a simple year gap filter set to the first age of

reproduction should suffice to remove most contaminating instances of aunt/niece pairs.

1304

S1.3. Accounting for a changing population

To interpret CKMR abundance estimates and appreciate its limitations, it is important to

understand that *each* pairwise comparison includes a specific year (y_j) for which abundance is

estimated. Crucially, y_j (the birth year of the younger individual) is different for each pairwise

comparison; however, pairwise comparisons can be grouped based on relevant covariates,

including y_j. If there are enough comparisons that share the same y_j, then it may be possible to

produce multiple independent estimates of $N_{(yj)}$, but doing so requires a very rich dataset. Instead,

most cases of CKMR application will benefit from leveraging all available data for a single

abundance estimate. This can be accomplished by specifying a population growth model that

links each instance of y_j. For example, we applied a simple exponential growth model to

Scenarios 2, 3, and 4, as well as to the small population simulations and Bimini dataset. To link

the exponential growth model to our CKMR model, we first selected a reference year (t_0), which

was the earliest instance of y_j in each dataset (i.e., the second oldest individual used in the half-

sibling probabilities). Then, for each pairwise comparison, abundance was estimated in y_j and

linked to t_0 via λ (see Equations 4 – 9). By including the parameter λ, all of the pairwise

comparisons were leveraged to produce a single estimate for $N_{(t0)}$. Then, N_t was derived from

estimates of $N_{(t0)}$ and λ using Equation 3.

1322

S1.4. Deriving Equation 7

Let a be the breeding interval, ψ the fraction of females that breed every a years, $N_{♀t}$

equal the total number of mature females in the population in year t, and $\widetilde{N}_t$ equal the number of

effective female breeders in year t. If we assume that $1/a$ of the non-annual breeders produce

pups each year, then

$\widetilde{N}_t = (\text{annual breeders}) + (1/a \text{ of non-annual breeders}) = (1 - \psi)N_t + \frac{\psi}{a}N_t =$

$\frac{a(1-\psi)N_t + \psi N_t}{a} = \frac{a + \psi - a\psi}{a}N_t.$

For individuals i and j born in years y_i and y_j, respectively, the probability that they have

the same mother d is the product of the probability that d is mother of i and the probability that d

is the mother of j. These probabilities are a function of relative reproductive effort and (when

$y_j > y_i$) survival. We define relative reproductive effort as the reproductive output of mother d in a

given year relative to the total reproductive output in that year. Assuming adult survival (ϕ) and

fecundity (f) are constant across mature ages, then the probability that i and j are half-siblings is

(from Bravington et al. 2016a):

$$\sum_{d \in \mathcal{F}_i} \frac{f}{f N_{(yi)}} \frac{\phi^\delta f}{f N_{(yj)}} = \frac{\phi^\delta}{N_{(yj)}}$$

1338

where the summation is across all breeding females at time i.

When breeding is intermittent, the denominators in the above equation (representing total

reproductive output in a given year t) are the product of fecundity (f) and effective female

breeders ($\widetilde{N}_t$), and the summation is across the number of effective breeders at time y_i. The first

quotient inside the summation becomes $\frac{f}{f\widetilde{N}_{(yi)}}$. Given that a female was alive and reproduced in

year y_i, the probability that same female survived and is the mother of j is now dependent on the

breeding cycle a and the fraction of the population that breeds every a years (ψ) or annually

$(1 - \psi)$. There are two cases to consider, namely whether or not δ is divisible by a. When δ is

divisible by a, then both annual and non-annual females breed; if δ is not divisible by a then only

the annual breeders are capable of being the mother. Thus, we have the probability of a maternal

half-sibling pair as

1350
$$
\begin{cases}
\sum_{d\in\tilde{\mathcal{F}}_i} \dfrac{f}{f\widetilde{N}_{(yi)}} \dfrac{\phi^\delta f(1-\psi)}{f\widetilde{N}_{(yj)}} = \dfrac{\phi^\delta(1-\psi)}{\widetilde{N}_{(yj)}} & \delta \text{ not evenly divisible by } a \\[4mm]
\sum_{d\in\tilde{\mathcal{F}}_i} \dfrac{f}{f\widetilde{N}_{(yi)}} \dfrac{\phi^\delta f}{f\widetilde{N}_{(yj)}} = \dfrac{\phi^\delta}{\widetilde{N}_{(yj)}} & \delta \text{ evenly divisible by } a
\end{cases}
$$

1351

1352   In the above, when δ is not divisible by a, the probability that mother d is an annual breeder is

1353 $(1 - \psi)$. When δ is divisible by a, the probability that mother d is either an annual or non-annual

1354   breeder is 1.

1355   Replacing $\widetilde{N}_t$ with $N_{\female t}$ we have equation (7):

1356

1357

$$
\begin{cases}
\dfrac{a\phi^{\delta}(1-\psi)}{(a+\psi-a\psi)N_t} & \delta \text{ not evenly divisible by } a \\[2ex]
\dfrac{a\phi^{\delta}}{(a+\psi-a\psi)N_t} & \delta \text{ evenly divisible by } a
\end{cases}
$$

1358

1359 When $\psi=1$ and $a>1$, this yields the obvious result that $\widetilde{N}_t = \frac{1}{a}N_t$. For example, if all females

1360 breed on a biennial cycle (i.e., $\psi = 1$ & $a = 2$), then all gaps δ are multiples of 2 and we have:

1361
$$
\frac{2\phi^{(\delta)}}{[2+1-2]N_t} = \frac{2\phi^{(\delta)}}{N_t} = \frac{\phi^{(\delta)}}{N_t/2} = \frac{\phi^{(\delta)}}{\widetilde{N}_t} \, .
$$

1362

1363 An alternative assumption for non-annual breeders is that *all* of them breed exactly every a years

1364 (rather than *1/a* of them breeding each year, as above). In this case, the number of effective

1365 breeders $\widetilde{N}_t$ in a given year depends on whether δ is divisible by a:

1366
$$
\widetilde{N}_t =
\begin{cases}
(1-\psi)N_t & \delta \text{ not divisible by } a \\
(1-\psi)N_t + \psi N_t & \delta \text{ divisible by } a
\end{cases}
$$

1367 Thus, every a years, all mature females breed, and only $(1-\psi)$ mature females breed in the

1368 intervening years.

1369

S1.5. Intermittent breeding and systemic differences in lifetime reproductive output

Our simulation results show that multiennial breeding causes bias in juvenile-focused

CKMR models that do not explicitly account for it. In circumstances where 100% of females

breed on a multiennial cycle, the positive bias likely arises from the inclusion of pairwise

comparisons that span year gaps without any HSPs (e.g., odd year gaps for biennial populations).

Such comparisons can be filtered before model fitting to produce unbiased parameter estimates,

including abundance, but the quantity ($N_{\female b(t)}$) will be different than if off-cycle comparisons are

included ($N_{\female(t)}$). If off-cycle comparisons are excluded (or, alternatively, given a probability of 0

in the model), $N_{\female(t)}$ can be derived by multiplying estimates of $N_{\female b(t)}$ by the reproductive

periodicity (e.g., 2 for biennial breeders). In most practical applications of CKMR, instances of

off-cycle breeding and/or skipped breeding cycles are likely to result in some positive

comparisons that do not align with the expected breeding cycle. Filtering comparisons from off-

cycle breeding years would then result in loss of valuable data at best, and biased parameter

estimates at worst (see Figure 4). In contrast, our multiennial model (Eq. 7) can leverage

pairwise comparisons regardless of whether the birth year gap is on- or off-cycle and produce

reliable estimates of total abundance and other parameters across a range of breeding cycles.

Using this approach, $N_{\female b(t)}$ can be derived from estimates of $N_{\female(t)}$ using Equation 6, as we have

done with Bimini lemon sharks.

Another circumstance that is related to differences in realized reproductive output and

known to cause issues for CKMR is persistent individual differences in fecundity that give rise to

lifetime differences in reproductive output. We confirmed this issue by fitting an annual and

multiennial model to simulated populations with variable proportions of biennial and annual

breeders *without* adjusting fecundity (Figure S10). This resulted in a population where annual

breeders produced more offspring on average over their lifetimes, but this difference in

reproductive output was not reflected in the kinship probabilities (Eq 7 assumes equal

reproductive output among all females). Abundance estimates (primarily for females) using the

multiennial model were more biased as a consequence of this unmodeled heterogeneity relative

to when lifetime fecundity was constant across the population (Figure 4), though we note that the

absolute bias was still less than 20% for two of the three sampling scenarios. Thus, our results

confirm that unmodeled persistent differences in lifetime fecundity across different sectors of the

population will, at a minimum, result in less reliable model performance, and we recommend

future CKMR model development to accommodate persistent individual differences in fecundity.

1402

S1.6 CKMR with a small population

The pseudo-likelihood that CKMR employs has the properties of a full likelihood when

sampling is sparse because each pairwise comparison among individuals is approximately

independent (Bravington et al. 2016a). However, when CKMR is applied to very small

populations (<~100 individuals) and sampling is non-sparse the assumption of independence

among samples is difficult to fulfill as capture of littermates, which may have similar genetic

fitness and/or early-life history environments, becomes increasingly common. In such cases, it is

important to account for or remove littermates to maintain independence among samples, since

comparisons between littermates and a third animal are not independent (Bravington et al.

2016a).

In our application to simulated and real lemon sharks with very small population sizes

(<100 total adult females), we found that applying CKMR to this small population without

substantial modification to the likelihood produced estimates of abundance that were still very

close to the true value. When we downsampled the real genetic dataset to 30% of the original

sample size, the five-year window and the window that included all available samples produced

parameter estimates that were similar to the full dataset but were predictably less precise (Figure

S8, Table S4, Table S5). The combination of downsampling and limiting comparisons to a three-

year window produced several instances of very few MHSPs arising from exceptionally small

sample sizes, and the associated abundance estimates were unrealistic (Table S4, Table S5).

Apart from those instances, the overall results remained similar whether we kept or excluded

littermates, but retaining littermates produced parameter estimates that were less biased. We

suspect that when populations are very small and heavily sampled such that pairwise

comparisons are not independent, excluding littermates may skew the fecundity evident in the

remaining samples, thereby increasing bias relative to when littermates are retained. By contrast,

in larger populations retaining littermates would violate the assumption of non-independence for

only a small subset of the potential comparisons, but a relatively large proportion of the

identified true kin pairs. This could result in a large effect on parameter estimation, which may

depend on relatively few kin pairs if sampling is sparse. Adapting CKMR for robust application

to small populations is an active area of research that will likely require modifications to the

statistical framework (Bravington et al. 2016a), including adding a parameter for capture

probability to PO kinship probabilities (see section 2.4.1 of this paper). However, our results

show that even in the absence of such modifications, CKMR can provide useful parameter

estimates for small populations.

1436

1437

1438

1439 **Table S1.** Parameters for the data generating model (DGM). Values in bold produced, on
1440 average, a stable population size at equilibrium over the 90-year time course.

Parameter	True value (DGM)
Female fecundity (total pups per year)	**5.8** (range: 2-9)
Breeding cycle	1, 2, or 3 years
Initial adult population size	**1000**
Young-of-year survival (age 0)	**0.7**
Juvenile survival (ages 1-11)	**0.80** (range: 0.72 – 0.81)
Adult survival (age 12-50)	**0.825** (range: 0.75 - 0.83)
Age at maturity	**12**
Sex ratio	**50/50**
Maximum age	**50**

1441

Table S2. An example pairwise comparison matrix (subsetted for first 20 rows) used as input to
a multiennial CKMR model (Eq. 6). ref.year is the birth year of the younger individual in the
pairwise comparison (i.e., y_j); all is the total number of comparisons that contain the same values
for ref.year, mort.yrs, and ref.year.gap; yes is the total number of positive comparisons with
those same values; type is either HS (half-sibling) or parent-offspring (PO); parent is mother or
father; ref.year.gap is the reference year gap i.e. ($y_j - t_0$); BI refers to the breeding interval, which
is either on-cycle or off (depending on a) for HS comparisons and irrelevant for PO comparisons.
There are myriad ways to construct a pairwise comparison dataframe for CKMR; we found that
producing the table below allowed us to store all comparisons in the same dataframe for input to
the CKMR model. Note that not all columns are necessary for each model: for instance, BI is
only used when the model accounts for intermittent breeding dynamics, while ref.year and
pop.growth.yrs only matter if using a population growth model.

ref.year	all	yes	mort.yrs	type	parent	ref.year.gap	BI
90	90	0	0	PO	mother	10	NA
90	449	2	1	HS	mother	10	off
90	90	0	1	PO	mother	10	NA
90	717	0	2	HS	mother	10	on
90	72	0	2	PO	mother	10	NA
90	899	1	3	HS	mother	10	off
90	108	1	3	PO	mother	10	NA
90	845	0	4	HS	mother	10	on
90	611	0	5	HS	mother	10	off
90	537	1	6	HS	mother	10	on
90	378	0	7	HS	mother	10	off
90	234	0	8	HS	mother	10	on
89	100	0	0	PO	mother	9	NA
89	125	2	0	PO	mother	9	NA

1454

1455

1456 **Table S3:** Year-specific abundance estimates for $N_{\female b(t)}$ from the real dataset for Bimini lemon
1457 sharks when only one individual was retained from each mother/father pairing.

Estimation year	All available samples				Five-year window				Three-year window			
	Median	95% HPDI	Total samples	MHSPs	Median	95% HPDI	Total samples	MHSPs	Median	95% HPDI	Total samples	MHSPs
1997	15	(10-20)	106	110	15	(10-20)	106	110	13	(5-20)	93	70
1998	16	(11-21)	122	147	15	(9-21)	117	122	15	(5-25)	79	31
1999	23	(16-28)	158	245	22	(15-29)	146	194	20	(8-32)	94	55
2000	22	(16-27)	180	310	20	(13-26)	137	140	12	(4-20)	80	30
2001	38	(28-47)	214	361	24	(12-40)	152	127	42	(14-72)	101	28
2002	40	(32-47)	252	474	35	(25-46)	155	117	36	(12-61)	105	30
2003	46	(38-54)	285	583	42	(31-54)	174	135	32	(11-54)	115	44
2004	50	(43-58)	318	686	37	(23-50)	171	129	27	(9-43)	108	41
2005	58	(47-69)	377	845	43	(30-55)	207	197	34	(12-53)	126	47
2006	62	(53-71)	428	1042	45	(34-56)	219	194	32	(12-51)	145	47
2007	52	(45-61)	473	1254	32	(24-39)	223	221	**27**	**(10 – 40)**	**157**	**78**
2008	60	(51-69)	495	1351	43	(28-56)	213	168	41	(12-70)	122	28
2009	52	(45-60)	540	1553	28	(17-40)	227	229	24	(9-36)	118	69
2010	53	(46-61)	560	1662	21	(12-34)	189	145	22	(7-40)	92	18
2011	45	(40-51)	612	1993	37	(27-45)	191	222	31	(11-46)	121	59
2012	47	(41-53)	620	2011	33	(21-48)	153	101	141	(17-442)	84	6
2013	42	(38-48)	657	2239	24	(13-38)	166	160	25	(9-38)	100	64
2014	39	(35-44)	678	2373	21	(16-26)	142	117	19	(5-43)	68	9
2015	34	(30-38)	718	2707	26	(18-34)	162	206	20	(7-30)	101	59

1458 *Note:* MHSPs refers to the number of maternal half-sibling pairs identified. Bold instances
1459 included one or more estimated parameter where Rhat = 1.01.
1460

Table S4: Year-specific abundance estimates for $N_{\female b(t)}$ from the real dataset for Bimini lemon sharks when the dataset was downsampled to 30% of the full dataset and all individuals were kept from each mother/father pairing. The numbers reported here represent the means over 50 iterations of random downsampling.

Estimation year	All available samples				Five-year window				Three-year window			
	Median	95% HPDI	Total samples	MHSPs	Median	95% HPDI	Total samples	MHSPs	Median	95% HPDI	Total samples	MHSPs
1997	15	(9-21)	88	80	15	(9-21)	88	80	14	(5-22)	79	56
1998	13	(10-17)	113	128	14	(10-18)	109	112	13	(4-22)	75	32
1999	22	(16-27)	150	212	22	(15-28)	141	168	21	(7-33)	95	46
2000	16	(12-20)	174	299	12	(8-17)	136	150	8	(3-13)	86	52
2001	29	(21-37)	199	337	17	(8-29)	144	119	61	(17-127)	86	16
2002	32	(25-39)	221	407	28	(19-37)	133	108	35	(9-74)	71	17
2003	35	(28-43)	244	491	35	(22-49)	131	86	27	(8-53)	70	22
2004	35	(28-43)	271	582	26	(16-38)	121	86	22	(7-40)	72	24
2005	43	(34-52)	317	692	31	(20-42)	143	115	31	(10-54)	96	29
2006	47	(38-56)	350	818	29	(19-40)	151	113	24	(8-41)	106	32
2007	39	(32-46)	386	999	24	(17-30)	165	157	21	(8-33)	115	60
2008	38	(32-46)	404	1083	24	(16-32)	160	139	31	(9-64)	87	19
2009	34	(29-40)	441	1279	21	(12-30)	170	174	21	(8-34)	91	51
2010	34	(29-40)	456	1318	21	(12-30)	139	105	42	(6-163)	70	11
2011	28	(25-33)	494	1579	26	(19-34)	144	170	20	(7-31)	90	55
2012	30	(26-35)	498	1570	22	(12-34)	112	74	953	(18-5203)	57	2
2013	28	(24-32)	522	1753	20	(10-31)	118	115	19	(7-30)	66	38
2014	26	(22-30)	532	1809	17	(10-24)	91	56	566	(4-3287)	38	3
2015	27	(23-30)	562	1998	26	(15-36)	106	99	40	(10-99)	64	20

Note: MHSPs refers to the number of maternal half-sibling pairs identified.

1467 **Table S5:** Year-specific abundance estimates for $N_{\female b(t)}$ from the real dataset for Bimini lemon
1468 sharks when the dataset was downsampled to 30% of the full dataset and only one individual was
1469 retained from each mother/father pairing. The numbers reported here represent the means over
1470 50 iterations of random downsampling.

Estimation year	All available samples				Five-year window				Three-year window			
	Median	95% HPDI	Total samples	MHSPs	Median	95% HPDI	Total samples	MHSPs	Median	95% HPDI	Total samples	MHSPs
1997	14	(9-22)	60	37	14	(9-22)	60	37	18	(10-29)	54	25
1998	13	(8-21)	72	53	16	(8-28)	70	45	39	(11-102)	47	11
1999	21	(14-30)	94	91	22	(13-34)	88	73	30	(15-54)	57	21
2000	17	(11-24)	106	116	17	(9-27)	82	55	16	(7-32)	47	12
2001	37	(25-51)	123	129	40	(21-63)	87	48	93	(26-260)	53	8
2002	34	(24-46)	141	162	30	(16-49)	83	40	93	(19-387)	49	8
2003	36	(27-46)	155	200	38	(20-65)	87	39	52	(17-129)	53	11
2004	39	(29-49)	171	223	39	(19-70)	82	35	59	(18-139)	51	11
2005	52	(41-64)	202	281	51	(25-86)	100	56	72	(27-143)	65	15
2006	58	(47-71)	229	333	70	(32-119)	108	52	61	(23-127)	74	15
2007	58	(47-69)	253	398	49	(25-81)	114	71	41	(19-77)	82	26
2008	56	(46-67)	265	433	41	(20-67)	109	55	54	(17-135)	61	9
2009	50	(42-59)	288	495	22	(16-30)	118	74	29	(17-47)	63	21
2010	52	(43-61)	299	535	21	(14-29)	97	44	41	(8-201)	48	7
2011	43	(37-49)	323	618	21	(16-27)	100	74	31	(18-50)	63	21
2012	43	(37-50)	332	654	20	(12-31)	79	33	1304	(20-6180)	43	2
2013	35	(30-40)	350	733	20	(14-27)	85	52	25	(14-42)	50	19
2014	35	(30-40)	363	790	15	(10-23)	68	28	825	(9-4807)	30	2
2015	31	(27-35)	372	839	19	(13-26)	79	51	38	(16-81)	48	13

1471 *Note:* MHSPs refers to the number of maternal half-sibling pairs identified.

Figure 1: Schematic of CKMR sensitivity tests, examined via individual-based simulation (see also Tables 1 and 2; Scenario 1 was model validation, and Scenario 5 involved real genetic data, so are not included here). Populations with distinct pedigrees were produced and sampled via an individual-based data-generating model (purple). Population parameters were individually varied for each of three scenarios. Each population was sampled in three ways, and each set of samples was used as input to two estimation models (green): one model was naïve to the added population dynamics of the DGM, and one model was adapted to account for them. The year of estimation (year t) was varied for Scenario 2; otherwise, simulation results that are discussed in the text used the model settings highlighted in bold.

Figure 2: Base CKMR model performance and kin pairs detected for three different sampling schemes at four different sampling intensities over 500 iterations. **a)** Relative bias of abundance estimates of adult females (*Nf*) and males (*Nm*) as a percentage of the truth (i.e. relative bias x 100). Bias was calculated from the median of each of 500 posterior distributions. **b)** CV on abundance estimates with log-scaled Y axis for visualization. **c)** Number of sex-specific half-sibling pairs detected by sampling scheme and sampling intensity. For each iteration, the number of half sibling pairs for each sex was calculated and averaged. **d)** Number of sex-specific parent-offspring pairs detected for the "sample all ages" sampling scheme.

Figure 3: Performance of CKMR models when confronted with a changing population **a-c)** Relative bias of CKMR abundance estimates for mature females ($N_{\female t}$) when applied to populations experiencing variable degrees of population growth or decline. Plots are split by sampling scheme (column), population growth pattern (row facet) and the year to which abundance estimates were targeted (year t). The dashed vertical blue line represents 0% relative bias. Scenarios assessed had population growth as **a)** slightly increasing (+1% per year), **b)** slightly declining (-1% per year), or **c)** severely declining (-5-10% per year over the final 10 years). Two different models were fit to 500 simulated populations for each scenario: a naïve model without a parameter for population growth (red) and an adapted model that included a parameter (λ) for population growth (blue). Plots were truncated at +/-100% for visualization because there were long tails of positive bias for the 10 year past scenarios. Note that for the target YOY sampling scenario, t_0 (the first instance of y_j in the dataset) occurred three years in the past, making this the only scenario where an estimation year of t_0 occurred more recently than when the estimation year was "present-5". **d)** Summary of age distribution of samples for all three sampling scenarios. The dashed vertical line represents t_0, which varied depending on the ages sampled. Many more samples were drawn during year t_0 for the target YOY sampling scheme than for any other. **e)** Relative bias of λ estimates over the modeled time period.

Figure 4: Relative bias of parameter estimates when naïve and adapted CKMR models were fit to samples from a population that breeds intermittently. The "naïve" model references a model that was included off-cycle comparisons, while the "naïve – filtered" model excluded off-cycle comparisons. The reference year (t_0) was set to the earliest instance of y_j and abundance was estimated in the present via Eq. 3 and compared to the true abundance in that year. For the "naïve – filtered" scenario, the quantity estimated by the model was the effective female breeders in year t ($\widetilde{N}_t$), while the "naïve" and "adapted" models estimated total females ($N_{\female t}$, or Nf). We derived total Nf for the "naïve – filtered" scenario by multiplying $\widetilde{N}_t$ by the breeding cycle (2 for biennial breeders and 3 for triennial breeders). On the x-axis, "biennial*" refers to a situation where 10% of biennial breeders bred off-cycle and 10% of on-cycle females failed to breed each year. **a)** Relative bias of abundance estimates for females (Nf). **b)** Relative bias of abundance estimates males (Nm). **c)** Relative bias of survival estimates.

Figure 5: Effect of aging error on CKMR parameter estimates. **a)** Amount of error introduced per age for one of the 500 iterations that were run to test the effects of age misassignment on CKMR parameter estimates. The iteration represented here was chosen randomly and assumed to be generally representative of all 500 iterations. **b)** Relative bias of abundance estimates for females (Nf) when uncertainty was introduced to length-based age assignments. **c)** Relative bias of survival estimates when uncertainty was introduced to length-based age assignments. There was no intentional model misspecification in these simulations; rather, annual models were fit to populations that bred annually (light blue), while multiennial models were fit to populations that bred biennially (dark blue), thereby isolating the effects of aging error on the resulting bias.

Figure 6: Time series of CKMR parameter estimates for simulated (**a-c**) and real (**d**) female lemon sharks at Bimini, Bahamas using all samples collected before the estimation year (green; solid trendline), all samples collected in a five year window prior to the estimation year (orange; dotted trendline) and all samples collected in a three year window prior to the estimation year (purple; dashed trendline). **a-c)** Relative bias from 100 distinct population simulations and model fits. **a)** Relative bias of abundance estimates for adult females (*Nf*) in each year of the time series. **b)** Relative bias of λ estimates relative to the observed population growth rate in the associated estimation year. **c)** Relative bias of survival (☐) relative to the observed survival rate in the associated estimation year. **d)** Abundance estimates for breeding females (*Nfb*) in the North Bimini Lagoon using real genetic data, derived from estimates of *Nf* using Eq. 6. Points represent the median of the posterior distribution, and error bars reflect the 95% highest posterior density interval (HPDI). The trend is visualized using a loess regression. The black line labeled as

"pedigree abundance" is a time-series of abundance estimates for *Nfb* that was independently derived for the population by Dibattista et. al. (2011).

Figure S1: Prevalence of aunt/niece (uncle/nephew, etc.) pairs and effects on model performance by sampling scheme and sampling intensity. **a)** Histogram summarizing the number of identified aunt/niece (uncle/nephew, etc.) pairs. The Y axis represents the number of iterations that contained the number of aunt/niece pairs specified by the X axis. Green represents the total number of aunt/niece etc. pairs identified; orange represents the number of pairs remaining after instituting a filter that removed pairwise comparisons with a year gap greater than the age of reproduction. Most simulations contained 0 aunt/niece pairs, so these were removed for visualization purposes. **b)** Effects of mistakenly including aunt/niece pairs as half-siblings before (green) and after (orange) a year gap filter when 2% of the population was sampled. The effect on bias was minimal, and the distributions overlapped almost perfectly (though there is a very slight observable difference for the sample all ages scenario).

Figure S2: Histogram of full siblings observed for different sampling schemes and intensities, collated over 500 iterations. The Y axis represents the number of iterations that contained the number of full sibling pairs specified by the X axis The total number of full siblings (green) includes within-cohort full siblings and represents almost all instances. The number of cross-cohort full siblings (orange) was substantially lower.

Figure S3: Estimates of **a)** female abundance (*Nf*) when a parameter for population growth (λ) was included in the model and realized population growth was stable, and **b)** survival (ϕ) under different population growth scenarios.

Figure S4: Comparison of relative bias for different population growth scenarios and approaches to integrating a population growth model. The orange density plots come from a model where abundance in year t was directly estimated in year t (i.e., the "Estimation year") using Eqs. 4 and 5. The blue density plots come from the primary model we used throughout our simulations, where t_0 represented the first instance of y_j in each dataset, and abundance in year t was derived from estimates of $N_{♀(t0)}$ using Eq. 3.

Figure S5: Estimates of ψ for multiennial breeding scenarios. The blue points represent the distribution of estimates, and the red crosses represent the mean realized value of psi across all 500 iterations, calculated as the proportion of total positive comparisons that came from on-cycle breeders.

Figure S6: A representative iteration of a time-series of abundance estimates for females from Bimini lemon shark simulations when full siblings are retained in the analysis. Estimates for each year for three different sampling windows are shown relative to the truth (black). Trends are visualized using a loess regression.

Figure S7: Time series of CKMR parameter estimates for simulated (**a-c**) and real (**d**) female lemon sharks at Bimini, Bahamas when only one individual from each mother/father pairing is retained from each dataset. The colors and lines are the same as Figure 6. **a-c)** Relative bias from 100 distinct population simulations and model fits. **a)** Relative bias of abundance estimates for adult females (*Nf*) in each year of the time series. **b)** Relative bias of λ estimates relative to the observed population growth rate in the associated estimation year. **c)** Relative bias of survival (□) relative to the observed survival rate in the associated estimation year. **d)** Abundance estimates for breeding females (*Nfb*), derived from estimates of total *Nf* using Eq.6, in the North Bimini Lagoon using real genetic data. Points represent the median of the posterior distribution, and error bars reflect the 95% highest posterior density interval (HPDI). The trend is visualized using a loess regression. The black line labeled as "pedigree abundance" is a time-series of abundance estimates for the population that was independently derived for the population by Dibattista et. al. (2011).

Figure S8: Estimates of abundance and survival for females in the North Bimini Lagoon using datasets that were downsampled and three different sampling windows. Estimates represent the means over 50 iterations of random downsampling. **a)** Estimates of abundance and survival for breeding females using 30% of the total samples for each year without removing full siblings. **b)** Estimates of abundance and survival for breeding females using 30% of the total samples for each year and also removing full siblings.

a) Retain full siblings

Sample window

- all available samples
- five year window
- three year window

b) Remove full siblings

Figure S9: CKMR-based survival estimates for Bimini lemon sharks using real data and three different sampling windows when **a)** full siblings were retained in the dataset and **b)** when full siblings were removed from the dataset. Trends were visualized using a loess regression.

Figure S10: Relative bias of parameter estimates in a simulated population with different ratios of biennial vs. annual female breeders, holding fecundity constant such that lifetime reproductive output varied among the different portions of the population. Both annual and biennial models were fit to the data. **a)** Relative bias of abundance estimates for females. **b)** Relative bias of abundance estimates for males. Note that males bred annually, but shared parameters for survival ($\square$) and population growth (λ) with females. **c)** Relative bias of shared survival estimates. **d)** Relative bias of estimates of ψ with different proportions of biennial breeders.