A tRNA modification in *Mycobacterium tuberculosis* facilitates optimal intracellular growth

Authors: Francesca G. Tomasi[1*], Satoshi Kimura[2,3,4*#], Eric J. Rubin[1], Matthew K. Waldor[1,2,,3,4]

Affiliations:

[1]Department of Immunology and Infectious Diseases Harvard T. H. Chan School of Public Health,

Boston, MA USA

[2]Division of Infectious Diseases, Brigham and Women's Hospital, Boston, MA, USA

[3]Department of Microbiology, Harvard Medical School, Boston, MA, USA

[4]Howard Hughes Medical Institute, Boston, MA, USA

*Equal contribution

#For correspondence: s.kimura.res@gmail.com

Abstract

Diverse chemical modifications fine-tune the function and metabolism of tRNA. Although tRNA modification is universal in all kingdoms of life, profiles of modifications, their functions, and physiological roles have not been elucidated in most organisms including the human pathogen, *Mycobacterium tuberculosis* (*Mtb*), the causative agent of tuberculosis. To identify physiologically important modifications, we surveyed the tRNA of *Mtb*, using tRNA sequencing (tRNA-seq) and genome-mining. Homology searches identified 23 candidate tRNA modifying enzymes that are predicted to create 16 tRNA modifications across all tRNA species. Reverse transcription-derived error signatures in tRNA-seq predicted the sites and presence of 9 modifications. Several chemical treatments prior to tRNA-seq expanded the number of predictable modifications. Deletion of *Mtb* genes encoding two modifying enzymes, TruB and MnmA, eliminated their respective tRNA modifications, validating the presence of modified sites in tRNA species. Furthermore, the absence of *mnmA* attenuated *Mtb* growth in macrophages, suggesting that MnmA-dependent tRNA uridine sulfation contributes to *Mtb* intracellular growth. Our results lay the foundation for unveiling the roles of tRNA modifications in *Mtb* pathogenesis and developing new therapeutics against tuberculosis.

Introduction

Introduction

tRNA is an adaptor molecule that enables protein synthesis by converting the triplet genetic

code in mRNA into amino acids. The fidelity of base paring between mRNA codons and tRNA

anticodons is monitored within ribosomes and is critical for properly incorporating the amino acids

bound to the 3' ends of tRNAs into growing polypeptides. For optimal translation, the abundances,

and properties of tRNA isoacceptors are fine-tuned by diverse mechanisms, including chemical

modifications[1-4] Dysregulation of tRNA abundance and/or structure leads to defective decoding

and results in ribosome pausing and collisions, protein misfolding, stress responses and can have

detrimental or lethal effects on the cell[5-10].

Chemical modifications of tRNA (tRNA modifications) are found in all kingdoms of life and

fine-tune tRNA properties including mRNA decoding efficiency, recognition by aminoacyl-tRNA

synthetases, half-life and structural stability[2, 10-12]. Modifications are prevalent in the anticodon

loop, particularly at the first letter of the anticodon. Modifications of the anticodon loop directly

modulate codon recognition, whereas modifications in the tRNA body region primarily stabilize tRNA

tertiary structure, protecting them from degradation in the cell. tRNA modifications are generated by

dedicated site-specific enzymes referred to as tRNA modifying enzymes. tRNA modifications have

been extensively characterized in a few model organisms[2, 13], but their profiles, regulation, and

functions in non-model organisms, including bacterial pathogens, are understudied[13]

tRNA sequencing (tRNA-seq) allows for the rapid and systematic prediction of many tRNA

modification sites[14, 15]. We recently developed a comparative tRNA-seq protocol to profile tRNA

modifications in organisms with uncharted tRNA modification profiles; in *Vibrio cholerae,* this

approach led to the discovery of a new RNA modification and RNA editing process[16]. tRNA-seq

enables rapid prediction of modified sites through detection of reverse transcription-derived

signatures, such as nucleotide misincorporation and early termination, both of which occur more

frequently at modified sites. Furthermore, several chemical treatments of tRNA can convert

modifications that are not recognizable as reverse transcription-derived signatures into detectable

signals, expanding the repertoire of modifications that can be distinguished by tRNA-seq[17-20].

Mycobacterium tuberculosis (*Mtb*), the agent of tuberculosis (TB), is a global pathogen that caused

>10.5 million cases and over 1.5 million deaths worldwide in 2020[21]. Several studies have uncovered

roles for non-*Mtb* mycobacterial tRNA modifications in stress responses, adaptation to environmental

changes, and persister cell formation[22]. *Mycobacterium bovis* BCG, an organism closely related to *Mtb*,

responds to hypoxia by reprogramming 40 ribonucleoside modifications in tRNA to facilitate

translation of a subset of proteins that promote survival in hypoxic conditions[22]. To date, studies of

the profiles and functions of *Mtb* tRNA modifications have been limited.

Here, we conducted tRNA-seq in *Mtb*. We assigned modifications to many of the reverse

transcription-derived signatures identified, using information on the presence of homologs of known

modifying enzymes in *Mtb*. Chemical treatments of tRNAs carried out prior to tRNA-seq increased

the detectability of certain modifications. We constructed two *Mtb* deletion mutant strains, with

deletions of *mnmA* and *truB*, and confirmed that the absence of these modification enzymes eliminated

the predicted signals in tRNA-seq data. Furthermore, while deletion of *mnmA* in *Mtb* did not affect

the pathogen's growth in *in vitro* laboratory growth conditions, the *mnmA* knockout strain's growth

was attenuated in a macrophage infection model. Our findings suggest that tRNA modifications

warrant further study as we unravel the complexity of *Mtb* infections, as they may serve as targets for

new therapeutics.

Results

***In silico* prediction of *Mtb* tRNA modifying enzymes**

To predict tRNA modifications in *Mtb*, we used Basic Local Alignment Search Tool

(BLAST)[23] to identify homologues of all tRNA modification enzymes registered in Modomics in

the *Mtb* genome[24] (Supplementary Table 1). With a stringent threshold (E-value <1×10^{-10}), 31 *Mtb*

genes homologous to genes encoding known RNA modification enzymes were identified

(Supplementary Table 2). 23 of these genes are predicted to synthesize 16 tRNA modifications in *Mtb*

(Supplementary Table 2 and 3), including *miaA* and *miaB* for 2-methylthio-6-isopentenyl-adenosine

(ms^2i^6A), *tsaD*, *tsaB*, *tsaE*, and *sua5* for N^6-threonylcarbamoyladenosine (t^6A), *mnmA* for 2-thiouridine

(s^2U), *truB*, *truA*, *rluA*, and *pus9* for pseudouridine (Ψ), *trm2* for 5-methyluridine (m^5U), *trmD* for 1-

methylguanosine (m^1G), *trmI* for 1-methyladenosine (m^1A), *trmL* for 2'-*O*-methylcytidine (Cm) or 2'-

O-methyluridine (Um), *trmB* for 7-methylguanosine (m^7G), two *trmH* for 2'-*O*-methylguanosine (Gm),

trmR for 5-methoxyuridine (mo^5U), *trcM* for 5-methylcytidine (m^5C), *dusB* for dihydrouridine (D), *tadA*

for inosine (I), and *tilS* for lysidine (k^2C). While eight additional genes met the threshold for homology

to modification enzymes, they exhibited greater similarity to a Glutamyl-tRNA synthase; Rv2992c,

ribosome associated GTPases (Der; Rv1713 and Era; Rv2364c), molybdopterin biosynthesis proteins

(MoeW; Rv2338c and MoeB2; Rv3116), thiosulfate sulfur transferases (CysA; Rv3117 and SseA;

Rv3283) and a riboflavin biosynthesis protein (RibG; Rv1409), and likely do not correspond to tRNA

modification enzymes (Supplementary Table 2).

We mined data from genome-wide Tn-seq and CRISPRi screens[25, 26] to assess the impacts

of *Mtb* tRNA modifications on its growth. Five genes encoding *Mtb* tRNA modifying enzymes, *trmD*,

tilS, *tadA*, *tsaC2*, and *miaA*, were reported to be essential for *Mtb* growth in both Tn-seq[25] and

CRISPRi screens[26] (Supplementary Table 3), suggesting that the modifications they produce are

critical for tRNA functions. Indeed, *E. coli trmD, tilS, tadA, and tsaC2* are also essential for growth and

critical for codon decoding, aminoacylation, and reading frame maintenance[27-30]. The *Mtb*

modifications synthesized by these enzymes likely have similar impacts on tRNA functions.

Unexpectedly, one modifying enzyme, MiaA, is non-essential in *E. coli* grown in nutrient rich medium,

but apparently essential in *Mtb*, suggesting that i^{6}A, the modification introduced by MiaA, may have

more profound roles in *Mtb* translation than in *E. coli*.

Several of the putative *Mtb* tRNA modifying enzymes are conserved across all three domains

of life (e.g., TruB and TsaD) (Fig. 1). By contrast, some enzymes were limited to bacterial species

closely related to *Mtb*, possibly suggesting their species-specific physiological roles, including

pathogenesis. For example, TrmI (Rv2118c) homologs are widely present in Archaea and Eukaryotes,

but are sparsely distributed in bacteria, where they are primarily limited to Actinomycetia, including

Mtb, and several thermophilic bacterial species (e.g., *Thermus thermophilus*). TrmI synthesizes m^{1}A at

position 58 in eukaryotes[31] and at position 57/58 in archaea[32]; indeed, TrmI in *Mtb* has been

proven to generate m^{1}A at position 58[33]. Pus9 (Rv3300c) has many homologs exhibiting weak

similarity across eukaryotes and bacteria. However, some species, including *Pseudomonas aeruginosa*,

Acinetobacter baumannii, *Neisseria gonorrhoeae*, and several species of Actinomycetia, encode proteins

showing strong homology to *Mtb* Pus9, suggesting that this set of pseudouridylases have a distinctive

property such as substrate specificity.

tRNA modifying enzymes predicted in *Mtb* were observed in mycobacteria species, including

Mycolicibacterium smegmatis and *Mycobacteroides abscessus* (Fig. 1), suggesting that tRNA modification

patterns are similar among mycobacterium species.

Fig. 1. Phylogenetic distribution of *Mtb* tRNA modifying enzyme homologs. Heat map of log10 E-values from BLAST search results. BLAST searches were conducted against 120 manually picked organisms using *Mtb* tRNA modifying enzymes as queries. When one organism has multiple hits, the lowest log10(Eval) values among hits is shown. iTol[34] was used to depict the results.

Profiling *Mtb* tRNA modification sites by tRNA sequencing

To begin profiling detectable *Mtb* tRNA modifications, we sequenced tRNAs isolated from

wild type *Mtb* strain H37Rv grown in 7H9 medium. In this protocol, tRNAs are first reversed

transcribed to cDNA[16]. During cDNA synthesis, chemical modifications on tRNA nucleotides

disrupt Watson-Crick base pairing and increase the frequency of reverse transcriptase errors, leading

to incorporation of the incorrect nucleotide or early termination of cDNA synthesis[35]. These

reverse transcription derived 'signatures' typically correspond to modified sites[14, 15] and are

depicted in the heat map in Fig. 2.

Comparison of the reverse transcription derived signatures observed in *Mtb* to *E. coli*, where

tRNA modifications have been well characterized[16] enables the prediction of the presence of

common modifications, including ms^2i^6A, m^1G, I, and k^2C (Fig 2 and Supplementary Fig. 1). These

predictions are strongly supported by the set of tRNA modification enzymes identified in the *Mtb*

genome (Fig. 1 and Supplementary Table 3), including *miaA* and *miaB* (ms^2i^6A), *trmD* (m^1G), *tadA* (I),

and *tilS* (k^2C). Some tRNA modifications were observed in *Mtb* but are not present in *E. coli*. In other

actinobacteria, A58 and A59 are likely modified to m^1A[36], and since *trmI*, the methylase that

generates this modification is present in *Mtb*[33], most *Mtb* tRNAs likely contain this modification as

well. Nucleoside variation in the sequence of tRNA genes can account for some of the variations in

modified sites between *E. coli* and *Mtb*. For example, in *Mtb*, termination signatures derived from G

at position 37 were detected in tRNA-Arg2, -Gln1, and -Gln2, whereas this position in these tRNAs

in *E. coli* are modified A, such as m^2A and m^6A, which are silent in tRNA-seq. Since m^1G induces

strong termination during reverse transcription, these positions are likely modified to m^1G, as

observed in the *Bacillus subtilis* tRNA-Arg2 gene position 37 G[37].

tRNA samples were also treated with several chemical treatments prior to sequencing, to

expand the set of tRNA modifications detectable by tRNAseq. These treatments included

iodoacetamide (IAA), for detection of sulfur modifications,1-cyclohexyl-(2-morpholinoethyl)

carbodiimide (CMC) for detection of Ψ, and alkali for detection of D and m^7G. The chemical

treatment protocols were first carried out with *E. coli*, to validate the methods.

IAA is a thiol-reactive compound that covalently attaches carboxyamidomethyl to thiolated

uridines via nucleophilic substitution[38] and modified s^4U is detected as C instead of U. In IAA-

treated samples, position 8 and 9, corresponding to s^4U in many tRNA species, had high

misincorporation frequencies, confirming that IAA treatment modifies s^4U, leading to elevated

misincorporation (Supplementary Fig. 2). Furthermore, we observed higher misincorporation and

termination signals at the positions corresponding to other sulfur modification, s^2C, s^2U and their

derivatives, such as position 32 in tRNA-Arg3, -Arg5, -Ser3, and Arg4, and 34 in tRNA-Glu, -Gln1,

and -Lys (Fig. 3 and Supplementary Fig. 2), revealing that IAA treatment facilitates the detection of

not only s^4U but also additional sulfur modifications, which are weakly detected without the IAA

treatment.

Figure 3. IAA treatment promotes detection of sulfur modifications on tRNAs by enhancing termination signals. (A, B) Heatmaps of the termination signals of *E. coli* tRNAs treated with (A) or without (B) IAA. Known modification sites, including sulfur modifications (s^4U, s^2C, s^2U in white) are shown. (C) Termination frequency at s^4U, s^2C, and s^2U sites of tRNAs treated with or without IAA.

Next, we applied IAA treatment to *Mtb* tRNA-seq. IAA treatment increased termination

signals from position 34 in tRNA-Glu1, -Gln1, and -Lys, which contain s^2U derivatives in *E. coli* (Fig.

4A). The 2-thiouridine modification is carried out by MnmA in *E. coli*[39], and a homolog, *Rv3024c*,

of this enzyme was identified in the *Mtb* genome (Supplementary Table 3)[40]. We used double

stranded DNA-based recombineering to delete Rv3024c in *Mtb,* yielding strain *MtbΔmnmA*[41].

Sequencing of tRNA isolated from *MtbΔmnmA* with prior IAA treatment showed reduced termination

signals from position 34 in tRNA-Glu1, -Gln1, and -Lys in treated samples, indicating that Rv3024c

plays a critical role in the modification responsible for increased termination frequency derived from

position 34 in these tRNAs (Fig. 4B and 4C). Together, these observations strongly suggest that

Rv3024c encodes an MnmA-like enzyme that sulfurates position 34 uridines in three *Mtb* tRNA

isoacceptors.

192
193
194

Fig. 4 Heat map and plot of early termination frequency from sequencing of tRNAs from wild
type and *MtbΔmnmA* with and without RNA alkylation. (A, B) Heat map of early termination
frequencies across tRNA molecules and positions for WT (A) and *MtbΔmnmA* (B). Sulfur
modification is shown in white. (C) Plot of termination frequencies at position 37 in wild type (WT)
Mtb and *MtbΔmnmA* for lysine_UUU, glutamate_UUG, and glutamine_UUC isoacceptors. IAA;
iodoacetamide.

As shown previously[42], CMC treatment increased both misincorporation and termination

signatures at a subset of Ψs in *E. coli* tRNAs (Fig 5, and Supplementary Fig. 3 and 4). In addition,

CMC-treated samples showed increased frequencies of both misincorporation and termination at

position 16, 17, 20, and 20A corresponding to D, and m^7G at position 46. Both modifications are

known to undergo base elimination in mild alkali conditions[43]. Since these signals were also

observed in the reaction condition in which CMC was not added, these signals are likely attributable

to the alkali treatment that is common to both conditions.

Reverse transcription-derived signatures derived from alkali-treated D showed a distinctive

pattern. With this treatment, when two Ds are at consecutive positions, e.g., D16 and D17, termination

signals were elevated at the following position, i.e., position 18 (Supplementary Fig. 3, 5 and 6).

Furthermore, alkali treatment also led to higher misincorporation frequencies at singlet Ds (Fig. 5,

Supplementary Fig. 5 and 6). Thus, termination and misincorporation signatures enabled the

prediction of known *E. coli* tRNA sites modified to D.

Fig. 5. CMC and alkali treatment facilitate detection of Ψ, D, and m⁷G modifications in *E. coli*. (A, B) Heatmaps of the misincorporation signals of *E. coli* tRNAs treated with (A) or without (B) CMC. In both conditions, tRNAs are incubated in alkali condition. Known Ψ, D, and m⁷G sites are shown. Ψ is shown in white. (C) Misincorporation frequency at known Ψ, D, m⁷G sites of tRNAs treated with CMC+/alkali+ or CMC-/alkali+, and tRNAs without treatment.

CMC/alkali treatment facilitated the identification of additional modifications in *Mtb* tRNAs.

Regardless of CMC treatment, alkali treated samples showed increased misincorporation and

termination frequencies derived from U located at position 16, 17, 20, and 20A (Fig. 6 and

Supplementary Fig. 7). As observed in *E. coli*, termination signals at position 18 and 21 likely

correspond to consecutive Ds at position 16 and 17, and 20 and 20A, respectively. *Mtb* Rv0823c is a

homolog of dihydrouridylase DusB (Supplementary Table 3), which likely accounts for the synthesis

of D at these positions. Furthermore, alkali treatment increased the misincorporation frequencies at

G at position 46 (Supplementary Fig. 8). Since Rv0208c is a homolog of TrmB, which synthesizes

m^7G at position 46 in *E. coli*, multiple *Mtb* tRNA species likely contain m^7G at position 46 (Fig. 6 and

Supplementary Fig. 8).

CMC treatment also increased the termination frequency at several sites. Termination

signatures derived from position 55, which is exclusively uridine in all tRNA species, increased in most

tRNA species, suggesting that *Mtb* tRNAs contain pseudouridines at this position[37]. Rv2793c is an

Mtb homologue of *E. coli* TruB and deletion of Rv2793c reduced the termination frequencies at this

position in tRNAs isolated from *Mtb_ΔtruB* (Fig. 6 and Supplementary Fig. 9)[41]. Together, these

observations suggest that Rv2793c encodes a TruB-like enzyme that modifies position 55 uridines to

Ψ across tRNA species. Furthermore, the presence of a TruA homolog in *Mtb* (Rv3455c) suggests

that in multiple tRNA species U at positions 38-40 can be modified to Ψ. Indeed, the termination

signatures derived from positions 38 and 39 increased depending on CMC treatment, strongly

suggesting that these positions are modified to Ψ (Fig. 6 and Supplementary Fig. 9).

In total, among 16 tRNA *Mtb* tRNA modifications predicted by the presence of tRNA

modifying enzymes (Fig 1 and Supplementary Table 3), 9 species of modifications were detected based

on reverse-transcription derived signatures (Fig. 2C).

Fig 6. Heat map of early termination frequency from sequencing of tRNAs isolated from WT and *MtbΔtruB* following CMC treatment. Heat map of early termination frequencies across tRNA molecules and positions for WT (left) and *MtbΔtruB* (right). Termination signals derived from position 55 are shown in white.

Growth of *MtbΔmnmA* is attenuated in a macrophage infection model

To address whether *Mtb* tRNA modifications impact the pathogen's growth in the host environment, we used the MtbTnDB transposon insertion sequencing (Tn-seq) database[44, 45] to determine if transposon insertions in genes encoding tRNA modifying enzymes have been associated with *in vivo* growth defects. Transposon insertions in *mnmA* were reported to attenuate *Mtb* growth in mice infected with a library of *Mtb* transposon mutants, suggesting that *mnmA* facilitates *Mtb* growth *in vivo*. We found that the growth of WT and *MtbΔmnmA* were similar in 7H9 medium (Fig. 7), suggesting the absence of s^2U modification at position 34 does not impair *Mtb* growth in culture. In contrast, the *MtbΔmnmA* mutant was significantly impaired for growth in a macrophage infection model[45] (Fig. 7). Defective growth of the *MtbΔmnmA* mutant was also observed in macrophages treated with all-trans retinoic acid (ATRA), which promotes macrophage control of *Mtb* infection[46].

These observations strongly suggest that modification of U to s^{2}U by MnmA facilitates *Mtb* growth in

macrophages.

Fig. 7. *MtbΔmnmA* is attenuated in a macrophage infection model. Top: Wild type and
MtbΔmnmA do not display growth differences in 7H9 medium. Bottom: Auto-luminescent wild type
and *ΔmnmA Mtb* strains were diluted to a multiplicity of infection of 2 bacteria per mouse bone
marrow-derived macrophage with or without all-trans retinoic acid (ATRA). Survival was measured
by luminescence and normalized to luminescence at time 0.

Discussion

Here, we profiled *Mtb* tRNA modifications with tRNA sequencing to provide the first maps of the tRNA modification landscape in this global pathogen. In total, nine modifications, including six modifications without chemical treatment, were identified based on reverse transcription derived signatures. CMC/alkali treatment and IAA treatment further identified Ψ and m^7G and sulfur modifications, respectively. Although we did not chemically validate the modifications predicted by tRNA-seq with mass spectrometry, the identification of *Mtb* homologs of tRNA modifying enzymes strongly bolsters the RT-signature-based predictions. Furthermore, the deletion of *truB* and *mnmA* genes in *Mtb* eliminated the respective modification signatures of pseudouridine and s^2U, validating that the enzymes encoded by these genes synthesize these modifications. Finally, the growth defect of the $\Delta mnmA$ strain within macrophages but not in a nutrient-rich medium suggests that s^2U tRNA modification facilitates *Mtb* adaptation to the host intracellular environment.

IAA treatment was developed for detecting s^4U modification in pulse-chase experiments to measure RNA turnover[38]. We found that this treatment enhances the RT-signatures of additional sulfur modifications as well, including s^2C and s^2U, indicating that IAA should have general utility in tRNA-seq-based profiling of sulfur modifications, including in studies tracking changes in tRNA sulfuration[47, 48].

Multiple modifications do not cause reverse transcriptase errors. Although tRNA-seq provides a simple and rapid method for profiling the landscape of modifications in all tRNA species, this approach does not enable comprehensive identification of modifications. Here, several modifications, including t^6A, m^5U, mo^5U, m^5C, Cm, Um, and Gm, predicted by the presence of *Mtb* homolog of tRNA modifying enzymes, such as TsaC2, TsaB, TsaD, TsaE, Trm2, TrcM, TrmR, TrmL, and TrmH, were not detected in tRNA-seq. Additionally, CMC treatment may not determine all Ψ positions, thus, the target of Rv3300 and Rv1540 remain unclear. Since these genes are similar to *E. coli rluA,* which

also targets rRNA, these genes may target rRNAs instead of tRNAs. Mapping and further analysis of

these modifications will require different approaches such as tRNA purification and RNA mass

spectrometric analysis.

Several strong RT signatures were detected in *Mtb* tRNAs but not in *E. coli* (Fig. 2A), including

G45 in tRNA-Gly2. Most of these signals are strict misincorporations of C or T at A or G position,

respectively. Since modifications at a purine position, e.g., m^1G and m^1A, generally cause random

misincorporation of the other three nucleosides, these *Mtb*-specific misincorporation signatures are

not likely to be derived from modification. Further mass spectrometric analysis of purified tRNAs will

be necessary to assess whether these signals are indeed derived from *Mtb*-specific modifications; such

Mtb-specific modifications could be related to *Mtb* pathogenesis and/or survival in the host.

The role of s^2U in *Mtb* appears to be unusual. s^2U is a universally conserved modification

observed in all three domains of life at the wobble position in the anticodon. This modification

enhances the stacking of s^2U with U35 to stabilize the anticodon structure, facilitating codon-

anticodon interactions. s^2U is also recognized by multiple aminoacyl-tRNA synthetases for efficient

amino acylation[12]. Although elimination of this sulfur modification causes severe growth

phenotypes in most organisms[39, 49], unexpectedly, the deletion of *mnmA* in *Mtb* did not attenuate

growth of the pathogen *in vitro*, suggesting that the *Mtb* requirement for s^2U modification differs from

other organisms. The marked growth retardation of Δ*mnmA* strain within macrophages indicates the

specific requirement of this modification within host cells. This modification may be necessary for

maintaining general translation efficiency inside host cells and/or facilitate the expression of specific

genes that are necessary for survival within macrophages. In fact, lack of *mnmA* is reported to sensitize

E. coli to oxidative stress, raising the possibility that s^2U modification promotes *Mtb* growth under

oxidative stress elicited by the host.

Differential codon usage between house-keeping genes and virulence genes could contribute

to the differential growth phenotypes observed *in vitro* and *in vivo* in the Δ*mnmA* mutant. Among

multiple codons encoding Glu, Gln, and Lys, s^2U modification dependent codon usage might be

preferentially distributed in genes associated with intracellular growth. For example, *Mtb* has two

tRNA isoacceptors, tRNA-Glu1(s^2UUC) and tRNA-Glu2(CUC), to decipher two Glu codons, GAA

and GAG. According to the wobble paring rule, GAA is only decoded by tRNA-Glu1(s^2UUC),

whereas GAG is decoded by both tRNA-Glu1(s^2UUC) and tRNA-Glu2(CUC); i.e., GAG can be

deciphered by an s^2U-independent tRNA. Thus, genes required for intracellular growth might be

enriched with GAA, an s^2U-dependent codon. Similar codon usage differences could be present in

Gln and Lys codons deciphered by s^2U-containing tRNAs.

In most organisms, s^2U is further modified into derivatives containing an additional chemical

moiety at position 5[2, 50, 51]. However, *Mtb* does not contain apparent homologs of the tRNA

modifying enzymes that introduce the additional modifications to s^2U. Thus, *Mtb* may contain s^2U or

s^2U derivatives synthesized by other types of enzymes. Additional analyses to elucidate the structures

of modified s^2U in *Mtb* are warranted.

Another unexpected finding is the presence of a TrmR homolog in *Mtb*. TrmR in *Bacillus*

subtilis is a methylase that converts a 5-hydroxyuridine (ho^5U) into a 5-methoxyuridine (mo^5U) at the

tRNA wobble position, suggesting that *Mtb* possesses mo^5U. However, *Mtb* does not encode an

apparent homolog for the enzyme that mediates the RNA hydroxylation reaction that yields ho^5U, the

substrate for TrmR. Although we cannot rule out the possibility that *Mtb* TrmR methylates a substrate

other than RNA, it is possible that *Mtb* utilizes an unknown hydroxylation pathway for synthesizing

ho^5U.

Our findings serve as a valuable starting point for the research community to continue

characterizing the physiological roles and mechanisms of *Mtb* tRNA modifications. Since tRNA-seq

offers an efficient and scalable platform for surveying changes in tRNA modifications, this approach

will be valuable across growth conditions, and may be extended to growth inside host cells. Finally,

further studies elucidating the mechanisms by which tRNA modifications facilitate *Mtb* growth in host

cells should be valuable for designing new therapeutics for tuberculosis.

Acknowledgement: We appreciate all the members of Waldor lab for fruitful discussion and

comments on the manuscript. We also thank Gregory Babunovic for his valuable assistance setting

up the macrophage infections used here. This work is supported by NIH/NIAID grants to M.K.W.

(R01AI-042347) and E.J.R. (P01AI-095208), a Dean's Innovation Award from Harvard Medical

School to E.J.R. and the Howard Hughes Medical Institute (HHMI) to M.K.W.

Materials and Methods

Bacterial strains and growth conditions: *Mtb* strains were grown from frozen stocks into

Middlebrook 7H9 medium supplemented with 0.2% glycerol, 0.05% Tween-80, and ADC (5 g/L

bovine serum albumin, 2 g/L dextrose, 3 μg/ml catalase). Cultures were incubated at 37 °C. Strains

were grown to mid log-phase for all experiments (OD_{600} 0.4-0.6). Growth was measured on a BioTek

plate reader for *in vitro* growth by measuring OD_{600} every 24 hours.

Bacterial strain construction: Supplementary Table 4 depicts the strains, plasmids, primers, and

recombinant DNA used for this study. Plasmids were built by restriction digest of a parental vector

and inserts were prepared by Gibson assembly[52]. Plasmids were isolated from *E. coli* and confirmed

via Sanger sequencing carried out by Genewiz, LLC (Massachusetts, USA).

370
Deletion mutants: The knockout strain *MtbΔmnmA*::zeo (zeocin) was built using double-stranded

recombineering in the parental *Mtb* strain H37Rv. A linear dsDNA fragment was constructed using

stitch PCR with the primers listed in Supplementary Table 4 which consisted of a 500bp region

upstream of *mnmA* (Rv3024c))*,* 500 bp downstream region, and a *lox*-zeo-*lox* fragment. This cassette

was transformed into an H37Rv recombineering strain as described[53] and plated on 7H10 + zeocin

plates.

Homology search

Local BLAST was performed to search for *Mtb* homologs of tRNA modifying enzymes. First, the

uniport IDs of tRNA modifying enzymes were obtained from Modomics[24], and 12 proteins were

manually added to the list [16,54−58], including Q47319/TapT, P24188/TrhO, P76403/TrhP,

O32034/TrhP1, O32035/TrhP2, P36566/CmoM, and Q87K36/TrcP, O34614/MnmM,

Q8N5C7/DTWD1, Q8NBA8/DTWD2, O32036/TrmR, Q9KV41/AcpA. Uniprot ID provides a

fasta file of tRNA modifying enzymes from the Uniprot database[59]. A blast database file was

generated by 'makeblastdb' script using a fasta file of *Mtb* proteins (H37Rv strain) retrieved from

NCBI. Then, the homologs of tRNA modifying enzymes were searched against the *Mtb* protein

database using the fasta file of tRNA modifying enzymes as a query. Output format is defined by the

following script: -outfmt "7 qacc sacc stitle score qcovs evalue pident" -evalue 1e-10. The output file

was modified by excel (Supplementary Table 4).

Phylogenetic analysis of *Mtb* tRNA modifying enzyme homologs

Local BLAST was conducted to search for homologs of *Mtb* tRNA modifying enzymes in 120

manually picked organisms across three domains of life. A custom database was generated by

combining the fasta files of organisms' proteins retrieved from NCBI. Homologs of eighteen *Mtb*

tRNA modifying enzymes were searched against the custom protein database using local blast.

Log10Evalues were visualized by iTol[34] with a phylogenetic tree generated by phyloT[60].

tRNA sequencing

<u>Extraction of total RNA:</u> Strains were grown to mid-log phase with the appropriate antibiotics and

inducing agents described above. RNA was collected at the same OD_{600} for each strain (between 0.4-

0.6). Cells were left on ice for 20 minutes, then pelleted by centrifuging at 4,000 rpm for 10 minutes

at 4 °C. Pellets were resuspended in 0.5-1 mL of TriZol (Life Technologies) and lysed using a BeadBug

microtube homogenizer (Millipore Sigma). 200 µL of chloroform was added to each tube, after which

samples obtained from *Mtb* strains were removed from biosafety level 3 precautions. Samples were

centrifuged at 15,000 rpm for 15 minutes at 4 °C and the aqueous layer was collected into a fresh tube.

To the original tube, 250 µL of sodium acetate buffer (300 mM sodium acetate pH 5.2 and 10 mM

EDTA pH 8.0) was added, and samples were vortexed at 4 °C for 5 minutes then centrifuged at 15,000

g for 15 minutes at 4 °C. The aqueous layer was added to the fresh sample-containing tubes. 400 μL

chloroform was added, and tubes were briefly vortexed and then centrifuged at 15,000 rpm for 1

minute at 4 °C. The aqueous phase was collected into a fresh tube and RNA recovered by ethanol

precipitation. RNA pellets were resuspended in 10 mM sodium acetate pH 5.2 and stored at -80 °C

until processed for sequencing. Total RNA samples were alkali-treated prior to tRNA extraction to

deacylate all tRNAs (1 hour at 37 °C in 100 mM Tris-HCl pH 9.0).

<u>Isolation of tRNA fraction</u>: 1−2 μg of total RNA was run on a 10% TBE-UREA gel (ThermoFisher

Scientific) at 250 V for 1 hour. Gels were stained with SYBR Gold (ThermoFisher Scientific), and

tRNA was excised. Excised gels containing tRNA fractions were mashed in RNAse-free tubes, and

300 μL elution buffer (300 mM NaOAc pH 5.5, 1 mM EDTA pH 8.0, 0.10% SDS) was added to each

tube. Samples were shaken on a thermoshaker (Eppendorf) for 1−4 hours at 37 °C and supernatant

was collected using an Ultrafree filter column (Millipore Sigma). tRNA was recovered by isopropanol

precipitation.

<u>tRNA dephosphorylation</u>: tRNA was dephosphorylated using QuickCIP (New England BioLabs)

according to manufacturer instructions, and tRNA was collected by phenol-chloroform extraction

followed by isopropanol precipitation.

<u>Iodoacetamide (IAA) treatment</u>: IAA treatment was performed as described[38]. Briefly, 500 ng of

total RNA is combined with 10 mM of iodoacetamide, 50 mM $NaPO_4$ pH 8.0, and 50% DMSO in a

final volume of 50 μL. Reactions were incubated at 50 °C for 15 minutes and quenched with DTT.

CMC treatment: CMC treatment was carried out as described in ref. Briefly, 2.5 µg tRNA fraction in

0.5 µl was mixed with 15 µl CMC-BEU buffer with or without CMC (0.34 M or 0 M CMC, 7 M urea,

4 mM EDTA, and 50 mM bicine pH 7.9) and incubated at 37 °C for 20 min. Adding 100 µl CMC

stop solution (0.3 M NaOAc pH 5.2 and 100 mM EDTA) quenched the reaction. RNA was desalted

with PD-10 desalting column (Cytiva) and recovered by ethanol precipitation. RNA was dissolved in

40 µl of 50 mM sodium carbonate buffer (pH 10.4) and incubated at 37 °C for 4 hours, followed by

ethanol precipitation.

Adapter ligation: 0.5 µL RNase inhibitor was added to 3.5µL dephosphorylated tRNA (200–250 ng

tRNA) and samples were boiled at 80 °C for 2 minutes. Boiled tRNA was mixed with 12 µL PEG

buffer mix (10 µL 50% PEG8000, 2 µL 10 × buffer B0216S; New England Biolabs). 3 µL of 5'

adenylated linkers (Supplementary Table 4) were added (33 pmol/µL) along with 1 µL T4 RNA ligase

2 truncated (New England BioLabs) and incubated at 25 °C for 2.5 hours. Samples were recovered by

isopropanol precipitation and run on a 10% TBE-Urea PAGE gel for 40 minutes at 250 V. Ligated

products were recovered by gel excision as described above.

Reverse transcription: Identical quantities of samples with different adapter sequences were pooled

for reverse transcription for a total of 200–250 ng tRNA. Reverse transcription was performed by

combining 2.1 µL dephosphorylated tRNA with 100 mM Tris-HCl pH 7.5, 0.5 mM EDTA, 1.25 µM

RT primer (Supplementary Table 4), 450 mM NaCl, 5 mM MgCl$_2$, 5 mM DTT, 500 nM TGIRT

(InGex), and 15% PEG8000 in a final volume of 9 µL. Samples were incubated at 25 °C for 30

minutes, after which 1 µL 10 mM dNTPs (New England BioLabs) were added and reactions incubated

at 60 °C for 1 hour. 1.15 μL NaOH was added, and samples were boiled for 15 minutes and run on a

10% TBE Urea PAGE gel at 250V for 1 hour. Reverse transcription products were excised from gels

and cDNA recovered by isopropanol precipitation. Linear single-stranded cDNA was circularized

using CircLigase II (Lucigen) in accordance with manufacturer instructions.

<u>PCR of tRNA libraries</u>: PCR reactions were set up using HF Phusion according to the manufacturer's

instructions using a universal reverse primer (Supplementary Table 4) and a different index primer for

each pool of samples. PCR reactions were aliquoted into 4 tubes and collected after 6, 8, 10, and 12

cycles. Samples were run on a Native TBE PAGE gel (ThermoFisher Scientific) at 180 V for 50

minutes, and amplified products were cut from the same cycle for each sequencing run. Samples were

recovered by gel excision and isopropanol precipitation.

<u>Sequencing</u>: Sequencing was performed on a MiSeq instrument (Illumina) using 150 bp single end

reads with a version 3, 150 cycle kit.

<u>Analysis</u>: 3' linker sequences and two nucleotides at the 5' end were trimmed using cutadapt and fastx-

trimmer. Bowtie v1.2.2 was used with default settings to map reads to reference *Mtb* tRNA sequences

retrieved from Mycobrowser[40] (Supplementary Table 5). Mpileup files were generated using

samtools (samtools mpileup -I -A --ff 4 -x -B -q 0 -d 10000000). For analysis of termination

frequencies, 5' end termini of mapped reads were piled up using bedtools genomecov (option, -d -5 -

ibam). The number of 5' termini at each tRNA position was divided by the total number of mapped

termini at that position plus all upstream (5') positions.

<u>Macrophage infection:</u> Auto-luminescent wild type and $\Delta mnmA$ *Mtb* strains grown to the same OD_{600}

were pelleted by centrifugation and prepared in RPMI media by soft spinning as described[61]. Briefly,

cells were washed, pelleted, resuspended, and centrifuged at 121 g, with the top half of the centrifuged

supernatant used. Suspensions were diluted to a multiplicity of infection of 2 bacteria per mouse bone

marrow-derived macrophage by determining the OD_{600}. Macrophages were infected for 6 hours,

followed by a PBS wash and addition of RPMI with or without all-trans retinoic acid (ATRA). ATRA

promotes macrophage control of *Mtb* infection[46] and was used to assess strain survival in an

increasingly restricted macrophage environment. Survival was measured by luminescence in a BioTek

plate reader and normalized to luminescence reads at time 0.

References

1. Huang, H.Y. and A.K. Hopper, *Multiple Layers of Stress-Induced Regulation in tRNA Biology.* Life (Basel), 2016. **6**(2).

2. Björk, G.R. and T.G. Hagervall, *Transfer RNA Modification: Presence, Synthesis, and Function.* EcoSal Plus, 2014. **6**(1).

3. Shepherd, J. and M. Ibba, *Bacterial transfer RNAs.* FEMS Microbiol Rev, 2015. **39**(3): p. 280-300.

4. Torrent, M., et al., *Cells alter their tRNA abundance to selectively regulate protein synthesis during stress conditions.* Sci Signal, 2018. **11**(546).

5. Orellana, E.A., E. Siegal, and R.I. Gregory, *tRNA dysregulation and disease.* Nat Rev Genet, 2022. **23**(11): p. 651-664.

6. Suzuki, T., *The expanding world of tRNA modifications and their disease relevance.* Nat Rev Mol Cell Biol, 2021. **22**(6): p. 375-392.

7. Liu, Y., et al., *tRNA-m(1)A modification promotes T cell expansion via efficient MYC protein synthesis.* Nat Immunol, 2022. **23**(10): p. 1433-1444.

8. Goodarzi, H., et al., *Modulated Expression of Specific tRNAs Drives Gene Expression and Cancer Progression.* Cell, 2016. **165**(6): p. 1416-1427.

9. Delaunay, S., et al., *Mitochondrial RNA modifications shape metabolic plasticity in metastasis.* Nature, 2022. **607**(7919): p. 593-603.

10. Nedialkova, D.D. and S.A. Leidel, *Optimization of Codon Translation Rates via tRNA Modifications Maintains Proteome Integrity.* Cell, 2015. **161**(7): p. 1606-18.

11. Kimura, S. and M.K. Waldor, *The RNA degradosome promotes tRNA quality control through clearance of hypomodified tRNA.* Proc Natl Acad Sci U S A, 2019. **116**(4): p. 1394-1403.

12. Giegé, R. and G. Eriani, *The tRNA identity landscape for aminoacylation and beyond.* Nucleic Acids Res, 2023.

13. de Crécy-Lagard, V. and M. Jaroch, *Functions of Bacterial tRNA Modifications: From Ubiquity to Diversity.* Trends Microbiol, 2021. **29**(1): p. 41-53.

14. Zhang, W., et al., *tRNA modification dynamics from individual organisms to metaepitranscriptomics of microbiomes.* Mol Cell, 2022. **82**(5): p. 891-906.

15. Zheng, G., et al., *Efficient and quantitative high-throughput tRNA sequencing.* Nat Methods, 2015. **12**(9): p. 835-837.

16. Kimura, S., P.C. Dedon, and M.K. Waldor, *Comparative tRNA sequencing and RNA mass spectrometry for surveying tRNA modifications.* Nat Chem Biol, 2020. **16**(9): p. 964-972.

17. Motorin, Y. and M. Helm, *Methods for RNA Modification Mapping Using Deep Sequencing: Established and New Emerging Technologies.* Genes (Basel), 2019. **10**(1).

18. Finet, O., et al., *Transcription-wide mapping of dihydrouridine reveals that mRNA dihydrouridylation is required for meiotic chromosome segregation.* Mol Cell, 2022. **82**(2): p. 404-419 e9.

19. Draycott, A.S., et al., *Transcriptome-wide mapping reveals a diverse dihydrouridine landscape including mRNA.* PLoS Biol, 2022. **20**(5): p. e3001622.

20. Dai, Q., et al., *Quantitative sequencing using BID-seq uncovers abundant pseudouridines in mammalian mRNA at base resolution.* Nat Biotechnol, 2022.

21. W.H.O., *Global tuberculosis report 2021.* 2021.

532   22.   Chionh, Y.H., et al., *tRNA-mediated codon-biased translation in mycobacterial hypoxic*
533   *persistence.* Nat Commun, 2016. **7**: p. 13302.

534   23.   Altschul, S.F., et al., *Basic local alignment search tool.* J Mol Biol, 1990. **215**(3): p. 403-
535   10.

536   24.   Boccaletto, P., et al., *MODOMICS: a database of RNA modification pathways. 2021*
537   *update.* Nucleic Acids Res, 2022. **50**(D1): p. D231-D235.

538   25.   DeJesus, M.A., et al., *Comprehensive Essentiality Analysis of the Mycobacterium*
539   *tuberculosis Genome via Saturating Transposon Mutagenesis.* mBio, 2017. **8**(1).

540   26.   Bosch, B., et al., *Genome-wide gene expression tuning reveals diverse vulnerabilities of*
541   *M. tuberculosis.* Cell, 2021. **184**(17): p. 4579-4592 e24.

542   27.   Masuda, I., et al., *tRNA methylation resolves codon usage bias at the limit of cell*
543   *viability.* Cell Rep, 2022. **41**(4): p. 111539. E.

544   28.   Soma, A., et al., *An RNA-modifying enzyme that governs both the codon and amino acid*
545   *specificities of isoleucine tRNA.* Mol Cell, 2003. **12**(3): p. 689-98.

546   29.   Wolf, J., A.P. Gerber, and W. Keller, *tadA, an essential tRNA-specific adenosine*
547   *deaminase from Escherichia coli.* EMBO J, 2002. **21**(14): p. 3841-51.

548   30.   El Yacoubi, B., et al., *The universal YrdC/Sua5 family is required for the formation of*
549   *threonylcarbamoyladenosine in tRNA.* Nucleic Acids Res, 2009. **37**(9): p. 2894-909.

550   31.   Anderson, J., et al., *The essential Gcd10p-Gcd14p nuclear complex is required for 1-*
551   *methyladenosine modification and maturation of initiator methionyl-tRNA.* Genes Dev,
552   1998. **12**(23): p. 3650-62.

553   32.   Guelorget, A., et al., *Insights into the hyperthermostability and unusual region-specificity*
554   *of archaeal Pyrococcus abyssi tRNA m1A57/58 methyltransferase.* Nucleic Acids Res,
555   2010. **38**(18): p. 6206-18.

556   33.   Varshney, U., et al., *Mycobacterium tuberculosis Rv2118c codes for a single-component*
557   *homotetrameric m1A58 tRNA methyltransferase.* Nucleic Acids Res, 2004. **32**(3): p.
558   1018-27.

559   34.   Letunic, I. and P. Bork, *Interactive Tree Of Life (iTOL) v5: an online tool for*
560   *phylogenetic tree display and annotation.* Nucleic Acids Res, 2021. **49**(W1): p. W293-
561   W296.

562   35.   Kellner, S., J. Burhenne, and M. Helm, *Detection of RNA modifications.* RNA Biol, 2010.
563   **7**(2): p. 237-47.

564   36.   Schwartz, M.H., et al., *Microbiome characterization by high-throughput transfer RNA*
565   *sequencing and modification analysis.* Nat Commun, 2018. **9**(1): p. 5353.

566   37.   Juhling, F., et al., *tRNAdb 2009: compilation of tRNA sequences and tRNA genes.*
567   Nucleic Acids Res, 2009. **37**(Database issue): p. D159-62.

568   38.   Herzog, V.A., et al., *Thiol-linked alkylation of RNA to assess expression dynamics.* Nat
569   Methods, 2017. **14**(12): p. 1198-1204.

570   39.   Kambampati, R. and C.T. Lauhon, *MnmA and IscS are required for in vitro 2-thiouridine*
571   *biosynthesis in Escherichia coli.* Biochemistry, 2003. **42**(4): p. 1109-17.

572   40.   Kapopoulou, A., J.M. Lew, and S.T. Cole, *The MycoBrowser portal: a comprehensive*
573   *and manually annotated resource for mycobacterial genomes.* Tuberculosis (Edinb),
574   2011. **91**(1): p. 8-13.

575   41.   van Kessel, J.C. and G.F. Hatfull, *Recombineering in Mycobacterium tuberculosis.* Nat
576   Methods, 2007. **4**(2): p. 147-52.

42.   Carlile, T.M., et al., *Pseudouridine profiling reveals regulated mRNA pseudouridylation*
in yeast and human cells. Nature, 2014. **515**(7525): p. 143-6.

43.   Marchand, V., et al., *Mapping of 7-methylguanosine (m(7)G), 3-methylcytidine (m(3)C),*
dihydrouridine (D) and 5-hydroxycytidine (ho(5)C) RNA modifications by AlkAniline-
Seq. Methods Enzymol, 2021. **658**: p. 25-47.

44.   Zhang, Y.J., et al., *Tryptophan biosynthesis protects mycobacteria from CD4 T-cell-*
mediated killing. Cell, 2013. **155**(6): p. 1296-308.

45.   Jinich, A., et al., *The Mycobacterium tuberculosis transposon sequencing database*
(MtbTnDB): a large-scale guide to genetic conditional essentiality. bioRxiv, 2021.

46.   Babunovic, G.H., et al., *CRISPR Interference Reveals That All-Trans-Retinoic Acid*
Promotes Macrophage Control of Mycobacterium tuberculosis by Limiting Bacterial
Access to Cholesterol and Propionyl Coenzyme A. mBio, 2022. **13**(1): p. e0368321.

47.   Edwards, A.M., K.A. Black, and P.C. Dos Santos, *Sulfur Availability Impacts*
Accumulation of the 2-Thiouridine tRNA Modification in Bacillus subtilis. J Bacteriol,
2022. **204**(5): p. e0000922.

48.   Laxman, S., et al., *Sulfur amino acids regulate translational capacity and metabolic*
homeostasis through modulation of tRNA thiolation. Cell, 2013. **154**(2): p. 416-29.

49.   Dewez, M., et al., *The conserved Wobble uridine tRNA thiolase Ctu1-Ctu2 is required to*
maintain genome integrity. Proc Natl Acad Sci U S A, 2008. **105**(14): p. 5459-64.

50.   Karlsborn, T., et al., *Elongator, a conserved complex required for wobble uridine*
modifications in eukaryotes. RNA Biol, 2014. **11**(12): p. 1519-28.

51.   Asano, K., et al., *Metabolic and chemical regulation of tRNA modification associated*
with taurine deficiency and human disease. Nucleic Acids Res, 2018. **46**(4): p. 1565-
1583.

52.   Gibson, D.G., et al., *Enzymatic assembly of DNA molecules up to several hundred*
kilobases. Nat Methods, 2009. **6**(5): p. 343-5.

53.   Murphy, K.C., K. Papavinasasundaram, and C.M. Sassetti, *Mycobacterial*
recombineering. Methods Mol Biol, 2015. **1285**: p. 177-99.

54.   Takakura, M., et al., *Biogenesis and functions of aminocarboxypropyluridine in tRNA.*
Nat Commun, 2019. **10**(1): p. 5542.

55.   Cho, G., J. Lee, and J. Kim, *Identification of a novel 5-aminomethyl-2-thiouridine*
methyltransferase in tRNA modification. Nucleic Acids Res, 2023. **51**(4): p. 1971-1983.

56.   Sakai, Y., S. Kimura, and T. Suzuki, *Dual pathways of tRNA hydroxylation ensure*
efficient translation by expanding decoding capability. Nat Commun, 2019. **10**(1): p.
2858.

57.   Sakai, Y., et al., *Biogenesis and growth phase-dependent alteration of 5-*
methoxycarbonylmethoxyuridine in tRNA anticodons. Nucleic Acids Res, 2016. **44**(2): p.
509-23.

58.   Kimura, S., et al., *Sequential action of a tRNA base editor in conversion of cytidine to*
pseudouridine. Nat Commun, 2022. **13**(1): p. 5994.

59.   UniProt, C., *UniProt: the Universal Protein Knowledgebase in 2023.* Nucleic Acids Res,
2023. **51**(D1): p. D523-D531.

60.   *PhyloT: a tree generator.*

61.   Saito, K., et al., *Rifamycin action on RNA polymerase in antibiotic-tolerant*
Mycobacterium tuberculosis results in differentially detectable populations. Proc Natl
Acad Sci U S A, 2017. **114**(24): p. E4832-E4840.