Cysteine-Rich Intestinal Protein 1 is a Novel Surface Marker for Myometrial Stem/Progenitor Cells.

Emmanuel N. Paul[1], Tyler J. Carpenter[1], Sarah Fitch[1,2], Rachael Sheridan[3], Kin H. Lau[3], Ripla Arora[1,2], Jose M. Teixeira[1*]

[1]Department of Obstetrics, Gynecology and Reproductive Biology, College of Human Medicine, Michigan State University, Grand Rapids, MI 48824, USA
[2]Institute for Quantitative Health Science and Engineering, East Lansing, MI 48824, USA
[3]Van Andel Research Institute, Grand Rapids, MI, USA.

*Corresponding Author:
Jose M. Teixeira, PhD, Michigan State University, 400 Monroe Ave NW, Grand Rapids, MI 49503. Email: teixei15@msu.edu

Acknowledgments:

The authors have no conflicts to declare.

Summary

Uterine fibroids are benign tumors that develop in the myometria of most women

with unknown etiology. Myometrial stem/progenitor cells (MyoSPCs) have been proposed

as the cells of origin for fibroids because uterine fibroids are almost always clonal. We

previously identified SUSD2 as a possible MyoSPC marker, but the relatively poor

enrichment in stem cell characteristics of SUSD2+ over SUSD2- cells compelled us to

find better discerning markers for more rigorous analysis. We combined bulk RNA-seq of

SUSD2+/- cells with single cell RNA-seq to identify markers capable of further enriching

for MyoSPCs. We observed seven distinct cell clusters within the myometrium, with the

vascular myocyte cluster most highly enriched for MyoSPC characteristics and markers,

including *SUSD2*. *CRIP1* expression was found highly upregulated in both techniques

and was used as a marker to sort CRIP1$^+$/PECAM1$^-$ cells that were enriched for colony

forming units and were able to differentiate into mesenchymal lineages, suggesting that

CRIP1$^+$/PECAM1$^-$ cells could be used to better understand the etiology of uterine fibroids.

Introduction

Uterine fibroids, also known as leiomyomas, are benign tumors found in the smooth muscle layer of the uterus, the myometrium. Although benign, uterine fibroids develop in up to 80% of women during their reproductive years and are often associated with debilitating symptoms such as menorrhagia, anemia, dysmenorrhea, fatigue, pelvic pain, urinary incontinence, and obstruction of the surrounding pelvic structures (1,2). Hormonal therapies, mainly used for alleviating fibroid symptoms, are generally short-term treatments due to long-term side effects or induced infertility (3), and hysterectomy, the most common and effective treatment for uterine fibroids, results in permanent infertility (4). Therefore, there is a clear need for effective medical therapies that preserve fertility and avoid invasive surgery. Understanding the disease etiology and pathogenesis could lead to alternative and less invasive therapies.

A dysregulated myometrial stem/progenitor cell (MyoSPC) has been proposed as the cell of origin for uterine fibroids. After embryonic development, tissue-specific stem cells remain throughout the body and play important roles in tissue homeostasis, including replacing dying cells and participating in tissue remodeling (5). The dramatic remodeling that occurs during pregnancy and following parturition in the uterus suggest a need for and the existence of myometrial stem cells (6). Uterine fibroids are thought to be a clonal disease (7-9), and since most clonal diseases have a single cell origin (10), we and others (11,12) have hypothesized that a mutated MyoSPC could be the cell of origin for uterine fibroids (13). The presence of cells with stem cell properties has been demonstrated using the label-retaining cells in a mouse model (14,15) and using the side population (SP) method in human myometrium (16,17). Putative MyoSPCs have been isolated and

studied by using a combination of cell surface markers, including CD44/Stro-1 (11) and

CD34/CD49f/b (18), or using the side population (SP) discrimination assay (16,17).

However, cell surface markers in these studies have been selected using stem cell

markers from other tissues or using the SP technique, which has multiple pitfalls. Indeed,

the SP assay is fallible to variation in staining conditions, as well as a heterogeneity of

the isolated cells, indicating that the SP phenotype is not exclusive to stem cells, and

more importantly, the dye used in SP assay has profound effects on cell survival (19).

Because the endometrial stroma and the myometrium originate from the same embryonic

tissue, the Müllerian duct mesenchyme (6), we recently proposed that SUSD2, an

endometrium stem cell marker (20) also enriches for MyoSPCs (21). While SUSD2+ cells

do have mesenchymal stem cell characteristics, SUSD2+ cells represent between 25-

40% of total myometrial cells. Additionally, colony formation is only increased 2-3 fold

increase in SUSD2[+] cells compared to the rest of the myometrium cells, suggesting that

further enrichment might be possible. The objective of the present study was to integrate

next-generation sequencing, including single cell RNA-seq and bulk RNA-seq, to identify

a more specific marker to significantly enrich for MyoSPCs from human myometrium,

which can then be used to better understand the molecular mechanisms underlying fibroid

etiology.

Results

SUSD2$^+$ are enriched for characteristic MSC genes compared with SUSD2$^-$ cells.

To determine how best to enrich for stem cell activities in the SUSD2+ MyoSPC population, we used SUSD2 to enrich for myometrium stem cells followed by RNA-seq to discover new cell surface markers for MyoSPCs in the human myometrium. Myometrial cells from non-fibroid patients (n = 5) were isolated and live SUSD2$^+$ and SUSD2$^-$ were sorted by flow cytometry. As with our previous results (21), 30-50% of the myometrium cells were SUSD2$^+$ (**Fig 1A**). Total RNA was isolated from the two cell populations and sequenced for differential gene expression analyses. Principal Component Analysis (PCA) plot showed that SUSD2$^+$ and SUSD2$^-$ cells were separated by principal component 1 with a variance of 39%, indicating a strong divergence in the transcriptomic profiles of these two cell populations (**Fig 1B**). A total of 6777 significant differentially expressed genes (DEGs) were detected between SUSD2$^+$ and SUSD2$^-$ myometrial cells with a false discovery rate (FDR) <0.05) (**Fig 1C and Table S1**). 3527 genes were down-regulated and 3250 were up-regulated in the SUSD2$^+$ population compared to the SUSD2$^-$ population. We confirmed that *SUSD2* was up-regulated in the SUSD2$^+$ sorted cells and were also enriched in other MSC markers such as *MCAM*, *PDGFRβ* and *CSPG4* (**Fig 1C and 1D**). A heatmap of the top 300 most DEGs from the SUSD2$^+$ compared to SUSD2$^-$ cells showed a good separation between cell types and included *SUSD2*, *MCAM*, *PDGFRβ* and *CSPG4* (**Fig 1E**).

Myometrium side population cells are not enriched in MSC markers.

The side population (SP) phenotype is another often used method to isolate cells with

stem cell characteristics that exploits the ability of some stem cells to efflux the DNA-

binding dye Hoechst 33342 via the ATP-binding cassette (ABC) transporters

(11,16,17,22,23). An average of 1.7% of the total myometrium cells were SP$^+$ (**Fig 2A**).

Addition of verapamil, a calcium channel blocker used as a negative control to validate

the SP, severely decreased of the number of the myometrium SP$^+$ cells (**Fig 2B**). SP$^+$

and SP$^-$ myometrium cells were sorted for total RNA sequencing and analyzed by PCA

plot, which showed that matched SP$^+$ and SP$^-$ cells segregated by the principal

component 2, accounting for 25% of the variance (**Fig 2C**). A total of 828 significant (FDR

<0.05) DEGs, including 478 upregulated genes and 350 downregulated genes, were

detected between the SP$^+$ and SP$^-$ myometrium cells (**Table S2**). The top 10 DEGs

enriched in the SP$^+$ compared to the SP$^-$ were associated with immune response (*XCL2*

(24), *CD69 (25)*, *IL7R (26)*, *KLRD1 (27)*, and *IL18R1 (28)*) apoptosis (*TNFRSF10A (29)*),

extracellular matrix (*SPOCK2 (30)*) and hematopoietic stem cell (*SELE (31)*, *GATA3 (32)*,

CD69 (33), and *VCAM1(34)*) (**Fig. 2D**). We confirmed an increase in expression of two

major ABC transporters, including *ABCB1*, and *ABCG2*, and a decrease in *PGR*, another

marker previously shown downregulated in the SP$^+$ compared to the SP$^-$ of human

myometrium cells (16) (**Fig 2E**). Surprisingly, SP$^+$ cells did not show increased expression

of the MSC markers, *SUSD2*, *MCAM*, *PDGFRβ*, *CSPG4*, *CD44*, *CD34* and *ITGA6* (also

known as *CD49f*) compared to the SP$^-$ (**Fig 2F**).

A putative MyoSPC cluster is determined by single cell RNA-seq.

Uniform Manifold Approximation and Projection (UMAP) of myometrial (n = 5) single cell

RNA-seq (scRNA-seq) revealed 7 main cell clusters (**Fig 3A**) with an equal cell

distribution across the five myometrium samples (**Fig S1**). Cluster identities were

assigned using the expression profiles of canonical markers for cell populations expected

to be found in the myometrium (**Fig 3B**) (35,36), including 4 different smooth muscle cell

types, vascular myocytes, myocytes, myofibroblasts, and fibroblast. The 4 MSC markers,

SUSD2, *MCAM*, *PDGFRβ* and *CSPG4*, were found highly expressed in the vascular

myocyte cluster, a known stem cell niche (**Fig 3C and 3D**). Known MSC characteristics

such as quiescent cell cycle state (G0) and the low gene regulation dynamics (37-39)

were determined by the cell cycle score and the velocity of the scRNA-seq data,

respectively. We identified a small group of cells within the vascular myocyte cluster in

the G1/G0 phase (**Fig 4A**) using a computational assignment of cell-cycle stage, as

described by Scialdone *et al.* (40). Cell velocity, which predicts the future state of

individual cells using the RNA splicing information from each cell (41), showed that the

same group of cells in G1/G0 phase in the vascular myocyte cluster are depicted with low

velocity vectors (**Fig 4B**) indicating low levels of transcriptional changes, another

characteristic of stem cells.

Combining bulk SUSD2 RNA-seq and myometrial scRNA-seq reveals a new

MyoSPC marker (CRIP1).

We defined cells within the vascular myocyte cluster presenting with high expression of

MSC markers in a G1/G0 phase and with low velocity as the "MyoSPC" cluster. A total of

6700 DEGs were found in the MyoSPC cluster compared to the rest of the myometrial

cells (**Table S3**). 54% (1929 DEGs) of the MyoSPC DEGs overlapped with the bulk RNA-

seq of SUSD2$^+$ compared to SUSD2$^-$ (**Fig 5A and Table S4**). Correlation analysis of the

log$_2$FC gene expression in the scRNA-seq analysis of the MyoSPC cluster compared to

the rest of the myometrial cells and the bulk RNA-seq of SUSD2+ compared with SUSD2-

cells by a scatter plot confirmed that the MSC markers, *SUSD2*, *MCAM*, *PDGFRβ* and

CSPG4 were upregulated in both. The most highly upregulated gene in the MyoSPC

cluster, Cysteine-Rich Intestinal Protein 1 (*CRIP1*), is also significantly upregulated in the

SUSD2+ cells (**Fig 5B**). UMAP plot showed that CRIP1 was highly expressed in the

vascular myocyte cluster, which includes MyoSPC cluster (**Fig 5C**), more particularly in

the defined MyoSPC cluster (**Fig 5D**). *CRIP1* expression wasn't differentially expressed

(log$_2$FC= -0.2, FDR p= 9.9x10^{-1}) in the RNA-seq results of the SP assay (**Fig S2**).

CRIP1+ cells have common stem/progenitor cell properties

We next investigated the CRIP1+ cells to establish their stem cell bona fides.

Immunofluorescence analysis using 3D imaging of the myometrial layer showed that

CRIP1+ cells are located surrounding the vascular endothelial cells (PECAM1$^+$, also

known as CD31), a known mesenchymal stem cell niche (42-44) (**Fig 6A**). Interestingly,

CRIP1$^+$ cell immunofluorescence was localized mostly near the larger blood vessels and

within a subset of SUSD2$^+$ cells. Flow cytometry revealed that CRIP1$^+$ cells represented

between 2 to 5% of the total myometrium cells (**Fig 6B**), a stem cell ratio expected to be

found in adult tissues (45). PECAM1 was used for negative selection of the smaller

population of endothelial cells that co-expressed CRIP1. CRIP1$^+$/PECAM1$^-$ cells and the

depleted cell population were sorted, and typical downstream stem cell assays were

performed to determine if the CRIP1$^+$/PECAM1$^-$ cells have stem/progenitor cell

proprieties. Colony formation assays indicated that CRIP1$^+$/PECAM1$^-$ sorted cells have a

greater self-renewal capacity compared to the depleted sorted population (**Fig 6C**), with

a significant increase of 4.5-fold greater number of colonies formed (**Fig 6D and 6E**), as

well as a significant increase of the size of the colonies (**Fig 6F**). After 5 days in smooth

muscle differentiation media, CRIP1$^+$/PECAM1$^-$ cells were positive for ACTA2, indicating

that they differentiated into smooth muscle cells (**Fig 6G**). What about depleted cells?

Similarly, CRIP1$^+$/PECAM1$^-$ cells were positive for the Oil Red O staining (**Fig 6H**), and

the alkaline phosphatase activity (**Fig 6I**) when grown in either in adipogenic or

osteogenic differentiation media, respectively, compared to CRIP1$^+$/PECAM1$^-$ cells

grown in control media, indicating that these putative MyoSPC cells have the capacity to

differentiate into adipocytes and osteocytes.

Discussion

We have identified CRIP1 as a novel cell surface marker that enriches for a

possible MyoSPC by combining the analyses of two next generation sequencing

techniques, bulk RNA-seq from SUSD2$^+$ and SUSD2$^-$ cells and scRNA-seq of total

myometrial samples. Bulk RNA-seq, enriched for known MSC markers, but the number

of DEGs were many, making it difficult to choose, identify and validate any possible

myometrium stem cell marker for further study. To reduce the number of candidate

markers for validation and follow up studies, we used the scRNA-seq to identify possible

stem cells based on MSC markers. Subsequent stem cell assays confirmed that CRIP1+

cells have MSC properties, and further studies are underway to determine whether these

cells could be a cell of origin for uterine fibroids.

CRIP1 was a highly upregulated gene in both SUSD2+ cells and the MyoSPC cluster we

identified by scRNA-seq. *CRIP1*, Cysteine-rich intestinal protein 1, is a member of the

LIM/double zinc-finger proteins that is predicted to be a novel biomarker in multiple

cancers and can promote several biological processes, including cell migration, invasion

and epithelial-mesenchymal transition by activating the Wnt/β-catenin signaling an

essential pathway in stem cell homeostasis (46-48). *CRIP1* was upregulated in the SP of

fibroid tissues along with 153 other DEGs by microarray analysis (49). In that study, the

authors used one of the stem cells characteristics, the increased expression ad activity

of ABC transporters, to isolate SP$^+$ cells from fibroid tissues. They demonstrated that the

SP$^+$ cells from fibroid tissues have stem cell characteristics, including self-renewal, and

differentiation into adipose and osteocyte cells. We performed similar experiments using

myometrial samples from non-fibroid patients (MyoN) and also found an increase in the

expression of major ABC transporters in the SP$^+$ compared to the SP$^-$ of human

myometrium cells We used MyoN samples because we recently reported that

myometrium from fibroid patients (MyoF) have a different transcriptomic profile compared

to MyoN samples, including an enrichment of DEGs in a leiomyoma disease ontology

panel (50). Here we have reported that *CRIP1* expression was not differentially expressed

and that the expression of the MSC markers, *SUSD2*, *MCAM*, *PDGFRβ*, and *CSPG4*

were decreased in SP$^+$ compared to the SP$^-$ from MyoN cells. The discordant results from

these studies could arise from the tissue type, that is, fibroid tumor versus MyoN, or

because the SP technique could be more applicable to other tissues (19) or hematopoietic

stem cells (HSC). It is worth noting that HSC exhibit a specific ABC transporter gene

expression profile distinct from other stem cells, including MSC (51). HSC expressed

higher level of most of the ABC transporters including *ABCB1*, *ABCC1* and *ABCG2*

compared to other stem cells. Indeed, the SP+ MyoN cells were enriched for ABC

transporters and HSC-associated genes. Additionally, the SP technique relies on an intact

cell metabolism and considerable variation in results is observed (19).

Our scRNA-seq results suggested that human myometrium has at least 7 different

cell types, including different types of smooth muscle cells, endothelial cells, and immune

cells. Similar clusters were reported in a scRNA-seq comparison of fibroids and

myometrium (36). The depth of sequencing for each cell was close to saturation allowing

us to identify a small cell population with stem cell characteristics, including the

expression of MSC markers *SUSD2*, *MCAM*, *PDGFRβ*, and *CSPG4*, a quiescent (G0)

cell cycle state (37) and low transcriptomic activity/low RNA velocity (37-39).

CRIP1$^+$/PECAM1$^-$ cells were primarily located in the perivascular region, a known stem

cell niche (42-44), particularly by the larger myometrial blood vessel. These results were

consistent with our scRNA-seq results showing that *CRIP1* expression was highly

expressed in the vascular myocyte cluster. Moreover, immunofluorescence staining

showed that CRIP1[+] cells were a subset the SUSD2[+] cells and that CRIP1[+]/PECAM1[-]

cells account for only 2 to 5% of the total human myometrium cells, a stem cell proportion

expected to be in an adult tissue (45). Interestingly, the depleted population was able to

form a few colonies, an indication that some cells in the depleted population still have a

self-renewal properties. Similar results were observed by our group using other MSC

markers to isolate myometrium stem/progenitor cells, including SUSD2, MCAM, or

PDGFRβ (21). This finding suggests that further enrichment of the MyoSPCs with some

of the other MSC markers could be done.

Study approval

The use of human tissue specimens was approved by the Spectrum Health Systems and

Michigan State University Institutional Review Boards (MSU IRB Study ID:

STUDY00003101, SR IRB #2017-198) as secondary use of biobank materials.

Author contributions

Experimental design (E.N.P, J.M.T) collected data and performed experiments (E.N.P,

T.J.C, S.F), analyzed data (E.N.P, R.S, K.H.L, R.A, J.M.T), wrote/reviewed manuscript

(E.N.P, T.J.C, S.F, R.S, K.H.L, R.A, J.M.T).

Acknowledgments

We would like to thank the patients who consented for the study, the Spectrum Health

Systems Universal Biorepository staff, and the Van Andel Institute Genomics Core

(RRID:SCR_022913), especially Marie Adams, Rebecca Siwicki and Julie Koeman, for

their assistance with the construction and sequencing of 10X libraries for the single cell

RNA-seq and bulk RNA-seq.This work was supported by grants HD096259 and

HD100959 from the Eunice Kennedy Shriver National Institute of Child Health and Human

Development (to J.M.T) and the SRI/Bayer Discovery/Innovation Grant (to E.N.P).

Methods

Sample collection and cell isolation

The use of human tissue specimens was approved by the Spectrum Health Systems Institutional Review Board as secondary use of biobank materials. Myometrial samples from non-fibroid patients (MyoN) were obtained following total hysterectomy from pre-menopausal, self-identified Caucasian women. No fibroids were detected by ultrasound prior to surgery. All patients who participated to the study gave consent to donate tissue through the Spectrum Health Biorepository. Tissues were washed with phosphate-buffered saline, dissected away from non-myometrial tissue, and minced. Cells were isolated in baffled flask with digestion media (DMEM/F12, Thermo Fisher, #SH3027101) containing 10% fetal bovine serum (FBS), collagenase type I (Sigma, #C0130), DNAse type I (Sigma, #10104159001) and $MgCl_2$ and incubated with agitation at 37°C. The resulting cell suspensions were strained through 100- and 40-µm cell strainers, washed with warm media (DMEM/F12) containing 15% FBS, and centrifuged. Isolated cells were then stored in freeze media (90% FBS, 10% DMSO, Thermo Fisher, #BP231-100) at -80 °C until needed.

Cell staining for FACS

Human primary myometrium cells were thawed and resuspended in 1% bovine serum albumin blocking buffer for 20 min at room temperature (RT). Cells were then incubated with the primary antibody for 45 min at RT; SUSD2-PE anti-human (Miltenyi Biotec, #130-117-682), PECAM1-FITC anti-human (Thermo Fisher, #11-0319-42), and CRIP1 rabbit anti-human (Thermo Fisher, #PA5-24643). For CRIP1/PECAM1 staining, cells were

incubated with a secondary antibody for 30 min at RT using an Alexa-647 anti-rabbit

(Thermo Fisher, #A21244). Stained myometrium cells were then wash with flow buffer,

and resuspend in 1 mL of flow buffer with 1 µg of 4′,6-diamidino-2-phenylindole (DAPI) or

Propidium Iodode (PI), depending on the experiment, for live dead discrimination. Cells

were sorted by the flow cytometry core at Van Andel Research Institute (VARI) using a

FACSymphony S6 cytometer (BD Biosciences) and analyzed with FlowJo Software (BD

Biosciences, version 10.8.1). Cells were sorted in media (DMEM/F12) at 4C.

The side population assay was conducted as described previously (17). Briefly, live cells

were labeled in a water bath at 37 °C for 90 min with 5 ug/mL of Hoechst 33342 dye

(Thermo Fisher). As a negative control, separate aliquots of cells from the same patients

were treated with 25 mg/ml of verapamil (Sigma) prior to addition of the Hoechst dye. PI

was added to stained cells with and without verapamil treatment and analyzed in a MoFlo

Astrios (Beckman Coulter) for side population gating by Hoechst red and blue filters and

sorting in media (DMEM/F12) at 4 °C.

RNA Isolation, Library Preparation and Sequencing

Sorted cells (SUSD2$^+$, SUSD2$^-$, SP$^+$ and SP$^-$) were lysed and isolated using RNeasy mini

kit (Qiagen, #74104). Isolated RNA was stored at −80 °C in nuclease-free water.

Nanodrop 1000 spectrophotometer (Thermo Fisher Scientific, Fair-lawn, NJ, USA) and

Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA) instruments were

used to measure RNA concentration and quality, according to the manufacturers'

protocols. RNA integrity values of ≥7.5 were required for library preparation and paired-

end (2 × 100 bp) RNA-sequencing on an Illumina NextSeq 6000 instrument (Illumina Inc.,

San Diego, CA, USA). Libraries were prepared using a Kapa RNA HyperPrep kit with

ribosomal reduction, pooled, and sequenced on flowcells to yield approximately 50–60

million reads/sample.

For single cell RNA-seq, dead cells from isolated myometrium cells were removed using

the Dead Cell Removal Kit (Miltenyi Biotec, #130-090-101) per manufacturer's

instructions. Live myometrium cells from 5 non-fibroid patients were then sequenced.

Libraries were generated and sequenced using the 10X Chromium Next GEM Single Cell

3' GEM kit (10X Genomics, v2) platform according to the manufacturer's instruction to

target an output of 7,000 cells per sample. Quality and quantity of the finished libraries

were assessed using a combination of Agilent DNA High Sensitivity chip (Agilent

Technologies, Inc.) and QuantiFluor® dsDNA System (Promega Corp., Madison, WI,

USA). 2 x 75 bp, paired end sequencing was performed on an Illumina NovaSeq 6000

sequencer using an S2, 100 cycle sequencing kit (v1.5) to a minimum depth of 50K reads

per cell (Illumina Inc., San Diego, CA, USA). Base calling was done by Illumina RTA3

and output was demultiplexed and converted to FastQ format with Cell Ranger (10X

Genomics, v3.1.0).

RNA Seq Analysis

For bulk RNA-seq, reads from 5 MyoN SUSD2+ and SUSD2+ cells and 3 MyoN SP[+] and

SP[-] cells, were trimmed for quality and adapters using TrimGalore (version 0.6.5), and

quality trimmed reads were assessed with FastQC (version 0.11.7). Trimmed reads were

mapped to Homo sapiens genome assembly GRCh38 (hg38) using STAR (version

2.7.9a) (72). Reads overlapping Ensembl annotations (version 99) were quantified with

STAR prior to model-based differential expression analysis using the edgeR-robust

method with paired samples. Genes with low counts per million (CPM) were removed

using the filterByExpr function from edgeR (72). Scatterplots of two selected principal

components was constructed with the pca function of the PCAtools R package (version

2.5.13) to verify sample separation prior to statistical testing. Generalized linear models

were used to determine if principal components were significantly associated with cell

type. Genes were considered differentially expressed if their respective edgeR-robust

false discovery rates (FDR) corrected p-values were less than 0.05. Differential

expression was calculated by comparing SUSD2$^+$ versus SUSD2$^-$ cells. Differentially

expressed genes (DEG) were visualized with volcano plots and heatmaps generated

using the EnhancedVolcano (version 1.8.0) and pheatmap (version 1.0.12) packages in

R. Box plots of the $\log_2$(CPM) values were generated using the R package ggplot2

(version 3.4.0).

A total of 9,775 total cells sent for sc-RNA-seq from 5 MyoN samples with ~512,000 reads

per cell passed the quality control. Demultiplexed raw sequencing reads were processed

and aligned to the Homo sapiens genome assembly GRCh38 (hg38) using STAR (version

2.7.9a) by the 10X Genomics Cell Ranger (version 3.1.0). Samples were merged using

the integration anchors function of the Seurat package (version 4.2.1) from R. Genes

expressed in fewer than three cells in a sample were excluded, as well as cells that

expressed fewer than 200 genes and mitochondrial gene content >5% of the total unique

molecular identifier count. Data were normalized using a global-scaling normalization

method that normalizes the feature expression measurements for each cell by the total

expression, multiplies this by a scale factor (10,000), and then log-transforms the results.

The top 2,000 most variable genes that were used for cell clustering were found using the

FindVariableFeatures function and were then normalized using the *ScaleData* function.

To determine the dimensionality of the dataset, an elbow plot was generated using the

Elbowplot function of Seurat. Based on the Elbow plot, we selected 15 principal

components (PC) for downstream analyses. Cell clusters were generated using

FindNeighbors and *FindClusters* functions. For visualization UMAPs were generated

using the *RunUMAP*, *FeaturePlot* and *DimPlot* fuctions. The *DotPlot* Seurat fuction was

used to generate dot plots to visualize gene expression for each assigned cluster. Cell

cycle score and velocity were determined using the functions *CellCycleScoring* from

Seurat and *RunVelocity* from SeuratWrappers (version 0.3.0), respectively. The "stem

cell" cluster was selected using the *CellSelector* function from Seurat. *RidgePlot* function

from the Seurat R package was use for the visualization of *CRIP1* gene expression in the

different myometrium clusters.

Venn diagrams of the overlapping DEG from the Bulk RNA-seq of SUSD2$^+$ and SUSD2$^-$

cells and DEG of scRNA-seq of the "stem cell" cluster vs the rest of the myometrium cells

were constructed using the eulerr package (version 6.1.1). The scatter plot of overlapping

DEG from bulk- and scRNA-seq was generated with ggplot2 (version 3.4.0).

Imaging

Whole mount immunofluorescent staining (52) was performed for human myometrial

samples that were fixed in a 4:1 solution of Methanol:DMSO and stored at -20 °C. Tissue

was removed from the fixative and rehydrated in (1:1) methanol: PBST (PBS, 1% triton)

solution for 15min followed by a 15min wash in 100% PBST. Samples were incubated in

a blocking solution (PBS, 2% powdered milk, 1% triton) for 1 h at room temperature and

then stained with a 1:500 dilution for primary antibodies in blocking solution for 7 nights

at 4°C. Primary antibodies used were Rabbit anti-human CRIP1 (Thermofisher, #PA5-

24643), Mouse anti-human SUSD2 (Biolegend #327401) and Mouse anti-human

PECAM1 (Abcam, #ab9498). Samples were then incubated with secondary antibodies,

Donkey anti-Rabbit IgG Alexa Fluor 555 (Invitrogen, #A31572), Goat anti-Mouse IgG

Alexa Fluor 647 (Invitrogen, #A21235) at a dilution of 1:300 and Hoechst dye (Sigma-

Aldrich, #B2261), for three nights at 4°C. Samples were transferred to a Methanol:PBST

(1:1) solution for 15 minutes, washed in 100% Methanol for 30 minutes, and incubated at

4°C overnight in a 3% H_2O_2 solution (Sigma-Aldrich, #216763) diluted in Methanol.

Tissues were then washed in 100% Methanol for 30 minutes and then cleared in Benzyl

Alcohol:Benzyl Benzoate (1:2) (Sigma-Aldrich, #108006, #B6630) overnight. Imaging

was performed using a Leica SP8 TCS white light laser confocal microscope utilizing 10x

air objective and a 7.0 µm Z stack (53). Imaris v9.2.1 (Bitplane) commercial software was

used to analyze confocal image files and create 3D renderings. The image files were

imported into Imaris 3D surpass mode and 3D renderings were created using the Surface

plugin. Images were captured using the Snapshot plugin of Imaris and videos were

generated using the Animation plugin.

Colony formation

After isolation from human myometrium tissues of CRIP1$^+$/PECAM1$^-$ cells and the

depleted population, cells were plated at colony forming density (50 cells/cm^2) in triplicate

in growth media (DMEM/F12, 10% FBS, 1× antibiotics/antimycotics) overnight and then

grown in MesenPro RS (Thermo Fisher, # 12746012) for 2 to 3 weeks. Cultures were

fixed in 4% paraformaldehyde (PFA) and stained with crystal violet for colony

visualization. Colonies were counted and total surface area was estimated using ImageJ

(version 1.53k), and the percent colony-forming units (CFUs) was calculated as (number

of colonies/number of cells plated) × 100 and averaged for triplicates. Matched

CRIP1$^+$/PECAM1$^-$ and depleted populations were cultured in different wells of the same

plate, and therefore culture was stopped for both cell types on the same day. Images

were taken using a Nikon SMZ18 microscope and Ds-Ri1 camera (Nikon Instruments

Inc.).

Mesenchymal lineage differentiation

For osteogenic and adipogenic differentiation, CRIP1$^+$/PECAM1$^-$ cells were plated in

triplicate at 50% to 80% confluency in growth media (DMEM/F12, 10% FBS, 1×

antibiotics/antimycotics) overnight and then cultured for 10 days in fresh StemPro

Adipogenesis Differentiation (Thermo Fisher Scientific, #A1007001) or StemPro

Osteogenesis Differentiation (Thermo Fisher Scientific, #A1007201) media according to

the manufacturer's instructions. Cells were cultured in regular growth media to serve as

differentiation controls. To assay adipogenic differentiation, cultures were fixed in 4%

PFA, stained using Oil Red O (Sigma, #01391) according to the manufacturer's

instructions. To assay osteogenic differentiation, cultures were stained for alkaline

phosphatase activity using the Alkaline Phosphatase (AP), Leukocyte kit (Sigma, #86R-

1KT) according to the manufacturer's instructions. For smooth muscle differentiation,

CRIP1$^+$/PECAM1$^-$ cells were plated on 1 mg/ml dried rat tail collagen (Corning, #354 236)

8 well chamber slide with growth media (DMEM/F12, 10% FBS, 1×

antibiotics/antimycotics) overnight and then cultured in Medium 231 with a smooth muscle

differentiation supplement (Thermo Fisher, #M231500). Cells were fixed in 4% PFA at the

indicated times (D0: before adding the differentiation media, D5: 5 days in differentiation

media) and stained using αSMA-Cy3 (Sigma, #C6198). Images were taken using a Nikon

Eclipse Ni-U or Nikon SMZ18 microscope and Ds-Qi1MC or Ds-Ri1 camera (Nikon

Instruments Inc.).

Statistical analyses

Bioinformatic statistics were performed using the listed packages in R (version 4.0.2).

Differentially expressed genes of the bulk RNA-seq from SUSD2$^+$ vs. SUSD2$^-$ and the

SP$^+$ vs. SP$^-$ were identified as those having an Benjamini–Hochberg FDR corrected p

<0.05 (54). Data with unequal variances were log transformed, and homogeneity of

variances verified before completion of analyses. Differential expressed genes of the

single cell RNA-seq were calculated using the non-parametric Wilcoxon Rank Sum test.

Adjusted p-value, based on Bonferroni correction using all features in the dataset was

used to determine significance. Genes with adjusted p-value < 0.05 were consider as

significant. For the colony formation assay, comparison of two means was performed with

a two-sided student t test, and significance was determined at p <0.05 after confirming

normal distribution using Graphpad Prism (version 9.4.1).

References

1. Zimmermann A, Bernuit D, Gerlinger C, Schaefers M, Geppert K. Prevalence, symptoms and management of uterine fibroids: an international internet-based survey of 21,746 women. *BMC Womens Health*. 2012;12:6.

2. Gupta S, Jose J, Manyonda I. Clinical presentation of fibroids. *Best Pract Res Clin Obstet Gynaecol*. 2008;22(4):615-626.

3. Farris M, Bastianelli C, Rosato E, Brosens I, Benagiano G. Uterine fibroids: an update on current and emerging medical treatment options. *Ther Clin Risk Manag*. 2019;15:157-178.

4. Sohn GS, Cho S, Kim YM, Cho CH, Kim MR, Lee SR, Working Group of Society of Uterine L. Current medical treatment of uterine fibroids. *Obstet Gynecol Sci*. 2018;61(2):192-201.

5. Ono M, Maruyama T. Stem Cells in Myometrial Physiology. *Semin Reprod Med*. 2015;33(5):350-356.

6. Teixeira J, Rueda BR, Pru JK. Uterine stem cells. StemBook. Cambridge (MA)2008.

7. Canevari RA, Pontes A, Rosa FE, Rainho CA, Rogatto SR. Independent clonal origin of multiple uterine leiomyomas that was determined by X chromosome inactivation and microsatellite analysis. *Am J Obstet Gynecol*. 2005;193(4):1395-1403.

8. Zhang P, Zhang C, Hao J, Sung CJ, Quddus MR, Steinhoff MM, Lawrence WD. Use of X-chromosome inactivation pattern to determine the clonal origins of uterine leiomyoma and leiomyosarcoma. *Hum Pathol*. 2006;37(10):1350-1356.

9. Holdsworth-Carson SJ, Zaitseva M, Vollenhoven BJ, Rogers PA. Clonality of smooth muscle and fibroblast cell populations isolated from human fibroid and myometrial tissues. *Mol Hum Reprod*. 2014;20(3):250-259.

10. Fialkow PJ. Clonal origin of human tumors. *Annu Rev Med*. 1979;30:135-143.

11. Mas A, Nair S, Laknaur A, Simon C, Diamond MP, Al-Hendy A. Stro-1/CD44 as putative human myometrial and fibroid stem cell markers. *Fertil Steril*. 2015;104(1):225-234 e223.

12. Yin P, Ono M, Moravek MB, Coon JSt, Navarro A, Monsivais D, Dyson MT, Druschitz SA, Malpani SS, Serna VA, Qiang W, Chakravarti D, Kim JJ, Bulun SE. Human uterine leiomyoma stem/progenitor cells expressing CD34 and CD49b initiate tumors in vivo. *J Clin Endocrinol Metab*. 2015;100(4):E601-606.

13. Bulun SE. Uterine fibroids. *N Engl J Med*. 2013;369(14):1344-1355.

14. Szotek PP, Chang HL, Zhang L, Preffer F, Dombkowski D, Donahoe PK, Teixeira J. Adult mouse myometrial label-retaining cells divide in response to gonadotropin stimulation. *Stem Cells*. 2007;25(5):1317-1325.

15. Patterson AL, George JW, Chatterjee A, Carpenter T, Wolfrum E, Pru JK, Teixeira JM. Label-Retaining, Putative Mesenchymal Stem Cells Contribute to Myometrial Repair During Uterine Involution. *Stem cells and development*. 2018.

16. Ono M, Maruyama T, Masuda H, Kajitani T, Nagashima T, Arase T, Ito M, Ohta K, Uchida H, Asada H, Yoshimura Y, Okano H, Matsuzaki Y. Side population in human uterine myometrium displays phenotypic and functional characteristics of myometrial stem cells. *Proc Natl Acad Sci U S A*. 2007;104(47):18700-18705.

17. Chang HL, Senaratne TN, Zhang L, Szotek PP, Stewart E, Dombkowski D, Preffer F, Donahoe PK, Teixeira J. Uterine leiomyomas exhibit fewer stem/progenitor cell characteristics when compared with corresponding normal myometrium. *Reprod Sci*. 2010;17(2):158-167.

18. Ono M, Kajitani T, Uchida H, Arase T, Oda H, Uchida S, Ota K, Nagashima T, Masuda H,
Miyazaki K, Asada H, Hida N, Mabuchi Y, Morikawa S, Ito M, Bulun SE, Okano H, Matsuzaki
Y, Yoshimura Y, Maruyama T. CD34 and CD49f Double-Positive and Lineage Marker-
Negative Cells Isolated from Human Myometrium Exhibit Stem Cell-Like Properties
Involved in Pregnancy-Induced Uterine Remodeling. *Biol Reprod*. 2015;93(2):37.

19. Golebiewska A, Brons NH, Bjerkvig R, Niclou SP. Critical appraisal of the side population
assay in stem cell and cancer stem cell research. *Cell Stem Cell*. 2011;8(2):136-147.

20. Masuda H, Anwar SS, Buhring HJ, Rao JR, Gargett CE. A novel marker of human
endometrial mesenchymal stem-like cells. *Cell Transplant*. 2012;21(10):2201-2214.

21. Patterson AL, George JW, Chatterjee A, Carpenter TJ, Wolfrum E, Chesla DW, Teixeira JM.
Putative human myometrial and fibroid stem-like cells have mesenchymal stem cell and
endometrial stromal cell properties. *Human reproduction*. 2020;35(1):44-57.

22. Redvers RP, Li A, Kaur P. Side population in adult murine epidermis exhibits phenotypic
and functional characteristics of keratinocyte stem cells. *Proc Natl Acad Sci U S A*.
2006;103(35):13168-13173.

23. Schienda J, Engleka KA, Jun S, Hansen MS, Epstein JA, Tabin CJ, Kunkel LM, Kardon G.
Somitic origin of limb muscle satellite and side population cells. *Proc Natl Acad Sci U S A*.
2006;103(4):945-950.

24. Fox JC, Nakayama T, Tyler RC, Sander TL, Yoshie O, Volkman BF. Structural and agonist
properties of XCL2, the other member of the C-chemokine subfamily. *Cytokine*.
2015;71(2):302-311.

25. Lopez-Cabrera M, Santis AG, Fernandez-Ruiz E, Blacher R, Esch F, Sanchez-Mateos P,
Sanchez-Madrid F. Molecular cloning, expression, and chromosomal localization of the
human earliest lymphocyte activation antigen AIM/CD69, a new member of the C-type
animal lectin superfamily of signal-transmitting receptors. *J Exp Med*. 1993;178(2):537-
547.

26. Akashi K, Kondo M, von Freeden-Jeffry U, Murray R, Weissman IL. Bcl-2 rescues T
lymphopoiesis in interleukin-7 receptor-deficient mice. *Cell*. 1997;89(7):1033-1041.

27. Bongen E, Vallania F, Utz PJ, Khatri P. KLRD1-expressing natural killer cells predict
influenza susceptibility. *Genome Med*. 2018;10(1):45.

28. Garlanda C, Dinarello CA, Mantovani A. The interleukin-1 family: back to the future.
Immunity. 2013;39(6):1003-1018.

29. Passarelli C, Selvatici R, Carrieri A, Di Raimo FR, Falzarano MS, Fortunato F, Rossi R, Straub
V, Bushby K, Reza M, Zharaieva I, D'Amico A, Bertini E, Merlini L, Sabatelli P, Borgiani P,
Novelli G, Messina S, Pane M, Mercuri E, Claustres M, Tuffery-Giraud S, Aartsma-Rus A,
Spitali P, T'Hoen PAC, Lochmuller H, Strandberg K, Al-Khalili C, Kotelnikova E, Lebowitz M,
Schwartz E, Muntoni F, Scapoli C, Ferlini A. Tumor Necrosis Factor Receptor SF10A
(TNFRSF10A) SNPs Correlate With Corticosteroid Response in Duchenne Muscular
Dystrophy. *Front Genet*. 2020;11:605.

30. Liu G, Ren F, Song Y. Upregulation of SPOCK2 inhibits the invasion and migration of
prostate cancer cells by regulating the MT1-MMP/MMP2 pathway. *PeerJ*. 2019;7:e7163.

31. Nabors LK, Wang LD, Wagers AJ, Kansas GS. Overlapping roles for endothelial selectins in
murine hematopoietic stem/progenitor cell homing to bone marrow. *Exp Hematol*.
2013;41(7):588-596.

32.   Tufa DM, Yingst AM, Shank T, Shim S, Trahan GD, Lake J, Woods R, Jones KL, Verneris MR.
532   Transient Expression of GATA3 in Hematopoietic Stem Cells Facilitates Helper Innate
533   Lymphoid Cell Differentiation. *Front Immunol*. 2019;10:510.
534   33.   Bujanover N, Goldstein O, Greenshpan Y, Turgeman H, Klainberger A, Scharff Y, Gazit R.
535   Identification   of   immune-activated   hematopoietic   stem   cells.   *Leukemia*.
536   2018;32(9):2016-2020.
537   34.   Pinho S, Wei Q, Maryanovich M, Zhang D, Balandran JC, Pierce H, Nakahara F, Di Staulo
538   A, Bartholdy BA, Xu J, Borger DK, Verma A, Frenette PS. VCAM1 confers innate immune
539   tolerance on haematopoietic and leukaemic stem cells. *Nat Cell Biol*. 2022;24(3):290-298.
540   35.   Pique-Regi R, Romero R, Garcia-Flores V, Peyvandipour A, Tarca AL, Pusod E, Galaz J,
541   Miller D, Bhatti G, Para R, Kanninen T, Hadaya O, Paredes C, Motomura K, Johnson JR,
542   Jung E, Hsu CD, Berry SM, Gomez-Lopez N. A single-cell atlas of the myometrium in human
543   parturition. *JCI Insight*. 2022;7(5).
544   36.   Goad J, Rudolph J, Zandigohar M, Tae M, Dai Y, Wei JJ, Bulun SE, Chakravarti D, Rajkovic
545   A. Single-cell sequencing reveals novel cellular heterogeneity in uterine leiomyomas. *Hum
546   Reprod*. 2022;37(10):2334-2349.
547   37.   O'Connor SA, Feldman HM, Arora S, Hoellerbauer P, Toledo CM, Corrin P, Carter L, Kufeld
548   M, Bolouri H, Basom R, Delrow J, McFaline-Figueroa JL, Trapnell C, Pollard SM, Patel A,
549   Paddison PJ, Plaisier CL. Neural G0: a quiescent-like state found in neuroepithelial-derived
550   cells and glioma. *Mol Syst Biol*. 2021;17(6):e9522.
551   38.   Riba A, Oravecz A, Durik M, Jimenez S, Alunni V, Cerciat M, Jung M, Keime C, Keyes WM,
552   Molina N. Cell cycle gene regulation dynamics revealed by RNA velocity and deep-
553   learning. *Nat Commun*. 2022;13(1):2865.
554   39.   Zywitza V, Misios A, Bunatyan L, Willnow TE, Rajewsky N. Single-Cell Transcriptomics
555   Characterizes Cell Types in the Subventricular Zone and Uncovers Molecular Defects
556   Impairing Adult Neurogenesis. *Cell Rep*. 2018;25(9):2457-2469 e2458.
557   40.   Scialdone A, Natarajan KN, Saraiva LR, Proserpio V, Teichmann SA, Stegle O, Marioni JC,
558   Buettner F. Computational assignment of cell-cycle stage from single-cell transcriptome
559   data. *Methods*. 2015;85:54-61.
560   41.   La Manno G, Soldatov R, Zeisel A, Braun E, Hochgerner H, Petukhov V, Lidschreiber K,
561   Kastriti ME, Lonnerberg P, Furlan A, Fan J, Borm LE, Liu Z, van Bruggen D, Guo J, He X,
562   Barker R, Sundstrom E, Castelo-Branco G, Cramer P, Adameyko I, Linnarsson S,
563   Kharchenko PV. RNA velocity of single cells. *Nature*. 2018;560(7719):494-498.
564   42.   Bautch VL. Stem cells and the vasculature. *Nat Med*. 2011;17(11):1437-1443.
565   43.   Crisan M, Yap S, Casteilla L, Chen CW, Corselli M, Park TS, Andriolo G, Sun B, Zheng B,
566   Zhang L, Norotte C, Teng PN, Traas J, Schugar R, Deasy BM, Badylak S, Buhring HJ,
567   Giacobino JP, Lazzari L, Huard J, Peault B. A perivascular origin for mesenchymal stem cells
568   in multiple human organs. *Cell Stem Cell*. 2008;3(3):301-313.
569   44.   Lv FJ, Tuan RS, Cheung KM, Leung VY. Concise review: the surface markers and identity of
570   human mesenchymal stem cells. *Stem Cells*. 2014;32(6):1408-1419.
571   45.   Bhartiya D. Adult tissue-resident stem cells-fact or fiction? *Stem Cell Res Ther*.
572   2021;12(1):73.

573   46.   Zhang LZ, Huang LY, Huang AL, Liu JX, Yang F. CRIP1 promotes cell migration, invasion and
574   epithelial-mesenchymal transition of cervical cancer by activating the Wnt/beta-catenin
575   signaling pathway. *Life Sci*. 2018;207:420-427.
576   47.   Fevr T, Robine S, Louvard D, Huelsken J. Wnt/beta-catenin is essential for intestinal
577   homeostasis and maintenance of intestinal stem cells. *Mol Cell Biol*. 2007;27(21):7551-
578   7559.
579   48.   Nusse R. Wnt signaling and stem cell control. *Cell Res*. 2008;18(5):523-527.
580   49.   Mas A, Cervello I, Gil-Sanchis C, Faus A, Ferro J, Pellicer A, Simon C. Identification and
581   characterization of the human leiomyoma side population as putative tumor-initiating
582   cells. *Fertil Steril*. 2012;98(3):741-751 e746.
583   50.   Paul EN, Burns GW, Carpenter TJ, Grey JA, Fazleabas AT, Teixeira JM. Transcriptome
584   Analyses of Myometrium from Fibroid Patients Reveals Phenotypic Differences Compared
585   to Non-Diseased Myometrium. *Int J Mol Sci*. 2021;22(7).
586   51.   Tang L, Bergevoet SM, Gilissen C, de Witte T, Jansen JH, van der Reijden BA, Raymakers
587   RA. Hematopoietic stem cells exhibit a specific ABC transporter gene expression profile
588   clearly distinct from other stem cells. *BMC Pharmacol*. 2010;10:12.
589   52.   Arora R, Fries A, Oelerich K, Marchuk K, Sabeur K, Giudice LC, Laird DJ. Insights from
590   imaging the implanting embryo and the uterine environment in three dimensions.
591   *Development*. 2016;143(24):4749-4754.
592   53.   Flores D, Madhavan M, Wright S, Arora R. Mechanical and signaling mechanisms that
593   guide pre-implantation embryo movement. *Development*. 2020;147(24).
594   54.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate - a Practical and Powerful
595   Approach to Multiple Testing. *J Roy Stat Soc B Met*. 1995;57(1):289-300.
596

 Figure Legends:

Figure 1. Bulk RNA-seq results of SUSD2$^-$ and SUSD2$^+$ cells from myometrial

samples. (**A**) Representative flow cytometry scatter plot of single cell SUSD2$^-$ and

SUSD2$^+$ sort showing over 30% of the live myometrial cells positive for the SUSD2

marker. Boxed areas indicate gating strategy. (**B**) Principal component analysis (PCA)

plot of RNA-seq results from SUSD2$^-$ (in black) and SUSD2$^+$ (in red) cells, each label

represents one sample (n = 5), variance for each PC is indicated in percentage. (**C**)

Volcano plot showing up (n = 3527) and down (n = 3250) DEGs with a false discovery

rate (FDR) p-value < 0.05 in SUSD2$^+$ vs. SUSD2$^-$ cells depicted as red dots. Grey dots

represent genes with an FDR p-value > 0.05. (**D**) Boxplot of mesenchymal stem cells

markers, *SUSD2*, *MCAM*, *PDGFRβ* and *CSPG4* in the SUSD2$^-$ (in grey) and SUSD2$^+$ (in

red) cell population (n = 5). SUSD2$^+$ sorted cells are significantly enriched for MSC

markers including *SUSD2* (log$_2$FC= 4.5, FDR p= 2.9x10^{-5}), *MCAM* (log$_2$FC= 3.1, FDR p=

8.8x10^{-16}), *PDGFRβ* (log$_2$FC= 2.8, FDR p= 1.5x10^{-11}) and *CSPG4* (log$_2$FC= 3.4, FDR p=

1.1x10^{-11}). (**E**) Gene expression is shown as log$_2$(CPM). Heatmap of the top 300 DEGs

from SUSD2$^+$ vs. SUSD2$^-$ cells comparison with unsupervised hierarchical clustering of

genes and samples (n = 5). Color gradient represents gene expression as z-score.

Figure 2. Transcriptomic analysis of the myometrium side population. (**A**) Scatter

plot of the gating strategy to sort the side population (SP$^+$) and the SP$^-$ cells from human

myometrium. (**B**) Verapamil pre-treatment of myometrium cells show a decrease of the

SP$^+$ cells from 1.99% to 0.1% of the total live single cells. (**C**) PCA plot of RNA-seq results

from SP$^-$ (in black) and SP$^+$ (in red) cells, each label represents one sample (n = 3),

variance for each PC is indicated in percentage. (**D**) Volcano plot showing up (n = 478)

and down (n = 350) DEGs with a false discovery rate (FDR) p-value < 0.05 in SP$^+$ vs. SP$^-$

cells depicted as red dots, including *SELE* (log$_2$FC= 4.0, FDR p= 6.0 x10^{-19}), *GATA3*

(log$_2$FC= 3.7, FDR p= 2.8 x10^{-15}), *XCL2* (log$_2$FC= 7.7, FDR p= 1.7 x10^{-15}), *IL7R* (log$_2$FC=

3.7, FDR p= 4.5 x10^{-13}), *CD69* (log$_2$FC= 3.5, FDR p= 7.6 x10^{-13}), *KLRD1* (log$_2$FC= 3.2,

FDR p= 1.3 x10^{-12}), *IL18R1* (log$_2$FC= 3.7, FDR p= 1.5 x10^{-12}), *VCAM1* (log$_2$FC= 3.1, FDR

p= 3.3 x10^{-12}), *TNFRSF10A* (log$_2$FC= 2.7, FDR p= 2.3 x10^{-10}), and *SPOCK2* (log$_2$FC=

3.2, FDR p= 2.4 x10^{-10}). Grey dots represent genes with an FDR p-value > 0.05. (**E**)

Boxplots of myometrium SP associated genes, *ABCB1* (log$_2$FC= 1.9, FDR p= 2.8x10^{-5}),

ABCC1 (log$_2$FC= 0.7, FDR p= 5.9 x10^{-1}), *ABCG2* (log$_2$FC= 1.1, FDR p= 9.5x10^{-2}), *ESR1*

(log$_2$FC= -1.1, FDR p= 2.3 x10^{-1}) and *PGR* (log$_2$FC= -1.5, FDR p= 6.4 x10^{-2}) and (**F**)

mesenchymal stem cells associated genes, *SUSD2* (log$_2$FC= -1, FDR p= 5.6x10^{-1}),

MCAM (log$_2$FC= -1.5, FDR p= 3.9x10^{-2}), *PDGFRβ* (log$_2$FC= -0.9, FDR p= 4.9x10^{-1}),

CSPG4 (log$_2$FC= -1.5, FDR p= 3.9x10^{-2}), *CD44* (log$_2$FC= -0.5, FDR p= 8.3x10^{-1}), *CD34*

(log$_2$FC= -0.2, FDR p= 4.9x10^{-1}), and *ITGA6* (log$_2$FC= 0.9, FDR p= 2.1x10^{-1}) in the SP$^-$

(in grey) and SP$^+$ (in red) cell population (n = 3), genes are expressed in log$_2$(CPM). ns =

non-significant, with FDR > 0.05.

Figure 3. Single cell RNA-seq analysis of isolated cells from human myometrial

samples. (**A**) Uniform manifold approximation and projection (UMAP) visualization of

9775 isolated cells from human myometrial samples (n = 5). Each cluster (n = 7) represent

a cell population with a similar transcriptomic profile. (**B**) Dotplot for cluster identification

using specific markers for cell types found in the myometrium. Mesenchymal stem cell

marker gene expression in the different myometrial cell clusters shown in a dotplot (**C**)

and by UMAP (**D**). Average gene expression and percentage of cells expressing the

specific gene in each cell cluster are shown by the color intensity and the diameter of the

dot, respectively, in B and C. Color gradient in the UMAP represents gene expression as

$\log_2$(CPM).

Figure 4. Identification of putative MyoSPCs from scRNA-seq

(**A**) Cell cycle score for each myometrium cells visualized in the UMAP plot. Cells in

G1/G0, S, and G2/M phases are plotted with the corresponding color. Boxed area is

shown at higher magnification. (**B**) Cell velocity predicting the future state of individual

cells illustrated in a UMAP plotted with the clusters as in panel A showed that in the

vascular myocyte cluster, the same group of cells in G1/G0 phase in panel A exhibit low

velocity. Boxed area is shown at higher magnification. Putative MyoSPCs are encircle in

black dots.

Figure 5. Overlapping analysis of the bulk RNA-seq of SUSD2+ vs SUSD2- and the

assigned stem cell cluster vs the rest of the myometrium cells from the single cell

RNA-seq. Venn diagrams illustrate the overlap DEG between the bulk RNA-seq of

SUSD2+ vs SUSD2- and the assigned stem cell cluster vs the rest of the myometrium

cells from the single cell RNA-seq analysis **(A)**. Scatter plot of $\log_2$(fold change) genes

from bulk RNA-seq (x axis) and single cell RNA-seq (y axis) **(B)**. Non-significant genes in

both analyses were represented in purple dots, DEG in the bulk RNA-seq only were

represented in blue dots, DEG in the scRNA-seq only were represented in green dots,

and the DEG in both analyses were represented in red dots. *CRIP1* is highly upregulated

in the MyoSPC cluster ($\log_2$FC=3.1, adjusted p value = 4 x 10^{-251}) and in the SUSD2+

cells ($\log_2$FC=1.9, FDR = 5 x 10^{-4}). UMAP (**C**) and Ridge plot (**D**) show *CRIP1* expression

in the different myometrium cell clusters by the color intensity or intensity level.

Figure 6. CRIP1$^+$ cells have stem/progenitor cell characteristics. (**A**)

Immunofluorescence imaging of human myometrium using PECAM1 as an endothelial

marker, SUSD2 as a mesenchymal stem cell marker, and CRIP1. Scale bar = 200 μm.

(**B**) Representative scatter plot of the gating strategy for CRIP1$^+$/PECAM$^-$ cell sort. (**C**)

Representative images of colonies formed by the CRIP1$^+$/PECAM$^-$ and depleted

myometrium cell population. (**D**) Plot with of number of colonies formed from

CRIP1$^+$/PECAM$^-$ and depleted myometrium cells after 2-3 weeks in culture (n = 6). (**E**)

Plot of colony forming efficiency represented as %CFUs (#CFU/cells seeded × 100) of

CRIP1$^+$/PECAM$^-$ and depleted myometrium cells (n = 6). (**F**) Total area of colony formed

in pixels from CRIP1$^+$/PECAM$^-$ and depleted myometrium cell population (n = 6). (**G**)

ACTA2 immunofluorescence in CRIP1$^+$/PECAM$^-$ myometrial cells after differentiation.

Scale bar = 100 μm. Representative images of CRIP1$^+$/PECAM$^-$ and depleted

myometrium cell population grown in control growth media and adipogenic (H) or

osteogenic (I) differentiation media. Adipogenic and control cultures were stained with Oil

Red O (red color, black arrows), and osteogenic and control cultures were stained for

alkaline phosphatase activity (purple color). Scale bar for the adipogenic and osteogenic

assays are 500 μm and 5000 μm, respectively. *p<0.05; **p< 0.01 by student t test.

Figure S1. Cell distribution across cell clusters in the single cell RNA-seq. Uniform

manifold approximation and projection (UMAP) visualization of 9775 isolated cells from 5

human myometrial samples (n = 5). Each color dots represents cells from a myometrium

sample. UMAP plot shows that patient's cells are equally distributed across clusters.

Figure S2. *CRIP1* expression in the side population. *CRIP1* expression in log_2CPM of

the RNA-seq results from the side population (SP$^+$) and the negative population (SP-)

showing no significant differences (FDR>0.05) between the two population.'

A

Cell cycle score:
Vascular myocyte cluster

MyoSPC

B

Cell velocity:
Vascular myocyte cluster

MyoSPC