Visual experience shapes functional connectivity between occipital and non-visual networks

Mengyu Tian[1,2] Xiang Xiao[3,4]。 Huiqing Hu[5] Rhodri Cusack[5] Marina Bedny[2]

Abstract

Comparisons of visual cortex function across blind and sighted adults reveals effects of experience on human brain function. Since almost all research has been done with adults, little is known about the developmental origins of plasticity. We compared resting state functional connectivity of visual cortices of blind adults ($n = 30$), blindfolded sighted adults ($n = 50$) to a large cohort of infants (Developing Human Connectome Project, $n = 475$). Visual cortices of sighted adults show stronger coupling with non-visual sensory-motor networks (auditory, somatosensory/motor), than with higher-cognitive prefrontal cortices (PFC). In contrast, visual cortices of blind adults show stronger coupling with higher-cognitive PFC than with nonvisual sensory-motor networks. Are infant visual cortices functionally like those of sighted adults, with blindness leading to functional change? We find that, on the contrary that secondary visual cortices of infants are functionally more like those of blind adults: stronger coupling with PFC than with nonvisual sensory-motor networks, suggesting that visual experience modifies elements of the sighted-adult long-range functional connectivity profile. Infant primary visual cortices are in-between blind and sighted adults i.e., more balanced PFC and sensory-motor connectivity than either adult group. The lateralization of occipital-to-frontal connectivity in infants resembles the sighted adults, consistent with the idea that blindness leads to functional change. These results suggest that both vision and blindness modify functional connectivity through experience-driven (i.e., activity-dependent) plasticity.

Author affiliations:

1 Center for Educational Science, Institute of Advanced Studies in Humanities and Social Sciences, Beijing Normal University, Zhuhai 519087, China

2 Department of Psychological and Brain Sciences, Johns Hopkins University, Baltimore,

Maryland 21218

3 Department of Psychology, Faculty of Art and Science, Beijing Normal University at

Zhuhai, Zhuhai, China

4 Neuroimaging Research Branch, National Institute on Drug Abuse, National Institutes of

Health, Baltimore, MD 21224, USA

5 Trinity College Institute of Neuroscience and School of Psychology, Trinity College Dublin,

Dublin 2, Ireland

Correspondence to: Mengyu Tian

Full address: No.18, Jinfeng Road, Tangjiawan, Zhuhai City, Guangdong Province, 519087,

P.R.China

E-mail: mengyutian@bnu.edu.cn

Keywords: Visual experience, infants, functional connectivity, visual cortex

Introduction

Relative to sighted adults, visual cortices of adults born blind show enhanced responses

during non-visual tasks, such as reading braille and localizing sounds as well as distinctive

patterns of long-range functional connectivity with non-visual networks (Abboud & Cohen,

2019; Bedny et al., 2011; Burton et al., 2012, 2014; Butt et al., 2013; Collignon et al., 2011;

Deen et al., 2015; Kanjlia et al., 2016, 2021; Lane et al., 2015; Liu et al., 2007; Sadato et al.,

1996; Striem-Amit et al., 2015; Watkins et al., 2012). Since almost all research thus far has

been done with adults, a key outstanding question concerns the developmental origins of

these experience-based differences.

One possibility is that at birth, infant visual cortices start out in the 'prepared' sighted adult

state and blindness modifies this functional connectivity. Alternatively, infant visual cortices

may start out functionally similar to those of blind adults and lifetime visual experience

shapes connectivity toward the sighted-adult pattern. To distinguish between these

possibilities, we compare the long-range function connectivity of visual cortices across blind

adults, sighted adults and a large cohort of 2-week-old infants (Developing Human

Connectome Project, dHCP, $n = 475$). Using resting state data provide a common measure of

cortical function across these diverse populations.

To our knowledge no prior studies have compared infants to multiple populations of adults

with different sensory experiences. Previous studies comparing infants to sighted adults have

largely reported similarity across groups (Barttfeld et al., 2018; Doria et al., 2010; Fransson et

al., 2009; Gao et al., 2009; Liu et al., 2008; Zhang et al., 2019). However, these studies

focused on whether large scale functional networks are present in infancy e.g., stronger

connectivity of regions within the visual network, than between visual and auditory regions.

Studies comparing blind and sighted adults find differences across groups in which non-

visual networks are most strongly coupled with the visual system i.e., visual cortices of

sighted adults show stronger coupling with non-visual sensory-motor networks (i.e., auditory,

somatosensory/motor) than higher-cognitive systems; by contrast, in blind adults, visual

cortex coupling is stronger with higher-cognitive prefrontal cortices (PFC) than with non-

visual sensory-motor networks (Abboud & Cohen, 2019; Bedny et al., 2011; Burton et al.,

2014; Deen et al., 2015; Kanjlia et al., 2021; Liu et al., 2007; Qin et al., 2013; Striem-Amit et

al., 2015; Watkins et al., 2012; Yu et al., 2008). In the current study, we compare this

experience-sensitive functional signature across infants, sighted and blind adults.

We measured the connectivity profile of four occipital 'visual' areas that show cross-modal

plasticity in blindness, i.e., are active during non-visual tasks in blind people and show

related changes in resting-state functional connectivity.

We focused on three functionally-distinct secondary visual areas (located in lateral, dorsal

and parts of the ventral occipital cortex) and the primary visual cortex (V1). The three

secondary visual areas have been found to respond to different non-visual tasks in blind

people: language tasks, numerical reasoning tasks and executive control tasks respectively.

Enhanced coupling with PFC is observed across all of three occipital regions in blind adults.

However, each region shows preferential coupling with a distinct subregion of PFC with

analogous functional profiles i.e., language responsive occipital areas are more coupled with

language responsive PFC (Kanjlia et al., 2016, 2021; Lane et al., 2015). These observations

suggest that resting-state and task-based functional profiles are related, although the

functional and developmental nature of this relationship remains an open question.

The precise visual functions of the studied secondary visual regions in sighted people are not known. Anatomically, these regions in sighted people approximately correspond to the locations of motion-sensitive V5/MT+ and the lateral occipital complex (LO), as well as ventral portions of occipito-temporal cortex including V4v and dorsal portions including V3a. The occipital ROI also extends ventrally into the middle portion of the ventral temporal lobe and dorsally into the intraparietal sulcus and superior parietal lobule (Tootell et al., 1997; Van Essen et al., 2001).

We also examined connectivity of anatomically defined primary visual cortex (V1), which likewise shows altered task-based responses and functional connectivity in congenitally blind adults (Amedi et al., 2003; Bedny et al., 2011; Burton et al., 2014; Butt et al., 2013; Lane et al., 2015; Raz et al., 2005; Sadato et al., 1996; Striem-Amit et al., 2015; Yu et al., 2008). Since many previous studies have found that blindness alters the balance of connectivity between visual cortex and higher-order prefrontal as opposed to sensory-motor regions, this was our primary outcome measure (Abboud & Cohen, 2019a; Bedny et al., 2011; Burton et al., 2014; Deen et al., 2015; Heine et al., 2015; Kanjlia et al., 2016, 2021; Lane et al., 2015; Y. Liu et al., 2007; Sen et al., 2022; Striem-Amit et al., 2015). We also examined changes in connectivity lateralization – i.e., the balance of between vs. within hemisphere connectivity, since prior task-based studies have found laterality changes in blindness as well as co-lateralization of occipital and non-occipital networks in this population (Kanjlia et al., 2021; Lane et al., 2017; Tian et al., 2023).

To preview the results, we find that, in infants, the long-range functional connectivity profile of secondary visual areas resembles that of blind adults, whereas V1 falls between blind and sighted adult populations. Relative to sighted adults, both blind adults and infants show stronger coupling between visual cortices and PFC and weaker coupling between visual cortices and non-visual sensory-motor networks. This suggests that vision contributes to modifying the balance of connectivity between occipital and non-visual networks after birth. In contrast, connectivity lateralization patterns appear to reflect blindness-related modification.

Results

Connectivity profile of secondary visual cortices in infants is more similar to that of blind than sighted adults

We first examined the long-range functional connectivity of (three) secondary visual areas

with sensory-motor areas on the one hand, and higher-order PFC networks on the other. In

sighted adults, all three secondary visual areas showed stronger functional connectivity with

non-visual sensory-motor areas (primary somatosensory and motor cortex, S1/M1, and

primary auditory cortex, A1) than with higher-cognitive prefrontal cortices (PFC). By

contrast, in blind adults, all secondary visual regions showed stronger functional connectivity

with PFC than with non-visual sensory-motor areas (S1/M1 and A1) (group (sighted adults,

blind adults) by ROI (PFC, non-visual sensory) interaction effect: $F_{(1, 78)} = 148.819$, $p < 0.001$;

post-hoc Bonferroni-corrected paired t-test, sighted adults: non-visual sensory > PFC: $t_{(49)} =$

9.722, $p < 0.001$; blind adults: non-visual sensory < PFC: $t_{(29)} = 8.852$, $p < 0.001$; Fig. 1).

Like in blind adults, in infants, secondary visual areas showed higher connectivity to PFC

than to non-visual sensory-motor areas (S1/M1 and A1) (non-visual sensory < PFC paired t-

test, $t_{(474)} = 20.144$, $p < 0.001$) (Fig. 1). The connectivity matrix of infants was more

correlated with that of blind than sighted adults, but strongly correlated with both adult

groups (secondary visual, PFC and non-visual sensory areas: infants correlated to blind adults:

$r = 0.721$, $p < 0.001$; to sighted adults: $r = 0.524$, $p < 0.001$; difference between correlations

of infants to blind vs. to sighted adults: z = 3.77, $p < 0.001$; see Supplementary Figure S1 for

the connectivity matrices).

These results suggests that vision is required to set up elements of the sighted adult functional

connectivity pattern, i.e., vision enhances occipital cortex connectivity to non-visual sensory-

motor networks and dampens connectivity to higher-cognitive prefrontal networks.

We checked the robustness of these results in a number of ways. We first compared the

effects across the three secondary visual regions and observed the same pattern across all.

(Supplementary results and Supplementary Figure S2). Next, to check the robustness of the

findings in infants we randomly split the infant dataset into two halves and did split-half

cross-validation. Across all comparisons the results of the two halves were highly similar,

suggesting the effects are robust (see Supplementary Figure S3 and S4). We performed this

validation procedure for all analyses reported below with similar results.

Figure 1 Functional connectivity of secondary visual cortices. (A) Violin plots show the
distributions of functional connectivity (r) of secondary visual cortices (blue) to non-visual
sensory-motor areas (purple) and prefrontal cortices (green), averaged across three occipital,
PFC, and sensory-motor ROIs (A1 and S1/M1) in sighted adults, blind adults, and infants.
Individual dots denote mean connectivity values per participant. Grey trend lines illustrate
within-participant changes across sensory–motor and prefrontal targets. Dark-blue horizontal
markers indicate group averages. Regions of interest (ROI) displayed on the left. Note that
regions extend to ventral surface, not shown. See Supplementary Figure S5 for the full views
of three occipital ROIs. **(B)** Circle plots represent the connectivity of secondary visual
cortices to non-visual networks, min-max normalized to [0,1], i.e., as a proportion. OC:
occipital cortices; MTH: math-responsive region; LG: language-responsive region; EF:
executive function-responsive (response-conflict) region. Asterisks (*) denote significant
Bonferroni-corrected pairwise comparisons (p < .05).

The connectivity pattern of V1 is influenced by early visual experience and blindness

V1 showed the same dissociation between sighted and blind adults as secondary visual areas:
in sighted adults, V1 has stronger functional connectivity with non-visual sensory-motor
areas than with PFC. By contrast, in blind adults, V1 shows stronger connectivity with PFC

than with non-visual sensory areas (group (sighted adults, blind adults) by ROI (PFC, non-

visual sensory) interaction: $F_{(1, 78)} = 125.775$, $p < 0.001$; post-hoc Bonferroni-corrected paired

t-test, sighted adults non-visual sensory > PFC: $t_{(49)} = 9.404$, $p < 0.001$; blind adults non-

visual sensory < PFC: $t_{(29)} = 7.128$, $p < 0.001$; Fig. 2).

The pattern for infants in V1 fell between that of sighted and blind adults. The connectivity

matrix of infants (V1, PFC, and non-visual sensory) was equally correlated with blind and

sighted adults (infants correlated to blind adults: $r = 0.654$, $p < 0.001$; to sighted adults: $r =$

0.594, $p < 0.001$; correlation of infants with blind vs. with sighted adults: $z = 0.832$, $p = 0.406$;

see Supplementary Figure S1 for the connectivity matrices). The difference in connectivity

strength between V1 to PFC and V1 to non-visual sensory regions was weaker in infants than

in sighted or blind adults (group (sighted adults, infants) by ROI (PFC, non-visual sensory)

interaction effect: $F_{(1, 523)} = 92.21$, $p < 0.001$; group (blind adults, infants) by ROI (PFC, non-

visual sensory) interaction effect: $F_{(1, 503)} = 57.444$, $p < 0.001$). V1 of infants showed

marginally stronger connectivity to non-visual sensory regions (S1/M1 and A1) than PFC

(non-visual sensory regions > PFC, paired *t*-test, $t_{(474)} = 1.95$, $p = 0.052$; Fig.2).

The dHCP cohort included both full-term neonates and preterm infants, scanned at their

equivalent gestational age. Visual exposure therefore varied somewhat in duration across

infants (from 0 to 19.71 weeks), with slightly longer exposure in preterm babies. This

variation did not affect connectivity patterns either in V1 or secondary visual cortices (V1: r

$= 0.06$, $p = 0.192$; secondary visual: $r = 0.004$, $p = 0.923$; see Supplementary Figure S6). We

also compared the connectivity patterns of preterm ($n = 90$) and full-term infants ($n = 385$)

and found no difference from each other or from the all-infant dataset (see Supplementary

Figure S7). A few weeks of vision after birth is therefore insufficient to influence

connectivity.

Figure 2 Functional connectivity of primary visual cortices (V1). Violin plots show the distributions of functional connectivity (r) of V1 to non-visual sensory-motor areas (purple) and prefrontal cortices (green), averaged across three PFC ROIs and sensory-motor ROIs (S1/M1 and A1). In sighted adults, blind adults, and infants. Individual dots denote mean connectivity values per participant. Grey trend lines illustrate within-participant changes across sensory–motor and prefrontal targets. Dark-blue horizontal markers indicate group averages. Asterisks (*) denote significant Bonferroni-corrected pairwise comparisons (p < .05). Cross (†) denote marginal difference in Bonferroni-corrected pairwise comparisons (0.05< p <0.1).

Evidence for blindness-related functional change in laterality of occipito-frontal connectivity

Compared to sighted adults, blind adults exhibit a stronger dominance of within-hemisphere connectivity over between-hemisphere connectivity. That is, in people born blind, left visual networks are more strongly connected to left PFC, whereas right visual networks are more strongly connected to right PFC. By contrast, in sighted adults this lateralized pattern is weaker: visual areas in each hemisphere show only a modest preference for ipsilateral

prefrontal cortices, and connectivity with the contralateral PFC remains comparatively strong.

This difference between adult groups is observed for both V1 and secondary visual cortices

(group (blind adults, sighted adults) by lateralization (within hemisphere, between

hemisphere) interaction in secondary visual cortices: $F_{(1, 78)} = 131.51$, $p < 0.001$; V1: $F_{(1, 78)} =$

87.211, $p < 0.001$). Secondary visual cortices showed a significant within > between

difference in both groups, with a larger effect in the blind group (post-hoc tests, Bonferroni-

corrected paired: t-test: sighted adults within hemisphere > between hemisphere: $t_{(49)} =$

7.441, $p = 0.012$, cohen'd = 0.817; blind adults within hemisphere > between hemisphere: t

$_{(29)} = 10.735$, $p < 0.001$, cohen'd = 1.96). In V1, only the blind group showed a significant

within > between hemisphere effect (post-hoc Bonferroni-corrected paired: t-test: sighted

adults within hemisphere < between hemisphere: $t_{(49)} = 3.251$, $p = 0.101$; blind adults within

hemisphere > between hemisphere: $t_{(29)} = 7.019$, $p < 0.001$).

With respect to laterality, infants resembled sighted more than blind adults (Fig. 3). For

secondary visual cortices, there was a significant difference between blind adults and infants

and no difference between sighted adults and infants (group (blind adults, infants) by

lateralization (within hemisphere, between hemisphere) interaction effect: $F_{(1, 503)} = 303.04$, p

< 0.001); (group (sighted adults, infants) by lateralization (within hemisphere, between

hemisphere) interaction effect: $F_{(1, 523)} = 2.244$, $p = 0.135$). A similar group by laterality

interaction were observed for V1 (group (blind adults, infants) by lateralization (within

hemisphere, between hemisphere) interaction: $F_{(1, 503)} = 123.608$, $p < 0.001$; group (sighted

adults, infants) by lateralization (within hemisphere, between hemisphere) interaction effect:

$F_{(1, 523)} = 2.827$, $p = 0.093$). This suggests that the enhancement of within over between

hemisphere long-range connectivity is related to blindness-driven functional change.

Task-based fMRI studies find that cross-modal responses in occipital cortex colateralize with

fronto-parietal networks with related functions (e.g., language, response selection) (Kanjlia et

al., 2021; Lane et al., 2017). For example, language-responsive occipital areas collateralize

with language responsive prefrontal areas across individuals (Lane et al., 2017). Recruitment

of visual cortices by cross-modal tasks (e.g., spoken language) may enhance within-

hemisphere connectivity in people born blind (Kanjlia et al., 2021; Lane et al., 2017; Tian et

al., 2022). Together, this evidence supports the hypothesis that, starting from the less

lateralized infant state, blindness increases lateralization of occipital long-range connectivity.

241
Figure 3 Within hemisphere vs. between hemisphere functional connectivity. Bar graph
shows within hemisphere (blue) and between hemisphere (orange) functional connectivity (r
coefficient of resting state correlations) of secondary visual (left) and V1 (right) to prefrontal
cortices in sighted adults, blind adults, and infants. Blind adults show a larger difference than
any of the other groups. Asterisks (*) denote significant Bonferroni-corrected pairwise
comparisons ($p < .05$).

Specialization of connectivity across different fronto-occipital networks: present in adults, absent at birth

In blind adults, different occipital areas show enhanced connectivity patterns with distinct

subregions of PFC and this specialization is aligned with the functional specialization

observed in task-based data (Bedny et al., 2011; Kanjlia et al., 2016, 2021). For example,

language-responsive subregions of occipital cortex show strongest functional connectivity

with language-responsive sub-regions of PFC, whereas math-responsive occipital areas show

stronger connectivity with math-responsive PFC. This pattern is most pronounced in blind

people but can be seen weakly even in sighted participants (Fig. 4) (Bedny et al., 2011;

Kanjlia et al., 2016, 2021; Lane et al., 2015). Is this fronto-occipital connectivity

specialization present in infancy, potentially enabling the task-based cross-modal

specialization?

We compared connectivity preferences across three prefrontal and three occipital regions

previous shown to activate preferentially in language (sentences > math), math (math >

sentences) and response-conflict (no-go > go with tones) (Kanjlia et al., 2016, 2021; Lane et

al., 2015). For ease of viewing, Fig. 4 shows results from two of the three regions, math and

language. See Supplementary Figure S8 for all three regions. Note that the statistical analyses
included all three areas.

Contrary to the hypothesis that specialization of functional connectivity across different
prefrontal/occipital areas is present from birth, infants showed a less differentiated fronto-
occipital connectivity pattern relative to both blind and sighted adults (Group (sighted adults,
blind adults, infants) by occipital regions (math, language, response-conflict) by PFC regions
(math, language, response-conflict) interaction $F_{(8, 2208)} = 16.323$, $p < 0.001$). Unlike in adults,
in infants, all the occipital regions showed stronger correlations with math- and response-
conflict related prefrontal areas than language-responsive prefrontal areas (Fig. 4 and
Supplementary Figure S8). However, the preferential correlation with math responsive PFC
was strongest in those occipital areas that go on to develop math responses in blind adults
(occipital regions (math, language, response-conflict) by PFC regions (math, language,
response-conflict) interaction in infants $F_{(4, 1896)} = 85.145$, $p < 0.001$, post-hoc Bonferroni-
corrected paired t-test see Supplementary Table S1).

Note that findings of reduced regional specialization in infants need to be interpreted with
caution. First, we do not know whether the same specialization of prefrontal sub-regions seen
in adults is present in infants, although, prior evidence suggests some prefrontal
specialization is already present (Raz & Saxe, 2020). Second, the more fine-grained
comparisons across occipital/frontal regions are more vulnerable to potential anatomical
alignment issues between adult and infant brains. In other words, lack of specialization in
infants could reflect the different location of the areas in this population.

Figure 4 Occipito-frontal functional connectivity. Bar graph shows across functional
connectivity of different sub-regions of prefrontal (PFC) and occipital cortex (OCC) in
sighted adults, blind adults, and infants. Sub-regions (regions of interest) were defined based
on task-based responses in a separate dataset of sighted (frontal) and blind (frontal and
occipital) adults (Kanjlia et al., 2016, 2021; Lane et al., 2015). PFC/OCC-MATH: math-
responsive regions were more active when solving math equations than comprehending
sentences. PFC/OC-LANG: language-responsive regions were more active when
comprehending sentences than solving math equations (Kanjlia et al., 2016, 2021; Lane et al.,
2015). In blind adults these regions show biases in connectivity related to their function i.e.,
language-responsive PFC is more correlated with language responsive OCC. No such pattern
is observed in infants. Asterisks (*) denote significant Bonferroni-corrected pairwise
comparisons ($p < .05$). See Supplementary Figure S9 for connectivity matrix.

Discussion

The present results provide insight into the developmental process of experience-based

functional specialization in human cortex. We find independent effects of visual experience

and blindness on the development of visual networks. Aspects of the sighted adult

connectivity pattern require visual experience. This is particularly striking for secondary

visual cortices, where, connectivity with nonvisual networks in infants resemble blind more

than sighted adults. Both in infants and blind adults, secondary occipital areas showed

stronger functional connectivity with higher-order prefrontal cortices than with other sensory-

motor networks (S1/M1, A1). Consistent with this observation, one previous study with a

small sample of infants found strong connectivity between lateral occipital and prefrontal

areas, although there was no comparison to blind adults in that study (Barttfeld et al., 2018).

In V1, infants fell somewhere in between sighted and blind adults, suggesting an effect both

of vision and of blindness on functional connectivity.

The present results reveal the effects of experience on development of functional connectivity

between infancy and adulthood, but do not speak to the precise time course of these effects.

Infants in the current sample had between 0 and 20 weeks of visual experience. Comparisons

across these infants suggests that several weeks of postnatal visual experience is insufficient

to produce a sighted-adult connectivity profile. The time course of development could be

anywhere between a few months and years and could be tested by examining data from

children of different ages.

We propose that vision, as well as temporally coordinated multi-modal experiences

contribute to establishing the sighted adult connectivity profile in visual cortices. For

example, coordinated visuo-motor activity during development may enhance connectivity

between visual and motor networks. Supporting this conjecture, in infants, motor competence
and early experience predict coupling between occipital and motor networks (Colomer et al.,
2023).

Many questions remain regarding the neurobiological mechanisms underlying experience-
based functional connectivity changes and their relationship to anatomical development.
Resting state functional correlations are an indirect measure of both function and anatomy
and differences in these measures are consistent with many possible underlying biological
mechanisms. Long-range anatomical connections between brain regions are already present
in infants—even prenatally—though they remain immature (Huang et al., 2009; Kostović et
al., 2019, 2021; Takahashi et al., 2012; Vasung, 2017). Functional connectivity changes may
stem from local synaptic modifications within these stable structural pathways, consistent
with findings that functional connectivity can vary independently of structural connection
strength (Fotiadis et al., 2024). Moreover, functional connectivity has been shown to
outperform structural connectivity in predicting individual behavioral differences, suggesting
that experience-based functional changes may reflect finer-scale synaptic or network-level
modulations not captured by macrostructural measures (Ooi et al., 2022). Prior studies also
suggest that, even in adults, coordinated sensory-motor experience can lead to enhancement
of functional connectivity across sensory-motor systems, indicating that large-scale changes
in functional connectivity do not necessarily require corresponding changes in anatomical
connectivity (Guerra-Carrillo et al., 2014; Li et al., 2018). Resting-state functional
connectivity captures synchrony in BOLD signal fluctuations rather than causal interactions
and differences in functional connectivity cannot on their own reveal how underlying
neurophysiological mechanisms are modified. Connectivity changes between two areas could
be mediated by 'third-party' hub regions. For example, posterior parietal cortex serves as a
cortical hub for multisensory integration and visuo-motor coordination and could mediate
occipital-to-sensory-motor communication (Rolls et al., 2023; Sereno & Huang, 2014).
Subcortical structures such as the thalamus could also play a mediating role (Vega-Zuniga et
al., 2025). Future studies will be needed to determine whether these functional changes are
accompanied by alterations in structural connectivity, and to probe causal interactions and
mechanistic underpinnings.

The current findings reveal both effects of vision and effects of blindness on the functional
connectivity patterns of the visual cortex. A further open question is whether visual
experience plays an instructive or permissive role in shaping neural connectivity patterns. An

instructive role implies that sensory experiences or patterns of neural activity directly shape

and organize neural circuitry. In contrast, a permissive role implies that sensory experience or

neural activity merely facilitates the influence of other factors—such as molecular signals—

on the formation and organization of neural circuits (Crair, 1999; Sur et al., 1999). Studies

with animals that manipulate the pattern or informational content of neural activity while

keeping overall activity levels constant could distinguish between these hypotheses (Crair,

1999; Roy et al., 2020; Stellwagen & Shatz, 2002). In humans, such manipulations are not

feasible, leaving us to study only the consequences of the presence or absence of vision.

Under an instructive account, visual and multisensory experience could strengthen coupling

between visual and other non-visual sensory-motor cortices through coordinated activity,

thereby establishing the sighted-adult connectivity pattern. In the absence of visual input, by

contrast, the lack of such coordinated activity may prevent these couplings from being

established. Alternatively, vision may act permissively, indirectly enabling maturational

processes that shift connectivity toward the sighted-adult configuration.

A further key question concerns the behavioral relevance of the connectivity signatures

observed in the current study. The capacity of occipital cortices to support visual and

multimodal behavior in sighted people may depend not only on local visual cortex function

but also on the capacity of the visual system to coordinate its function with non-visual

networks. Does enhanced connectivity between visual and non-visual sensory motor

networks facilitate multimodal integration for sighted people e.g., when catching a ball?

Potentially consistent with this possibility, recent evidence suggests that people who grew up

blind but recover sight in adulthood show multimodal integration deficits (Badde et al., 2020;

Guerreiro et al., 2015; Mowad et al., 2020; Putzar et al., 2007) and distinct occipital

oscillations (Pant et al., 2023).

Conversely, for people who remain blind throughout life, visual-PFC connectivity could

enable recruitment of visual cortices for higher-order non-visual functions, such as language

and executive control (Bedny et al., 2011; Kanjlia et al., 2021). Our results suggest that the

pattern of connectivity observed in blind adults may build on connectivity patterns already

present in infancy: like blind adults, infants show stronger occipital–PFC than occipital–

sensory–motor coupling. Repeated engagement of occipital networks during higher cognitive

tasks in early development could further enhance connectivity and enable specialization of

visual networks for different non-visual higher-order functions.

Some prior studies have measured resting-state and task-based functional profiles in the
same participants. These studies find that within visual cortices of blind people, the task-
based profile of a cortical area is related to its resting state connectivity pattern (Abboud &
Cohen, 2019b; Deen et al., 2015; Kanjlia et al., 2016, 2021). This suggests that these two
measures are related. However, the timecourse of this relationship, the developmental
trajectory and mechanism of plasticity is not known. Primarily this is because there is very
little relevant developmental evidence. For example, in the current study we find that the
resting state profile of secondary visual networks in infants is similar to that of blind adults.
However, we do not know whether the visual cortices of infants show enhanced task-based
cross modal responses, relative to sighted adults and how this compares to responses
observed in blind adults. Future work with infants and children would be able to address this
question.

In the current study, the clearest evidence for functional change driven by blindness was
observed for laterality. Connectivity lateralization in sighted infants resembles that of sighted
adults, in both V1 and secondary visual cortices. Relative to both sighted infants and sighted
adults, blind adults show more lateralized connectivity patterns between occipital and
prefrontal cortices. Previous studies suggest that in people born blind occipital and non-
occipital language responses are co-lateralized (Lane et al., 2017; Tian et al., 2023). We
speculate that habitual activation of visual cortices by higher-cognitive tasks, such as
language, which are themselves highly lateralized, contributes to this biased connectivity
pattern of occipital cortex in blindness. Taken together, these results suggest a developmental
framework in which intrinsic connectivity present in infancy provides a scaffold that is
subsequently shaped and reinforced by experience-dependent recruitment, through either
visual experience or the lifelong absence of vision in blindness. Longitudinal work across
successive developmental stages will be crucial to test how the alternative trajectories shaped
by visual experience versus blindness unfold over development.

Materials and methods

Participants

Fifty sighted adults and thirty congenitally blind adults contributed the resting state
data (sighted: n = 50; 30 females; mean age = 35.33 years, standard deviation (SD) = 14.65;

mean years of education = 17.08, SD = 3.1; blind: n = 30; 19 females; mean age = 44.23

years, SD = 16.41; mean years of education = 17.08, SD = 2.11; blind vs. sighted age, $t_{(78)}$ =

2.512, $p < 0.05$; blind vs. sighted years of education, $t_{(78)} = 0.05$, $p = 0.996$). Since blind

participants were on average older, we also performed analyses in an age-matched subgroups

of sighted controls ($n = 29$) and found similar results to the full sample (see Supplementary

Figure S10). Blind and sighted participants had no known cognitive or neurological

disabilities (screened through self-report). All adult anatomical images were read by a board-

certified radiologist, and no gross neurological abnormalities were found. All the blind

participants had at most minimal light perception from birth. Blindness was caused by

pathology anterior to the optic chiasm (i.e., not due to brain damage). All participants gave

written informed consent under a protocol approved by the Institutional Review Board of

Johns Hopkins University.

Neonate data were from the third release of the Developing Human Connectome Project

(dHCP) ($n = 783$) (https ://www.developing connectome.org). Ethical approval was obtained

from the UK Health Research Authority (Research Ethics Committee reference number:

14/LO/1169). After quality control procedures (described below), 475 subjects were included

in data analysis, with one scan per subject. The average age from birth at scan = 2.79 weeks

(SD = 3.77, median = 1.57, range = 0 – 19.71); average gestational age at scan = 41.23 weeks

(SD = 1.77, median = 41.29, range = 37 – 45.14); average gestational age at birth = 38.43

weeks (SD = 3.73, median = 39.71, range = 23 – 42.71). We only included infants who were

full-term or scanned at term-equivalent age if preterm, while not being flagged by the dHCP

project team as not passing quality control for functional MRI (fMRI) images ($n = 634$).

Infants with more than 160 motion outliers were exclude ($n = 116$ dropped). Motion-outlier

volumes were defined as DVARS (the root mean square intensity difference between

successive volumes) higher than 1.5 interquartile range above the 75th centile, after motion

and distortion correction. Infants with signal drop-out in regions of interest (ROI) were also

excluded ($n = 43$ dropped). To identify signal dropout, we first averaged blood oxygen level-

dependent (BOLD) signal intensity for all time point, for each subject, in each of 100 parcel

defined by Schaefer's atlas (Schaefer et al., 2018). For each ROI ($n = 18$ ROIs) in the current

study, signal dropout was then identified as BOLD intensity lower than -3 standard deviations,

where the mean and standard deviations were identified across all 100 cortical parcels.

Participants were excluded if any of the ROIs showed a signal dropout. The same signal

dropout assessment was also applied to the blind and sighted adults to ensure consistent

quality control across groups. One participant in the sighted adult group and two participants

in the blind adult group exhibited signal dropout in one ROI each. Excluding these

participants did not alter the group-level results (see Figure S13). The infants' structural

images were reviewed by a pediatric neuroradiologist from the dHCP team, who assigned

scores on a scale from 1 to 5. A score of 1 indicated a normal appearance for the subject's age,

while scores of 4 or 5 suggested potential or likely clinical significance, or both clinical and

imaging relevance. We repeated our analysis after excluding infants with a radiology score of

4 or 5, and the results remained consistent (see Figure S11).

Image acquisition

Blind and sighted adult MRI anatomical and functional images were collected on a 3T

Phillips scanner at the F. M. Kirby Research Center. T1-weighted anatomical images were

collected using a magnetization-prepared rapid gradient-echo (MP-RAGE) in 150 axial slices

with 1 mm isotropic voxels. Resting state fMRI data were collected in 36 sequential

ascending axial slices for 8 minutes. TR = 2 s, TE = 0.03 s, flip angle = 70°, voxel size = 2.4

$\times$ 2.4 $\times$ 2.5 mm, inter-slice gap = 0.5 mm, field of view (FOV) = 192 $\times$ 172.8 $\times$ 107.5.

Participants completed 1 to 4 scans of 240 volume each (average scan time = 710.4 second

per person). During the resting state scan, participants were instructed to relax but remain

awake. Sighted participants wore light-excluding blindfolds to equalize the light conditions

across the groups during the scans.

Infants (dHCP) Anatomical and functional images were collected on a 3T Phillips scanner at

the Evelina Newborn Imaging Centre, St Thomas' Hospital, London, UK. A dedicated

neonatal imaging 219 system including a neonatal 32-channel phased-array head coil was

used. T2w multi-slice fast spin-echo images were acquired with in-plane resolution 0.8x0.8

mm^2 and 1.6 mm slices overlapped by 0.8 mm (TR = 12000 ms, TE = 156 ms, SENSE factor

2.11 axial and 2.6 sagittal). In infants, T2w images were used as the anatomical image

because the brain anatomy is more clearly in T2w than in T1w images. Fifteen minutes of

resting state fMRI data were collected using a used multiband 9x accelerated echo-planar

imaging (TR = 392 ms, TE = 38 ms, 2300 volumes, with an acquired resolution of 2.15 mm

isotropic). Single-band reference scans were acquired with bandwidth-matched readout,

along with additional spin-echo acquisitions with both AP/PA fold-over encoding directions.

Data analysis

Resting state data were preprocessed using FSL version 5.0.9 (Smith et al., 2004), DPABI version 6.1 (Yan et al., 2016), FreeSurfer (Dale et al., 1999), and in-house code (https://github.com/NPDL/Resting-state_dHCP). The functional data for all groups were linearly detrended and low-pass filtered (0.08 Hz).

For adults, functional images were registered to the T1-weighted structural images, motion corrected using MCFLIRT (Jenkinson et al., 2002), and temporally high-pass filtering (150 s). No subject had excessive head movement ($> 2mm$) or rotation ($> 2°$) at any timepoint. Resting state data are known to include artifacts related to physiological fluctuations such as cardiac pulsations and respiratory-induced modulation of the main magnetic field. A component-based method, CompCor (Behzadi et al., 2007), was therefore used to control for these artifacts. Particularly, following the procedure described in Whitfield-Gabrieli *et al.*, nuisance signals were extracted from 2-voxel eroded masks of spinal fluid (CSF) and white matter (WM), and the first 5 principal components analysis (PCA) components derived from these signals was regressed out from the processed BOLD time series (Whitfield-Gabrieli & Nieto-Castanon, 2012). In addition, a scrubbing procedure was applied to further reduce the effect of motion on functional connectivity measures (Power et al., 2012, 2014). Frames with root mean square intensity difference exceeding 1.5 interquartile range above the 75th centile, after motion and distortion correction, were censored as outliers.

The infants resting state functional data were pre-processed by the dHCP group using the project's in-house pipeline (Fitzgibbon et al., 2020), This pipeline uses a spatial independent component analysis (ICA) denoising step to minimize artifact due to multi-band artefact, residual head-movement, arteries, sagittal sinus, CSF pulsation. For infants, ICA denoising is preferable to using CSF/WM regressors. Because it is challenging to accurately define anatomical boundaries of CSF/WM due to the low imaging resolution comparing with the brain size and the severe partial-volume effect in the neonate (Fitzgibbon et al., 2020). Like in the adults, frames with root mean square intensity difference exceeding 1.5 interquartile range above the 75th centile, after motion and distortion correction, were considered as motion outliers. Out from the 2300 frames, a subset of continuous 1600 with minimum number of motion outliers was kept for each subject. Motion outliers were censored from the subset of continuous 1600, and a subject was excluded from further analyses when the number of outlier exceeded 160 (10% of the continues subset) (Hu et al., 2022). While infant

connectivity estimates may be less robust at the individual level compared to adults due to

shorter scan durations and higher motion, our cohort's large sample size (n=475) and

rigorous motion censoring mitigate these limitations for group-level analyses. Substantial

differences between the groups exist in this study, including the number of subjects, brain

sizes, imaging parameters, and data preprocessing, all of which are likely to have an impact

on the overall signal quality. To assess the reliability of the ROI-wise FC used in all analyses,

we computed split-half noise ceilings for each participant. The rs-fMRI time series was

divided into two equal halves, ROI-wise FC profiles were calculated separately for each half,

and the noise ceiling for each ROI was defined as the Pearson correlation between the two

profiles following Lage-Castellanos et al. (2019). We then averaged the ROI-wise noise

ceilings for each group. The resulting values (infants: 0.90 ± 0.056; blind adults: 0.88 ± 0.041;

sighted adults: 0.90 ± 0.055) did not differ significantly (one-way ANOVA: $F_{(2,552)} = 2.348$, p

$= 0.097$). We also examined noise ceilings for each ROI separately. All ROIs showed high

absolute reliability (noise ceiling > 0.80) across groups. Although many ROIs showed

statistically significant group differences in noise ceiling (one-way ANOVA, $p < 0.05$), the

effect sizes were small to moderate (partial $\eta^2 < 0.14$). These findings indicate that reliability

may vary modestly across groups at the ROI level. We cannot determine whether such

variability contributes to the group differences reported in this study, but the consistently high

absolute reliability suggests that the FC estimates are generally robust. The full ROI-wise

statistical results are provided in the Supplementary Table S2. For both groups of adult and

infants, we performed a temporal low-pass filter (0.08 Hz low-pass cutoff) and a linear

detrending. ROI-to-ROI connectivity was calculated using Pearson's correlation between

ROI-averaged BOLD timeseries (ROI definition see below). The All t-tests and F-tests are

two-sided. The comparison of correlation coefficients was done using cocor software package

and Pearson and Filon's z (Diedenhofen & Musch, 2015; Pearson & Filon, 1898).

ROI definition

Frontal and secondary visual ROIs were defined functionally based on data from a separate

task-based fMRI experiments with blind and sighted adults (Kanjlia et al., 2016, 2021; Lane

et al., 2015). Three separate experiments were conducted with the same group of blind and

sighted subjects (sighted n=18; blind n=23). The language ROIs in the occipital and frontal

cortices were identified by sentence $>$ nonwords contrast in an auditory language

comprehension task (Lane et al., 2015). The math ROIs were identified by math $>$ sentence

contrast in an auditory task where participants judged equivalence of pairs of math equations

and pairs of sentences (Kanjlia et al., 2016). The executive function ROIs were identified by

no-go > frequent go contrast in an auditory go/no-go task with non-verbal sounds (Kanjlia et

al., 2021). All ROI files are available at openICPSR (https://doi.org/10.3886/E198832V1).

Occipital secondary 'visual' ROIs were defined based on group comparisons blind > sighted

in a whole-cortex analysis (Figure 1, Figure S5.) The occipital language ROI was defined as

the cluster that responded more to auditory sentence than auditory nonwords conditions in

blind, relative to sighted, in a whole-cortex analysis, likewise the occipital math ROI was

defined as math>sentences, blind>sighted interaction and the occipital executive ROI as no-

go > frequent go, blind>sighted. All three occipital ROIs were defined in the right

hemisphere. Left-hemisphere occipital ROIs were created by flipping the right-hemisphere

ROIs to the left hemisphere. Each functional ROI spans multiple anatomical regions and

together the secondary visual ROIs tile large portions of lateral occipital, occipito-temporal,

dorsal occipital and occipito-parietal cortices. In sighted people, the secondary visual

occipital ROIs include the anatomical locations of functional regions such as motion area

V5/MT+, the lateral occipital complex (LO), as well as ventral portions encompassing

category-selective ventral occipito-temporal cortices, including V4v, and dorsal portions

including V3a. The occipital ROI also extended ventrally into the middle portion of the

temporal lobe and dorsally into the intraparietal sulcus and superior parietal lobule. The

frontal PFC ROIs were defined functionally, based on a whole-cortex analysis which

combined all blind and sighted adult data. The frontal language ROI was defined as

responded more auditory sentence than auditory nonwords conditions across all blind and

sighted subjects, constrained to the prefrontal cortex. Likewise, math responsive PFC was

defined as math>sentences and executive no-go > frequent go. For frontal ROIs, the language

ROI was defined in the left, and the math and executive function ROI were defined in the

right hemisphere, then flip to the other hemisphere.

The V1 ROI was defined from a previously published anatomical surface-based atlas (PALS-

B12) (Van Essen, 2005). The primary somatosensory and motor cortex (S1/M1) ROI was

selected as the area that responds more to the go than no-go trials in the auditory go/no-go

task across both blind and sighted groups, constrained to the hand area in S1/M1 search-space

from neurosynth.org (term "hand movements") (Kanjlia et al., 2021). The primary auditory

cortex (A1) ROI was defined as the transverse temporal portion of a gyral-based atlas

(Desikan et al., 2006; Morosan et al., 2001).

All the ROIs were defined in standard space and then transformed into each subject's native

space. For adults this was done by employing the deformation field estimated by FreeSurfer.

For infants, the ROIs were transformed into each subject's native space using a two-step

approach. First, the ROIs were converted from the adult's MNI space into the 40-week dHCP

template (Bozek et al., 2018). ANTS, previously shown to be effective in pediatric studies

(Avants et al., 2014; Cabral et al., 2022; Jain et al., 2012; Lawson et al., 2013), was utilized

to estimate the deformation field between these two spaces. In this step, the infant's scalp and

cerebellum were masked, as these structures in the infant brain greatly differ from those in

the adult and can introduce bias into the registration process, as outlined in a study by Cabral

et al (Cabral et al., 2022). Secondly, the ROIs were further transformed from the 40-week

template space into each individual's native spaces, employing the deformation field

provided by the dHCP group. Nearest neighbor interpolation was applied in both steps

(Examples of the resulting ROI alignment on individual brains are shown in Supplementary

Figure S12). For both adults and infants, any overlapping voxels between ROIs were

removed and not counted toward any ROIs.

Data availability

593   Neonate data were from the third release of the Developing Human Connectome Project

594   (https://www.developing connectome.org). The de-identified blind and sighted adults' data

595   were posted on openICPSR (https://doi.org/10.3886/E198832V1).

Acknowledgements

We would like to thank all the blind and sighted participants, the blind community and the

National Federation of the Blind. Without their support, this study would not be possible. We

would also like to thank the F. M. Kirby Research Center for Functional Brain Imaging at the

Kennedy Krieger Institute for their assistance in data collection. Xiang Xiao was supported

by the Intramural Research Program of the National Institute on Drug Abuse, the National

Institute of Health, United States.

Funding

This work was supported by grants from the National Eye Institute at the National Institutes of Health (R01EY027352-01 and R01EY033340). RC was supported by the ERC Advanced Grant "Foundations of Cognition" (FOUNDCOG) 787981. MT was supported by grants from the National Natural Science Foundation of China [32400891]. XX was supported by grants from the National Natural Science Foundation of China [32400844] and the Fundamental Research Funds for the Central Universities.

Competing interests

The authors report no competing interests.

References

Abboud, S., & Cohen, L. (2019a). Distinctive Interaction Between Cognitive Networks and the Visual Cortex in Early Blind Individuals. *Cerebral Cortex*, *33*(0), 4725–4742. https://doi.org/10.1093/cercor/bhz006

Abboud, S., & Cohen, L. (2019b). Distinctive Interaction between Cognitive Networks and the Visual Cortex in Early Blind Individuals. *Cerebral Cortex*, *29*(11), 4725–4742. https://doi.org/10.1093/cercor/bhz006

Amedi, A., Raz, N., Pianka, P., Malach, R., & Zohary, E. (2003). Early "visual" cortex activation correlates with superior verbal memory performance in the blind. *Nature Neuroscience*, *6*(7), 758–766. https://doi.org/10.1038/nn1072

Avants, B. B., Tustison, N. J., Stauffer, M., Song, G., Wu, B., & Gee, J. C. (2014). The Insight ToolKit image registration framework. *Frontiers in Neuroinformatics*, *8*. https://www.frontiersin.org/articles/10.3389/fninf.2014.00044

Badde, S., Ley, P., Rajendran, S. S., Shareef, I., Kekunnaya, R., & Röder, B. (2020). Sensory experience during early sensitive periods shapes cross-modal temporal biases. *eLife*, *9*, e61238. https://doi.org/10.7554/eLife.61238

Barttfeld, P., Abboud, S., Lagercrantz, H., Adén, U., Padilla, N., Edwards, A. D., Cohen, L., Sigman, M., Dehaene, S., & Dehaene-Lambertz, G. (2018). A lateral-to-mesial

organization of human ventral visual cortex at birth. *Brain Structure and Function*,

223(7), 3107–3119. https://doi.org/10.1007/s00429-018-1676-3

Bedny, M., Pascual-Leone, A., Dodell-Feder, D., Fedorenko, E., & Saxe, R. (2011).

Language processing in the occipital cortex of congenitally blind adults. *Proceedings*

of the National Academy of Sciences of the United States of America, *108*(11), 4429–

4434. https://doi.org/10.1073/pnas.1014818108

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction

method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, *37*(1), 90–

101. https://doi.org/10.1016/j.neuroimage.2007.04.042

Bozek, J., Makropoulos, A., Schuh, A., Fitzgibbon, S., Wright, R., Glasser, M. F., Coalson,

T. S., O'Muircheartaigh, J., Hutter, J., Price, A. N., Cordero-Grande, L., Teixeira, R.

P. A. G., Hughes, E., Tusor, N., Baruteau, K. P., Rutherford, M. A., Edwards, A. D.,

Hajnal, J. V., Smith, S. M., … Robinson, E. C. (2018). Construction of a neonatal

cortical surface atlas using Multimodal Surface Matching in the Developing Human

Connectome Project. *NeuroImage*, *179*, 11–29.

https://doi.org/10.1016/j.neuroimage.2018.06.018

Burton, H., Sinclair, R. J., & Agato, A. (2012). Recognition memory for Braille or spoken

words: An fMRI study in early blind. *Brain Research*, *1438*, 22–34.

https://doi.org/10.1016/j.brainres.2011.12.032

Burton, H., Snyder, A. Z., & Raichle, M. E. (2014). Resting state functional connectivity in

early blind humans. *Frontiers in Systems Neuroscience*, *8*(1 APR), 1–13.

https://doi.org/10.3389/fnsys.2014.00051

Butt, O. H., Benson, N. C., Datta, R., & Aguirre, G. K. (2013). The fine-scale functional

correlation of striate cortex in sighted and blind people. *Journal of Neuroscience*,

33(41), 16209–16219. https://doi.org/10.1523/JNEUROSCI.0363-13.2013

Cabral, L., Zubiaurre-Elorza, L., Wild, C. J., Linke, A., & Cusack, R. (2022). Anatomical

correlates of category-selective visual regions have distinctive signatures of

connectivity in neonates. *Developmental Cognitive Neuroscience*, *58*, 101179.

https://doi.org/10.1016/j.dcn.2022.101179

Collignon, O., Vandewalle, G., Voss, P., Albouy, G., Charbonneau, G., Lassonde, M., &

Lepore, F. (2011). Functional specialization for auditory-spatial processing in the

occipital cortex of congenitally blind humans. *Proceedings of the National Academy*

of Sciences, *108*(11), 4435–4440. https://doi.org/10.1073/pnas.1013928108

Colomer, M., Chung, H., Meyer, M., Debnath, R., Morales, S., Fox, N. A., & Woodward, A.

(2023). Action experience in infancy predicts visual☐motor functional connectivity

during action anticipation. *Developmental Science*, *26*(3), e13339.

https://doi.org/10.1111/desc.13339

Crair, M. C. (1999). Neuronal activity during development: Permissive or instructive?

Current Opinion in Neurobiology, *9*(1), 88–93. https://doi.org/10.1016/S0959-

4388(99)80011-7

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis: I.

Segmentation and surface reconstruction. *NeuroImage*, *9*(2), 179–194.

https://doi.org/10.1006/nimg.1998.0395

Deen, B., Saxe, R., & Bedny, M. (2015). Occipital Cortex of Blind Individuals Is

Functionally Coupled with Executive Control Areas of Frontal Cortex. *Journal of*

Cognitive Neuroscience, *27*(8), 1633–1647. https://doi.org/10.1162/jocn_a_00807

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner,

R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J.

(2006). An automated labeling system for subdividing the human cerebral cortex on

MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980.

https://doi.org/10.1016/j.neuroimage.2006.01.021

Diedenhofen, B., & Musch, J. (2015). cocor: A Comprehensive Solution for the Statistical

Comparison of Correlations. *PLOS ONE*, *10*(4), e0121945.

https://doi.org/10.1371/journal.pone.0121945

Doria, V., Beckmann, C. F., Arichi, T., Merchant, N., Groppo, M., Turkheimer, F. E.,

Counsell, S. J., Murgasova, M., Aljabar, P., Nunes, R. G., Larkman, D. J., Rees, G., &

Edwards, A. D. (2010). Emergence of resting state networks in the preterm human

brain. *Proceedings of the National Academy of Sciences of the United States of*

America, *107*(46), 20015–20020. https://doi.org/10.1073/pnas.1007921107

Fitzgibbon, S. P., Harrison, S. J., Jenkinson, M., Baxter, L., Robinson, E. C., Bastiani, M.,

Bozek, J., Karolis, V., Cordero Grande, L., Price, A. N., Hughes, E., Makropoulos,

A., Passerat-Palmbach, J., Schuh, A., Gao, J., Farahibozorg, S. R., O'Muircheartaigh,

J., Ciarrusta, J., O'Keeffe, C., … Andersson, J. (2020). The developing Human

Connectome Project (dHCP) automated resting-state functional processing framework

for newborn infants. *NeuroImage*, *223*(August), 117303.

https://doi.org/10.1016/j.neuroimage.2020.117303

Fotiadis, P., Parkes, L., Davis, K. A., Satterthwaite, T. D., Shinohara, R. T., & Bassett, D. S.

(2024). Structure–function coupling in macroscale human brain networks. *Nature*

Reviews Neuroscience, *25*(10), 688–704. https://doi.org/10.1038/s41583-024-00846-6

Fransson, P., Skiöld, B., Engström, M., Hallberg, B., Mosskin, M., Åden, U., Lagercrantz,

H., & Blennow, M. (2009). Spontaneous brain activity in the newborn brain during

natural sleep-an fMRI study in infants born at full term. *Pediatric Research*, *66*(3),

301–305. https://doi.org/10.1203/PDR.0b013e3181b1bd84

Gao, W., Zhu, H., Giovanello, K. S., Smith, J. K., Shen, D., Gilmore, J. H., & Lin, W. (2009).

Evidence on the emergence of the brain's default network from 2-week-old to 2-year-

old healthy pediatric subjects. *Proceedings of the National Academy of Sciences*,

106(16), 6790–6795. https://doi.org/10.1073/pnas.0811221106

Guerra-Carrillo, B., MacKey, A. P., & Bunge, S. A. (2014). Resting-state fMRI: A window

into human brain plasticity. *Neuroscientist*, *20*(5), 522–533.

https://doi.org/10.1177/1073858414524442

Guerreiro, M. J. S., Putzar, L., & Röder, B. (2015). The effect of early visual deprivation on

the neural bases of multisensory processing. *Brain*, *138*(6), 1499–1504.

https://doi.org/10.1093/brain/awv076

Heine, L., Bahri, M. A., Cavaliere, C., Soddu, A., Laureys, S., Ptito, M., & Kupers, R.

(2015). Prevalence of increases in functional connectivity in visual, somatosensory

and language areas in congenital blindness. *Frontiers in Neuroanatomy*, *9*.

https://doi.org/10.3389/fnana.2015.00086

Hu, H., Cusack, R., & Naci, L. (2022). Typical and disrupted brain circuitry for conscious

awareness in full-term and preterm infants. *Brain Communications*, *4*(2), fcac071.

https://doi.org/10.1093/braincomms/fcac071

Huang, H., Xue, R., Zhang, J., Ren, T., Richards, L. J., Yarowsky, P., Miller, M. I., & Mori,

S. (2009). Anatomical Characterization of Human Fetal Brain Development with

Diffusion Tensor Magnetic Resonance Imaging. *The Journal of Neuroscience*, *29*(13),
4263–4273. https://doi.org/10.1523/JNEUROSCI.2769-08.2009

Jain, V., Duda, J., Avants, B., Giannetta, M., Xie, S. X., Roberts, T., Detre, J. A., Hurt, H.,
Wehrli, F. W., & Wang, D. J. J. (2012). Longitudinal Reproducibility and Accuracy
of Pseudo-Continuous Arterial Spin–labeled Perfusion MR Imaging in Typically
Developing Children. *Radiology*, *263*(2), 527–536.
https://doi.org/10.1148/radiol.12111509

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the
robust and accurate linear registration and motion correction of brain images.
NeuroImage, *17*(2), 825–841. https://doi.org/10.1016/s1053-8119(02)91132-8

Kanjlia, S., Lane, C., Feigenson, L., & Bedny, M. (2016). Absence of visual experience
modifies the neural basis of numerical thinking. *Proceedings of the National Academy
of Sciences of the United States of America*, *113*(40), 11172–11177.
https://doi.org/10.1073/pnas.1524982113

Kanjlia, S., Loiotile, R. E., Harhen, N., & Bedny, M. (2021). 'Visual' cortices of congenitally
blind adults are sensitive to response selection demands in a go/no-go task.
NeuroImage, *236*(April), 118023. https://doi.org/10.1016/j.neuroimage.2021.118023

Kostović, I., Katušić, A., & Kostović Srzentić, M. (2021). Linking histology and neurological
development of the fetal and infant brain. In *Factors Affecting Neurodevelopment* (pp.
213–225). Elsevier. https://doi.org/10.1016/B978-0-12-817986-4.00019-5

Kostović, I., Sedmak, G., & Judaš, M. (2019). Neural histology and neurogenesis of the
human fetal and infant brain. *NeuroImage*, *188*, 743–773.
https://doi.org/10.1016/j.neuroimage.2018.12.043

Lage-Castellanos, A., Valente, G., Formisano, E., & Martino, F. D. (2019). Methods for
computing the maximum performance of computational models of fMRI responses.
PLOS Computational Biology, *15*(3), e1006397.
https://doi.org/10.1371/journal.pcbi.1006397

Lane, C., Kanjlia, S., Omaki, A., & Bedny, M. (2015). "Visual" cortex of congenitally blind
adults responds to syntactic movement. *Journal of Neuroscience*, *35*(37), 12859–
12868. https://doi.org/10.1523/JNEUROSCI.1256-15.2015

Lane, C., Kanjlia, S., Richardson, H., Fulton, A., Omaki, A., & Bedny, M. (2017). Reduced

Left Lateralization of Language in Congenitally Blind Individuals. *Journal of*

Cognitive Neuroscience, *29*(1), 65–78. https://doi.org/10.1162/jocn_a_01045

Lawson, G. M., Duda, J. T., Avants, B. B., Wu, J., & Farah, M. J. (2013). Associations

between children's socioeconomic status and prefrontal cortical thickness.

Developmental Science, *16*(5), 641–652. https://doi.org/10.1111/desc.12096

Li, Q., Wang, X., Wang, S., Xie, Y., Li, X., Xie, Y., & Li, S. (2018). Musical training

induces functional and structural auditory☐motor network plasticity in young adults.

Human Brain Mapping, *39*(5), 2098–2110. https://doi.org/10.1002/hbm.23989

Liu, W. C., Flax, J. F., Guise, K. G., Sukul, V., & Benasich, A. A. (2008). Functional

connectivity of the sensorimotor area in naturally sleeping infants. *Brain Research*,

1223, 42–49. https://doi.org/10.1016/j.brainres.2008.05.054

Liu, Y., Yu, C., Liang, M., Li, J., Tian, L., Zhou, Y., Qin, W., Li, K., & Jiang, T. (2007).

Whole brain functional connectivity in the early blind. *Brain*, *130*(8), 2085–2096.

https://doi.org/Neuronal%2520recycling

Morosan, P., Rademacher, J., Schleicher, A., Amunts, K., Schormann, T., Zilles, K., & Zilles,

K. (2001). Human primary auditory cortex: Cytoarchitectonic subdivisions and

mapping into a spatial reference system. *NeuroImage*, *13*(4), 684–701.

https://doi.org/10.1006/nimg.2000.0715

Mowad, T. G., Willett, A. E., Mahmoudian, M., Lipin, M., Heinecke, A., Maguire, A. M.,

Bennett, J., & Ashtari, M. (2020). Compensatory Cross-Modal Plasticity Persists

After Sight Restoration. *Frontiers in Neuroscience*, *14*, 291.

https://doi.org/10.3389/fnins.2020.00291

Ooi, L. Q. R., Chen, J., Zhang, S., Kong, R., Tam, A., Li, J., Dhamala, E., Zhou, J. H.,

Holmes, A. J., & Yeo, B. T. T. (2022). Comparison of individualized behavioral

predictions across anatomical, diffusion and functional connectivity MRI.

NeuroImage, *263*, 119636. https://doi.org/10.1016/j.neuroimage.2022.119636

Pant, R., Ossandón, J., Stange, L., Shareef, I., Kekunnaya, R., & Röder, B. (2023). Stimulus-

evoked and resting-state alpha oscillations show a linked dependence on patterned

visual experience for development. *NeuroImage: Clinical*, *38*, 103375.

https://doi.org/10.1016/j.nicl.2023.103375

Pearson, K., & Filon, L. N. G. (1898). Mathematical contributions to the theory of evolution.
785   IV. On the probable errors of frequency constants and on the influence of random
786   selection on variation and correlation. *Proceedings of the Royal Society of London*,
787   *62*(379–387), 173–176. https://doi.org/10.1098/rspl.1897.0091

788 Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012).
789   Spurious but systematic correlations in functional connectivity MRI networks arise
790   from subject motion. *NeuroImage*, *59*(3), 2142–2154.
791   https://doi.org/10.1016/j.neuroimage.2011.10.018

792 Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E.
793   (2014). Methods to detect, characterize, and remove motion artifact in resting state
794   fMRI. *NeuroImage*, *84*, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

795 Putzar, L., Goerendt, I., Lange, K., Rösler, F., & Röder, B. (2007). Early visual deprivation
796   impairs multisensory interactions in humans. *Nature Neuroscience*, *10*(10), 1243–
797   1245. https://doi.org/10.1038/nn1978

798 Qin, W., Liu, Y., Jiang, T., & Yu, C. (2013). The Development of Visual Areas Depends
799   Differently on Visual Experience. *PLoS ONE*, *8*(1), 1–10.
800   https://doi.org/10.1371/journal.pone.0053784

801 Raz, G., & Saxe, R. (2020). Learning in Infancy Is Active, Endogenously Motivated, and
802   Depends on the Prefrontal Cortices. *Annual Review of Developmental Psychology*,
803   *2*(1), 247–268. https://doi.org/10.1146/annurev-devpsych-121318-084841

804 Raz, N., Amedi, A., & Zohary, E. (2005). V1 activation in congenitally blind humans is
805   associated with episodic retrieval. *Cerebral Cortex*, *15*(9), 1459–1468.
806   https://doi.org/10.1093/cercor/bhi026

807 Rolls, E. T., Deco, G., Huang, C.-C., & Feng, J. (2023). The human posterior parietal cortex:
808   Effective connectome, and its relation to function. *Cerebral Cortex*, *33*(6), 3142–
809   3170. https://doi.org/10.1093/cercor/bhac266

810 Roy, A., Wang, S., Meschede-Krasa, B., Breffle, J., & Van Hooser, S. D. (2020). An early
811   phase of instructive plasticity before the typical onset of sensory experience. *Nature*
812   *Communications*, *11*(1), 11. https://doi.org/10.1038/s41467-019-13872-1

Sadato, N., Pascual-Leone, A., Grafman, J., Ibañez, V., Deiber, M., Dold, G., & Hallett, M.

(1996). Activation of the primary visual cortex by Braille reading in blind subjects.

Nature, *380*(6574), 526–528. https://doi.org/10.1038/380526a0

Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J., Eickhoff,

S. B., & Yeo, B. T. T. (2018). Local-Global Parcellation of the Human Cerebral

Cortex from Intrinsic Functional Connectivity MRI. *Cerebral Cortex*, *28*(9), 3095–

3114. https://doi.org/10.1093/cercor/bhx179

Sen, S., Khalsa, N. N., Tong, N., Ovadia-Caro, S., Wang, X., Bi, Y., & Striem-Amit, E.

(2022). The Role of Visual Experience in Individual Differences of Brain

Connectivity. *The Journal of Neuroscience*, *42*(25), 5070–5084.

https://doi.org/10.1523/JNEUROSCI.1700-21.2022

Sereno, M. I., & Huang, R.-S. (2014). Multisensory maps in parietal cortex. *Current Opinion

in Neurobiology*, *24*, 39–46. https://doi.org/10.1016/j.conb.2013.08.014

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-

Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K.,

Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M.

(2004). Advances in functional and structural MR image analysis and implementation

as FSL. *NeuroImage*, *23*(SUPPL. 1), 208–219.

https://doi.org/10.1016/j.neuroimage.2004.07.051

Stellwagen, D., & Shatz, C. J. (2002). An Instructive Role for Retinal Waves in the

Development of Retinogeniculate Connectivity. *Neuron*, *33*(3), 357–367.

https://doi.org/10.1016/S0896-6273(02)00577-9

Striem-Amit, E., Ovadia-Caro, S., Caramazza, A., Margulies, D. S., Villringer, A., & Amedi,

A. (2015). Functional connectivity of visual cortex in the blind follows retinotopic

organization principles. *Brain*, *138*(6), 1679–1695.

https://doi.org/10.1093/brain/awv083

Sur, M., Angelucci, A., & Sharma, J. (1999). Rewiring cortex: The role of patterned activity

in development and plasticity of neocortical circuits. *Journal of Neurobiology*, *41*(1),

33–43. https://doi.org/10.1002/(SICI)1097-4695(199910)41:1%253C33::AID-

NEU6%253E3.0.CO;2-1

Takahashi, E., Folkerth, R. D., Galaburda, A. M., & Grant, P. E. (2012). Emerging Cerebral

Connectivity in the Human Fetal Brain: An MR Tractography Study. *Cerebral*

Cortex, *22*(2), 455–464. https://doi.org/10.1093/cercor/bhr126

Tian, M., Saccone, E. J., Kim, J. S., Kanjlia, S., & Bedny, M. (2022). Sensory modality and

spoken language shape reading network in blind readers of Braille. *Cerebral Cortex*.

Tian, M., Saccone, E. J., Kim, J. S., Shipra Kanjlia, & Bedny, M. (2023). Sensory modality

and spoken language shape reading network in blind readers of Braille. *Cerebral*

Cortex, *33*(6), 2426–2440. https://doi.org/10.1093/cercor/bhac216

Tootell, R. B. H., Mendola, J. D., Hadjikhani, N. K., Ledden, P. J., Liu, A. K., Reppas, J. B.,

Sereno, M. I., & Dale, A. M. (1997). Functional Analysis of V3A and Related Areas

in Human Visual Cortex. *Journal of Neuroscience*, *17*(18), 7060–7078.

https://doi.org/10.1523/JNEUROSCI.17-18-07060.1997

Van Essen, D. C. (2005). A Population-Average, Landmark- and Surface-based (PALS) atlas

of human cerebral cortex. *NeuroImage*, *28*(3), 635–662.

https://doi.org/10.1016/j.neuroimage.2005.06.058

Van Essen, D. C., Lewis, J. W., Drury, H. A., Hadjikhani, N., Tootell, R. B. H., Bakircioglu,

M., & Miller, M. I. (2001). Mapping visual cortex in monkeys and humans using

surface-based atlases. *Vision Research*, *41*(10), 1359–1378.

https://doi.org/10.1016/S0042-6989(01)00045-1

Vasung, L. (2017). Spatiotemporal Relationship of Brain Pathways during Human Fetal

Development Using High-Angular Resolution Diffusion MR Imaging and Histology.

Frontiers in Neuroscience, *11*.

Watkins, K. E., Cowey, A., Alexander, I., Filippini, N., Kennedy, J. M., Smith, S. M., Ragge,

N., & Bridge, H. (2012). Language networks in anophthalmia: Maintained hierarchy

of processing in "visual" cortex. *Brain*, *135*(5), 1566–1577.

https://doi.org/10.1093/brain/aws067

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A Functional Connectivity

Toolbox for Correlated and Anticorrelated Brain Networks. *Brain Connectivity*, *2*(3),

125–141. https://doi.org/10.1089/brain.2012.0073

Yan, C.-G., Wang, X.-D., Zuo, X.-N., & Zang, Y.-F. (2016). DPABI: Data Processing &

873   Analysis for (Resting-State) Brain Imaging. *Neuroinformatics*, *14*(3), 339–351.

874   https://doi.org/10.1007/s12021-016-9299-4

875   Yu, C., Liu, Y., Li, J., Zhou, Y., Wang, K., Tian, L., Qin, W., Jiang, T., & Li, K. (2008).

876   Altered functional connectivity of primary visual cortex in early blindness. *Human*

877   *Brain Mapping*, *29*(5), 533–543. https://doi.org/10.1002/hbm.20420

878   Zhang, H., Shen, D., & Lin, W. (2019). Resting-state functional MRI studies on infant brains:

879   A decade of gap-filling efforts. *NeuroImage*, *185*(July 2018), 664–684.

880   https://doi.org/10.1016/j.neuroimage.2018.07.004