Visual experience shapes functional connectivity between occipital and non-visual networks

Mengyu Tian[1,2] Xiang Xiao[3] Huiqing Hu[4] Rhodri Cusack[4] Marina Bedny[1]

Abstract

Comparisons of visual cortex function across blind and sighted adults reveals effects of experience on human brain function. Since almost all research has been done with adults, little is known about the developmental origins of plasticity. We compared resting state functional connectivity of visual cortices of blind adults ($n = 30$), blindfolded sighted adults ($n = 50$) to a large cohort of sighted infants (Developing Human Connectome Project, $n = 475$). Visual cortices of sighted adults show stronger coupling with non-visual sensory-motor networks (auditory, somatosensory/motor), than with higher-cognitive prefrontal cortices (PFC). In contrast, visual cortices of blind adults show stronger coupling with higher-cognitive PFC than with nonvisual sensory-motor networks. Are infant visual cortices functionally like those of sighted adults? Alternatively, do infants start like blind adults, with vision required to set up the sighted adult pattern? Remarkably, we find that, in infants, secondary visual cortices are more like those of blind adults: stronger coupling with PFC than with nonvisual sensory-motor networks, suggesting that visual experience establishes elements of the sighted-adult long-range connectivity. Infant primary visual cortices are in-between blind and sighted adults i.e., equal PFC and sensory-motor connectivity. The lateralization of occipital-to-frontal connectivity in infants resembles the sighted adults, consistent with reorganization by blindness. These results reveal instructive effects of vision and reorganizing effects of blindness on functional connectivity.

Author affiliations:

1 Center for Educational Science, Institute of Advanced Studies in Humanities and Social Sciences, Beijing Normal University, Zhuhai 519087, China

2 Department of Psychological and Brain Sciences, Johns Hopkins University, Baltimore, Maryland 21218

3 Neuroimaging Research Branch, National Institute on Drug Abuse, National Institutes of
Health, Baltimore, MD 21224, USA

4 Trinity College Institute of Neuroscience and School of Psychology, Trinity College Dublin,
Dublin 2, Ireland

Correspondence to: Mengyu Tian

Full address: No.18, Jinfeng Road, Tangjiawan, Zhuhai City, Guangdong Province, 519087,
P.R.China

E-mail: mengyutian@bnu.edu.cn

Keywords: Visual experience, infants, functional connectivity, visual cortex

Introduction

Relative to sighted adults, visual cortices of adults born blind show enhanced responses during
non-visual tasks, such as reading braille and localizing sounds as well as distinctive patterns of
long-range functional connectivity with non-visual networks (Abboud & Cohen, 2019; Bedny
et al., 2011; Burton et al., 2012, 2014; Butt et al., 2013; Collignon et al., 2011; Deen et al.,
2015; Kanjlia et al., 2016, 2021; Lane et al., 2015; Liu et al., 2007; Sadato et al., 1996; Striem-
Amit et al., 2015; Watkins et al., 2012). Since almost all research thus far has been done with
adults, a key outstanding question concerns the developmental origins of these experience-
based differences.

One possibility is that at birth, infant visual cortices start out in the 'prepared' sighted adult
state and blindness leads to functional reorganization. Alternatively, infant visual cortices may
start out functionally similar to those of blind adults and lifetime visual experience establishes
the sighted adult pattern. To distinguish between these possibilities, we compare the function
of visual cortices across blind adults, sighted adults and a large cohort of 2-week-old sighted
infants (Developing Human Connectome Project, dHCP, $n = 475$). We examined functional
connectivity using resting state data, which provide a common measure of cortical function
across these diverse populations.

To our knowledge no prior studies have compared infants to multiple populations of adults

with different sensory experiences. Previous studies comparing sighted infants to sighted adults

have largely reported similarity across groups (Barttfeld et al., 2018; Doria et al., 2010;

Fransson et al., 2009; Gao et al., 2009; Liu et al., 2008; Zhang et al., 2019). However, these

studies focused on whether large scale functional networks are present in infancy e.g., stronger

connectivity of regions within the visual network, than between visual and auditory regions.

Studies comparing blind and sighted adults find differences across groups in which non-visual

networks are most strongly coupled with the visual system i.e., visual cortices of sighted adults

show   stronger   coupling   with   non-visual   sensory-motor   networks   (i.e.,   auditory,

somatosensory/motor) than higher-cognitive systems; by contrast, in blind adults, visual cortex

coupling is stronger with higher-cognitive prefrontal cortices (PFC) than with non-visual

sensory-motor networks (Abboud & Cohen, 2019; Bedny et al., 2011; Burton et al., 2014; Deen

et al., 2015; Kanjlia et al., 2021; Liu et al., 2007; Qin et al., 2013; Striem-Amit et al., 2015;

Watkins et al., 2012; Yu et al., 2008). In the current study, we compare this experience-

sensitive functional signature across infants, sighted and blind adults.

We measured the connectivity profile of four occipital 'visual' areas that show cross-modal

plasticity in blindness, i.e., respond to non-visual tasks in blind people: Three secondary visual

areas (located in lateral, dorsal and parts of the ventral occipital cortex) and the primary visual

cortex (V1).

The three secondary visual areas have been found to respond to different non-visual tasks in

blind people: language tasks, numerical reasoning tasks and executive control tasks

respectively. Enhanced coupling with PFC in blindness is observed across all of three occipital

regions. However, in blind people each region shows preferential coupling with distinct

subregions of PFC with analogous functional profiles (e.g., language responsive occipital areas

are more coupled with language responsive PFC) (Kanjlia et al., 2016, 2021; Lane et al., 2015).

The precise visual functions of these secondary visual regions in sighted people are not known.

Anatomically, these regions correspond roughly to the location of areas such as motion area

V5/MT+, the lateral occipital complex (LO), V3a and V4v in sighted people (Tootell et al.,

1997; Van Essen et al., 2001).

We also examined connectivity of anatomically defined primary visual cortex (V1), which

likewise shows altered task-based responses and functional connectivity in congenitally blind

adults (Amedi et al., 2003; Bedny et al., 2011; Burton et al., 2014; Butt et al., 2013; Lane et al.,

2015; Raz et al., 2005; Sadato et al., 1996; Striem-Amit et al., 2015; Yu et al., 2008). Since
many previous studies have found that blindness alters the balance of connectivity between
visual cortex and higher-order prefrontal as opposed to sensory-motor regions, this was our
primary outcome measure. We also examined changes in connectivity lateralization – i.e., the
balance of between vs. within hemisphere connectivity.

To preview the results, we find that, in infants, the long-range functional connectivity profile
of secondary visual areas resembles that of blind adults, whereas V1 falls between blind and
sighted adult populations. Relative to sighted adults, both blind adults and infants show
stronger coupling between visual cortices and PFC and weaker coupling between visual
cortices and non-visual sensory-motor networks. This suggests that vision plays an instructive
role in setting up the balance of connectivity between occipital cortex and non-visual networks.
In contrast, connectivity lateralization patterns appear to reflect blindness-related
reorganization.

Results

Connectivity profile of secondary visual cortices in sighted infants is more similar to that of blind than sighted adults

We first examined the long-range functional connectivity of (three) secondary visual areas with
sensory-motor areas on the one hand, and higher-order PFC networks on the other. In sighted
adults, all three secondary visual areas showed stronger functional connectivity with non-visual
sensory-motor areas (primary somatosensory and motor cortex, S1/M1, and primary auditory
cortex, A1) than with higher-cognitive prefrontal cortices (PFC). By contrast, in blind adults,
all secondary visual regions showed stronger functional connectivity with PFC than with non-
visual sensory-motor areas (S1/M1 and A1) (group (sighted adults, blind adults) by ROI (PFC,
non-visual sensory) interaction effect: $F_{(1, 78)} = 148.819$, $p < 0.001$; post-hoc Bonferroni-
corrected paired t-test, sighted adults: non-visual sensory > PFC: $t_{(49)} = 9.722$, $p < 0.001$; blind
adults: non-visual sensory < PFC: $t_{(29)} = 8.852$, $p < 0.001$; Fig. 1).

Like in blind adults, in sighted infants, secondary visual areas showed higher connectivity to
PFC than to non-visual sensory-motor areas (S1/M1 and A1) (non-visual sensory < PFC paired
t-test, $t_{(474)} = 20.144$, $p < 0.001$) (Fig. 1). The connectivity matrix of sighted infants was more
correlated with that of blind than sighted adults, but strongly correlated with both adult groups
(secondary visual, PFC and non-visual sensory areas: sighted infants correlated to blind adults:

$r = 0.721, p < 0.001$; to sighted adults: $r = 0.524, p < 0.001$; difference between correlations of infants to blind vs. to sighted adults: $z = 3.77, p < 0.001$; see Supplementary Figure S1 for the connectivity matrices).

These results suggests that vision is required to set up elements of the sighted adult functional connectivity pattern, i.e., vision enhances occipital cortex connectivity to non-visual sensory-motor networks and dampens connectivity to higher-cognitive prefrontal networks.

We checked the robustness of these results in a number of ways. We first compared the effects across the three secondary visual regions and observed the same pattern across all. (Supplementary results and Supplementary Figure S2). Next, to check the robustness of the findings in infants we randomly split the infant dataset into two halves and did split-half cross-validation. Across all comparisons the results of the two halves were highly similar, suggesting the effects are robust (see Supplementary Figure S3 and S4). We performed this validation procedure for all analyses reported below with similar results.

Figure 1 Functional connectivity of secondary visual cortices. (A) Bar graph shows functional connectivity (*r*) of secondary visual cortices (blue) to non-visual sensory motor areas (purple) and prefrontal cortices (green), averaged across occipital, PFC and sensory-motor ROIs (A1 and S1/M1) in sighted adults, blind adults and sighted infants. Regions of interest (ROI) displayed on the left. Note that regions extend to ventral surface, not shown.

See Supplementary Figure S5 for the full views of three occipital ROIs. **(B)** Circle plots represent the connectivity of secondary visual cortices to non-visual networks, min-max normalized to [0,1], i.e., as a proportion. OC: occipital cortices; MTH: math-responsive region; LG: language-responsive region; EF: executive function-responsive (response-conflict) region. Asterisks (*) denote significant Bonferroni-corrected pairwise comparisons ($p < .05$).

The connectivity pattern of V1 is influenced by early visual experience and blindness

V1 showed the same dissociation between sighted and blind adults as secondary visual areas: in sighted adults, V1 has stronger functional connectivity with non-visual sensory-motor areas than with PFC. By contrast, in blind adults, V1 shows stronger connectivity with PFC than with non-visual sensory areas (group (sighted adults, blind adults) by ROI (PFC, non-visual sensory) interaction: $F_{(1, 78)} = 125.775$, $p < 0.001$; post-hoc Bonferroni-corrected paired t-test, sighted adults non-visual sensory > PFC: $t_{(49)} = 9.404$, $p < 0.001$; blind adults non-visual sensory < PFC: $t_{(29)} = 7.128$, $p < 0.001$; Fig. 2).

The pattern for sighted infants in V1 fell between that of sighted and blind adults. The connectivity matrix of sighted infants (V1, PFC, and non-visual sensory) was equally correlated with blind and sighted adults (infants correlated to blind adults: $r = 0.654$, $p < 0.001$; to sighted adults: $r = 0.594$, $p < 0.001$; correlation of infants with blind vs. with sighted adults: z = 0.832, $p = 0.406$; see Supplementary Figure S1 for the connectivity matrices). The difference in connectivity strength between V1 to PFC and V1 to non-visual sensory regions was weaker in sighted infants than in sighted or blind adults (group (sighted adults, infants) by ROI (PFC, non-visual sensory) interaction effect: $F_{(1, 523)} = 92.21$, $p < 0.001$; group (blind adults, infants) by ROI (PFC, non-visual sensory) interaction effect: $F_{(1, 503)} = 57.444$, $p < 0.001$). V1 of sighted infants showed marginally stronger connectivity to non-visual sensory regions (S1/M1 and A1) than PFC (non-visual sensory regions > PFC, paired t-test, $t_{(474)} = 1.95$, $p = 0.052$; Fig.2).

The dHCP cohort included both full-term neonates and preterm infants, scanned at their equivalent gestational age. Visual exposure therefore varied somewhat in duration across infants (from 0 to 19.71 weeks), with slightly longer exposure in preterm babies. This variation did not affect connectivity patterns either in V1 or secondary visual cortices (V1: $r = 0.06$, $p = 0.192$; secondary visual: $r = 0.004$, $p = 0.923$; see Supplementary Figure S6). We also compared the connectivity patterns of preterm ($n = 90$) and full-term infants ($n = 385$) and

found no difference from each other or from the all-infant dataset (see Supplementary Figure

S7). A few weeks of vision after birth is therefore insufficient to influence connectivity.

Figure 2 Functional connectivity of primary visual cortices (V1). Regions of interest
(ROI) displayed on the upper. Bar graph shows functional connectivity (*r*) of V1 to non-
visual sensory motor areas (purple) and prefrontal cortices (green), averaged across three
PFC ROIs and sensory-motor ROIs (S1/M1 and A1). Asterisks (*) denote significant
Bonferroni-corrected pairwise comparisons ($p < .05$). Cross (†) denote marginal difference in
Bonferroni-corrected pairwise comparisons ($0.05 < p < 0.1$).

Evidence for blindness-related reorganization in laterality of occipito-frontal connectivity

Compared to sighted adults, blind adults exhibit a stronger dominance of within-hemisphere

connectivity over between-hemisphere connectivity. That is, in people born blind, left visual

networks are more strongly connected to left PFC, whereas right visual networks are more

strongly connected to right PFC. By contrast, in sighted adults, this cross-hemisphere

difference is weak or absent. This difference between adult groups is observed for both V1 and

secondary visual cortices (group (blind adults, sighted adults) by lateralization (within

hemisphere, between hemisphere) interaction in secondary visual cortices: $F_{(1, 78)} = 131.51$, p
< 0.001). Secondary visual cortices showed a significant within > between difference in both
groups, with a lager effect in the blind group (post-hoc tests, Bonferroni-corrected paired: t-
test: sighted adults within hemisphere > between hemisphere: $t_{(49)} = 7.441$, $p = 0.012$; blind
adults within hemisphere > between hemisphere: $t_{(29)} = 10.735$, $p < 0.001$; V1: $F_{(1, 78)} = 87.211$,
$p < 0.001$). In V1, only the blind group showed a significant within > between hemisphere
effect (post-hoc Bonferroni-corrected paired: t-test: sighted adults within hemisphere <
between hemisphere: $t_{(49)} = 3.251$, $p = 0.101$; blind adults within hemisphere > between
hemisphere: $t_{(29)} = 7.019$, $p < 0.001$).

With respect to laterality, infants resembled sighted more than blind adults (Fig. 3). For
secondary visual cortices, there was a significant difference between blind adults and sighted
infants and no difference between sighted adults and sighted infants (group (blind adults,
infants) by lateralization (within hemisphere, between hemisphere) interaction effect: $F_{(1, 503)}$
$= 303.04$, $p < 0.001$); (group (sighted adults, infants) by lateralization (within hemisphere,
between hemisphere) interaction effect: $F_{(1, 523)} = 2.244$, $p = 0.135$). A similar group by
laterality interaction were observed for V1 (group (blind adults, infants) by lateralization
(within hemisphere, between hemisphere) interaction: $F_{(1, 503)} = 123.608$, $p < 0.001$; group
(sighted adults, infants) by lateralization (within hemisphere, between hemisphere) interaction
effect: $F_{(1, 523)} = 2.827$, $p = 0.093$). This suggests that the enhancement of within over between
hemisphere long-range connectivity is related to blindness-driven reorganization.

Task-based fMRI studies find that cross-modal responses in occipital cortex colateralize with
fronto-parietal networks with related functions (e.g., language, response selection) (Kanjlia et
al., 2021; Lane et al., 2017). For example, language-responsive occipital areas collateralize
with language responsive prefrontal areas across individuals (Lane et al., 2017). Recruitment
of visual cortices by cross-modal tasks (e.g., spoken language) may enhance within-hemisphere
connectivity in people born blind (Kanjlia et al., 2021; Lane et al., 2017; Tian et al., 2022).
Together, this evidence supports the hypothesis that, starting from the less lateralized infant
state, blindness increases lateralization of occipital long-range connectivity.

219
Figure 3 Within hemisphere vs. between hemisphere functional connectivity. Bar graph
shows within hemisphere (blue) and between hemisphere (orange) functional connectivity (r
coefficient of resting state correlations) of secondary visual (left) and V1 (right) to prefrontal
cortices in sighted adults, blind adults, and sighted infants. Blind adults show a larger difference
than any of the other groups. Asterisks (*) denote significant Bonferroni-corrected pairwise
comparisons ($p < .05$).

Specialization of connectivity across different fronto-occipital networks: present in adults, absent at birth

In blind adults, different occipital areas show enhanced connectivity patterns with distinct
subregions of PFC and this specialization is aligned with the functional specialization observed
in task-based data (Bedny et al., 2011; Kanjlia et al., 2016, 2021). For example, language-
responsive subregions of occipital cortex show strongest functional connectivity with
language-responsive sub-regions of PFC, whereas math-responsive occipital areas show
stronger connectivity with math-responsive PFC. This pattern is most pronounced in blind
people but can be seen weakly even in sighted participants (Fig. 4) (Bedny et al., 2011; Kanjlia
et al., 2016, 2021; Lane et al., 2015). Is this fronto-occipital connectivity specialization present
in infancy, potentially enabling the task-based cross-modal specialization?

We compared connectivity preferences across three prefrontal and three occipital regions
previous shown to activate preferentially in language (sentences > math), math (math >
sentences) and response-conflict (no-go > go with tones) (Kanjlia et al., 2016, 2021; Lane et
al., 2015). For ease of viewing, Fig. 4 shows results from two of the three regions, math and
language. See Supplementary Figure S8 for all three regions. Note that the statistical analyses
included all three areas.

Contrary to the hypothesis that specialization of functional connectivity across different
prefrontal/occipital areas is present from birth, sighted infants showed a less differentiated
fronto-occipital connectivity pattern relative to both blind and sighted adults (Group (sighted
adults, blind adults, sighted infants) by occipital regions (math, language, response-conflict)
by PFC regions (math, language, response-conflict) interaction $F_{(8, 2208)} = 16.323$, $p < 0.001$).
Unlike in adults, in infants, all the occipital regions showed stronger correlations with math-
and response-conflict related prefrontal areas than language-responsive prefrontal areas (Fig.
4 and Supplementary Figure S8). However, the preferential correlation with math responsive
PFC was strongest in those occipital areas that go on to develop math responses in blind adults
(occipital regions (math, language, response-conflict) by PFC regions (math, language,
response-conflict) interaction in infants $F_{(4, 1896)} = 85.145$, $p < 0.001$, post-hoc Bonferroni-
corrected paired t-test see Supplementary Table 1).

Note that findings of reduced regional specialization in infants need to be interpreted with
caution. First, we do not know whether the same specialization of prefrontal sub-regions seen
in adults is present in infants, although, prior evidence suggests some prefrontal specialization
is already present (Raz & Saxe, 2020). Second, the more fine-grained comparisons across
occipital/frontal regions are more vulnerable to potential anatomical alignment issues between
adult and infant brains. In other words, lack of specialization in infants could reflect the
different location of the areas in this population.

Figure 4 Occipito-frontal functional connectivity. Bar graph shows across functional
connectivity of different sub-regions of prefrontal (PFC) and occipital cortex (OCC) in sighted
adults, blind adults, and sighted infants. Sub-regions (regions of interest) were defined based
on task-based responses in a separate dataset of sighted (frontal) and blind (frontal and
occipital) adults (Kanjlia et al., 2016, 2021; Lane et al., 2015). PFC/OCC-MATH: math-
responsive regions were more active when solving math equations than comprehending
sentences. PFC/OC-LANG: language-responsive regions were more active when
comprehending sentences than solving math equations (Kanjlia et al., 2016, 2021; Lane et al.,
2015). In blind adults these regions show biases in connectivity related to their function i.e.,
language-responsive PFC is more correlated with language responsive OCC. No such pattern
is observed in infants. Asterisks (*) denote significant Bonferroni-corrected pairwise
comparisons ($p < .05$). See Supplementary Figure S9 for connectivity matrix.

Discussion

The present results provide insight into the developmental process of experience-based

functional specialization in human cortex. We find independent effects of visual experience

and blindness on the development of visual networks. Aspects of the sighted adult connectivity

pattern require visual experience to become established. This is particularly striking for

secondary visual cortices, where, connectivity with nonvisual networks in sighted infants

resemble blind more than sighted adults. Both in infants and blind adults, secondary occipital

areas showed stronger functional connectivity with higher-order prefrontal cortices than with

other sensory-motor networks (S1/M1, A1). Consistent with this observation, one previous

study with a small sample of infants found strong connectivity between lateral occipital and

prefrontal areas, although there was no comparison to blind adults in that study (Barttfeld et

al., 2018). In V1, infants fell somewhere in between sighted and blind adults, suggesting an

effect both of vision and of blindness on functional connectivity.

The present results reveal the effects of experience on development of functional connectivity

between infancy and adulthood, but do not speak to the precise time course of these effects.

Infants in the current sample had between 0 and 20 weeks of visual experience. Comparisons

across these infants suggests that several weeks of postnatal visual experience is insufficient to

produce a sighted-adult connectivity profile. The time course of development could be

anywhere between a few months and years and could be tested by examining data from children

of different ages.

We propose that vision, as well as temporally coordinated multi-modal experiences

contribute to establishing the sighted adult connectivity profile in visual cortices. For

example, coordinated visuo-motor activity during development may enhance connectivity

between visual and motor networks. Supporting this idea, evidence shows that in infants,

motor competence and early experience predict coupling between occipital and motor

networks (Colomer et al., 2023).Despite these insights, many questions remain regarding the

neurobiological mechanisms underlying experience-based functional connectivity changes

and their relationship to anatomical development. Long-range anatomical connections

between brain regions are already present in infants—even prenatally—though they remain

immature (Huang et al., 2009; Kostović et al., 2019, 2021; Takahashi et al., 2012; Vasung,

2017). Functional connectivity changes may stem from local synaptic modifications within

these stable structural pathways, consistent with findings that functional connectivity can

vary independently of structural connection strength (Fotiadis et al., 2024). Moreover,

functional connectivity has been shown to outperform structural connectivity in predicting

individual behavioral differences, suggesting that experience-based functional changes may

reflect finer-scale synaptic or network-level modulations not captured by macrostructural

measures (Ooi et al., 2022). Prior studies also suggest that, even in adults, coordinated

sensory-motor experience can lead to enhancement of functional connectivity across sensory-

motor systems, indicating that large-scale changes in functional connectivity do not

necessarily require corresponding changes in anatomical connectivity (Guerra-Carrillo et al.,

2014; Li et al., 2018). Another possibility is some connectivity changes are mediated by

'third-party' regions, including subcortical mechanisms such as those involving the thalamus

(Vega-Zuniga et al., 2025).

The current findings reveal both effects of vision and effects of blindness on the functional

connectivity patterns of the visual cortex. A further open question is whether visual

experience plays an instructive or permissive role in shaping neural connectivity patterns. An

instructive role suggests that specific sensory experiences or patterns of neural activity

directly shape and organize neural circuitry. In contrast, a permissive role implies that

sensory experience or neural activity merely facilitates the influence of other factors—such as

molecular signals—on the formation and organization of neural circuits (Crair, 1999; Sur et

al., 1999). Studies with animals that manipulate the pattern or informational content of neural

activity while keeping overall activity levels constant could distinguish between these

hypotheses (Crair, 1999; Roy et al., 2020; Stellwagen & Shatz, 2002).

A further key question concerns the behavioral relevance of the connectivity signatures

observed in the current study. The capacity of occipital cortices to support visual and

multimodal behavior in sighted people may depend not only on local visual cortex function but

also on the capacity of the visual system to coordinate its function with non-visual networks.

Does enhanced connectivity between visual and non-visual sensory motor networks facilitate

multimodal integration for sighted people e.g., when catching a ball? Potentially consistent

with this possibility, recent evidence suggests that people who grew up blind but recover sight

in adulthood show multimodal integration deficits (Ashtari, 2020; Badde et al., 2020; Guerreiro

et al., 2015; Putzar et al., 2007) and distinct occipital oscillations (Pant et al., 2023).

Conversely, for people who remain blind throughout life, visual-PFC connectivity could enable

recruitment of visual cortices for higher-order non-visual functions, such as language and

executive control. Habitual activation of occipital networks during higher cognitive tasks in

early development could intern enhance connectivity and specialization of visual networks for

non-visual functions.

In the current study, the clearest evidence for functional change driven by blindness was

observed for laterality. Connectivity lateralization in sighted infants resembles that of sighted

adults, in both V1 and secondary visual cortices. Relative to both sighted infants and sighted

adults, blind adults show more lateralized connectivity patterns between occipital and

prefrontal cortices. Previous studies suggest that in people born blind occipital and non-

occipital language responses are co-lateralized (Lane et al., 2017). We speculate that habitual

activation of visual cortices by higher-cognitive tasks, such as language, which are themselves

highly lateralized, contributes to this biased connectivity pattern of occipital cortex in blindness.

Materials and methods

Participants

Fifty sighted adults and thirty congenitally blind adults contributed the resting state

data (sighted: n = 50; 30 females; mean age = 35.33 years, standard deviation (SD) = 14.65;

mean years of education = 17.08, SD = 3.1; blind: n = 30; 19 females; mean age = 44.23 years,

SD = 16.41; mean years of education = 17.08, SD = 2.11; blind vs. sighted age, $t_{(78)}$ = 2.512,

$p < 0.05$; blind vs. sighted years of education, $t_{(78)}$ = 0.05, $p = 0.996$). Since blind participants

were on average older, we also performed analyses in an age-matched subgroups of sighted

controls ($n = 29$) and found similar results to the full sample (see Supplementary Figure S10).

Blind and sighted participants had no known cognitive or neurological disabilities (screened

through self-report). All adult anatomical images were read by a board-certified radiologist, and no gross neurological abnormalities were found. All the blind participants had at most minimal light perception from birth. Blindness was caused by pathology anterior to the optic chiasm (i.e., not due to brain damage). All participants gave written informed consent under a protocol approved by the Institutional Review Board of Johns Hopkins University.

Neonate data were from the third release of the Developing Human Connectome Project (dHCP) ($n = 783$) (https ://www.developing connectome.org). Ethical approval was obtained from the UK Health Research Authority (Research Ethics Committee reference number: 14/LO/1169). After quality control procedures (described below), 475 subjects were included in data analysis, with one scan per subject. The average age from birth at scan = 2.79 weeks (SD = 3.77, median = 1.57, range = 0 – 19.71); average gestational age at scan = 41.23 weeks (SD = 1.77, median = 41.29, range = 37 – 45.14); average gestational age at birth = 38.43 weeks (SD = 3.73, median = 39.71, range = 23 – 42.71). We only included infants who were full-term or scanned at term-equivalent age if preterm, while not being flagged by the dHCP project team as not passing quality control for functional MRI (fMRI) images ($n = 634$). Infants with more than 160 motion outliers were exclude ($n = 116$ dropped). Motion-outlier volumes were defined as DVARS (the root mean square intensity difference between successive volumes) higher than 1.5 interquartile range above the 75th centile, after motion and distortion correction. Infants with signal drop-out in regions of interest (ROI) were also excluded ($n = 43$ dropped). To identify signal dropout, we first averaged blood oxygen level-dependent (BOLD) signal intensity for all time point, for each subject, in each of 100 parcel defined by Schaefer's atlas (Schaefer et al., 2018). For each ROI ($n = 18$ ROIs) in the current study, signal dropout was then identified as BOLD intensity lower than -3 standard deviations, where the mean and standard deviations were identified across all 100 cortical parcels. Participants were excluded if any of the ROIs showed a signal dropout. The infants' structural images were reviewed by a pediatric neuroradiologist from the dHCP team, who assigned scores on a scale from 1 to 5. A score of 1 indicated a normal appearance for the subject's age, while scores of 4 or 5 suggested potential or likely clinical significance, or both clinical and imaging relevance. We repeated our analysis after excluding infants with a radiology score of 4 or 5, and the results remained consistent (see Figure S11).

Image acquisition

Blind and sighted adult MRI anatomical and functional images were collected on a 3T Phillips scanner at the F. M. Kirby Research Center. T1-weighted anatomical images were collected using a magnetization-prepared rapid gradient-echo (MP-RAGE) in 150 axial slices with 1 mm isotropic voxels. Resting state fMRI data were collected in 36 sequential ascending axial slices for 8 minutes. TR = 2 s, TE = 0.03 s, flip angle = 70°, voxel size = 2.4 × 2.4 × 2.5 mm, inter-slice gap = 0.5 mm, field of view (FOV) = 192 × 172.8 × 107.5. Participants completed 1 to 4 scans of 240 volume each (average scan time = 710.4 second per person). During the resting state scan, participants were instructed to relax but remain awake. Sighted participants wore light-excluding blindfolds to equalize the light conditions across the groups during the scans.

Infants (dHCP) Anatomical and functional images were collected on a 3T Phillips scanner at the Evelina Newborn Imaging Centre, St Thomas' Hospital, London, UK. A dedicated neonatal imaging 219 system including a neonatal 32-channel phased-array head coil was used. T2w multi-slice fast spin-echo images were acquired with in-plane resolution 0.8x0.8 mm^2 and 1.6 mm slices overlapped by 0.8 mm (TR = 12000 ms, TE = 156 ms, SENSE factor 2.11 axial and 2.6 sagittal). In infants, T2w images were used as the anatomical image because the brain anatomy is more clearly in T2w than in T1w images. Fifteen minutes of resting state fMRI data were collected using a used multiband 9x accelerated echo-planar imaging (TR = 392 ms, TE = 38 ms, 2300 volumes, with an acquired resolution of 2.15 mm isotropic). Single-band reference scans were acquired with bandwidth-matched readout, along with additional spin-echo acquisitions with both AP/PA fold-over encoding directions.

Data analysis

Resting state data were preprocessed using FSL version 5.0.9 (Smith et al., 2004), DPABI version 6.1 (Yan et al., 2016), FreeSurfer (Dale et al., 1999), and in-house code (https://github.com/NPDL/Resting-state_dHCP). The functional data for all groups were linearly detrended and low-pass filtered (0.08 Hz).

For adults, functional images were registered to the T1-weighted structural images, motion corrected using MCFLIRT (Jenkinson et al., 2002), and temporally high-pass filtering (150 s). No subject had excessive head movement (> 2mm) or rotation (> 2°) at any timepoint. Resting state data are known to include artifacts related to physiological fluctuations such as cardiac pulsations and respiratory-induced modulation of the main magnetic field. A component-based

method, CompCor (Behzadi et al., 2007), was therefore used to control for these artifacts.

Particularly, following the procedure described in Whitfield-Gabrieli *et al*., nuisance signals

were extracted from 2-voxel eroded masks of spinal fluid (CSF) and white matter (WM), and

the first 5 principal components analysis (PCA) components derived from these signals was

regressed out from the processed BOLD time series (Whitfield-Gabrieli & Nieto-Castanon,

2012). In addition, a scrubbing procedure was applied to further reduce the effect of motion on

functional connectivity measures (Power et al., 2012, 2014). Frames with root mean square

intensity difference exceeding 1.5 interquartile range above the 75th centile, after motion and

distortion correction, were censored as outliers.

The infants resting state functional data were pre-processed by the dHCP group using the

project's in-house pipeline (Fitzgibbon et al., 2020), This pipeline uses a spatial independent

component analysis (ICA) denoising step to minimize artifact due to multi-band artefact,

residual head-movement, arteries, sagittal sinus, CSF pulsation. For infants, ICA denoising is

preferable to using CSF/WM regressors. Because it is challenging to accurately define

anatomical boundaries of CSF/WM due to the low imaging resolution comparing with the brain

size and the severe partial-volume effect in the neonate (Fitzgibbon et al., 2020). Like in the

adults, frames with root mean square intensity difference exceeding 1.5 interquartile range

above the 75th centile, after motion and distortion correction, were considered as motion

outliers. Out from the 2300 frames, a subset of continuous 1600 with minimum number of

motion outliers was kept for each subject. Motion outliers were censored from the subset of

continuous 1600, and a subject was excluded from further analyses when the number of outlier

exceeded 160 (10% of the continues subset) (Hu et al., 2022). While infant connectivity

estimates may be less robust at the individual level compared to adults due to shorter scan

durations and higher motion, our cohort's large sample size (n=475) and rigorous motion

censoring mitigate these limitations for group-level analyses. Substantial differences between

the groups exist in this study, including the number of subjects, brain sizes, imaging parameters,

and data preprocessing, all of which are likely to have an impact on the overall signal quality.

To address this concern, we compared the split-half noise ceiling across the groups (infants,

sighted adults, and blind adults). For each participant, we first split the rs-fMRI time series into

two halves, then calculated the ROI-wise rsFC pattern from the two splits. The split-half noise

ceiling was estimated according to Lage-Castellanos et al (Lage-Castellanos et al., 2019). The

noise ceilings of the three groups (infants: 0.90 ± 0.056, blind adults: 0.88 ± 0.041, sighted

adults: 0.90 ± 0.055) showed no significant difference (One-way ANOVA, F (2,552) = 2.348, p = 0.097). Therefore, overall signal quality is unlikely to impact our results.

For both groups of adult and infants, we performed a temporal low-pass filter (0.08 Hz low-pass cutoff) and a linear detrending. ROI-to-ROI connectivity was calculated using Pearson's correlation between ROI-averaged BOLD timeseries (ROI definition see below). The All t-tests and F-tests are two-sided. The comparison of correlation coefficients was done using cocor software package and Pearson and Filon's z (Diedenhofen & Musch, 2015; Pearson & Filon, 1898).

ROI definition

Frontal and secondary visual ROIs were defined functionally based on data from a separate task-based fMRI experiments with blind and sighted adults (Kanjlia et al., 2016, 2021; Lane et al., 2015). Three separate experiments were conducted with the same group of blind and sighted subjects (sighted n=18; blind n=23). The language ROIs in the occipital and frontal cortices were identified by sentence > nonwords contrast in an auditory language comprehension task (Lane et al., 2015). The math ROIs were identified by math > sentence contrast in an auditory task where participants judged equivalence of pairs of math equations and pairs of sentences (Kanjlia et al., 2016). The executive function ROIs were identified by no-go > frequent go contrast in an auditory go/no-go task with non-verbal sounds (Kanjlia et al., 2021). All ROI files are available at openICPSR (https://doi.org/10.3886/E198832V1).

Occipital secondary 'visual' ROIs were defined based on group comparisons blind > sighted in a whole-cortex analysis (Figure 1, Figure S5.) The occipital language ROI was defined as the cluster that responded more to auditory sentence than auditory nonwords conditions in blind, relative to sighted, in a whole-cortex analysis, likewise the occipital math ROI was defined as math>sentences, blind>sighted interaction and the occipital executive ROI as no-go > frequent go, blind>sighted. All three occipital ROIs were defined in the right hemisphere. Left-hemisphere occipital ROIs were created by flipping the right-hemisphere ROIs to the left hemisphere. Each functional ROI spans multiple anatomical regions and together the secondary visual ROIs tile large portions of lateral occipital, occipito-temporal, dorsal occipital and occipito-parietal cortices. In sighted people, the secondary visual occipital ROIs include the anatomical locations of functional regions such as motion area V5/MT+, the lateral occipital complex (LO), category specific ventral occipitotemporal cortices and dorsally, V3a and V4v.

The occipital ROI also covers the middle of the ventral temporal lobe. Dorsally, it extended to
the intraparietal sulcus and superior parietal lobule.

The frontal PFC ROIs were defined functionally, based on a whole-cortex analysis which
combined all blind and sighted adult data. The frontal language ROI was defined as responded
more auditory sentence than auditory nonwords conditions across all blind and sighted subjects,
constrained to the prefrontal cortex. Likewise, math responsive PFC was defined as
math>sentences and executive no-go > frequent go. For frontal ROIs, the language ROI was
defined in the left, and the math and executive function ROI were defined in the right
hemisphere, then flip to the other hemisphere.

The V1 ROI was defined from a previously published anatomical surface-based atlas (PALS-
B12) (Van Essen, 2005). The primary somatosensory and motor cortex (S1/M1) ROI was
selected as the area that responds more to the go than no-go trials in the auditory go/no-go task
across both blind and sighted groups, constrained to the hand area in S1/M1 search-space from
neurosynth.org (term "hand movements") (Kanjlia et al., 2021). The primary auditory cortex
(A1) ROI was defined as the transverse temporal portion of a gyral-based atlas (Desikan et al.,
2006; Morosan et al., 2001).

All the ROIs were defined in standard space and then transformed into each subject's native
space. For adults this was done by employing the deformation field estimated by FreeSurfer.
For infants, the ROIs were transformed into each subject's native space using a two-step
approach. First, the ROIs were converted from the adult's MNI space into the 40-week dHCP
template (Bozek et al., 2018). ANTS, previously shown to be effective in pediatric studies
(Avants et al., 2014; Cabral et al., 2022; Jain et al., 2012; Lawson et al., 2013), was utilized to
estimate the deformation field between these two spaces. In this step, the infant's scalp and
cerebellum were masked, as these structures in the infant brain greatly differ from those in the
adult and can introduce bias into the registration process, as outlined in a study by Cabral *et al*
(Cabral et al., 2022). Secondly, the ROIs were further transformed from the 40-week template
space into each individual's native spaces, employing the deformation field provided by the
dHCP group. Nearest neighbor interpolation was applied in both steps (Examples of the
resulting ROI alignment on individual brains are shown in Supplementary Figure S12). For
both adults and infants, any overlapping voxels between ROIs were removed and not counted
toward any ROIs.

Data availability

Neonate data were from the second and third release of the Developing Human Connectome Project (https://www.developing connectome.org). The de-identified blind and sighted adults' data were posted on openICPSR (https://doi.org/10.3886/E198832V1).

Acknowledgements

We would like to thank all the blind and sighted participants, the blind community and the National Federation of the Blind. Without their support, this study would not be possible. We would also like to thank the F. M. Kirby Research Center for Functional Brain Imaging at the Kennedy Krieger Institute for their assistance in data collection. Xiang Xiao was supported by the Intramural Research Program of the National Institute on Drug Abuse, the National Institute of Health, United States.

Funding

This work was supported by grants from the National Eye Institute at the National Institutes of Health (R01EY027352-01 and R01EY033340). RC was supported by the ERC Advanced Grant "Foundations of Cognition" (FOUNDCOG) 787981.

Competing interests

The authors report no competing interests.

References

Abboud, S., & Cohen, L. (2019). Distinctive Interaction Between Cognitive Networks and the Visual Cortex in Early Blind Individuals. *Cerebral Cortex*, *33*(0), 4725–4742. https://doi.org/10.1093/cercor/bhz006

Amedi, A., Raz, N., Pianka, P., Malach, R., & Zohary, E. (2003). Early "visual" cortex activation correlates with superior verbal memory performance in the blind. *Nature Neuroscience*, *6*(7), 758–766. https://doi.org/10.1038/nn1072

Ashtari, M. (2020). Compensatory Cross-Modal Plasticity Persists After Sight Restoration.
Frontiers in Neuroscience, *14*, 16.

Avants, B. B., Tustison, N. J., Stauffer, M., Song, G., Wu, B., & Gee, J. C. (2014). The Insight
ToolKit image registration framework. *Frontiers in Neuroinformatics*, *8*.
https://www.frontiersin.org/articles/10.3389/fninf.2014.00044

Badde, S., Ley, P., Rajendran, S. S., Shareef, I., Kekunnaya, R., & Röder, B. (2020). Sensory
experience during early sensitive periods shapes cross-modal temporal biases. *eLife*, *9*,
e61238. https://doi.org/10.7554/eLife.61238

Barttfeld, P., Abboud, S., Lagercrantz, H., Adén, U., Padilla, N., Edwards, A. D., Cohen, L.,
Sigman, M., Dehaene, S., & Dehaene-Lambertz, G. (2018). A lateral-to-mesial
organization of human ventral visual cortex at birth. *Brain Structure and Function*, *223*(7),
3107–3119. https://doi.org/10.1007/s00429-018-1676-3

Bedny, M., Pascual-Leone, A., Dodell-Feder, D., Fedorenko, E., & Saxe, R. (2011). Language
processing in the occipital cortex of congenitally blind adults. *Proceedings of the National*
Academy of Sciences of the United States of America, *108*(11), 4429–4434.
https://doi.org/10.1073/pnas.1014818108

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction
method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, *37*(1), 90–101.
https://doi.org/10.1016/j.neuroimage.2007.04.042

Bozek, J., Makropoulos, A., Schuh, A., Fitzgibbon, S., Wright, R., Glasser, M. F., Coalson, T.
S., O'Muircheartaigh, J., Hutter, J., Price, A. N., Cordero-Grande, L., Teixeira, R. P. A.
G., Hughes, E., Tusor, N., Baruteau, K. P., Rutherford, M. A., Edwards, A. D., Hajnal, J.
V., Smith, S. M., … Robinson, E. C. (2018). Construction of a neonatal cortical surface
atlas using Multimodal Surface Matching in the Developing Human Connectome Project.
NeuroImage, *179*, 11–29. https://doi.org/10.1016/j.neuroimage.2018.06.018

Burton, H., Sinclair, R. J., & Agato, A. (2012). Recognition memory for Braille or spoken
words: An fMRI study in early blind. *Brain Research*, *1438*, 22–34.
https://doi.org/10.1016/j.brainres.2011.12.032

Burton, H., Snyder, A. Z., & Raichle, M. E. (2014). Resting state functional connectivity in
early blind humans. *Frontiers in Systems Neuroscience*, *8*(1 APR), 1–13.
https://doi.org/10.3389/fnsys.2014.00051

Butt, O. H., Benson, N. C., Datta, R., & Aguirre, G. K. (2013). The fine-scale functional
correlation of striate cortex in sighted and blind people. *Journal of Neuroscience*, *33*(41),
16209–16219. https://doi.org/10.1523/JNEUROSCI.0363-13.2013

Cabral, L., Zubiaurre-Elorza, L., Wild, C. J., Linke, A., & Cusack, R. (2022). Anatomical
correlates of category-selective visual regions have distinctive signatures of connectivity
in neonates. *Developmental Cognitive Neuroscience*, *58*, 101179.
https://doi.org/10.1016/j.dcn.2022.101179

Collignon, O., Vandewalle, G., Voss, P., Albouy, G., Charbonneau, G., Lassonde, M., &
Lepore, F. (2011). Functional specialization for auditory-spatial processing in the
occipital cortex of congenitally blind humans. *Proceedings of the National Academy of*
Sciences, *108*(11), 4435–4440. https://doi.org/10.1073/pnas.1013928108

Colomer, M., Chung, H., Meyer, M., Debnath, R., Morales, S., Fox, N. A., & Woodward, A.
(2023). Action experience in infancy predicts visual-motor functional connectivity during
action anticipation. *Developmental Science*, *26*(3), e13339.
https://doi.org/10.1111/desc.13339

Crair, M. C. (1999). Neuronal activity during development: Permissive or instructive? *Current*
Opinion in Neurobiology, *9*(1), 88–93. https://doi.org/10.1016/S0959-4388(99)80011-7

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis: I.
Segmentation and surface reconstruction. *NeuroImage*, *9*(2), 179–194.
https://doi.org/10.1006/nimg.1998.0395

Deen, B., Saxe, R., & Bedny, M. (2015). Occipital Cortex of Blind Individuals Is Functionally
Coupled with Executive Control Areas of Frontal Cortex. *Journal of Cognitive*
Neuroscience, *27*(8), 1633–1647. https://doi.org/10.1162/jocn_a_00807

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner,
R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006).
An automated labeling system for subdividing the human cerebral cortex on MRI scans
into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980.
https://doi.org/10.1016/j.neuroimage.2006.01.021

Diedenhofen, B., & Musch, J. (2015). cocor: A Comprehensive Solution for the Statistical
Comparison of Correlations. *PLOS ONE*, *10*(4), e0121945.
https://doi.org/10.1371/journal.pone.0121945

Doria, V., Beckmann, C. F., Arichi, T., Merchant, N., Groppo, M., Turkheimer, F. E., Counsell, S. J., Murgasova, M., Aljabar, P., Nunes, R. G., Larkman, D. J., Rees, G., & Edwards, A. D. (2010). Emergence of resting state networks in the preterm human brain. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(46), 20015–20020. https://doi.org/10.1073/pnas.1007921107

Fitzgibbon, S. P., Harrison, S. J., Jenkinson, M., Baxter, L., Robinson, E. C., Bastiani, M., Bozek, J., Karolis, V., Cordero Grande, L., Price, A. N., Hughes, E., Makropoulos, A., Passerat-Palmbach, J., Schuh, A., Gao, J., Farahibozorg, S. R., O'Muircheartaigh, J., Ciarrusta, J., O'Keeffe, C., … Andersson, J. (2020). The developing Human Connectome Project (dHCP) automated resting-state functional processing framework for newborn infants. *NeuroImage*, *223*(August), 117303. https://doi.org/10.1016/j.neuroimage.2020.117303

Fotiadis, P., Parkes, L., Davis, K. A., Satterthwaite, T. D., Shinohara, R. T., & Bassett, D. S. (2024). Structure–function coupling in macroscale human brain networks. *Nature Reviews Neuroscience*, *25*(10), 688–704. https://doi.org/10.1038/s41583-024-00846-6

Fransson, P., Skiöld, B., Engström, M., Hallberg, B., Mosskin, M., Åden, U., Lagercrantz, H., & Blennow, M. (2009). Spontaneous brain activity in the newborn brain during natural sleep-an fMRI study in infants born at full term. *Pediatric Research*, *66*(3), 301–305. https://doi.org/10.1203/PDR.0b013e3181b1bd84

Gao, W., Zhu, H., Giovanello, K. S., Smith, J. K., Shen, D., Gilmore, J. H., & Lin, W. (2009). Evidence on the emergence of the brain's default network from 2-week-old to 2-year-old healthy pediatric subjects. *Proceedings of the National Academy of Sciences*, *106*(16), 6790–6795. https://doi.org/10.1073/pnas.0811221106

Guerra-Carrillo, B., MacKey, A. P., & Bunge, S. A. (2014). Resting-state fMRI: A window into human brain plasticity. *Neuroscientist*, *20*(5), 522–533. https://doi.org/10.1177/1073858414524442

Guerreiro, M. J. S., Putzar, L., & Röder, B. (2015). The effect of early visual deprivation on the neural bases of multisensory processing. *Brain*, *138*(6), 1499–1504. https://doi.org/10.1093/brain/awv076

Hu, H., Cusack, R., & Naci, L. (2022). Typical and disrupted brain circuitry for conscious awareness in full-term and preterm infants. *Brain Communications*, *4*(2), fcac071. https://doi.org/10.1093/braincomms/fcac071

Huang, H., Xue, R., Zhang, J., Ren, T., Richards, L. J., Yarowsky, P., Miller, M. I., & Mori, S.
(2009). Anatomical Characterization of Human Fetal Brain Development with Diffusion
Tensor Magnetic Resonance Imaging. *The Journal of Neuroscience*, *29*(13), 4263–4273.
https://doi.org/10.1523/JNEUROSCI.2769-08.2009

Jain, V., Duda, J., Avants, B., Giannetta, M., Xie, S. X., Roberts, T., Detre, J. A., Hurt, H.,
Wehrli, F. W., & Wang, D. J. J. (2012). Longitudinal Reproducibility and Accuracy of
Pseudo-Continuous Arterial Spin–labeled Perfusion MR Imaging in Typically
Developing Children. *Radiology*, *263*(2), 527–536.
https://doi.org/10.1148/radiol.12111509

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the
robust and accurate linear registration and motion correction of brain images. *NeuroImage*,
17(2), 825–841. https://doi.org/10.1016/s1053-8119(02)91132-8

Kanjlia, S., Lane, C., Feigenson, L., & Bedny, M. (2016). Absence of visual experience
modifies the neural basis of numerical thinking. *Proceedings of the National Academy of
Sciences of the United States of America*, *113*(40), 11172–11177.
https://doi.org/10.1073/pnas.1524982113

Kanjlia, S., Loiotile, R. E., Harhen, N., & Bedny, M. (2021). 'Visual' cortices of congenitally
blind adults are sensitive to response selection demands in a go/no-go task. *NeuroImage*,
236(April), 118023. https://doi.org/10.1016/j.neuroimage.2021.118023

Kostović, I., Katušić, A., & Kostović Srzentić, M. (2021). Linking histology and neurological
development of the fetal and infant brain. In *Factors Affecting Neurodevelopment* (pp.
213–225). Elsevier. https://doi.org/10.1016/B978-0-12-817986-4.00019-5

Kostović, I., Sedmak, G., & Judaš, M. (2019). Neural histology and neurogenesis of the human
fetal and infant brain. *NeuroImage*, *188*, 743–773.
https://doi.org/10.1016/j.neuroimage.2018.12.043

Lage-Castellanos, A., Valente, G., Formisano, E., & Martino, F. D. (2019). Methods for
computing the maximum performance of computational models of fMRI responses. *PLOS
Computational Biology*, *15*(3), e1006397. https://doi.org/10.1371/journal.pcbi.1006397

Lane, C., Kanjlia, S., Omaki, A., & Bedny, M. (2015). "Visual" cortex of congenitally blind
adults responds to syntactic movement. *Journal of Neuroscience*, *35*(37), 12859–12868.
https://doi.org/10.1523/JNEUROSCI.1256-15.2015

Lane, C., Kanjlia, S., Richardson, H., Fulton, A., Omaki, A., & Bedny, M. (2017). Reduced Left Lateralization of Language in Congenitally Blind Individuals. *Journal of Cognitive Neuroscience*, *29*(1), 65–78. https://doi.org/10.1162/jocn_a_01045

Lawson, G. M., Duda, J. T., Avants, B. B., Wu, J., & Farah, M. J. (2013). Associations between children's socioeconomic status and prefrontal cortical thickness. *Developmental Science*, *16*(5), 641–652. https://doi.org/10.1111/desc.12096

Li, Q., Wang, X., Wang, S., Xie, Y., Li, X., Xie, Y., & Li, S. (2018). Musical training induces functional and structural auditory-motor network plasticity in young adults. *Human Brain Mapping*, *39*(5), 2098–2110. https://doi.org/10.1002/hbm.23989

Liu, W. C., Flax, J. F., Guise, K. G., Sukul, V., & Benasich, A. A. (2008). Functional connectivity of the sensorimotor area in naturally sleeping infants. *Brain Research*, *1223*, 42–49. https://doi.org/10.1016/j.brainres.2008.05.054

Liu, Y., Yu, C., Liang, M., Li, J., Tian, L., Zhou, Y., Qin, W., Li, K., & Jiang, T. (2007). Whole brain functional connectivity in the early blind. *Brain*, *130*(8), 2085–2096. https://doi.org/Neuronal recycling

Morosan, P., Rademacher, J., Schleicher, A., Amunts, K., Schormann, T., Zilles, K., & Zilles, K. (2001). Human primary auditory cortex: Cytoarchitectonic subdivisions and mapping into a spatial reference system. *NeuroImage*, *13*(4), 684–701. https://doi.org/10.1006/nimg.2000.0715

Ooi, L. Q. R., Chen, J., Zhang, S., Kong, R., Tam, A., Li, J., Dhamala, E., Zhou, J. H., Holmes, A. J., & Yeo, B. T. T. (2022). Comparison of individualized behavioral predictions across anatomical, diffusion and functional connectivity MRI. *NeuroImage*, *263*, 119636. https://doi.org/10.1016/j.neuroimage.2022.119636

Pant, R., Ossandón, J., Stange, L., Shareef, I., Kekunnaya, R., & Röder, B. (2023). Stimulus-evoked and resting-state alpha oscillations show a linked dependence on patterned visual experience for development. *NeuroImage: Clinical*, *38*, 103375. https://doi.org/10.1016/j.nicl.2023.103375

Pearson, K., & Filon, L. N. G. (1898). Mathematical contributions to the theory of evolution. IV. On the probable errors of frequency constants and on the influence of random selection on variation and correlation. *Proceedings of the Royal Society of London*, *62*(379–387), 173–176. https://doi.org/10.1098/rspl.1897.0091

Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage*, *59*(3), 2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*, *84*, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Putzar, L., Goerendt, I., Lange, K., Rösler, F., & Röder, B. (2007). Early visual deprivation impairs multisensory interactions in humans. *Nature Neuroscience*, *10*(10), 1243–1245. https://doi.org/10.1038/nn1978

Qin, W., Liu, Y., Jiang, T., & Yu, C. (2013). The Development of Visual Areas Depends Differently on Visual Experience. *PLoS ONE*, *8*(1), 1–10. https://doi.org/10.1371/journal.pone.0053784

Raz, G., & Saxe, R. (2020). Learning in Infancy Is Active, Endogenously Motivated, and Depends on the Prefrontal Cortices. *Annual Review of Developmental Psychology*, *2*(1), 247–268. https://doi.org/10.1146/annurev-devpsych-121318-084841

Raz, N., Amedi, A., & Zohary, E. (2005). V1 activation in congenitally blind humans is associated with episodic retrieval. *Cerebral Cortex*, *15*(9), 1459–1468. https://doi.org/10.1093/cercor/bhi026

Roy, A., Wang, S., Meschede-Krasa, B., Breffle, J., & Van Hooser, S. D. (2020). An early phase of instructive plasticity before the typical onset of sensory experience. *Nature Communications*, *11*(1), 11. https://doi.org/10.1038/s41467-019-13872-1

Sadato, N., Pascual-Leone, A., Grafman, J., Ibañez, V., Deiber, M., Dold, G., & Hallett, M. (1996). Activation of the primary visual cortex by Braille reading in blind subjects. *Nature*, *380*(6574), 526–528. https://doi.org/10.1038/380526a0

Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J., Eickhoff, S. B., & Yeo, B. T. T. (2018). Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cerebral Cortex*, *28*(9), 3095–3114. https://doi.org/10.1093/cercor/bhx179

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K.,

Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M.
(2004). Advances in functional and structural MR image analysis and implementation as
FSL. *NeuroImage*, *23*(SUPPL. 1), 208–219.
https://doi.org/10.1016/j.neuroimage.2004.07.051

Stellwagen, D., & Shatz, C. J. (2002). An Instructive Role for Retinal Waves in the
Development of Retinogeniculate Connectivity. *Neuron*, *33*(3), 357–367.
https://doi.org/10.1016/S0896-6273(02)00577-9

Striem-Amit, E., Ovadia-Caro, S., Caramazza, A., Margulies, D. S., Villringer, A., & Amedi,
A. (2015). Functional connectivity of visual cortex in the blind follows retinotopic
organization principles. *Brain*, *138*(6), 1679–1695. https://doi.org/10.1093/brain/awv083

Sur, M., Angelucci, A., & Sharma, J. (1999). Rewiring cortex: The role of patterned activity in
development and plasticity of neocortical circuits. *Journal of Neurobiology*, *41*(1), 33–43.
https://doi.org/10.1002/(SICI)1097-4695(199910)41:1<33::AID-NEU6>3.0.CO;2-1

Takahashi, E., Folkerth, R. D., Galaburda, A. M., & Grant, P. E. (2012). Emerging Cerebral
Connectivity in the Human Fetal Brain: An MR Tractography Study. *Cerebral Cortex*,
22(2), 455–464. https://doi.org/10.1093/cercor/bhr126

Tian, M., Saccone, E. J., Kim, J. S., Kanjlia, S., & Bedny, M. (2022). Sensory modality and
spoken language shape reading network in blind readers of Braille. *Cerebral Cortex*.

Tootell, R. B. H., Mendola, J. D., Hadjikhani, N. K., Ledden, P. J., Liu, A. K., Reppas, J. B.,
Sereno, M. I., & Dale, A. M. (1997). Functional Analysis of V3A and Related Areas in
Human Visual Cortex. *Journal of Neuroscience*, *17*(18), 7060–7078.
https://doi.org/10.1523/JNEUROSCI.17-18-07060.1997

Van Essen, D. C. (2005). A Population-Average, Landmark- and Surface-based (PALS) atlas
of human cerebral cortex. *NeuroImage*, *28*(3), 635–662.
https://doi.org/10.1016/j.neuroimage.2005.06.058

Van Essen, D. C., Lewis, J. W., Drury, H. A., Hadjikhani, N., Tootell, R. B. H., Bakircioglu,
M., & Miller, M. I. (2001). Mapping visual cortex in monkeys and humans using surface-
based atlases. *Vision Research*, *41*(10), 1359–1378. https://doi.org/10.1016/S0042-
6989(01)00045-1

Vasung, L. (2017). Spatiotemporal Relationship of Brain Pathways during Human Fetal
Development Using High-Angular Resolution Diffusion MR Imaging and Histology.
Frontiers in Neuroscience, *11*.

Vega-Zuniga, T., Sumser, A., Symonova, O., Koppensteiner, P., Schmidt, F. H., & Joesch, M.
(2025). A thalamic hub-and-spoke network enables visual perception during action by
coordinating visuomotor dynamics. *Nature Neuroscience*.
https://doi.org/10.1038/s41593-025-01874-w

Watkins, K. E., Cowey, A., Alexander, I., Filippini, N., Kennedy, J. M., Smith, S. M., Ragge,
N., & Bridge, H. (2012). Language networks in anophthalmia: Maintained hierarchy of
processing in "visual" cortex. *Brain*, *135*(5), 1566–1577.
https://doi.org/10.1093/brain/aws067

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A Functional Connectivity
Toolbox for Correlated and Anticorrelated Brain Networks. *Brain Connectivity*, *2*(3),
125–141. https://doi.org/10.1089/brain.2012.0073

Yan, C.-G., Wang, X.-D., Zuo, X.-N., & Zang, Y.-F. (2016). DPABI: Data Processing &
Analysis for (Resting-State) Brain Imaging. *Neuroinformatics*, *14*(3), 339–351.
https://doi.org/10.1007/s12021-016-9299-4

Yu, C., Liu, Y., Li, J., Zhou, Y., Wang, K., Tian, L., Qin, W., Jiang, T., & Li, K. (2008). Altered
functional connectivity of primary visual cortex in early blindness. *Human Brain Mapping*,
29(5), 533–543. https://doi.org/10.1002/hbm.20420

Zhang, H., Shen, D., & Lin, W. (2019). Resting-state functional MRI studies on infant brains:
A decade of gap-filling efforts. *NeuroImage*, *185*(July 2018), 664–684.
https://doi.org/10.1016/j.neuroimage.2018.07.004