Title: Plasticity of gene expression in the nervous system by exposure to environmental
odorants that inhibit HDACs

AUTHORS: Sachiko Haga-Yamanaka[*,1], Rogelio Nuñez-Flores[*,1,5], Christi Ann Scott[*,2], Sarah
Perry[3], Stephanie Turner Chen[2], Crystal Pontrello[1], Meera Goh Nair[5] and Anandasankar Ray[1,2,3]

ADDRESS:

[1]Department of Molecular, Cell and Systems Biology

[2]Cell, Molecular and Developmental Biology Program

[3]Genetics, Genomics and Bioinformatics Program

[4]Department of Nematology

[5]Division of Biomedical Sciences

University of California, Riverside, CA 92521, USA

* these authors contributed equally

CORRESPONDING AUTHOR: Anandasankar Ray

Department of Molecular Cell and Systems Biology, University of California Riverside, 3401
Watkins Drive, Riverside, CA 92521, USA

CONTACT INFORMATION : Phone : +1(951) 827-5998, Email : anand.ray@ucr.edu

24
25
26

ABSTRACT

Eukaryotes are often exposed to microbes and respond to their secreted metabolites, such as the microbiome in animals or commensal bacteria in roots. Little is known about the effects of long-term exposure to volatile chemicals emitted by microbes, or other volatiles that we are exposed to over a long duration. Using the model system *Drosophila melanogaster,* we evaluate a yeast emitted volatile, diacetyl, found in high levels around fermenting fruits where they spend long periods of time. We find that exposure to just the headspace containing the volatile molecules can alter gene expression in the antenna. Experiments showed that diacetyl and structurally related volatile compounds inhibited human histone-deacetylases (HDACs), increased histone-H3K9 acetylation in human cells, and caused wide changes in gene expression in both *Drosophila* and mice. Diacetyl crosses the blood-brain barrier and exposure causes modulation of gene expression in the brain, therefore has potential as a therapeutic. Using two separate disease models known to be responsive to HDAC-inhibitors, we evaluated physiological effects of volatile exposure. First, we find that the HDAC inhibitor also halts proliferation of a neuroblastoma cell line in culture as predicted. Next, exposure to vapors slows progression of neurodegeneration in a *Drosophila* model for Huntington's disease. These changes strongly suggest that unbeknown to us, certain volatiles in the surroundings can have profound effects on histone acetylation, gene expression and physiology in animals.

INTRODUCTION

Multicellular organisms evolve in the presence of microbial volatiles and metabolites (1, 2), often for prolonged periods of time. Human-associated microbes have incredible metabolic diversity, encoding 100x as many genes as the human genome itself(1). It is estimated that about half of the thousands of metabolites found in mammalian blood are either produced or altered by microbes (3). Microbial metabolites are an important mechanism by which the microbiome influences host immunity amongst other things (2, 4). Many volatiles are generated from the metabolism of a variety of food components, including triglycerides, sugars and amino acids (5). Some chemicals are also produced by microorganisms such as yeasts and lactic bacteria during fermentation in many foods and beverages (5).

In the model system *Drosophila melanogaster*, prolonged exposure to CO_2 showed an increase in volume of the CO_2-sensing glomerulus in the antennal lobe, which can lead to adaptations in fly behavior (6, 7, 8). Carbon dioxide is a common byproduct of yeast fermentation in overripe fruits, the preferred food source. In order to study the physiological effects of long-term exposure to other volatiles emitted during fermentation, we exposed flies for 5 days to diacetyl, a buttery odorant made at high levels by yeast in fermenting fruits (1350 ppb in ripe banana), that is also found in many commonly consumed foods such as butter, yoghurt, wine, fruits, beer, popcorn (5, 9-11). Diacetyl was selected for two reasons: first, *Drosophila melanogaster* use fermenting fruits as their favored food source, oviposition site and spend long durations of time exposed to it. Second, diacetyl is known to be an inhibitory odorant of the CO_2 receptor and therefore an opposite effect is expected on the V-glomerulus from the previous study (12). Follow up experiments revealed that diacetyl acts as an HDAC inhibitor in vitro. We found that volatiles structurally like diacetyl have similar HDAC-inhibitory properties as well.

Subsequent analyses with two different model systems: *Drosophila melanogaster* and *Mus*

musculus showed gene expression changes in tissues caused by odor exposure from a distant

source. We found expression of hundreds of genes to be modulated upon exposure to the volatile,

as would be expected from an inhibitor of HDACs. HDACs are histone-modifying enzymes

involved in the removal of acetyl groups from lysine residues and the remodeling of chromatin

structure to regulate gene expression (13, 14). With large-scale roles in gene regulation, HDACs

are promising targets in drug development for many diseases such as cancers and

neurodegenerative disorders (15-17). Several classes of orally administered HDAC inhibitors have

been found to attenuate the progression of certain cancers and neurodegenerative diseases

including Alzheimer's disease and Huntington's disease (17). We determined that exposure to

diacetyl volatiles substantially altered gene expression in the mouse brain, presumably by crossing

the blood-brain barrier. Diacetyl exposure also slowed degeneration of photoreceptor cells in a

Huntington's disease model in *Drosophila*. Last, exposure to diacetyl caused dramatic reduction

in proliferation of a neuroblastoma cell line in culture. Our discovery of a family of volatile

odorants that also act as HDAC inhibitors highlights the need to understand how small molecules

present in our environment interact with and alter eukaryotic gene expression.

RESULTS

Activity-dependent expression modulation of chemosensory receptors

Previous studies demonstrated that prolonged exposure of ~5 days to elevated CO_2,

causes an increase in volume of the CO_2-sensitive V-glomerulus in the antennal lobe, which can

lead to adaptations in fly behavior (6, 7, 8). To assess the effect of CO_2 response inhibitor

diacetyl on the antennal lobe, flies were exposed to the odorant during the first days immediately

following eclosion. The CO_2-detecting ab1C neuron was labelled using the promoters of the CO_2

receptors, *Gr63a-Gal4; UAS-mcd8GFP* and *Gr21a-Gal4; UAS-mcd8GFP*, and the flies were

exposed to headspace above a 1 % (V/V) diacetyl solution in an air-tight container. The flies

were then imaged for expression of GFP in the ab1C neurons of the antenna, as well as the V

glomerulus, which receives axonal input from ab1C neurons in the antennal lobe (18).

Surprisingly, the *Gr63a* promoter driven GFP signal in the V glomerulus was completely

abolished after 5 days of exposure to the odor for all brains imaged (Figure 1A,1B). However,

there was also a concomitant decrease in GFP signal in ab1C neurons in the antenna (Figure 1A).

Similar results were seen for *Gr21a* driven expression, where expression of GFP was lost in both

the V-glomerulus and ab1C neurons (Figure S1A, S1B). Interestingly, after five days of recovery

in clean air the GFP expression was regained in the V-glomerulus as well as in ab1C neurons in

the antenna, indicating that both *Gr21a* and *Gr63a* expression had been restored (Figure S2).

Because olfactory neurons do not regenerate in the fruit fly (19, 20), these results suggest that

changes in GFP levels are likely due to changes in the promoter expression and not due to

neuronal cell death.

In order to test whether expression of *Gr63a* and *Gr21a* is being downregulated by

diacetyl we relied on quantitative-PCR along with a few members of the *Odorant receptor (Or)*

family genes. Apart from *Gr63a* and *Gr21a,* we also evaluated receptors where diacetyl has no

activity (*Or47a*, *Or88a*) and the broadly expressed co-receptor (*Or83b*). Wild-type flies that

were exposed to diacetyl for 6 days showed dramatic downregulation of all genes tested.

However, after a 5-day recovery period, expression of all but one (*Or88a*) of the genes was

recovered, including *Gr63a* (Figure S1C). This form of plasticity in gene expression is

unexpected and therefore mechanistically interesting to investigate.

Diacetyl and structurally related odorants act as HDAC inhibitors in vitro

Diacetyl is structurally related to the soluble metabolite secreted by the gut microbiome and liver, β-hydroxybutyrate, which is known to inhibit zinc-dependent HDACs (21). It is also structurally related to another known HDAC inhibitor Sodium butyrate. Diacetyl is present widely in nature as a pH-neutral fermentation product of microorganisms such as yeasts in over-ripe fruit, the favorite food for *Drosophila*, and also from lactic acid bacteria, and is found in many foods and beverages (5). In humans, diacetyl is also produced by microbes on the skin, in the oral cavity, and is found in breath (22). Interestingly, this volatile is known to traverse the cell membrane (23).

Soluble metabolites from the gut microbiome, such as short chain fatty acids, are known to affect histone modifying enzymes, gene expression, and physiology in various parts of the host (1, 24). It is presumed that these dissolved compounds absorbed in the blood circulate to various parts of the body. Microbes in our skin microbiome, mouth microbiome, and environment also emit a diverse repertoire of volatile metabolites that can traverse much greater distances away from the source (25). To examine if diacetyl can directly modulate HDACs like β-hydroxybutyrate, we performed in vitro acetylation assays with recombinant human HDACs. We found that indeed, diacetyl inhibited all four purified human Class I HDACs (HDAC1, 2, 3 and 8) and the one Class II HDAC (HDAC6) tested. The inhibition occurred in a dose-dependent manner. The IC_{50} values for HDAC1, 2, 3, 8 and 6 were 7.3 mM, 23.1 mM, 7.5 mM, 14.3 mM, and 24.5 mM respectively (Figure 1C). The levels of inhibition as observed from IC_{50} values for HDAC1 and HDAC3 were comparable to those of β-hydroxybutyrate whose IC_{50} values for HDAC1 and 3 were previously reported as 5.3 and 2.4 mM, respectively (21).

To test other odorants for their ability to inhibit HDACs, we identified a list of volatile

odorants structurally similar to diacetyl using the similarity search feature on PubChem (Figure

1D) and are volatile and tested them on four purified human HDACs, from Class I (HDAC1 and

HDAC3) and Class II (HDAC4 and HDAC6). Each compound was tested at 15 mM, the

concentration at which diacetyl inhibits HDAC1, 3, and 6 with more than 50% efficacy (Figure

1C, D). All compounds tested reduced the activity of at least one of the HDACs (Figure 1D). Ethyl

pyruvate, methyl pyruvate, 2,3-hexanedione, 2,3-heptandione, 2,3-pentandione inhibited HDAC1,

3 and 6 with more than 70% efficacy. Allyl butyrate also similarly inhibited those HDACs but

with lower efficacy. 1-acetoxyacetone, and 2,3-butanediol inhibited HDAC6 more than 70% but

inhibitory effects against HDAC1 and 4 were lower. Moreover, propyl formate strongly inhibited

HDAC 6 but did not inhibit HDAC1, 3, and 4. Taken together, these results indicate that microbial

volatiles and structurally related odorants can inhibit human HDACs, and that each compound has

specific HDAC inhibitory properties.

Microbial volatile increases histone acetylation in human cell nuclei

In order to directly examine histone acetylation at the cellular level, we used human

HEK293 cells, which offer a tractable system to prepare nuclear extracts, and picked the volatile

diacetyl to test in more detail. We exposed the cells to different doses of diacetyl for 2 or 6 hours

and monitored histone acetylation levels within nuclear extracts in Western Blot assays. Compared

to the mock treatment, 10 mM diacetyl significantly increased H3K9 acetylation levels within 2

hours of treatment, whereas the acetylation levels of H3K14 and H4K5 were not affected (Figure

2A). This specificity for an increase of the H3K9 mark is consistent with previous observations

for β-hydroxybutyrate. Preferential acetylation of H3K9 was also observed previously in HEK293

cells with treatment of the structurally related β-hydroxybutyrate (21). After 6 hours of treatment,

the H3K9 acetylation induced by 10 mM diacetyl was further increased (Figure 2B). The increase

in H3K9 acetylation with diacetyl treatment was dependent both on the duration of exposure and

concentration of the odorant.

Organisms are constantly exposed to volatiles commonly found in their food and

environment for prolonged periods of time. In order to test the effect of a lower concentration of

the volatiles, we selected a 5-day exposure time at a 100-fold lower concentration. When we

treated HEK293 cells with this lower dose of diacetyl (100 μM), H3K9 acetylation level increased

after 96 hours of exposure and reached significantly higher levels than control after 120 hours

(Figure 2C,D). These results demonstrate that even prolonged exposure to low levels of diacetyl

can greatly impact the epigenetic landscape inside the nucleus. More importantly, a 5-day exposure

was sufficient to alter the epigenetic state of nuclei at concentrations that are present in some food

sources. Taken together, these results demonstrate that diacetyl can act as an HDAC inhibitor, with

the potential of causing broad modulations of gene expression and histone acetylation in cells.

Transcriptional response to odor exposure is conserved in invertebrates and vertebrates

We next performed in vivo experiments to determine whether eukaryotic cells alter gene

expression in response to diacetyl exposure, as would be expected with HDAC-inhibition.

Epigenetic changes occur slowly over days, and to test its effects on gene expression we used two

model systems for eukaryotes: invertebrates (*Drosophila melanogaster)* and vertebrates *(Mus*

musculus). We placed *Drosophila* adult males in vials and housed these vials within a closed

container exposed to headspace from a 1% diacetyl solution (in paraffin oil) for 5 days, similar to

a previous long-term odor-exposure study (7) (Figure 3A). The transcriptome of the primary

olfactory organ, the antenna, was compared with that of the control group of age-matched flies

that were exposed to the solvent paraffin oil (PO). The antennal transcriptional profile of diacetyl-

exposed flies showed substantial changes in gene expression when compared to the solvent control

(Figure 3B). We identified 1234 differentially expressed genes (DEGs) (false discovery rate, FDR

< 0.05) in the antennal transcriptome of diacetyl-exposed flies compared to control animals. Of

these, 645 genes were significantly up-regulated ($\log_2$ fold-change > 1; red dots in Figure 3B) and

589 genes were significantly down-regulated ($\log_2$ fold-change < -1; blue dots in Figure 3B). A

broad range of genes was significantly altered, with several biological process GO terms

significantly enriched in the up-regulated gene list including "response to biotic stimulus" ($p <$

3×10^{-7}),"response to bacterium" ($p < 3 \times 10^{-7}$) and "defense response" ($p < 3 \times 10^{-7}$). In the down-

regulated gene set the GO term most enriched was "sensory perception of chemical stimulus" (p<

6×10^{-72}). The GO molecular functions that were most common in the upregulated sets included

enzymes like hydrolases and oxidoreductases, as well as nucleic acid binding genes (Figure 3C).

Down-regulated genes include receptors, hydrolases, and transporters (Figure 3C).

The first tissue that would have contact with airborne volatiles in terrestrial vertebrates are

the airways and lungs. We therefore performed transcriptome analyses on lung tissue of mice

exposed to diacetyl headspace at different doses for a period of 5 days, as was done in *Drosophila*

(Figure 3D). The dose of the volatile in the chamber was measured in parallel experiments by

tracking volatile loss and airflow and determined to range between a peak 1920 ppm at the end of

hour 2, tapering off to 28 ppm by hour 120 (Supplemental Figure S3B). Indeed, expression of a

substantial number of genes was modulated in the diacetyl-exposed lungs compared to the control.

The changes were dose-dependent and more pronounced in mice exposed to headspace from a 1%

diacetyl source compared to those exposed similarly to 0.1% (v/v) diacetyl source (Figure 3E,

Supplemental Figure S3C). Among these diverse sets of regulated genes in the lung tissue, a

significant overlap was found between 1% (v/v) and 0.1% (v/v) headspace for both up-regulated

genes ($p=3 \times 10^{-3}$) and down-regulated genes ($p=6 \times 10^{-3}$, Figure 4F), further supporting a dose-

dependent effect of diacetyl on gene expression in the mouse lung.

The GO enrichment analysis of the lung transcriptome from mice exposed to diacetyl

revealed several interesting sets of genes that were significantly altered (Figure 3G,H). Among

these genes is the SARS-CoV-2 entry receptor ortholog, ACE2. The level of this gene in the lungs

of diacetyl exposed mice showed a reduction in expression levels ($\log_2$ fold-change= - 0.62, FDR=

2.02×10^{-5}). The level of ACE2 in the nasal tissue, lungs and other organs play a critical role in

infection by the SARS-CoV-2 virus, suggesting that epigenetic mechanisms may contribute to

expression. While we do not know whether expression of ACE2 in humans is regulated similarly,

these results highlight the importance of understanding volatile-induced modulation of gene

expression in the lungs and other tissues.

HDAC family members are highly conserved across eukaryotes, spanning both animal and

insects. Volatile microbial metabolites have been present throughout the evolution of animals,

insects, and plants, and have potential as signals for multi-domain communication because they

can travel wide ranges; plants detect root infections and time their immune response using volatiles

(26), and microbial volatiles elicit olfactory behaviors in insects and nematodes. Our results

indicate that a volatile compound, capable of directly inhibiting HDACs and altering acetylation

levels, can cause significant changes in gene expression in a variety of organisms following

exposure from a distance, through the air.

DEGs overlap those for known HDAC inhibitors and upregulation is partially reversible

One of the hallmarks of epigenetic regulators which makes them attractive candidates for

therapeutics is plasticity. Upon removal of the inhibitors, we expect that some of the changes in

gene expression are reversible. To test for reversibility, we performed a recovery experiment

following diacetyl exposure using the *Drosophila* model. We maintained 5-day diacetyl-exposed

flies in clean air for 5 additional days (Figure 4A). In parallel, we performed age-matched mock

experiments with paraffin oil solvent exposure alone. A large number of genes were down-

regulated following the recovery in comparison to the 10-day-old flies in the mock condition

(Figure 4B). Interestingly, there was a significant overlap of these down-regulated genes with the

set that was up-regulated in the diacetyl treatment (Figure 4C). These results suggest that the

effects of HDAC inhibitory odorant exposure are not permanent but dynamic, and removal of the

odorant leads to subsequent changes in gene expression of the up-regulated set.

Drugs that inhibit HDACs are in various stages of approval as treatments for several

diseases. The approved drugs include sodium butyrate and valproic acid. To compare the gene-

regulatory effects of these drugs to diacetyl, we performed RNA-seq after raising the flies on food

containing sodium butyrate (SB) or valproic acid (VA) (27) and compared gene expression in the

antennae of flies raised on untreated food for 5 days. We next compared the up-regulated gene

profiles following each treatment to the one induced by exposure to diacetyl. As expected, feeding

SB and VA induced significant changes in expression levels of several genes (Figure 4D,E).

Interestingly we found that 113 of diacetyl up-regulated genes were also up-regulated in either SB,

VA or both treatment conditions (Figure 4F). Pairwise statistical analysis of each gene set revealed

a significant overlap of diacetyl-induced genes with SB-induced genes ($p=6\times10^{-11}$) and with VA-

induced genes ($p=2\times10^{-65}$) (Figure 4F). There was, as expected, also a significant overlap between

SB- and VA- induced genes ($p=1\times10^{-52}$) (Figure 4F). This highly significant overlap among up-

regulated genes lends further support to our model that diacetyl vapors act as an HDAC inhibitor

in vivo. As expected, each of the 3 treatments also modulated a substantial number of unique genes

(Figure 4G, H), suggesting that differences in oral vs. vapor delivery, molecular structure and

inhibition profile across the repertoire of HDACs may contribute to differences in gene regulation.

These results suggest that some volatile odorants, or the microbes that emit them, may impart

therapeutic effects like other HDAC inhibitors from a distance, through the air.

257 Volatile Diacetyl alters gene expression in the mouse brain

One of the categories of challenging diseases that HDAC inhibitors are being tested for

drugs are neurodegenerative diseases (28) where volatile odorants may provide a new class of

inhaled therapeutics. A major design challenge for nervous system drugs is the ability to cross the

blood-brain barrier. Due to the volatility and small size of volatile odorants, there is a possibility

that they could diffuse through the intranasal route to the brain directly (29). We cannot tell *a*

priori whether odorants like diacetyl can travel across the nasal membrane to the brain. In order to

test whether cells in the brain of mice respond when presented with diacetyl vapors from a distance,

we used gene expression as the final readout to measure it. We performed RNA-seq experiments

on mice exposed only to aroma of diacetyl in the air for 5 days as before. Littermate controls were

exposed in a similar manner to the solvent (PO) in the air. Interestingly, several genes were

differentially expressed upon exposure to headspace from 0.1% (v/v) diacetyl passaged through

the air in the holding cage where it is further diluted (49 up-regulated, 32 down regulated, $|\log_2$

fold-change| >1, FDR<0.05) or to headspace from 1% (v/v) diacetyl diluted in the holding cage

(748 up-regulated, 1031 down regulated, $|\log_2$ fold-change| >1, FDR <0.05) (Figure 5B,C). GO

analysis of the regulated genes in the exposed mouse brain transcriptome revealed several

interesting sets of genes were significantly altered in each set (Figure 5E). These results indicate

that a volatile odorant like diacetyl can reach the brain in mammals, and presumably alter gene

expression in neuronal cells. Although the overall DEG sets were different across the lungs and

brain, there was a statistically significant overlap, particularly in the 1% exposure, as would be

expected from using the same mechanism of action. There was a highly significant overlap

($p=2 \times 10^{-101}$) between genes upregulated in the brain compared with the lungs (248 of 748), as well

as for downregulated genes ($p=3 \times 10^{-163}$; 477 of 1031 genes) (Figure 5D).

Diacetyl prevents proliferation of neuroblastoma cells

Among the significantly downregulated genes was *MYCN*, which is known to show

increased expression and plays a central role in many types of cancers, suggesting that epigenetic

regulation may affect tumor cells. Broad-spectrum HDAC inhibitors have been approved by the

FDA for treatment of a variety of cancers. To test the exciting possibility that a volatile HDAC

inhibitor, which affects gene expression in the brain, would have an effect against cancers as well,

we performed a cell-based assay. We picked neuroblastoma as a candidate since it has a strong

dependence on *MYCN* and develop from fetal adrenal neuroblasts (30). Several genetic and

"omics" studies have identified genes that are upregulated in neuroblastomas, some of which are

considered to be key causative factors (31). We evaluated the DEGs from the diacetyl-exposed

mouse brain transcriptome for several of these key genes. Three of the six key genes identified

across multiple studies were significantly altered, two of which were significantly downregulated,

MYCN (P=2.06E-13, FDR=1.76E-12) and *Alk* (P=3.3E-5, FDR=9.48E-5) (Figure 5F)**.** An

unbiased computational study evaluated omics datasets and identified 4 genes that were the best

predictors of clinical outcome (32), all of which are significantly different in the brains of the

diacetyl exposed mice, strikingly 3 of which were significantly downregulated, *NCAN (p=1.33E-*

16, FDR=1.59E-15), STK33 (p=4.6E-7, FDR=1.79E-6) and *ERCC6L2 (p=4.57E-5,*

FDR=0.00012) (Figure 5F). These findings are extremely provocative, given that the

downregulated genes, particularly MYCN play a role in neuroblastomas and in several other

cancers.

We tested diacetyl for the ability to inhibit proliferation of a SH-SY5Y

neuroblastoma cell line from a metastatic bone tumor. A significant reduction in cell count was

seen in diacetyl treatments in the SH-SY5Y cancer cell lines (Figure 5G). To validate the initial

result, we retested diacetyl with at two concentrations across 3 different cancer cell lines. Diacetyl

showed a significant reduction in cell numbers for the neuroblastoma cell line specifically, and not

any other cell lines, relative to the control (Figure 5G). These results demonstrate that the effect is

specific to the SH-SY5Y cancer cell line, and unlikely to be caused by a non-specific general toxic

effect. It also suggests that the potential epigenetic regulation of cancer promoting genes may be

caused by exposure to volatiles like diacetyl in the environment.

Testing therapeutic effect in neuroblastoma and Huntington's neurodegeneration

Since diacetyl can affect changes in a mammalian brain by crossing the blood brain barrier,

we wanted to test if it can impart therapeutic effects in disorders of the brain like neurodegeneration

for which HDAC inhibitors have been proposed as therapeutics. In order to test diacetyl in the

context of neurodegenerative disease, we used the well-established *Drosophila* model of

Huntington's disease, which has previously been used to demonstrate the efficacy of sodium

butyrate in slowing neurodegeneration (27, 33). The human Huntingtin protein with expanded

poly-Q repeats is expressed in the neurons of the compound eye in *Drosophila*, causing progressive

degeneration of the seven visible photoreceptor rhabdomere cells in each ommatidium (34). We

selected this model in particular since polyglutamine disorders are well suited for targeting by

inhibitors of HDAC1 and HDAC3 (35) such as diacetyl (Figure 1B). Moreover, previous studies

have shown that orally administered HDAC inhibitors such as sodium butyrate and SAHA can

significantly reduce photoreceptor degeneration in this model (27). When the transgenic flies

expressing two copies of the human Huntingtin with poly-Q repeats (HTTQ120) under control of

the eye-specific GMR promoter were raised at 18^{o}C, the number of rhabdomeres in each

ommatidium was similar to that normally observed in wild-type flies immediately post-eclosion

(day 1, Figures 6A-C). When these HTTQ120 flies were moved to 25^{o} C following eclosion

(Figure 6A), they showed dramatic degeneration of rhabdomeres over a period of 10 days (Figures

6B, D-G). The mean number of rhabdomeres was reduced from 7 to ~1 by day 10. The dose of the

volatile in the chamber was measured in parallel experiments by tracking volatile loss and airflow

(0.5 L/min) and determined to range from 260-95 ppm, an average of 125 ppm at the end of hour

1, tapering off to 0 ppm by hour 24. The odorant was replaced each day, leading to a dosage profile

that would seem like a once-a-day treatment regimen (Supplemental Figure S3A).

Remarkably, when the Huntingtin (HTTQ120)-expressing flies were exposed immediately

after eclosion to volatile headspace of 1% diacetyl (Figure 6A), they showed a substantial (~50%)

reduction of rhabdomere loss (Figures 6B, D-G). The majority of ommatidia retained 6-7

rhabdomeres at day 5 (Figures 6D and F). Even after 10 days, the majority of the ommatidia still

had 2-3 rhabdomeres left in the odor-exposed flies, while the solvent controls had only ~1-2

rhabdomeres (Figures 6E and G). These results demonstrate that exposure to volatile diacetyl in

air slows down the photoreceptor degeneration caused in these Huntington's disease model flies.

Thus, odorants like diacetyl may have potential as a prophylactic against this neurodegenerative

disorder. Therefore, the volatile odorant can both significantly alter gene expression from a

distance as well as cause a phenotypic change in the fly with a human disease gene.

Discussion

In this study, we discovered that eukaryotic cells could alter gene expression in response to an odorant, in a manner independent of traditional neuronal activity-induced pathways. Members of a conserved family of HDACs detected the concentration of diacetyl differentially. Diacetyl exposure increased levels of H3K9 acetylation in the nucleus in a dose-dependent manner and led to changes in gene expression. The odorant diacetyl occurs naturally and is generated from the metabolism of a variety of food components, including triglycerides, sugars and amino acids (5). Moreover, the chemical is produced by the activity of microorganisms such as yeasts and lactic bacteria during fermentation in many foods and beverages (5). Although beyond the scope of this study, the discovery that cells can alter gene expression upon prolonged exposure to a common naturally occurring odorant raises important questions about the potential physiological consequences of the expression changes. Given our repeated exposure to particular flavors and fragrances, the findings outlined here highlight a new consideration for evaluating the safety of certain volatile chemicals that can cross the cell membrane.

In mammals, upregulation of circulating β-hydroxybutyrate during fasting or calorie restriction induces changes in the expression of a set of genes (21). The acetylation mark specificity and IC_{50} of β-hydroxybutyrate is similar to that of diacetyl. Any beneficial physiological effects of odor-detection at lower odor concentrations would also be countered by studies of potential risks at higher concentrations. In fact, the deleterious effect of exposure to high levels to diacetyl causing bronchiolitis obliterans, or "popcorn lung", and its toxicity in cultured cells is already known (36). Even so it is present in several foods we eat and is on the GRAS (Generally Regarded as Safe) list of Flavor and Extract Manufacturers Association (FEMA) for use as a flavoring ingredient at low concentrations. While the specific mechanism of "popcorn lung" is unknown, it

has previously been proposed that carbonyl groups of diacetyl can react to modify amino acid side

chains on proteins or generate reactive dicarbonyl and reactive oxygen species, leading to

excessive cytokine production and inflammation (37). Our results raise the possibility that the

molecular mechanisms underlying this disorder could also be partially attributed to the unusually

high HDAC-mediated genetic response to diacetyl by cells in the lungs. Several studies using

rodent animal models have shown toxicity of high levels of diacetyl in the respiratory tracts

including the nose and lungs (38, 39). The toxic effect of higher dosages has been observed for

many types of HDAC inhibitors. For example, in one *Drosophila* study, feeding of a high dosage

of 4-phenylbutyrate reduces survival rate, while a lower dosage extends longevity (40). Another

concern is that one of the highest known levels of diacetyl exposure to humans is through smoking

cigarettes, where exposure from even a single cigarette (250-361ppm) is high (41, 42): Salem (128

µg/cigarette), Camel (128.1 µg/cigarette), Carlton (45.6 µg/cigarette) (42). Different types of diets

and lifestyle choices have been known to have effects on health and well-being due to differences

in balance of different health promoting ingredients, yet little is known about role of such flavors,

fragrances and metabolites acting on epigenetic marks and altering gene expression. Our findings

raise the possibility that diacetyl and structurally related odors in the environment may affect gene

expression directly through absorption/transport into these cells. In mammals, the epithelium-lined

lungs and nasal/oral pathway is directly exposed to odorants presenting an opportunity to affect

gene expression and physiology.

Eukaryotes primarily detect volatile odorants in the environment using olfactory neurons

that express a variety of transmembrane receptors, a family of genes that have independently

evolved multiple times in different phyla. Examples of these unrelated genes include: the

ionotropic 7-transmembrane (TM) odorant receptor *(*Or*)* family, which is insect-specific; 3-TM

ionotropic receptors (IRs), which are present across most arthropods; the nematode-divergent 7-

TM GPCRs belonging to the *str, sra, srg, srw, srz, srbc, srsx* and *srr* families; the mammalian 7-

TM GPCR olfactory receptor (OR) family; and the trace amine-associated receptor (TAAR) family

(43-47). The activation of these receptors induces neuronal action potentials, and this information

is conveyed to higher brain centers where olfactory perception is generated (48). This second

response pathway which we have discovered is much slower and has an enzymatic mechanism

which draws parallels with the multiple pathways in place to detect light. In the specialized cells

of the eye, detection of light occurs via rhodopsins (7-transmembrane GPCRs) and leads to

neuronal activity and behavioral responses. However, other cells, including in plants, are also able

to respond to changes in light intensity of certain wavelengths using the ancient, conserved

cryptochrome proteins that are related to photolyases (49). This suggests that vital sensory

modalities can be detected via multiple pathways. Our observations raise the question whether

HDACs represent a conserved detection mechanism for odorants like diacetyl that link odor-

detection to a specific gene expression response. HDAC proteins, however, are an ancient family

of genes that predate even histones themselves (50, 51). It is conceivable that odor detection

mechanisms that emerged in ancient forms may not have involved specialized transmembrane

receptors, or neurons, or resulted in rapid behavioral movements.

Our data suggests that an HDAC inhibitory odorant also affects gene expression in the lungs

and brain of mice, including significantly decreasing the expression level of the ACE2 receptor,

whose ortholog in humans is the entry receptor for the SARS-CoV-2 virus that causes COVID-19.

The applied impact for the future could be far-reaching, as HDACs are known to affect a wide

variety of pathways in eukaryotes, and inhibitors are candidates for novel inhaled therapeutics as

treatment of many conditions such as COVID-19, cancers, neurodegeneration and a variety of

infectious diseases (52-60).

Our studies also show that one of the volatiles is protective against the neurodegenerative

effects of Huntington's disease model and neuroblastoma. Exposure to diacetyl volatiles in the fly

model of Huntington's disease reduces cell degeneration, as has been previously observed with

orally administered HDAC inhibitors like sodium butyrate and SAHA in this genetic model (27).

Previous studies indicate that the inhibition of HDACs counter the acetyltransferase inhibitory

activity of the polyglutamine-domain of the Htt protein which binds to p300, P/CAF and CBP (27).

This newly discovered activity in this class of volatile chemicals will also allow us to

understand how their presence in our environment, diet and microbiome shapes the epigenetic

landscape in the lungs and affects inflammation and immune responses. Our approach to

investigate consequences in model systems allows for a foundation for future research and allow

identification of key molecular biomarkers that can be associated with observation of epigenetic

effects, toxic or not, in individuals. The public health impact of these findings can be far-reaching,

both from the perspective of toxicity as well as therapeutics. Investigating the health safety

concerns of exposure to such odorants at high doses or prolonged periods is essential to understand

potential for damage. Conversely, at non-toxic concentrations HDAC inhibitors can be used for

treatment of many debilitating conditions in the lungs such as COPD, inflammation, and COVID-

19 infection (61).

Taken together, our discovery that cells can alter gene expression in response to an odor that

acts as an HDAC inhibitor and reprogram gene expression, promises the pursuit of new types of

odor-based therapeutics already approved for human consumption as prophylactics for a myriad

of diseases. Simultaneously, they raise the concern of a new type of environmental agent which

can reprogram gene expression and have widespread effects that are yet unknown.

METHODS

Drosophila Stocks and Manipulations

438   Fly stocks were maintained on conventional fly food under a 12-hr. light:12 hr. dark cycle at 18°C

439   or 25°C. The fly strain of *w1118* backcrossed 5 times to *Canton-S* (*wCS*) was used in all the

440   *Drosophila* transcriptome experiments. P{GMR-HTT.Q120}2.4 (Bloomington # 8533) were used

441   for neurodegeneration experiments. For imaging *Gr63a-Gal4; UAS-mcd8:GFP, Gr21a-Gal4;*

442   *UAS-mcd8:GFP* stocks were obtained as a kind gift from the Carlson lab, Yale.

Odor exposure experiments for imaging

Newly eclosed male wCS flies were sorted in groups of 30 into vials with standard cornmeal medium.

Vials were closed with two overlapping 3''x3''polypropylene mesh squares and tied with cotton twine.

Vials were used within 6 hours. Four vials were placed in a 1000ml Nalgene straight sided jar with a

10ml glass beaker with 10ml of diacetyl 10^{-2} in paraffin oil. Jars were closed for exposure periods

ranging from 2-6 days. In recovery experiments, vials were removed from the jars and were placed in a

25°C incubator for the remainder of the recovery period. Flies were anesthetized on ice, sacrificed in

ethanol, and then immediately put into 1X phosphate buffered saline with 0.3% triton-x (PTX). Brains or

antenna were put in 4% paraformaldehyde in PTX and incubated for 30min while rotating at 25°C.

Samples were washed 5 times in PTX, and blocked in 5% natural goat serum in PTX for 1-hr while

rotating at 25°C. Samples were then incubated in primary antibody with 5% goat serum in PTX, mouse-

nc82 (1:20) (Development Studies Hybridoma Bank, University of Iowa) and rabbit-antiGFP (1:150)

(Invitrogen) in PTX for 48 hours in 4°C while rotating. After washing 5 times in PTX samples were

incubated in secondary antibody with 5% goat serum in PTX, rabbit-anti-Alexa488 (1:400) (Invitrogen)

and mouse-antiAlexa568 (1:400) (Invitrogen) for 48 hours in 4°C. Samples are washed 5 times in PTX

and stored in 70% glycerol in PTX. Images were taken using a Ziess 510 laser scanning confocal

microscope, and image analysis was done using Image J software.

QRT-PCR

Drosophila heads (with antenna) were collected on liquid nitrogen and stored at -80°C until processing.

300 heads were collected for each sample, and ground on liquid nitrogen. mRNA was extracted using a

PolyATract kit (Promega) and was reverse transcribed using Superscript III (Invitrogen) according to the

manufacturer's protocol. The quality of the cDNA was assessed on a 1.5% agarose gel. cDNA was

added to individual reactions of SYBR green master mix (BioRad) and run on a BioRad-MyQ

thermocycler. The program began with a single cycle for 3-min at 95°C, followed by 40 cycles of 15-sec

at 95°C, 30-sec at 58°C and 30-sec at 72°C. Afterward, the PCR products were heated to 95°C for 1-min

and cooled to 55°C for 1-min, to measure the dissociation curves. The efficiency of each primer set was

first validated by constructing a standard curve and three 10x serial dilutions of first strand cDNA. For

each serial dilution, the CT value was plotted against the log (template dilution) and the slope and r^2 value

of each regression line calculated. Expression of each *Or* or *Gr* gene was assessed in triplicate.

Dissociation curves were used to assess the purity of the PCR reactions. *Or* and *Gr* transcript levels were

normalized by using the transcript levels of *ribosomal protein 49*. Expression levels were calculated using

the Pfaffl equation.

In general, primers were designed by using coding sequences close to the 3' end of the gene, and where

possible, primers spanned an intron.

Primers:

Gr21a F: CGATCGTCTTTCCGAATCTC

Gr21a R: GGCTCAGATCCACCCATAGA

Gr63a F: AAATGAACTCCGCCTCCTTT

Gr63a R: CGCAATTTCAGAGGCAAACT

RP49 F: CTGCCCACCGGATTCAAG

RP49 R: GTTTCATGCGGCGAGATCG

Or47a F: ATCACAGGCCACATTGAACA

Or47a R: TCCCCGCAGTAGCAGTAGAT

Or88a F: TTAAAGTGGCCTTCCTGGTG

Or88a R: ATGCGGCAATAAAGTTCCAC

Or83b F: TTCTTGGCATTCGCTTTTCT

Or83b R: TCCCTGGATTTGTTTGCTTC

Odor Exposure Protocol for Transcriptome Analysis

Flies were exposed to diacetyl (B85307, Sigma-Aldrich, St. Louis, MO) by placing them in vials

in a cylindrical closed container (112 mm diameter x 151 mm height) along with an odor-

containing glass vial. The odorant was dissolved in 5 mL paraffin oil at 1% dilution. For a given

exposure protocol, two groups of flies were prepared: those exposed to 1% diacetyl headspace and

those exposed to paraffin oil headspace alone (control flies). Adult male flies aged 1 d were

transferred to fly vials containing fresh medium and put into the container with the odor vial. At

the end of the fifth day of exposure, flies were collected, and their antennae were dissected for

RNA extraction. All treatments and experiments were performed at room temperature. For the

recovery experiment, flies were transferred to a container with a glass vial of paraffin oil after 5

days of diacetyl exposure. At the end of the fifth day of recovery, flies were collected, and their

antennae were dissected for total RNA extraction. The second and third antennal segments from

40-60 male flies after treatment were carefully hand-dissected from the head and collected in 1.5

ml microfuge tubes kept cold in liquid nitrogen. Antennae were mechanically crushed with

disposable RNAse-free plastic pestles, and total RNA was isolated using a Trizol-based protocol.

cDNA libraries were prepared from total RNA using the Illumina TruSeq RNA Sample

Preparation Kit (v2) and 50 bps single- and paired-end sequencing was done using the

HighSeq2000. Two biological replicates were sequenced for each condition, with an average of

27 million reads / replicate, and with an average of 84% mapped.

Two-month-old C57BL/6 male mice (2-3 for each condition in a single cage) were

continually exposed to air flowing over headspace of paraffin oil (solvent control), 0.1%, or 1%

diacetyl over a period of 5 days, then euthanized for collecting the lung and brain tissues and

processing for mRNA isolation. All protocols for animal use and euthanasia were approved by the

Institutional Animal Care and Use Committee (#20150028) and were in accordance with the

provisions established by the Animal Welfare Act and the Public Health Services (PHS) Policy.

In the transcriptome analysis, two replicates were performed for each condition, with an average

of 123,687,411 reads / replicate, with an average of 88% mapped. Multiplexed libraries were

made from total RNA input using the Illumina TruSeq RNA sample preparation kit (v2) and 50

bps single-end sequencing was done using the NextSeq500.

Calculation ppm in air based on weight loss of odorant over time

The odor solution in PO in each setup (Drosophila and mouse) was weighed using a sensitive

balance. PO is non-volatile and therefore weight loss is attributable to diacetyl volatilization.

Volatile Headspace exposure (ppm) is calculated through odor cartridge weight change from pure

odor compound volatizing out (Inamdar et al., 2012).

ppm=(10^6 w/MW)/(v/V$_m$)

Where ppm is the concentration of tested VOC in parts per million (v/v); w is the weight of the

tested chemical in grams; MW is the molecular weight of the tested VOC (grams/mole); v is the

total volume of the respective exposure chambers; and Vm is the molecular volume under the

tested condition.

•V$_m$=24.45(760/P)((t+273.15)/298.15)

Where 24.45 = gram molecular volume, under standard Lab conditions of 760 mm Hg, 25°C; P =

ambient pressure, in mm Hg; and t = ambient temperature, in °C. Odor cartridges, consisting of

varying odor volume/volume dilutions, were weighed periodically under standard ambient

temperature and pressure. Under these conditions, the pure volatile odor compounds volatize out

of the odor cartridge and the evaporation rates were used for our exposure calculations. Based on

Ideal Gas Law and experimental conditions, ppm is calculated based on volatized odor compound

and experimental chamber volume based on airflow.

HDAC inhibitor Treatment Protocol for Transcriptome Analysis

Sodium butyrate (B5887, Sigma-Aldrich) or valproic acid (P4543, Sigma-Aldrich) were dissolved

in normal fly food medium at the final concentration of 10 mM. Three groups of flies were

prepared: those treated with one of the HDAC inhibitors and those without HDAC inhibitor

treatment (control flies). Adult flies aged 1 d were transferred to fly vials containing medium with

or without a HDAC inhibitor. At the end of the fifth day of treatment, flies were collected, and

their antennae were dissected for RNA extraction. All treatments and experiments were performed

at room temperature. Two biological replicates with 60 flies/replicate were performed for each

condition, with an average of 23 million reads / replicate, and with an average of 92% mapped.

Multiplexed libraries were made from total RNA input using the Illumina TruSeq RNA sample

preparation kit (v2) and 50 bps paired-end sequencing was done using the HighSeq2000.

Bioinformatic analysis of RNA-seq experiments

Reads aligned to the latest release of each of the genomes used (dm6 for the *Drosophila* genome,

GRCm38 for the *Mus Musculus* genome) and quantified with kallisto (version 0.43.1) (62) Only

libraries for which we obtained >75 % alignment were used for downstream analysis. Transcript

counts were summarized to gene-level using tximport package (version 1.4.0)(63). For any

instances of detected batch effects, we removed unwanted variation using RuvR in the RuvSeq

package (version 1.10.0)(64). Differentially expressed gene (DEG) analysis was performed with

the edgeR package (version 3.18.1)(65) using low count filtering (cpm >0.5) and TMM

normalization. Protein classification analysis was performed with PANTHER (version 13.1)(66).

All significance analyses of gene overlap were done using the GeneOverlap package in R package

(version 1.14.0). GO enrichment analysis was performed using clusterProfiler (version 3.6.0)(67).

HDAC Activity Assays

HDAC activity of class I HDACs (HDAC1, 2, 3 and 8) was measured with the fluorometric HDAC

Activity Assay kit: HDAC1 (10011563, Cayman Chemical, Ann Arbor, MI), HDAC2, HDAC 3,

and HDAC 8 (50062, 50073 and 50068, BPS Bioscience, San Diego, CA), according to the

manufacturer's instructions. HDAC activity of class II HDACs (HDAC4 and HDAC6) were

measured with the fluorometric HDAC Activity Assay kit: HDAC4, HDAC 4 (50064 and 50076,

BPS Bioscience, San Diego, CA), according to the manufacturer's instructions. HDAC activity in

these 96-well format kits are directly correlated with fluorescence produced by the enzyme and

substrate interaction. Each test compound was tested in duplicates, minimum. Due to interference,

concentrations of >30 mM were not used in the IC50 calculation. To account for any potential

absorbance or fluorescence contributed by the test chemicals interacting with the assay substrate,

all baseline background measurements of test compound + substrate control wells (everything, but

no HDAC enzyme) were performed and subtracted against the respective test compound wells. It

is fair to note that no significant changes in background fluorescence were seen with tested

compounds in comparison to control background fluorescence. HDAC activity was calculated

based on averaged maximum fluorescence from negative control wells and the averaged

fluorescence in test compound wells.

$$\frac{Fluorescence_{Negative\ control} - Fluorescence_{HDACi}}{Fluorescence_{Negative\ control}} x100\% = Remaining\ HDAC\ activity\ \%$$

HDAC % inhibition was directly calculated from obtained HDAC activity %.

Cell Culture and Treatment

Human embryonic kidney 293 (HEK293) cells were grown in 100 mm cell culture dishes with

Dulbecco's modified Eagle's medium (DMEM) (10-013, Corning, Manassas, VA), supplemented

with 10% fetal bovine serum (FBS) (26140-079, Gibco, Carlsbad, CA) at 37°C with 5% CO_2.

Cells that were ~80% confluent were treated with freshly prepared medium supplemented with

diacetyl at concentrations indicated. The cells for mock controls were handled in the same manner

without adding diacetyl to the medium. To prevent diffusion of diacetyl odor from the treatment

dishes to the ones of mock control, the cell culture dishes in different conditions were cultured in

separate CO_2 incubators.

Cell proliferation assays were performed to assess effects of compounds against cancer cell lines.

Cells were maintained in a 37°C incubator with 5% CO_2 throughout the whole experiment.

Compound stocks are dissolved in 10% DMSO, and final concentrations made fresh in complete

media as needed. 10% DMSO solvent is used as control in complete media. Cells were seeded

onto 12-well plates and allowed 12-24 hours until treatment to allow cells to adhere to wells.

This is followed by minimum 5-day treatment of compounds, changing the media with treatment

every 48 hours. Cell lines A549, PC3, and SK-MEL-5 will be seeded at 8000 cells/well, which

was determined to provide 80-100% confluency after 6-day growth. SH-SY5Y was seeded at

8000 cells/well and 16,000 cells/well due to their known slower division rate and given at least

10 days for sufficient growth. After allotted growth times, cell counts were performed using the

Countess^{TM} automated cell counter. All assays had a minimum of three replicates for all

conditions and analyzed through GraphPad Prism One-way ANOVA for significance.

Odor Exposure Protocol for Huntington's Disease Model Flies

Flies were exposed to diacetyl in a cylindrical container (112 mm diameter x 151 mm height).

Each container was tightly closed but had 2 holes, one of which connected to an air suction port,

and the other to a vial containing either of 5 mL paraffin oil or 5 mL 1% diacetyl in paraffin oil. A

gentle suction was applied to pull the headspace from the odor or paraffin oil vials into the

cylindrical structure. pGMR-HTTQ120 flies were maintained at 18°C. Adult flies aged 1 d were

transferred to fly vials containing fresh medium and put into the odor-filled container at room

temperature. Paraffin oil and 1% diacetyl solution were prepared and replaced every day. At the

end of the fifth day of exposure, half of the flies were collected and subjected to pseudopupil

analysis as before (27). The remaining flies were transferred to fresh medium and exposed to the

odors for an additional 5 days. All treatments and experiments were performed at room

temperature. The number of rhabdomeres in each ommatidium was counted using the pseudopupil

analysis, and statistical significance was determined by unpaired t test (two-tailed) against PO at

each time point.

Preparation of Nuclear Extracts from HEK293 Cells

Nuclear extracts of HEK293 cells were prepared according to a protocol described previously (68),

with minor modifications. In brief, HEK293 cells were washed twice with cold phosphate-buffered

saline (PBS) and lysed with hypotonic buffer (10 mM HEPES-KOH [pH 7.9], 1.5 mM $MgCl_2$, 10

mM KCl, protease inhibitor cocktail (04693159001, Roche, Indianapolis, IN), 1 mM DTT, 1 mM

TSA). Following a brief centrifugation, the pellet was resuspended in hypertonic buffer (20 mM

HEPES-KOH [pH 7.9], 25% glycerol, 420 mM NaCl 1.5 mM $MgCl_2$, 0.2 mM EDTA, protease

inhibitor cocktail, 1 mM DTT, 1 mM TSA). The supernatant was recovered as nuclear extract.

Western Blot Analysis

Proteins in the nuclear extracts (60 µg protein) were separated by SDS–PAGE gels (456-1043,

Bio-Rad, Hercules, CA), transferred onto PVDF membranes (162-0174, Bio-Rad), and incubated

with anti-histone antibodies: acetylated H3K9 (1/2000: ab4441, abcam, Cambridge, MA),

acetylated H3K14 (1/5000: 06-911, EMD Millipore, Billerica, MA), acetylated H4K5 (1/2000:

07-327, EMD Millipore). Bound antibody was detected by horseradish peroxidase-conjugated

anti-rabbit secondary antibody (1/20000: 1705046, Bio-Rad) and developed using Clarity™

Western ECL Substrate (1705060, Bio-Rad). Signals were detected and captured by

ImageQuant™ LAS 4000 mini (GE healthcare, Pittsburgh, PA), and band intensities were

quantified with ImageJ software. H3K9 acetylation intensity in individual lanes was reported

relative to the normalized Mock treatment and calculated using this formula: Relative H3K9ace

intensity for each timepoint = (H3K9ace / PCNA) / averaged (Mock H3K9ace / Mock PCNA).

Statistical significance was determined by unpaired *t* test (two-tailed) against mock at each time

point.

Data availability

All transcriptome data sets in this publication have been deposited in the GEO repository

with accession number GSE116502. All other data is presented in the figures, tables, and

supplementary materials.

Author contributions: S. H.Y. performed the biochemical assays, cell culture assays,

neurodegeneration assays and wrote the first draft. R.N.F performed biochemical assays and the

cancer cell line exposure experiments and helped write the paper. C.A.S performed the

bioinformatics and transcriptome analysis, parts of the biochemical assays, and helped write the

paper. S.P. performed some of the transcriptome analysis. S.T.C performed the imaging studies

and RT-PCR. C.P. performed some biochemical analyses. M.G.N. helped plan and assisted with

the mouse exposure experiments. A.R. conceptualized the study, planned, and supervised the

experiments and wrote the final draft of the paper.

Competing interests: A.R. is founder of startups: Sensorygen working on computational

neurobiology of olfaction and taste, and Remote Epigenetics working on small molecule

enhancement of agricultural production. A.R., S.T.C., S.H.Y., R.N-F., are listed as inventors on

patents filed by University of California.

Acknowledgments: We would like to thank Dr. Larry Marsh for advice about the fly eye assay

and HDAC inhibitors, Joel Kowalewski for help with critical comments, Dr. Thomas Girke for

guidance in bioinformatics analysis, Drs. Anupama Dahanukar, Naoki Yamanaka, Chih-cheng

Tsai, Karine LeRoch and Adler R. Dillman for sharing equipment and reagents. This work was

partially supported by funds provided by the National Institutes of Health, NIAID (R01AI153195

to M.G.N and R01AI087785 to A.R), and HATCH funds from the Agricultural Experimentation

Station UC Riverside to A.R.

References

1.   McFall-Ngai M*, et al.* (2013) Animals in a bacterial world, a new imperative for the life sciences.
Proceedings of the National Academy of Sciences 110(9):3229-3236 %U
http://www.pnas.org/content/3110/3229/3229.
2.   Dorrestein PC, Mazmanian SK, & Knight R (2014) Finding the missing links among metabolites,
microbes, and the host. *Immunity* 40(6):824-832.
3.   Wikoff WR*, et al.* (2009) Metabolomics analysis reveals large effects of gut microflora on
mammalian blood metabolites. *Proceedings of the National Academy of Sciences* 106(10):3698-
3703 %U http://www.pnas.org/content/3106/3610/3698.
4.   Rooks MG & Garrett WS (2016) Gut microbiota, metabolites and host immunity. *Nat Rev*
Immunol 16(6):341-352.
5.   Shibamoto T (2014) Diacetyl: occurrence, analysis, and toxicity. *J Agric Food Chem*
62(18):4048-4053.
6.   Devaud JM, Acebes A, & Ferrus A (2001) Odor exposure causes central adaptation and
morphological changes in selected olfactory glomeruli in Drosophila. *Journal of Neuroscience*
21(16):6274-6282.
7.   Sachse S*, et al.* (2007) Activity-dependent plasticity in an olfactory circuit. *Neuron* 56(5):838-
850.
8.   Devaud JM, Acebes A, & Ferrus A (2001) Odor exposure causes central adaptation and
morphological changes in selected olfactory glomeruli in Drosophila. *J Neurosci* 21(16):6274-
6282.
9.   Martineau B, Henick-Kling, T., Acree, T. (1995) Reassessment of the Influence of Malolactic
Fermentation on the Concentration of Diacetyl in Wines. *Am J Enol Vitic* 46(3):385-388.
10.   Nykanen L, Nykanen, I. (1991) Distilled beverages. *Volatile Compounds in Foods and*
Beverages, ed Maarse H (Dekker Inc, New York), pp 548-580.
11.   Hughes PS, Baxter, E.D. (2001) Flavour Determinants of Beer Quality. *Beer: Quality, Safety and*
Nutritional Aspects, (The Royal Society od Chemistry, Camdridge, UK), pp 40-73.
12.   Turner SL & Ray A (2009) Modification of CO2 avoidance behaviour in Drosophila by
inhibitory odorants. *Nature* 461(7261):277-281.
13.   Shahbazian MD & Grunstein M (2007) Functions of site-specific histone acetylation and
deacetylation. *Annu Rev Biochem* 76:75-100.
14.   Graff J & Tsai LH (2013) Histone acetylation: molecular mnemonics on the chromatin. *Nat Rev*
Neurosci 14(2):97-111.
15.   Minucci S & Pelicci PG (2006) Histone deacetylase inhibitors and the promise of epigenetic (and
more) treatments for cancer. *Nat Rev Cancer* 6(1):38-51.
16.   Kazantsev AG & Thompson LM (2008) Therapeutic application of histone deacetylase inhibitors
for central nervous system disorders. *Nat Rev Drug Discov* 7(10):854-868.
17.   Chuang DM, Leng Y, Marinova Z, Kim HJ, & Chiu CT (2009) Multiple roles of HDAC
inhibition in neurodegenerative conditions. *Trends Neurosci* 32(11):591-601.
18.   Jones WD, Cayirlioglu P, Kadow IG, & Vosshall LB (2007) Two chemosensory receptors
together mediate carbon dioxide detection in Drosophila. *Nature* 445:86-90.
19.   Berdnik D, Chihara T, Couto A, & Luo L (2006) Wiring Stability of the Adult Drosophila
Olfactory Circuit after Lesion. *J. Neurosci.* 26(13):3367-3376.

20.   Sigg D, Thompson CM, & Mercer AR (1997) Activity-Dependent Changes to the Brain and
Behavior of the Honey Bee, Apis mellifera (L.). *J. Neurosci.* 17(18):7148-7156.
21.   Shimazu T*, et al.* (2013) Suppression of oxidative stress by beta-hydroxybutyrate, an endogenous
histone deacetylase inhibitor. *Science* 339(6116):211-214.
22.   Whiteson KL*, et al.* (2014) Breath gas metabolites and bacterial metagenomes from cystic
fibrosis airways indicate active pH neutral 2,3-butanedione fermentation. *ISME J* 8(6):1247-1258.
23.   Krogerus K & Gibson BR (2013) Influence of valine and other amino acids on total diacetyl and
2,3-pentanedione levels during fermentation of brewer's wort. *Appl Microbiol Biotechnol*
97(15):6919-6930.
24.   Krautkramer KA, Rey FE, & JM. D (2017) Chemical signaling between gut microbiota and host
chromatin: What is your gut really saying? . *J Biol Chem* 292(21):8582-8593.
25.   James A, Austin C, Cox D, Taylor D, & Calvert R (2013) Microbiological and biochemical
origins of human axillary odour. *FEMS Microbiol Ecol* 83(3):527-540.
26.   Effantin G, Rivasseau C, Gromova M, Bligny R, & Hugouvieux-Cotte-Pattat N (2011) Massive
production of butanediol during plant infection by phytopathogenic bacteria of the genera
Dickeya and Pectobacterium. *Molecular microbiology* 82(4):988-997.
27.   Steffan JS*, et al.* (2001) Histone deacetylase inhibitors arrest polyglutamine-dependent
neurodegeneration in Drosophila. *Nature* 413(6857):739-743.
28.   Fischer A (2010) [HDAC inhibitors as therapy for neural disorders. Discovery of a new therapy].
Pharm Unserer Zeit 39(3):204-209.
29.   Chauhan MB & Chauhan NB (2015) Brain Uptake of Neurotherapeutics after Intranasal versus
Intraperitoneal Delivery in Mice. *J Neurol Neurosurg* 2(1).
30.   Jansky S*, et al.* (2021) Single-cell transcriptomic analyses provide insights into the developmental
origins of neuroblastoma. *Nature Genetics* 53(5):683-693.
31.   Pugh TJ*, et al.* (2013) The genetic landscape of high-risk neuroblastoma. *Nature Genetics*
45(3):279-284.
32.   Zhong X*, et al.* (2018) Identification of Potential Prognostic Genes for Neuroblastoma. *Front*
Genet 9.
33.   Marsh JL & Thompson LM (2006) Drosophila in the study of neurodegenerative disease. *Neuron*
52(1):169-178.
34.   Jackson GR*, et al.* (1998) Polyglutamine-expanded human huntingtin transgenes induce
degeneration of Drosophila photoreceptor neurons. *Neuron* 21(3):633-642.
35.   Thomas EA (2014) Involvement of HDAC1 and HDAC3 in the Pathology of Polyglutamine
Disorders: Therapeutic Implications for Selective HDAC1/HDAC3 Inhibitors. *Pharmaceuticals*
(Basel) 7(6):634-661.
36.   More SS, Raza A, & Vince R (2012) The butter flavorant, diacetyl, forms a covalent adduct with
2-deoxyguanosine, uncoils DNA, and leads to cell death. *J Agric Food Chem* 60(12):3311-3317.
37.   Starek-Swiechowicz B & Starek A (2014) Diacetyl exposure as a pneumotoxic factor: a review.
Rocz Panstw Zakl Hig 65(2):87-92.
38.   Morgan DL, Flake GP, Kirby PJ, & Palmer SM (2008) Respiratory toxicity of diacetyl in
C57BL/6 mice. *Toxicol Sci* 103(1):169-180.
39.   Palmer SM*, et al.* (2011) Severe airway epithelial injury, aberrant repair and bronchiolitis
obliterans develops after diacetyl instillation in rats. *PLoS One* 6(3):e17644.
40.   Kang HL, Benzer S, & Min KT (2002) Life extension in Drosophila by feeding a drug. *Proc Natl*
Acad Sci U S A 99(2):838-843.
41.   Rigler MW & Longo WE (2010) Emission of diacetyl (2,3 butanedione) from natural butter,
microwave popcorn butter flavor powder, paste, and liquid products. *Int J Occup Environ Health*
16(3):291-302.
42.   Pierce JS, Abelmann A, Spicer LJ, Adams RE, & Finley BL (2014) Diacetyl and 2,3-
pentanedione exposures associated with cigarette smoking: implications for risk assessment of
food and flavoring workers. *Crit Rev Toxicol* 44(5):420-435.

43.   Buck L & Axel R (1991) A Novel Multigene Family May Encode Odorant Receptors: A
754   Molecular Basis for Odor Recognition. *Cell* 65:175-187.
755   44.   Clyne PJ*, et al.* (1999) A novel family of divergent seven-transmembrane proteins: candidate
756   odorant receptors in *Drosophila*. *Neuron* 22(2):327-338.
757   45.   Troemel ER, Chou JH, Dwyer ND, Colbert HA, & Bargmann CI (1995) Divergent seven
758   transmembrane receptors are candidate chemosensory receptors in *C. elegans*. *Cell* 83(2):207-
759   218.
760   46.   Benton R, Vannice K, Gomezdiaz C, & Vosshall L (2009) Variant Ionotropic Glutamate
761   Receptors as Chemosensory Receptors in Drosophila. *Cell* 136(1):149-162.
762   47.   Vosshall LB, Amrein H, Morozov PS, Rzhetsky A, & Axel R (1999) A spatial map of olfactory
763   receptor expression in the *Drosophila* antenna. *Cell* 96(5):725-736.
764   48.   Knaden M & Hansson BS (2014) Mapping odor valence in the brain of flies and mice. *Curr Opin*
765   *Neurobiol* 24(1):34-38.
766   49.   Fogle KJ*, et al.* (2015) CRYPTOCHROME-mediated phototransduction by modulation of the
767   potassium ion channel beta-subunit redox sensor. *Proc Natl Acad Sci U S A* 112(7):2245-2250.
768   50.   Gregoretti IV, Lee YM, & Goodson HV (2004) Molecular evolution of the histone deacetylase
769   family: functional implications of phylogenetic analysis. *J Mol Biol* 338(1):17-31.
770   51.   Postberg J, Forcob S, Chang WJ, & Lipps HJ (2010) The evolutionary history of histone H3
771   suggests a deep eukaryotic root of chromatin modifying mechanisms. *BMC Evol Biol* 10:259.
772   52.   Witt O & Lindemann R (2009) HDAC inhibitors: magic bullets, dirty drugs or just another
773   targeted therapy. *Cancer Lett* 280(2):123-124.
774   53.   Wang J*, et al.* (2013) Potential advantages of CUDC-101, a multitargeted HDAC, EGFR, and
775   HER2 inhibitor, in treating drug resistance and preventing cancer cell migration and invasion.
776   *Mol Cancer Ther* 12(6):925-936.
777   54.   Wang J*, et al.* (2012) Sirtinol, a class III HDAC inhibitor, induces apoptotic and autophagic cell
778   death in MCF-7 human breast cancer cells. *Int J Oncol* 41(3):1101-1109.
779   55.   Yeung A*, et al.* (2012) HDAC inhibitor M344 suppresses MCF-7 breast cancer cell proliferation.
780   *Biomed Pharmacother* 66(3):232-236.
781   56.   Park JH*, et al.* (2012) A new synthetic HDAC inhibitor, MHY218, induces apoptosis or
782   autophagy-related cell death in tamoxifen-resistant MCF-7 breast cancer cells. *Invest New Drugs*
783   30(5):1887-1898.
784   57.   Cai X*, et al.* (2010) Discovery of 7-(4-(3-ethynylphenylamino)-7-methoxyquinazolin-6-yloxy)-N-
785   hydroxyheptanamide (CUDc-101) as a potent multi-acting HDAC, EGFR, and HER2 inhibitor
786   for the treatment of cancer. *J Med Chem* 53(5):2000-2009.
787   58.   Shankar S & Srivastava RK (2008) Histone deacetylase inhibitors: mechanisms and clinical
788   significance in cancer: HDAC inhibitor-induced apoptosis. *Adv Exp Med Biol* 615:261-298.
789   59.   Platta CS, Greenblatt DY, Kunnimalaiyaan M, & Chen H (2007) The HDAC inhibitor trichostatin
790   A inhibits growth of small cell lung cancer cells. *J Surg Res* 142(2):219-226.
791   60.   Komatsu N*, et al.* (2006) SAHA, a HDAC inhibitor, has profound anti-growth activity against
792   non-small cell lung cancer cells. *Oncol Rep* 15(1):187-191.
793   61.   Shakespear MR, Halili MA, Irvine KM, Fairlie DP, & Sweet MJ (2011) Histone deacetylases as
794   regulators of inflammation and immunity. *Trends Immunol* 32(7):335-343.
795   62.   Bray NL, Pimentel H, Melsted P, & Pachter L (2016) Near-optimal probabilistic RNA-seq
796   quantification. *Nat Biotechnol* 34(5):525-527.
797   63.   Soneson C, Love MI, & Robinson MD (2015) Differential analyses for RNA-seq: transcript-level
798   estimates improve gene-level inferences. *F1000Res* 4:1521.
799   64.   Risso D, Ngai J, Speed TP, & Dudoit S (2014) Normalization of RNA-seq data using factor
800   analysis of control genes or samples. *Nat Biotechnol* 32(9):896-902.
801   65.   Robinson MD, McCarthy DJ, & Smyth GK (2010) edgeR: a Bioconductor package for
802   differential expression analysis of digital gene expression data. *Bioinformatics* 26(1):139-140.

66. Mi H, *et al.* (2017) PANTHER version 11: expanded annotation data from Gene Ontology and
Reactome pathways, and data analysis tool enhancements. *Nucleic Acids Res* 45(D1):D183-D189.
67. Yu G, Wang LG, Han Y, & He QY (2012) clusterProfiler: an R package for comparing biological
themes among gene clusters. *OMICS* 16(5):284-287.
68. Andrews NC & Faller DV (1991) A rapid micropreparation technique for extraction of DNA-
binding proteins from limiting numbers of mammalian cells. *Nucleic Acids Res* 19(9):2499.

Figure Legends

Figure 1. Volatiles found in microbes can inhibit HDACs in vitro

(**A**) Antennal and whole mount brain staining of *Gr63a-Gal4* flies. Flies were exposed to diacetyl

(headspace from 10^{-2} soln) or air for 2 to 5 days. Brains and antenna were dissected on the

indicated days, fixed, and then stained for neuropil marker nc82 (red) and anti-GFP (green). (**B**)

Mean of ab1C neurons expressing GFP after indicated days of odor exposure. d4on=2,3-

butanedione. n=6, error bars=s.e.m. Schematic chemical structures of diacetyl, β-hydroxybutyrate

and Sodium butyrate. (**C**) Dose-activity curves of class I HDACs: HDAC1, HDAC2, HDAC3,

HDAC8 and class II HDAC6 treated with various concentrations of diacetyl. Percentage of HDAC

activity is relative to the activity of each enzyme without diacetyl. IC50s are indicated in the chart

areas. Error bars, S.E.M., n = 4-5. (**E**) Representative structures of odorants that inhibit HDACs,

and (**F**) Average percentage inhibition of class I HDACs: HDAC1, HDAC3 and class II HDAC4,

HDAC6 treated with 15mM of indicated volatiles. Error bar= S.D., each tested in duplicate.

Figure 2. Diacetyl increases level of histone acetylation in HEK293 cells

(**A and B**) Representative images from Western blots showing acetylation levels of H3K9 (left),

H3K14 (middle) and H4K5 (right) in HEK293 cells after 2 hours (**A**) and 6 hours (**B**) of diacetyl

exposure. PCNA (Proliferating cell nuclear antigen) is a 29 kDa nuclear protein used as a loading

control for nuclear protein extracts. (**C**) Western blots showing acetylation levels of H3K9 in

HEK293 cells treated with 100 μM diacetyl for 72, 96, and120 hours. PCNA is used for a loading

control. (**D**) Bar graph showing the relative intensities of acetylated H3K9 in HEK293 cells treated

with 100 μM diacetyl for 72 -120 hours. n = 4 samples, * $p < 0.05$, and **** $p < 0.0001$.

Figure 3. Remote exposure to HDACi volatile alters gene expression in an invertebrate, and

vertebrate.

(**A**) Schematic of odor exposure protocol for transcriptome analysis from the antennae. (**B**) Plot

highlighting up- and down-regulated genes in the diacetyl-exposed group. Red and blue dots

represent up-regulated genes (false discovery rate (FDR) < 0.05, $\log_2$ fold change (LFC) > 1) and

down-regulated genes (FDR < 0.05, LFC < 1), respectively. (**C**) Bar graphs denoting the protein

classification of the genes up- and down-regulated after odor exposure. **(D)** Schematic of diacetyl

exposure protocol for transcriptome analysis of mouse lung tissue. **(E)** Plot highlighting up- and

down-regulated genes in the diacetyl-exposed groups. Red and blue dots represent up-regulated

genes (FDR < 0.05, LFC > 1) and down-regulated genes (FDR < 0.05, LFC < 1), respectively in

lungs. **(F)** Table showing pairwise tests of significance of overlap between gene sets. P-values

from Fisher's exact test, colored with associated odds ratio (strength of association). **(G and H)**

Bar graphs denoting the protein classification of the genes up- and down-regulated in the lung after

exposure to headspace above 0.1% (v/v) **(G)** and 1% (v/v) **(H)** of diacetyl solutions.

Figure 4. Gene expression changes are partly reversible, and overlaps with known HDACi

drugs

(**A**) Schematic of odor exposure and recovery protocol for transcriptome analysis from the

antennae. (**B**) Plot highlighting up- and down-regulated genes in the recovery from diacetyl

exposure group. Red and blue dots represent up-regulated genes (FDR < 0.05, LFC > 1) and down-

regulated genes (FDR < 0.05, LFC < 1), respectively. (**C**) Table showing pairwise tests of

significance of overlap between gene sets. P-values from Fisher's exact test, colored with

associated odds ratio (strength of association). (**D** and **E**) Plots showing enrichment of up- and

down-regulated genes in sodium butyrate- (**D**) and valproic acid-treated (**E**) groups. Red and blue

dots represent up-regulated genes (FDR < 0.05, LFC > 1) and down-regulated genes (FDR < 0.05,

LFC < 1), respectively. (**F**) *Left:* Venn diagrams showing the overlaps of up-regulated genes

among diacetyl-, sodium butyrate- and valproic acid-treated groups. *Right:* Table showing

pairwise tests of significance of overlap between gene sets. P-values from Fisher's exact test. (**G**

and H) GO-enrichment analysis of the common Up-regulated and down-regulated genes across

all 3 treatments in (**F**).

Figure 5. Exposure to diacetyl vapor alters gene expression in the mouse brain

(**A**) Schematic of diacetyl exposure protocol for transcriptome analysis of mouse brain tissues. (**B**

and C) Plot showing up- and down-regulated genes in the diacetyl-exposed groups. Red and blue

dots represent up-regulated genes (FDR < 0.05, LFC > 1) and down-regulated genes (FDR < 0.05,

LFC < 1), respectively in the brain. (**D**) *Left:* Table showing pairwise tests of significance of

overlap between gene sets. P-values from Fisher's exact test. *Right:* Venn diagrams showing the

overlaps of DEGs between lung and brain groups. (**E**) Bar graphs denoting the protein

classification of the brain genes up- and down-regulated after 1% diacetyl exposure. (**F**) Mean fold

change of select neuroblastoma related genes in the brain of mice exposed to diacetyl vapors. (**G**)

Cell counts of indicated cancer cell lines in tissue culture treated with solvent control or indicated

concentration of diacetyl. N=3-6, p<0.001.

Figure 6. Odor exposure slows Huntington's neurodegeneration model in fly eye

(**A**) Schematic diagram showing temperature of experimental condition and timing of the eye

examination in pGMR-HTTQ120 flies. (**B**) Bar graph showing mean number of rhabdomeres in

each ommatidium in solvent paraffin oil (PO, blue) and diacetyl-exposed (red) pGMR-HTTQ120

flies at 1, 5, and 10 days after eclosion (AE). n = 600 ommatidia from 15 flies, **** $p < 0.0001$.

(C) A representative image of ommatidia of pGMR-HTTQ120 flies at 1 day AE. **(D and E)**

Representative images of ommatidia of pGMR-HTTQ120 flies exposed to paraffin oil (PO) or

diacetyl at 5 **(D)** and 10 **(E)** days AE. **(F and G)** Histogram showing the percent of the ommatidium

with a given number of rhabdomeres indicated on the x axis at 5 **(F)** and 10 **(G)** days AE.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6