The evolution of morphological castes under decoupled control

Lewis Flintham[1,2] and Jeremy Field[1,3,4]

1. College of Life and Environmental Sciences, University of Exeter, Penryn, Cornwall TR10 9FE, UK.
2. 2. Lf367@exeter.ac.uk
3. 3. J.P.Field@exeter.ac.uk
4. ORCID: orcid.org/0000-0003-0663-4031

Funding Statement: This work was funded partially by the University of Exeter

Acknowledgements: We would like to thank Alex Wild for kindly allowing use of the photograph for Figure 1.

Keywords: "Castes", "Eusociality", "Morphological caste", "Major evolutionary transition", "Parent offspring conflict", "Hymenoptera".

Manuscript information: 5 figures, 1 table, word count: 5841.

Abstract

The evolution of eusociality is regarded as a major evolutionary transition, where units that previously reproduced independently function as one complex entity. Advanced eusocial societies are characterised by morphologically differentiated castes and reduced conflict. We explore conditions under which morphological castes may arise and the factors constraining their evolution. Control over offspring morphology and behaviour may often be decoupled. Queens and provisioners can influence morphology directly, through the nutrition they provide, while offspring control their own behaviour as adults. Queens and provisioners may, however, influence worker behaviour indirectly, if offspring modify their behaviour in response to their morphology. Our results suggest that the evolution of a morphologically differentiated worker caste depends on the prior presence of a behavioural caste: specialist worker morphology will be mismatched with behaviour unless some offspring already choose to work. A mother's certainty about her offspring's behaviour should also be critical – less certainty results in greater mismatch. Decoupled control is important in maintaining a worker caste, and may result in reduced or no conflict between offspring and provisioners. We also show how worker productivity in the absence of a morphological trait can affect the likelihood of that trait being favoured by natural selection.

1.   Introduction

The evolution of obligate eusociality (Crespi and Yanega, 1995; Boomsma, 2013;

Boomsma and Gawne, 2018) has long been considered a major evolutionary

transition (Szathmary and Smith, 1995; West et al, 2015), characterised by

morphologically differentiated castes and reduced conflict over reproduction (Bourke,

2011; Quinones and Pen, 2017; Boomsma and Gawne, 2018). Morphological castes

are startlingly diverse, ranging from simple differences in body size (Trible and

Kronaeur, 2017; Kerr et al, 2019) to elaborate task-specific morphologies in taxa such

as turtle ants, in which the head armour of soldiers is shaped so as to seal the nest

entrance (Powell, 2016; Figure 1). Boomsma and Gawne (2018) regard irreversibility

as a key component of evolutionary transitions and identify morphological castes as

representing that evolutionarily irreversible step. Greater morphological specialisation

tends to be found in larger groups (Bourke, 1999; Bourke, 2011). This is thought to

reflect reduced reproductive conflict and increased efficiency returns from task

specialisation at larger group sizes (Bourke, 1999; Bourke, 2011).

Figure 1. The two morphological worker castes of the turtle ant Cephalotes varians;
(A) the soldier, or major worker, sporting the impressive head armour it uses to seal
the nest entrance and (B) the much smaller minor worker which conducts all the
foraging duties for the nest. Photo ©Alex Wild, used by permission.

Decoupling of control over offspring morphology and behaviour

Eusocial societies exhibiting morphological caste are characterised by a shift away

from individual control. This is to say that individuals often have limited control over

their own morphologies. Morphology is instead primarily controlled by either the queen

(and if present, the King) or the workforce at large (Wade, 2001; Schwander et al,

2008; Korb, 2008; Russell and Lummaa, 2009). This is most clearly seen in the social

insects, in particular the bees, wasps and ants, which include a significant majority of

eusocial organisms. These taxa have a holometabolic life history, in which skeletal

development occurs in a controlled environment prior to adult emergence from the

pupa (Anderson, 1972). The controlled environment is often a nest where young are

provisioned and cared for by the mother or workforce (Peters et al, 2017). The use of

a nest is ancestral for all social Hymenoptera (Boomsma and Gawne, 2018).

Combined with a holometabolic life history, nests give provisoners a huge amount of

control over the development of immature offspring by enabling them to manipulate

the environment in which offspring develop and the provisions that offspring receive

(Reinhold, 2002; Karsai and Hunt, 2002; Schwander et al, 2008; Kovacs et al, 2010;

Brand and Chapusiat, 2012). While the importance of nest use in driving eusociality

has been noted in the past (Hunt, 2007; Nowak et al, 2010; Boomsma and Gawne,

2018), its role in enabling elevated maternal control over development has received

less attention.

If we consider individuals to be fitness maximisers (Lehmann and Rousset, 2020), we

would expect those controlling phenotype to manipulate it, and hence caste, to their

own benefit (Michener and Brothers, 1974; Alexander 1974; Dawkins, 1976; Kapheim

et al 2015). Previous work addressing the evolution of such maternally influenced

traits has focused predominantly on the ability for individuals to resist manipulation by

provisioners (Keller and Nonacs, 1993; Uller and Pen, 2011), and on how provisioners

that can manipulate both offspring behaviour and morphology might influence the

origin of eusociality (Craig, 1979; Gonzalez-Forero, 2015; Gonzalez-Forero and Pena,

2021; Liu et al, 2022). Here we instead consider when morphological castes will

evolve assuming split control, with provisioners controlling the morphology of offspring

and offspring retaining control of their behavioural phenotype. We believe this to be

generally representative of the facultatively eusocial and solitary Hymenoptera (Hunt,

2007; Kapheim et al, 2011; Brand and Chapusiat, 2012; Kapheim et al 2015). In the

following text, "behavioural worker caste" refers to all individuals in a group that

behave as workers irrespective of their morphology. The term "morphological worker

caste" refers to all the individuals in a group that possess specialist morphology

irrespective of whether they behave as workers or reproductives. Specialist

morphology enhances their ability to carry out worker functions while decreasing their

fitness as reproductives. We consider a "worker caste" to be present in a group if there

is either a behavioural worker caste or a morphological worker caste.

We will start by considering a singly mated, solitary, female hymenopteran. We

assume that this female controls the morphology of her offspring, but that her offspring

control their own behaviour, so that control over behaviour and morphology is

decoupled. To begin with, we will also assume that any individuals which become

workers will remain workers (behavioural caste is immutable) and that the mother

produces an equal sex ratio. In this situation, a simple implementation of Hamilton's

rule (Hamilton, 1964; Grafen, 1985) indicates that a focal daughter, being equally

related to her siblings and her offspring (r=0.5), would favour becoming a worker when

she would be more productive as a worker than if she dispersed as a reproductive. In

contrast, because the relatedness of a mother to her offspring (r=0.5) is twice her

relatedness to her grand-offspring (r=0.25), we would expect the mother to favour her

daughter becoming a worker, and hence the imposition of a worker caste, when

workers are only half as productive as reproductives. The productivity of a worker

relative to a reproductive is therefore a limiting factor in the arisal of a worker caste

(Charnov, 1978; Bourke and Ratnieks, 1999; Reuter and Keller, 2001) and we would

expect the mother to favour imposing a worker caste - perhaps by manipulating

daughter morphology - before her daughter would favour being a member of it

(Michener and Brothers, 1974; Alexander 1974; Dawkins, 1976; Craig, 1979; Kapheim

et al 2015). Accordingly we might expect the first workers to be morphologically

specialised. In fact, however, behavioural castes always precede the evolution of

morphological castes in the social insects (Boomsma, 2013; Peters et al, 2017;

Boomsma and Gawne, 2018).

Behavioural castes are often mutable, so that individuals can switch behavioural

castes over their lifetimes. The inclusion of effects such as worker takeover, through

which a worker can become a reproductive, may reduce the threshold worker

productivity required for a daughter to favour joining a behavioural worker caste (Field

et al., 1999; Toyoizumi and Field, 2014; Price and Field, 2022). Similarly, if a daughter

can help raise more reproductive sisters (r=0.75) than reproductive brothers (r=0.25)

by staying as a worker, it will further lower the productivity of a worker relative to a

reproductive required for the daughter to favour joining a behavioural worker caste

(Trivers and Hare, 1976). The fact that we see strong evidence for behavioural worker

castes preceding morphological worker castes in the social insects (Boomsma, 2013; Peters et al, 2017; Boomsma and Gawne, 2018) therefore indicates that at least one of the following must be true: (1) the mutability of behavioural worker castes causes daughters to favour joining them at a lower worker productivity than required for mothers to favour imposing a morphological worker caste, (2) workers help raise broods which are sufficiently female biased to mean that daughters favour joining behavioural worker castes at a lower worker productivity than required for mothers to favour imposing a morphological worker caste or (3) the evolution of a morphological worker caste is somehow reliant on the prior presence of a behavioural worker caste.

On its own, the mutability of behavioural castes (explanation 1) cannot reasonably affect the argument that mothers will impose a worker caste before daughters prefer to join it. This would require workers to always immediately inherit the queen position or to produce an infinite number of offspring whenever they do (Supplementary materials Part 1). Similarly, mechanisms such as unmatedness (Godfray and Grafen, 1988; Godfray, 1990) and split sex ratios (Seger, 1983; Grafen, 1986; Gruber and Field, 2022), are unlikely to result in sex ratios that are female-biased enough to reverse the argument (Godfray and Hardy, 1990; Grafen, 1986; Boomsma and Eickwort, 1993). Even when daughters raise entirely sisters, they receive only 1.5 times the fitness benefit of raising the same number of their own offspring, less than the doubled benefit that mothers receive. Thus, we find that our proposed explanations 1 and 2, even acting in tandem, could not cause daughters to favour the arisal of a worker caste before mothers do (Supplementary materials Part 1).

Productivity in the absence of specialist morphology

Across the social Hymenoptera, worker productivity varies widely (Michener, 1964;

Jeanne et al., 2022), depending on factors such as group size (Michener, 1964;

Jeanne et al., 2022), time constraints (Lucas and Field, 2013) and environmental

constraints (Bourke, 1999; Jeanne et al., 2022), due to seasonal variations in resource

availability (Poitrineau et al., 2009). Variations in worker productivity can occur both

between species (Field and Toyoizumi, 2020) but also within them over time

(Michener, 1964; Jeanne et al., 2022). The increase in worker productivity that

specialist worker morphological traits represent for a given worker will often change as

a function of the productivity of that worker. To illustrate, consider a worker whose

primary function is nest defence. If a large worker is twice as likely to successfully

repel an attack on the nest when compared as a small worker, then as the frequency

of attacks on the nest increases the difference in number of attacks successfully

repelled by a large and small worker will also increase. We might then expect large

worker size to be most likely to evolve when attacks are most frequent, i.e. when

workers engaging in nest defence would be most productive.

Summary

To summarize, there is significant evidence in the nest building Hymenoptera that

provisioners can control the morphology of offspring (Reinhold, 2002; Karsai and

Hunt, 2002; Schwander et al, 2008; Kovacs et al, 2010; Brand and Chapusiat, 2012).

Provisioners (queens or the workforce) should favour daughters becoming workers at

a lower worker productivity than daughters themselves, so that we might expect the

first workers be morphologically specialised. Yet the evidence suggests that daughters

choose to become workers evolutionarily before morphological castes arise. Here, we

examine our proposed explanation 3, that the evolution of a morphological worker

caste is reliant on the prior presence of a behavioural worker caste. Drawing on the

relationship between form and function, we propose a simple model illustrating the

relationship between the evolution of morphological caste and the certainty the

individual controlling morphology has over the behaviour of individuals occupying that

caste. We then use this model to explore how worker productivity in the absence of

specialist morphology might influence the likelihood of that morphology evolving.

2.   The Model

We imagine a bivoltine life cycle such as that of many temperate zone small colony bees and wasps (e.g. Seger, 1983; Field et al., 2010; Couchoux & Field 2019). A single female reproductive will found a nest then produce a first brood of offspring (B1) which she will provision herself. A fraction of the female offspring from this B1 brood will immediately disperse as reproductives and found new nests, while the remainder stay and help raise a second brood (B2) of their mother's offspring. The females from the second brood all disperse as reproductives. Daughters from B1 which stay and help represent a behavioural worker caste. While in many small colony bees and wasps, behavioural castes are mutable, with worker takeover being common, we will start by assuming that they are immutable and that once an individual becomes a worker it will remain a worker.

2.1.   The role of dispersal

We wish to determine the point at which it will be beneficial for a mother to impart a novel morphological trait m onto a daughter in B1, for example through the nutrition she provides the daughter during immature development. We are primarily interested in traits that benefit one behavioural caste and are detrimental to the other. Examples might be enlarged ovaries (benefitting reproductives) and sterilisation (benefitting workers). We model the expected fitness per unit investment a mother will receive from a given daughter in the presence f' and absence f of morphological trait m (Table 1).

213　*Table 1. Parameters used in our models.*

	With morphological trait m	Without morphological trait m ("baseline value")
Fitness a worker daughter will provide to the mother	v'_w	v_w
Fitness a reproductive daughter will provide to the mother	v'_r	v_r
Cost to mother of producing a daughter	c'	c
Probability a daughter disperses (independent of morphology)	d	d
Expected fitness per unit investment a daughter will provide to the mother	f'	f
Probability a daughter disperses when trait m decreases the likelihood of daughters staying as workers.	$1 - (1 - d)(1 + a)$	d
Probability a daughter disperses when trait m increases the likelihood of daughters staying as workers.	$d(1 - a)$	d

215　*Equation 1:*

216
$$f' = \frac{(1 - d)v'_w + dv'_r}{c'}, \qquad f = \frac{(1 - d)v_w + dv_r}{c}$$

217　Trait m will be selected for when;

219　*Equation 2:*

220
$$f' > f$$

Thus the expected fitness per unit investment a mother will receive from a given

daughter is dependent on the probability of that daughter dispersing as a

reproductive d, as well as on the cost of producing the daughter. Since m here is

considered to be a novel trait, we assume that daughters do not respond to the

presence of the trait behaviourally, and have the same probability of dispersing as

reproductives with or without the trait (Table 1).

Substituting Equation 1 into Equation 2 we obtain the parameter space where we

expect trait m to be selected for (Figure 2):

Equation 3:

$$d[c(v'_r - v'_w) + c'(v_w - v_r)] + cv'_w - c'v_w > 0$$

When m is a trait conferring specialist worker morphology, we would expect it to

increase the fitness the mother receives from workers per unit cost more than the

fitness she receives from reproductives per unit cost ($\frac{v'_w}{c'} - \frac{v_w}{c} > \frac{v'_r}{c'} - \frac{v_r}{c}$)(Fisher, 1930).

When no daughters from B1 disperse as reproductives ($d = 0$), a trait which increases

the fitness a mother receives from a worker per unit cost   ($\frac{v'_w}{c'} - \frac{v_w}{c} > 0$) will always be

selected for. More generally, there will be a negative linear relationship between

probability of dispersal and the likelihood of trait m being favoured (Figure 2): the

smaller the proportion of B1 daughters that disperse as reproductives, the less often

morphology is mismatched with behaviour, and the more likely that it will be

advantageous for a mother to impart worker morphology. Because we are primarily

interested in worker morphology which decreases the fitness per unit cost of a

reproductive daughter, and behaviour here is considered independent of morphology,

we do not expect such specialist worker morphology to evolve without the prior

evolution of helping behaviour: if all B1 daughters become reproductives, trait $\mathbf{m}$

reduces offspring fitness.

Figure 2. The linear relationship between the net fitness increase a provisioner will receive from imparting morphological trait m onto a daughter ($f' - f$), and the baseline probability of a daughter dispersing (see Table 1 for parameter definitions). We have plotted this for $-1 \leq a \leq 1$ at 0.1 increments. The thick central lines represent $a = 0$ and hence Equation 3. A Daughters with trait m give more fitness as workers than reproductives ($v'_w > v'_r$) for parameter values ($v'_w = 0.55$, $v'_r = 0.5$, $v_w = 0.7$, $v_r = 1$, $c' = 0.6$, $c = 1$). B Daughters with trait m give less fitness as workers than reproductives ($v'_w < v'_r$) for parameter values ($v'_w = 0.6$, $v'_r = 0.7$, $v_w = 0.7$, $v_r = 1$, $c' = 0.8$, $c = 1$).

 2.2. Behaviour as a response to morphology

The ability to respond plastically to traits can have significant impacts on individual

fitness (El Mouden and Gardner, 2008). Morphological traits may not always be novel,

and if we assume that trait m was historically present in our population, it seems likely

that B1 daughters will have evolved to alter their dispersal decisions in response to the

trait's presence. To model this new situation, we introduce parameter a (Table 1;

Figure 3), which represents the effect of trait m on the probability of B1 daughters

dispersing. When $a > 0$, daughters with trait m are more likely to stay as workers, and

when $a = 1$ they always stay. Conversely, when $a < 0$, daughters with trait m are

more likely to disperse as reproductives, and when $a = -1$ they always disperse

(Figure 3). Incorporating parameter a gives us the following equation:

Equation 4:

262
$$f = \frac{(1-d)v_w + dv_r}{c},$$

264
265
$$f' = \begin{cases} \dfrac{[1-d(1-a)]v'_w + d(1-a)v'_r}{c'}, & a \in \{0 \le a \le 1\} \\[4mm] \dfrac{(1-d)(1+a)v'_w + [1-(1-d)(1+a)]v'_r}{c'}, & a \in \{-1 \le a \le 0\} \end{cases}$$

When $a = 0$, the presence of trait m has no effect on the dispersal decisions of

daughters, and Equation 4 reduces to Equation 1. When daughters can respond to the

presence of morphological traits ($a \ne 0$), mothers indirectly obtain a degree of control

over daughter behaviour through manipulating their morphology. The extent of this

control is represented by the magnitude of a. In order to determine when m will be

favoured by selection now that daughters can respond to it, we can substitute

Equation 4 into Equation 2 and rearrange to give the parameter space where we

expect trait m to evolve as:

Equation 5:

$$0 < \begin{cases} d[c(1-a)(v'_r - v'_w) + c'(v_w - v_r)] + cv'_w - c'v_w, & a \in \{0 \le a \le 1\} \\ d[c(1+a)(v'_r - v'_w) + c'(v_w - v_r)] + c(1+a)(v'_w - v'_r) + cv'_r - c'v_w, & a \in \{-1 \le a \le 0\} \end{cases}$$

We again find a linear relationship between d, the baseline probability of dispersal,

and the likelihood of trait m evolving (Figure 2). The relationship is again negative

when m confers specialist worker morphology, but the gradient is now dependant on

the value of a. When a is less than zero (daughters with trait m are more likely to

disperse), our y intercept is also dependant on a, such that a is moderating our

relationship between baseline dispersal and the likelihood of trait m evolving (Figure

2). This is because a causes the realised dispersal choices of daughters to correlate

with their morphologies, altering the degree of mismatch between form and function.

Parameter a therefore introduces a mechanism through which mothers can use their

control over daughter morphology to influence daughter behaviour.

To take an example, trait m might represent a simple reduction in the size of B1

daughters. Smaller daughters will be cheaper to produce (Brand and Chapusiat,

2012), so that if worker performance is independent of size (e.g. Couchoux and Field,

2019), the fitness per unit investment received by a mother from a worker will

increase. However, we might also expect trait m to decrease the fitness per unit

investment from reproductives: reproductives must found new nests and lay eggs, and

smaller reproductives may perform less well with these tasks (Couchoux and Field,

2019). If there has historically been variation in the size of daughters independent of

trait m, perhaps environmentally induced, it seems plausible that daughters will have

already evolved to alter their dispersal decisions according to their size, with smaller

daughters being more likely to stay as workers (Brand and Chapusiat, 2012). Our

value of a in this instance would be positive (since daughters with trait m are more

likely to stay as workers), and so long as $v'_w > v'_r$, trait m will be favoured by selection

over a larger range of values of d (see Figure 2A). Conversely, if $v'_w < v'_r$, so that

even with trait m daughters are less productive as workers than reproductives, a

positive value of a will reduce the likelihood of trait m evolving (Supplementary

materials Part 3, Figure 2B). Note that specialist worker morphology, which decreases

the fitness per unit cost of a reproductive daughter, can now be selected for when $d =$

1 (all offspring disperse) and $a > 0$ (Figure 2). However, it seems unlikely that

daughters would have the ability to respond to the presence of trait m in a population

where no daughters stay as workers in its absence.

We can extend this idea of maternal certainty over offspring behaviour to instances

where workers have the possibility of inheriting the queen role by considering time

spent as an inheriting worker as equivalent to time spent as a reproductive

(Supplementary materials Part 2). This changes the proportion of time a mother

expects a daughter to spend as a worker or reproductive for a given probability of

dispersal (Toyoizumi and Field, 2014). Thus the likelihood of a mismatch between

form and function when a mother imparts specialist worker morphology onto a

daughter also changes, making specialist worker morphology less likely to be

favoured. However, since time spent as a reproductive or inheriting worker still

depends on the probability of dispersal (d), the qualitative nature of the relationship

between dispersal and the parameter space where we might expect to see the

evolution of specialist morphology remains similar (Figure S1).

(a) Degree to which daughters change behaviour in presence of trait m

Figure 3. The relationship between a, the degree to which daughters change their dispersal behaviour in the presence of trait m, and offspring dispersal. Different lines represent different values of baseline dispersal (dispersal in the absence of trait m; $0 \leq d \leq 1$) in 0.25 increments. When $a > 0$ daughters are more likely to work, when $a < 0$ daughters are more likely to disperse (Table 1).

2.3.   Trait benefit as a function of baseline productivity

The fitness represented by a worker offspring bearing trait m (v'_w) is likely to be

correlated with the fitness represented by a worker that lacks trait m (v_w): trait m

represents a single modification of worker morphology, leaving the remaining

morphology unchanged. Thus, when the productivity of a worker without trait m

changes we expect the productivity of workers with trait m to change as well, so that

selection on trait m will depend on v_w. Now it may be the case that the fitness benefit

conferred by trait m varies with v_w such as in the relationship $v'_w = 2v_w$, where trait m

doubles the fitness represented by a worker. Again taking the example of worker size,

this would be the case if a large worker is twice as productive as a small worker. The

difference in productivity between large and small workers (the fitness benefit of being

large) would then be equal to the productivity of a small worker. Whether the

relationship between size and benefit to productivity is linear (Figure 4) or increases

with diminishing returns, the benefit of being a large worker ($v'_w - v_w$) increases as

the productivity of a small worker increases. We note that the productivity and fitness

of a worker may not always be equivalent, for example when the mortality of immature

offspring also changes in the presence of a morphological trait (Cant and Field, 2001;

Couvillon and Dornhaus, 2010). Substituting v'_w for $2v_w$ in Equation 5 we can

rearrange to give the following:

Equation 6:

$$0 < \begin{cases} v_w(d[\,c' + (a-1)2c] + 2c - c') + d[cv'_r(1-a) - c'v_r], & a \in \{0 \le a \le 1\} \\ v_w(d[\,c' - (1+a)2c] + 2c(1+a) - c') + [1 - (1-d)(1+a)]cv'_r - dc'v_r, \\ \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad a \in \{-1 \le a \le 0\} \end{cases}$$

As discussed above, when $v'_w = 2v_w$ we tend to see a strong positive relationship

between v_w and the likelihood of trait m being favoured by selection (Figure 4).

However, Equation 6 also highlights how this is not always the case. If our value of a

is very low ($a < -0.5$; Figure 4), or the cost of producing a daughter with trait m ($c' -$

c) is very large ($c' > 2c$), then we can in fact generate negative relationships between

v_w and the likelihood of trait m being selected, despite setting $v'_w =$

$2v_w$ (Supplementary materials Part 3).

Figure 4. *The relationship between the fitness of a worker in the absence of a trait and the likelihood of that trait being selected, when workers with that trait are twice as productive as those without it ($v'_w = 2v_w$). We have plotted this for $-1 \leq a \leq 1$ at 0.1 increments with parameter values ($v'_r = 0.8$, $v_r = 1$, $d = 0.5$, $c' = 1$, $c = 1$). The thick central line represents $a = 0$.*

3.   Discussion

The presence of a morphological worker caste is thought to be a key component of what qualifies eusociality as a major evolutionary transition (Boomsma, 2013). However, unless a behavioural worker caste has already evolved, a new morphological trait that increases worker fitness but decreases the fitness of reproductives cannot invade, for the simple reason that all offspring will be reproductives. Thus, although we might expect queens to use their control over offspring morphology to impose worker behaviour at a lower threshold benefit/cost ratio than the threshold at which offspring themselves would work voluntarily, worker behaviour will not evolve until the worker's own threshold is reached. Our analysis can therefore explain why behavioural castes always precede morphological castes in the social Hymenoptera (Boomsma, 2013; Peters et al, 2017; Boomsma and Gawne, 2018). We have primarily considered queens as being the provisioners controlling the morphology of offspring. However, once a behavioural caste has been established as we see in the social Hymenoptera, it becomes increasingly common for the workforce to instead control offspring morphology (Cameron and Robinson, 1990; Inoue et al., 1996; O'Donnell and Jeanne, 1993; Giray et al., 2005). The worker productivity at which the workforce favours offspring helping will depend on the number of times the queen has mated and also on the sex ratio of offspring produced (Reuter and Keller, 2001). However, we would still expect to see a disparity between workforce and offspring over the worker productivity at which helping is preferred (Stubblefield and Charnov, 1986).

More generally our findings suggest that in the social Hymenoptera, where it is likely that provisioners control offspring morphology while offspring control their own

behaviour (Reinhold, 2002; Karsai and Hunt, 2002; Kapheim et al, 2011; Kovacs et al,

2010; Brand and Chapusiat, 2012; Kapheim et al, 2015), the certainty that

provisioners have about offspring dispersal decisions is a key constraint on the

evolution of morphological castes, and by extension on a major evolutionary transition

(Equation 3). Reduced certainty results in greater mismatch between form and

function. These conclusions are expected to hold even when behavioural castes are

mutable (Supplementary materials Part 2). It is important to emphasise that the

certainty provisioners have over offspring dispersal decisions can be increased

through mechanisms other than a direct offspring response to morphology, for

example where there is seasonal variation in offspring dispersal. A clear example of

this is in temperate primitively eusocial bees and wasps, where daughters from the

first brood (B1) are more likely to remain at the nest as helpers than daughters from

the second brood (B2), which always disperse as reproductives (section 2.1).

Provisioners therefore have greater certainty about second brood daughter behaviour.

The ability for offspring to adjust their dispersal decisions as a response to their

morphology can also give provisioners a significant degree of indirect control over the

behaviour of female offspring and allow traits to be selected for at lower baseline rates

of dispersal than we might otherwise expect (Equation 5).

Baseline worker productivity can affect the likelihood of specialist morphology evolving

We have shown that the baseline productivity of workers (in the absence of

specialised morphology) can be important in determining when specialist morphology

is likely to evolve. The gain in worker productivity (and hence provisioner fitness)

provided by a morphological trait may vary as a function of baseline worker

productivity, an example being a trait that increases worker size, where larger workers

are always twice as productive as smaller workers. In this case, when smaller workers

have a productivity value of 5, larger workers would have a productivity of 10 (5

greater), but when smaller workers have a productivity value of 7, larger workers have

a productivity of 14 (7 greater). A trait causing larger worker size will then be most

likely to evolve when baseline worker productivity is high, since this is when larger

workers provide the largest gain. Generally, if the extra fitness provided by a

specialised morphological trait increases with baseline worker productivity (linearly or

otherwise), the trait is most likely to be favoured when baseline productivity is high. In

practice, determining the relationship between morphology and productivity can be

complex, with morphological traits not always enhancing all worker tasks equally, and

epistasis for fitness (Bank, 2022) often making the extraction of simple empirical

correlations difficult (Gimm et al, 2005; Kugonza et al, 2011; Wen et al 2015).

Decoupled control and loss of conflict between offspring and provisioners

We suggest that, with decoupled control, morphological castes become less likely to

be lost as they become more specialised, and conflict between offspring and

provisioners may disappear. After a worker caste has evolved, it might be lost when

the environment changes so that worker productivity decreases. Indeed, phylogenetic

work shows that the largely behavioural worker caste seen in sweat bees has

frequently been lost (Wcislo and Danforth, 1997), probably when bees entered

environments where the season was too short for workers to be productive (Lucas and

Field, 2013). Once a morphological worker caste is present, however, we expect

workers to evolve to respond to their morphology when making decisions about their

behaviour (section 2.2). Figure 5 illustrates the implications of this. For simplicity, we

focus on the case when offspring with specialist worker or reproductive morphology

are equally costly to produce – an example might be where worker morphology

involves just mandibles shaped specifically for defensive or foraging tasks. Initially,

offspring prefer to disperse at higher worker productivities than provisioners prefer –

productivities at which provisioners would prefer them to work (Figure 5A; see

Introduction. With each additional morphological specialisation the behavioural caste

that an offspring chooses to join should depend more and more on its morphology:

offspring will incur larger and larger costs if they mismatch their behaviour to their

morphology. The effect is almost ratchet-like: each additional morphological

specialism reduces the fitness of daughters that mismatch their behaviour to their

morphology by dispersing, so that worker productivity must drop progressively lower

for offspring with worker morphology to nevertheless prefer dispersal (Figure 5B).

Figure 5. The relationship between the level of worker morphological specialisation and the threshold worker productivity at which offspring dispersal is favoured by two parties, offspring and provisioners. Preferences under decoupled control are represented by shaded (bold-edged) and unshaded boxes. The worker productivities at which offspring will begin to disperse, so that the worker caste is lost, are represented by shaded/bold-edged boxes. Red arrows linked to faded boxes indicate the possibility that offspring regain control of their morphology, recoupling it with control of behaviour, so that they do not incur mismatch costs when dispersing.

Note, however, that because provisioners control offspring morphology, the worker

productivity at which provisioners prefer offspring to disperse remains constant (figure

5). If the fitness that provisioners obtain from workers with specialist worker

morphology falls below what they obtain from dispersers with reproductive

morphology, provisioners are expected to impart reproductive morphology – offspring

should respond to this by dispersing, thus avoiding any mismatch. Eventually, once

morphological castes are sufficiently specialised, the ratchet-like effect will mean that

the worker productivity at which offspring with worker morphology themselves prefer

dispersal drops below that at which provisioners prefer their offspring to disperse

(Figure 5C). It will then be provisioners, not offspring, that prefer the loss of a worker

caste first during evolution. At this point realised conflict between offspring and

provisioners over dispersal is lost, with both parties preferring the same outcome (the

dispersal of offspring).

If at any point individual offspring are able to regain control of their morphology during

evolution (recoupling control over morphology and behaviour), as seems to be the

case in stingless bees (Meliponini) and a few other obligately eusocial taxa (Ishay and

Landau, 1972; Wenseleers et al., 2003; Kaptein et al, 2005; Mas and Kolliker, 2008),

the threshold worker productivity required for a worker caste to be lost becomes wholly

determined by individual offspring (red arrows in Figure 5). Offspring can then choose

whether to work or disperse as reproductives without incurring costs through

mismatching morphology and behaviour. This too represents a loss of conflict over

dispersal, since now only one party is able to influence the outcome, this time resulting

in dispersal being preferred at a higher level of worker productivity. This apparently

offspring controlled caste determination seen in the Meliponini (Wenseleers et al,

2003; Wenseleers and Ratnieks, 2004) could therefore represent a vulnerability to

selection away from a worker caste, though other aspects of meliponine life history,

such as the prevalence of swarm founding, make such a reversal unlikely.

Shared control of offspring morphology, rather than control solely by provisioners, is a

possible future extension of our ideas. Shared control seems especially likely in

progressively provisioning taxa, with substantial evidence that offspring begging

influences provisioning decisions in obligately eusocial Hymenoptera (Ishay and

Landau, 1972; Creemers et al, 2003; Kaptein et al, 2005; den Boer and Duchateau,

2006; Mas and Kolliker, 2008; He et al, 2016). However, evidence for offspring

resistance to provisioner manipulation in the Hymenoptera is limited outside of

obligately eusocial species (Ishay and Landau, 1972; Wenseleers and Ratnieks, 2003;

Kaptein et al, 2005; Mas and Kolliker, 2008; Gonzalez-Forero, 2015). Begging

behaviours in solitary or facultatively eusocial progressively provisioning taxa have not

to our knowledge been reported, suggesting that our assumption of queen control over

morphology but offspring control of behaviour is reasonable before morphological

castes evolve. It is possible, however, that the absence of begging outside the

obligately eusocial Hymenoptera instead reflects a disparity in the volume of research

conducted.

2.3.1. Conclusions

(1) There is strong evidence for the evolution of a morphological worker caste being

reliant on the presence of a behavioural worker caste in the social Hymenoptera. (2)

We suggest that maternal certainty of offspring dispersal decisions is one of the main

constraints on the evolution of a morphological worker caste in the facultatively

eusocial Hymenoptera. (3) The decoupling of control over morphology and behaviour

could be a key factor maintaining a morphological worker caste and could lead to

reduced conflict between offspring and provisioners. (4) The productivity of workers in

the absence of specialist worker morphology can affect the likelihood of specialist

morphology being favoured by selection.

Bibliography

Alexander, R. D. (1974). The Evolution of Social Behavior. *Annual review of ecology and systematics, 5*, 325-383. doi:https://doi.org/10.1146/annurev.es.05.110174.001545

Anderson, D. T. (1972). The development of holometabolous insects. In S. J. Counce, & C. H. Waddington (Eds.), *Developmental systems: Insects* (Vol. 1, pp. 165-242). Academic Press.

Bank, C. (2022). Epistasis and adaptation on fitness landscapes. *Annual Review of Ecology, Evolution, and Systematics, 53*, 457-479. doi:https://doi.org//10.1146/annurev-ecolsys-102320-112153

Boomsma, J. J. (2013). Beyond promiscuity: mate-choice commitments in social breeding. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences, 368*, 20120050. doi:https://doi.org//10.1098/rstb.2012.0050

Boomsma, J. J., & Eickwort, G. C. (1993). Colony structure, provisioning and sex allocation in the sweat bee Halictus ligatus (Hymenoptera: Halictidae). *Biological Journal of the Linnean Society, 48*, 355-377. doi:https://doi.org//10.1111/j.1095-8312.1993.tb02097.x

Boomsma, J. J., & Gawne, R. (2018). Superorganismality and caste differentiation as points of no return: how the major evolutionary transitions were lost in translation. *Biological Reviews of the Cambridge Philosophical Society, 93*, 28-54. doi:https://doi.org/10.1111/brv.12330

Bourke, A. F. (1999). Colony size, social complexity and reproductive conflict in social insects. *Journal of evolutionary biology, 12*, 245-257. doi:https://doi.org//10.1046/j.1420-9101.1999.00028.x

Bourke, A. F. (2011). *Principles of social evolution.* Oxford University Press. doi:https://doi.org//10.1093/acprof:oso/9780199231157.001.0001

Bourke, A. F., & Ratnieks, F. L. (1999). Kin conflict over caste determination in social Hymenoptera. *Behavioral Ecology and Sociobiology, 46*, 287-297. doi:https://doi.org//10.1007/s002650050622

Brand, N., & Chapuisat, M. (2012). Born to be bee, fed to be worker? The caste system of a primitively eusocial insect. *Frontiers in zoology, 9*, 35. doi:https://doi.org//10.1186/1742-9994-9-35

Cameron, S. A., & Robinson, G. E. (1990). Juvenile hormone does not affect division of labor in bumble bee colonies (hymenoptera: apidae). *Annals of the Entomological Society of America, 83*, 626-631. doi:https://doi.org//10.1093/aesa/83.3.626

Cant, M. A., & Field, J. (2001). Helping effort and future fitness in cooperation animal societies. *Proceedings. Biological Sciences / the Royal Society, 268*, 1959-1964. doi:https://doi.org//10.1098/rspb.2001.1754

Charnov, E. L. (1978). Evolution of eusocial behavior: offspring choice or parental parasitism? *Journal of Theoretical Biology, 75*, 451-465. doi:https://doi.org//10.1016/0022-5193(78)90356-9

Couchoux, C., & Field, J. (2019). Parental manipulation of offspring size in social groups: a test using paper wasps. *Behavioral Ecology and Sociobiology, 73*, 36. doi:https://doi.org//10.1007/s00265-019-2646-3

Couvillon, M. J., & Dornhaus, A. (2010). Small worker bumble bees (Bombus impatiens) are hardier against starvation than their larger sisters. *Insectes sociaux, 57*, 193-197. doi:https://doi.org//10.1007/s00040-010-0064-7

Craig, R. (1979). Parental manipulation, kin selection, and the evolution of altruism. *Evolution, 33*, 319.
doi:https://doi.org//10.2307/2407622

Creemers, B., Billen, J., & Gobin, B. (2003). Larval begging behaviour in the ant Myrmica rubra.
Ethology Ecology & Evolution, 15, 261-272.
doi:https://doi.org//10.1080/08927014.2003.9522671

Crespi, B. J., & Yanega, D. (1995). The definition of eusociality. *Behavioral Ecology, 6*, 109-115.
doi:https://doi.org//10.1093/beheco/6.1.109

Dawkins, R. (1976). *The Selfish Gene.* Oxford University Press.

den Boer, S. P., & Duchateau, M. J. (2006). A larval hunger signal in the bumblebee Bombus terrestris.
Insectes sociaux, 53, 369-373. doi:https://doi.org//10.1007/s00040-006-0883-8

El Mouden, C., & Gardner, A. (2008). Nice natives and mean migrants: the evolution of dispersal-
dependent social behaviour in viscous populations. *Journal of Evolutionary Biology, 21*, 1480-
1491. doi:https://doi.org//10.1111/j.1420-9101.2008.01614.x

Field, J., & Toyoizumi, H. (2020). The evolution of eusociality: no risk-return tradeoff but the ecology
matters. *Ecology Letters, 23*, 518-526. doi:https://doi.org//10.1111/ele.13452

Field, J., Shreeves, G., & Sumner, S. (1999). Group size, queuing and helping decisions in facultatively
eusocial hover wasps. *Behavioral Ecology and Sociobiology, 45*, 378-385.
doi:https://doi.org//10.1007/s002650050574

Fisher, R. A. (1930). *The genetical theory of natural selection.* Oxford: Clarendon Press.
doi:https://doi.org//10.5962/bhl.title.27468

Giray, T., Giovanetti, M., & West-Eberhard, M. J. (2005). Juvenile hormone, reproduction, and worker
behavior in the neotropical social wasp Polistes canadensis. *Proceedings of the National*
Academy of Sciences of the United States of America, 102, 3330-3335.
doi:https://doi.org//10.1073/pnas.0409560102

Godfray, H. C. (1990). The causes and consequences of constrained sex allocation in haplodiploid
animals. *Journal of Evolutionary Biology, 3*, 3-17. doi:https://doi.org//10.1046/j.1420-
9101.1990.3010003.x

Godfray, H. C., & Grafen, A. (1988). Unmatedness and the evolution of eusociality. *The American*
Naturalist, 131, 303-305. doi:https://doi.org//10.1086/284791

Godfray, H. C., & Hardy, I. C. (1990). Estimating the frequency of constrained sex allocation in field
populations of hymenoptera. *Behaviour, 114*, 137-147.
doi:https://doi.org//10.1163/156853990X00086

González-Forero, M. (2015). Stable eusociality via maternal manipulation when resistance is costless.
Journal of Evolutionary Biology, 28, 2208-2223. doi:https://doi.org//10.1111/jeb.12744

González-Forero, M., & Peña, J. (2021). Eusociality through conflict dissolution. *Proceedings. Biological*
Sciences / the Royal Society, 288, 20210386. doi:https://doi.org//10.1098/rspb.2021.0386

Grafen, A. (1985). A geometric view of relatedness. *Oxford surveys in evolutionary biology, 2*, 28-89.

Grafen, A. (1986). Split sex ratios and the evolutionary origins of eusociality. *Journal of Theoretical*
Biology, 122, 95-121. doi:https://doi.org//10.1016/S0022-5193(86)80227-2

Grimm, L. H., Kelly, S., Krull, R., & Hempel, D. C. (2005). Morphology and productivity of filamentous
fungi. *Applied Microbiology and Biotechnology, 69*, 375-384.
doi:https://doi.org//10.1007/s00253-005-0213-5

Gruber, J., & Field, J. (2022). Male survivorship and the evolution of eusociality in partially bivoltine
sweat bees. *Plos One, 17*, e0276428. doi:https://doi.org//10.1371/journal.pone.0276428

Hamilton, W. D. (1964). The genetical evolution of social behaviour. I. *Journal of Theoretical Biology, 7*,
1-16. doi:https://doi.org//10.1016/0022-5193(64)90038-4

He, X. J., Zhang, X. C., Jiang, W. J., Barron, A. B., Zhang, J. H., & Zeng, Z. J. (2016). Starving honey bee
(Apis mellifera) larvae signal pheromonally to worker bees. *Scientific Reports, 6*, 22359.
doi:https://doi.org//10.1038/srep22359

Hunt, J. H. (2007). *The Evolution of Social Wasps.* Oxford University Press.

Inoue, T., Salmah, S., & Sakagami, S. (1996). Individual Variations in Worker Polyethism of the
Sumatran Stingless Bee, Trigona (Tetragonula) minangkabau (Apidae, Meliponinae). *Japanese
journal of entomology, 64*, 641-668.

Ishay, J., & Landau, E. M. (1972). Vespa larvae send out rhythmic hunger signals. *Nature, 237*, 286-287.
doi:https://doi.org//10.1038/237286a0

Jeanne, R. L., Loope, K. J., Bouwma, A. M., Nordheim, E. V., & Smith, M. L. (2022). Five decades of
misunderstanding in the social Hymenoptera: a review and meta-analysis of Michener's
paradox. *Biological Reviews of the Cambridge Philosophical Society, 97*, 1559-1611.
doi:https://doi.org//10.1111/brv.12854

Kapheim, K. M., Bernal, S. P., Smith, A. R., Nonacs, P., & Wcislo, W. T. (2011). Support for maternal
manipulation of developmental nutrition in a facultatively eusocial bee, Megalopta genalis
(Halictidae). *Behavioral Ecology and Sociobiology, 65*, 1179-1190.
doi:https://doi.org//10.1007/s00265-010-1131-9

Kapheim, K. M., Nonacs, P., Smith, A. R., Wayne, R. K., & Wcislo, W. T. (2015). Kinship, parental
manipulation and evolutionary origins of eusociality. *Proceedings. Biological Sciences / the
Royal Society, 282*, 20142886. doi:https://doi.org//10.1098/rspb.2014.2886

Kaptein, N., Billen, J., & Gobin, B. (2005). Larval begging for food enhances reproductive options in the
ponerine ant Gnamptogenys striatula. *Animal Behaviour, 69*, 293-299.
doi:https://doi.org//10.1016/j.anbehav.2004.04.012

Karsai, I., & Hunt, J. H. (2002). Food Quantity AffectTraits of Offspring in the Paper Wasp Polistes
metricus (Hymenoptera: Vespidae). *Environmental Entomology, 31*, 99-106.
doi:https://doi.org//10.1603/0046-225X-31.1.99

Keller, L., & Nonacs, P. (1993). The role of queen pheromones in social insects: queen control or queen
signal? *Animal Behaviour, 45*, 787-794. doi:https://doi.org//10.1006/anbe.1993.1092

Kerr, N. Z., Crone, E. E., & Williams, N. M. (2019). Integrating vital rates explains optimal worker size
for resource return by bumblebee workers. *Functional Ecology, 33*, 467-478.
doi:https://doi.org//10.1111/1365-2435.13251

Korb, J. (2008). The ecology of social evolution in termites. In J. Korb, & J. Heinze (Eds.), *Ecology of*
social evolution (pp. 151-174). Berlin, Heidelberg: Springer Berlin Heidelberg.
doi:https://doi.org/10.1007/978-3-540-75957-7

Kovacs, J. L., Hoffman, E. A., Marriner, S. M., Rekau, J. A., & Goodisman, M. A. (2010). Environmental
and genetic influences on queen and worker body size in the social wasp Vespula maculifrons.
Insectes sociaux, 57, 53-65. doi:https://doi.org//10.1007/s00040-009-0050-0

Kugonza, D. R., Nabasirye, M., Mpairwe, D., Hanotte, O., & Okeyo, A. M. (2011). Productivity and
morphology of Ankole cattle in three livestock production systems in Uganda. *Animal Genetic*
Resources/Ressources génétiques animales/Recursos genéticos animales, 48, 13-22.
doi:https://doi.org//10.1017/S2078633611000038

Lehmann, L., & Rousset, F. (2020). When do individuals maximize their inclusive fitness? *The American*
Naturalist, 195, 717-732. doi:https://doi.org//10.1086/707561

Liu, M., West, S. A., & Wild, G. (2022). The evolution of manipulative cheating. *eLife, 11*.
doi:https://doi.org//10.7554/eLife.80611

Lucas, E. R., & Field, J. (2013). Caste determination through mating in primitively eusocial societies.
Journal of Theoretical Biology, 335, 31-39. doi:https://doi.org//10.1016/j.jtbi.2013.05.032

Mas, F., & Kölliker, M. (2008). Maternal care and offspring begging in social insects: chemical
signalling, hormonal regulation and evolution. *Animal Behaviour, 76*, 1121-1131.
doi:https://doi.org//10.1016/j.anbehav.2008.06.011

Michener, C. D. (1964). Reproductive efficiency in relation to colony size in hymenopterous societies.
Insectes sociaux, 11, 317-341. doi:https://doi.org//10.1007/BF02227433

Michener, C. D., & Brothers, D. J. (1974). Were workers of eusocial hymenoptera initially altruistic or
oppressed? *Proceedings of the National Academy of Sciences of the United States of America,*
71, 671-674. doi:https://doi.org//10.1073/pnas.71.3.671

Nowak, M. A., Tarnita, C. E., & Wilson, E. O. (2010). The evolution of eusociality. *Nature, 466*, 1057-
1062. doi:https://doi.org//10.1038/nature09205

O'Donnell, S., & Jeanne, R. L. (1993). Methoprene accelerates age polyethism in workers of a social
wasp (Polybia occidentalis). *Physiological entomology, 18*, 189-194.
doi:https://doi.org//10.1111/j.1365-3032.1993.tb00467.x

Peters, R. S., Krogmann, L., Mayer, C., Donath, A., Gunkel, S., Meusemann, K., . . . Niehuis, O. (2017).
Evolutionary history of the hymenoptera. *Current Biology, 27*, 1013-1018.
doi:https://doi.org//10.1016/j.cub.2017.01.027

Poitrineau, K., Mitesser, O., & Poethke, H. J. (2009). Workers, sexuals, or both? Optimal allocation of
resources to reproduction and growth in annual insect colonies. *Insectes sociaux, 56*, 119-129.
doi:https://doi.org//10.1007/s00040-009-0004-6

Powell, S. (2016). A comparative perspective on the ecology of morphological diversification in
complex societies: nesting ecology and soldier evolution in the turtle ants. *Behavioral Ecology*
and Sociobiology, 70, 1075-1085. doi:https://doi.org//10.1007/s00265-016-2080-8

Powell, S., Price, S. L., & Kronauer, D. J. (2020). Trait evolution is reversible, repeatable, and decoupled
in the soldier caste of turtle ants. *Proceedings of the National Academy of Sciences of the*
United States of America, 117, 6608-6615. doi:https://doi.org//10.1073/pnas.1913750117

Price, T. N., & Field, J. (2022). Sisters doing it for themselves: extensive reproductive plasticity in
workers of a primitively eusocial bee. *Behavioral Ecology and Sociobiology, 76*, 85.
doi:https://doi.org//10.1007/s00265-022-03196-4

Quiñones, A. E., & Pen, I. (2017). A unified model of Hymenopteran preadaptations that trigger the
evolutionary transition to eusociality. *Nature Communications, 8*, 15920.
doi:https://doi.org//10.1038/ncomms15920

Reinhold, K. (2002). Maternal effects and the evolution of behavioral and morphological characters: a
literature review indicates the importance of extended maternal care. *The Journal of Heredity,*
93, 400-405. doi:https://doi.org//10.1093/jhered/93.6.400

Reuter, M., & Keller, L. (2001). Sex ratio conflict and worker production in eusocial hymenoptera. *The*
American Naturalist, 158, 166-177. doi:https://doi.org//10.1086/321311

Russell, A. F., & Lummaa, V. (2009). Maternal effects in cooperative breeders: from hymenopterans to
humans. *Philosophical Transactions of the Royal Society of London. Series B, Biological*
Sciences, 364, 1143-1167. doi:https://doi.org//10.1098/rstb.2008.0298

Schwander, T., Humbert, J.-Y., Brent, C. S., Cahan, S. H., Chapuis, L., Renai, E., & Keller, L. (2008).
Maternal effect on female caste determination in a social insect. *Current Biology, 18*, 265-269.
doi:https://doi.org//10.1016/j.cub.2008.01.024

Seger, J. (1983). Partial bivoltinism may cause alternating sex-ratio biases that favour eusociality.
Nature, 301, 59-62. doi:https://doi.org//10.1038/301059a0

Stubblefield, J. W., & Charnov, E. L. (1986). Some conceptual issues in the origin of eusociality.
Heredity, 57 (Pt 2), 181-187. doi:https://doi.org//10.1038/hdy.1986.108

Szathmáry, E., & Smith, J. M. (1995). The major evolutionary transitions. *Nature, 374*, 227-232.
doi:https://doi.org//10.1038/374227a0

Toyoizumi, H., & Field, J. (2014). Dynamics of social queues. *Journal of Theoretical Biology, 346*, 16-22.
doi:https://doi.org//10.1016/j.jtbi.2013.12.017

Trible, W., & Kronauer, D. J. (2017). Caste development and evolution in ants: it's all about size.
Journal of Experimental Biology, 220, 53-62. doi:https://doi.org//10.1242/jeb.145292

Trivers, R., & Hare, H. (1976). Haplodiploidy and the Evolution of the Social Insects. *Science, 191*, 249-
263. doi:https://doi.org//10.1126/science.1108197

Uller, T., & Pen, I. (2011). A theoretical model of the evolution of maternal effects under parent-
offspring conflict. *Evolution, 65*, 2075-2084. doi:https://doi.org//10.1111/j.1558-
5646.2011.01282.x

Wade, M. J. (2001). Maternal effect genes and the evolution of sociality in haplo-diploid organisms.
Evolution, 55, 453-458. doi:https://doi.org//10.1554/0014-
3820(2001)055[0453:megate]2.0.co;2

Wcislo, W. T., & Danforth, B. N. (1997). Secondarily solitary: the evolutionary loss of social behavior.
Trends in Ecology & Evolution, 12, 468-474. doi:https://doi.org//10.1016/s0169-
5347(97)01198-1

Wen, W., Li, K., Alseekh, S., Omranian, N., Zhao, L., Zhou, Y., . . . Fernie, A. R. (2015). Genetic
determinants of the network of primary metabolism and their relationships to plant
performance in a maize recombinant inbred line population. *The Plant Cell, 27*, 1839-1856.
doi:https://doi.org//10.1105/tpc.15.00208

Wenseleers, T., & Ratnieks, F. L. (2004). Tragedy of the commons in Melipona bees. *Proceedings.*
Biological Sciences / the Royal Society, 271 Suppl 5, S310-S312.
doi:https://doi.org//10.1098/rsbl.2003.0159

Wenseleers, T., Ratnieks, F. L., & Billen, J. (2003). Caste fate conflict in swarm-founding social
hymenoptera: an inclusive fitness analysis. *Journal of Evolutionary Biology, 16*, 647-658.
doi:https://doi.org//10.1046/j.1420-9101.2003.00574.x

West, S. A., Fisher, R. M., Gardner, A., & Kiers, E. T. (2015). Major evolutionary transitions in
individuality. *Proceedings of the National Academy of Sciences of the United States of America,*
112, 10112-10119. doi:https://doi.org//10.1073/pnas.1421402112