# The evolution of morphological castes under decoupled control

Lewis Flintham[1,2,3] and Jeremy Field[1]

1. Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Penryn, Cornwall TR10 9EZ, United Kingdom
2. Division of Biosciences, University College London, London, United Kingdom
3. School of Biological Sciences, Royal Holloway University of London, Egham, Surrey, TW20 0EX, United Kingdom

**Contact information:**

Lewis Flintham. Email: <Lewis.Flintham.23@ucl.ac.uk>.

Jeremy Field. Email: <J.P.Field@exeter.ac.uk>. Mailing address: Jeremy Field, Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Penryn, Cornwall TR10 9EZ, United Kingdom.

*Acknowledgements*

This work was funded by the University of Exeter. We would like to thank Alex Wild for kindly allowing use of the photograph for Figure 1. We would also like to thank Andy Gardner, Peter Nonacs, and two anonymous reviewers for their comments on the manuscript, and Vincent Jansen for helpful discussions about the models.

*Data Availability Statement*

This study did not generate any new data. Additional information supporting this publication is available as supplementary information accompanying this publication.

**Abstract**

Eusociality, where units that previously reproduced independently function as one entity, is of major interest in evolutionary biology. Obligate eusociality is characterised by morphologically differentiated castes and reduced conflict. We explore conditions under which morphological castes may arise in the Hymenoptera and factors constraining their evolution. Control over offspring morphology and behaviour seem likely to be decoupled. Provisioners (queens and workers) can influence offspring morphology directly through the nutrition they provide, while adult offspring control their own behaviour. Provisioners may, however, influence worker behaviour indirectly if offspring modify their behaviour in response to their morphology. If manipulation underlies helping, we should not see helping evolve before specialised worker morphology, yet empirical observations suggest that behavioural castes precede morphological castes. We use evolutionary invasion analyses to show how the evolution of a morphologically differentiated worker caste depends on the prior presence of a behavioural caste: specialist worker morphology will be mismatched with behaviour unless some offspring already choose to work. A mother's certainty about her offspring's behaviour is also critical – less certainty results in greater mismatch. We show how baseline worker productivity can affect the likelihood of a morphological trait being favoured by natural selection. We then show how under a decoupled control scenario, morphologically differentiated castes should be less and less likely to be lost as they become more specialised. We also suggest that for eusociality to be evolutionarily irreversible, workers must be unable to functionally replace reproductives and reproductives must be unable to reproduce without help from workers.

## 1.    Introduction

Eusociality, where some individuals in a social group forfeit their own reproduction to help others to reproduce, is a phenomenon of major interest in evolutionary biology. Division of labour is most extreme in so-called obligate eusocial groups, where workers are not able to functionally replace reproductives and reproductives cannot produce reproductive offspring without help from a worker caste (see also Crespi and Yanega, 1995). These groups are characterised by morphologically differentiated castes and reduced conflict over reproduction (Bourke, 2011; Quinones and Pen, 2017; Boomsma and Gawne, 2018). In contrast, in facultatively eusocial groups, either reproductives cannot produce reproductive offspring without help from a worker caste (Crespi and Yanega, 1995), or workers are not able to functionally replace reproductive roles, but not both. In this instance, caste differences tend to be mainly behavioural (Boomsma and Gawne, 2018), and the functional constraints on caste members are typically the result of ecological conditions and life history traits (e.g.

*Figure 1. The two morphological worker castes of the turtle ant Cephalotes varians; (A) the soldier, or major worker, sporting the impressive head armour it uses to seal the nest entrance and (B) the much smaller minor worker which conducts all the foraging duties for the nest. Photo ©Alex Wild, used with permission. This image is not covered by the terms of the Creative Commons license of this publication. For permission to reuse, please contact the rights holder.*

mating opportunities or swarm founding) rather than morphology. Finally, cooperatively

breeding groups are those where workers can functionally replace reproductives and

reproductives are capable of producing reproductive offspring without help from a

worker caste (Crespi and Yanega, 1995). Different authors define these categories in

slightly different ways, and the above working definitions are provided for clarity.

The significance of specialised castes as a means of social organisation in natural

systems has long been understood (Oster and Wilson, 1978), and much previous work

has focused on their implications and how they might evolve (Oster and Wilson, 1978;

Bourke, 2011; Quinones and Pen, 2017; Cooper and West, 2018; Boomsma and

Gawne, 2018). Morphological castes are startlingly diverse, ranging from simple

differences in body size (Trible and Kronaeur, 2017; Kerr et al., 2019) to elaborate

task-specific morphologies in taxa such as turtle ants, in which the head armour of

soldiers is shaped so as to seal the nest entrance (Powell, 2016; Figure 1). Greater

morphological specialisation tends to be found in larger groups (Bourke, 1999;

Bourke, 2011; Ferguson-Gow et al., 2014). This is thought to reflect both reduced

reproductive conflict, favouring the evolution of caste specific morphology when

individuals control the expression of their own traits, and increased efficiency returns

from task specialisation, favouring the evolution of morphological subcastes (Bourke,

1999; Bourke, 2011). But it may not always be reasonable to assume that individuals

control the expression of their own traits. Here, we use a pairwise evolutionary

invasion analysis (Avila and Mullon, 2023) to  investigate how a morphological worker

caste will evolve when control over worker phenotype is split between offspring and

provisioners (individuals providing resources to offspring), such that manipulation can

potentially occur.

*Decoupling of control over offspring morphology and behaviour*

Eusocial groups exhibiting morphological caste are characterised by a shift away from individual control. This is to say that individuals often have limited control over their own morphologies. Morphology is instead primarily determined by either the queen (and if present, the King) or the workforce as a whole (Wade, 2001; Schwander et al., 2008; Korb, 2008; Russell and Lummaa, 2009). This is most clearly seen in the social insects, in particular the bees, wasps and ants, which include a significant majority of eusocial organisms. These taxa have a holometabolic life history, in which exoskeletal development occurs in a controlled environment prior to adult emergence from the pupa (Anderson, 1972). The controlled environment is often a nest where young are provisioned and cared for by the mother or workforce (Peters et al., 2017). The use of a nest is ancestral for all social Hymenoptera (Boomsma and Gawne, 2018). Combined with a holometabolic life history, nests give provisoners a huge amount of control over the development of immature offspring by enabling them to manipulate the environment in which offspring develop and the provisions that offspring receive (Reinhold, 2002; Karsai and Hunt, 2002; Schwander et al., 2008; Kovacs et al., 2010; Brand and Chapusiat, 2012). While the importance of nest use in driving eusociality is widely accepted (Hunt, 2007; Nowak et al., 2010; Boomsma and Gawne, 2018), its role in enabling elevated maternal control over development, and hence a greater degree of maternal manipulation, has received less attention.

If we consider individuals to be fitness maximisers (Lehmann and Rousset, 2020), we would expect those controlling phenotype to manipulate it, and hence caste, to their own benefit (Michener and Brothers, 1974; Alexander 1974; Dawkins, 1976; Kapheim

et al., 2015). Previous work addressing the evolution of such maternally influenced traits has focused predominantly on the ability of individuals to resist manipulation by provisioners (Keller and Nonacs, 1993; Wenseleers et al., 2003; Uller and Pen, 2011), and on how provisioners that can manipulate both offspring behaviour and morphology might influence the origin of eusociality (Craig, 1979; Gonzalez-Forero, 2015; Gonzalez-Forero and Pena, 2021; Rodrigues and Gardner, 2021; Liu et al., 2022). Our focus in this paper, however, is not the origin of eusociality or helping behaviour, but the evolution of morphological castes *per se*, and we consequently aim to characterise when manipulation of offspring phenotype by provisioners will be favoured by selection. It has previously been noted that there may be costs as well as benefits to manipulating offspring, such as manipulator error resulting in offspring death (Wenseleers et al., 2003). We aim to identify key factors which influence the costs and benefits of manipulation and provide insight into how these factors might shape the evolution of morphological castes in the social Hymenoptera. To do so we assume a state of split control, with provisioners controlling the morphology of offspring but offspring retaining control of their behavioural phenotype. We believe this assumption to be generally representative of worker traits found in the facultatively eusocial, cooperatively breeding, and solitary Hymenoptera (Hunt, 2007; Kapheim et al., 2011; Brand and Chapusiat, 2012; Kapheim et al., 2015), from which obligate eusociality evolved. The terminology surrounding eusociality can be confusing, with different authors advocating different usages. For the purposes of this paper, "behavioural worker caste" refers to all individuals in a group that behave as workers irrespective of their morphology. The term "morphological worker caste" refers to all the individuals in a group that possess specialist worker morphology, irrespective of whether they behave as workers or reproductives. Specialist worker morphology

enhances individuals' ability to carry out worker functions while decreasing their fitness

as reproductives. We consider a "worker caste" to be present in a group if there is

either a behavioural worker caste or a morphological worker caste.

We will start by considering a singly mated, solitary, female hymenopteran. We

assume that this female controls the morphology of her offspring, but that her offspring

control their own behaviour, so that control over behaviour and morphology is

decoupled. To begin with, we will also assume that any individuals which become

workers will remain workers (behavioural caste is immutable) and that the mother

produces an equal sex ratio. In this situation, a simple implementation of Hamilton's

rule (Hamilton, 1964; Grafen, 1985) indicates that a focal daughter, being equally

related to her siblings and her offspring (r=0.5), would favour becoming a worker when

she would be more productive as a worker than if she dispersed as a reproductive. In

contrast, because the relatedness of a mother to her offspring (r=0.5) is twice her

relatedness to her grand-offspring (r=0.25), we would expect the mother to favour her

daughter becoming a worker, and hence her imposing a worker caste through altering

the environment she provides for offspring development, when workers are only half

as productive as reproductives. The productivity (number of offspring produced over a

lifetime) of a worker relative to a reproductive is therefore critical for the arisal of a

worker caste (Charnov, 1978; Bourke and Ratnieks, 1999; Reuter and Keller, 2001)

and we would expect the mother to favour imposing a worker caste – perhaps by

manipulating daughter morphology – before her daughter would favour being a

member of it, i.e. at a lower worker/reproductive productivity ratio (Michener and

Brothers, 1974; Alexander 1974; Dawkins, 1976; Craig, 1979; Kapheim et al., 2015).

Inducing morphological specialization in daughters, so that they themselves will be

more productive if they choose to become workers, is an obvious mechanism by which

mothers could impose a worker caste. Accordingly, under our assumed state of

decoupled control, we might expect the first workers to be morphologically specialised.

While some authors argue that this is sometimes indeed the case (Piekarski et al.,

2018; Boomsma 2022 Box 6.2), others consider that behavioural castes preceded the

evolution of morphological castes in the social insects (Bourke, 2011; Quinones and

Pen, 2017; Beekman and Oldroyd, 2019; Bourke 2023)), suggesting that the evolution

of a morphological worker caste is somehow reliant on the prior presence of a

behavioural worker caste (see Discussion and Supplementary materials Part 1).

*Productivity in the absence of specialist morphology*

Across the social Hymenoptera, worker productivity varies widely (Michener, 1964;

Jeanne et al., 2022), depending on factors such as group size (Michener, 1964;

Jeanne et al., 2022), time constraints (Lucas and Field, 2013) and environmental

constraints (Bourke, 1999; Jeanne et al., 2022) due to seasonal variations in resource

availability (Poitrineau et al., 2009). Variation in worker productivity can occur between

species (Field and Toyoizumi, 2020) but also within them over time (Michener, 1964;

Jeanne et al., 2022).  The increase in worker productivity that a specialist worker

morphological trait represents for a given worker will often depend on baseline worker

productivity (worker productivity without that morphological trait). To illustrate, consider

a small worker whose primary function is nest defence. If a large worker is twice as

likely to successfully repel an attack on the nest, then as the frequency of attacks

increases, the difference in number of attacks successfully repelled by a large and

small worker will also increase. We might then expect large worker size to be most

likely to evolve when attacks are most frequent, i.e. when workers engaging in nest

defence would be most productive.

*Summary of aims*

To summarise, there is significant evidence in the nest building Hymenoptera that

provisioners can control the morphology of offspring (Reinhold, 2002; Karsai and

Hunt, 2002; Schwander et al., 2008; Kovacs et al., 2010; Brand and Chapusiat, 2012).

Under the additional assumption that individuals control their own behaviour, we aim

to characterise some of the costs and benefits determining when provisioners would

favour manipulating the morphology of offspring. Drawing on the relationship between

form and function, we propose a simple model illustrating the relationship between the

evolution of morphological caste and the certainty the individual controlling

morphology has over the behaviour of individuals occupying that caste. We then go on

to use this model to explore how worker productivity in the absence of specialist

morphology might influence the benefit that morphology provides, and hence the

likelihood of that morphology evolving. Finally we extend our perspective of decoupled

control to the evolution of eusociality more generally, with particular reference to the

topics of reversibility and the arisal of helping behaviour via maternal manipulation.

## 2.    The Model

We imagine a bivoltine life cycle such as that of many temperate zone cooperatively breeding bees and wasps (e.g. Seger, 1983; Field et al., 2010; Couchoux & Field 2019). A single female reproductive founds a nest then provisions a first brood of offspring (B1) herself. A fraction of the female offspring from this B1 brood will immediately disperse as reproductives and found new nests, while the remainder stay and help raise a second brood (B2) of their mother's offspring. The females from the second brood all disperse as reproductives. Daughters from B1 which stay and help represent a behavioural worker caste. While in many small colony cooperatively breeding bees and wasps, behavioural castes are mutable, with worker takeover being common, we will start by assuming that they are immutable and that once an individual becomes a worker it will remain a worker. We formulate the following models in the style of a pairwise evolutionary invasion analysis where we ignore non-selective effects and we make the additional simplifying assumption that mutants are rare (Avila and Mullon, 2023). In addition, since we are primarily concerned with when a given manipulation trait can first be expected to spread, we focus on finding invasion conditions as opposed to evolutionary stability. A more formal derivation of the equations in this section can be found in Supplementary materials Part 4

### 2.1.    The role of dispersal

We wish to determine when it will be beneficial for a mother (queen) to impart a novel morphological trait $m$ onto a daughter in B1, for example through the nutrition she provides the daughter during immature development. We are primarily interested in traits that provide a fitness benefit to the mother when imparted on offspring of one

behavioural caste but a fitness cost to the mother when imparted to the other caste.

Examples might be enlarged ovaries – benefitting reproductives but representing

wasted investment on workers - and sterilisation – detrimental to reproductives but

beneficial to the mother when imparted to workers as it avoids colony provisions being

used to raise the offspring of workers (with which the mother shares lower

relatedness). We model the expected fitness (copies of genetic material contributed to

a future generation) per unit investment a mother will receive from a given daughter in

the presence $f'$ and absence $f$ of morphological trait $m$ (Table 1). It may be helpful to

envisage trait $m$ as a change in offspring morphology resulting from a mutant allele for

manipulation in the mother. $f'$ and $f$ then represent the fitness a mother, with ($f'$) and

without ($f$) this mutant allele, will expect to receive from a given daughter.

Table 1. Parameters used in our models.

| | With morphological trait $m$ | Without morphological trait $m$ ("baseline value") |
|---|---|---|
| Fitness a worker daughter will provide to the mother | $v'_w$ | $v_w$ |
| Fitness a reproductive daughter will provide to the mother | $v'_r$ | $v_r$ |
| Cost to mother of producing a daughter | $c'$ | $c$ |
| Probability a daughter disperses (independent of morphology) | $d$ | $d$ |
| Expected fitness per unit investment a daughter will provide to the mother | $f'$ | $f$ |
| Probability a daughter disperses when trait $m$ increases the likelihood of daughters staying as workers. | $d(1-a)$ | $d$ |

250  *Equation 1:*

$$f' = \frac{(1-d)v'_w + dv'_r}{c'}, \qquad f = \frac{(1-d)v_w + dv_r}{c}$$

252  Trait $m$ will be expected to invade when;

254   *Equation 2:*

255
$$f' > f$$

Thus the expected fitness per unit investment a mother will receive from a given

daughter is dependent on the probability of that daughter dispersing as a

reproductive $d$, as well as on the cost of producing the daughter. Since $m$ here is

considered to be a novel trait, we assume that daughters do not respond to the

presence of the trait behaviourally, and have the same probability of dispersing with or

without the trait (Table 1). We will later relax this assumption to see how morphology

will evolve when offspring do change their behaviour in response.

Substituting Equation 1 into Equation 2 we obtain the parameter space where we

expect imparting trait $m$ to be selected for (Figure 2):

*Equation 3:*

267
$$d[c(v'_r - v'_w) + c'(v_w - v_r)] + cv'_w - c'v_w > 0$$

When $m$ is a trait conferring specialist worker morphology, we would expect it to

increase the fitness the mother receives from workers per unit cost more than the

fitness she receives from reproductives per unit cost ($\frac{v'_w}{c'} - \frac{v_w}{c} > \frac{v'_r}{c'} - \frac{v_r}{c}$). When no

daughters from B1 disperse as reproductives ($d = 0$), a trait which increases the

fitness a mother receives from a worker per unit cost ($\frac{v'_w}{c'} - \frac{v_w}{c} > 0$) will always be

selected for. More generally, there will be a negative linear relationship between

probability of dispersal and the likelihood of manipulation for trait $m$ being favoured

(Figure 2): the smaller the proportion of B1 daughters that disperse as reproductives,

the more likely that it is advantageous for a mother to impart worker morphology,

because morphology will less often be mismatched with behaviour. Because we are

primarily interested in worker morphology which decreases the fitness per unit cost of

a reproductive daughter, and behaviour here is considered independent of

morphology, we do not expect such specialist worker morphology to evolve without the

prior evolution of helping behaviour: if all B1 daughters become reproductives, trait **m**

inevitably reduces offspring fitness.

*Figure 2. The linear relationship between the net fitness increase a provisioner will receive from imparting morphological trait **m** onto a daughter ($f' - f$), and the baseline probability of a daughter dispersing (see Table 1 for parameter definitions). We have plotted this for $0 \le a \le 1$ at 0.1 increments. The thick line represents $a = 0$ and hence Equation 3. For parameter values ($v'_w$ = 0.55, $v'_r$ = 0.5, $v_w$ = 0.7, $v_r$ = 1, $c'$ = 0.6, $c$ = 1). Dashed line at y=0 represents the threshold for manipulators imparting trait **m** to invade (Equation 2)*

 *2.2.   Behaviour as a response to morphology*

In the previous section, morphological evolution responds to behaviour in the sense

that worker morphology cannot evolve without the prior presence of a behavioural

worker caste. Next, we consider the possibility that individuals can respond plastically

to their morphology (El Mouden and Gardner, 2008) by choosing to disperse as

reproductives or not. Morphological traits may not always be novel, and if we assume

that trait $m$ was historically present in our population, it seems likely that B1 daughters

will have evolved to alter their dispersal decisions in response to the trait's presence.

To model this new situation, we introduce parameter $a$ (Table 1; Figure 3), which

represents the effect of trait $m$ on the probability of B1 daughters dispersing.

When $a > 0$, daughters with trait $m$ are more likely to stay as workers, and when $a =$

$1$ they always stay. Conversely, when $a < 0$, daughters with trait $m$ are more likely to

disperse as reproductives - this area of the parameter space has limited relevance to

the evolution of specialist worker traits, and is explored in Supplementary materials

Parts 2 and 3.

Incorporating parameter $a$ gives us the following equation:

*Equation 4:*

$$f = \frac{(1 - d)v_w + dv_r}{c},$$

$$f' = \frac{[1 - d(1 - a)]v'_w + d(1 - a)v'_r}{c'}, \qquad a \in \{0 \le a \le 1\}$$

306
307

When $a = 0$, the presence of trait $m$ has no effect on the dispersal decisions of

daughters, and Equation 4 reduces to Equation 1. When daughters can respond to the

presence of morphological traits ($a \neq 0$), mothers obtain a degree of indirect control

over daughter behaviour through manipulating their morphology. The extent of this

control is represented by the magnitude of $a$. In order to determine when imparting $m$

will be favoured by selection now that daughters can respond to it behaviourally, we

can substitute Equation 4 into Equation 2 and rearrange to give the parameter space

where we expect manipulation for trait $m$ to invade as:

*Equation 5:*

$$0 < d[c(1-a)(v_r' - v_w') + c'(v_w - v_r)] + cv_w' - c'v_w, \qquad a \in \{0 \leq a \leq 1\}$$

We again find a linear relationship between $d$, the baseline probability of dispersal,

and the likelihood of manipulation for trait $m$ invading (Figure 2). The relationship is

again negative when $m$ confers specialist worker morphology, but the gradient is now

dependant on the value of $a$, such that  $a$ is moderating the relationship between

baseline dispersal and the likelihood of manipulation for trait $m$ invading (Figure 2).

This is because $a$ causes the realised dispersal choices of daughters to correlate with

their morphologies, altering the degree of mismatch between form and function.

Parameter $a$ therefore introduces a mechanism through which mothers can use their

control over daughter morphology to influence daughter behaviour, and consequently

moderate the certainty a mother has about how an individual daughter will behave.

To take an example, trait $m$ might represent a simple reduction in the size of B1

daughters. Smaller daughters will be cheaper to produce (Brand and Chapusiat,

2012), so that if worker performance is independent of size (e.g. Couchoux and Field,

2019), the fitness per unit investment received by a mother from a worker will

increase. However, we might also expect trait $m$ to decrease the fitness per unit

investment from daughters that disperse as reproductives: reproductives must found

new nests and lay eggs, and smaller reproductives may perform less well with these

tasks (Couchoux and Field, 2019).  If there has historically been variation in the size of

daughters independent of trait $m$, perhaps environmentally induced, it seems plausible

that daughters will have already evolved to alter their dispersal decisions according to

their size, with smaller daughters being more likely to stay as workers (Brand and

Chapusiat, 2012). Our value of $a$ in this instance would be positive (since daughters

with trait $m$ are more likely to stay as workers; Supplementary materials Part 3), and

manipulation for trait $m$ will be favoured by selection over a larger range of values of $d$

than if daughters are unable to respond plastically to morphology (see Figure 2).

We can also extend the idea of maternal certainty over offspring behaviour to

instances where workers have the possibility of inheriting the queen role. This

changes the proportion of time a mother expects a daughter to spend as a worker or

reproductive for a given probability of dispersal (Toyoizumi and Field, 2014). Time

spent as an inheriting worker is assumed equivalent to time spent as a reproductive in

the sense that the optimal morphology in both cases is reproductive morphology

(Supplementary materials Parts 1 and 2). Thus the likelihood of a mismatch between

form and function when a mother imparts specialist worker morphology onto a

daughter increases, making specialist worker morphology less likely to be favoured.

However, since time spent as a reproductive or inheriting worker still depends on the

probability of dispersal ($d$), the qualitative nature of the relationship between dispersal

and the parameter space where we might expect to see the evolution of specialist

morphology remains similar (Figure S1).

We note that when $a > 0,$ the sex ratio of dispersing offspring that a mother with the

trait for manipulation produces will be more male biased than the sex ratio of the

offspring produced by a mother without the trait: some of the female offspring which

would have dispersed now stay and help. This will introduce an element of frequency

dependence that is not taken into account in our analysis and would manifest as

changes in $v_r$ relative to $v'_w$. The more common the manipulation trait becomes, the

more male biased the mating pool, lowering the reproductive value of male offspring

and hence the expected fitness of a mother who manipulates her offspring relative to

one who does not. However, so long as the population has a sufficiently large, well

mixed mating pool and an evolutionarily stable sex ratio, this will not impact the

invasion conditions for a rare manipulation trait (Avila and Mullon, 2023).

(***a***) Degree to which daughters change behaviour in presence of trait ***m***

*Figure 3. The relationship between **a**, the degree to which daughters change their dispersal behaviour in the presence of trait **m**, and realised offspring dispersal. Different lines represent different values of baseline dispersal (dispersal in the absence of trait **m**; $0 \leq d \leq 1$) in 0.25 increments. When $a > 0$ daughters are more likely to work (Table 1).*

### 2.3. *Trait benefit as a function of baseline productivity*

The fitness represented by a worker offspring bearing trait $m$ ($v'_w$) is likely to be correlated with the fitness represented by a worker that lacks trait $m$ ($v_w$): trait $m$ represents a single modification of worker morphology, leaving the remaining morphology unchanged. Thus, when the productivity of a worker without trait $m$ changes we expect the productivity of workers with trait $m$ to change as well, so that selection on trait $m$ will depend on $v_w$. Now it may be the case that the fitness benefit conferred by trait $m$ varies with $v_w$ such as in the relationship $v'_w = 2v_w$, where trait $m$ doubles the fitness represented by a worker. Again taking the example of worker size, this would be the case if a large worker is twice as productive as a small worker. The difference in productivity between large and small workers (the fitness benefit of being large) would then be equal to the productivity of a small worker. We use changes in productivity and fitness interchangeably here, but note that the change in a worker's productivity caused by a worker trait may not always be equivalent to the change in the mother's fitness, if the offspring produced are not all equivalent (e.g. if larger workers invest less into producing their own male offspring). Even when changes are equivalent, this may not be apparent in the field, for example if immature or adult workers with a morphological trait have higher mortality, but the productivity of only surviving workers is measured (Cant and Field, 2001; Couvillon and Dornhaus, 2010). Substituting $v'_w$ for $2v_w$ in Equation 5 we can rearrange to give the following:

*Equation 6:*

$$0 < \; v_w(d[\, c' + (a-1)2c] + 2c - c') + d[cv'_r(1-a) - c'v_r], \qquad\qquad a \in \{0 \le a \le 1\}$$

As discussed above, when $v'_w = 2v_w$ we tend to see a strong positive relationship between $v_w$ and the likelihood that manipulation for trait $m$ will invade (Figure 4). However, Equation 6 also highlights how this is not always the case. For example,

since it is the benefit per unit cost which primarily determines whether a trait will be

selected for, if the cost of producing a daughter with trait $m$ ($c' - c$) is very large ($c' >$

$2c$), then we can still generate negative relationships between $v_w$ and the likelihood

that manipulation for trait $m$ will invade, despite setting $v'_w = 2v_w$ (Supplementary

materials Part 3).

More generally we find that if there is a positive relationship between the productivity

benefit of a specialist worker trait and baseline worker productivity; so long as the

increased benefit provided by the trait remains proportionally greater than the cost of

producing it, and if the trait does not significantly increase the likelihood of daughters

dispersing, an increase in baseline worker productivity will lead to increased selection

for that trait (Supplementary materials Part 3). This effect is not dependent on

decoupled control (Supplementary materials Part 2).

*Figure 4. The relationship between the fitness of a worker in the absence of a trait and the likelihood of that trait being selected, when workers with that trait are twice as productive as those without it ($v'_w = 2v_w$). We have plotted this for $0 \leq a \leq 1$ at 0.1 increments with parameter values ($v'_r = 0.8$, $v_r = 1$, $d = 0.5$, $c' = 1$, $c = 1$). Dashed line at y=0 represents the threshold for when manipulators imparting trait $m$ will invade (Equation 2).*

## 3.    Discussion

Our analysis shows that unless a behavioural worker caste has already evolved, a new morphological trait that increases worker fitness but decreases the fitness of reproductives cannot invade, for the simple reason that all offspring will be reproductives. This may seem obvious, but it means that although we might expect queens to use their control over offspring morphology to impose worker behaviour at a lower threshold benefit/cost ratio than the threshold at which offspring themselves would work voluntarily, worker behaviour will not evolve unless the worker's own threshold has already been reached. Thus, contrary to some previous suggestions (Michener and Brothers, 1974; Alexander 1974; Craig, 1979; Kapheim et al., 2015; Nonacs and Denton 2023), maternal manipulation of offspring morphology cannot induce helping behaviour where it was not previously present: the origin of helping is instead constrained by the productivity threshold at which offspring themselves will choose to work voluntarily. We note that in Kapheim et al.'s (2015) model, daughters are assumed to respond to morphological manipulation by staying at their natal nests and helping more frequently i.e. plasticity for helping behaviour is assumed to already be present in the population, and must have evolved historically (as in our second model when $a > 0$).

Our analysis thus suggests that behavioural castes should always precede morphological castes in the social Hymenoptera, and that the evolution of morphological castes relies on provisioners having a high degree of certainty over which offspring will help. If morphological manipulation (of worker traits) cannot be limited to those offspring which are less likely to disperse as reproductives, morphological castes would not be expected to evolve (Supplementary materials Part

4). We do not attempt a systematic review here, but several empirical patterns seem broadly consistent with this. First, behavioural castes are often observed when no morphological caste differentiation has been reported, whereas the opposite is not true. While the possibility that there are subtle morphological differences between castes has not been examined in most apparently facultatively eusocial and cooperatively breeding taxa, there are clear examples of behavioural castes exhibiting no size or other morphological differentiation (e.g. Field 2008; Field et al., 2010). Second, we never see extreme morphological caste specialisation when offspring have a low likelihood of helping, but we do sometimes see limited morphological caste specialisation when offspring have a high likelihood of helping (O'Donnell and Jeanne, 1993; Tibbetts, 2006; Toth et al., 2009), consistent with the evolution of morphological castes being reliant on provisioners having a high degree of certainty of offspring behaviour. An additional pattern we might expect to see is that taxa with morphologically differentiated castes should be nested within lineages that exhibit only behavioural castes. Here, the evidence is mixed. Allodapine bees may represent an example of the predicted pattern (Schwarz et al., 2007), but it has been argued that the phylogeny of vespid wasps indicates that eusociality and morphological castes evolved simultaneously (Piekarski et al., 2018; see also Boomsma 2022 Box 6.20). Our analysis suggests that this is unlikely. One possibility is that morphologically undifferentiated ancestral taxa are now extinct and therefore absent from phylogenetic reconstructions, as is presumably the case for ants, all of which are eusocial. Another possibility is that once unmanipulated offspring began to help, selection favoured mothers conferring morphological specializations on them, which in turn favoured manipulated offspring being even more likely to help (as in our second model, and

discussed in Supplementary materials Part 3). The resulting feedback loop could

make helping and morphological specialization appear to have evolved together.

We have primarily considered queens as being the provisioners controlling the

morphology of offspring. However, once a behavioural caste has been established as

we see in the social Hymenoptera, it becomes increasingly common for the workforce

as a whole to instead control offspring morphology (Cameron and Robinson, 1990;

Inoue et al., 1996; O'Donnell and Jeanne, 1993; Giray et al., 2005). The worker

productivity at which the workforce favours offspring helping will depend on the

number of times the queen has mated and also on the sex ratio of offspring produced

(Reuter and Keller, 2001). However, we would still expect to see a disparity between

workforce and offspring over the worker productivity at which helping is preferred

(Stubblefield and Charnov, 1986).

More importantly, our findings suggest that in the social Hymenoptera, where it is likely

that provisioners control offspring morphology while offspring control their own

behaviour (Reinhold, 2002; Karsai and Hunt, 2002; Kapheim et al., 2011; Kovacs et

al., 2010; Brand and Chapusiat, 2012; Kapheim et al., 2015), the certainty that

provisioners have about offspring dispersal decisions is a key constraint on the

evolution of increasingly differentiated morphological castes (Equation 3). Reduced

certainty results in greater mismatch between form and function. These conclusions

are expected to hold even when individuals can switch behavioural castes within a

lifetime (Supplementary materials Part 2). It is important to emphasise that the

certainty provisioners have over offspring dispersal decisions can be increased

through mechanisms other than a direct offspring response to morphology, for

example where there is seasonal variation in offspring dispersal. A clear example of

this is in temperate cooperatively breeding bees and wasps, where daughters from the

first brood (B1) are more likely to become helpers than daughters from the second

brood (B2), which always disperse as reproductives (section 2.1). Provisioners

therefore have greater certainty about daughter behaviour, and parental manipulation

of offspring morphology might evolve if it could be limited to first brood offspring. The

ability for offspring to adjust their dispersal decisions as a response to their

morphology can also give provisioners a significant degree of indirect control over the

behaviour of female offspring and allow morphological traits to be selected for at lower

baseline rates of dispersal than we might otherwise expect (Equation 5).

*Baseline worker productivity can affect the likelihood of specialist morphology evolving*

We have shown that the baseline productivity of workers (in the absence of

specialised morphology) can be important in determining when specialist morphology

is likely to evolve. The gain in worker productivity (and hence provisioner fitness)

provided by a morphological trait may vary as a function of baseline worker

productivity, an example being a trait that increases worker size, where larger workers

are always twice as productive as smaller workers. In this case, when smaller workers

have a productivity value of 5, larger workers would have a productivity of 10 (5

greater), but when smaller workers have a productivity value of 7, larger workers have

a productivity of 14 (7 greater). A trait causing larger worker size will then be most

likely to evolve when baseline worker productivity is high, since this is when larger

workers provide the largest gain. Generally, if the extra fitness provided by a

specialised morphological trait increases with baseline worker productivity (linearly or

otherwise), the trait is most likely to be favoured when baseline productivity is high -

unless that trait is especially costly (Supplementary materials Part 3). In practice,

determining the relationship between morphology and productivity can be complex,

with morphological traits not always enhancing all worker tasks equally, and epistasis

for fitness (Bank, 2022) often making the extraction of simple empirical correlations

difficult (Grimm et al., 2005; Kugonza et al., 2011; Wen et al., 2015).

*Decoupled control, reversibility, and loss of conflict between offspring and provisioners*

We now extend our analysis by considering how it might relate to the evolutionary

reversibility of morphological castes. It has previously been suggested that the

presence of a morphologically specialised worker caste, in particular a sterile worker

caste, represents a state of irreversibility wherein the worker caste can no longer be

lost evolutionarily (Boomsma 2013; Boomsma and Gawne 2018).  The perspective we

have taken on control structure reveals a mechanism that could underlie reduced

reversibility: with decoupled control, morphological castes become less likely to be lost

as they become more specialised, and conflict between offspring and provisioners

over offspring caste fate may disappear (see Bourke 2011: 178 for a related

discussion).

After a worker caste has evolved, it might be lost when the environment changes so

that worker productivity decreases. Indeed, phylogenetic work shows that the largely

behavioural worker caste seen in sweat bees has frequently been lost (Wcislo and

Danforth, 1997), probably when bees entered environments where the season was

too short for workers to be productive (Lucas and Field, 2013). Once a morphological

worker caste controlled by provisioners is present, however, we expect workers to

evolve to respond to their morphology when making decisions about their behaviour

(section 2.2). Figure 5 illustrates the implications of this. For simplicity, we focus on the

case when offspring with specialist worker or reproductive morphology are equally

costly to produce – an example might be where worker morphology simply involves

mandibles shaped specifically for defensive or foraging tasks. Initially, offspring prefer

to disperse at higher worker productivities than provisioners prefer – productivities at

which provisioners, because they are more closely related to the queen's offspring

than to her grand-offspring, would prefer them to work (Figure 5A; see Introduction).

With each additional morphological specialisation, however, the behavioural caste that

an offspring chooses should be constrained more and more by its morphology:

offspring will incur larger and larger costs if they mismatch their behaviour to their

morphology. The effect is almost ratchet-like: each additional morphological

specialism reduces the fitness of daughters that mismatch their behaviour to their

morphology by dispersing, so that worker productivity must drop progressively lower

for offspring with worker morphology to nevertheless prefer dispersal (Figure 5B).

In contrast, because provisioners control offspring morphology, the worker productivity

at which provisioners prefer offspring to disperse remains constant (figure 5): if the

fitness that provisioners obtain from workers with specialist worker morphology falls

*Figure 5. The relationship between the level of worker morphological specialisation and the threshold worker productivity at which offspring dispersal is favoured by two parties, offspring and provisioners. Preferences under decoupled control are represented by shaded (bold-edged) and unshaded boxes. The worker productivities at which offspring will begin to disperse, so that the worker caste is lost, are represented by shaded/bold-edged boxes. Red arrows linked to faded boxes indicate the possibility that offspring regain control of their morphology, recoupling it with control of behaviour, so that they do not incur mismatch costs when dispersing.*

below what they obtain from dispersers with reproductive morphology, provisioners

are expected to impart reproductive morphology – offspring should respond to this by

dispersing, thus avoiding any mismatch. Eventually, once morphological castes are

sufficiently specialised, the ratchet-like effect will mean that the worker productivity at

which offspring with worker morphology themselves prefer dispersal drops below that

at which provisioners prefer their offspring to disperse (Figure 5C). It should then be

provisioners, not offspring, that first cause the loss of a worker caste during evolution,

should worker productivity drop sufficiently. The worker caste is then no longer 'self-

determined' (Bourke and Ratnieks, 1999) by the offspring and can instead be

considered to be 'manipulator enforced' by the provisioner. At this point, realised

conflict between offspring and provisioners over dispersal is lost, with both parties

preferring the same outcome (the dispersal of offspring).

If at any point individual offspring are able to regain control of their morphology during

evolution (recoupling control over morphology and behaviour), as seems to be the

case in stingless bees (Meliponini) and a few other obligately eusocial taxa (Ishay and

Landau, 1972; Wenseleers et al., 2003; Kaptein et al., 2005; Mas and Kolliker, 2008),

the threshold worker productivity required for a worker caste to be lost becomes wholly

determined by individual offspring (returning the worker caste to being 'self-

determined'; red arrows in Figure 5). Offspring can then choose whether to work or

disperse as reproductives without incurring costs through mismatching morphology

and behaviour. This too represents a loss of realised conflict over dispersal, since now

only one party is able to influence the outcome, this time resulting in dispersal being

preferred at a higher level of worker productivity. The apparently offspring-controlled

caste determination seen in the Meliponini (Wenseleers et al., 2003; Wenseleers and

Ratnieks, 2004) could therefore represent a vulnerability to selection for loss of the

worker caste, although so long as other aspects of meliponine life history remain, such

as swarm founding that requires workers, such a reversal seems unlikely.

*Extreme worker traits and evolutionary irreversibility*

It is useful to consider where extreme morphological specializations such as worker

sterility might appear in the above sequence (Figure 5). The very high costs of

mismatch that such specializations entail means that they should either be associated

with a recoupling of control over morphology and behaviour, such that mismatch can

be avoided, or with a manipulator enforced worker caste. This is because, if control

has not been recoupled and mismatch cannot be avoided, we would expect less

extreme specialisations, with lower costs of mismatch, to evolve before extreme traits

like sterility – when provisioners are less certain about offspring behaviour. These less

extreme traits might be sufficient on their own to cause the worker caste to become

manipulator enforced (Figure 5C). Additional, more extreme morphological

specialisations such as sterility would not then further reduce the likelihood of a worker

caste being lost, since the point at which workers themselves might prefer to disperse

would already have been passed (Figures 5C and 5D).

In the previous section, we suggested that the worker caste is less likely to be lost as it

becomes more specialised. We here consider more generally what evolutionary

‘irreversibility’ means in the context of eusociality (Boomsma and Gawne, 2018; see

also Nonacs and Denton 2023). If reproductives are dependent on help from a worker

caste to produce reproductive offspring – for example if the life cycle requires swarm

founding – then that worker caste (morphologically specialised or otherwise) cannot be

lost evolutionarily following a reduction in worker productivity, without resulting in

extinction (Crespi and Yanega, 1995).  It might then seem reasonable to consider the

worker caste evolutionarily irreversible at this point. Irreversibility for the worker caste

is then conditional on the capabilities of individuals of the reproductive caste, not on

how morphologically specialised the worker caste itself has become. However, the

capabilities of individuals in the reproductive caste could change – for example if the

environment changes so that swarm founding is no longer essential. Likewise, if

workers are unable to functionally replace reproductives in producing both male and

female reproductive offspring, so long as worker capabilities remaining unchanged the

reproductive caste cannot be lost evolutionarily without resulting in extinction. Thus,

both of the criteria for obligate eusociality (Crespi and Yanega, 1995) must be

maintained for eusociality to be considered evolutionarily irreversible, at least at a

given point in time: workers must be unable to functionally replace reproductives, and

reproductives must be unable to produce reproductive offspring without help from a

worker caste. We note that while being strongly correlated with morphological

specialisation (Bourke, 2011; Quinones and Pen, 2017; Boomsma and Gawne, 2018),

both criteria could in principle be maintained entirely through ecological conditions and

life history traits, such as when workers lack mating opportunities (inability for workers

to functionally replace reproductives) and swarm founding is favoured (reliance of

reproductives on the worker caste to produce reproductive offspring).

*The prevalence of decoupled control*

While we have focused primarily on the evolution of a morphological worker caste in

the facultatively eusocial and cooperatively breeding Hymenoptera, the relationships

we have discussed underlying the evolution of morphological traits will hold for any

level of social organisation or study system so long as our assumptions regarding the

immutability of morphological castes and split control are met (Supplementary

materials Part 2). Our assumption that provisioners control the morphology of offspring

may not be equally valid for all morphological traits (e.g. body size vs body shape),

and must be evaluated on a trait by trait basis. Relaxing these assumptions represents

a possible future extension of our ideas and would greatly generalise our results,

though it seems likely that the key linear relationships we identify involving behavioural

certainty and baseline productivity will remain important. Shared control seems

especially likely in progressively provisioning taxa, with substantial evidence that

offspring begging influences provisioning decisions in obligately eusocial

Hymenoptera (Ishay and Landau, 1972; Creemers et al., 2003; Kaptein et al., 2005; den Boer and Duchateau, 2006; Mas and Kolliker, 2008; He et al., 2016). However, evidence for offspring resistance to provisioner manipulation in the Hymenoptera is limited outside of obligately eusocial species (Ishay and Landau, 1972; Wenseleers and Ratnieks, 2004; Kaptein et al., 2005; Mas and Kolliker, 2008; Gonzalez-Forero, 2015). Begging behaviours in solitary, cooperatively breeding, or facultatively eusocial progressively provisioning taxa have not to our knowledge been reported, suggesting that our assumption of queen control over morphology but offspring control of behaviour is reasonable before morphological castes evolve. It is possible, however, that the absence of reported begging outside the obligately eusocial Hymenoptera instead reflects a disparity in the volume of research conducted.

***Conclusions***

(1) There is evidence that in the social Hymenoptera, the evolution of a morphological worker caste is reliant on the prior presence of a behavioural worker caste. (2) We suggest that maternal certainty of offspring dispersal decisions is one of the main constraints on the evolution of a morphological worker caste in the facultatively eusocial and cooperatively breeding Hymenoptera. (3) The decoupling of control over morphology and behaviour could be a key factor maintaining a morphological worker caste, and can lead to reduced conflict between offspring and provisioners. (4) The productivity of workers in the absence of specialist worker morphology can affect the likelihood of specialist morphology being favoured by selection.

## *Bibliography*

Alexander, R. D. (1974). The Evolution of Social Behavior. *Annual review of ecology and systematics, 5*, 325-383. doi:https://doi.org/10.1146/annurev.es.05.110174.001545

Anderson, D. T. (1972). The development of holometabolous insects. In S. J. Counce, & C. H. Waddington (Eds.), *Developmental systems: Insects* (Vol. 1, pp. 165-242). Academic Press.

Avila, P., & Mullon, C. (2023). Evolutionary game theory and the adaptive dynamics approach: adaptation where individuals interact. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences, 378*, 20210502. doi:10.1098/rstb.2021.0502

Bank, C. (2022). Epistasis and adaptation on fitness landscapes. *Annual Review of Ecology, Evolution, and Systematics, 53*, 457-479. doi:https://doi.org//10.1146/annurev-ecolsys-102320-112153

Beekman, M., & Oldroyd, B. P. (2019). Conflict and major transitions — why we need true queens. *Current Opinion in Insect Science, 34*, 73-79. doi:10.1016/j.cois.2019.03.009

Boomsma, J. J. (2013). Beyond promiscuity: mate-choice commitments in social breeding. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences, 368*, 20120050. doi:https://doi.org//10.1098/rstb.2012.0050

Boomsma, J. J. (2022). *Domains and Major Transitions of Social Evolution.* New York: Oxford University Press. Retrieved February 26, 2024

Boomsma, J. J., & Gawne, R. (2018). Superorganismality and caste differentiation as points of no return: how the major evolutionary transitions were lost in translation. *Biological Reviews of the Cambridge Philosophical Society, 93*, 28-54. doi:https://doi.org/10.1111/brv.12330

Bourke, A. F. (1999). Colony size, social complexity and reproductive conflict in social insects. *Journal of evolutionary biology, 12*, 245-257. doi:https://doi.org/10.1046/j.1420-9101.1999.00028.x

Bourke, A. F. (2011). *Principles of social evolution.* Oxford University Press. doi:https://doi.org//10.1093/acprof:oso/9780199231157.001.0001

Bourke, A. F. (2023). Conflict and conflict resolution in the major transitions. *Proceedings. Biological Sciences / the Royal Society, 290*, 20231420. doi:10.1098/rspb.2023.1420

Bourke, A. F., & Ratnieks, F. L. (1999). Kin conflict over caste determination in social Hymenoptera. *Behavioral Ecology and Sociobiology, 46*, 287-297. doi:https://doi.org//10.1007/s002650050622

Brand, N., & Chapuisat, M. (2012). Born to be bee, fed to be worker? The caste system of a primitively eusocial insect. *Frontiers in zoology, 9*, 35. doi:https://doi.org//10.1186/1742-9994-9-35

Cameron, S. A., & Robinson, G. E. (1990). Juvenile hormone does not affect division of labor in bumble bee colonies (hymenoptera: apidae). *Annals of the Entomological Society of America, 83*, 626-631. doi:https://doi.org//10.1093/aesa/83.3.626

Cant, M. A., & Field, J. (2001). Helping effort and future fitness in cooperation animal societies. *Proceedings. Biological Sciences / the Royal Society, 268*, 1959-1964. doi:https://doi.org//10.1098/rspb.2001.1754

Charnov, E. L. (1978). Evolution of eusocial behavior: offspring choice or parental parasitism? *Journal*
*of Theoretical Biology, 75*, 451-465. doi:https://doi.org//10.1016/0022-5193(78)90356-9

Couchoux, C., & Field, J. (2019). Parental manipulation of offspring size in social groups: a test using
paper wasps. *Behavioral Ecology and Sociobiology, 73*, 36.
doi:https://doi.org//10.1007/s00265-019-2646-3

Couvillon, M. J., & Dornhaus, A. (2010). Small worker bumble bees (Bombus impatiens) are hardier
against starvation than their larger sisters. *Insectes sociaux, 57*, 193-197.
doi:https://doi.org//10.1007/s00040-010-0064-7

Craig, R. (1979). Parental manipulation, kin selection, and the evolution of altruism. *Evolution, 33*, 319.
doi:https://doi.org//10.2307/2407622

Creemers, B., Billen, J., & Gobin, B. (2003). Larval begging behaviour in the ant Myrmica rubra.
*Ethology Ecology & Evolution, 15*, 261-272.
doi:https://doi.org//10.1080/08927014.2003.9522671

Crespi, B. J., & Yanega, D. (1995). The definition of eusociality. *Behavioral Ecology, 6*, 109-115.
doi:https://doi.org//10.1093/beheco/6.1.109

Dawkins, R. (1976). *The Selfish Gene.* Oxford University Press.

den Boer, S. P., & Duchateau, M. J. (2006). A larval hunger signal in the bumblebee Bombus terrestris.
*Insectes sociaux, 53*, 369-373. doi:https://doi.org//10.1007/s00040-006-0883-8

El Mouden, C., & Gardner, A. (2008). Nice natives and mean migrants: the evolution of dispersal-
dependent social behaviour in viscous populations. *Journal of Evolutionary Biology, 21*, 1480-
1491. doi:https://doi.org//10.1111/j.1420-9101.2008.01614.x

Ferguson-Gow, H., Sumner, S., Bourke, A. F., & Jones, K. E. (2014). Colony size predicts division of
labour in attine ants. *Proceedings. Biological Sciences / the Royal Society, 281*.
doi:10.1098/rspb.2014.1411

Field, J. (2008). The ecology and evolution of helping in hover wasps (hymenoptera: stenogastrinae). In
J. Korb, & J. Heinze (Eds.), *Ecology of social evolution* (pp. 85-107). Berlin, Heidelberg: Springer
Berlin Heidelberg. doi:10.1007/978-3-540-75957-7\_4

Field, J., & Toyoizumi, H. (2020). The evolution of eusociality: no risk-return tradeoff but the ecology
matters. *Ecology Letters, 23*, 518-526. doi:https://doi.org//10.1111/ele.13452

Field, J., Paxton, R. J., Soro, A., & Bridge, C. (2010). Cryptic plasticity underlies a major evolutionary
transition. *Current Biology, 20*, 2028-2031. doi:10.1016/j.cub.2010.10.020

Giray, T., Giovanetti, M., & West-Eberhard, M. J. (2005). Juvenile hormone, reproduction, and worker
behavior in the neotropical social wasp Polistes canadensis. *Proceedings of the National*
*Academy of Sciences of the United States of America, 102*, 3330-3335.
doi:https://doi.org//10.1073/pnas.0409560102

González-Forero, M. (2015). Stable eusociality via maternal manipulation when resistance is costless.
*Journal of Evolutionary Biology, 28*, 2208-2223. doi:https://doi.org//10.1111/jeb.12744

González-Forero, M., & Peña, J. (2021). Eusociality through conflict dissolution. *Proceedings. Biological*
*Sciences / the Royal Society, 288*, 20210386. doi:https://doi.org//10.1098/rspb.2021.0386

Grafen, A. (1985). A geometric view of relatedness. *Oxford surveys in evolutionary biology, 2*, 28-89.

Grimm, L. H., Kelly, S., Krull, R., & Hempel, D. C. (2005). Morphology and productivity of filamentous
743            fungi. *Applied Microbiology and Biotechnology, 69*, 375-384.
744            doi:https://doi.org//10.1007/s00253-005-0213-5

745      Hamilton, W. D. (1964). The genetical evolution of social behaviour. I. *Journal of Theoretical Biology, 7*,
746            1-16. doi:https://doi.org//10.1016/0022-5193(64)90038-4

747      He, X. J., Zhang, X. C., Jiang, W. J., Barron, A. B., Zhang, J. H., & Zeng, Z. J. (2016). Starving honey bee
748            (Apis mellifera) larvae signal pheromonally to worker bees. *Scientific Reports, 6*, 22359.
749            doi:https://doi.org//10.1038/srep22359

750      Hunt, J. H. (2007). *The Evolution of Social Wasps.* Oxford University Press.

751      Inoue, T., Salmah, S., & Sakagami, S. (1996). Individual Variations in Worker Polyethism of the
752            Sumatran Stingless Bee, Trigona (Tetragonula) minangkabau (Apidae, Meliponinae). *Japanese
753            journal of entomology, 64*, 641-668.

754      Ishay, J., & Landau, E. M. (1972). Vespa larvae send out rhythmic hunger signals. *Nature, 237*, 286-287.
755            doi:https://doi.org//10.1038/237286a0

756      Jeanne, R. L., Loope, K. J., Bouwma, A. M., Nordheim, E. V., & Smith, M. L. (2022). Five decades of
757            misunderstanding in the social Hymenoptera: a review and meta-analysis of Michener's
758            paradox. *Biological Reviews of the Cambridge Philosophical Society, 97*, 1559-1611.
759            doi:https://doi.org//10.1111/brv.12854

760      Kapheim, K. M., Bernal, S. P., Smith, A. R., Nonacs, P., & Wcislo, W. T. (2011). Support for maternal
761            manipulation of developmental nutrition in a facultatively eusocial bee, Megalopta genalis
762            (Halictidae). *Behavioral Ecology and Sociobiology, 65*, 1179-1190.
763            doi:https://doi.org//10.1007/s00265-010-1131-9

764      Kapheim, K. M., Nonacs, P., Smith, A. R., Wayne, R. K., & Wcislo, W. T. (2015). Kinship, parental
765            manipulation and evolutionary origins of eusociality. *Proceedings. Biological Sciences / the
766            Royal Society, 282*, 20142886. doi:https://doi.org//10.1098/rspb.2014.2886

767      Kaptein, N., Billen, J., & Gobin, B. (2005). Larval begging for food enhances reproductive options in the
768            ponerine ant Gnamptogenys striatula. *Animal Behaviour, 69*, 293-299.
769            doi:https://doi.org//10.1016/j.anbehav.2004.04.012

770      Karsai, I., & Hunt, J. H. (2002). Food Quantity AffectTraits of Offspring in the Paper Wasp Polistes
771            metricus (Hymenoptera: Vespidae). *Environmental Entomology, 31*, 99-106.
772            doi:https://doi.org//10.1603/0046-225X-31.1.99

773      Keller, L., & Nonacs, P. (1993). The role of queen pheromones in social insects: queen control or queen
774            signal? *Animal Behaviour, 45*, 787-794. doi:https://doi.org//10.1006/anbe.1993.1092

775      Kerr, N. Z., Crone, E. E., & Williams, N. M. (2019). Integrating vital rates explains optimal worker size
776            for resource return by bumblebee workers. *Functional Ecology, 33*, 467-478.
777            doi:https://doi.org//10.1111/1365-2435.13251

778      Korb, J. (2008). The ecology of social evolution in termites. In J. Korb, & J. Heinze (Eds.), *Ecology of
779            social evolution* (pp. 151-174). Berlin, Heidelberg: Springer Berlin Heidelberg.
780            doi:https://doi.org/10.1007/978-3-540-75957-7

Kovacs, J. L., Hoffman, E. A., Marriner, S. M., Rekau, J. A., & Goodisman, M. A. (2010). Environmental
and genetic influences on queen and worker body size in the social wasp Vespula maculifrons.
*Insectes sociaux, 57*, 53-65. doi:https://doi.org//10.1007/s00040-009-0050-0

Kugonza, D. R., Nabasirye, M., Mpairwe, D., Hanotte, O., & Okeyo, A. M. (2011). Productivity and
morphology of Ankole cattle in three livestock production systems in Uganda. *Animal Genetic
Resources/Ressources génétiques animales/Recursos genéticos animales, 48*, 13-22.
doi:https://doi.org//10.1017/S2078633611000038

Lehmann, L., & Rousset, F. (2020). When do individuals maximize their inclusive fitness? *The American
Naturalist, 195*, 717-732. doi:https://doi.org//10.1086/707561

Liu, M., West, S. A., & Wild, G. (2022). The evolution of manipulative cheating. *eLife, 11*.
doi:https://doi.org//10.7554/eLife.80611

Lucas, E. R., & Field, J. (2013). Caste determination through mating in primitively eusocial societies.
*Journal of Theoretical Biology, 335*, 31-39. doi:https://doi.org//10.1016/j.jtbi.2013.05.032

Mas, F., & Kölliker, M. (2008). Maternal care and offspring begging in social insects: chemical
signalling, hormonal regulation and evolution. *Animal Behaviour, 76*, 1121-1131.
doi:https://doi.org//10.1016/j.anbehav.2008.06.011

Michener, C. D. (1964). Reproductive efficiency in relation to colony size in hymenopterous societies.
*Insectes sociaux, 11*, 317-341. doi:https://doi.org//10.1007/BF02227433

Michener, C. D., & Brothers, D. J. (1974). Were workers of eusocial hymenoptera initially altruistic or
oppressed? *Proceedings of the National Academy of Sciences of the United States of America,
71*, 671-674. doi:https://doi.org//10.1073/pnas.71.3.671

Nonacs, P., & Denton, K. K. (2023). Eusociality is not a major evolutionary transition, and why that
matters. *Insectes sociaux*. doi:10.1007/s00040-023-00942-3

Nowak, M. A., Tarnita, C. E., & Wilson, E. O. (2010). The evolution of eusociality. *Nature, 466*, 1057-
1062. doi:https://doi.org//10.1038/nature09205

O'Donnell, S., & Jeanne, R. L. (1993). Methoprene accelerates age polyethism in workers of a social
wasp (Polybia occidentalis). *Physiological entomology, 18*, 189-194.
doi:https://doi.org//10.1111/j.1365-3032.1993.tb00467.x

Oster, G. F., & Wilson, E. O. (1978). *Caste and Ecology in the Social Insects. (MPB-12), Volume 12.*
Princeton University Press. Retrieved February 26, 2024, from
https://www.degruyter.com/document/doi/10.12987/9780691209326/html

Peters, R. S., Krogmann, L., Mayer, C., Donath, A., Gunkel, S., Meusemann, K., . . . Niehuis, O. (2017).
Evolutionary history of the hymenoptera. *Current Biology, 27*, 1013-1018.
doi:https://doi.org//10.1016/j.cub.2017.01.027

Piekarski, P. K., Carpenter, J. M., Lemmon, A. R., Moriarty Lemmon, E., & Sharanowski, B. J. (2018).
Phylogenomic evidence overturns current conceptions of social evolution in wasps (vespidae).
*Molecular Biology and Evolution, 35*, 2097-2109. doi:10.1093/molbev/msy124

Poitrineau, K., Mitesser, O., & Poethke, H. J. (2009). Workers, sexuals, or both? Optimal allocation of
resources to reproduction and growth in annual insect colonies. *Insectes sociaux, 56*, 119-129.
doi:https://doi.org//10.1007/s00040-009-0004-6

Powell, S. (2016). A comparative perspective on the ecology of morphological diversification in
complex societies: nesting ecology and soldier evolution in the turtle ants. *Behavioral Ecology*
*and Sociobiology, 70*, 1075-1085. doi:https://doi.org//10.1007/s00265-016-2080-8

Powell, S., Price, S. L., & Kronauer, D. J. (2020). Trait evolution is reversible, repeatable, and decoupled
in the soldier caste of turtle ants. *Proceedings of the National Academy of Sciences of the*
*United States of America, 117*, 6608-6615. doi:https://doi.org//10.1073/pnas.1913750117

Quiñones, A. E., & Pen, I. (2017). A unified model of Hymenopteran preadaptations that trigger the
evolutionary transition to eusociality. *Nature Communications, 8*, 15920.
doi:https://doi.org//10.1038/ncomms15920

Reinhold, K. (2002). Maternal effects and the evolution of behavioral and morphological characters: a
literature review indicates the importance of extended maternal care. *The Journal of Heredity,*
*93*, 400-405. doi:https://doi.org//10.1093/jhered/93.6.400

Reuter, M., & Keller, L. (2001). Sex ratio conflict and worker production in eusocial hymenoptera. *The*
*American Naturalist, 158*, 166-177. doi:https://doi.org//10.1086/321311

Rodrigues, A. M., & Gardner, A. (2022). Reproductive value and the evolution of altruism. *Trends in*
*Ecology & Evolution, 37*, 346-358. doi:10.1016/j.tree.2021.11.007

Russell, A. F., & Lummaa, V. (2009). Maternal effects in cooperative breeders: from hymenopterans to
humans. *Philosophical Transactions of the Royal Society of London. Series B, Biological*
*Sciences, 364*, 1143-1167. doi:https://doi.org//10.1098/rstb.2008.0298

Schwander, T., Humbert, J.-Y., Brent, C. S., Cahan, S. H., Chapuis, L., Renai, E., & Keller, L. (2008).
Maternal effect on female caste determination in a social insect. *Current Biology, 18*, 265-269.
doi:https://doi.org//10.1016/j.cub.2008.01.024

Schwarz, M. P., Richards, M. H., & Danforth, B. N. (2007). Changing paradigms in insect social
evolution: insights from halictine and allodapine bees. *Annual Review of Entomology, 52*, 127-
150. doi:10.1146/annurev.ento.51.110104.150950

Seger, J. (1983). Partial bivoltinism may cause alternating sex-ratio biases that favour eusociality.
*Nature, 301*, 59-62. doi:https://doi.org//10.1038/301059a0

Stubblefield, J. W., & Charnov, E. L. (1986). Some conceptual issues in the origin of eusociality.
*Heredity, 57 ( Pt 2)*, 181-187. doi:https://doi.org//10.1038/hdy.1986.108

Szathmáry, E., & Smith, J. M. (1995). The major evolutionary transitions. *Nature, 374*, 227-232.
doi:https://doi.org//10.1038/374227a0

Tibbetts, E. A. (2006). Badges of status in worker and gyne Polistes dominulus wasps. *Annales zoologici*
*Fennici, 43*, 575-582. Retrieved March 5, 2024, from https://www.jstor.org/stable/23736763

Toth, A. L., Bilof, K. B., Henshaw, M. T., Hunt, J. H., & Robinson, G. E. (2009). Lipid stores, ovary
development, and brain gene expression in Polistes metricus females. *Insectes sociaux, 56*, 77-
84. doi:10.1007/s00040-008-1041-2

Toyoizumi, H., & Field, J. (2014). Dynamics of social queues. *Journal of Theoretical Biology, 346*, 16-22.
doi:https://doi.org//10.1016/j.jtbi.2013.12.017

Trible, W., & Kronauer, D. J. (2017). Caste development and evolution in ants: it's all about size.
*Journal of Experimental Biology, 220*, 53-62. doi:https://doi.org//10.1242/jeb.145292

Uller, T., & Pen, I. (2011). A theoretical model of the evolution of maternal effects under parent-
offspring conflict. *Evolution, 65*, 2075-2084. doi:https://doi.org//10.1111/j.1558-
5646.2011.01282.x

Wade, M. J. (2001). Maternal effect genes and the evolution of sociality in haplo-diploid organisms.
*Evolution, 55*, 453-458. doi:https://doi.org//10.1554/0014-
3820(2001)055[0453:megate]2.0.co;2

Wcislo, W. T., & Danforth, B. N. (1997). Secondarily solitary: the evolutionary loss of social behavior.
*Trends in Ecology & Evolution, 12*, 468-474. doi:https://doi.org//10.1016/s0169-
5347(97)01198-1

Wen, W., Li, K., Alseekh, S., Omranian, N., Zhao, L., Zhou, Y., . . . Fernie, A. R. (2015). Genetic
determinants of the network of primary metabolism and their relationships to plant
performance in a maize recombinant inbred line population. *The Plant Cell, 27*, 1839-1856.
doi:https://doi.org//10.1105/tpc.15.00208

Wenseleers, T., & Ratnieks, F. L. (2004). Tragedy of the commons in Melipona bees. *Proceedings.*
*Biological Sciences / the Royal Society, 271 Suppl 5*, S310-S312.
doi:https://doi.org//10.1098/rsbl.2003.0159

Wenseleers, T., Ratnieks, F. L., & Billen, J. (2003). Caste fate conflict in swarm-founding social
hymenoptera: an inclusive fitness analysis. *Journal of Evolutionary Biology, 16*, 647-658.
doi:https://doi.org//10.1046/j.1420-9101.2003.00574.x