The allotetraploid *Nicotiana tabacum* genome and GenBank genomics highlight the genomic features, genetic diversity and regulation of morphological, metabolic and disease-resistance traits

Yanjun Zan[1#], Shuai Chen[1#], Min Ren[1#], Guoxiang Liu[1#], Yutong Liu[1#], Huan Si[1], Zhengwen Liu[1], Dan Liu[1], Xingwei Zhang[1], Ying Tong[1], Yuan Li[1], Caihong Jiang[1], Liuying Wen[1], Zhiliang Xiao[1], Yangyang Sun[1], Ruimei Geng[1], Quanfu Feng[1], Yuanying Wang[1], Guoyou Ye[2,3], Lingzhao Fang[4], Yong Chen[5]*, Lirui Cheng[1]*, Aiguo Yang[1]*

[1]Tobacco Research Institute, Chinese Academy of Agricultural Sciences, Qingdao, 266000, China

[2]CAAS-IRRI Joint Laboratory for Genomics-assisted Germplasm Enhancement, Agricultural Genomics Institute in Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen 518120, China.

[3]Strategic Innovation Platform, International Rice Research Institute, Metro Manila, 1301, Philippines.

[4]Center for Quantitative Genetics and Genomics, Aarhus University, Aarhus, 8000, Denmark.

[5]China National Tobacco Corporation, Beijing, 100045, China

[#]These authors contributed equally

[*]Corresponding authors

Aiguo Yang

Email: yangaiguo@caas.cn

Lirui Cheng

Email: chenglirui@caas.cn

Yong Chen

Email: chenyong@tobacco.gov.cn

Abstract:

Nicotiana tabacum is a model organism in plant molecular and pathogenic research and has
significant potential in the production of biofuels and active pharmaceutical compounds in
synthetic biology. Because of the large allotetraploid genome of tobacco, its genomic features,
genetic diversity and genetic regulation of many complex traits remain unknown. In this study,
we present a nearly complete chromosome-scale assembly of *N. tabacum* and provide
evidence that homoeologous exchange between subgenomes and epigenetic remodelling are
likely mechanisms of genome stabilization and subgenome coordination following
polyploidization. By leveraging GenBank-scale sequencing and phenotyping data from 5196
lines, geography at the continent scale, rather than types assigned on the basis of curing crop
practices, was found to be the most important correlate of genetic structure. Using 178
marker□trait associations detected in genome-wide association analysis, a reference
genotype-to-phenotype map was built for 39 morphological, developmental, and disease-
resistance traits. A novel gene, auxin response factor 9 (*Arf9*), associated with wider leaves
after being knocked out, was fine-mapped to a single nucleotide polymorphism (SNP). This
point mutation alters the translated amino acid from Ala_{203} to Pro_{203}, likely preventing
homodimer formation during DNA binding. Our analysis also revealed signatures of positive
and polygenic selection for multiple traits during the process of selective breeding. Overall,
this study demonstrated the power of leveraging GenBank genomics to gain insights into the
genomic features, genetic diversity, and regulation of complex traits in *N. tabacum*, laying a
foundation for future research on plant functional genomics, crop breeding, and the
production of biopharmaceuticals and biofuels.

Introduction

Nicotiana species have served as models for a series of cytogenetic and molecular genetic
breakthroughs, including the discovery of protoplast fusion[1], the introduction of foreign
genetic material through interspecies hybridization[2,3], T-DNA mediated plant transformation[4],
virus-induced genetic transformation[5], the first edible vaccine[6,7], and plant pathogen studies[8].
The rich diversity in plant architecture, leaf morphology, flowering time and metabolic traits,
ease of genetic manipulation, and fairly high number of curated pathways in *Nicotiana*
tabacum highlight its central importance in plant development research and its significant
potential in the field of synthetic biology[6,7].

Nicotiana tabacum (2n=4x=48, ~4.30 Gb) is an interspecific hybrid derived from two
progenitors, *Nicotiana sylvestris* (2n=2x=24, 2.59 Gb) and *Nicotiana tomentosiformis*
(2n=2x=24, 2.22 Gb) approximately 0.2 Mya ago[9-11]. Several attempts have been made to
assemble its large allotetraploid genome; however, the latest assembly included only 81% of
the genomic sequences and was very fragmented[9,11,12]. As a consequence of the large and
repetitive polyploid genome, obtaining a chromosome-level assembly constitutes a significant
challenge, as there are difficulties not only in bridging highly repetitive repeat regions but
also in computationally disentangling homologous sequences with high similarity from two
progenitor species[9,11,12]. In the absence of a high-quality genome assembly and with nearly all
genetic studies being performed in the private sector, the genomic features, global genetic
diversity and genetic basis of many important traits in this model plant remain poorly
understood. These factors highlight the necessity for a complete genome assembly and
GenBank-scale holistic investigation of the genetic diversity and regulation of complex traits
that are highly relevant for future plant functional genomics, crop improvement, and plant
synthetic biology research.

In this study, we present the first complete chromosome-level assembly of the allotetraploid
Nicotiana tabacum genome and genetic and phenotypic data for an entire *Nicotiana tabacum*
GenBank with 5196 germplasms hosted at the Chinese Academy of Agriculture Sciences. The
complete genome assembly demonstrated that chromosome rearrangement between
subgenomes and epigenetic remodeling are likely mechanisms in genome stabilization and

subgenome coordination following polyploidization. By utilizing this valuable resource,

insights were gained into the genomic features, genetic diversity and genetic regulation of

plant architecture, morphological, metabolic, and disease-resistance traits in this model

species. A novel gene, auxin response factor 9 (*Arf9*), resulting in wider leaves after being

knocked out, was fine-mapped to a single nucleotide polymorphism (SNP) altering the

translated amino acid from Ala_{203} to Pro_{203}, which in turn prevented the formation of a

homodimer during DNA binding. Results from our study, together with the released data and

seeds, will likely accelerate aspects of tobacco research that were previously hindered by the

complexity of the polyploid genome and a lack of publicity for research driven by the private

sector, with benefits in plant functional genomics, crop improvement research, and the

production of biopharmaceutical compounds and biofuel.

Results

Chromosome-scale assembly and genomic features of the allotetraploid Nicotiana tabacum

genome

Based on k-mer analysis with 52X Illumina short reads and flow cytometry analysis, the *N.*

tabacum genome was estimated to be 4.29-4.30 Gb, representing a downsizing of

approximately 10% from the sum of the two parental genomes (Fig 1A, Supplementary Fig

S1). To overcome the challenge of assembling a large polyploid genome, we generated 52X

Illumina short reads, 123.35 X PacBio Sequel II reads, 124.93X 10X genomic linked reads,

120.12 X Hi-C reads and 192.27 X Bionano reads from *N. tabacum L.* var *ZY300* and adopted

a hybrid assembly approach (Materials and methods). The final assembly included 4.17 Gb

(Fig 1B) of sequences with contig N50 at 27.17 Mb, and 96.98% of the sequences were

anchored to 24 pseudochromosomes (Supplementary Table 1, Supplementary Fig S2). This

represented a 10-15 percent increase in the assembled genome size and a 10-fold increase in

contig N50 length compared to those of the latest assembly (Supplementary Table 1) [9,11,12].

The continuity and completeness of the assembled genome were supported by a high mapping

rate (99.72%) and high genome coverage (99.47%) of Illumina short reads, 99.2% single-

copy orthologues from the Benchmarking Universal Single-Copy Orthologues (BUSCO)

analysis and a high long terminal repeat (LTR) assembly index (LAI) score (LAI=9.74).

The complete assembly of a complex allotetraploid presents the opportunity to study

subgenome evolution and coordination following polyploidization. In total, 57.14% and 42.85%

of the genome was partitioned to *N. sylvestris* (S) and *N. tomentosiformis* (T), respectively,

with 12 chromosome rearrange events pinpointed with approximately 1 kb resolution

(Supplementary Table 2). This suggested that chromosomal rearrangements produced

significant changes in genome structure, which likely stabilized chromosome pairing after

polyploidization. To evaluate the mechanisms contributing to genome downsizing and

subgenome coordination, we annotated the transposable elements (TEs) and homologous gene

pairs and quantified their sequence and expression divergence. Based on a de novo-

constructed species-specific TE library, 82.75% of the genome was masked. Estimation of the

TE insertion time showed that approximately 7.56% of the TEs were inserted after merging of

the two subgenomes 0.2 Mya ago (Fig 1C), suggesting that polyploidization did not stimulate

extensive reactivation of TEs. The overall sequence similarity of 35740 homologous gene

pairs between subgenomes was high (median = 92%, Fig 1D), and expression divergence was

low (median fold change estimated from 400 RNA-seq data points of 0.98, Fig 1E,

Supplementary Table 3). However, over a thousand homologous genes with high sequence

similarity displayed considerable expression divergence (median fold change = 3.92, Fig 1E,

Supplementary Table 3, Supplementary Fig S3), indicating a likely role of epigenetic

remodelling in subgenome gene expression coordination for a small proportion of

homologous genes. A detailed summary of the TE annotation and gene annotation statistics is

available in Supplementary Tables 4-5.

Fig 1. High-quality assembly and features of the *N. tabacum* genome. A) Phenotype diversity and genome size variation among two diploid progenitors, *N. sylvestris* and *N. tomentosiformis*, and one of the allotetraploid *N. tabacum* samples. **B)** Genomic features of *N. tabacum.* Tracks from innermost to outermost are sequence coverage obtained by mapping Illumina short reads from *N. tomentosiformis* (a) and *N. sylvestris* (b), gene density (c), and transposable element (TE) density (d). Synteny blocks between the two subgenomes are represented by coloured linkers across the centre of the plot. **C)** Estimated insertion time for TEs from the two subgenomes. Histogram of sequence divergence **D)** and expression divergence **E)** of 35740 protein-coding homologous gene pairs from the two subgenomes.

GenBank genomics highlights the diversity of a global collection of 5196 tobacco lines

To understand the global distribution of genetic variation and genetic structure, we genotyped an entire GenBank collection hosted at the Chinese Academy of Agriculture Sciences using a genotype-by-sequencing approach (Materials and methods). This is one of the world's largest collections of tobacco germplasm resources, with 2582 sun-cured tobacco, 2152 flue-cured tobacco, 223 burley tobacco, 126 cigar tobacco and 113 oriental tobacco germplasms, amounting to 5196 germplasms in total (Supplementary Table 6). These five types of tobacco were named after their roles in agronomical practices (Supplementary Table 7) and characterized by distinctive characteristics in plant architecture, leaf morphology, flowering time and metabolic traits (Fig 2A, Supplementary Fig S4).

Using 95,308 SNPs covering 98% of the genomic bins at a 1 Mb resolution (Fig 2B), we

found that geography at the continent scale was the most important correlate of genetic

structure and that genetic structure did not always match the conventional types defined on

the basis of agronomical practices. For example, the first principle component (PC1)

separated flue-cured tobacco from sun-cured tobacco to a large extent. However, the majority

of samples located on the left side of Fig 2C were of flue-cured tobacco originating from

North America (NA) or derived materials belonging to the NA lineage, while plants located

on the opposite side were sun-cured tobacco landraces collected in China (Fig 2C/2D). In

addition, a latitudinal cline was observed along the second principal component (PC2). Sun-

cured tobacco landraces from northern China (NC) with a significantly shorter life cycle,

colder growth temperature and longer day length were located in the upper right corner

(Cluster NC, Fig 2C), while plants sampled in southern China (SC) with opposite climate

signatures were clustered at the bottom of Fig 2C. According to historical records, burley

tobacco originated from a single mutated variety, known as White Burley, characterized by a

chlorophyll deficiency phenotype controlled by double homozygous recessive alleles at the

Yellow burley 1 (*YB1*) and Yellow burley 2 (*YB2*) loci[13]. Consistent with this, burley tobacco

samples were clustered together near the NA group and showed the highest pairwise isolation

by distance (IBS) value (median=0.88, Fig 2E). Cigar and oriental tobacco samples were

scattered among sun-cured and flue-cured tobacco samples, indicating a likely admixed

breeding history (Fig 2C/D).

Pairwise IBS values for the five tobacco types ranged from 0.75-0.88, suggesting that the

gene pool used to develop each type of tobacco was very narrow. However, the distribution of

IBS values for flue-cured, sun-cured and cigar tobacco was multimodal, indicating the

cooccurrence of divergent gene pools likely due to introgression from foreign materials.

Despite the marked phenotypic differentiation in plant morphology, flowering and metabolic

traits (Fig 2A/Supplementary Fig S4), genome-wide differentiation among the five types and

three genetic groups was low, with pairwise F_{st} values ranging from 0.04 to 0.23 (Fig 2F/G)

and no selective sweeps detected (data not shown here). This is most likely due to the short

and special selective breeding history of less than a few hundred years aiming to balance

quality among many leaf characteristics.

Fig 2. The genetic and phenotypic diversity of a global collection of 5,196 tobacco germplasm resources. A). A heatmap of mean phenotype rank for 20 traits (x-axis) among five types of tobacco named on the basis of their role in agronomical practices. **B).** Genomic distribution of 95,308 SNPs in the *N. tabacum* genome called with the genotype-by-sequencing approach. **C), D)** Genetic structure of 5196 *N. tabacum* lines inferred from principal component analysis. Samples are coloured according to the type assigned on the basis of agronomical practices. **E).** Distribution of pairwise IBS values in different tobacco types. Fixation index (F_{st}) between different tobacco types **F).** Groups defined on the basis of genetic structure **G).**

A comprehensive catalogue of the allelic variations underlying variation of plant morphological, physiological, metabolic and disease-resistance traits

With the availability of genome-wide markers and historical phenotypic records for over 5000

lines, we used the genome-wide association study (GWAS) approach to generate a

comprehensive catalogue of the allelic variation underlying differences in 43 traits (Fig

2A/Fig 3A) with intermediate to high heritability (h^2 = 0.15-0.82, Supplementary Tables 8-9).

A total of 178 significant marker☐trait associations were identified for 39 traits (Fig 3B,

Supplementary Fig S5), and several high-potential associations were detected and explored

further. P values and additive effects for the quantitative trait loci (QTLs) are summarized in

Supplementary Table 10. Although linkage disequilibrium (LD) is too extensive to directly

pinpoint the candidate genes underlying each association peak, we found a few genes

previously identified as QTLs[12,14–18] or annotated as potential candidates near eight peaks

(dashed lines in Fig 3B). In addition, we conducted a review of the literature[12,14–18] to

determine the relative position of previously reported QTLs (Supplementary Table 11) in the

ZY300 genome assembly and discovered that 95% of the QTLs detected in our study were

novel. Altogether, these marker☐trait associations constitute a comprehensive genotype-to-

phenotype map (Fig 3D, Supplementary Table 12), providing a roadmap for future genetic

studies in this model plant species.

To make these resources accessible to a wide range of researchers, we selected 310 accessions

to cover the majority of the genetic and phenotypic diversity, and have made their seeds

available for ordering (See for detailed information). Our intention is for this collection to

remain actively curated as ever more genomic data are produced and a wide range of

phenotypic data are generated (not only by us, but also by the community— see for detailed

information on those accessions and on how to contribute).

231
232

Fig 3. Association results for 43 plant morphological, physiological, metabolic and disease-resistance traits. A). Phenotype diversity of plant architecture, leaf morphology and flowering morphology and colour among five selected plants. The vertical dashed grey lines highlight the genomic position of potential candidate genes. **B).** Manhattan plots of GWAS results from 43 GWAS scans. The red horizontal dashed lines indicate the Bonferroni-corrected genome-wide significance thresholds. The vertical dashed grey lines highlight the genomic position of 16 SNPs associated with more than two traits. **C).** A heatmap illustrating the p values of 16 SNPs detected for more than two traits. Each cell represents the -log10 (p value) of a particular SNP (x-axis) associated with a specific trait (y-axis on the right). D). The reference genotype-to-phenotype map includes marker☐trait associations detected in our study (orange), previous reports (sky blue), and both (dark green). Previously reported QTLs spanning more than 10 Mb and QTLs without probe sequences are not included here.

We detected 32 QTLs (Supplementary Tables S10/S12) for four plant architecture traits,

namely, plant height (PH), internode distance (ID), leaf number (LN) and stem diameter (SD),

that were moderately correlated (Fig 3B, Supplementary Fig S6). Overall, taller plants tended

to have more leaves, wider stems and larger ID values (Fig 4A, Supplementary Fig S4),

indicating a shared molecular basis for plant architecture development. Consistent with this

finding, four QTLs were associated with variation in two or more plant architecture traits (Fig

3B/3C). For example, the QTL (Chr7: 28 405 285 bp) located at the proximal end of

chromosome 7 was associated with variation in PH, LN and SD (Fig 4B). Allele G at this

locus showed a dominance effect over allele A (Fig 4D), whereby the GG genotype at this

locus increased ID, LN and SD by 0.22 ± 0.04 cm (P= 6.32 x 10^{-6}), 0.79 ± 0.15 (P= 4.02 x 10^{-8})

and 0.29 ± 0.05 cm (P= 8.60 x 10^{-11}), respectively (Fig 4D/E/F). Although the LD in this

region extends for several megabases, we discovered a potential candidate, NtZY07G00865,

homologous to *Arabidopsis thaliana* gibberellin 20-oxidase (AT4G25420, GA5), which

regulates plant developmental processes[19], located within the association peaks.

260
261
Fig 4. GWASs identify potential candidate genes for plant architecture, leaf morphology
and flowering time variation. Graphical illustration of the variation in plant architecture **A),**
leaf morphology **G),** and flowering time **K)** traits in *Nicotiana tobacco*. Association results
for internode distance, stem diameter and leaf number **B)**, leaf width, leaf flatness **H),**
flowering time and bolting time **L). C).** The LD heatmap and literature review pinpointed
NtZY07G00865, a homologue of *A. thaliana* AT4G25420 (*GA5*) involved in the later steps of
the gibberellin biosynthetic pathway, as a potential candidate underlying the association peak
at chromosome 7: 28 405 285 bp. **I).** The LD heatmap and literature review pinpointed
NtZY23G02972, a homologue of *A. thaliana* auxin response factor 9 (*Arf9*), which regulates

cell division activity, as a potential candidate underlying the association peak at chromosome

23: 148 211 202 bp. **M), N),** The LD heatmap and literature review pinpointed

NtZY04G03098 and NtZY06G00215, both homologous to *A. thaliana* flowering time locus T

involved in flowering time regulation, as potential candidates underlying the association peak

at Chr6:14 129 418 bp and Chr4:141 102 779 bp, respectively. **D), E), F), J), O), P)**

Genotype-to-phenotype map for the QTLs associated with the corresponding traits.

Strong agronomical interests in leaf usage resulted in an excess amount of diversity in leaf

morphology (Fig 3A, Fig 4G) traits. Here, we detected 65 QTLs for 12 leaf morphology traits

(Supplementary Table S8), including leaf width (LW), leaf length (LL), vein diameter (VD),

leaf stem angle (LSA), leaf vein angle (LVA), leaf auricle (LA), leaf thickness (LT), leaf

shape (LS), leaf tip (LTP), leaf colour (LC), leaf serration (LSR), and leaf flatness (LF).

Overall, the correlations between these traits were relatively low (median Spearman

correlation = 0.05, Fig S3). However, 3 QTLs were simultaneously associated with LW and

LF (Fig 3B/Fig 4H). One of these QTLs (Chr23: 148 211 202 bp) is located at the end of

chromosome 23. Allele T at this locus increased LW by 1.11 ± 0.18 (P= 1.66 x 10^{-9}, Fig 4J)

and increased LF by 0.2 ± 0.02 units (P = 1.58 x 10^{-18}). Similar to that for the peak described

in the previous section, the LD in this region was too extensive to directly pinpoint candidates;

however, we found NtZY23G02972, a homologue of *A. thaliana* auxin response factor 9

(*Arf9*), which regulates cell division activity, to be a potential candidate for further validation

(Fig 4I).

Regulation of flowering time variation has attracted broad interest due to its importance in

plant development, ecological adaptation and breeding applications. Here, four QTLs were

detected for flowering time (FT) and budding time (BT) variation (Fig 4 K-L). Among these,

two QTLs located at the end of chromosome 4 (Chr4:141 102 779 bp) and proximal end of

chromosome 6 (Chr6:14 129 418 bp) attracted our attention (Fig 4L). Both QTLs included a

candidate gene homologous to *A. thaliana* flowering time locus T (*FT*), and these two

chromosome regions were synteny blocks from the S and T subgenomes (Fig 4 M/N, Fig 1B).

This suggests that the function of both homologous genes was retained after merging of the

two subgenomes.

In addition to the association peaks described above, we detected additional QTLs associated

with leaf metabolites and resistance to several virus-/bacterium-/fungus-induced diseases

(Supplementary Table S8). A few of them included genes with known functions reported in
previous studies[3,20]. For example, the QTL located on the proximal end of chromosome 11
(Chr11:2 116 418 bp) associated with TMV resistance was associated with the *N* gene, a TIR-
NBS-LRR protein introgressed through interspecific hybridization with *N. glutinosa*[3,20]. In
addition, two QTLs, located on chromosome 7: 175, 557, 338 bp and chromosome 14: 2,809,
268 bp, associated with resistance to black shank (RBSH), were found to include *Sar8.2a* and
Sar8.2b (Fig 3B), which were detected and fine-mapped previously[17]. Overall, these
marker☐trait associations represent a comprehensive catalogue of the allelic variations
underlying differences in plant morphological, physiological, metabolic and disease-
resistance traits, providing a deeper understanding of complex regulation and several
candidates for future genetics research in this model plant species.

Fine-mapping the causal variants underlying LW variation and functional characterization
of the candidate gene Arf9

Motivated by broad interest in plant leaf development and its importance in light capture
efficiency, we attempted to fine-map the QTL at Chr23: 148 211 202 bp associated with LW
variation. A population of near-isogenic lines (NILs) was created by crossing two parents
segregating at this QTL, Samsum (P1, T/T at Chr23: 148 211 202 bp) with wide leaves and
K326 (P2, C/C at Chr23: 148 211 202 bp) with narrow leaves. Four generations of
backcrossing were performed with K326 as the recurrent parent to create a BC4 population,
and another four generations of self-pollination were performed to generate BC4F4
individuals. Based on the pattern of recombination and phenotype measurements for
recombinant genotypes from BC4F2 and BC4F4 (Fig 5A, Supplementary Tables S13/S14),
the QTL was fine-mapped to a 134 kb region between two markers, Chr23: 147 231 844 bp
and Chr23: 147 366 149 (Fig 5A). There were only two SNPs between the two parents (Fig
5B) and one gene, NtZY23G02972, in this region. NtZY23G02972 is a homologue of *A.*
thaliana auxin response factor 9 (*Arf9*), which widely exists in a number of crops but has not
been functionally characterized. CRISPR☐Cas9 knockout lines of NtZY23G02972 in *N.*
tabacum var. *XHJ* showed an increase in LW by 6 cm (2.74×10^{-3}; Fig 6 C-E, Supplementary
Fig S7), suggesting that the nonfunctional allele of NtZY23G02972 was associated with
wider leaves. Previously, the structure of ARF5 was characterized, and ARF DNA-binding
domains are known to form a homodimer to generate cooperative DNA binding, which is
critical for in vivo ARF5 function[21]. Here, we found that the second SNP on chromosome 23,

147 311 281 bp, located in the eighth exon of NtZY23G02972, altered the translated amino

acid sequence from alanine (Ala_{203}) to proline (Pro_{203}), and the first SNP was located in the

first exon of *Arf9*, altering the amino acid sequence from glutamic acid (Glu_{29}) to glycine

(Gly_{29}). This second SNP was located inside the functional domain forming a homodimer (Fig

5F), while the first SNP was at the proximal end of the protein, which is less likely to have an

impact on protein activity. Altogether, it is highly likely that a causal variant at chromosome

23, 147 311 281 bp, affected the formation of homodimers by changing the amino acid

sequence from alanine (Ala_{203}) to proline (Pro_{203}) and increased LW. Our results revealed a

novel gene regulating plant leaf development and provided a likely causal candidate for future

functional genetic research.

349
350

Fig 5. Fine-mapping and functional characterization of the QTL located at chromosome

23, 148 211 202 bp, associated with leaf width variation. A). Fine-mapping using near-

isogenic lines (NILs). Each bar represents a unique genotype at this QTL region. Red

represents the genotype of the donor parent (P1; Samsum) with wider leaves, while sky blue

highlights the genotype of the recurrent parent (P2; K326) with narrower leaves. Unique

offspring genotype classes from two generations, BC4F2 and BC4F3, are illustrated using

coloured bars, and the line of origin is highlighted using red, sky blue, and grey for P1

homozygous, P2 homozygous, and heterozygous, respectively. **B).** Gene structure in the fine-

mapped region and position of the two SNPs segregating in this population. A highly likely

causal SNP is highlighted using red stars. **C).** Targeted sequences of NtZY23T02972 (*Arf9*) in

CRISPR☐Cas9 knockout lines. **D).** Illustration of the leaf width for a control line (Benhart

1000-1) and targeted line before (XHJ) and after (*arf9*-XHJ-20, *arf9*-XHJ-33) CRISPR☐Cas9

knockout. **E).** Boxplot of the leaf width for a control line (Benhart 1000-1) and targeted line

before (XHJ) and after (*arf9*-XHJ-20, *arf9*-XHJ-33) CRISPR☐Cas9 knockout. **F).** A SNP

altered the protein sequence of ARF from P > M, potentially affecting the formation of a

homodimer to generate cooperative DNA binding.

Signature of positive selection and polygenic selection during the breeding process

When the germplasms were classified into landraces, introduced varieties (varieties developed

from abroad) and Chinese local varieties (abbreviated as varieties hereafter) developed over a

period of several hundred years, seven traits, LW, LL, FT, SL, RS, TS and RBSH, were found

to have been continuously increased/decreased during the process of breeding (Fig 6 A-H). To

test for evidence of selection, we evaluated whether the frequencies of favourable alleles

associated with each trait increased over time. In total, 46.42% of the QTLs (highlighted

using black/red stars in Fig 6I) showed an increase in allele frequency during the process of

breeding. However, none of the alleles reached fixation (frequency >0.95), and only 17.00%

of the QTLs showed allele frequencies greater than 0.8, indicating that there is great potential

for future genetic improvement. Among these, 3 QTLs (10.71%) displayed lower allele

frequencies in landraces than in introduced varieties, followed by varieties (highlighted using

red stars in Fig 6I). For example, one QTL located at chromosome 23: 148, 211, 202 bp is

associated with LW variation. Allele T at this locus increased LW by 1.11 cm (Fig 6J; P value

$= 1.67 \times 10^{-9}$), and the frequency of the T allele continuously increased from 0.15 (landraces)

to 0.38 (introduced varieties) and then to 0.47 (varieties; Fig 6J), suggesting that this allele is

under positive selection.

In addition, polygenic scores were calculated using alleles with a -log10(p value) greater than

2.5, representing aggregated effects of minor-effect alleles. Overall, these minor-effect alleles

(less than 0.5% of all SNPs) accounted for 23.12% to 95.56% (median = 33.40%, Fig 6K) of

the kinship heritability, suggesting an important contribution of the polygenic background to

the variation in these traits. Taking RS as an example, although no significant association was

detected at the genome-wide significance threshold, aggregated effects of 229 (0.23%) alleles

with -log10(p values) above 2.5 (Fig 6L) explained 43% of the phenotypic variation and

95.56% of the kinship heritability (Fig 6L/M). In total, 37.55% (86) of the alleles showed an

increase in allele frequency from the landraces to the introduced varieties or varieties (Fig 6N).

Although the magnitude of the frequency increases was small (mean = 0.07, Fig 6N), together

they increased the level of reducing sugar from 8.07% to 14.71% from landraces to varieties

(Fig 6O), demonstrating the power of polygenic adaptation in response to artificial selection.

Overall, these results demonstrated the power of leveraging GenBank genomics for dissecting

the genetic basis of complex trait variation and evolution during the process of selective

breeding and provided a blueprint for future crop improvement.

Fig 6. Signature of selection during selective breeding. A-H) Phenotype distribution of

eight traits among different categories of materials (landraces, introduced varieties, and

varieties) developed during the process of breeding. **I).** Frequency shifts of 28 QTLs

associated with the variation in eight traits. **J).** Frequency shifts of the QTL associated with

leaf width. **K).** The proportion of variance explained (PVE) by the alleles with -log10(P)

values above 2.5 and all the alleles, estimated in a linear mixed model. **L).** Manhattan plot for

RS association analysis. **M).** Scatterplot between predicted RS values based on 229 alleles

with -log10(P) values above 2.5 and measured alleles. **N).** Frequency shifts of the 229 alleles

with -log10(P) values above 2.5 in genome-wide association analysis of RS. **O).** Relationship

between the mean of predicted RS level based on 229 alleles with -log10(P) values above 2.5

and measured RS values for three categories of plant materials developed during the process

of breeding.

Discussion

N. benthamiana, an ancient allotetraploid species (~ 6 Mya) in the *Nicotiana* genus,

experienced a significant reduction in chromosome number (24 >19) and genome size

(~50%)[22] after polyploidization. In contrast, over 90% of the parental genomic sequences are

retained in the *N. tabacum* genome, and only 12 chromosome rearrangement events have been

identified. This suggests that genome downsizing occurs on an evolutionary scale of several

million years, and the stabilization of chromosome pairing during meiosis by chromosome

rearrangement was likely to be the primary process in such a young polyploid (0.2 Mya). The

high sequence similarity and high expression divergence between 1276 homologous genes

(Supplementary Fig S3) indicate likely significant epigenetic remodelling for subgenome

coordination.

In curing crops, such as tobacco and coffee, a primary target of breeding is balancing quality

with certain morphological and phenological traits, whereby directional selection is relatively

rare in comparison to that in field or horticulture crops. In addition, the history of selective

breeding in tobacco is considerably short, being only ~ 200 years for flue-cured varieties and

~ 500 years for all other varieties. These factors likely explain the observed low divergence

between major tobacco types, which in turn facilitated high statistical power for GWASs in *N.*

tabacum. Future research with resequencing data is required to investigate the signatures of

selection at a finer resolution.

We provided a comprehensive genotype-to-phenotype map in this model plant species and

demonstrated the power of this roadmap in plant functional genetics by fine-mapping one of

the detected QTLs. For example, LW is an important morphological trait associated with

photosynthesis and is of great importance in light capture efficiency. By leveraging the

genotype-to-phenotype map, we detected, fine-mapped and functionally validated a novel

gene, *Arf9*, that regulates LW. Homologues of *Arf9* are widely present in a number of field

crops, such as maize, rice, and wheat, and but not been functionally characterized. It is

worthwhile to evaluate the function and potential of *Arf9* in major crop molecular genetic

research and breeding applications.

In summary, we presented the first chromosome-level assembly of the *N. tabacum*

allotetraploid genome and detected global genetic and phenotypic polymorphisms by

sequencing and phenotyping an entire GenBank collection of 5196 *N. tabacum* germplasm

resources. The extensive genotype and phenotype datasets, marker□trait associations, and

candidate genes presented in this study will serve as a community resource for accelerating

future comparative genomics, plant functional genomics and crop improvement research.

Materials and Methods

Plant materials, DNA extraction and sequencing

Total genomic DNA was collected and extracted from fresh leaves of *N. tabacum L.* var

ZY300 using the CTAB method. For PacBio long-read sequencing, 20 kb insertion

SMARTbell libraries were constructed and sequenced on the PacBio Sequel II platform

(Pacific Biosciences). For 10x Genomics sequencing, 1 ng of genomic DNA with a long

sequence length (approximately 50 kb) was partitioned by a microfluidic chip on the

Chromium platform, and 16-bp barcodes were introduced into droplets. For Illumina short-

read sequencing, paired-end libraries with insert sizes of 350□bp were constructed and

sequenced on the HiSeq PE150 platform. For Hi-C sequencing, DpnII-digested and cross-

linked DNA was labelled with biotin and proximity-ligated to form chimeric junctions.

Biotin-labelled samples were captured and sheared into 350-bp fragments. After terminal

repair, A addition, joint connection, and library construction, Illumina sequencing was

performed on the HiSeq PE150 platform. For Bionano sequencing, high-molecular weight

DNA with a fragment distribution greater than 150 kb was isolated using Bionano sample

preparation kits (Bionano Genomics). Genomic DNA was labelled using Direct Label

Enzyme (DLE-1) and stained following BioNano protocols. Then, the labelled DNA was

loaded into a nanochannel (Saphyr Chip, Bionano Genomics) and imaged using the Saphyr

system (Bionano Genomics) following the Saphyr System User Guide.

Low-quality paired reads (reads with ≥ 10% unidentified nucleotides (N); > 10 nt aligned to

the adapter, > 20% bases having a phred quality score <5 and putative PCR duplicates

generated during the library construction process), which resulted mainly from base-calling

duplicates and adapter contamination, were removed. In total, these steps yielded 1.79 Tb of

high-quality data for chromosome-scale scaffold analysis.

Genome assembly and scaffolding

For initial genome assembly, PacBio long sequence data were self-corrected to generate

preassembled reads and assembled by the overlap-layout consensus algorithm[23]. The

assembled contigs were further polished with Illumina short-read sequence data by the Pilon

pipeline[24]. The polished contigs were further connected to generate super scaffolds using 10x

Genomics linked-reads data using fragScaff software[25]. Scaffolds were anchored by linkage

information, restriction enzyme site, and string graph formulation with the package

LACHESIS[26]. Hi-C data were used to further improve the quality of the assemblies with HiC-

Pro software (v.2.10.0). The placement and orientation errors exhibiting obvious discrete

chromatin interaction patterns were adjusted manually. A final assembly was generated with

Bionano Solve (v3.5.1). Chromosomes were named by mapping SSR markers from a widely

used linkage map[27] to unify the physical map and linkage map.

Repeat and gene annotation

Extensive de novo TE Annotator (EDTA v1.9.3)[28], a pipeline combining homology-based and

de novo prediction methods using LTRharvest[29], LTR FINDER[29], LTR retriever[30], TIR-

learner[31], Helitron Scanner[32], and Repeat Modeler[33], was used to annotate TEs, estimate TE

insertion time and generate a species-specific library for gene annotation. Structural

annotation of genes was conducted through a combination of homology-based, transcriptome-

based and ab initio-based methods. For homologue prediction, protein sequences of plants,

including   *N.   tabacum*

(https://solgenomics.net/ftp/genomes/Nicotiana_tabacum/edwards_et_al_2017/annotation/),

Solanum tuberosum (GCF_000226075.1), *Solanum lycopersicum* (GCF_000188115.4),

Coffea canephora (PRJEB4211_v1), *Gossypium hirsutum* (GCF_000987745.1), *A. thaliana*

(GCA_000001735.1)   and   *Vitis   vinifera*   (http://jul2018-

plants.ensembl.org/Vitis_vinifera/Info/Index), were used as queries to search against the

ZY300 genome using GeneWise (v.2.4.1)[34]. For transcriptome-based gene prediction,

trimmed RNA-sequencing reads from stems, roots, leaves, anthers, flowers, and axillary buds

were de novo assembled using Trinity (v2.1.1) [35]. To further improve the annotation, RNA-

Seq reads from different tissues were aligned to the ZY300 genome using TopHat (v2.0.11)

with default parameters to identify exons and splice junctions. The alignment was used as

input for Cufflinks (v2.2.1) [36] for transcript assembly with default parameters. For ab initio

prediction, we used Augustus[37], GlimmerHMM[38], SNAP[39], GeneID (v1.4)[40] and Genscan[41] to

predict gene structure. Finally, all predictions of gene models generated from these

approaches were integrated into a consensus gene set using EVidenceModeler (v1.1.170)[42].

After prediction, PASA[43] was used to update isoforms to gene models and to produce a final

gff3 file with three rounds of iteration.

For functional annotation, predicted protein-coding genes were aligned to multiple public

databases, including NR, Swiss-Prot, TrEMBL75, COG, and KOG, using NCBI

BLAST□+□v.2.2.31[44]. Motifs and domains were annotated using InterProScan (release 5.32-

71.0)[45]. Gene Ontology (GO) terms and KEGG pathways of predicted sequences were

assigned by InterProScan and KEGG Automatic Annotation Server[45,46], respectively.

Subgenome partitioning and synteny analysis

Illumina short-read sequences from *N. sylvestris* (accession ID: ERR274529) and *N.*

tomentosiformis (accession ID: ERR274543) were downloaded from the NCBI SRA database

and aligned to the ZY300 reference genome using bwa-mem[47]. SAMtools[48] was used to count

the number of reads mapped to the ZY300 genome from each of the two parental genomes,

and the average read depth in 1-Mb windows was calculated using a customized R script.

First, a preliminary parental origin assignment was made using the window-averaged read

ratio. Second, for windows containing breakpoints, manual curation was performed in

Integrative Genomics Viewer[49] (IGV) to pinpoint the exact location. WGDI[50] was used to

detect synteny blocks based on collinear genes, and the R package circlize[51] was used to

illustrate blocks between subgenomes after merging large synteny blocks.

Subgenome expression bias analysis

Four hundred RNA sequencing datasets were randomly selected from the NCBI SRA

database (accession ID available in Supplementary Table S16) to quantify the level of

expression bias for homologous genes from two parental genomes. Raw data were
downloaded and trimmed using fastp[52] with the default parameters and subsequently aligned
to the ZY300 reference genome using HISAT2[53]. The expression level was quantified as
FPKM using Stringtie[54]. Fold change between homologous genes was calculated as the ratio
between averaged FPKM values across the 400 samples.

Genotyping by sequencing, read mapping, and variant calling

DNA extraction was performed using CTAB methods. After quality control using Nanodrop
and Qubit, DNA samples were digested with *NlaIII* + *MseI*, barcoded and purified using
AMPure XP beads. Quality-controlled libraries were pooled and sequenced on the Illumina
NovaSeq platform. Raw reads were trimmed using fastp[52] following the default parameters
and subsequently aligned to the ZY300 reference genome using bwa[47]. Samtools[48] mpileup
was used for variant calling. The called SNPs were initially filtered for read depth, call rate,
and minor allele frequency with vcftools[55] on the basis of average read $DP > 2$ and $MAF >$
0.03, individual missing rate < 0.5, and site missing rate <0.5. The filtered genotypes were
then imputed using beagle[56]. The imputed SNPs were further filtered for a MAF greater than
0.03, and 95,308 SNPs remained for downstream analysis.

Population structure analysis

To further assess the relatedness between individuals, principal component analysis was
performed using the identity-by-state (IBS) genetic distance matrix calculated in PLINK
(v1.90)[57,] and principal component analysis was performed using the *princomp* function in the
R base package. After grouping samples, VCFtools (v0.1.16)[55] was used to calculate the
population differentiation coefficient (F_{st}) among major tobacco types and genetic clusters.

Genome-wide association analysis

Genome-wide association (GWA) analysis was conducted for 43 traits (phenotype
measurements are available in Supplementary Table S17) using a linear mixed model
implemented in the *mlma* module of GCTA[58]. A subsequent conditional analysis implemented
in the *cojo* module of GCTA[58] was performed to screen for independent association signals.
As the LD was extensive in this population, assuming that all tested markers were statistically
independent and deriving a Bonferroni-corrected significance threshold would have been too
conservative. Therefore, we estimated the effective number of independent markers (Me)[59]

and derived a less conservative threshold following 0.05/Me (1. x 10^{-5} equivalent to $-\log_{10}$ (p

value) = 5).

Polygenic selection analysis and genomic prediction

We predicted the aggregated effects of all the SNPs with a -log10 (p value) above 2.5 using

the following model.

Y= Xβ + Zu + e (1)

where Y is a column vector of length n containing phenotype measurements. X is a matrix of

n rows and 1 column on 1s, representing the population mean. u is a random effect vector of

the polygenic effects (polygenic scores) representing the aggregated effects of all the SNPs

for the n individuals. Z is the corresponding design matrix obtained from Cholesky

decomposition of the kinship matrix G, estimated on the basis of the markers with a -log10 (p

value) above 2.5 using the *A.mat* function in the R package rrBLUP [60]. The Z matrix satisfies

ZZ'=G; therefore, u ~ N (0, $I\sigma_g^2$). e is the residual variance with e ~ N (0, $I\sigma_e^2$). The proportion

of variance explained by these SNPs was estimated as $\sigma_g^2/(\sigma_g^2+ \sigma_e^2)$. Model fit was assessed in

the R package rrBLUP [60].

Fine mapping and experimental validation of candidate genes

A population of NILs was developed by marker-assisted backcrossing using Samsun (P1, T/T

at Chr23: 148 211 202 bp) as the wide-leaved donor parent and K326 (P2, C/C at Chr23: 148

211 202 bp) as the narrow-leaved recurrent parent. First, a total of 1694 BC$_4$F$_2$ individuals

from the K326×Samsun population were genotyped using five markers from Chr23: 145 182

211 bp and Chr23:149 453 746 bp (Supplementary Table 13), and then selected recombinants

were self-pollinated to create BC$_4$F$_3$ lines for further evaluation. This narrowed the QTL to a

region between Chr23: 147 119 955 and Chr23: 147 360 000bp. An additional 3017 BC$_4$F$_3$

individuals derived from the population were further screened using seven SNP markers in

this narrow region (Supplementary Table 13), and the BC$_4$F$_4$ lines of recombinants were

phenotyped for further fine mapping. Two sgRNAs (small guide RNAs, Fig 5C) were

designed to target the candidate gene NtZY23T02972 based on the assembled ZY300 genome.

The vectors were constructed and transformed into the receptor XHJ. The genotype of gene-

edited lines was identified by PCR amplification and Sanger sequencing. For fine mapping,

LW was recorded by measuring at least 15 plants per line during flowering in Sanya (E

108.56, N 18.09), China. For transgenic experiments, phenotypes of knockout lines and the

wild type were investigated in Sanya (E 108.56, N 18.09), China. Each experiment had two
replicates with at least five independent plants per replicate. The mean LW of at least five
plants per replicate was used for further analyses.

Funding

This research was funded by the National Science Foundation of China (32200503),
Agricultural Science and Technology Innovation Program (ASTIP-TRIC01) from Chinese
Academy Agriculture Sciences, the Crop Germplasm Resources Protection and Utilization
Special Grant from the Ministry of Agriculture and Rural Affairs of the People's Republic of
China, and the National Science and Technology Resource Sharing Service Platform
(NCGRC-2023-18).

Author Contributions

A.Y., L.C. and Y.C. designed and supervised this study. Y.Z., S.C., M.R., G.L., and Y.L.
collected the data and implemented the analysis. H.S. contributed to the statistical analysis.
Y.L., X.Z., Y.T., C.J., Z.L., D.L., Y.S., Z.X., and R.G. conducted the field experiments and
phenotype measurements. Q.F. and Y.W. collected the germplasms. G.Y. and L.F. offered
valuable input during the design and implementation of this study. Y.Z., S.C. and L.C. wrote
the manuscript.

All authors have read and approved the manuscript and have no conflicts of interest to declare.

Acknowledgements

We thank Yan Ji for her efforts in improving the figures and tables. We thank Wei Zhao and
Ying Guo for their constructive comments to improve the manuscript. We thank Shilin Tian
for his valuable input during the design and implementation of this study.

Data availability
All sequence data will be available from the NCBI upon acceptance of this manuscript.
All phenotype measurements and metadata of samples are included in supplementary tables.

References

1. Eriksson, T. & Jonasson, K. Nuclear division in isolated protoplasts from cells of
higher plants grown in vitro. *Planta* **89**, 85–89 (1969).

2. Clausen, R. E. & Mann, M. C. Inheritance in Nicotiana Tabacum: V. The Occurrence
of Haploid Plants in Interspecific Progenies. *Proc. Natl. Acad. Sci.* 121–124 (1924).

3. Holmes, F. O. Inheritance in tobacco of an improved resistance to infection by tobacco
mosaic virus. *Virology* **12**, 59–67 (1960).

4. Venkateswarlu, K. & Nazar, R. N. Evidence for T–DNA Mediated Gene Targeting to
Tobacco Chloroplasts. *Bio/Technology* **9**, 1103–1105 (1991).

5. Angell, S. M. Consistent gene silencing in transgenic plants expressing a replicating
potato virus X RNA. *EMBO J.* **16**, 3675–3684 (1997).

6. Gunasekaran, B. & Gothandam, K. M. A review on edible vaccines and their prospects.
Brazilian J. Med. Biol. Res. = Rev. Bras. Pesqui. medicas e Biol. **53**, e8749 (2020).

7. Yao, J., Weng, Y., Dickey, A. & Wang, K. Plants as Factories for Human
Pharmaceuticals: Applications and Challenges. *Int. J. Mol. Sci.* **16**, 28549–28565
(2015).

8. Goodin, M. M., Zaitlin, D., Naidu, R. A. & Lommel, S. A. Nicotiana benthamiana□:
Its History and Future as a Model for Plant–Pathogen Interactions. *Mol. Plant-Microbe*
Interact. **21**, 1015–1026 (2008).

9. Sierro, N. *et al.* The tobacco genome sequence and its comparison with those of tomato
and potato. *Nat. Commun.* **5**, 3833 (2014).

10. Sierro, N. *et al.* Reference genomes and transcriptomes of Nicotiana sylvestris and
Nicotiana tomentosiformis. *Genome Biol.* **14**, R60 (2013).

11. Leitch, I. J. *et al.* The Ups and Downs of Genome Size Evolution in Polyploid Species
of Nicotiana (Solanaceae). *Ann. Bot.* **101**, 805–814 (2008).

12. Edwards, K. D. *et al.* A reference genome for Nicotiana tabacum enables map-based
cloning of homeologous loci implicated in nitrogen utilization efficiency. *BMC*
Genomics **18**, 448 (2017).

13. Cameron, D. R. & Moav, R. M. INHERITANCE IN NICOTIANA TABACUM
XXVII. POLLEN KILLER, AN ALIEN GENETIC LOCUS INDUCING ABORTION
OF MICROSPORES NOT CARRYING IT. *Genetics* **42**, 326–335 (1957).

14. Lan, T. *et al.* Mapping of quantitative trait loci conferring resistance to bacterial wilt in
tobacco (Nicotiana tabacum L.). *Plant Breed.* **133**, 672–677 (2014).

15. Yuan, G. *et al.* Development of a MAGIC population and high-resolution quantitative
trait mapping for nicotine content in tobacco. *Front. Plant Sci.* **13**, 1086950 (2022).

16.   Liu, Y. *et al.* Identification of QTLs Associated With Agronomic Traits in Tobacco via

a Biparental Population and an Eight-Way MAGIC Population. *Front. Plant Sci.* **13**,

878267 (2022).

17.   Shi, R., Jin, J., Nifong, J. M., Shew, D. & Lewis, R. S. Homoeologous chromosome

exchange explains the creation of a QTL affecting soil☐borne pathogen resistance in

tobacco. *Plant Biotechnol. J.* **20**, 47–58 (2022).

18.   Sallaud, C. *et al.* Characterization of two genes for the biosynthesis of the labdane

diterpene Z-abienol in tobacco (Nicotiana tabacum) glandular trichomes. *Plant J.* **72**,

1–17 (2012).

19.   Zheng, F., Wang, Y., Gu, D. & Liu, X. Histone Deacetylase HDA15 Restrains PHYB-

Dependent Seed Germination via Directly Repressing GA20ox1/2 Gene Expression.

Cells **11**, 3788 (2022).

20.   Marathe, R., Anandalakshmi, R., Liu, Y. & Dinesh-Kumar, S. P. The tobacco mosaic

virus resistance gene, N. *Mol. Plant Pathol.* **3**, 167–172 (2002).

21.   Boer, D. R. *et al.* Structural basis for DNA binding specificity by the auxin-dependent

ARF transcription factors. *Cell* **156**, 577–89 (2014).

22.   Bombarely, A. *et al.* A draft genome sequence of Nicotiana benthamiana to enhance

molecular plant-microbe biology research. *Mol. Plant. Microbe. Interact.* **25**, 1523–

1530 (2012).

23.   Pendleton, M. *et al.* Assembly and diploid architecture of an individual human genome

via single-molecule technologies. *Nat. Methods* **12**, 780–786 (2015).

24.   Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant

detection and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

25.   Adey, A. *et al.* In vitro, long-range sequence information for de novo genome assembly

via transposase contiguity. *Genome Res.* **24**, 2041–2049 (2014).

26.   Burton, J. N. *et al.* Chromosome-scale scaffolding of de novo genome assemblies

based on chromatin interactions. *Nat. Biotechnol.* **31**, 1119–1125 (2013).

27.   Bindler, G. *et al.* A high density genetic map of tobacco (Nicotiana tabacum L.)

obtained from large scale microsatellite marker development. *Theor. Appl. Genet.* **123**,

219–230 (2011).

28.   Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a

streamlined, comprehensive pipeline. *Genome Biol.* **20**, 275 (2019).

29.   Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible

software for de novo detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18

(2008).

30. Ou, S. & Jiang, N. LTR_retriever: A Highly Accurate and Sensitive Program for

Identification of Long Terminal Repeat Retrotransposons. *Plant Physiol.* **176**, 1410–

1422 (2018).

31. Su, W., Gu, X. & Peterson, T. TIR-Learner, a New Ensemble Method for TIR

Transposable Element Annotation, Provides Evidence for Abundant New Transposable

Elements in the Maize Genome. *Mol. Plant* **12**, 447–460 (2019).

32. Xiong, W., He, L., Lai, J., Dooner, H. K. & Du, C. HelitronScanner uncovers a large

overlooked cache of Helitron transposons in many plant genomes. *Proc. Natl. Acad.*

Sci. **111**, 10263–10268 (2014).

33. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable

element families. *Proc. Natl. Acad. Sci.* **117**, 9451–9457 (2020).

34. Birney, E., Clamp, M. & Durbin, R. GeneWise and Genomewise. *Genome Res.* **14**,

988–995 (2004).

35. Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the

Trinity platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494–1512

(2013).

36. Ghosh, S. & Chan, C.-K. K. Analysis of RNA-Seq Data Using TopHat and Cufflinks.

Methods Mol. Biol. **1374**, 339–361 (2016).

37. Stanke, M. & Morgenstern, B. AUGUSTUS: a web server for gene prediction in

eukaryotes that allows user-defined constraints. *Nucleic Acids Res.* **33**, W465–W467

(2005).

38. Majoros, W. H., Pertea, M. & Salzberg, S. L. TigrScan and GlimmerHMM: two open

source ab initio eukaryotic gene-finders. *Bioinformatics* **20**, 2878–9 (2004).

39. Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 59 (2004).

40. Guigó, R. Assembling genes from predicted exons in linear time with dynamic

programming. *J. Comput. Biol.* **5**, 681–702 (1998).

41. Burge, C. & Karlin, S. Prediction of complete gene structures in human genomic DNA.

J. Mol. Biol. **268**, 78–94 (1997).

42. Haas, B. J. *et al.* Automated eukaryotic gene structure annotation using

EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biol.* **9**,

R7 (2008).

43. Haas, B. J. *et al.* Improving the Arabidopsis genome annotation using maximal

transcript alignment assemblies. *Nucleic Acids Res.* **31**, 5654–66 (2003).

44.   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**,

421 (2009).

45.   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.

Bioinformatics **30**, 1236–40 (2014).

46.   Aoki-Kinoshita, K. F. & Kanehisa, M. Gene Annotation and Pathway Mapping in

KEGG. in *Comparative Genomics* 71–92 (Humana Press, 2007). doi:10.1385/1-59745-

515-6:71.

47.   Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler

transform. *Bioinformatics* **26**, 589–595 (2010).

48.   Li, H. A statistical framework for SNP calling, mutation discovery, association

mapping and population genetical parameter estimation from sequencing data.

Bioinformatics **27**, 2987–2993 (2011).

49.   Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer

(IGV): high-performance genomics data visualization and exploration. *Brief. Bioinform.*

14, 178–92 (2013).

50.   Sun, P. *et al.* WGDI: A user-friendly toolkit for evolutionary analyses of whole-

genome duplications and ancestral karyotypes. *Mol. Plant* **15**, 1841–1851 (2022).

51.   Gu, Z., Gu, L., Eils, R., Schlesner, M. & Brors, B. Circlize implements and enhances

circular visualization in R. *Bioinformatics* **30**, 2811–2812 (2014).

52.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

53.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**,

907–915 (2019).

54.   Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from

RNA-seq reads. *Nat. Biotechnol.* **33**, 290–5 (2015).

55.   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

(2011).

56.   Browning, B. L. & Browning, S. R. Genotype Imputation with Millions of Reference

Samples. *Am. J. Hum. Genet.* **98**, 116–126 (2016).

57.   Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-

Based Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

58.   Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: A Tool for Genome-

wide Complex Trait Analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).

59.   Li, M.-X., Yeung, J. M. Y., Cherny, S. S. & Sham, P. C. Evaluating the effective

774   numbers of independent tests and significant p-value thresholds in commercial

775   genotyping arrays and public imputation reference datasets. *Hum. Genet.* **131**, 747–56

776   (2012).

777   60.   Endelman, J. B. Ridge Regression and Other Kernels for Genomic Selection with R

778   Package rrBLUP. *Plant Genome* **4**, 250–255 (2011).

779