Socio-cultural practices affect sexual dimorphism in stature in Early Neolithic Europe

Samantha L Cox[1,2*], Nicole Nicklisch[3], Michael Francken[4], Joachim Wahl[5], Harald Meller[6], Wolfgang Haak[7], Kurt W Alt[3], Eva Rosenstock[8], and Iain Mathieson[1*]

[1]*Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA*

[2]*Physical Anthropology Section, Penn Museum, University of Pennsylvania, Philadelphia, PA, USA*

[3]*Center of Natural and Cultural Human History, Danube Private University, Krems-Stein, Austria*

[4]*Landesamt für Denkmalpflege, Dienstsitz Konstanz, Konstanz, Germany*

[5]*Institut für Naturwissenschaftliche Archäologie, AG Paläoanthropologie, Tübingen, Germany*

[6]*State Office for Heritage Management and Archaeology Saxony-Anhalt, State Museum of Prehistory, Halle/Saale, Germany*

[7]*Department of Archaeogenetics, Max Planck Institute for the Science of Human History, Jena, Germany*

[8]*Bonn Center for ArchaeoSciences, Universität Bonn, Bonn, Germany*

Correspondence to SLC (coxsl@sas.upenn.edu) or IM (mathi@pennmedicine.upenn.edu)

Abstract

The rules and structure of human culture impact health and disease as much as genetics or the natural environment. Studying the origins and evolution of these patterns in the archaeological record is challenging as it is difficult to tease apart the effects of genetics, culture, and environment. We take a multidisciplinary approach by combining published ancient DNA, skeletal metrics, paleopathology, and dietary stable isotopes to analyze cultural, environmental, and genetic contributions to variation in stature in four geographically defined populations of Early Neolithic Europe: North Central, South Central, Southern (Mediterranean), and Southeastern (Balkan) Europeans. In individuals from Central Europe, female stature is low, despite polygenic scores for height identical to males and to neighboring regions. Dietary and skeletal stress markers indicate environmental stress that is equal in both sexes, but the high stature sexual dimorphism ratio suggests that these stresses were exacerbated in females by cultural factors, likely associated with male preference and sex-biased allocation of resources. In contrast, shorter average stature in Mediterranean Neolithic populations have been previously reported to be associated with genetic differences; however, this is likely an artifact of residual population structure in the genome-wide association studies (GWAS). Instead, we suggest that reduced sexual dimorphism in the region indicates a degree of male vulnerability in response to general environmental stress. We conclude that while population-level stature trends may in some cases reflect genetic factors, differences in sexual dimorphism are largely driven by culture, or the interaction of culture and environment. Our analysis indicates that biological effects of sex-specific inequities can be linked to cultural influences at least as early as 7000 years before present. Understanding these patterns is key to interpreting the evolution of genetic and socio-cultural determinants of health and our results show that culture, more than environment or genetics, drove height disparities in Early Neolithic Europe.

1 Introduction

Human skeletal variation reflects varying combinations of genetic, cultural, and environmental factors. While there are many links between culture and health in the modern world, the history and evolution of these relationships are not always well established. Due to the entanglement of these factors, our ability to draw conclusions about their effects has been limited in archaeological data. With the recent advent of ancient DNA sequencing technology, genetic information from ancient populations has become increasingly available, but attempting to analyze changing patterns of variation based solely on genetic data is equally difficult as genotypes do not necessarily equate to phenotypes due to the effect of environment (Harpak & Przeworski, 2021). Similarly, while it is tempting to predict phenotypic changes in ancient people based on their genetic variation, this is currently challenging as genetic effects are not always transferable across populations (Martin et al., 2019). Our solution is to integrate these complementary fields to construct multidisciplinary analyses with phenotype, genotype, culture, and environment data from ancient human populations. This approach allows us to begin to separate the effects of these variables and reveal the interactions between genes, environment, and culture which are critical in shaping human health and variation.

Many traits of interest, including height, are highly polygenic with thousands of independent genetic variants contributing significantly to heritability. One common approach to addressing the role of genetics in morphological change is to compare patterns of phenotypic variation with genetic ancestry or genome-wide patterns of genetic variation (e.g., Roseman, 2004; Whitlock, 2008; Savell et al., 2016). However, even for highly polygenic traits like height, genome-wide variation may not be directly relevant to a particular trait, leading to spurious associations between genetic effects, ancestry, and environmental confounds. For example, if a population is tall and has a high proportion of ancestry from Neolithic sources, it could be concluded that Neolithic ancestry is associated with "genetic tallness"; however, the effects could equally be non-genetic and related to lifestyle changes associated with agriculture. An alternative approach is to focus only on genetic variation that is known to be associated with a specific trait (e.g., Berg & Coop, 2014; Mathieson et al., 2015). Effect sizes for these trait-related variants estimated from genome-wide association studies (GWAS) of present-day individuals can be combined with genetic data from ancient individuals to calculate polygenic scores (PRS), which can be thought of as estimated genetic values for the phenotype. In European ancestry populations, polygenic scores for height can explain up to 25% of phenotypic variation in present-day individuals (Yengo et al., 2018), and 6-8% of variation in ancient individuals (Cox et al., 2022; Marciniak et al., 2022). On a broad scale, temporal changes in polygenic score over time in Europe are qualitatively consistent with changes in stature as inferred from the skeletal record (Cox et al., 2019) while local deviations from this pattern provide evidence of environmental effects (Cox et al., 2019; Marciniak et al., 2022).

Analyses of human populations over tens of thousands of years involve individuals that are diverse in genetic ancestry, environment, and culture and it is challenging to exclude the possibility

of confounding by unmeasured variables. We therefore focus specifically on the European Early Neolithic. One of the most studied periods in prehistory, it represents a fundamental shift in technology, culture, and genetics. In particular, the *Linearbandkeramik* (LBK) culture of Central Europe is one of the most comprehensively documented Early Neolithic cultures, with an abundance of excavated settlements and cemeteries (i.e, Bickle & Whittle, 2013). The Mesolithic hunter-gatherer population in this region made a limited genetic contribution to the LBK population, whose members harbor only traces of hunter-gatherer admixture (Haak et al., 2015; Mathieson et al., 2015; Lipson et al., 2017). Contemporary populations from southeastern Europe have similarly low levels of hunter-gatherer ancestry (Mathieson et al., 2018). In contrast, Neolithic southern European populations associated with the Cardial and Impressed Ware cultures followed a separate migration route, occupied a milder climate zone, and carry more Mesolithc ancestry (Haak et al., 2015; Antonio et al., 2019) (Fig. 1). Individuals in this region tend to be shorter than those from Central Europe and this combination of factors has led to suggestions of a genetic basis for decreased statures in this region (Mathieson et al., 2015; Martiniano et al., 2017).

By comparing and contrasting these three closely related archaeological populations, we aim to investigate how differences in environment and genetics combine to produce observed phenotypes. We collected genetic data, skeletal metrics, paleopathology, and dietary stable isotopes, to begin separating the effects of each on Neolithic stature trends. By specifically investigating and controlling for the effects of genetics in these samples, we are able to provide novel, nuanced interpretations of height variation, gain a better understanding of the aspects of height which are controlled by genetics or environment, and show evidence for sex-specific cultural effects which modify the genetically predicted patterns. We illustrate the strengths of leveraging multidisciplinary datasets, and indicate caution when analyzing genotype-phenotype relationships without complete data, especially for traits which are not preserved in the archaeological record and cannot be directly tested. This integrated analysis highlights the role of plasticity in morphology, and establishes culturally mediated disparities at least as early as the European Neolithic.

2 Materials and Methods

We collected a combination of genetic, dietary stable isotope, skeletal metric, and paleopathological (stress) data from 1282 individuals from the Central European Early Neolithic associated with the archaeological LBK culture, approximately 7700-6900BP (Figure 2, Supplementary Table 1). We divided these individuals into two regions based on geographical location, those to the north of 50°N latitude (North Central) and those to the south (South Central) (Figure 2A-B; North Central n=203, n femur length=131, n isotopes=100, n aDNA=67, n stress=97; South Central n=1107, n femur length=188, n isotopes=670, n aDNA=72, N stress=526). Each individual has at least one of the data types, and while some individuals have multiple data types, the overlaps are small (Supplementary Figure 1).

Figure 1: *This schematic illustration highlights the two main migration routes from Anatolia into Europe during the Early Neolithic period. Population movements followed two routes: southern, along the Mediterranean coast (including sea routes, generalized here by dashed blue lines) where they admixed with existing hunter-gatherer populations; or northern, through the Balkans and into Central Europe, with only limited hunter-gatherer admixture. We also analyze patterns within the Linearbandkeramik culture, dividing it into Northern and Southern Central European groups.*

To provide wider context, we also compared Central individuals to other Neolithic populations from southern European (Mediterranean) and southeastern European (Balkan) regions, and restricted to individuals dated to between 8000-6000BP. We chose these regions as the Neolithic transition occurs at a similar time and is associated with populations closely related to Central Europe. The acceptable date range for inclusion in the study was expanded from that which defines the LBK as these dates encompass comparable Early Neolithic phases in other parts of Europe while maximizing the number of eligible individuals. There could be a possibility that the later Balkan and Mediterranean individuals were more adapted to Neolithic life than the Central European groups, as these samples cover a longer time period, but we found no statistical within-population differences in our variables between the early and late ranges of our time span (minimum p=0.08). We excluded areas such as Scandinavia and Britain where Neolithic technologies were not generally adopted until a later date. For the final analysis, we included 160 Mediterranean (n femur length=67, n isotopes=26, n aDNA=42) and 135 Balkan (n femur length=12, n isotopes=97, n aDNA=49) individuals (Figure 2). Unfortunately, there is a wide range of recording and reporting used for skeletal stress indicators, and it was not possible to build a statistically powerful dataset in these two populations for comparison; as a result, we did not analyze paleopathology in these populations. Finally, we collected genetic data from Mesolithic hunter-gatherer (n=25, 14000-7080BP, south of 48°N) and Anatolian Neolithic (n=21) individuals for additional comparison.

2.1 Genetic data

We obtained genetic data for a total of 276 individuals (Antonio et al., 2019; Brunel et al., 2020; Childebayeva et al., 2022; Fernandes et al., 2020; Fregel et al., 2018; Fu et al., 2016; Gamba et al., 2014; González-Fortes et al., 2017; Hofmanová et al., 2016; Jones et al., 2015; Lazaridis et al., 2017; Lipson et al., 2017; Marcus et al., 2020; Mathieson et al., 2015, 2018; Nikitin et al., 2019; Olalde et al., 2015, 2019; Rivollat et al., 2020; Valdiosera et al., 2018; Villalba-Mouco et al., 2019). Most data were generated by targeting a set of 1.24 million SNPs (the "1240" capture reagent, Haak et al., 2015; Fu et al., 2016). For those individuals with shotgun sequence data we randomly selected a single allele from each of the 1240k sites. Coverage in our dataset is low (median coverage=0.33; coverage above 0.60 n=71), and typically, it is not possible to directly infer diploid genotypes, potentially limiting PRS performance. Cox et al. (2022) showed that imputation of missing genotypes helps to improve polygenic predictions for low coverage ancient samples, and we imputed using the two-stage method described in that paper, restricting to SNPs in the 1240k set.

We calculated polygenic scores as described in Cox et al. (2022). Briefly, we used standing height summary statistics generated by *fastGWA* from 456,000 individuals of European ancestry in the UK Biobank (Jiang et al., 2019) for analyses of combined-sex PRS, and summary statistics from male- and female-only UK Biobank GWAS generated by the Neale Lab (Neale Lab, 2018). To test the potential effects of residual population structure in our data, we also computed PRS using additional summary statistics from the Howe et al. (2022) sibling cohort (n=99,997). We

intersected the sites from each of these datasets with those on the 1240k array and then further
restricted to HapMap3 SNPs (SNPs n=405,000). We computed polygenic scores using both a
clumping/thresholding approach (r^2=0.3, p-value cutoff=10^{-6}, 100kb windows in *plink2* (Chang
et al., 2015)), and an infinitesimal *LDpred2* model using their pre-computed LD reference panel
(Privé et al., 2020). Finally, we computed polygenic scores using the --score command in *plink2*.
In order to maximize the possibility of detecting sex-specific effects, we generated sex-specific PRS
using three different approaches: 1) calculating PRS for all individuals using the female summary
statistics; 2) calculating PRS for all individuals using the male summary statistics; and 3) calculating
PRS for males and females separately using their respective summary statistics. While approach
3 seems at first to be the best for detecting these effects, observed patterns potentially become
difficult to interpret due to differences in scaling between male and female PRS calculated as separate
datasets.

We computed principal components for both unimputed and imputed data using *smartpca* (Pat-
terson et al., 2006), projecting ancient individuals onto principal component axes defined by 777
present-day West Eurasian individuals (Lazaridis et al., 2014). We also estimated K=2 ADMIX-
TURE (Alexander et al., 2009) components for unimputed ancient individuals after first LD pruning
using the command --pairwise-indep 200 25 0.4 in *plink2*.

166 2.2   Osteology and stable isotope data

We aggregated skeletal metric data from both published (Hujić, 2016; Nicklisch, 2017; Ruff, 2018;
Meyer et al., 2018; Rosenstock et al., 2019a,b) and unpublished (new n=28) sources. Maximum
femur lengths were recorded when available, otherwise we estimated femur length from published
stature estimates as described by Cox et al. (2022). Estimated femur lengths correlate highly
with stature estimates but decrease the error that results from combining different estimations
methods. The method from Ruff et al (2012) provides separate equations for estimating the statures
of northern vs. southern Europeans when using the tibia due to differences in body proportions
between the regions. There are two Mediterranean samples for which we estimated the length of
the femur based on statures which used Ruff et al's southern tibia equation. Ruff et al do not
provide regional equations for femur estimation, so for these two individuals, we estimated femur
length using the reverse of this region-agnostic femur equation. For individuals without DNA, we
relied upon their published sex estimations; sex for unpublished individuals was determined using
Acsádi et al. (1970) (see the Supplement for details). Similarly, ages were determined based on
the average of the age range reported for each individual in their original publications. Finally, for
the paleopathological data in Central Europe, we took data from published sources (Lillie, 2008;
Whittle et al., 2013a,b; Ash et al., 2016; Hujić, 2016; Nicklisch, 2017), recorded as presence/absence
of linear enamel hypoplasia, porotic hyperostosis, and cribra orbitalia.
While affected by many confounding factors such as climate, vegetation, and individual metabolism
(Scheibner, 2016), δ^{13}C and δ^{15}N stable isotope data can be used to reconstruct aspects of diet

(O'Brien, 2015). Here, carbon values are indicative of dietary plant resources, and of the terrestrial
vs marine vs limnic provenance of food, while nitrogen values are mainly associated with dietary
protein intake and generally indicate proportions of plant- vs animal-based diets (O'Brien, 2015;
Scheibner, 2016). We collected dietary stable isotopes δ^{13}C and δ^{15}N from published (Dürrwächter
et al., 2006; Richards et al., 2008; Oelze et al., 2011; Bentley et al., 2013; Bickle et al., 2013b,a;
Hofmann et al., 2013; Whittle et al., 2013a,b; Ash et al., 2016; Hujić, 2016; Scheibner, 2016; Meyer
et al., 2018; Münster et al., 2018; Knipper, 2020) and unpublished (new n=38) reports. We ex-
cluded atomic mass spectrometer (AMS) values derived from radiocarbon dating, as they may not
be comparable to isotope-ratio mass spectrometer (IRMS) measurements, as well as values from
children below the age of three, due to increased nitrogen values from breastfeeding. Stable isotope
values from older children were included in population-wide diet analyses as the isotope ranges fall
within those of adults; however, we only included adults with estimated sexes in the sex-based diet
analyses. If information on the sampled material was available, we chose values measured from
rib collagen, as these samples are most plentiful, though they only reflect the last few years of the
individual's life.

All previously unpublished osteological data was collected and analyzed by co-authors with
permission from the necessary regulating organizations and in accordance with German laws and
policies.

2.3 Statistical models

We tested the effects of PRS, femur length, and isotope data on stature using linear regression
models including sex and geographic region as covariates in combination with other variables as
appropriate (e.g., femur $\sim$ sex + region + PRS; δ^{15}N $\sim$ sex + region + femur). We included
interaction terms to test the relationships between geographic regions and sex (e.g., femur $\sim$ region
* sex) and used t-tests to test within-sex differences between regions. We used logistic regression
with the same covariates to test for factors affecting presence/absence of paleopathologies. We
carried out all statistical tests using the the base functions in R version 4.0 (R Core Team, 2021).

3 Results

3.1 Distribution of stature, polygenic scores and stable isotope values

Polygenic scores are very similar between all populations (pairwise t-test p > 0.9) using the clump-
ing/thresholding PRS construction (Figure 3A). PRS constructed with LDPred show Mediterranean
individuals to be shorter than the other populations (p=0.002; Supplementary Fig 5). However,
PRS constructed using summary statistics derived from between-sibling analysis finds similar ge-
netic values in all populations, with both PRS construction methods, so we conclude that apparent
lower Mediterranean PRS may be due to population stratification in the GWAS data and may not

Figure 2: *Upper row: Locations of sites used for genetic (A) and skeletal (B) data in the analysis. The Central European population is split into Northern and Southern groups at 50°N latitude (emphasized). Lower row: (C) imputed genetic data projected into the PCA space of 777 modern Eurasian individuals (grey points). (D) plot of sample numbers by time period in years before present (years BP) for skeletal (right) and genetic (left) data.*

220 reflect a true genetic difference (Figure 5D). Despite the similarities in PRS, observed patterns of
221 femur length vary between sexes and populations. Males show no apparent difference between the
222 Central and Balkan femora (p=0.56), but the Mediterranean male population is significantly shorter
223 (p=5.5×10^{-7}, β=-1.44cm). Conversely, female femora show a different pattern, with no significant
224 difference between Mediterranean, South Central (p=0.97), and Balkan (p=0.54) populations, but
225 substantially shorter values in the North Central (β=-2.0cm, p=9×10^{-07}) (Figure 3B). There are
226 no significant differences between male and female PRS in any population (Fig. 3A), providing no
227 evidence for a genetic basis to this dimorphism.

228   Signatures of δ^{13}C and δ^{15}N indicate different dietary patterns in each of the analyzed groups.
229 Both the Mediterranean and Balkan groups are significantly distinct from the Central in δ^{13}C
230 (p<4×10^{-16}) and δ^{15}N (Balkans: p<7.7×10^{-13}; Mediterranean: 1.2×10^{-122}) values. Generally,
231 the Balkan population is characterized by high δ^{15}N values, while Mediterranean populations show
232 high δ^{13}C relative to the Central Europeans (Figure 3C). The exception to this pattern is a cluster
233 of individuals, classified as Balkan in our analysis, which overlap with the North Central population
234 as well as some of the Mediterranean values. These points represent individuals from present-day
235 Greece and indicate that the diets of these peoples might better be classified as Mediterranean
236 rather than Balkan. Nitrogen values are generally elevated in males compared to females, but this
237 difference is only significant in the Mediterranean (p=0.035).

3.2   Patterns of non-genetic factors in Central Europe

239 The most dramatic observation is the difference in female stature and consequent sexual dimorph-
240 ism in Northern compared to Southern Central Europe. Female femora in Northern Central
241 Europe are significantly shorter than female femora in the South (p=2.7×10^{-6}, β=1.7cm), while
242 male femora are not significantly different (p=0.35) (Figure 3B). On average, male femora from the
243 North are about 13% longer than female femora, Southern Central and Balkan males are about 9%
244 and 11% longer respectively, and Mediterranean males are only 5% longer (Figure 3B). These val-
245 ues are reduced slightly when calculated using estimated statures instead of femora (North Central:
246 10%, South Central: 7%, Balkans: 8%, Mediterranean: 4%), possibly due to error associated with
247 stature estimation (see Cox et al., 2022) and body proportions, or because the relationship between
248 femur length and stature is different between males and females. Where we have both genetic and
249 metric data for the same individuals, there is a qualitative relationship between femur length and
250 PRS; PRS tends to increase as femur lengths increase (Supplementary Figure 3B). However, the
251 effect of PRS on femur length is barely statistically significant (p=0.05), likely due to the small
252 number of individuals in the sample (n=55).

253   Overall, trends in dietary stable isotopes show both males and females in Southern Central
254 Europe to have significantly higher δ^{15}N (Male p=1.3×10^{-9}, β=0.83‰, Female p=5.3×10^{-9},
255 β=0.87‰) and lower δ^{13}C (Male p=0.0003, β=-0.30‰, Female p=8.1×10^{-7}, β=-0.38‰) as
256 compared to the North. However, while males qualitatively have higher nitrogen, the interaction

Figure 3: *Solid bars across the tops of plots indicate significant differences between male populations by pairwise t-test; solid bars below plots indicate significant differences between female populations by pairwise t-test; p-values < 0.05 (*), < 0.01 (**) and < 0.001 (***). Black points indicate the mean of each group and vertical bars show mean standard error. A) Polygenic scores for the four populations show similar scores for individuals across all regions. Differences between male and female PRS are not significant. B) Femur length in the four populations. Values to the right of the vertical dotted line are the difference between the mean male and female femora; values to the left of the vertical dotted line are the sexual dimorphism ratios of male/female femur lengths for each population. C) Plot of $\delta^{13}C$ (x-axis) and $\delta^{15}N$ (y-axis) dietary stable isotopes for the four populations. Individuals from the Balkans are distinguished by high nitrogen values, while those in the Mediterranean generally have higher carbon. D) Sex differences in $\delta^{15}N$ values by sex for each population. $\delta^{15}N$ values are slightly higher for males in all our populations but this difference is only significant in the Mediterranean (p = 0.035).*

Figure 4: *Evidence of environmental stress in Northern Central Europe. A) Differences in $\delta^{13}C$ (right) and $\delta^{15}N$ (left) values. Overall, the South has higher nitrogen values than the North (p=6.8 × 10^{-13}), and less carbon (p=5.3 × 10^{-15}); within each population, the difference in isotopes between sexes is not significant. B) Proportion of linear enamel hypoplasias. The South has significantly less than the North (p=0.001). C) Presence of linear enamel hypoplasia is significantly associated with shorter femora (p=0.02); differences in prevalence between sexes are not significant.*

effect between sexes is not significant in either region (Figure 4A). There is no difference in carbon
values between sexes. For individuals with both stature and stable isotope values, we find no
statistically significant relationship between femur length and δ^{15}N or δ^{13}C in either Central group,
separately or combined, though the sample is small.

We do, however, find a statistically significant relationship between presence of linear enamel
hypoplasias and shorter femora, suggesting that LEH may reflect some underlying variable in child-
hood that also affects stature (p=0.021, β=-1.0cm)(Figure 4C). Both males and females from the
north are more likely to have LEH than individuals living in the south (p=0.002). Indeed, over
50% of the Northern sample have LEH while they are only present in about 20% of the Southern
(Figure 4B). There is no significant difference between the number of males and females with LEH in
either region. Though the interaction effect between sex and LEH on femur length is not significant,
qualitatively the effect of LEH on femur length appears greater in females than in males (Figure
4C). When the sexes are analyzed separately, females with LEH do have significantly shorter femora
than those without (p=0.018, β=-1.46), which is not the case for males (p=0.479). We hypothesize
that the relationship between LEH and femur length is driven by females but we lack an adequate
sample size to detect the interaction effect in the full model. Incidence of cribra orbitalia is also
significantly higher in the Northern region than in the Southern (p=1.8 × 10^{-6}), though there is no
relationship with femur length. There are no significant trends related to the presence of porotic
hyperostosis.

In summary, comparison of Northern and Southern Central Europe identifies no predicted genetic
difference in stature, which is consistent with male but not female femur length. This suggests a
non-genetic basis for reduced female stature. Stable isotope data and skeletal stress indicators

suggest less protein intake and more general stress in the North; however, males and females overall appear equally affected by these variables. Despite a similar number of hypoplasias in both sexes, shorter femora in females suggests increased general stress leading to a female-specific reduction in stature due to other unmeasured environmental or cultural factors.

3.3 Patterns of genetic ancestry in the Mediterranean

In contrast to Northern Central Europe, Mediterranean Neolithic males are shorter than other groups, but females are not. PCA indicates that individuals from the Central regions and the Balkans share similar genetic ancestry while those from the Mediterranean are distinct (Figure 2C; unimputed PCA in Supplementary Figure 4A), a known difference due to higher levels of hunter-gatherer ancestry in the Mediterranean (Haak et al., 2015). We therefore additionally compared our samples to Mesolithic individuals of Western Hunter-Gatherer (WHG) ancestry, as well as individuals from early Neolithic Anatolia. These two groups represent source populations for the two largest ancestry components in Europe at this time (Haak et al., 2015; Mathieson et al., 2015).

On the PCA plots of these extended data, Neolithic Anatolians cluster with the Central and Balkan groups. While Mediterraneans are near the farmer cluster, they are shifted towards the WHG (Figure 5A; unimputed PCA in Supplementary Figure 4B). ADMIXTURE analysis on all six populations supports this conclusion, showing significantly increased proportions of WHG ancestry in the Neolithic Mediterranean as compared with the other groups (maximum p=0.002 vs the Balkans, Fig. 5C). The average proportion of WHG ancestry in the Mediterranean is 11.4%; in the Balkans, 5.3%; in the South Central, 4.1%; and in the North Central, 1.1%. If there are significant PRS differences between Mediterranean and other populations, they are likely linked to this greater WHG ancestry and reflect genetic differences between WHG and other populations.

Computing PRS using clumping/thresholding, we find that the WHG have the lowest PRS of any population in our data (maximum p=0.002 vs Mediterranean), while Anatolians are similar to the Balkan and Central Europeans. Among individuals, proportions of WHG ancestry are strongly associated with decreased PRS (p=1.6×10^{-06}, β=-0.08cm/%). However, when we compute PRS with an infinitesimal LDpred2 model, Mediterranean PRS is intermediate between Neolithic Europeans and WHG. When we repeat the LDpred analysis using summary statistics computed from between-sibling GWAS (Howe et al., 2022) we find that the direction of Hunter-Gatherer PRS flips, and they have significantly greater PRS than the other groups (p=0.002) (Supplementary Figure 3A). The inconsistency of these results shows that the apparent PRS difference between WHG and Neolithic populations is highly sensitive to the PRS construction and summary statistics. This may indicate uncorrected population stratification in the non-sibling GWAS (Sohail et al., 2019; Berg et al., 2019). We therefore conclude that there is no strong evidence for a genetic difference in stature between Mediterranean and other Neolithic populations.

Figure 5: *A) Plot of ancient individuals projected into modern PCA space, including those of Mesolithic Western Hunter-Gatherer (WHG) and Anatolian Early Farmer ancestry. B) ADMIXTURE plot of K=2 ancestry groups showing the increased proportion of WHG ancestry in the Mediterranean individuals. C) Polygenic scores for each region including Mesolithic Hunter-Gatherers and Anatolian Early Farmers. D) Polygenic scores calculated from between-sibling summary statistics.*

4 Discussion

Interpretations of human stature variation through time remain confounded by the difficulty of separating genetic and environmental effects, obscuring trends. Accurately interpreting past stature variation not only allows us to understand these communities, but provides us with insights into the origin and evolution of modern health patterns. Height, while interesting in its own right, can also serve as a model trait for how to incorporate genetics and anthropological data into studies of human morphology and variation. Here, by integrating genetic, cultural, and environmental data, we are able to begin teasing apart the contributions of genetic and non-genetic factors in producing the observed phenotypic variation. We also illustrate the existing limitations of interpreting genetic data.

Overall, the Central and Balkan groups are genetically homogenous with similar levels of WHG admixture and polygenic scores, while Mediterranean individuals have more WHG ancestry (consistent with previous observations: Haak et al. (2015); Mathieson et al. (2015); Lipson et al. (2017)), which may be associated with lower PRS, though this relationship is not robust and may simply reflect residual population stratification. None of our populations show evidence for substantial genetic differences in height between sexes, which is expected given that there is little evidence for sex-specific genetic effects (Randall et al., 2013; Bernabeu et al., 2021; Zhu et al., 2022) (Supplemetary Figure 2). We can therefore largely exclude a genetic contribution to differences in stature between North Central individuals and other groups, while we find no strong evidence for a genetic contribution to shorter Mediterranean femora.

4.1 Sexual dimporphism in Central Europe reflects the effects of culture

Dietary differences between Southern and Northern Central European populations may indicate environmental stress in the north. In the early European Neolithic, the expansion of agriculture is thought to have been largely limited by poor soils and climate, as colder temperatures and decreased daylight made it increasingly difficult to grow early cereals (wheat, barley) and pulses (peas) (Bogaard, 2004; Schier, 2022), resulting in inconsistent, small harvests. The boundary to which these plants could be grown has been estimated to coincide with the northern limits of the LBK culture (Betti et al., 2020), and the majority of our Northern sites are concentrated near this climate edge in areas of fertile loess soils. However, examination of isotope values from herbivorous animals in our study regions (Dürrwächter et al., 2006; Oelze et al., 2011; Bentley et al., 2013; Denaire et al., 2017; Gillis et al., 2020; Knipper, 2020, data from) indicates that baseline values of δ^{15}N are elevated in the South Central region as compared to the North potentially due to differences in climate and the use of manure as fertilizer, and therefore at least some of the difference between Northern and Southern nitrogen values might be attributable to differences in climate and farming practices rather than diet. On the other hand, while remains of cattle and dairy production are documented in Northern LBK sites (Gillis et al., 2017; Salque et al., 2013), there is less archaeological

evidence for the presence of other wild or domesticated animals, indicating the people of this region
were highly reliant on plant foods (Nicklisch et al., 2016; Münster et al., 2018). Isotopic values
from other studies confirm the zooarchaeological evidence and show a higher proportion of plant
vs animal foods in the North, particularly domesticated cereal grains (Nicklisch et al., 2016; Oelze
et al., 2011). We conclude that our observed differences in Northern and Southern stable isotope
values probably reflects both dietary factors and differences in climate or farming practices.

It is therefore not surprising that the people living in the Northern Central region exhibit
evidence of increased stress due to potentially unreliable and lower quality food resources. Lower
protein consumption has been linked to decreased stature (Ghosh, 2016), and could be an indicator
of dietary stress. Diet can in some cases affect dimorphism (Gray & Wolfe, 1980), but while males
in the Southern Central and Balkan regions have higher nitrogen levels than females, the isotopic
signatures of North Central males and females are very similar, suggesting that this factor alone does
not explain reduced female stature in the North. Femur length and isotope values for individuals are
not significantly associated in our data, an indication that either diet has little effect on Neolithic
stature or stable isotopes do not capture the elements of diet relevant to height. Alternatively,
it is possible the range of variation in our data is too small to see this effect, or our sample of
individuals with both metric and isotopic data is not large enough. In addition, we only analyzed
adult samples and while the isotopic values of weaned children in the LBK fall within the range of
adults (Münster et al., 2018), it is possible that there could be sex differences in childhood diets
affecting femur growth.

Paleopathological analysis also indicates increased stress in the Northern population in the form
of increased incidence of linear enamel hypoplasia and cribra orbitalia. The causes of LEH formation
are varied and their appearance in the bioarchaeological record is generally interpreted as a non-
specific indication of childhood stress (Guatelli-Steinberg & Lukacs, 1999). Other archaeological
sites have reported a high instance of LEH with high sexual dimorphism ratios in areas of Neolithic
Europe, though the cause and meaning of these patterns was not explored (e.g, Lubell et al., 1990,
and references therein). It has been suggested that cribra orbitalia might also reflect childhood
stress, specifically anaemias, even when seen in adults (McFadden & Oxenham, 2020). Incidence of
both these stress indicators is higher in Northern compared to Southern Central Europe but is not
different between sexes in either group. The association between shorter femora and presence of
LEH appears to be driven by the females, suggesting a moderating factor causing a female-specific
effect despite equal incidence of LEH in both sexes.

While we see a general increase in stress shared between sexes in North Central Europe, typical
population-level stress responses usually show male vulnerability and female buffering effects (Ger-
man & Hochberg, 2020; Stini, 1985; Brauer, 1982). Our data indicate an opposite pattern in Central
Europe, and no evidence of a variable which acts upon females alone. However, the Northern pop-
ulation shows sexual dimorphism that is extreme by present-day standards. In modern samples,
the ratio of male to female height is 1.06-1.08 in most global populations (Gaulin & Boster, 1992),

though it is difficult to know how to compare height versus femur length ratios as the transformation
from metrics to stature scales differently in males and females. Based on 147 European individuals
from the past 100 years (data from Ruff, 2018), we find that the height ratio is very similar to
the ratio of femur length—typically within 1%. We therefore conclude that dimporphism ratios
in Southern Central (1.09) and Balkan (1.11) Europeans are elevated and the ratio in the North
Central region is exceptionally high (1.14). Few modern populations have height dimorphism ratios
as high as 1.10, and those that we could find in the literature come from India (Kanwar et al., 2011)
and the United Arab Emirates (Abdulrazzaq et al., 2008), both of which are countries known for
their cultural preferences and biases for male children.

We therefore hypothesize that the effects of high environmental stress in the North were mod-
ulated by culture. Other researchers have noted specific situations in which culture buffers males
against environmental effects and creates vulnerability in females: there is an association between
decreased female stature and polygyny in cultures around the globe (see Kanazawa & Novak, 2005);
female height was more influenced by economic conditions during infancy and early childhood than
males in lower-class 19th-century Europe (Baten & Murray, 2000); sexual dimorphism ratios in
modern Chile decreased after the institution of social and government programs to combat gender
inequality (Castellucci et al., 2021); and 20th-century female stature decreased in India during times
of environmental stress due to sexually disproportionate investment of scarce resources (Moradi &
Guntupalli, 2009). Strontium isotope values from LBK sites identify a large portion of LBK females
across the region as being non-local to those sites, though patterns in specific locations vary, broadly
indicating patrilocality and the potential for differential cultural treatments of females as compared
to males (Bentley et al., 2002, 2012; Hrnčíř et al., 2020). We suggest that culturally mediated
differences led to sex-specific stress responses in Neolithic Central Europe, *via* cultural practices
which either directly decrease female stature or, more likely, support catch-up growth preferentially
in males. Though dimorphism ratios in the South Central and Balkan regions are not as extreme
as in the North, they are elevated and also consistent with this pattern of male-bias, but response
is likely less exaggerated due to lower environmental stress conditions.

4.2 Mediterranean differences may have both genetic and environmental bases

In the Early Neolithic Mediterranean population we see decreased male stature and very low di-
morphism ratios (1.05) relative to other Neolithic populations. Mediterranean populations are
genetically distinct from other Early Neolithic groups with a higher proportion of WHG ancestry.
In some analyses, WHG ancestry proportion correlates with lower PRS for height. However, PRS
in the Mediterranean and WHG populations are sensitive to PRS construction method likely due to
residual population stratification in the GWAS. These inconsistent results mean that we can neither
confirm nor exclude the possibility of a genetic contribution to differences in stature between the
Mediterranean and other Early Neolithic populations. Even so, as with Central European popula-
tions, genetic effects alone would not explain the reduced dimorphism ratio, so we also need to

consider cultural/environmental effects.

The dimorphism ratio in the Mediterranean Neolithic is low, though not outside the range of
present-day populations (Gaulin & Boster, 1992). In fact, while males are relatively short, the
longest average female femur lengths of our data are in the Mediterranean. This reduction in
dimorphism is commonly seen in populations where the sexes experience an equal stress burden as
males tend to be more sensitive, decreasing their height, while females are biologically buffered and
stature remains consistent (German & Hochberg, 2020; Stini, 1985; Brauer, 1982; Garvin, 2012).
Although we do not have paleopathological stress data for the Mediterranean individuals in our
sample, published values for other Neolithic Mediterranean populations are generally similar to
those for South Central Europe (e.g., Papathanasiou, 2011, 2005; Silva & Cunha, 2001, except,
Cucina 2002). Dietary isotopes indicate that the Mediterranean diet differs in some aspects, with
increased δ^{13}C values compared to the other Neolithic populations but similar δ^{15}N values. Our
data indicates similar protein intake and low-level stress as other Neolithic populations, but do not
suggest any clear hypothesis for the difference in male stature between the Mediterranean and other
Neolithic groups. Possible differences in Mediterranean body proportions which are not captured
by femur length should also be mentioned as a caveat, though this likely would not be enough to
account for the differences in stature compared to the rest of Europe, and would not affect observed
dimorphism within the population. One possibility is that the Mediterranean experienced similar
levels of environmental stress as other Neolithic groups, but that they did not share the cultural
practices which preferentially supported males and increased female vulnerability.

4.3   Conclusion

By integrating genetic and anthropological data, we are able to begin to understand the contribu-
tions of genetics and environment to human variation, allowing us to better interpret the genetic,
environmental, and cultural landscapes of Neolithic Europe. Using this approach, we gain a deeper
understanding of the relationship between phenotypic plasticity and genetic architecture, which con-
strains the mechanisms by which human biology adapts to environment, culture, and genetic drift.
Our results are consistent with a model in which sexually dimorphic differences in femur length are
culturally and environmentally driven: relatively low dimorphism in the Mediterranean caused by
female buffering to environmental stress and less cultural male-preference, and high dimorphism
in Northern Central Europe caused by the interaction of relatively high environmental stress and
strong cultural male-preference. Some analyses suggest that differences in average femur length
between Central/Southeastern Europe and the Mediterranean are associated with differing genetic
ancestries, but less robust results, uncertainty about the transferrability of polygenic scores, and
questions of residual population stratification prevent us from interpreting this conclusively. In this
study we focused on the European Early Neolithic because of relative genetic, cultural, and environ-
mental homogeneity but, with more data, these approaches could be extended to other populations,
traits, and timescales to further explore the effects of human culture on biological variation.

462   **Data Availability** All data used in this analysis is provided in Supplementary Table 1. Original

463   ancient DNA data files can be downloaded from the resources provided in their cited publications.

464   Previously published osteological data can be found in their cited sources (Supplementary Table 1).

465   **Code Availability** R code used in this analysis is available for download on the Mathieson Lab

466   GitHub: $https://github.com/mathilab/Neolithic_height.git$.

References

Abdulrazzaq, Y., Moussa, M., & Nagelkerke, N. (2008). National growth charts for the United Arab Emirates. *Journal of Epidemiology*, *18*(6), 295–303.

Acsádi, G., Nemeskéri, J., & Balás, K. (1970). *History of human life span and mortality*. Akademiai kiado Budapest.

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Res*, *19*(9), 1655–64. doi: 10.1101/gr.094052.109.

Antonio, M. L., Gao, Z., Moots, H. M., Lucci, M., Candilio, F., Sawyer, S., et al. (2019). Ancient Rome: A genetic crossroads of Europe and the Mediterranean. *Science*, *366*(6466), 708–714. doi: 10.1126/science.aay6826.

Ash, A., Francken, M., Pap, I., Tvrdỳ, Z., Wahl, J., & Pinhasi, R. (2016). Regional differences in health, diet and weaning patterns amongst the first Neolithic farmers of central Europe. *Scientific reports*, *6*(1), 1–10.

Baten, J., & Murray, J. E. (2000). Heights of men and women in 19th-century Bavaria: economic, nutritional, and disease influences. *Explorations in Economic History*, *37*(4), 351–369.

Bentley, R. A., Bickle, P., Fibiger, L., Nowell, G. M., Dale, C. W., Hedges, R. E. M., et al. (2012). Community differentiation and kinship among Europe's first farmers. *Proceedings of the National Academy of Sciences*, *109*(24), 9326–9330. doi: 10.1073/pnas.1113710109.

Bentley, R. A., Bickle, P., Francken, M., Gerling, C., Hamilton, J., Hedges, R., et al. (2013). Baden-Württemberg. In P. Bickle, & A. Whittle (Eds.) *The First Farmers in Central Europe: Diversity in LBK Lifeways*. Oxford, UK: Oxbow Books.

Bentley, R. A., Price, T. D., Lning, J., Gronenborn, D., Wahl, J., & Fullagar, P. (2002). Prehistoric migration in Europe: strontium isotope analysis of early Neolithic skeletons. *Current Anthropology*, *43*(5), 799–804.

Berg, J. J., & Coop, G. (2014). A population genetic signal of polygenic adaptation. *PLoS Genet*, *10*(8), e1004412. doi: 10.1371/journal.pgen.1004412.

Berg, J. J., Harpak, A., Sinnott-Armstrong, N., Joergensen, A. M., Mostafavi, H., Field, Y., et al. (2019). Reduced signal for polygenic adaptation of height in UK Biobank. *Elife*, *8*, e39725. doi: 10.7554/eLife.39725.

Bernabeu, E., Canela-Xandri, O., Rawlik, K., & et al. (2021). Sex differences in genetic architecture in the UK Biobank. *Nature Genetics*, *53*, 1283–1289. doi: 10.1038/s41588–021–00912–0.

Betti, L., Beyer, R. M., Jones, E. R., Eriksson, A., Tassi, F., Siska, V., et al. (2020). Climate shaped
how Neolithic farmers and European hunter-gatherers interacted after a major slowdown from
6,100 BCE to 4,500 BCE. *Nature Human Behaviour*, *4*(10), 1004–1010.

Bickle, P., Arbogast, R.-M., Bentley, R. A., Fibiger, L., Hamilton, J., Hedges, R., et al. (2013a).
Alsace. In P. Bickle, & A. Whittle (Eds.) *The First Farmers in Central Europe: Diversity in*
LBK Lifeways. Oxford, UK: Oxbow Books.

Bickle, P., Bentley, R. A., Blesl, C., Fibiger, L., Hamilton, J., Hedges, R., et al. (2013b). Austria. In
P. Bickle, & A. Whittle (Eds.) *The First Farmers in Central Europe: Diversity in LBK Lifeways*.
Oxford, UK: Oxbow Books.

Bickle, P., & Whittle, A. (Eds.) (2013). *The First Farmers of Central Europe*. Oxford, UK: Oxbow.

Bogaard, A. (2004). *Neolithic farming in central Europe: an archaeobotanical study of crop hus-*
bandry practices. Routledge.

Brauer, G. (1982). Size sexual dimorphism and secular trend: indicators of subclinical malnutrition.
In R. Hall (Ed.) *Sexual Dimorphism in Homo sapiens: A Question of Size*. New York, NY:
Praeger.

Brunel, S., Bennett, E. A., Cardin, L., Garraud, D., Barrand Emam, H., Beylier, A., et al. (2020).
Ancient genomes from present-day France unveil 7,000 years of its demographic history. *Proceed-*
ings of the National Academy of Sciences, *117*(23), 12791–12798.

Castellucci, H., Viviani, C., Boccardo, G., Arezes, P., Bartsch, Á., Martínez, M., et al. (2021).
Gender inequality and sexual height dimorphism in Chile. *Journal of Biosocial Science*, *53*(1),
38–54.

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-
generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience*, *4*, 7. doi:
10.1186/s13742–015–0047–8.

Childebayeva, A., Rohrlach, A. B., Barquera, R., Rivollat, M., Aron, F., Szolek, A., et al. (2022).
Population genetics and signatures of selection in Early Neolithic European farmers. *Molecular*
Biology and Evolution, *39*(6), msac108.

Cox, S. L., Moots, H. M., Stock, J. T., Shbat, A., Bitarello, B. D., Nicklisch, N., et al. (2022).
Predicting skeletal stature using ancient DNA. *American Journal of Biological Anthropology*,
177(1), 162–174.

Cox, S. L., Ruff, C. B., Maier, R. M., & Mathieson, I. (2019). Genetic contributions to variation in
human stature in prehistoric Europe. *Proceedings of the National Academy of Sciences*, *116*(43),
21484–21492.

Cucina, A. (2002). Brief communication: diachronic investigation of linear enamel hypoplasia in prehistoric skeletal samples from Trentino, Italy. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, *119*(3), 283–287.

Denaire, A., Lefranc, P., Wahl, J., Bronk Ramsey, C., Dunbar, E., Goslar, T., et al. (2017). The cultural project: Formal chronological modelling of the early and middle neolithic sequence in lower alsace. *Journal of Archaeological Method and Theory*, *24*, 1072–1149.

Dürrwächter, C., Craig, O. E., Collins, M. J., Burger, J., & Alt, K. W. (2006). Beyond the grave: variability in Neolithic diets in Southern Germany. *Journal of Archaeological Science*, *33*(1), 39–48.

Fernandes, D. M., Mittnik, A., Olalde, I., Lazaridis, I., Cheronet, O., Rohland, N., et al. (2020). The spread of steppe and Iranian-related ancestry in the islands of the western Mediterranean. *Nature Ecology & Evolution*, *4*(3), 334–345.

Fregel, R., Méndez, F. L., Bokbot, Y., Martín-Socas, D., Camalich-Massieu, M. D., Santana, J., et al. (2018). Ancient genomes from North Africa evidence prehistoric migrations to the Maghreb from both the Levant and Europe. *Proceedings of the National Academy of Sciences*, *115*(26), 6774–6779.

Fu, Q., Posth, C., Hajdinjak, M., Petr, M., Mallick, S., Fernandes, D., et al. (2016). The genetic history of Ice Age Europe. *Nature*, *534*(7606), 200–205. doi: 10.1038/nature17993.

Gamba, C., Jones, E. R., Teasdale, M. D., McLaughlin, R. L., Gonzalez-Fortes, G., Mattiangeli, V., et al. (2014). Genome flux and stasis in a five millennium transect of European prehistory. *Nature communications*, *5*(1), 1–9.

Garvin, H. M. (2012). *The effects of living conditions on human cranial and postcranial sexual dimorphism*. Ph.D. thesis, The Johns Hopkins University.

Gaulin, S. J., & Boster, J. S. (1992). Human marriage systems and sexual dimorphism in stature. *American journal of physical anthropology*, *89*(4), 467–475.

German, A., & Hochberg, Z. (2020). Sexual dimorphism of size ontogeny and life history. *Frontiers in Pediatrics*.

Ghosh, S. (2016). Protein quality in the first thousand days of life. *Food and Nutrition Bulletin*, *37*, S14–S21.

Gillis, R. E., Eckelmann, R., Filipović, D., Müller-Scheeßel, N., Cheben, I., Furholt, M., et al. (2020). Stable isotopic insights into crop cultivation, animal husbandry, and land use at the Linearbandkeramik site of Vráble-Vel'ké Lehemby (Slovakia). *Archaeological and Anthropological Sciences*, *12*, 1–15.

Gillis, R. E., Kovačiková, L., Bréhard, S., Guthmann, E., Vostrovská, I., Nohálová, H., et al. (2017). The evolution of dual meat and milk cattle husbandry in Linearbandkeramik societies. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1860), 20170905.

González-Fortes, G., Jones, E. R., Lightfoot, E., Bonsall, C., Lazar, C., Grandal-d'Anglade, A., et al. (2017). Paleogenomic evidence for multi-generational mixing between Neolithic farmers and Mesolithic hunter-gatherers in the Lower Danube Basin. *Current Biology*, *27*(12), 1801–1810.

Gray, J. P., & Wolfe, L. D. (1980). Height and sexual dimorphism of stature among human societies. *American Journal of Physical Anthropology*, *53*(3), 441–456.

Guatelli-Steinberg, D., & Lukacs, J. R. (1999). Interpreting sex differences in enamel hypoplasia in human and non-human primates: Developmental, environmental, and cultural considerations. *American journal of physical anthropology*, *110*(S29), 73–126.

Haak, W., Lazaridis, I., Patterson, N., Rohland, N., Mallick, S., Llamas, B., et al. (2015). Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature*, *522*(7555), 207–11. doi: 10.1038/nature14317.

Harpak, A., & Przeworski, M. (2021). The evolution of group differences in changing environments. *PLoS Biol*, *19*(1), e3001072. doi: 10.1371/journal.pbio.3001072.

Hofmann, D., Pechtl, J., Bentley, R. A., Bickle, P., Fibiger, L., Grupe, G., et al. (2013). Southern Bavaria. In P. Bickle, & A. Whittle (Eds.) *The First Farmers in Central Europe: Diversity in LBK Lifeways*. Oxford, UK: Oxbow Books.

Hofmanová, Z., Kreutzer, S., Hellenthal, G., Sell, C., Diekmann, Y., Díez-del Molino, D., et al. (2016). Early farmers from across Europe directly descended from Neolithic Aegeans. *Proceedings of the National Academy of Sciences*, *113*(25), 6886–6891.

Howe, L. J., Nivard, M. G., Morris, T. T., Hansen, A. F., Rasheed, H., Cho, Y., et al. (2022). Within-sibship genome-wide association analyses decrease bias in estimates of direct genetic effects. *Nature genetics*, *54*(5), 581–592.

Hrnčíř, V., Vondrovskỳ, V., & Květina, P. (2020). Post-marital residence patterns in LBK: Comparison of different models. *Journal of Anthropological Archaeology*, *59*, 101190.

Hujić, A. (2016). *Das Kind in uns unter der Lupe der Isotopie, Allometrie und Pathologie. Zusammenhang zwischen $\delta^{15}N$ und $\delta^{13}C$ als Eiweißproxy und dem Knochenwachstum bei linienbandkeramischen Individuen aus Stuttgart-Mühlhausen, "Viesenhäuser Hof" und Schwetzingen unter Berücksichtigung verschiedener Indikatoren für Nährstoffversorgung*. Ph.D. thesis, Freie Universität. doi: 10.17169/refubium-5564.

Jiang, L., Zheng, Z., Qi, T., Kemper, K. E., Wray, N. R., Visscher, P. M., et al. (2019). A resource-efficient tool for mixed model association analysis of large-scale data. *Nat Genet*, *51*(12), 1749–1755. doi: 10.1038/s41588–019–0530–8.

Jones, E. R., Gonzalez-Fortes, G., Connell, S., Siska, V., Eriksson, A., Martiniano, R., et al. (2015). Upper Palaeolithic genomes reveal deep roots of modern Eurasians. *Nature communications*, *6*(1), 1–8.

Kanazawa, S., & Novak, D. L. (2005). Human sexual dimorphism in size may be triggered by environmental cues. *Journal of Biosocial Science*, *37*, 657–665.

Kanwar, S. C., Sabharwal, A., Bhadra, K., & Narang, A. (2011). Nationwide reference data for height, weight and body mass index of Indian schoolchildren. *Natlonal Medical Journal of India*, *24*(5), 269–77.

Knipper, C. (2020). Kohlenstoff- und Stickstoff-Isotopenanalysen an bandkeramischen Bestattungen vom Viesenhäuser Hof bei Stuttgart-Mühlhausen: Implikationen zur Ernährungsrekonstruktion, Geschlechtsspezifik und Siedlungsdynamik. In C. Meyer, P. Held, C. Knipper, & N. Nicklisch (Eds.) *Der Zahn der Zeit – Mensch und Kultur im Spiegel interdisziplinärer Forschung. Festschrift für Kurt W. Alt*. Halle(Saale): Veröffentlichungen des LDA und Archäologie Sachsen-Anhalt.

Lazaridis, I., Mittnik, A., Patterson, N., Mallick, S., Rohland, N., Pfrengle, S., et al. (2017). Genetic origins of the Minoans and Mycenaeans. *Nature*, *548*(7666), 214–218.

Lazaridis, I., Patterson, N., Mittnik, A., Renaud, G., Mallick, S., Kirsanow, K., et al. (2014). Ancient human genomes suggest three ancestral populations for present-day europeans. *Nature*, *513*(7518), 409–13. doi: 10.1038/nature13673.

Lillie, M. (2008). Vedrovice: demography and palaeopathology in an early farming population. *Anthropologie*, *46*(2/3), 135–152.

Lipson, M., Szecsenyi-Nagy, A., Mallick, S., Posa, A., Stegmar, B., Keerl, V., et al. (2017). Parallel palaeogenomic transects reveal complex genetic history of early European farmers. *Nature*, *551*(7680), 368–372. doi: 10.1038/nature24476.

Lubell, D., Jackes, M., & Meiklejohn, C. (1990). Archaeology and human biology of the Mesolithic-Neolithic transition in southern Portugal: a preliminary report. In *The Mesolithic in Europe. International Symposium. 3*, (pp. 632–640).

Marciniak, S., Bergey, C. M., Silva, A. M., Hałuszko, A., Furmanek, M., Veselka, B., et al. (2022). An integrative skeletal and paleogenomic analysis of stature variation suggests relatively reduced health for early European farmers. *Proceedings of the National Academy of Sciences*, *119*(15), e2106743119. doi: 10.1073/pnas.2106743119.

Marcus, J. H., Posth, C., Ringbauer, H., Lai, L., Skeates, R., Sidore, C., et al. (2020). Genetic history from the Middle Neolithic to present on the Mediterranean island of Sardinia. *Nature communications*, *11*(1), 1–14.

Martin, A. R., Kanai, M., Kamatani, Y., Okada, Y., Neale, B. M., & Daly, M. J. (2019). Clinical use of current polygenic risk scores may exacerbate health disparities. *Nat Genet*, *51*(4), 584–591. doi: 10.1038/s41588–019–0379–x.

Martiniano, R., Cassidy, L. M., O'Maolduin, R., McLaughlin, R., Silva, N. M., Manco, L., et al. (2017). The population genomics of archaeological transition in west iberia: Investigation of ancient substructure using imputation and haplotype-based methods. *PLoS Genet*, *13*(7), e1006852. doi: 10.1371/journal.pgen.1006852.

Mathieson, I., Alpaslan-Roodenberg, S., Posth, C., Szecsenyi-Nagy, A., Rohland, N., Mallick, S., et al. (2018). The genomic history of southeastern Europe. *Nature*, *555*(7695), 197–203. doi: 10.1038/nature25778.

Mathieson, I., Lazaridis, I., Rohland, N., Mallick, S., Patterson, N., Roodenberg, S. A., et al. (2015). Genome-wide patterns of selection in 230 ancient Eurasians. *Nature*, *528*(7583), 499–503. doi: 10.1038/nature16152.

McFadden, C., & Oxenham, M. F. (2020). A paleoepidemiological approach to the osteological paradox: Investigating stress, frailty and resilience through cribra orbitalia. *American Journal of Physical Anthropology*, *173*(2), 205–217.

Meyer, C., Knipper, C., Nicklisch, N., Münster, A., Kürbis, O., Dresely, V., et al. (2018). Early Neolithic executions indicated by clustered cranial trauma in the mass grave of Halberstadt. *Nature Communications*, *9*(1), 1–11.

Moradi, A., & Guntupalli, A. M. (2009). What does gender dimorphism in stature tell us about discrimination in rural India, 1930-1975. *Gender Bias: Health, Nutrition and Work*.

Münster, A., Knipper, C., Oelze, V. M., Nicklisch, N., Stecher, M., Schlenker, B., et al. (2018). 4000 years of human dietary evolution in central Germany, from the first farmers to the first elites. *PloS one*, *13*(3), e0194862.

Neale Lab (2018). UK Biobank GWAS Results. http://www.nealelab.is/uk-biobank. Accessed: 2021-10-20.

Nicklisch, N. (2017). *Spurensuche am Skelett: Paläodemografische und epidemiologische Untersuchungen an neolithischen und frühbronzezeitlichen Bestattungen aus dem Mittelelbe-Saale-Gebiet im Kontext populationsdynamischer Prozesse*. Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum.

Nicklisch, N., Ganslmeier, R., Siebert, A., Friederich, S., Meller, H., & Alt, K. W. (2016). Holes in teeth–dental caries in Neolithic and early bronze age populations in central Germany. *Annals of Anatomy-Anatomischer Anzeiger*, *203*, 90–99.

Nikitin, A., Stadler, P., & Kotova, e. a., N. (2019). Interactions between earliest Linearbandkeramik farmers and central European hunter gatherers at the dawn of European Neolithization. *Scientific Reports*, *9*, 19544. doi: 10.1038/s41598–019–56029–2.

O'Brien, D. M. (2015). Stable Isotope Ratios as Biomarkers of Diet for Health Research. *Annu Rev Nutr*, *35*, 565–594.

Oelze, V. M., Siebert, A., Nicklisch, N., Meller, H., Dresely, V., & Alt, K. W. (2011). Early Neolithic diet and animal husbandry: stable isotope evidence from three Linearbandkeramik (LBK) sites in Central Germany. *Journal of Archaeological Science*, *38*(2), 270–279.

Olalde, I., Mallick, S., Patterson, N., Rohland, N., Villalba-Mouco, V., Silva, M., et al. (2019). The genomic history of the Iberian Peninsula over the past 8000 years. *Science*, *363*(6432), 1230–1234.

Olalde, I., Schroeder, H., Sandoval-Velasco, M., Vinner, L., Lobón, I., Ramirez, O., et al. (2015). A common genetic origin for early farmers from Mediterranean Cardial and Central European LBK cultures. *Molecular biology and evolution*, *32*(12), 3132–3142.

Papathanasiou, A. (2005). Health status of the Neolithic population of Alepotrypa Cave, Greece. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, *126*(4), 377–390.

Papathanasiou, A. (2011). Health, diet and social implications in Neolithic Greece from the study of human osteological material. *Human bioarchaeology of the transition to agriculture*, (pp. 87–106).

Patterson, N., Price, A. L., & Reich, D. (2006). Population structure and eigenanalysis. *PLoS Genet*, *2*(12), e190. doi: 10.1371/journal.pgen.0020190.

Privé, F., Arbel, J., & Vilhjálmsson, B. J. (2020). LDpred2: better, faster, stronger. *Bioinformatics*, *36*(22-23), 5424–5431.

R Core Team (2021). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Randall, J. C., Winkler, T. W., Kutalik, Z., Berndt, S. I., Jackson, A. U., Monda, K. L., et al. (2013). Sex-stratified genome-wide association studies including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. *PLOS Genetics*, *9*(6), 1–19. doi: 10.1371/journal.pgen.1003500.

Richards, M. P., Montgomery, J., Nehlich, O., & Grimes, V. (2008). Isotopic analysis of humans and animals from Vedrovice. *Anthropologie*, *46*(2/3), 185–194.

Rivollat, M., Jeong, C., Schiffels, S., Küçükkalıpçı, İ., Pemonge, M.-H., Rohrlach, A. B., et al. (2020). Ancient genome-wide DNA from France highlights the complexity of interactions between Mesolithic hunter-gatherers and Neolithic farmers. *Science advances*, *6*(22), eaaz5344.

Roseman, C. C. (2004). Detecting interregionally diversifying natural selection on modern human cranial form by using matched molecular and morphometric data. *Proceedings of the National Academy of Sciences*, *101*(35), 12824–12829. doi: 10.1073/pnas.0402637101.

Rosenstock, E., Ebert, J., Martin, R., & Groß, M. (2019a). LiVES COstA Digest: checked and updated entries from the LiVES collection of Osteological Anthropometry. doi: 10.17171/2-12-2-1.

Rosenstock, E., Ebert, J., Martin, R., Hicketier, A., Walter, P., & Groß, M. (2019b). Human stature in the near east and europe ca. 10,000–1000 bc: its spatiotemporal development in a bayesian errors-in-variables model. *Archaeological and Anthropological Sciences*, *11*(10), 5657–5690. doi: 10.1007/s12520–019–00850–3.

Ruff, C. B. (Ed.) (2018). *Skeletal variation and adaptation in Europeans: Upper Paleolithic to the twentieth century*. Hoboken, NJ: John Wiley & Sons. (data available at https://www.hopkinsmedicine.org/fae/CBR.html).

Salque, M., Bogucki, P. I., Pyzel, J., Sobkowiak-Tabaka, I., Grygiel, R., Szmyt, M., et al. (2013). Earliest evidence for cheese making in the sixth millennium BC in northern Europe. *Nature*, *493*(7433), 522–525.

Savell, K. R., Auerbach, B. M., & Roseman, C. C. (2016). Constraint, natural selection, and the evolution of human body form. *Proc Natl Acad Sci U S A*, *113*(34), 9492–7. doi: 10.1073/pnas.1603632113.

Scheibner, A. (2016). *Prähistorische Ernährung in Vorderasien und Europa. Eine kulturgeschichtliche Synthese auf der Basis ausgewählter Quellen. Schriften zum Lebensstandard in der Vorgeschichte 1*. Leidorf: Rahden: Berliner Archäologische Forschungen.

Schier, W. (2022). Modes and Models of Neolithization in Europe: Comments to an Ongoing Debate. In P. Biehl, & E. Rosenstock (Eds.) *6000 BC: Transformation and Change in the Near East and Europe*, (pp. 372–392). Cambridge, UK: Cambridge University Press. doi: 10.1017/9781107337640.026.

Silva, A. M., & Cunha, E. (2001). Paleopathological study of the community exhumed from the Hipogeu of Monte Canelas I (Alcalar, Portugal). In *Actas del V Congreso Nacional de Paleopatologia*, (pp. 353–356).

728 Sohail, M., Maier, R. M., Ganna, A., Bloemendal, A., Martin, A. R., Turchin, M. C., et al. (2019).
729   Polygenic adaptation on height is overestimated due to uncorrected stratification in genome-wide
730   association studies. *Elife*, *8*, e39702. doi: 10.7554/eLife.39702.

731 Stini, W. (1985). Growth rates and sexual dimorphism in evolutionary perspective. In W. Stini
732   (Ed.) *The Analysis of Prehistoric Diets*. Orlando, FL: Academic Press.

733 Valdiosera, C., Günther, T., Vera-Rodríguez, J. C., Ureña, I., Iriarte, E., Rodríguez-Varela, R.,
734   et al. (2018). Four millennia of Iberian biomolecular prehistory illustrate the impact of prehistoric
735   migrations at the far end of Eurasia. *Proceedings of the National Academy of Sciences*, *115*(13),
736   3428–3433.

737 Villalba-Mouco, V., van de Loosdrecht, M. S., Posth, C., Mora, R., Martínez-Moreno, J., Rojo-
738   Guerra, M., et al. (2019). Survival of Late Pleistocene hunter-gatherer ancestry in the Iberian
739   Peninsula. *Current Biology*, *29*(7), 1169–1177.

740 Whitlock, M. C. (2008). Evolutionary inference from QST. *Mol Ecol*, *17*(8), 1885–96. doi:
741   10.1111/j.1365–294X.2008.03712.x.

742 Whittle, A., Ander, A., Bentley, R. A., Bickle, P., Cramp, L., Domboróczki, L., et al. (2013a).
743   Hungary. In P. Bickle, & A. Whittle (Eds.) *The First Farmers in Central Europe: Diversity in*
744   *LBK Lifeways*. Oxford, UK: Oxbow Books.

745 Whittle, A., Bentley, R. A., Bickle, P., Dočkalová, Fibiger, L., Hamilton, J., et al. (2013b). Moravia
746   and western Slovakia. In P. Bickle, & A. Whittle (Eds.) *The First Farmers in Central Europe:*
747   *Diversity in LBK Lifeways*. Oxford, UK: Oxbow Books.

748 Yengo, L., Sidorenko, J., Kemper, K. E., Zheng, Z., Wood, A. R., Weedon, M. N., et al. (2018).
749   Meta-analysis of genome-wide association studies for height and body mass index in approx-
750   imately 700000 individuals of European ancestry. *Hum Mol Genet*, *27*(20), 3641–3649. doi:
751   10.1093/hmg/ddy271.

752 Zhu, C., Ming, M. J., Cole, J. M., Kirkpatrick, M., & Harpak, A. (2022). Amplification is the primary
753   mode of gene-by-sex interaction in complex human traits. *bioRxiv*. doi:10.1101/2022.05.06.490973.