Socio-cultural practices may have affected sexual dimorphism in stature in Early Neolithic Europe

Samantha L Cox[1,2*], Nicole Nicklisch[3], Michael Francken[4], Joachim Wahl[5], Harald Meller[6], Wolfgang Haak[7], Kurt W Alt[3], Eva Rosenstock[8], and Iain Mathieson[1*]

[1]*Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA*

[2]*Physical Anthropology Section, Penn Museum, University of Pennsylvania, Philadelphia, PA, USA*

[3]*Center of Natural and Cultural Human History, Danube Private University, Krems-Stein, Austria*

[4]*State Office for Cultural Heritage Management Baden-Württemberg, Osteology, Konstanz, Germany*

[5]*Institut für Naturwissenschaftliche Archäologie, AG Paläoanthropologie, Tübingen, Germany*

[6]*State Office for Heritage Management and Archaeology Saxony-Anhalt, State Museum of Prehistory, Halle, Germany*

[7]*Department of Archaeogenetics, Max Planck Institute for the Science of Human History, Jena, Germany*

[8]*Bonn Center for ArchaeoSciences, Universität Bonn, Bonn, Germany*

*Correspondence to SLC (coxsl@sas.upenn.edu) or IM (mathi@pennmedicine.upenn.edu)

Abstract

The rules and structure of human culture impact health and disease as much as genetics or the natural environment. To study the origin and evolution of these patterns, we take a multidisciplinary approach combining ancient DNA, skeletal metrics, paleopathology, and stable isotopes. Our analysis focuses on cultural, environmental, and genetic contributions to variation in stature in four populations of Early Neolithic Europe. In Central Europe, low female stature is likely due to male preference in resource allocation under conditions of stress. In contrast, shorter male stature in Mediterranean populations may reflect a lack of preference. Our analysis suggests that biological consequences of sex-specific inequities can be linked to culture as early as 7000 years before present. Understanding these patterns is key to interpreting the evolution of genetic and socio-cultural determinants of health, and our results show that culture, more than environment or genetics, drove height disparities in Early Neolithic Europe.

1 Introduction

Human skeletal variation reflects varying combinations of genetic, cultural, and environmental factors. While there are many links between culture and health in the modern world, the history and evolution of these relationships are not always well established. Due to the entanglement of these factors, our ability to draw conclusions about their effects has been limited in archaeological data. With the recent advent of ancient DNA sequencing technology, genetic information from ancient populations has become increasingly available. However, attempting to analyze changing patterns of

variation based solely on genetic data is difficult—genotypes do not necessarily equate to phenotypes
due to the effect of the environment.[1] Similarly, while it is tempting to predict phenotypic changes
in ancient people based on their genetic variation, this is currently challenging as genetic effects are
not always transferable across populations.[2] Our solution is to integrate these complementary fields
to construct multidisciplinary analyses with phenotype, genotype, culture, and environment data
from ancient human populations. This approach allows us to begin to separate the effects of these
variables and reveal the interactions between genes, environment, and culture which are critical in
shaping human health and variation.

Many traits of interest, including height, are highly polygenic, with thousands of independent
genetic variants contributing significantly to heritability. One common approach to addressing the
role of genetics in morphological change is to compare patterns of phenotypic variation with genetic
ancestry or genome-wide patterns of genetic variation.[3–5] However, even for highly polygenic traits
like height, genome-wide variation may not be directly relevant, leading to spurious associations
between genetic effects, ancestry, and environmental confounds. For example, if a population is tall
and has a high proportion of ancestry from Neolithic sources, it could be concluded that Neolithic
ancestry is associated with "genetic tallness"; however, the effects could equally be non-genetic
and related to lifestyle changes associated with agriculture. An alternative approach is to focus
only on genetic variation that is known to be associated with a specific trait.[6,7] Effect sizes for
these trait-related variants estimated from genome-wide association studies (GWAS) of present-day
individuals can be combined with genetic data from ancient individuals to calculate polygenic risk
scores (PRS), which can be thought of as estimated genetic values for the phenotype. In European
ancestry populations, polygenic scores for height can explain up to 25% of phenotypic variation
in present-day individuals,[8] and 6-8% of variation in ancient individuals.[9,10] On a broad scale,
temporal changes in polygenic score over time in Europe are qualitatively consistent with changes
in stature as inferred from the skeletal record,[11] while local deviations from this pattern provide
evidence of environmental effects.[10,11]

Analyses of human populations over tens of thousands of years involve individuals that are
diverse in genetic ancestry, environment, and culture and it is challenging to exclude the possibility
of confounding by unmeasured variables. We therefore focus specifically on the European Early
Neolithic. One of the most studied periods in prehistory, it represents a fundamental shift in
technology, culture, and genetics. In particular, the *Linearbandkeramik* (LBK) culture of Central
Europe is one of the most comprehensively documented Early Neolithic cultures, with an abundance
of excavated settlements and cemeteries.[12] LBK groups tended to choose settlement locations based
on the presence of rich loess soils for farming, and the northern edge of these soils appears to
delineate the northern limit of LBK sites.[13,14] Bioarchaeological evidence indicates broad regional
differences between individuals from northern settlements in this agricultural boundary zone vs
southern settlements in a climate zone that was more comfortable for Neolithic crops.[15,16] Based
on this, we divided our Central European group into Northern (above 50°N latitude) and Southern

(below 50°N) populations. The Mesolithic hunter-gatherer population in Central Europe made a
limited genetic contribution to the LBK population, whose members harbor only traces of hunter-
gatherer admixture.[7,17,18] Contemporary populations from southeastern Europe have similarly low
levels of hunter-gatherer ancestry.[19] In contrast, Neolithic southern European populations associated
with the Cardial and Impressed Ware cultures followed a separate migration route (Figure 1),
occupied a milder climate zone, and carried more Mesolithic ancestry.[17,20] Individuals in this region
tend to be shorter than those from Central Europe and combined with their admixed ancestry this
has led to suggestions of a genetic basis for decreased statures in this region.[7,21]

By comparing and contrasting these four closely related archaeological populations, we aim to
investigate how differences in environment and genetics combine to produce observed phenotypes.
We collected genetic data, skeletal metrics, paleopathology, and dietary stable isotopes to begin
separating the effects of each on Neolithic stature trends. By specifically investigating and con-
trolling for the effects of genetics in these samples, we are able to provide nuanced interpretations
of height variation, gain a better understanding of the aspects of height which are controlled by
genetics or environment, and show evidence for sex-specific cultural effects which modify the ge-
netically predicted patterns. We illustrate the strengths of leveraging multidisciplinary datasets,
and indicate caution when analyzing genotype-phenotype relationships without complete data, es-
pecially for traits which are not preserved in the archaeological record and cannot be directly tested.
This integrated analysis highlights the role of plasticity in morphology, and establishes culturally
mediated disparities at least as early as the European Neolithic.

92   2   Results

93   2.1   Distribution of stature, polygenic scores, and stable isotope values

We collected either genetic, dietary stable isotope, paleopathology, or skeletal metric data from
1282 individuals associated with the archaeological LBK culture in Northern and Southern Central
Europe dated to between 7700-6900 years before present (BP), as well as 135 individuals from
the southeastern (Balkan) and 160 individuals from the southern (Mediterranean) regions dated
between 8000-6000 BP. All individuals included in the analysis have at least one of the four data
types available (Materials and Methods; Figure 2; Supplementary Figure 1).

Observed patterns of femur length vary between sexes and populations. Male femora show no
apparent difference between the Central and Balkan regions (p=0.56), but Mediterranean males are
significantly shorter (p=5.5×10^{-7}, β=-1.44cm). Conversely, female femora show a different pattern,
with no significant difference between Mediterranean, South Central (p=0.97), and Balkan (p=0.54)
populations, but substantially shorter values in the North Central population (p=9×10^{-07}, β=-
2.0cm) (Figure 3A). Differences between male and female femur lengths are highly significant in all
popultions (p<2×10^{-16}). In contrast to the differences in femoral lengths, polygenic scores for height
are very similar between all populations (pairwise t-tests p > 0.9) using the clumping/thresholding

Figure 1: *This schematic illustration highlights the two main migration routes from Anatolia to Europe during the Early Neolithic period.[22] Populations followed two routes: southern, along the Mediterranean coast (including sea routes, generalized here by dashed blue lines) where they admixed with existing hunter-gatherer populations; or northern, through the Balkans and into Central Europe, with only limited hunter-gatherer admixture. We analyze patterns within the Linearbandkeramik culture, dividing it into Northern and Southern Central European groups.*

Figure 2: *Upper row: Sites used for genetic (A) and skeletal (B) data in the analysis. The Central European population is split into Northern and Southern groups at 50°N latitude (emphasized). Lower row: (C) imputed genetic data projected into the PCA space of 777 modern Eurasian individuals (grey points). (D) sample numbers by years before present (years BP) for skeletal (right) and genetic (left) data.*

108 PRS construction (Figure 3B). PRS constructed with LDpred show Mediterranean individuals to be
109 shorter than the other populations (p=0.002; Supplementary Figure 5). However, PRS constructed
110 using summary statistics derived from between-sibling analysis finds similar genetic values in all
111 populations with both PRS construction methods, so we conclude that apparent lower Mediter-
112 ranean PRS may be due to population stratification in the GWAS data and may not reflect a
113 true genetic difference. There are no significant differences between male and female PRS in any
114 population (Figure 3B), providing no evidence for a genetic basis to this dimorphism.

115   Signatures of δ^{13}C and δ^{15}N suggest different dietary patterns in each of the analyzed groups
116 (Figure 3C, D). Both the Mediterranean and Balkan groups are significantly distinct from the
117 Central in δ^{13}C (multiple comparisons, maximum p=4.0×10^{-16}) and δ^{15}N (multiple comparisons,
118 maximum p=7.7×10^{-13}) values. Generally, the Balkan population is characterized by high δ^{15}N
119 values, while Mediterranean populations show high δ^{13}C relative to the Central Europeans (Figure
120 3C). The exception to this pattern is a cluster of individuals, classified as Balkan in our analysis,
121 which overlaps with the North Central population as well as some of the Mediterranean. These
122 points represent individuals from present-day Greece and indicate that the diets of these peoples
123 might better be classified as Mediterranean than Balkan. Nitrogen values are generally elevated in
124 males compared to females (Figure 3D), but this difference is only significant in the Mediterranean
125 (p=0.035).

2.2   Patterns of non-genetic factors in Central Europe

127 The most dramatic observation is the difference in female stature and consequent sexual dim-
128 porphism in Northern compared to Southern Central Europe. Female femora in the North are
129 significantly shorter than female femora in the South (p=2.7×10^{-6}, β=1.7cm), while male femora
130 are highly similar (p=0.35) (Figure 3A). On average, male femora from the North are about 13%
131 longer than female femora, Southern Central and Balkan male femora are about 9% and 11% longer
132 respectively, and Mediterranean male femora are only 5% longer (Figure 3A). These values are
133 reduced slightly when calculated using estimated statures instead of femora (North Central: 10%,
134 South Central: 7%, Balkans: 8%, Mediterranean: 4%), possibly due to error associated with stature
135 estimation (see Ref. 9) and body proportions, or because the relationship between femur length and
136 stature is different between males and females. Where we have both genetic and metric data for
137 the same individuals, there is a qualitative relationship between femur length and PRS; PRS tends
138 to increase as femur lengths increase (Supplementary Figure 3B). However, the effect of PRS on
139 femur length is only marginally significant (p=0.05), likely due to the small number of individuals
140 with both types of data available in the sample (n=55).

141   Overall, trends in dietary stable isotopes show that both males and females in Southern Cen-
142 tral Europe have significantly higher δ^{15}N (male p=1.3×10^{-9}, β=0.83‰; female p=5.3×10^{-9},
143 β=0.87‰) and lower δ^{13}C (male p=3×10^{-4}, β=-0.30‰; female p=8.1×10^{-7}, β=-0.38‰) as
144 compared to the North. However, while males in both regions qualitatively have higher nitrogen,

Figure 3: *Solid bars across the tops of plots indicate significant differences between male populations by pairwise t-test; solid bars below plots indicate significant differences between female populations by pairwise t-test; p-values < 0.05 (*), < 0.01 (**) and < 0.001 (***); black points indicate the mean of each group; and vertical bars show mean standard error. A) Femur length in the four population: values to the right of the vertical dotted line are the difference between the mean male and female femora; values to the left are the sexual dimorphism ratios of male/female femur lengths for each population. B) Polygenic scores for the four populations show similar scores for individuals across all regions. Differences between male and female PRS are not significant. C) Plot of $\delta^{13}C$ (x-axis) and $\delta^{15}N$ (y-axis) dietary stable isotopes for the four populations: individuals from the Balkans are distinguished by high nitrogen values, while those in the Mediterranean generally have higher carbon. D) Sex differences in $\delta^{15}N$ values by sex for each population: $\delta^{15}N$ values are slightly higher for males in all populations, but this difference is only significant in the Mediterranean (p=0.035).*

Figure 4: *Evidence of environmental stress in Northern Central Europe. A) Differences in $\delta^{13}C$ (right) and $\delta^{15}N$ (left) values. Overall, the South has higher nitrogen values than the North ($p=6.8 \times 10^{-13}$), and lower carbon ($p=5.3 \times 10^{-15}$); within each population, the difference in isotopes between sexes is not significant. B) Proportion of linear enamel hypoplasias. The South has significantly less than the North ($p=0.001$). C) Presence of linear enamel hypoplasia is significantly associated with shorter femora ($p=0.02$); differences in prevalence between sexes are not significant.*

the interaction effect between sexes is not significant in either North or South (Figure 4A), indicat-
ing that the difference between male and female values in each region is not significant. There is
no difference in carbon values between sexes. For individuals with both stature and stable isotope
values, we find no statistically significant relationship between femur length and δ^{15}N or δ^{13}C in
either Central group, separately or combined, but the sample is small.

We do, however, find a statistically significant relationship between presence of linear enamel
hypoplasias (LEH) and shorter femora, suggesting that LEH may reflect an underlying variable in
childhood that also affects stature (p=0.021, β=-1.0cm)(Figure 4C). Both males and females from
the North are more likely to have LEH than individuals living in the South (p=0.002). Indeed, over
50% of the Northern sample have LEH while they are only present in about 20% of the Southern
(Figure 4B). There is no significant difference between the number of males and females with LEH in
either region. Though the interaction effect between sex and LEH on femur length is not significant,
qualitatively the effect of LEH on femur length appears greater in females than in males (Figure
4C). When the sexes are analyzed separately, females with LEH do have significantly shorter femora
than those without (p=0.018, β=-1.46), which is not the case for males (p=0.479). We hypothesize
that the relationship between LEH and femur length is driven by females, but we lack an adequate
sample size to detect the interaction effect in the full model. Incidence of cribra orbitalia is also
significantly higher in the Northern region than in the Southern (p=1.8×10^{-6}), though there is no
relationship with femur length. There are no significant trends related to the presence of porotic
hyperostosis.

In summary, comparison of Northern and Southern Central Europe identifies no predicted genetic
difference in stature, which is consistent with male but not female femur length. This suggests a

non-genetic basis for reduced female stature. Stable isotope data and skeletal stress indicators
suggest lower protein intake and more general stress in the North; however, males and females
overall appear equally affected by these variables. Despite a similar number of hypoplasias in both
sexes, shorter femora in females suggest that increased general stress, due to other unmeasured
environmental or cultural factors, leads to a female-specific reduction in stature.

172 2.3 Patterns of genetic ancestry in the Mediterranean

In contrast to Northern Central Europe, Mediterranean Neolithic males are shorter than other
groups, but females are not. PCA indicates that individuals from the Central regions and the
Balkans share similar genetic ancestry while those from the Mediterranean are distinct (Figure 2C;
unimputed PCA in Supplementary Figure 4A), a difference known to be due to higher levels of
hunter-gatherer ancestry in the Mediterranean.[17] We therefore additionally compared our samples
to Mesolithic individuals of Western Hunter-Gatherer (WHG) ancestry, as well as individuals from
early Neolithic Anatolia. These two groups represent source populations for the two largest ancestry
components in Europe at this time.[7,17]

On the PCA plots of these extended data, Neolithic Anatolians cluster with the Central and
Balkan groups. While Mediterraneans are near the farmer cluster, they are shifted towards the
WHG (Figure 5A; unimputed PCA in Supplementary Figure 4B). ADMIXTURE analysis of all six
populations supports this conclusion, showing significantly increased proportions of WHG ancestry
in the Neolithic Mediterranean as compared with the other groups (maximum p=0.002 vs the
Balkans, Fig. 5C). The average proportion of WHG ancestry in the Mediterranean is 11.4%; in the
Balkans, 5.3%; in the South Central, 4.1%; and in the North Central, 1.1%. If there are significant
PRS differences between Mediterranean and other populations, they are likely linked to this greater
WHG ancestry and reflect genetic differences between WHG and other populations.

Computing PRS using clumping/thresholding, we find that the WHG have the lowest PRS of
any population in our data (maximum p=0.002 vs Mediterranean), while Anatolians are similar to
the Balkan and Central Europeans. Among individuals, proportions of WHG ancestry are strongly
associated with decreased PRS (p=1.6×10^{-06}, β=-0.08cm/%). However, when we compute PRS
with an infinitesimal LDpred2 model, Mediterranean PRS is intermediate between Neolithic Euro-
peans and WHG. When we repeat the LDpred analysis using summary statistics computed from
between-sibling GWAS,[23] we find that the direction of Hunter-Gatherer PRS flips, and they have
significantly greater PRS than the other groups (p=0.002) (Supplementary Figure 3A). The in-
consistency of these results shows that the apparent PRS difference between WHG and Neolithic
populations is highly sensitive to the PRS construction and summary statistics. This may indicate
uncorrected population stratification in the non-sibling GWAS.[24,25] We therefore conclude that
there is no strong evidence for a genetic difference in stature between Mediterranean and other
Neolithic populations.

Figure 5: *A) Ancient individuals projected into modern PCA space, including those of Mesolithic Western Hunter-Gatherer (WHG) and Anatolian Early Farmer ancestry. B) ADMIXTURE plot of K=2 ancestry groups showing the increased proportion of WHG ancestry in Mediterranean individuals. C) Polygenic scores for each region including Mesolithic Hunter-Gatherers and Anatolian Early Farmers. D) Polygenic scores calculated from between-sibling summary statistics.*

3 Discussion

Understanding the causes of past stature variation not only allows us to understand ancient communities, but may also provide us with insights into the origin and evolution of modern health patterns. However, interpretations of human stature variation through time remain confounded by the difficulty of separating genetic and environmental effects, obscuring trends. Recently, several researchers have begun to compile multivariate datasets for the purposes of understanding human stature.[10,26–28]; however, many of these analyses do not directly take genetic effects into account,[26,28] or cover very broad temporal or geographic ranges.[10,27] In contrast, we aim to understand these processes on a finer scale to better interpret the outcomes of biological and environmental/cultural interactions. For instance, previous studies of stature variation from the Mesolithic to Neolithic indicated that Neolithic individuals were not achieving their genetic height potential,[10] but our analyses suggest that this effect might be heterogeneous, pertaining more to some locations and portions of society than others. Finally, while interesting in its own right, height can also serve as a model trait for how to incorporate genetics and anthropological data into studies of human morphology and variation. Here, by integrating genetic, cultural, and environmental data, we are able to begin teasing apart the contributions of genetic and non-genetic factors in producing the observed phenotypic variation. We also illustrate the existing limitations of interpreting genetic data.

Overall, the Central and Balkan groups are genetically homogenous with similar levels of hunter-gatherer admixture and polygenic scores, while Mediterranean individuals have more hunter-gatherer ancestry (consistent with previous observations[7,17,18]). This may be associated with lower PRS, though this relationship is not robust and may simply reflect residual population stratification in the GWAS. None of our populations show evidence for substantial genetic differences in height between sexes (Supplemetary Figure 2), which is expected given that there is little evidence for sex differences in ancestry, or of sex-specific genetic effects on stature.[29–31] We can therefore largely exclude a genetic contribution to differences in stature between North Central individuals and other groups, while we find no strong evidence for a genetic contribution to shorter Mediterranean stature.

3.1 Sexual dimorphism in Central Europe reflects the effects of culture

Dietary differences between Southern and Northern Central European populations may indicate environmental stress in the North. In the early European Neolithic, the expansion of agriculture is thought to have been largely limited by poor soils and climate, as colder temperatures and decreased daylight made it increasingly difficult to grow early cereals (wheat, barley) and pulses (peas).[32,33] The boundary to which these plants could be grown has been estimated to coincide with the northern limits of the LBK culture,[13,34] and the majority of our Northern sites are concentrated near this climate edge in areas of fertile loess soils.[14] However, as there are many nuances which affect the interpretation of stable isotope values, especially between populations, differences between our

239 Northern and Southern groups may not be as dramatic as they appear. An examination of isotope
240 values from herbivorous animals in our study regions (using data from Refs. 16,35–39) indicates
241 that baseline values of $\delta^{15}N$ are elevated in the South Central region as compared to the North,
242 potentially due to differences in climate and the use of manure as fertilizer. Therefore some portion
243 of the difference between Northern and Southern nitrogen values might be attributable to variation
244 in climate and farming practices rather than diet. Differences in carbon values between populations
245 can be similarly sensitive to environment and we feel that interpreting the carbon results would
246 be difficult without a more in-depth isotopic analysis. However, isotopic values from other studies
247 show a higher proportion of plant vs animal foods in the North, particularly domesticated cereal
248 grains.[15,36,40] Additionally, the available archaeological evidence supports some level of dietary
249 difference between the Northern and Southern regions. While remains of cattle and dairy production
250 are documented in Northern LBK sites,[41,42] there is less archaeological evidence for the presence of
251 other wild or domesticated animals that are seen in the South, indicating the people of this region
252 were highly reliant on plant foods.[40,43] We conclude that our observed differences in Northern and
253 Southern stable isotope values probably reflect both dietary factors and differences in climate or
254 farming practices.

255 It is therefore not surprising that people of the Northern Central region exhibit evidence of
256 increased stress potentially due to unreliable and lower quality food resources. Lower protein con-
257 sumption could be an indicator of dietary stress and has been linked to decreased stature.[44] Diet
258 can affect dimorphism in some cases,[45] but the isotopic signatures of males and females in the North
259 Central, South Central, and Balkan regions are very similar, suggesting that this factor alone does
260 not explain reduced female stature in the North. Femur length and isotope values for individuals are
261 not significantly associated in our data, an indication that either diet has little effect on Neolithic
262 stature or stable isotopes do not capture the elements of diet relevant to height. Alternatively, it is
263 possible the range of variation in our data is too small to see this effect, or our sample of individuals
264 with both metric and isotopic data is not large enough. In addition, we only analyzed adult samples
265 and while the isotopic values of weaned children in the LBK fall within the range of adults,[43] it
266 is possible that there could be sex differences in childhood diets affecting femur growth. Future
267 studies incorporating collagen from long bones or teeth, rather than from ribs as we have here,
268 would give dietary evidence with greater time depth, and might be able to provide more nuanced
269 interpretations in the absence of a known-sex sub-adult population.

270 Paleopathological analysis also indicates increased stress in the Northern population in the form
271 of increased incidence of linear enamel hypoplasia and cribra orbitalia. The causes of LEH formation
272 are varied and their appearance in the bioarchaeological record is generally interpreted as a non-
273 specific indication of childhood stress.[46] Other archaeological sites have reported a high instance of
274 LEH with high sexual dimorphism ratios in areas of Neolithic Europe, though the cause and meaning
275 of these patterns was not explored (e.g. Ref. 47 and references therein). It has been suggested that
276 cribra orbitalia might also reflect childhood stress, specifically anaemias, even when seen in adults.[48]

Our results are consistent with others who have considered the same paleopathologies and found
a qualitative relationship between presence of paleopathology and shorter femora.[10] In our data,
incidence of both LEH and cribra orbitalia are higher in Northern compared to Southern Central
Europe, but are not different between sexes in either group. The association between shorter femora
and presence of LEH appears to be driven by females, suggesting a moderating factor causing a
female-specific effect despite equal incidence of LEH in both sexes.

While we see a general increase in stress shared between sexes in North Central Europe, typical
population-level stress responses usually show male vulnerability and female buffering effects.[49–51]
Though the exact causes and mechanisms are not well understood, female biology tends to have a
less extreme response, or is "buffered", to many diseases[52–54] and environmental changes[55] compared
to males. Our data indicate an opposite pattern in Central Europe, and no evidence of a variable
which acts upon females alone. However, the Northern population shows sexual dimorphism that
is extreme by present-day standards. In most modern global populations the ratio of male to
female height is 1.06-1.08[56] (ratios in Ref. 56 range up to 1.12, but population locations or cultural
affiliations are not given, see Ref. 45), though it is difficult to know how to compare height versus
femur length ratios as the transformation from metrics to stature scales differently in males and
females. Based on 147 European individuals from the past 100 years (using data from Ref. 57),
we find that the height ratio is very similar to the ratio of femur length—typically within 1%. We
therefore conclude that dimporphism ratios in Southern Central (1.09) and Balkan (1.11) Europeans
are elevated and the ratio in the North Central region is exceptionally high (1.14). Few modern
populations have height dimorphism ratios as high as 1.10, and those that we could find in the
literature come from India[58] and the United Arab Emirates,[59] both countries known for their
cultural preferences and biases for male children.[60]

We therefore hypothesize that the effects of high environmental stress in the North were mod-
ulated by culture. Other researchers have noted specific situations in which culture buffers males
against environmental effects and creates vulnerability in females: there is an association between
decreased female stature and polygyny in cultures around the globe;[61] female height was more
influenced by economic conditions during infancy and early childhood than males in lower-class
19th-century Europe;[62] sexual dimorphism ratios in modern Chile decreased after the institution
of social and government programs to combat gender inequality;[63] and 20th-century female stature
decreased in India during times of environmental stress due to sexually disproportionate investment
of scarce resources.[60] In LBK sites, strontium isotope values show that females are more likely to
be non-local compared males, suggesting patrilocality and potential differences in cultural treat-
ment of females.[14,64] In parallel to our evidence for higher biological variation in females, ongoing
discussion about the relationship between biological sex and the formation of gendered identities
in the LBK suggests more variation in the roles and identities of females compared to males.[65] We
therefore suggest that culturally mediated differences led to sex-specific stress responses in Neolithic
Central Europe *via* cultural practices which either directly decreased female stature or, more likely,

supported catch-up growth preferentially in males. Though dimorphism ratios in the South Central
and Balkan regions are not as extreme as in the North, they are elevated and also consistent with
this pattern of male bias, but response is likely less exaggerated due to lower environmental stress
conditions.

3.2 Mediterranean differences may have both genetic and environmental bases

In the Early Neolithic Mediterranean population we see decreased male stature and low dimorphism
ratios (1.05) relative to other Neolithic populations. Mediterranean populations are genetically dis-
tinct from other Early Neolithic groups with a higher proportion of WHG ancestry. In some analyses,
WHG ancestry proportion correlates with lower PRS for height. However, PRS in the Mediterranean
and WHG populations are sensitive to PRS construction method likely due to residual population
stratification in the GWAS. These inconsistent results mean that we can neither confirm nor exclude
the possibility of a genetic contribution to differences in stature between the Mediterranean and
other Early Neolithic populations, though on balance we find the likelihood for a substantial genetic
contribution to be low. Even if it were not, the genetic effects alone could not explain the reduced
dimorphism ratio, emphasizing the need to also consider cultural/environmental effects.

While the dimorphism ratio in the Mediterranean Neolithic is low, it is not outside the range of
present-day populations.[56] In fact, while males are relatively short, the longest average female femur
lengths of our data are in the Mediterranean. This reduction in dimorphism is commonly seen in
populations where the sexes experience an equal stress burden: as males tend to be more sensitive,
decreasing their height, females are biologically buffered and stature remains consistent.[49–51,66]
Although we do not have paleopathological stress data for the Mediterranean individuals in our
sample, published values for other Neolithic Mediterranean populations are generally similar to those
for South Central Europe,[67–69] with exceptions.[70] Dietary isotopes indicate that the Mediterranean
diet differs in some aspects, with increased δ^{13}C values compared to the other Neolithic populations,
but similar δ^{15}N values. Our data indicates similar protein intake and low-level stress as other
Neolithic populations, but do not suggest any clear hypothesis for the difference in male stature
between the Mediterranean and other Neolithic groups. Possible differences in Mediterranean body
proportions which are not captured by femur length should be mentioned as a caveat, though
this likely would not be enough to account for the differences in stature compared to the rest of
Europe, and would not affect observed dimorphism within the population. Our hypothesis is that
the Mediterranean experienced similar levels of environmental stress as other Neolithic groups, but
that they did not share the cultural practices which preferentially supported males and increased
female vulnerability.

3.3 Conclusion

By integrating genetic and anthropological data, we are able to begin to understand the contribu-
tions of genetics and environment to human variation, allowing us to better interpret the genetic,

environmental, and cultural landscapes of Neolithic Europe. Our results are consistent with a model
in which sexually dimorphic differences in femur length are culturally and environmentally driven:
relatively low dimorphism in the Mediterranean caused by female buffering to environmental stress
and less cultural male preference, and high dimorphism in Northern Central Europe caused by the
interaction of relatively high environmental stress and strong cultural male preference. Some analy-
ses suggest that differences in average femur length between Central/Southeastern Europe and the
Mediterranean are associated with differing genetic ancestries, but lack of robustness, uncertainty
about the transferrability of polygenic scores, and questions of residual population stratification
prevent us from interpreting this conclusively. In this study we focused on the European Early
Neolithic because of its relative genetic, cultural, and environmental homogeneity, but, with more
data, these methods could be extended to other populations, traits, and timescales to further ex-
plore the effects of human culture on biological variation. Using this approach, we gain a deeper
understanding of the relationship between phenotypic plasticity, culture and genetic architecture,
which constrain the mechanisms by which human biology adapts to environment.

4   Materials and Methods

We collected a combination of genetic, dietary stable isotope, skeletal metric, and paleopathological
(stress) data from 1282 individuals from the Central European Early Neolithic associated with the
archaeological LBK culture, approximately 7700-6900 BP (Figure 2, Supplementary Table 1). As
there is archaeological evidence for broad regional variation within the LBK and our sampled sites
form clear geographic groups, we divided these individuals into two regions based on geographical
location, those to the north of 50°N latitude (North Central) and those to the south (South Central)
(Figure 2A-B; North Central n=203, n femur length=133, n isotopes=100, n aDNA=67, n stress=83;
South Central n=1067, n femur length=187, n isotopes=670, n aDNA=72, n stress=523). Each
individual has at least one of the data types, and while some individuals have multiple data types,
the overlaps are small (Supplementary Figure 1).
To provide wider context, we also compared Central individuals to other Neolithic populations
from southern European (Mediterranean) and southeastern European (Balkan) regions, and re-
stricted to individuals dated to 8000-6000 BP. We chose these regions as the Neolithic transition
occurs at similar times and is associated with populations closely related to Central Europe. The
acceptable date range for inclusion in the study was expanded from that which defines the LBK as
these dates encompass comparable Early Neolithic phases in other parts of Europe while maximizing
the number of eligible individuals. There could be a possibility that the later Balkan and Mediter-
ranean individuals were more adapted to Neolithic life than the Central European groups, as these
samples cover a longer time period, but we found no statistical within-population differences in
our variables between the early and late ranges of our time span (minimum p=0.08). We excluded
areas such as Scandinavia and Britain, where Neolithic technologies were not generally adopted

until a later date. For the final analysis, we included 127 Mediterranean (n femur length=60, n
isotopes=25, n aDNA=42) and 139 Balkan (n femur length=12, n isotopes=78, n aDNA=49) indi-
viduals (Figure 2). Unfortunately, there is a wide range of recording and reporting used for skeletal
stress indicators, and it was not possible to build a statistically powerful dataset in these two popu-
lations for comparison; as a result, we did not analyze paleopathology in these populations. Finally,
we collected genetic data from Mesolithic hunter-gatherer (n=25, 14000-7080BP, south of 48°N)
and Anatolian Neolithic (n=21) individuals for additional comparison.

4.1 Genetic data

We obtained genetic data for a total of 276 individuals.[7,18–20,71–87] Most data were generated by
targeting a set of 1.24 million SNPs (the "1240k" capture reagent).[17,75] For each individual, we
randomly selected a single allele from each of the 1240k sites. Coverage in our dataset is low
(median coverage=0.33; coverage above 0.60 n=71), and typically, it is not possible to directly infer
diploid genotypes, potentially limiting PRS performance. Imputation of missing genotypes has been
shown to help improve polygenic predictions for low coverage ancient samples,[9] and we therefore
imputed diploid genotypes using the two-stage method described in that paper, restricting to SNPs
in the 1240k set.

We calculated polygenic scores as previously described.[9] Briefly, we used standing height sum-
mary statistics generated by *fastGWA* from 456,000 individuals of European ancestry in the UK
Biobank[88] for analyses of combined-sex PRS, and summary statistics from male- and female-only
UK Biobank GWAS generated by the Neale Lab.[89] To test the potential effects of residual pop-
ulation structure in our data, we also computed PRS using additional summary statistics from a
between-sibling GWAS (n=99,997).[23] We intersected the sites from each of these datasets with those
on the 1240k array and then further restricted to HapMap3 SNPs (SNPs n=405,000). We computed
polygenic scores using both a clumping/thresholding approach (r^2=0.3, p-value cutoff=10^{-6}, 100kb
windows in *plink2*[90]), and an infinitesimal *LDpred2* model using their pre-computed LD reference
panel.[91] Finally, we computed polygenic scores using the --score command in *plink2*. In order to
maximize the possibility of detecting sex-specific effects, we generated sex-specific PRS using three
different approaches: 1) calculating PRS for all individuals using the female summary statistics; 2)
calculating PRS for all individuals using the male summary statistics; and 3) calculating PRS for
males and females separately using their respective summary statistics. While approach 3 seems
at first to be the best for detecting these effects, observed patterns potentially become difficult to
interpret due to differences in scaling between male and female PRS calculated as separate datasets.

We computed principal components for both unimputed and imputed data using *smartpca*,[92] pro-
jecting ancient individuals onto principal component axes defined by 777 present-day West Eurasian
individuals.[93] We also estimated K=2 unsupervised ADMIXTURE[94] components for unimputed
ancient individuals after first LD pruning using the command --pairwise-indep 200 25 0.4 in *plink2*.

4.2 Osteology and stable isotope data

We aggregated skeletal metric data from both published[57,95–99] and unpublished (n=28) sources.
Maximum femur lengths were recorded when available, otherwise we estimated femur length from
published stature estimates.[9] Estimated femur lengths correlate highly with stature estimates, but
decrease the error that results from combining different estimations methods. The method from
Ref. 100 provides separate equations for estimating the statures of northern vs. southern Europeans
when using the tibia, due to differences in body proportions between the regions. There are two
Mediterranean samples for which we estimated the length of the femur based on statures which used
the southern tibia equation. Ref. 100 does not provide regional equations for femur estimation, so
for these two individuals, we estimated femur length using the reverse of this region-agnostic femur
equation.

For the individuals in this study who do not have genetic data, morphology was used to estimate
sex. The majority of individuals have been taken from previous publications, and we used the sexes
which had been estimated by those authors. For the individuals in our study which have not
been previously published, sex was determined by co-authors using a 5-point scale on the cranium
and pelvis as described by Ref. 101. For all individuals, sexes determined as probable male or
probable female were coded in our study as either male or female as appropriate. Subadults and
those with indeterminate morphologies were coded as NA, resulting in these individuals being
dropped from the sex-specific analyses. The majority of sexes for individuals with metric data were
determined by, or supervised by, co-authors and the remainder (n=13) either have genetic sexes
or come from Ref. 57 which we consider a reliable source. Despite generally high accuracy for
morphological sex determination, some level of uncertainty always remains, mainly due to variation
in sexual dimorphism and preservation of the remains.[102] Sex estimations for our sample have
all been performed in the last 20 years, and the majority within the last 5 years, meaning the
researchers who performed them should be aware of avoiding the biases which can affect sex-ratios
in the estimations of older data. Our dataset is large enough that small errors in classification
of sex should not make substantial differences to results or interpretation, but the potential for
inaccurate morphological sex estimations must always be considered in any osteological analysis. A
large portion of our paleopathology data comes from tables S3 and S6 of Ref. 103, in which there
are many instances of the same individual listed in both tables, but with discordant sex estimations.
As we could not determine the reason for these discrepancies, we used the sex which was reported
in the original publications cited as sources for their data. The few individuals (n=3) for whom this
could not be resolved were treated as indeterminate and coded as NA. Ages were determined based
on the average of the age range reported for each individual in their original publications.

For the paleopathological data in Central Europe, we took data from published sources,[95,96,103–106]
as presence/absence of linear enamel hypoplasia (LEH), porotic hyperostosis, and cribra orbitalia.
These three pathologies are often used by anthropologists as indicators of general, non-specific stress
experienced by individuals or populations. While the exact eitiologies of these pathologies are gen-

erally not known, they have been shown to change through time within and between populations,
and often correlate with environmental, social, or cultural shifts. Linear enamel hypoplasias are
horizontal defects in tooth enamel that form during episodes of childhood stress severe enough to
interrupt growth for some period of time, usually associated with dietary deficiency or infectious
disease.[46] Individuals can exhibit one or multiple LEH on single or multiple teeth and in order
to minimize errors from differences in reporting LEH in the literature, we have simply recorded
whether an individual had any LEH (present) or none (absent). Porotic hyperostosis and cribra
orbitalia are both porous lesions that are distinguished by their appearance on either the cranial
vault or roof of the eye orbit respectively. The eitiologies of these are mostly unknown and though
they are traditionally associated with amaemias, there are also a number of other conditions that
can produce the same type of lesions. Medically, there is little evidence of these pathological changes
despite their prevalence in the bioarchaeological record.[107] Similar to LEH, we have recorded these
as either present or absent for each individual in order to standardize between reporting conventions
across publications.

While sensitive to confounding factors such as climate, vegetation, and individual metabolism,[108]
δ^{13}C and δ^{15}N stable isotope data can be used to reconstruct aspects of diet.[109] Here, carbon values
are indicative of dietary plant resources and of the terrestrial vs marine vs limnic provenance of
food, while nitrogen values are mainly associated with dietary protein intake and generally indi-
cate proportions of plant- vs animal-based diets.[108,109] We collected dietary stable isotopes δ^{13}C
and δ^{15}N from published[16,35–37,43,95,97,103,105,106,108,110–113] and unpublished (n=38) reports. We
excluded atomic mass spectrometer (AMS) values derived from radiocarbon dating, as they may
not be comparable to isotope-ratio mass spectrometer (IRMS) measurements, as well as values from
children below the age of three, due to increased nitrogen values from breastfeeding. Stable isotope
values from older children were included in population-wide diet analyses as the isotope ranges fall
within those of adults; however, we only included adults with estimated sexes in the sex-based diet
analyses. If information on the sampled material was available, we chose values measured from
rib collagen, as these samples are most plentiful, though they only reflect the last few years of the
individual's life.

All previously unpublished osteological data was collected and analyzed by co-authors with
permission from the necessary regulating organizations and in accordance with German laws and
policies.

4.3 Statistical models

We tested the effects of PRS, femur length, and isotope data on stature using linear regression
models including sex and geographic region as covariates in combination with other variables as
appropriate (e.g., femur $\sim$ sex + region + PRS; δ^{15}N $\sim$ sex + region + femur). We included
interaction terms to test the relationships between geographic regions and sex (e.g., femur $\sim$ region
* sex) and used t-tests to test within-sex differences between regions. We used logistic regression

with the same covariates to test for factors affecting presence/absence of paleopathologies. We
carried out all statistical tests using the base functions in R version 4.0.[114]

4.4 Data Availability

501 All non-genetic data and polygenic scores used in this analysis are provided in Supplementary Table
502 1. Original ancient DNA data files can be downloaded from the resources provided in their cited
503 publications. Previously published osteological data can be found in their cited sources (Supple-
504 mentary Table 1).

4.5 Code Availability

506 R code used in this analysis is available at `https://github.com/mathilab/Neolithic_height.git`.

4.6 Acknowledgements

This work was supported by a grant from the National Science Foundation [BCS2123627] to IM.
Metric skeletal data collection from Stuttgart-Mühlhausen was funded by the Deutsche Forschungs-
gemeinschaft (DFG; German Research Foundation) [RO 4148/1-1]. The investigations of the skeletal
collections from Derenburg, Karsdorf, and Halberstadt were supported by the DFG [Al 287/7-1 and
7-3; Me 3245/1-1 and 1-3]. The content is the responsibility of the authors and does not necessarily
represent the official view of the NSF or other funders.

4.7 Author Contributions

515 S.L.C. designed the study, collected data, performed analysis, and wrote the manuscript; I.M.
516 designed the study and wrote the manuscript; N.N. and K.W.A., contributed data and archaeological
517 background; E.R. contributed data and performed analysis; M.F., J.W., H.M., and W.H. contributed
518 data. All authors edited and approved the final version.

4.8 Competing Interests

520 The authors declare no competing interests.

References

522 1. Harpak, A. & Przeworski, M. The evolution of group differences in changing environments.
523   *PLoS Biol* **19**, e3001072. doi: 10.1371/journal.pbio.3001072 (2021).

524 2. Martin, A. R. *et al.* Clinical use of current polygenic risk scores may exacerbate health dis-
525   parities. *Nat Genet* **51**, 584–591. doi: 10.1038/s41588–019–0379–x (2019).

3. Roseman, C. C. Detecting interregionally diversifying natural selection on modern human cranial form by using matched molecular and morphometric data. *Proceedings of the National Academy of Sciences* **101**, 12824–12829. doi: 10.1073/pnas.0402637101 (2004).

4. Whitlock, M. C. Evolutionary inference from QST. *Mol Ecol* **17**, 1885–96. doi: 10.1111/j.1365–294X.2008.03712.x (2008).

5. Savell, K. R., Auerbach, B. M. & Roseman, C. C. Constraint, natural selection, and the evolution of human body form. *Proc Natl Acad Sci U S A* **113**, 9492–7. doi: 10.1073/pnas.1603632113 (2016).

6. Berg, J. J. & Coop, G. A population genetic signal of polygenic adaptation. *PLoS Genet* **10**, e1004412. doi: 10.1371/journal.pgen.1004412 (2014).

7. Mathieson, I. *et al.* Genome-wide patterns of selection in 230 ancient Eurasians. *Nature* **528**, 499–503. doi: 10.1038/nature16152 (2015).

8. Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and body mass index in approximately 700000 individuals of European ancestry. *Hum Mol Genet* **27**, 3641–3649. doi: 10.1093/hmg/ddy271 (2018).

9. Cox, S. L. *et al.* Predicting skeletal stature using ancient DNA. *American Journal of Biological Anthropology* **177**, 162–174 (2022).

10. Marciniak, S. *et al.* An integrative skeletal and paleogenomic analysis of stature variation suggests relatively reduced health for early European farmers. *Proceedings of the National Academy of Sciences* **119**, e2106743119. doi: 10.1073/pnas.2106743119 (2022).

11. Cox, S. L., Ruff, C. B., Maier, R. M. & Mathieson, I. Genetic contributions to variation in human stature in prehistoric Europe. *Proceedings of the National Academy of Sciences* **116**, 21484–21492. doi: 10.1073/pnas.1910606116 (2019).

12. Bickle, P. & Whittle, A. (eds.) *The First Farmers of Central Europe* (Oxbow, Oxford, UK, 2013).

13. Bakels, C. C. *The Western European loess belt: agrarian history, 5300 BC-AD 1000* (Springer Science & Business Media, 2009).

14. Bentley, R. A. *et al.* Community differentiation and kinship among Europe's first farmers. *Proceedings of the National Academy of Sciences* **109**, 9326–9330. doi: 10.1073/pnas.1113710109 (2012).

15. Nicklisch, N., Oelze, V. M., Schierz, O., Meller, H. & Alt, K. W. A healthier smile in the past? dental caries and diet in early neolithic farming communities from central germany. *Nutrients* **14**, 183. doi: 10.3390/nu14091831 (2022).

16. Knipper, C. Kohlenstoff- und Stickstoff-Isotopenanalysen an bandkeramischen Bestattungen
vom Viesenhäuser Hof bei Stuttgart-Mühlhausen: Implikationen zur Ernährungsrekonstruk-
tion, Geschlechtsspezifik und Siedlungsdynamik. In Meyer, C., Held, P., Knipper, C. & Nick-
lisch, N. (eds.) *Der Zahn der Zeit – Mensch und Kultur im Spiegel interdisziplinärer Forschung.*
Festschrift für Kurt W. Alt (Veröffentlichungen des LDA und Archäologie Sachsen-Anhalt,
Halle(Saale), 2020).

17. Haak, W. *et al.* Massive migration from the steppe was a source for Indo-European languages
in Europe. *Nature* **522**, 207–11. doi: 10.1038/nature14317 (2015).

18. Lipson, M. *et al.* Parallel palaeogenomic transects reveal complex genetic history of early
European farmers. *Nature* **551**, 368–372. doi: 10.1038/nature24476 (2017).

19. Mathieson, I. *et al.* The genomic history of southeastern Europe. *Nature* **555**, 197–203. doi:
10.1038/nature25778 (2018).

20. Antonio, M. L. *et al.* Ancient Rome: A genetic crossroads of Europe and the Mediterranean.
Science **366**, 708–714. doi: 10.1126/science.aay6826 (2019).

21. Martiniano, R. *et al.* The population genomics of archaeological transition in west iberia:
Investigation of ancient substructure using imputation and haplotype-based methods. *PLoS*
Genet **13**, e1006852. doi: 10.1371/journal.pgen.1006852 (2017).

22. Gronenborn, D., Horejs, B., Börner & Ober.   Map: Expansion of farming in west-
ern Eurasia, 9600 - 4000 cal BC (update vers. 2021.2) (RGZM/ÖAI).   online (2021).
https://www.academia.edu/9424525/Map_Expansion_of_farming_in_western_Eurasia_
9600_4000_cal_BC_update_vers_2021_2_.

23. Howe, L. J. *et al.* Within-sibship genome-wide association analyses decrease bias in estimates
of direct genetic effects. *Nature genetics* **54**, 581–592 (2022).

24. Sohail, M. *et al.* Polygenic adaptation on height is overestimated due to uncorrected stratifi-
cation in genome-wide association studies. *Elife* **8**, e39702. doi: 10.7554/eLife.39702 (2019).

25. Berg, J. J. *et al.* Reduced signal for polygenic adaptation of height in UK Biobank. *Elife* **8**,
e39725. doi: 10.7554/eLife.39725 (2019).

26. Vercellotti, G. *et al.* Exploring the multidimensionality of stature variation in the past through
comparisons of archaeological and living populations. *American Journal of Physical Anthro-*
pology **155**, 229–242 (2014).

27. Roseman, C. C. & Auerbach, B. M. Ecogeography, genetics, and the evolution of human body
form. *J Hum Evol* **78**, 80–90. doi: 10.1016/j.jhevol.2014.07.006 (2015).

28. Stock, J. T. *et al.* Long-term trends in human body size track regional variation in subsistence
transitions and growth acceleration linked to dairying. *Proceedings of the National Academy*
of Sciences **120**, e2209482119 (2023).

29. Randall, J. C. *et al.* Sex-stratified genome-wide association studies including 270,000 individu-
als show sexual dimorphism in genetic loci for anthropometric traits. *PLOS Genetics* **9**, 1–19.
doi: 10.1371/journal.pgen.1003500 (2013).

30. Bernabeu, E., Canela-Xandri, O., Rawlik, K. & et al. Sex differences in genetic architecture in
the UK Biobank. *Nature Genetics* **53**, 1283–1289. doi: 10.1038/s41588–021–00912–0 (2021).

31. Zhu, C., Ming, M. J., Cole, J. M., Kirkpatrick, M. & Harpak, A. Amplification is
the primary mode of gene-by-sex interaction in complex human traits. *bioRxiv* (2022).
doi:10.1101/2022.05.06.490973.

32. Bogaard, A. *Neolithic farming in central Europe: an archaeobotanical study of crop husbandry*
practices (Routledge, 2004).

33. Schier, W. Modes and Models of Neolithization in Europe: Comments to an Ongoing De-
bate. In Biehl, P. & Rosenstock, E. (eds.) *6000 BC: Transformation and Change in the*
Near East and Europe, 372–392 (Cambridge University Press, Cambridge, UK, 2022). doi:
10.1017/9781107337640.026.

34. Betti, L. *et al.* Climate shaped how Neolithic farmers and European hunter-gatherers interacted
after a major slowdown from 6,100 BCE to 4,500 BCE. *Nature Human Behaviour* **4**, 1004–1010
(2020).

35. Dürrwächter, C., Craig, O. E., Collins, M. J., Burger, J. & Alt, K. W. Beyond the grave:
variability in Neolithic diets in Southern Germany. *Journal of Archaeological Science* **33**,
39–48 (2006).

36. Oelze, V. M. *et al.* Early Neolithic diet and animal husbandry: stable isotope evidence from
three Linearbandkeramik (LBK) sites in Central Germany. *Journal of Archaeological Science*
38, 270–279 (2011).

37. Bentley, R. A. *et al.* Baden-Württemberg. In Bickle, P. & Whittle, A. (eds.) *The First Farmers*
in Central Europe: Diversity in LBK Lifeways (Oxbow Books, Oxford, UK, 2013).

38. Denaire, A. *et al.* The cultural project: Formal chronological modelling of the early and middle
neolithic sequence in lower alsace. *Journal of Archaeological Method and Theory* **24**, 1072–1149
(2017).

39. Gillis, R. E. *et al.* Stable isotopic insights into crop cultivation, animal husbandry, and land
use at the Linearbandkeramik site of Vráble-Vel'ké Lehemby (Slovakia). *Archaeological and*
Anthropological Sciences **12**, 1–15 (2020).

40. Nicklisch, N. *et al.* Holes in teeth–dental caries in Neolithic and early bronze age populations
in central Germany. *Annals of Anatomy-Anatomischer Anzeiger* **203**, 90–99 (2016).

41. Gillis, R. E. *et al.* The evolution of dual meat and milk cattle husbandry in Linearbandkeramik
societies. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20170905 (2017).

42. Salque, M. *et al.* Earliest evidence for cheese making in the sixth millennium BC in northern
Europe. *Nature* **493**, 522–525 (2013).

43. Münster, A. *et al.* 4000 years of human dietary evolution in central Germany, from the first
farmers to the first elites. *PloS one* **13**, e0194862 (2018).

44. Ghosh, S. Protein quality in the first thousand days of life. *Food and Nutrition Bulletin* **37**,
S14–S21 (2016).

45. Gray, J. P. & Wolfe, L. D. Height and sexual dimorphism of stature among human societies.
American Journal of Physical Anthropology **53**, 441–456 (1980).

46. Guatelli-Steinberg, D. & Lukacs, J. R. Interpreting sex differences in enamel hypoplasia in
human and non-human primates: Developmental, environmental, and cultural considerations.
American journal of physical anthropology **110**, 73–126 (1999).

47. Lubell, D., Jackes, M. & Meiklejohn, C. Archaeology and human biology of the Mesolithic-
Neolithic transition in southern Portugal: a preliminary report. In *The Mesolithic in Europe.*
International Symposium. 3, 632–640 (1990).

48. McFadden, C. & Oxenham, M. F. A paleoepidemiological approach to the osteological para-
dox: Investigating stress, frailty and resilience through cribra orbitalia. *American Journal of*
Physical Anthropology **173**, 205–217 (2020).

49. German, A. & Hochberg, Z. Sexual dimorphism of size ontogeny and life history. *Frontiers in*
Pediatrics (2020).

50. Stini, W. Growth rates and sexual dimorphism in evolutionary perspective. In Stini, W. (ed.)
The Analysis of Prehistoric Diets (Academic Press, Orlando, FL, 1985).

51. Brauer, G. Size sexual dimorphism and secular trend: indicators of subclinical malnutrition.
In Hall, R. (ed.) *Sexual Dimorphism in Homo sapiens: A Question of Size* (Praeger, New
York, NY, 1982).

52. Vlassoff, C. Gender differences in determinants and consequences of health and illness. *Journal*
of Health, Population, and Nutrition **25**, 47–61 (2007).

53. Dehingia, N. & Raj, A. Sex differences in covid-19 case fatality: do we know enough? *The*
Lancet Global Health **9**, e14–e15 (2021).

54. Takahashi, T. *et al.* Sex differences in immune responses that underlie covid-19 disease outcomes. *Nature* **588**, 315–320 (2020).

55. Frayer, D. W. & Wolpoff, M. H. Sexual dimorphism. *Annual Review of Anthropology* **14**, 429–473 (1985).

56. Gaulin, S. J. & Boster, J. S. Human marriage systems and sexual dimorphism in stature. *American journal of physical anthropology* **89**, 467–475 (1992).

57. Ruff, C. B. (ed.) *Skeletal variation and adaptation in Europeans: Upper Paleolithic to the twentieth century* (John Wiley & Sons, Hoboken, NJ, 2018). (data available at https://www.hopkinsmedicine.org/fae/CBR.html).

58. Kanwar, S. C., Sabharwal, A., Bhadra, K. & Narang, A. Nationwide reference data for height, weight and body mass index of Indian schoolchildren. *Natlonal Medical Journal of India* **24**, 269–77 (2011).

59. Abdulrazzaq, Y., Moussa, M. & Nagelkerke, N. National growth charts for the United Arab Emirates. *Journal of Epidemiology* **18**, 295–303 (2008).

60. Moradi, A. & Guntupalli, A. M. What does gender dimorphism in stature tell us about discrimination in rural India, 1930-1975. *Gender Bias: Health, Nutrition and Work* (2009).

61. Kanazawa, S. & Novak, D. L. Human sexual dimorphism in size may be triggered by environmental cues. *Journal of Biosocial Science* **37**, 657–665 (2005).

62. Baten, J. & Murray, J. E. Heights of men and women in 19th-century Bavaria: economic, nutritional, and disease influences. *Explorations in Economic History* **37**, 351–369 (2000).

63. Castellucci, H. *et al.* Gender inequality and sexual height dimorphism in Chile. *Journal of Biosocial Science* **53**, 38–54 (2021).

64. Bentley, R. A. *et al.* Prehistoric migration in Europe: strontium isotope analysis of early Neolithic skeletons. *Current Anthropology* **43**, 799–804 (2002).

65. Bickle, P. Thinking gender differently: New approaches to identity difference in the central european neolithic. *Cambridge Archaeological Journal* **30**, 201–218 (2020).

66. Garvin, H. M. *The effects of living conditions on human cranial and postcranial sexual dimorphism.* Ph.D. thesis, The Johns Hopkins University (2012).

67. Papathanasiou, A. Health, diet and social implications in Neolithic Greece from the study of human osteological material. *Human bioarchaeology of the transition to agriculture* 87–106 (2011).

68. Papathanasiou, A. Health status of the Neolithic population of Alepotrypa Cave, Greece.
American Journal of Physical Anthropology: The Official Publication of the American Associ-
ation of Physical Anthropologists **126**, 377–390 (2005).

69. Silva, A. M. & Cunha, E. Paleopathological study of the community exhumed from the Hipogeu
of Monte Canelas I (Alcalar, Portugal). In *Actas del V Congreso Nacional de Paleopatologia*,
353–356 (2001).

70. Cucina, A. Brief communication: diachronic investigation of linear enamel hypoplasia in
prehistoric skeletal samples from Trentino, Italy. *American Journal of Physical Anthropology*
119, 283–287 (2002).

71. Brunel, S. *et al.* Ancient genomes from present-day France unveil 7,000 years of its demographic
history. *Proceedings of the National Academy of Sciences* **117**, 12791–12798 (2020).

72. Childebayeva, A. *et al.* Population genetics and signatures of selection in Early Neolithic
European farmers. *Molecular Biology and Evolution* **39**, msac108 (2022).

73. Fernandes, D. M. *et al.* The spread of steppe and Iranian-related ancestry in the islands of the
western Mediterranean. *Nature Ecology & Evolution* **4**, 334–345 (2020).

74. Fregel, R. *et al.* Ancient genomes from North Africa evidence prehistoric migrations to the
Maghreb from both the Levant and Europe. *Proceedings of the National Academy of Sciences*
115, 6774–6779 (2018).

75. Fu, Q. *et al.* The genetic history of Ice Age Europe. *Nature* **534**, 200–205. doi: 10.1038/na-
ture17993 (2016).

76. Gamba, C. *et al.* Genome flux and stasis in a five millennium transect of European prehistory.
Nature communications **5**, 1–9 (2014).

77. González-Fortes, G. *et al.* Paleogenomic evidence for multi-generational mixing between Ne-
olithic farmers and Mesolithic hunter-gatherers in the Lower Danube Basin. *Current Biology*
27, 1801–1810 (2017).

78. Hofmanová, Z. *et al.* Early farmers from across Europe directly descended from Neolithic
Aegeans. *Proceedings of the National Academy of Sciences* **113**, 6886–6891 (2016).

79. Jones, E. R. *et al.* Upper Palaeolithic genomes reveal deep roots of modern Eurasians. *Nature*
Communications **6**, 1–8 (2015).

80. Lazaridis, I. *et al.* Genetic origins of the Minoans and Mycenaeans. *Nature* **548**, 214–218
(2017).

81. Marcus, J. H. *et al.* Genetic history from the Middle Neolithic to present on the Mediterranean
island of Sardinia. *Nature Communications* **11**, 1–14 (2020).

82. Nikitin, A., Stadler, P. & Kotova, e. a., N. Interactions between earliest Linearbandkeramik
farmers and central European hunter gatherers at the dawn of European Neolithization. *Sci-*
entific Reports **9**, 19544. doi: 10.1038/s41598–019–56029–2 (2019).

83. Olalde, I. *et al.* A common genetic origin for early farmers from Mediterranean Cardial and
Central European LBK cultures. *Molecular Biology and Evolution* **32**, 3132–3142 (2015).

84. Olalde, I. *et al.* The genomic history of the Iberian Peninsula over the past 8000 years. *Science*
363, 1230–1234 (2019).

85. Rivollat, M. *et al.* Ancient genome-wide DNA from France highlights the complexity of in-
teractions between Mesolithic hunter-gatherers and Neolithic farmers. *Science Advances* **6**,
eaaz5344 (2020).

86. Valdiosera, C. *et al.* Four millennia of Iberian biomolecular prehistory illustrate the impact
of prehistoric migrations at the far end of Eurasia. *Proceedings of the National Academy of*
Sciences **115**, 3428–3433 (2018).

87. Villalba-Mouco, V. *et al.* Survival of Late Pleistocene hunter-gatherer ancestry in the Iberian
Peninsula. *Current Biology* **29**, 1169–1177 (2019).

88. Jiang, L. *et al.* A resource-efficient tool for mixed model association analysis of large-scale
data. *Nature Genetics* **51**, 1749–1755. doi: 10.1038/s41588–019–0530–8 (2019).

89. Neale Lab. UK Biobank GWAS Results. http://www.nealelab.is/uk-biobank (2018). Accessed:
2021-10-20.

90. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer
datasets. *Gigascience* **4**, 7. doi: 10.1186/s13742–015–0047–8 (2015).

91. Privé, F., Arbel, J. & Vilhjálmsson, B. J. LDpred2: better, faster, stronger. *Bioinformatics*
36, 5424–5431 (2020).

92. Patterson, N., Price, A. L. & Reich, D. Population structure and eigenanalysis. *PLoS Genetics*
2, e190. doi: 10.1371/journal.pgen.0020190 (2006).

93. Lazaridis, I. *et al.* Ancient human genomes suggest three ancestral populations for present-day
europeans. *Nature* **513**, 409–13. doi: 10.1038/nature13673 (2014).

94. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in
unrelated individuals. *Genome Research* **19**, 1655–64. doi: 10.1101/gr.094052.109 (2009).

95. Hujić, A. *Das Kind in uns unter der Lupe der Isotopie, Allometrie und Pathologie. Zusammenhang zwischen $\delta^{15}N$ und $\delta^{13}C$ als Eiweißproxy und dem Knochenwachstum bei linienbandkeramischen Individuen aus Stuttgart-Mühlhausen, "Viesenhäuser Hof" und Schwetzingen unter Berücksichtigung verschiedener Indikatoren für Nährstoffversorgung.* Ph.D. thesis, Freie Universität (2016). doi: 10.17169/refubium-5564.

96. Nicklisch, N. *Spurensuche am Skelett: Paläodemografische und epidemiologische Untersuchungen an neolithischen und frühbronzezeitlichen Bestattungen aus dem Mittelelbe-Saale-Gebiet im Kontext populationsdynamischer Prozesse* (Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum, 2017).

97. Meyer, C. *et al.* Early Neolithic executions indicated by clustered cranial trauma in the mass grave of Halberstadt. *Nature Communications* **9**, 1–11 (2018).

98. Rosenstock, E., Ebert, J., Martin, R. & Groß, M. LiVES COstA Digest: checked and updated entries from the LiVES collection of Osteological Anthropometry. doi: 10.17171/2-12-2-1 (2019).

99. Rosenstock, E. *et al.* Human stature in the near east and europe ca. 10,000–1000 bc: its spatiotemporal development in a bayesian errors-in-variables model. *Archaeological and Anthropological Sciences* **11**, 5657–5690. doi: 10.1007/s12520–019–00850–3 (2019).

100. Ruff, C. B. *et al.* Stature and body mass estimation from skeletal remains in the european holocene. *American journal of physical anthropology* **148**, 601–617 (2012).

101. Acsádi, G., Nemeskéri, J. & Balás, K. *History of human life span and mortality* (Akademiai kiado Budapest, 1970).

102. Krishan, K. *et al.* A review of sex estimation techniques during examination of skeletal remains in forensic anthropology casework. *Forensic Science International* **261**, 165–e1 (2016).

103. Ash, A. *et al.* Regional differences in health, diet and weaning patterns amongst the first Neolithic farmers of central Europe. *Scientific Reports* **6**, 1–10 (2016).

104. Lillie, M. Vedrovice: demography and palaeopathology in an early farming population. *Anthropologie* **46**, 135–152 (2008).

105. Whittle, A. *et al.* Hungary. In Bickle, P. & Whittle, A. (eds.) *The First Farmers in Central Europe: Diversity in LBK Lifeways* (Oxbow Books, Oxford, UK, 2013).

106. Whittle, A. *et al.* Moravia and western Slovakia. In Bickle, P. & Whittle, A. (eds.) *The First Farmers in Central Europe: Diversity in LBK Lifeways* (Oxbow Books, Oxford, UK, 2013).

107. Brickley, M. B. Cribra orbitalia and porotic hyperostosis: A biological approach to diagnosis. *American Journal of Physical Anthropology* **167**, 896–902 (2018).

108. Scheibner, A. *Prähistorische Ernährung in Vorderasien und Europa. Eine kulturgeschichtliche*
Synthese auf der Basis ausgewählter Quellen. Schriften zum Lebensstandard in der
Vorgeschichte 1 (Berliner Archäologische Forschungen, Leidorf: Rahden, 2016).

109. O'Brien, D. M. Stable Isotope Ratios as Biomarkers of Diet for Health Research. *Annu Rev*
Nutr **35**, 565–594 (2015).

110. Richards, M. P., Montgomery, J., Nehlich, O. & Grimes, V. Isotopic analysis of humans and
animals from Vedrovice. *Anthropologie* **46**, 185–194 (2008).

111. Bickle, P. *et al.* Austria. In Bickle, P. & Whittle, A. (eds.) *The First Farmers in Central*
Europe: Diversity in LBK Lifeways (Oxbow Books, Oxford, UK, 2013).

112. Bickle, P. *et al.* Alsace. In Bickle, P. & Whittle, A. (eds.) *The First Farmers in Central*
Europe: Diversity in LBK Lifeways (Oxbow Books, Oxford, UK, 2013).

113. Hofmann, D. *et al.* Southern Bavaria. In Bickle, P. & Whittle, A. (eds.) *The First Farmers in*
Central Europe: Diversity in LBK Lifeways (Oxbow Books, Oxford, UK, 2013).

114. R Core Team. *R: A Language and Environment for Statistical Computing.* R Foundation for
Statistical Computing, Vienna, Austria (2021). https://www.R-project.org/.