Internalization-Responsive Nano-PLGA For Image-Guided Photodynamic Therapy Against HER2-Positive Breast Cancer

Anna S. Sogomonyan[a], Sergey M. Deyev[a], Victoria O. Shipunova[a,b,*]

[a] Shemyakin–Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, Moscow, 117997, Russia

[b] Sirius University of Science and Technology, 1 Olympic Ave., 354340 Sochi, Russia

[*]e-mail: viktoriya.shipunova@phystech.edu

Abstract Oncotheranostic nanoagents are powerful artificial compounds that significantly outperform traditional antitumor therapies. Among the wide variety of nanoagents, polymer nanoparticles have proven to be one of the most successful platforms for translation into the clinic. Here we show the development of an oncotheranostic nanoagent based on poly(lactic-co-glycolic acid) (PLGA) with red/green light dual-activated diagnostic/therapeutic properties for targeted photodynamic therapy of HER2-positive aggressive breast cancer. PLGA particles were loaded with red light-activated Nile Blue dye with pronounced solvatochromic properties for diagnostic applications and with green light-activated photodynamic sensitizer Rose Bengal for photodynamic therapy. The nanoparticles were decorated with anti-HER2 antibodies during the synthesis without chemical conjugation which is very important for further large-scale biotechnological production. *In vitro* and *in vivo studies* proved the red-light mediated HER2-specific imaging and green-light induced cytotoxicity of obtained anti-HER2 PLGA. Interestingly, these particles fluoresce only when internalized into cells, which minimizes background signals until the complete elimination from the organism, thus facilitating real-time diagnostics. We have shown that such particles allow efficient and selective localization of the HER2-positive primary tumor node, as well as metastases, and achieve complete remission in BALB/c Nu/Nu mice with the HER2-positive xenografts using the single session of photodynamic therapy.

Keywords nanoparticles, PLGA, targeted delivery, photosensitizers, theranostics, PDT, HER2, dual-modality

1. Introduction

The creation of theranostic agents for the diagnosis and treatment of cancer is at the forefront of modern biomedicine due to many serious problems in this field [1,2]. Traditional methods of both diagnostics and therapy lead to a whole range of undesirable side effects that crucially affect a significantly weakened organism. Namely, chemotherapy with doxorubicin leads to high cardio- and hepatotoxicity, various surgical methods break the integrity of blood vessels, increasing the rate of metastasis, and various methods of radiation therapy lead to irreversible changes in the bone marrow [3,4]. The development of safe and effective methods of systemic influence on the organism with oncological processes is one of the greatest challenges facing modern natural sciences [5,6].

These challenges led to the development of a relatively new direction in biomedicine, theranostics. Theranostics involves the combination of diagnostic and therapeutic tools on a single platform that allows both

visualizing the target tissue and, if necessary, performing the treatment and therapy monitoring, thus implementing the concept of a "magic bullet", formulated by the founder of chemotherapy, Paul Ehrlich [7,8]. The most suitable platform for the creation of theranostic agents is biocompatible and biodegradable nanoparticles (polymer and protein nanoparticles, liposomes), which, due to their structure, can serve as carriers of both diagnostic and therapeutic modalities [9]. At the same time, decorating the surface of nanoparticles with targeting molecules makes it possible to implement the concept of targeted delivery of therapeutic and/or diagnostic compounds directly to the tissue of interest [10]. For example, targeted therapy for HER2-overexpressing cancers remains one of the most serious problems of modern molecular oncology [11]. In particular, nanoparticles based on a copolymer of lactic and glycolic acids, poly(lactic-co-glycolic acid) or PLGA, are widely used in the clinic; more than 20 PLGA-based medications have already been approved by the FDA for human administration [12]. Such particles are used as carriers of therapeutic compounds for the treatment of prostate and breast cancer (Suprefact Depot, Lupron Depot, Trelstar, Eligard, Decapeptyl), neurologic disorders (Risperdal Consta), opioid dependence treatment (Vivitrol, Sublocade) and other diseases.

However, to date, there are no *targeted* PLGA nanoparticles for clinical use, which is caused, in our opinion, by the insufficient effectiveness of existing formulations, the complexity of chemical conjugation with targeting molecules on an industrial scale, and low batch-to-batch reproducibility. These problems stimulate the rise of fundamental research in the field of designing targeted theranostic PLGA nanoparticles with improved efficiency.

Thus, different PLGA-based nanoformulations have been previously developed for imaging and therapy of HER2-positive cancer. HER2-targeted PLGA were developed for the targeted delivery of chemotherapic drugs, namely doxorubicin [13] and tamoxifen [14]. Moreover, several PLGA-based nanoplatforms combining diagnostic and therapeutic functions were designed for HER2-targeted delivery *in vivo*. Namely, perfluoropentane-loaded PLGA conjugated to trastuzumab was developed for ultrasound and magnetic resonance imaging and photothermal therapy of HER2-positive xenografts [15], PLGA-based nanoparticles were developed for photoacoustic&MRI imaging and gold-nanoparticle induced photothermal therapy [16].

However, today photodynamic therapy (PDT) is considered to be more promising and safer than photothermal therapy (PTT), which is confirmed by the presence of a sufficiently large number of PDT methods in clinical practice and only a few cases of PTT [17,18]. This fact leads to the understanding that the design of HER2-targeted theranostic agents with the property of light-inducible PDT is an urgent need for personalized medicine. But, to date, there are no such PLGA-based nanoparticles that have shown their effectiveness *in vivo*. There are several *in vitro* studies directed toward the development of HER2-targeted nanoparticles for PDT and testing their efficacy on cell culture [19,20]; and non-targeted structures loaded with a photosensitizer have also been generated [21].

Here we show the creation of anti-HER2 PLGA nanoparticles possessing fluorescent diagnostic properties and light-inducible PDT of HER2-overexpressing tumors. Such theranostic nanoparticles make it possible to

localize the tumor and metastases when exposed to red laser radiation and to eliminate HER2-overexpressing tumors *in vivo* when exposed to green laser radiation. Interestingly, these particles fluoresce only when internalized into cells, which minimizes background signals when they are not yet completely cleared from the organism, thus facilitating real-time diagnostics.

These properties of the particles are achieved through the loading of the fluorescent dye Nile Blue (NB) with pronounced solvatochromic properties and the photodynamic sensitizer, Rose Bengal (RB), which produces reactive oxygen species (ROS) upon exposure to green light. Nanoparticles are decorated with anti-HER2 antibody trastuzumab, which is incorporated into the nanoparticle structure during the synthesis process without the need for chemical modification with cross-linking reagents such as EDC/NHS, which is of great importance for potential biotechnological production. We have shown that such particles allow efficient and selective localization of the HER2+ primary tumor node, as well as metastases, and achieve complete remission in the xenograft HER2-positive tumor model in BALB/c Nu/Nu mice.

2. Results

2.1. Design of experiment

HER2-specific cancer diagnostics and therapy were realized according to the following experimental scheme (**Fig. 1**). Co-polymer of lactic and glycolic acids, PLGA, was used as the matrix for the synthesis of theranostic nanoagent. PLGA nanoparticles were synthesized by the "water-oil-water" microemulsion method with the incorporation of water-soluble Rose Bengal into the first "water" phase and water-insoluble Nile Blue dye into the "oil" phase. The nanoparticles were equipped with the anti-HER2 antibody trastuzumab for HER2-specific cancer cell targeting *in vivo*. When administered systemically into the organism, these anti-HER2 PLGA nanoparticles selectively bind to HER2-overexpressing cancer cells in a tumor through the interaction of trastuzumab within the composition of nanoparticles with the receptor HER2 on the cancer cell surface.

The diagnostic modality of anti-HER2 PLGA particles was realized through Nile Blue loading and red light irradiation. Such nanoparticles are efficiently internalized into cells and fluoresce only after cell uptake upon exposure to red light due to extensive solvatochromic properties of Nile Blue dye (NB, 5-amino-9-(diethylamino)benzo[a]phenoxazin-7-ium perchlorate). NB is water insoluble dye (maximum excitation/emission in chloroform is 624 nm/647 nm) that fluoresces only in the lipophilic microenvironment inside cells with weak fluorescence outside cells in water buffer systems. This solves the problem of background nanoparticle fluorescence when they are administered systemically *in vivo* and have not yet been completely eliminated from the bloodstream.

The image-guided therapeutic modality was realized through the incorporation of Rose Bengal dye in the first water phase during the nanoparticle synthesis. Rose Bengal (RB, 4,5,6,7-tetrachloro-2',4',5',7'-tetraiodofluorescein) is a photosensitizer that produces reactive oxygen species upon exposure to green light. It possesses weak fluorescence upon exposure to both green and red light, does not interfere with the fluorescence

of NB, and does not generate ROS under red light thus being an ideal PDT agent within the structure of nanoparticle that is activated only on demand after the bioimaging results.

Figure 1. Internalization-responsive nano-PLGA for HER2+ breast cancer theranostics *in vivo*. Polymer nanoparticles are loaded with the photosensitizer Rose Bengal and fluorescent dye Nile Blue and equipped with the HER2-targeting IgG trastuzumab. The diagnostic dye, Nile Blue, visualizes HER2+ tumor *in vivo* under red light irradiation only after cell uptake due to its solvatochromic properties, while the therapeutic dye, generates reactive oxygen species only after green light irradiation. The differential exposure of PLGA to NIR or green light allows sequentially carrying the diagnostics and image-guided therapy of HER2+ cancer thus realizing the theranostic concept.

Thus both diagnostic and therapeutic modalities are included in the single nanoparticle, each of which is activated with the light of certain wavelengths to perform image-guide HER2-specific PDT treatment. The effective theranostic functions of anti-HER2 PLGA nanoparticles were tested on HER2-overexpressing xenograft tumors showing selective accumulation of nanoparticles in primary tumor tissue and metastases, as well as for light-induced photodynamic therapy with the achievement of complete remission.

2.2. Anti-HER2 PLGA nanoparticles synthesis and characterization

Polymer nanoparticles were synthesized by the double emulsion method as schematically illustrated in **Fig. 2a**. The first emulsion was obtained by sonication of the mixture of PLGA and Nile Blue (NB) solution in

chloroform with the solution of Rose Bengal (RB) in water. The second emulsion was obtained by the sonication of the mixture of the first emulsion and poly(vinyl alcohol) and chitosan oligosaccharide lactate in PBS.

Figure 2. Synthesis and characterization of theranostic anti-HER2 PLGA nanoparticles. (**a**) Schematic illustration of anti-HER2 targeted polymer nanoparticle synthesis *via* "water-oil-water" double emulsion method. Particles are loaded with Nile Blue as a diagnostic modality and with Rose Bengal photosensitizer as a therapeutic modality. (**b**) Scanning electron micrographs and physical size distribution of unmodified PLGA nanoparticles and anti-HER2 PLGA. Scale bar, 1 µm. (**c**) Hydrodynamic size distribution of pristine PLGA particle and anti-HER2 PLGA). (**d**) ζ-potential distribution of unmodified PLGA particles and anti-HER2 PLGA obtained with electrophoretic light scattering method.

To incorporate targeting modalities into the nanoparticle structure, the anti-HER2 antibody, trastuzumab, was added to the second water phase thus most probably incorporating into the nanoparticle structure with its slightly hydrophobic Fc fragment. The morphology of as-synthesized nanoparticles with and without trastuzumab was investigated with scanning electron microscopy (**Fig. 2b**). Data presented in **Fig. 2b** demonstrate that the synthesized nanoparticles are monodisperse structures with a shape close to spherical. According to SEM image processing using ImageJ software, the average diameter of nanoparticles was found to be 235 ± 68 nm and 232 ± 67 nm for anti-HER2 PLGA and PLGA, respectively.

The dynamic and electrophoretic light scattering measurements demonstrate the hydrodynamic size of nanoparticles to be equal to 369 ± 166 nm and 323 ± 113 nm for anti-HER2 PLGA and PLGA, respectively (**Fig. 2c**) and ζ-potential –0.66 ± 13.9 m and –3.60 ± 14.8 mV for anti-HER2 PLGA and PLGA, respectively (**Fig. 2d**).

The data obtained confirm colloidal stability of as-synthesized nanoparticles. For further *in vivo* use, the long-term colloidal stability was confirmed with the measurements of the hydrodynamic sizes of nanoparticles for 1 month in different buffer systems with pH ranging from 2 to 9. To avoid any interference of Nile Blue dye with the 633 nm laser wavelength of the ZetaSizer NanoZS analyzer, particles synthesized in the same way without NB were used (PLGA/RB). The DLS measurements showed the stability for a month of such nanoparticles in buffer solutions with a pH range of 2 to 9 – nanoparticles did not form aggregates for at least 35 days under all tested conditions (**Fig. S1**).

2.3. Theranostic anti-HER2 PLGA particles specifically target HER2-positive cells *in vitro*

The targeting abilities of theranostic anti-HER2 PLGA nanoparticles were confirmed *in vitro* by specific labeling of cancer cells with HER2 overexpression. BT-474 cells with stable expression of luciferase NanoLuc, BT-NanoLuc were used for these tests [13,22,23]. To verify the HER2 overexpression on the BT-NanoLuc cells' surface, flow cytometry tests were performed with cell labeling with fluorescent anti-HER2 antibody, trastuzumab-FITC. Chinese hamster ovary, CHO, cells without HER2 were used for control experiments [24].

Figure 3. Cancer cell targeting with anti-HER2 PLGA theranostic nanoparticles *in vitro*. (**a**) Flow cytometry analysis of HER2 expression on the surface of BT-NanoLuc and CHO cells. Blue histograms – cells labeled with trastuzumab-FITC, grey histograms – autofluorescence. Data received in FL1 channel (λ_{ex} = 488 nm, λ_{em} = 525/20 nm). (**b**) Confocal microscopy of BT-NanoLuc and CHO cells labeled with trastuzumab-FITC (λ_{ex} = 488 nm, λ_{em} = 492-500 nm) and Hoechst 33342 for nuclei staining only (λ_{ex} = 405 nm, λ_{em} = 410-520 nm). Rows (I) – cells labeled Hoechst 33342 only, rows (II) – cells labeled with trastuzumab-FITC and Hoechst 33342. Scale bars, 50 µm. (**c**) Flow cytometry test on anti-HER2 PLGA nanoparticles targeting efficiency: blue and pink histograms:

cells labeled with anti-HER2 PLGA in different concentrations, black open histograms – cells' autofluorescence. Flow cytometry data were acquired in the FL4 channel (λ_{ex} = 640 nm, λ_{em} = 675/25 nm).

Data presented in **Fig. 3a** confirm the overexpression of HER2 on BT-NanoLuc cells, namely, the median fluorescence intensities (MFI) of cell populations labeled with trastuzumab-FITC was found to equal to 207 441 and 12 706, BT-NanoLuc and CHO cells, respectively. The flow cytometry data are supported by the imaging tests: cells were labeled with trastuzumab-FITC and Hoechst 33342 and visualized with confocal laser scanning microscopy (**Fig. 3b**).

Next, we evaluated the targeting ability of anti-HER2 PLGA using the flow cytometry test. Data presented in **Fig. 3c** (and **Fig. S2** with the wide nanoparticle concentration range) confirm the specificity of targeted anti-HER2 PLGA nanoparticles interaction with HER2-overexpressing cells: MFI of BT-NanoLuc labeled with anti-HER2 PLGA is *ca.* 4.5 higher than that for CHO cells.

2.4. Fluorescence of anti-HER2 PLGA nanoparticles is cell uptake-responsive

The lipophilic nature of Nile Blue and its highly solvatochromic properties, meaning that the fluorescence emission is highly dependent on the solvent used, especially its polarity, make this dye an excellent candidate for nanoparticle imaging only after cellular uptake. This fact was confirmed using both cell-free and *in vitro* tests.

First of all, the fluorescence spectroscopy measurements were performed. Particles loaded with Nile Blue – PLGA/NB, with Rose Bengal – PLGA/RB, and with both dyes – PLGA/NB/RB were excited with λ_{ex} = 500-530 nm suitable for Rose Bengal excitation and with λ_{ex} = 590-630 nm suitable for Nile Blue excitation in water buffer system (PBS) as well as in polar aprotic solvent, namely, dimethyl sulfoxide (DMSO).

Figure 4. Nile Blue-guided solvatochromic nature of anti-HER2 PLGA nanoparticles. Fluorescence emission spectra of targeted anti-HER2 nanoparticles with different dye loadings in PBS and DMSO upon λ_{ex} = 500 nm to 630 nm were registered.

Data presented in **Fig. 4** confirm that the fluorescence significantly increases when the water buffer system is changed to polar aprotic solvent, moreover, the incorporation of both dyes into nanoparticle structure

(PLGA/NB/RB) significantly enhances this effect compared to nanoparticle loaded with Nile Blue only (PLGA/NB).

Next, imaging of HER2-overexpressing cells anti-HER2 PLGA nanoparticles was performed. Cells were seeded at 96-well plates and anti-HER2 PLGA particles with different dye loadings – with RB only, with NB, and with NB/RB were added to cells in equal concentrations. Cells were incubated with nanoparticles for 1 h and then imaged with confocal laser scanning microscopy without any washing steps under the following excitation: λ_{ex} = 561 nm (the best for RB excitation) and λ_{ex} = 640 nm (the best for NB excitation) and emission: λ_{em} = 600/52 nm (the best for RB emission) and λ_{em} = 647LP nm (the best for NB emission). Data presented in **Fig. 5** show that the most effective fluorescence is observed for PLGA/NB and PLGA/NB/RB inside cells under the irradiation of cells with 640 nm in the fluorescence channel corresponding to Nile Blue (647 LP). Meanwhile, the fluorescence of nanoparticles outside cells is very weak and observed outside cells due to RB dye only under irradiation with 561 nm in the fluorescence channel corresponding to RB (600/52 nm).

Moreover, similarly to the fluorescence spectroscopy tests, the fluorescence of nanoparticles inside the polar microenvironment (here, cell compartments) is significantly higher for particles loaded with both NB and RB dyes in comparison with particles loaded with RB only.

Figure 5. Cell uptake-responsive fluorescence of anti-HER2 PLGA nanoparticles due to Nile Blue loading. Confocal microscopy of BT-NanoLuc cells incubated with anti-HER2 PLGA nanoparticles loaded with fluorescent dyes: Rose Bengal (left), Nile Blue (center), and Nile Blue-Rose Bengal (right) without any washing steps from non-bound nanoparticles. Scale bars, 50 µm.

2.5. Anti-HER2 PLGA nanoparticles provide light-induced tunable cytotoxicity to HER2-overexpressing cancer cells *in vitro* through ROS generation

Along with an effective imaging dye, NB, anti-HER2 PLGA nanoparticles were loaded with RB dye, which is a photosensitizer and is used as an antibacterial agent due to the ROS generation, as well as in studies for melanoma treatment. To find the most effective RB concentration during the nanoparticle synthesis and

prevent issues associated with dye aggregation and stacking inside the nanoparticle architecture, the ROS generation efficacy study was performed. Particles were synthesized with dyes at a concentration of 1 g/L NB and a serial 2-fold dilution of RB within the $0.2 \div 3$ g/L range, as well as particles synthesized with only one dye, NB or RB.

For the preliminary assessment of the therapeutic properties of the targeted particles, the ROS generation was measured when the targeted particles were exposed to green laser light, $\lambda_{ex} = 532$ nm. Due to the known properties of Nile Blue to act as a photosensitizer under the 808 nm laser irradiation [25], $\lambda_{ex} = 808$ nm and the combination of $\lambda_{ex} = 532$ and 808 nm were used for nanoparticle irradiation to verify the combination effect of green and NIR irradiation for photosensitization. The total reactive oxygen species assay kit, 520 nm (Thermo) was used to assess the efficacy of ROS generation *via* the fluorescence intensity measurement. According to the data obtained (**Fig. 6a**), the higher the RB concentration, the higher ROS production under $\lambda_{ex} = 532$ nm irradiation. Interestingly, the addition of NB into nanoparticles does not affect the ROS generation under the green light irradiation and slightly decreases ROS when the combination of green&NIR light is used (**Fig. 6a**).

Next, the viability of HER2-overexpressing cells labeled with anti-HER2 PLGA nanoparticles was measured. To assess the percentage of viable cells when exposed to targeted particles and external light irradiation, a series of MTT tests and colony formation assays were performed (**Fig. 6b,c,d, Fig. S3**). To test the effectiveness of the cytotoxic effects of targeted particles with dual dye loading, the cytotoxicity was compared with the effect of targeted particles with single dye loading. Particles were synthesized with NB, RB and NB/RB. Next, the BT-NanoLuc cells were incubated with particles and irradiated with a green laser and, with a combination of green and NIR lasers. Cells incubated with particles without irradiation were used as controls. Data presented in **Fig. 6b** demonstrate that in contrast to the cell-free ROS study, particles loaded with both dyes exhibit a stronger cytotoxic effect in comparison with particles with NB only under the irradiation with both green and green&NIR light (**Fig. 6b**). This is probably due to the additional cytotoxic effects of the NB dye when exposed to a NIR light that is not associated with the release of reactive oxygen species; no enhanced cytotoxic effect was observed when exposed to the NIR light on particles with RB, which confirms the effect of NIR light only on NB (**Fig. 6b**).

In view of the findings for NB cytotoxicity not entirely correlated with *in vitro* ROS production, we further investigated the effect of RB concentration (during the synthesis of PLGA) on light-induced cytotoxicity. The data presented in **Fig. 6c** confirm that the concentration of 3 g/L during the synthesis is still optimal, which was used in further experiments, namely in concentration- and time-dependent studies. BT-NanoLuc cells were incubated with anti-HER2 PLGA in different concentrations (**Fig. 6d**) and for different incubation times (**Fig. S4**), washed from non-bound particles, and irradiated with the external laser light. Data presented in **Fig. 6d** show that the higher concentration, the higher the light-induced cytotoxicity, moreover the anti-HER2 PLGA-induced cytotoxicity increased significantly when the nanoparticles enter the cell – the cytotoxicity is significantly higher

after the prolonged incubation of nanoparticles with cells at +37 ºC, and the cytotoxicity can be slightly increased using the combination of green and NIR light irradiation (**Fig. S4**).

Figure 6. Light-induced cytotoxicity of anti-HER2 PLGA nanoparticles to HER2-overexpressing cancer cells *in vitro* through ROS generation. (**a**) ROS generation by targeted PLGA nanoparticles upon the irradiation with lasers: λ_{ex} = 532 nm, 808 nm, or 532/808 nm, dependence on RB and NB loading into PLGA nanoparticle during synthesis. ROS was measured with total reactive oxygen species assay kit, 520 nm (Thermo). (**b**) BT-NanoLuc (HER2-overexpressing cells) viability was assessed with the MTT test. Cells were labeled with different nanoparticle formulations and irradiated with lasers: λ_{ex} = 532 nm, 808 nm or 532/808 nm for 10 min. Unlabeled and non-irradiated cells were used as control. (**c**) BT-NanoLuc viability assay: the influence of RB concentration during the synthesis of PLGA on the nanoparticle cytotoxicity was studied. (**d**) BT-NanoLuc viability assay: the concentration-dependent cytotoxicity of anti-HER2 PLGA nanoparticles loaded with NB (1 g/L) and RB (3 g/L) was studied under the irradiation with lasers: λ_{ex} = 532 nm, 808 nm, or 532/808 nm for 10 min.

2.6. Light irradiation of cells labeled with anti-HER2 PLGA causes cell death *via* apoptosis

To study the mechanism of HER2-positive cell death after light irradiation, we first examined the dynamics of cell growth. The growth rate of HER2-positive cells was investigated using a real-time cell analysis (RTCA) E16 xCelligence system (**Fig. 7**, **Fig. S5**). BT-NanoLuc cells were incubated with anti-HER2 PLGA particles, and irradiated with a green laser. Irradiated and non-irradiated cells with particles were seeded into E16 plates and allowed to grow for 150 h (6.2 days). The dynamics of cell growth presented in **Fig. 7a** demonstrate that under irradiation, the dynamics of cell growth depend significantly on the particle concentration. However, in the absence of irradiation, the cell growth is slightly affected with targeted nanoparticles, the only significant

decrease is observed for the highest concentration used. The data presented in **Fig. S5** demonstrate that the cell growth dynamic reduction after the irradiation is related mainly to RB loading into the nanoparticle and slightly affected by NB loading (**Fig. S5**).

Figure 7. Cell death mechanism study. (**a**) Real-time cell analysis of BT-NanoLuc cell growth rate after the incubation with anti-HER2 PLGA nanoparticles and green light irradiation: the cell index dependence on the time of cultivation. (**b**) Annexin V/PI flow cytometry analysis of cells treated with anti-HER2 PLGA and irradiated with a green light for 0 – 40 min. Flow cytometry dot plots indicate the intensity of PI (a marker of late apoptosis/necrosis) fluorescence on Annexin V fluorescence (a marker of apoptosis). The viable cells, early apoptotic and late apoptotic/necrotic cells are represented in the bottom left quadrant (Annexin V⁻/PI⁻), bottom right (Annexin V⁺/PI⁻), and upper right (Annexin V⁺/PI⁺) quadrants, respectively.

The obtained data indicate that cell death is caused most probably by the quite slow mechanism of programmed cell death (apoptosis) and not by immediate death due to injury (necrosis). To clearly determine the mechanism of cell death, the Annexin V-Alexa Fluor 488/propidium iodide flow cytometry test was used. BT-NanoLuc cells were incubated with anti-HER2 PLGA nanoparticles at various concentrations, irradiated with a green laser for 0 to 40 min, and seeded into 6-well plates. Cells were cultured for 12 h, and then stained with Annexin V-Alexa Fluor 488/propidium iodide and analyzed with flow cytometry (**Fig. 7b, Fig. S6**). The data presented in **Fig. 7b** demonstrate the presence of apoptotic (right bottom quadrant) and late apoptotic/necrotic (right top quadrant of flow cytometry dot plot) cells depending on the nanoparticle concentration and irradiation time thus confirming the cell death route *via* apoptosis.

2.7. Fluorescent bioimaging of HER2-overexpressing tumors with anti-HER2 PLGA nanoparticles *in vivo*

The diagnostic properties of anti-HE2 PLGA were studied by their ability to visualize the localization of HER2-overexpressing tumors in vivo. BALB/c Nu/Nu mice underwent subcutaneous transplantation with BT-NanoLuc cells that possess HER2 overexpression. The stable expression of luciferase NanoLuc allows tracking the tumor growth progress and metastasis spreading non-invasively in vivo through the intravenous injection of h-coelenterazine – specific substrate of NanoLuc luciferase. When the tumor volume reaches 60 mm^3, mice were injected with h-coelenterazine and the bioluminescence of the primary tumor on the right flank and metastases on the opposite flank was registered with the IVIS Spectrum CT bioimaging device (**Fig. 8a**). Next, anti-HER2 PLGA were injected at different doses and the fluorescence (λ_{ex} = 640 nm, λ_{em} = 680 nm) was registered after 1h, 2h, 6h, 12h, 18h of injection. Data presented in **Fig. 8a** illustrate the accumulation of anti-HER2 PLGA within the primary tumor area as well as in metastases with significant fluorescence quenching after 2-6 h of injection depending on the injected dose. However, the highest injected dose resulted in stable fluorescence even up to 18 h post-injection. However, the dose of 3 mg was chosen as the optimal one for further tests since it results in stable and informative fluorescence for more than 2 h after injection.

Figure 8. Bioimaging properties of anti-HER2 PLGA nanoparticles *in vivo*. (**a**) BALB/c Nu/Nu mice with s.c. BT-NanoLuc tumors possessing HER2 overexpression were injected with anti-HER2 PLGA and visualized 1, 2, 6, 12, and 18 h post-injection (λ_{ex} = 640 nm, λ_{em} = 680 nm). The primary BT-NanoLuc tumor and metastases are visualized through the i.v. injection of h-coelenterazine and subsequent bioluminescence *in vivo* imaging (left

panel). (b) Mice with two tumors: 1) MCF7-iRFP tumors (left flank, pink circle, imaged *via* fluorescence detection) and 2) BT-NanoLuc tumors (right flank, blue circle, imaged *via* bioluminescence detection) were injected with anti-HER2 PLGA and 1h post-injection the fluorescence of anti-HER2 PLGA was registered ($\lambda_{ex} = $ 640 nm, $\lambda_{em} = $ 680 nm).

Since the HER2 receptor is typically expressed in normal tissues of the human body but in a significantly lower amount compared to breast cancer cells, the issue of the selectivity of as-synthesized anti-HER2 PLGA in terms of *in vivo* HER2-binding should be resolved prior to therapy study. To address this question, we used BALB/c Nu/Nu mice with two tumors possessing different HER2 expression. BT-NanoLuc cells with HER2 overexpression were s.c. injected into the right flank, and MCF-7-iRPF fluorescent cells with normal HER2 expression were injected into the left flank. The distribution of BT-NanoLuc cells is registered through the bioluminescence detection, while the localization of MCF-7-iRFP cells is registered through the fluorescence of iRFP protein detection. These mice underwent the injection of anti-HER2 PLGA followed by fluorescent bioimaging (**Fig. 8b**). Data presented in **Fig. 8b** confirm that anti-HER2 PLGA localized predominantly in BT-NanoLuc tumors and metastasis with non-significant accumulation in MCF-7-iRFP tumors.

2.9. Anti-HER2 PLGA-based photodynamic image-guided therapy eradicates HER2+ xenogeneic ductal carcinoma *in vivo*

The therapeutic efficacy of anti-HER2 PLGA was assessed in vivo. Treatment of xenograft tumors was carried out on day 14 when the tumor volume reached ca. 60 mm^3. Mice were randomly divided into three groups (n = 4) and treated with i) PBS, ii) anti-HER2 PLGA and iii) anti-HER2 PLGA with green light irradiation (**Fig. 9a**). The primary tumors were irradiated for 10 min and areas with metastasis for 3 min. The tumor growth measurements with the caliper (**Fig. 9b**) were accompanied by the bioluminescent in vivo imaging (**Fig. 9c**).

In vivo bioimaging of experimental animals demonstrate the high therapeutic potential of anti-HER2 PLGA. Tumor growth was fully inhibited by the combination of ani-HER2 PLGA and light irradiation – no luminescence of tumors and metastases was observed until the end of the experiment (40 days from the treatment start).

3. Discussion

Theranostic agents are powerful synthetic compounds that are capable of selectively killing tumor-cell populations and/or inhibiting their growth and proliferation. The development of a new generation of these agents is critical for the advancement of cancer diagnosis and treatment, as well as for the treatment of other diseases [26,27]. At the same time, the development and translation into clinical practice of oncotheranostic agents are significantly limited by a number of intractable problems. Such agents are complex multicomponent systems that are not only difficult to synthesize but also difficult to thoroughly characterize, obtaining drug samples with acceptable batch-to-batch reproducibility. In particular, methods of chemical conjugation of nanoparticles with targeting molecules, using, for example, standard cross-linkers (e.g., EDC/NHS), lead to non-reproducible results of the conjugation reaction due to highly competitive and yet uncontrolled hydrolysis reactions.

Figure 9. Photodynamic therapy of HER2-positive tumors realized with theranostic anti-HER2 PLGA nanoparticles leading to cancer remission. (**a**) A treatment scheme based on anti-HER2 PLGA. BALB/c Nu/Nu mice were s.c. implanted with $2 \cdot 10^6$ BT474-NanoLuc cells. After 14 d of tumor inoculation, mice were randomized into groups (n = 4) and received either the injection of anti-HER2 PLGA or a combination of anti-HER2 PLGA with light irradiation. (**b**) Tumor growth dynamics: control group, group with the single injection anti-HER2 PLGA, and group with single injection anti-HER2 PLGA and the green light irradiation of primary tumor and metastases. (**c**) Bioimaging of BT474-NanoLuc tumor xenografts: (1) before treatment, (2) and (3) – after treatment. BALB/c Nu/Nu mice were injected with 5 µg of h-coelenterazine, and bioluminescence was recorded with IVIS Spectrum CT.

These problems are solved by the development of nanoagents that are modified by targeting molecules already during the synthesis process [28–30], or by sterically oriented self-assembly of components through various interfaces, such as streptavidin/biotin [31], SpyTag/SpyCatcher [32,33], barnase/barstar [34,35]. However, considering *in vivo* applications, such approaches, on the one hand, add multifunctionality to drug delivery systems by the possibility of changing self-assembled modules, on the other hand, they significantly affect the size of systems and their immunogenicity, in particular, streptavidin possesses significant immunogenicity [36]. An alternative

solution is the development of fully genetically encoded self-assembling protein nanoparticles with targeted modality which is introduced into the protein's genetic construct [37].

However, from the point of view of loading with therapeutic and diagnostic compounds, protein particles are not as versatile as, for example, liposomes or polymer matrices: upon loading, the tertiary structure of the protein can be disturbed, which affects the efficiency of self-assembly and subsequent possibilities in target recognition. At the same time, the simultaneous loading of hydrophilic and hydrophobic substances into protein particles is a really difficult task while maintaining protein functionality.

To overcome described issues, we developed the method for the synthesis of theranostic polymer nanoparticles equipped with HER2-targeting antibodies without chemical conjugation. For this purpose, a hydrophobic polymer, PLGA, loaded with a therapeutic and diagnostic compound was used. The surface of PLGA particles was equipped by incorporating the anti-HER2 antibody into the particle during the microemulsion synthesis – it is most likely, that the Fc fragment of the antibody, due to its inner hydrophobic region, is incorporated into the hydrophobic structure of the PLGA polymer thus covering the surface of nanoparticle with HER2-targeting molecules without the need for chemical conjugation.

PLGA polymer is widely used for extended drug release both in fundamental life science and in clinical practice, and now more than 20 PLGA-based medications are already introduced into the clinic for the treatment of breast and prostate cancer, acromegaly, osteoarthritis, diabetes, growth hormone deficiency treatment, and ocular diseases [12,38]. PLGA is a biodegradable copolymer possessing by hydrolyzing in water ester bonds forming lactic and glycolic acids, which are by-products of *in vivo* metabolism [38,39]. Here we use poly(D,L-lactide-co-glycolide), ester terminated, lactide:glycolide 85:15, Mw 190,000-240,000 polymer. The 85:15 ratio was chosen based on the screening of previously published works. Namely, the ratio of monomers of lactic and glycolic acids in the copolymer affects the rate of hydrolysis, and, accordingly, the time of the release of the loaded substance in the particle. When using polymers, this parameter must be taken into account so that the particle with the substance does not degrade prematurely in the bloodstream before reaching the targeted organ. It has been shown that with a ratio of lactic and glycolic acids in the polymer of 85:15, the release time of the drug in the body increases, compared to 75:25 and 50:50 ratios [26]. Moreover, the loading of therapeutic substances in a polymer with a ratio of 85:15 is higher than with a 50:50 ratio [40].

The "water-oil-water" double microemulsion synthesis method of PLGA nanoparticles proved its versatility for loading different substances – both hydrophobic and hydrophilic [13,20,41,42]. To incorporate diagnostic modality into nanoparticle structure, we used hydrophobic Nile Blue dye from the benzophenoxazine family with high fluorescence, high quantum yield, and excellent photostability [43]. The HER2-positive imaging process was guided using the fluorescent properties of Nile Blue dye and infrared light that excites it. Nile Blue (NB, Nile Blue A, 5-amino-9-(diethylamino)benzo[a]phenoxazin-7-ium perchlorate) is a water-insoluble dye, possessing maximum excitation and emission in chloroform 624 nm/647 nm, thus entering the NIR-I biological transparency window. NB is being used as a histological stain for *in vitro* and *ex vivo* imaging of lysosomes and

lipids [43], pH sensorics [44], and several *in vivo* studies demonstrated the effective NB accumulation in the tumor after systemic administration [25,45]. The extensive solvatochromic properties of NB make this dye an excellent candidate for imaging specific cancer cells only after cell internalization. We showed that in water buffer systems the fluorescence of these nanoparticles is very weak, however, after cell internalization, the fluorescence significantly increases in the lipophilic microenvironment inside cells. Considering *in vivo* applications of this property it is worth mentioning that this feature solves the problem of background fluorescence of nanoparticles after systemic administration: when the particles are not completely eliminated from the organism, and the tumor accumulation still did not reach the plateau, the real-time diagnostics is already possible since background fluorescence of circulating particles is not an obstacle for tumor accumulation registration. We showed that the synthesized targeted particles are unique diagnostic agents, since a single injection of particles into a mouse with the HER2-positive tumor model it is possible to specifically visualize HER2-positive tumors for a long time (up to 18 h).

To incorporate diagnostic modality into nanoparticle structure, we used hydrophilic fluorescent dye, Rose Bengal. Rose Bengal (RB, 4,5,6,7-tetrachloro-2',4',5',7'-tetraiodofluorescein) is a photosensitizer that produces ROS upon exposure to green light [20,46]. Previously, we experimentally proved that cancer cells can be selectively eliminated when exposed to RB in the PLGA particle and green light irradiation [20]. RB is a promising diagnostic and therapeutic agent as it is already being used in the treatment of eye diseases [47,48] as well as in the treatment of skin cancer [49,50]. The water-soluble photosensitizer RB fluoresces at 575 nm when exposed to the green light at a wavelength of 532 nm. Light with a wavelength of 532 nm is located on the border with the transparency window of biological tissue, however, green light in contrast to red light causes fewer side effects under prolonged exposure [51,52]. In our work, we showed that when exposed to targeted anti-HER2 PLGA nanoparticles loaded with RB and NB and irradiated with a green laser at a wavelength of 532 nm (100 mW), mice with xenograft HER2-positive tumors were completely cured. These data demonstrate the safe and effective use of green light in combination with anti-HER2 PLGA in anticancer PDT since no necrotizing of both tumor and healthy tissue was observed during the treatment.

The obtained nanoparticles were shown to be effective theranostic agents possessing both HER2-specific bioimaging properties and light-inducible photodynamic therapy *in vivo* when it is necessary. The *in vitro* and *in vivo* efficacy of anti-HER2 PLGA nanoparticles were thoroughly tested and 100% inhibition of HER2-positive tumor growth in all mice from the experimental group was demonstrated, thus confirming the great potential of the synthesized nanomedication for personalized medicine.

4. Materials and Methods

4.1. Materials

Chemicals: Sigma, Germany: poly (lactic-co-glycolic) acid (PLGA) (lactide: glycolide 85:15, MW 190-240 kDa), chitosan oligosaccharide lactate, poly (vinyl alcohol) (Mowiol 4-88, PVA, MW 31 kDa), Rose Bengal,

Nile Blue A perchlorate; Helicon, Russia: BSA; Chimmed, Russia: chloroform; Roche, Switzerland: Herceptin (Trastuzumab); PanReac Applichem, Germany: dimethylsulfoxide (DMSO).

4.2. Synthesis of anti-HER2 PLGA nanoparticles

The targeted particles were synthesized by the «water-oil-water» double emulsion method. The first emulsion was formed by adding 150 µL of 3 g/L Rose Bengal in Milli-Q water and 10 µL of 1 g/L Nile Blue to 300 µL of 40 g/L PLGA in chloroform. The first emulsion was subjected to ultrasound with 750 W Sonicator Vibra-Cell (Sonics & Materials, Inc, USA) for 1 min with 40 % power and 1 min with 60 % power. The second emulsion was prepared by adding the first emulsion to 3 mL of 3 % PVA in water Milli-Q containing 0.3 g/L chitosan oligosaccharide lactate and 200 µL of 10 g/L trastuzumab. The solution was sonicated for 1 min with 40 % power and 1 min with 60 % power at +4 ℃. The chloroform was then evaporated using the rotator at 20 rpm for 3 h at room temperature. The particles were then washed with PBS, pH 7.4 by triple centrifugation and finally resuspended in 100 µL of PBS, pH 7.4. The final concentration of nanoparticles was determined by drying 70 µL of particles at 60 °C.

4.3. Scanning electron microscopy

The morphology of particles was evaluated with scanning electron microscopy at an accelerating voltage with a MAIA3 electron microscope (Tescan, Czech Republic) at an accelerating voltage of 5 kV. The physical size of the targeted particles was determined with ImageJ software to get the particle size distribution.

4.4. Dynamic and electrophoretic light scattering measurements

The hydrodynamic size of PLGA nanoparticles was determined using a Zetasizer Nano ZS analyzer (Malvern Instruments Ltd, UK) in Milli-Q water at 25 ℃. Z-potential was measured using Zetasizer Nano ZS (Malvern Instruments Ltd, UK) in 10 mM KNO_3 at 25 °C. Particle stability was measured with a Zetasizer Nano ZS (Malvern Instruments Ltd, UK) through consecutive measurements of the size within one month. The particles were diluted in buffer solutions with pH from 2 to 9 and the hydrodynamic size of the particles was measured. The following buffer systems were used: pH 2 (0.1 M glycine-HCl), pH 3 (0.1 M glycine-HCl), pH 4 (0.15 M citrate-phosphate buffer), pH 5 (0.15 M citrate-phosphate buffer), pH 6 (0.1 M HEPES buffer), pH 7 (0.15 M phosphate buffer), pH 8 (0.1 M TRIS buffer), pH 9 (0.1 M TRIS buffer).

4.5. Cell culture

The source of cell lines is the collection of the laboratory of molecular immunology, Institute of Bioorganic Chemistry, Russian Academy of Sciences: human breast cancer cells BT-474 with stable NanoLuc expression, BT-NanoLuc [13], MCF-7 with iRFP protein expression, MCF-7-iRFP, Chinese hamster ovary, CHO. BT-NanoLuc cells were cultured in DMEM growth medium (Gibco, USA) supplemented with 10% fetal bovine

serum (Capricorn, Germany), 2 mM L-glutamine (PanEko, Russia), 50 U/mL/50 µg/mL penicillin-streptomycin (PanEko, Russia), 1 µg/mL puromycin (Sigma, Germany) and 10 µg/mL ciprofloxacin (Elfa, India) in culture flasks (Nunc, Denmark) in a CO_2 incubator (BINDER, Germany) at 37 °C with 5 % CO_2. MCF-7-iRFP cells were cultured in FluoroBright DMEM growth medium (Gibco, USA) with 10% fetal bovine serum (Capricorn, Germany), 2 mM L-glutamine (PanEko, Russia), 50 U/mL/50 µg/mL penicillin-streptomycin (PanEko, Russia) and 10 µg/mL ciprofloxacin (Elfa, India) in culture flasks (Nunc, Denmark) in a CO_2 incubator (BINDER, Germany) at 37 °C with 5% CO_2. Cells were removed from the surface of culture flasks using a 2 mM EDTA solution (PanEko, Russia).

4.6. Confocal microscopy

Visualization of HER2-overexpressing cells labeled with targeted nanoparticles was performed on the Leica DMI6000B (Leica Microsystems, Germany) microscope equipped with confocal microscopy upgrade (Thorlabs, USA) at the following conditions. BT-NanoLuc cells were seeded at 96-well flat-bottomed culture plates in 100 µL of DMEM growth medium, incubated overnight under a humidified atmosphere with 5 % CO_2 at 37 °C and anti-HER2 PLGA targeted particles were added in 100 µL of full culture medium to get a final concentration of nanoparticles 10 µg/mL. Cells were incubated with nanoparticles for 1 h at 37 °C, and washed from unbound particles with subsequent microscopy tests. Cells were imaged with green and red lasers separately and in combination, namely: 561 nm green laser excites Rose Bengal dye; 640 nm red laser excites Nile Blue dye. Emission filters 600/52 nm and 647LP nm were used for visualization for all lasers.

Trastuzumab-FITC binding was visualized using an LSM 980 (Zeiss) confocal microscope. Cells were treated with 1 µg/mL Hoechst33342 and 2 µg/mL trastuzumab-FITC in PBS with 1% BSA for 30 min, washed with PBS and imaged under the following conditions: (λ_{ex} = 488 nm, λ_{em} = 492-500 nm for trastuzumab-FITC detection and λ_{ex} = 405 nm, λ_{em} = 410-520 nm for Hoechst 33342 detection.

4.7. Fluorescence spectroscopy of targeted nanoparticles

Targeted nanoparticles loaded with Rose Bengal and Nile Blue dyes were resuspended in 100 µL of PBS at 10 µg/mL, pH 7.4, and/or in 100 µL of DMSO and added to the 96-well plate. The particle fluorescence was measured using an Infinite M100 Pro spectrophotometer (Tecan, Austria) within excitation wavelengths 500-600 nm range.

4.8. ROS study

The generation of reactive oxygen species by targeted particles was measured using irradiation laser wavelengths of 532 nm and/or 808 nm. Particles were synthesized with different concentrations of dyes. To perform the ROS generation test, 600 µL of PBS pH 7.4 was added to 1.5 mL tubes, next 20 µL nanoparticles at 30 g/L were added, the sample was divided into 4 tubes, and the Invitrogen reagent (Total Reactive Oxygen

Species (ROS) assay kit, 520 nm) was added to these samples according to the manufacturer's recommendations. Then samples were irradiated with lasers individually and in combination (532 nm, 808 nm, 532&808 nm) for 5 min. Then, the fluorescence of irradiated nanoparticles was measured in 96-well plates with 488 nm excitation and 525 nm emission with a gain of 149 on an Infinite M100 Pro spectrophotometer (Tecan, Austria).

4.9. MTT assay

The cytotoxicity of nanoparticles under different conditions was determined using the MTT test.

i) The efficacy of dual dye loading into nanoparticles was determined as follows. $75 \cdot 10^3$ BT-NanoLuc cells were added to 1.5 mL tubes in 300 µL of 1% fat-free milk in PBS, next 25 µL 30 g/L of particles were added, samples were incubated for 15 min at +4 ℃, and washed from unbound particles by triple centrifugation. Finally, cells were resuspended in 600 µL of 1% BSA and divided into 200 µL samples in 1.5 mL tubes. Cells were irradiated for 1) 532 nm laser, 100 mW (KLM-532-200 Optronic, Russia), 10 min, room temperature; 2) 808 nm laser (KLM-H808-1200-5 Optronic, Russia), 1200 mW, 10 min, 30 ℃, 3) the combination of both lasers. Next, cells were seeded into 96-well plates at a concentration of 5000 cells per well in 200 µL of complete DMEM medium and cultured for 48 h.

Then the medium was removed, 100 µL of MTT solution at 0.5 g/L was added to each well, and plates were incubated for 1 h at 37 ℃ and 5% CO_2. Next, the MTT solution was removed and 100 µL of DMSO was added. The absorbance was measured at a wavelength of $\lambda = 570$ nm (reference $\lambda = 630$ nm) using an Infinite M100 Pro plate reader (Tecan, Austria).

ii) The concentration-dependent cytotoxicity of anti-HER2 PLGA/NB/RB was determined similarly by incubating cells with 15, 7.5, 3.75, 1.8, 0.9, 0.45 g/L anti-HER2 PLGA/NB/RB.

iii) The cytotoxicity of nanoparticles loaded with different concentrations of RB was determined similarly by incubating cells with particles synthesized with different RB concentrations (3, 1.5, 0.75, 0.4, 0.2 g/L).

iv) The influence of incubation time on cytotoxicity was determined similarly by irradiation cells 1) immediately after the washing from non-bound particles, 2) after 1 h of incubation at 37 ℃, 3) after 4 h after incubation at 37 ℃.

4.10. Colony formation assay

BT-NanoLuc cells were incubated with targeted anti-HER2 PLGA/NB/RB particles in different concentrations (15, 7.5, 3.75, 1.8, 0.9, 0.45 g/L) in 1.5 mL test tubes at a concentration of $75 \cdot 10^3$ in 100 µL of colorless DMEM medium. Incubation was carried out at +4 ℃ on a rotator for 15 min. Cells were washed from unbound particles and resuspended in 100 µL of PBS with 1% BSA. Next, the cells with particles were irradiated for 10 min with 532 nm and/or 808 nm for 10 min. Irradiated BT-NanoLuc cells with targeted particles were added to the wells of a 12-well plate: 10^3 cells in 1200 µL of complete DMEM medium. The plates were incubated at 37 ℃ and 5% CO_2 for 7 days until colonies of more than 50 cells were formed. After incubation, the medium

was removed, and the wells were washed with 600 µL of PBS and 600 µL of 70% ethanol in PBS. Next, 600 µL of 95% ethanol was added, and wells were incubated for 15 min. Next, after 95% ethanol was removed, the wells were washed with tap water and 600 µL of 1% crystal violet was added. Wells were incubated for 30 min at room temperature. The crystal violet was then removed and the wells were washed 10 times with tap water. The plates were scanned (EPSON Perfection 2400 Photo, Indonesia).

4.11. Real-time cell survival analysis

Real-time cell survival analysis was performed on the xCELLigence instrument (Agilent, USA). $3 \cdot 10^3$ of BT-NanoLuc cells were incubated with nanoparticles PLGA/RB, PLGA/NB, PLGA/NB/RB, and PLGA/NB/RB) at various concentrations in 100 µL of PBS for 15 min at +4 ℃. Next, the cells were washed from unbound particles 3 times with 200 µL of PBS and resuspended in 300 µL of phenol red-free complete growth medium.

Next, samples were irradiated for 10 min with a 532 nm laser (100 mW). Then, 150 µL of irradiated and non-irradiated cells with particles were added to each well of the E16 xCELLigence plate, into which 50 µL of a complete phenol red-free growth medium DMEM were previously added. The binding kinetics and cell growth were measured for 150 h at 37 ℃ and 5% CO_2.

4.12. Apoptosis test

The Invitrogen reagent kit (Molecular Probes Dead Cell Apoptosis Kit with Annexin V Alexa Fluor 488 & Propidium Iodide, PI) was used to study the mechanism of cell death. $400 \cdot 10^3$ of BT-NanoLuc cells were incubated with anti-HER2 nanoparticles at 2 concentrations, washed from non-bound particles by centrifugation and resuspended in 400 µL of colorless FluoroBrite DMEM medium (Gibco, USA). Next, 500 µL of a FluoroBrite DMEM medium was added to 6-well plates, and 100 µL of $10 \cdot 10^4$ cells incubated with nanoparticles were added per well. Next, the cells were irradiated for 0, 1, 3, 10, 20, 40 min with a 532 nm. Cells without the addition of particles and without irradiation were used as controls. The plates were incubated for 12 h at 37 ℃ and 5% CO_2. After incubation, the medium was removed from the plates and the cells were detached with 2 mM EDTA solution. Annexin V and PI reagents were added to the obtained samples according to the manufacturer's recommendations. The samples were then evaluated on a NovoCyte 3000 VYB cytometer (ACEA Biosciences, USA) in the BL1 channel (excitation laser 488 nm, emission filter 530/30 nm) for Annexin V and the BL3 channel (excitation laser 488 nm, emission filter 615/20 nm) for PI.

4.13. Tumor-bearing mice

Female BALB/c Nu/Nu mice (20-25 g) were obtained from the vivarium of the Institute of Bioorganic Chemistry, (Pushchino, Russia). The mice were housed in specific pathogen-free facilities with free access to food and water. All procedures were approved by the IBCh RAS Institutional Animal Care and Use Committee (protocol No. 298/2020, dated 29 May 2020). Experiments were performed using Zoletil (Virbak, France) and

Rometar (Bioveta, Czech Republic) anesthesia at a dose of 25/5 mg/kg. Aerrane (Isoflurane, Baxter) inhalation was performed for anesthesia using the RAS-4 Rodent Anesthesia System (Perkin Elmer, Waltham, MA, USA) for experiments with IVIS Spectrum CT living imaging system (Perkin Elmer, Waltham, MA, USA).

Xenograft tumors were obtained by inoculating $2 \cdot 10^6$ BT-NanoLuc cells with 30% Matrigel (Corning, USA) in 100 µL of complete culture medium subcutaneously into the right flank of BALB/c Nu/Nu mice. $2 \cdot 10^6$ MCF-7-iRFP cells were s.c. inoculated into the left flank of BALB/c Nu/Nu mice.

4.14. *In vivo* bioimaging

Bioluminescent images of xenograft tumors were obtained through i.p. injection of h-coelenterazine (200 µL, 25 µg/mL in PBS). Next, bioluminescent images were captured with an open filter on the IVIS Spectrum CT system (Perkin Elmer, Waltham, MA, USA) IS1803N7357 iKon camera (Andor, USA) using 10-s exposure time (f⁄stop = 1, binning = 8, the field of view 13.4 cm x 13.4 cm) and normalized to photons per second per cm^2 per steradian (p/sec/cm^2/sr). For nanoparticle localization studies, fluorescent images were obtained respectively under the following conditions: excitation 640 nm, emission 680 nm, exposure = 1s, f⁄stop = 2, binning = 8, the field of view 13.4 cm x 13.4 cm. MCF-7-iRPF tumor imaging was performed under the following conditions: excitation 605 nm, emission 660 nm, exposure = 1s, f⁄stop = 2, binning = 8, the field of view 13.4 cm x 13.4 cm.

4.15. Therapy study

For the treatment experiments, mice were challenged with cancer cells, then randomly divided into three groups (n = 4 in each). The tumor growth dynamic was assessed with the tumor volume measurement using the caliper and tumor volume calculation with the formula $V = a^2 \cdot b/2$, where a is the width and b is the length of the tumor. When the tumor volume reached 60 mm^3, mice received an i.p. injection of nanoparticles in PBS. 1h after the injection, the fluorescent images were obtained and mice were irradiated with the light (laser, 532 nm, 100 mW) according to the obtained images. Namely, primary tumors (10 min) and lymph nodes (3 min) with the localization of targeted nanoparticles were irradiated.

4.16. Statistical analysis

All experiments were performed independently three times. Numerical data are shown as mean ± standard deviation. Significant differences were determined using a two-tailed Welch t-test. Statistically significant – P <0.05 (*), 0.01 (**), 0.001 (***).

References

1. Vlamos, P. *GeNeDis 2016. Geriatrics / Panayiotis Vlamos, Editor* (Springer, Cham, 2017).
2. Kelkar, S. S. & Reineke, T. M. Theranostics: combining imaging and therapy. *Bioconjugate chemistry* **22,** 1879–1903; 10.1021/bc200151q (2011).
3. Safra, T. *et al.* Pegylated liposomal doxorubicin (doxil): reduced clinical cardiotoxicity in patients reaching or exceeding cumulative doses of 500 mg/m2. *Annals of oncology : official journal of the European Society for Medical Oncology* **11,** 1029–1033; 10.1023/A:1008365716693 (2000).
4. O'Brien, M. E. R. *et al.* Reduced cardiotoxicity and comparable efficacy in a phase III trial of pegylated liposomal doxorubicin HCl (CAELYX/Doxil) versus conventional doxorubicin for first-line treatment of metastatic breast cancer. *Annals of oncology : official journal of the European Society for Medical Oncology* **15,** 440–449; 10.1093/annonc/mdh097 (2004).
5. Nikitin, M. P. *et al.* Enhancement of the blood-circulation time and performance of nanomedicines via the forced clearance of erythrocytes. *Nature biomedical engineering* **4,** 717–731; 10.1038/s41551-020-0581-2 (2020).
6. Shramova, E. *et al.* Dual Targeting of Cancer Cells with DARPin-Based Toxins for Overcoming Tumor Escape. *Cancers* **12**; 10.3390/cancers12103014 (2020).
7. Strebhardt, K. & Ullrich, A. Paul Ehrlich's magic bullet concept: 100 years of progress. *Nature reviews. Cancer* **8,** 473–480; 10.1038/nrc2394 (2008).
8. Jong, W. H. de & Borm, P. J. A. Drug delivery and nanoparticles:applications and hazards. *International journal of nanomedicine* **3,** 133–149; 10.2147/ijn.s596 (2008).
9. Anselmo, A. C. & Mitragotri, S. Nanoparticles in the clinic: An update. *Bioengineering & translational medicine* **4,** e10143; 10.1002/btm2.10143 (2019).
10. Chiu, H. I., Samad, N. A., Fang, L. & Lim, V. Cytotoxicity of targeted PLGA nanoparticles: a systematic review. *RSC advances* **11,** 9433–9449; 10.1039/D1RA00074H (2021).
11. Meric-Bernstam, F. *et al.* Advances in HER2-Targeted Therapy: Novel Agents and Opportunities Beyond Breast and Gastric Cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research* **25,** 2033–2041; 10.1158/1078-0432.CCR-18-2275 (2019).
12. Lim, Y. W. *et al.* Challenges and Complications of Poly(lactic-co-glycolic acid)-Based Long-Acting Drug Product Development. *Pharmaceutics* **14**; 10.3390/pharmaceutics14030614 (2022).
13. Shipunova, V. O. *et al.* Dual Regioselective Targeting the Same Receptor in Nanoparticle-Mediated Combination Immuno/Chemotherapy for Enhanced Image-Guided Cancer Treatment. *ACS nano* **14,** 12781–12795; 10.1021/acsnano.0c03421 (2020).
14. Vivek, R. *et al.* Multifunctional HER2-antibody conjugated polymeric nanocarrier-based drug delivery system for multi-drug-resistant breast cancer therapy. *ACS applied materials & interfaces* **6,** 6469–6480; 10.1021/am406012g (2014).
15. Deng, L. *et al.* A Laser-Activated Biocompatible Theranostic Nanoagent for Targeted Multimodal Imaging and Photothermal Therapy. *Theranostics* **7,** 4410–4423; 10.7150/thno.21283 (2017).
16. Zheng, D. *et al.* Her2-Targeted Multifunctional Nano-Theranostic Platform Mediates Tumor Microenvironment Remodeling and Immune Activation for Breast Cancer Treatment. *International journal of nanomedicine* **15,** 10007–10028; 10.2147/IJN.S271213 (2020).
17. Sarbadhikary, P., George, B. P. & Abrahamse, H. Recent Advances in Photosensitizers as Multifunctional Theranostic Agents for Imaging-Guided Photodynamic Therapy of Cancer. *Theranostics* **11,** 9054–9088; 10.7150/thno.62479 (2021).
18. Mallidi, S. *et al.* Beyond the Barriers of Light Penetration: Strategies, Perspectives and Possibilities for Photodynamic Therapy. *Theranostics* **6,** 2458–2487; 10.7150/thno.16183 (2016).
19. Lee, Y.-H. & Chang, D.-S. Fabrication, characterization, and biological evaluation of anti-HER2 indocyanine green-doxorubicin-encapsulated PEG-b-PLGA copolymeric nanoparticles for targeted photochemotherapy of breast cancer cells. *Scientific reports* **7,** 46688; 10.1038/srep46688 (2017).
20. Shipunova, V. O., Sogomonyan, A. S., Zelepukin, I. V., Nikitin, M. P. & Deyev, S. M. PLGA Nanoparticles Decorated with Anti-HER2 Affibody for Targeted Delivery and Photoinduced Cell Death. *Molecules (Basel, Switzerland)* **26**; 10.3390/molecules26133955 (2021).
21. Yang, Y. *et al.* Biodegradable Polymer Nanoparticles for Photodynamic Therapy by Bioluminescence Resonance Energy Transfer. *Biomacromolecules* **19,** 201–208; 10.1021/acs.biomac.7b01469 (2018).
22. Proshkina, G. M. *et al.* DARPin_9-29-Targeted Gold Nanorods Selectively Suppress HER2-Positive Tumor Growth in Mice. *Cancers* **13**; 10.3390/cancers13205235 (2021).
23. Shramova, E. I. *et al.* Genetically encoded BRET-activated photodynamic therapy for the treatment of deep-seated tumors. *Light, science & applications* **11,** 38; 10.1038/s41377-022-00729-4 (2022).
24. Shipunova, V. O., Nikitin, M. P., Nikitin, P. I. & Deyev, S. M. MPQ-cytometry: a magnetism-based method for quantification of nanoparticle-cell interactions. *Nanoscale* **8,** 12764–12772; 10.1039/C6NR03507H (2016).
25. Lin, C. W., Shulok, J. R., Kirley, S. D., Cincotta, L. & Foley, J. W. Lysosomal localization and mechanism of uptake of Nile blue photosensitizers in tumor cells. *Cancer research* **51,** 2710–2719 (1991).
26. Swider, E. *et al.* Customizing poly(lactic-co-glycolic acid) particles for biomedical applications. *Acta biomaterialia* **73,** 38–51; 10.1016/j.actbio.2018.04.006 (2018).
27. Wischke, C. & Schwendeman, S. P. Principles of encapsulating hydrophobic drugs in PLA/PLGA microparticles. *International journal of pharmaceutics* **364,** 298–327; 10.1016/j.ijpharm.2008.04.042 (2008).
28. Kotelnikova, P. A. *et al.* Synthesis of Magnetic Nanoparticles Stabilized by Magnetite-Binding Protein for Targeted Delivery to Cancer Cells. *Doklady. Biochemistry and biophysics* **481,** 198–200; 10.1134/S1607672918040051 (2018).

29. Shipunova, V. O. *et al.* Self-assembling nanoparticles biofunctionalized with magnetite-binding protein for the targeted delivery to HER2/neu overexpressing cancer cells. *Journal of Magnetism and Magnetic Materials* **469,** 450–455; 10.1016/j.jmmm.2018.09.015 (2019).

30. Shipunova, V. O. *et al.* Versatile Platform for Nanoparticle Surface Bioengineering Based on SiO2-Binding Peptide and Proteinaceous Barnase*Barstar Interface. *ACS applied materials & interfaces* **10,** 17437–17447; 10.1021/acsami.8b01627 (2018).

31. Aghayeva, U. F., Nikitin, M. P., Lukash, S. V. & Deyev, S. M. Denaturation-resistant bifunctional colloidal superstructures assembled via the proteinaceous barnase-barstar interface. *ACS nano* **7,** 950–961; 10.1021/nn302546v (2013).

32. Zakeri, B. *et al.* Peptide tag forming a rapid covalent bond to a protein, through engineering a bacterial adhesin. *Proceedings of the National Academy of Sciences of the United States of America* **109,** E690-7; 10.1073/pnas.1115485109 (2012).

33. Bae, Y. *et al.* Engineering Tunable Dual Functional Protein Cage Nanoparticles Using Bacterial Superglue. *Biomacromolecules* **19,** 2896–2904; 10.1021/acs.biomac.8b00457 (2018).

34. Komedchikova, E. N. *et al.* Two-Step Targeted Drug Delivery via Proteinaceous Barnase-Barstar Interface and Doxorubicin-Loaded Nano-PLGA Outperforms One-Step Strategy for Targeted Delivery to HER2-Overexpressing Cells. *Pharmaceutics* **15,** 52; 10.3390/pharmaceutics15010052 (2023).

35. Stepanov, A. V. *et al.* Switchable targeting of solid tumors by BsCAR T cells. *Proceedings of the National Academy of Sciences of the United States of America* **119,** e2210562119; 10.1073/pnas.2210562119 (2022).

36. Förster, G. J., Santos, E. B., Smith-Jones, P. M., Zanzonico, P. & Larson, S. M. Pretargeted radioimmunotherapy with a single-chain antibody/streptavidin construct and radiolabeled DOTA-biotin: strategies for reduction of the renal dose. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* **47,** 140–149 (2006).

37. Shipunova, V. O. & Deyev, S. M. Artificial Scaffold Polypeptides As an Efficient Tool for the Targeted Delivery of Nanostructures In Vitro and In Vivo. *Acta naturae* **14,** 54–72; 10.32607/actanaturae.11545 (2022).

38. Lü, J.-M. *et al.* Current advances in research and clinical applications of PLGA-based nanotechnology. *Expert review of molecular diagnostics* **9,** 325–341; 10.1586/erm.09.15 (2009).

39. Sadat Tabatabaei Mirakabad, F. *et al.* PLGA-based nanoparticles as cancer drug delivery systems. *Asian Pacific journal of cancer prevention : APJCP* **15,** 517–535; 10.7314/APJCP.2014.15.2.517 (2014).

40. Niwa, T., Takeuchi, H., Hino, T., Kunou, N. & Kawashima, Y. Preparations of biodegradable nanospheres of water-soluble and insoluble drugs with D,L-lactide/glycolide copolymer by a novel spontaneous emulsification solvent diffusion method, and the drug release behavior. *Journal of Controlled Release* **25,** 89–98; 10.1016/0168-3659(93)90097-O (1993).

41. Kovalenko, V. L. *et al.* Lectin-Modified Magnetic Nano-PLGA for Photodynamic Therapy In Vivo. *Pharmaceutics* **15,** 92; 10.3390/pharmaceutics15010092 (2023).

42. Shipunova, V. O. *et al.* Targeting Cancer Cell Tight Junctions Enhances PLGA-Based Photothermal Sensitizers' Performance In Vitro and In Vivo. *Pharmaceutics* **14**; 10.3390/pharmaceutics14010043 (2021).

43. Martinez, V. & Henary, M. Nile Red and Nile Blue: Applications and Syntheses of Structural Analogues. *Chemistry (Weinheim an der Bergstrasse, Germany)* **22,** 13764–13782; 10.1002/chem.201601570 (2016).

44. Madsen, J. *et al.* Nile Blue-based nanosized pH sensors for simultaneous far-red and near-infrared live bioimaging. *Journal of the American Chemical Society* **135,** 14863–14870; 10.1021/ja407380t (2013).

45. LEWIS, M. R., GOLAND, P. P. & SLOVITER, H. A. The action of oxazine dyes on tumors in mice. *Cancer research* **9,** 736–740 (1949).

46. Nsubuga, A., Mandl, G. A. & Capobianco, J. A. Investigating the reactive oxygen species production of Rose Bengal and Merocyanine 540-loaded radioluminescent nanoparticles. *Nanoscale advances* **3,** 1375–1381; 10.1039/D0NA00964D (2021).

47. Doughty, M. J. Rose bengal staining as an assessment of ocular surface damage and recovery in dry eye disease-a review. *Contact lens & anterior eye : the journal of the British Contact Lens Association* **36,** 272–280; 10.1016/j.clae.2013.07.008 (2013).

48. Toshida, H. *et al.* Efficacy and safety of retinol palmitate ophthalmic solution in the treatment of dry eye: a Japanese Phase II clinical trial. *Drug design, development and therapy* **11,** 1871–1879; 10.2147/DDDT.S137825 (2017).

49. Mittal, N., Palaskar, S. & Shankari, M. Rose Bengal staining - diagnostic aid for potentially malignant and malignant disorders: a pilot study. *Indian journal of dental research : official publication of Indian Society for Dental Research* **23,** 561–564; 10.4103/0970-9290.107326 (2012).

50. Read, T. A. *et al.* Intralesional PV-10 for the treatment of in-transit melanoma metastases-Results of a prospective, non-randomized, single center study. *Journal of surgical oncology* **117,** 579–587; 10.1002/jso.24921 (2018).

51. Grosjean, P., Wagnieres, G., Fontolliet, C., van den Bergh, H. & Monnier, P. Clinical photodynamic therapy for superficial cancer in the oesophagus and the bronchi: 514 nm compared with 630 nm light irradiation after sensitization with Photofrin II. *British journal of cancer* **77,** 1989–1995; 10.1038/bjc.1998.330 (1998).

52. Fritsch, C. *et al.* Green light is effective and less painful than red light in photodynamic therapy of facial solar keratoses. *Photodermatology, photoimmunology & photomedicine* **13,** 181–185; 10.1111/j.1600-0781.1997.tb00226.x (1997).