Dynamical states of self-organized waves in a giant single-celled organism feeding on light

Eldad Afik,[1,2]* Toni J.B. Liu,[1] Elliot M. Meyerowitz[1,2]*

[1]Division of Biology and Biological Engineering, California Institute of Technology,
1200 E California Blvd., Pasadena, CA 91125, USA
[2]Howard Hughes Medical Institute

*To whom correspondence should be addressed;
E-mail: eldad.afik@gmail.com, meyerow@caltech.edu

Living systems often appear to act upon potential outcomes, exhibiting anticipatory behavior. Here we study *Caulerpa brachypus*, a marine green alga consisting of differentiated organs resembling leaves, stems and roots. While an individual can exceed a meter in size, it is a single multinucleated giant cell. Our experimental work reveals self-organized waves of greenness — chloroplasts — that propagate throughout the alga and anticipate the day-night light cycle. Using dynamical systems analysis we show that these waves are coupled to a self-sustained oscillator, and demonstrate their entrainment to light. Under constant conditions we find that light intensity affects the natural period and can drive transition to temporal disorder. We find also that the rate of development of organs, and their shape and size, depend on light intensity and its temporal distribution. The findings suggest the waves and their environmental control as a link among biological oscillators, metabolism and morphogenesis.

Introduction

A free-falling ball follows a trajectory determined by external forces. When we aim to catch the ball, in contrast, our dynamics follows an intrinsic predictive model of the world, exhibiting anticipatory behavior (*1, 2, 3*); such behavior has been observed in living systems down to cellular levels (*4, 5, 6, 7*).

Time-keeping and synchronization with the external world, as well as within the organism, seem to be essential for homeostasis, the ability of living systems to maintain essential variables within physiological limits (*1, 8, 9*). This is one of the ways by which living organisms manifest self-organization (*10*), defined here as networks of processes which are self-stabilizing far from thermodynamic equilibrium.

Biological oscillators are found throughout the living world, constituting a mechanism for organismal synchronization. Their manifestations, such as pulse rate and pressure, are measures of homeostasis. Biological oscillators are amenable to analytic approaches from dynamical systems (*11, 9, 12*). Here we study active rhythmic transport in *Caulerpa*, a marine green alga which presents complex morphology while being a giant single cell. *Caulerpa* challenges central paradigms in developmental biology, as it exhibits pattern formation and morphogenesis without multicellularity (*13*). While tracking morphogenesis of regenerating algal segments we observed that the growing tips of stem-like (stolon) and leaf-like (frond) regions exhibit temporal variations in intensity of their green color. During night distal regions typically appear more transparent, while during day the green coverage is more homogeneous. This has been attributed to long-distance chloroplast migration (*14*). Whether rhythmic transport in this single cell is light induced, or is of autonomous nature, is unknown.

Our findings show that the waves of greenness exhibit anticipatory behavior, usually starting toward the daylight state before dawn, and toward the night state before dark. Moreover, the analysis shows that anticipatory behavior in this system is explained by entrainment — adjustment of the rhythm of an organismal self-sustained oscillator by interaction with another oscillator (*15*), the driving illumination in this case. Using a scalable and affordable methodology, varying two control parameters — the temporal period of the illumination and its intensity — we identify distinct dynamical states of the self-organized waves. Our experimental results demonstrate that the waves propagating throughout the organism are coupled to, or part of, a self-sustained oscillator, and are entrained by time-dependent illumination. In constant illumination, the intrinsic period is intensity dependent, and under higher illumination intensities analysis indicates a transition to states of increased temporal disorder.

Results

To study quantitatively the self-organized green wave dynamics in *Caulerpa brachypus*, we developed a set of protocols for algal culture and propagation, as well as automated illumination, live imaging and image analysis. Using Raspberry Pi-controlled illumination and cameras, we track over weeks the morphogenesis of tens of samples concurrently, while tracing at resolution of tens of seconds the variation of the green coverage. A diagram of the experimental flow is shown in Fig. 1A; snapshots exemplifying the relevant time scales are presented in Fig. 1B–D. Our observations of samples cultured under cycles of 12 hours light followed by 12 hours dark (12hL-12hD, T=24h)[1] have allowed analysis of waves of greenness that propagate over

---

[1]*Light* and *Dark* denote reference illumination intensities, the latter is $1/200 \cdot I_L$, for $I_L \approx 4.5 \mu\mathrm{mol} \cdot \mathrm{s}^{-1} \cdot \mathrm{m}^{-2}$

centimeters, within a few hours, at a whole-organism scale; a time-lapse video is presented in movie S1. By coarse-graining in space we achieve a reduced description to a dynamic macroscopic observable. The time-series indicates that the initiation of the waves anticipates the external change in illumination, as exemplified in Fig. 1E. This has led us to ask whether the period of the waves follows that of the driving illumination.

To quantify the temporal frequency content of the biological response, we apply power spectral analysis (*16, 17*). While the time series of individual samples exhibit high variability, the spectral decomposition using power spectra captures and highlights the similarities among them. Examples from three samples and their corresponding power spectra are plotted in Fig. 2A and B, respectively. The pronounced local maxima in the power spectra correspond to the fundamental frequency $f_{r,0}$, centered at 1/24h for these samples, and its higher harmonics — integer multiples of $f_{r,0}$. The relative power of the higher harmonics characterizes the non-sinusoidal waveforms shown in Fig. 2A.

Under driving periods within 18h to 30h the response of the green pulses $f_{r,0}$ matches the driving frequency f_d in 1:1 correspondence, as shown in Fig. 2D. Moreover, the power spectra suggest an equivalence of the dynamics when rescaled by the control parameter f_d. That is, presenting the power spectra from various f_d as a function of f_r / f_d reveals the similarity in dynamics in this range; see Fig. 2E. The corresponding curves prior to rescaling are shown in Fig. 2C.

Subject to driving periods away from the 18h to 30h range, we find in addition to the driving frequency evidence for a distinct circadian mode — an indication that the waves are coupled to, or part of, an intrinsic self-sustained oscillator. For example, under driving period T=48h time series exhibit a steady phase relation to the illumination pattern; at the same time, at mid-Light intervals the dynamics resemble the day to night transitions observed under T=24h; see Fig. 3A. Under a driving period T=3h, eight times faster than earth's rotation, the time series show fast undulations attributed to a response to the driving illumination; these ride over an undulation of a longer period, as shown in Fig. 3B.

Entrainment of a nonlinear oscillator is predicted to occur at driving frequencies near rational ratios of its natural frequency and the driving one, in which case *m* cycles of the response would match *n* cycles of the driving oscillator (*15*). Within an *n:m*-entrainment range, the response fundamental frequency $f_{r,0}$ and its higher harmonics are predicted to be modified accordingly. To test for entrainment, we have measured samples subject to f_d about rational ratios of 24h, namely T=48h, 6h, 3h, and 1.5h, corresponding to 2^1, 2^{-1}, 2^{-3} and 2^{-4} multiples of 24h, as well as longer ones T=94h and 54h. The results support a 1:1-entrainment range, as well as evidence for other *n:m*-entrainment regions. The corresponding power spectra are presented as heatmaps, where the columns correspond to f_d, including those from f_d near 1/24h; see Fig. 3C [2].

Finally, we test whether alternating environmental conditions are necessary to sustain the waves, to complete the argument for an intrinsic self-sustained oscillator, and measure its natu-

---

[2]visualization inspired by (*18*)

Figure 1: **Self-organized macroscopic green waves: experimental design and coarse-graining.** (**A**) Schematic of the Cut & Regenerate experimental cycle. A *Caulerpa* cell is cut and segments are let to regenerate in individual wells; custom-built controllers apply the individually assigned illumination protocols per dish and control the time-lapse imaging; samples regenerating over weeks provide tens of segments for the next experiment; thus the experiments provide living material for the next cycle of experiments. (**B**) Snapshots from morphogenesis dynamics over the course of 35 days. (**C**) A developmental trajectory of a frond, focusing on a sub-interval of 13 days; these correspond to the frond marked by red rectangles in (**B**). (**D**) A temporal sub-interval spanning 48h, showing two periods of the green waves; the gray-scale frames represent the intensity in the Blue channel from the RGB images. (**E**) Coarse-graining of the spatio-temporal dynamics achieved by counting pixels whose intensity in the Blue channel falls below a preset threshold; the white and gray shadings correspond to 12h intervals of Light and Dark illumination levels, respectively.

Figure 2: **Dynamical equivalence under driving periods near 24h.** (**A**) Time series from three samples regenerating under 12hL-12hD (T=24h); area fraction corresponds to the measured green area within a whole well, normalized by the area of the well. (**B**) Power spectra corresponding to the time series in (A); the arrows annotate the 1st,2nd,3rd and 10th local maxima, corresponding to the response fundamental frequency $f_{r,0}$ and higher harmonics, namely integer multiples of $f_{r,0}$. (**C**) Power spectra of samples regenerating under illumination periods within 18h to 30h; the curves result from averaging over samples grouped by driving illumination frequencies, $f_d = 1/T$. (**D**) Response fundamental frequencies $f_{r,0}$ plotted as function of the driving frequency $f_d = 1/T$; the guiding dashed line represents a 1:1 correspondence. (**E**) Power spectra presented as a function of f_r/f_d , the response frequency rescaled by the driving frequency.

Figure 3: **The waves are coupled to, or part of, an intrinsic self-sustained oscillator, in addition to the driving illumination for periods far from 24h.** (**A**) Time-series from three samples regenerating under driving illumination of T=24h (12hL-12hD) switching to T=48h (24hL-24hD); while each is distinct, the curves reveal a steady phase relation to the illumination pattern. (**B**) Time series from three samples regenerating under T=24h (12hL-12hD) switching to T=3h (1.5hL-1.5hD); note that the fast undulations, which follow the driving illumination, are riding on a longer period one, close to 24h. (**C**) Power spectra presented as heatmaps; each column results from averaging over samples, grouped by driving illumination frequencies, $f_d = 1/T$; the guiding dashed line represents a 1:1 correspondence; the columns near f_d of 1/24h overlap with those in Fig. 2C, presented here to demonstrate 1:1-entrainment range; away from this region, in addition to the spectral content associated with the driving frequency f_d, local maxima are found near 1/24h (gray horizontal line), an indication of an intrinsic oscillator of circadian frequency.

ral period. To this end, we let samples regenerate under constant illumination. While we indeed find self-sustained circadian oscillations, subject to constant illumination the observations indicate new dynamical states. Examples of time series regenerating under low driving intensity $I_d = 2^{-5} \cdot I_L$ show a reduced amplitude of the oscillations, yet these persist over weeks; see Fig. 4A. Samples subject to high driving intensity $I_d = 2 \cdot I_L$ show signs of intermittent oscillations, presented in Fig. 4B. Power spectral analysis shows that: (i) the response fundamental frequency increases with decreasing illumination intensity, see Fig. 4C; (ii) under higher illumination intensities the local maxima in the power spectra 'drown' in a rising continuum of frequencies, and (iii) compared with the entrained states, the higher harmonic content is less pronounced; these findings are presented in Fig. 4D. Thus, under constant photon flux, the energy source for this organism, we find that the system can be driven towards aperiodic dynamical states by high fluxes. This is a manifestation of the nonlinear nature of the dynamics, and a plausible signature of chaotic dynamics.

How does the photon flux affect the developmental dynamics in this organism? It is expected that the energy injection rate would be a limiting factor to growth. Inferring the apparent area of samples from our macroscopic dynamical variable, we find that both the central tendency and the dispersion of growth rates increase with the illumination intensity; that is, algal samples regenerating under high photon flux cover larger areas faster as a population, while being less predictable individually, compared with those regenerating under a low photon flux. Moving averages, as estimates of apparent sample area, are presented in Fig. 4E. However, it turns out that the temporal pattern of the illumination leads to distinguishable morphological traits, even when the mean photon flux is kept the same. For example, we compare two mean photon flux levels, $2^{-1} \cdot I_L$ and I_L, each at constant illumination and T=24h, presented in Fig. 5. The comparison hints that the temporal distribution of photons, analogous to time-restricted feeding, impacts development as a control parameter which is independent of the mean photon flux.

Discussion

Our study reveals self-organized macroscopic waves in a unicellular organism. Using coarse-graining analysis and experimental perturbations we have identified distinct dynamical states of the waves. This report provides evidence for coupling to an intrinsic self-sustained oscillator, which is entrained by the time-dependent driving illumination, thus explaining the anticipatory behavior of the waves. Moreover, we find that under constant conditions the natural frequency depends on the photon flux, and the dynamics exhibits a transition to temporal disorder.

Caulerpa feeds on photons, so metabolism is likely to be a key to understanding the observed entertainment by light, via photosynthesis. This is the case in the KaiABC system — a molecular circadian oscillator of certain cyanobacteria (*19, 20*). Moreover, considering photosynthesis sheds new light on our identification of distinct dynamical states, subject to the two control parameters — illumination intensity and period. Constant illumination by itself does not drive the system to temporal disorder. Two of the driving protocols we have studied here

Figure 4: **Free-running at constant-in-time illumination reveals intensity dependent dynamical states.** (**A**) Time series from three samples regenerating under driving illumination of T=24h switching to constant intensity $I_d = 2^{-5} \cdot I_L$. (**B**) Time-series from three samples regenerating under driving illumination of T=24h switching to constant-in-time intensity $I_d = 2 \cdot I_L$. (**C**) Response fundamental frequencies $f_{r,0}$ plotted as function of the driving intensity, I_d ; the data indicate an intensity dependence of the free-running fundamental frequency. (**D**) Power spectra inferred from samples regenerating under driving illumination intensities within $2^{-5} \cdot I_L$-to-$2^{+1} \cdot I_L$; the curves result from averaging over samples grouped by driving illumination intensities, I_d; under higher driving intensities the data reveal an increase in power distribution across a continuum of frequencies. (**E**) Moving averages of the time-series for samples regenerating under constant illumination; in addition to intensity-dependent growth rates, the data reveal a sample dispersion which increases with the driving intensity.

Figure 5: **Developmental effect of photon flux intensity and temporal distribution.** Snapshots from four driving illumination conditions, comparing mean photon flux levels $2^0 \cdot I_L$ and $2^{-1} \cdot I_L$, at T=24h and constant illumination. Samples regenerating under time dependent illumination at T=24h share qualitative morphological features, which are distinguishable from those under constant illumination, for example, the typical frond size and aspect ratio. The panels are cropped to show three wells out of eight from each culture dish; following 13 days subject to T=24h, $2^{-1} \cdot I_L$ (12hL-12hD) the new driving illumination was applied; panels correspond to 25 days after this transition.

are equivalent in their average photon flux, namely 12hL-12hD and constant $I_d = 2^{-1} \cdot I_L$. The latter leads to chaotic dynamics, hinting that relaxation in the dark is essential for the typical organismal dynamics.

How do tissues and organs emerge on centimeter scales in the absence of cellularization? Active transport has been hypothesized to play a key role in *Caulerpa* regeneration ([21, 13]). The observation of the self-organized waves leads us to postulate that their synchronization with light-driven metabolic switching may hold the answer. This hypothesis assumes two key elements: (i) spatial distribution of chloroplasts would result in local variation in light-harvesting efficiency, and (ii) light-dependent metabolic states of chloroplasts. Under these assumptions, redistribution of chloroplasts in anticipation of the temporal changes in photon flux could create and stabilize metabolic sub-environments, akin to localized organs. An additional contribution of the waves to the dynamics may come in the form of an effective pump: cortical mass migration of chloroplasts could result in bulk flow, which in turn drives cytoplasmic streaming; dynamics of this kind has been demonstrated in other macroscopic cells ([22]).

Ashby ([10]) has proposed a paradigm for self-organization, where a subset of variables in a dynamical system can undergo abrupt transitions between two values. Such transitions would result in dynamical switching among distinct behaviors of the whole system. In this light we propose that biological oscillators, morphogenesis, and metabolism are sub-systems in a network of processes, whose dynamics and inter-relations give rise to self-stabilization far from thermodynamic equilibrium in this natural system which is alive — a generalized homeostasis.

References

1. W. Ross Ashby. *Design for a Brain: The origin of adaptive behaviour*. Springer Netherlands, Dordrecht, 1960.

2. Robert Rosen. On anticipatory systems: I. When can a system contain a predictive model of another? *Journal of Social and Biological Structures*, 1(2):155–162, April 1978.

3. Naftali Tishby and Daniel Polani. Information Theory of Decisions and Actions. *Perception-Action Cycle*, pages 601–636, 2011.

4. Eric D. Siggia and Massimo Vergassola. Decisions on the fly in cellular sensory systems. *Proceedings of the National Academy of Sciences*, 110(39):E3704–E3712, September 2013. Publisher: National Academy of Sciences Section: PNAS Plus.

5. Antonio Celani, Emmanuel Villermaux, and Massimo Vergassola. Odor Landscapes in Turbulent Environments. *Physical Review X*, 4(4):041015, October 2014. Publisher: American Physical Society.

6. Kabir Husain, Weerapat Pittayakanchit, Gopal Pattanayak, Michael J. Rust, and Arvind Murugan. Kalman-like Self-Tuned Sensitivity in Biophysical Sensing. *Cell Systems*, 9(5):459–465.e6, November 2019.

7. Monika Scholz, Aaron R. Dinner, Erel Levine, and David Biron. Stochastic feeding dynamics arise from the need for information and energy. *Proceedings of the National Academy of Sciences*, 114(35):9261–9266, August 2017. Publisher: Proceedings of the National Academy of Sciences.

8. Walter B. Cannon. Organization for physiological homeostasis. *Physiological Reviews*, 9(3):399–431, July 1929. Publisher: American Physiological Society.

9. Leon Glass. Synchronization and rhythmic processes in physiology. *Nature*, 410(6825):277–284, March 2001. Number: 6825 Publisher: Nature Publishing Group.

10. W. Ross Ashby. Principles of the Self-Organizing Dynamic System. *The Journal of General Psychology*, 37(2):125–128, October 1947.

11. Arthur T. Winfree. *The Geometry of Biological Time*. Interdisciplinary Applied Mathematics. Springer, New York, NY, 2001.

12. Steven H Strogatz. *Sync: How order emerges from chaos in the universe, nature, and daily life*. Hachette UK, 2012.

13. William P. Jacobs. Caulerpa. *Scientific American*, 271(6):100–105, December 1994.

14. Clinton J Dawes and D Craig Barilotti. Cytoplasmic organization and rhythmic streaming in growing blades of Caulerpa prolifera. *American Journal of Botany*, pages 8–15, 1969. https://doi.org/10.1002/j.1537-2197.1969.tb07499.x.

15. Arkady Pikovsky, Michael Rosenblum, and Jurgen Kurths. *Synchronization: A universal concept in nonlinear sciences*. Number 12 in Cambridge Nonlinear Science Series. Cambridge University Press, 2001.

16. Josef Honerkamp. *Statistical Physics*. Graduate Texts in Physics. Springer Berlin Heidelberg, Berlin, Heidelberg, 2012.

17. Adrián E. Granada, Trinitat Cambras, Antoni Díez-Noguera, and Hanspeter Herzel. Circadian desynchronization. *Interface Focus*, 1(1):153–166, February 2011. Publisher: Royal Society.

18. Benjamín Herrmann, Philipp Oswald, Richard Semaan, and Steven L. Brunton. Modeling synchronization in forced turbulent oscillator flows. *Communications Physics*, 3(1):1–9, October 2020. Number: 1 Publisher: Nature Publishing Group.

19. Michael J. Rust, Susan S. Golden, and Erin K. O'Shea. Light-Driven Changes in Energy Metabolism Directly Entrain the Cyanobacterial Circadian Oscillator. *Science*, 331(6014):220–223, January 2011. Publisher: American Association for the Advancement of Science Section: Report.

20. Yi Liao and Michael J. Rust. The circadian clock ensures successful DNA replication in cyanobacteria. *Proceedings of the National Academy of Sciences*, 118(20), May 2021. Publisher: National Academy of Sciences Section: Biological Sciences.

21. J. M. Janse. An investigation on polarity and organ-formation with Caulerpa prolifera. *Koninklijke Nederlandse Akademie van Wetenschappen Proceedings Series B Physical Sciences*, 7:420–435, 1904.

22. F. G. Woodhouse and R. E. Goldstein. Cytoplasmic streaming in plant cells emerges naturally by microfilament self-organization. *Proceedings of the National Academy of Sciences*, 110(35):14132–14137, August 2013.

23. James C. W. Chen and W. P. Jacobs. Quantitative Study of Development of the Giant Coenocyte, Caulerpa prolifera. *American Journal of Botany*, 53(5):413–423, 1966.

24. DD Sabnis and WP Jacobs. Cytoplasmic streaming and microtubules in the coenocytic marine alga, Caulerpa prolifera. *Journal of cell science*, 2(4):465–472, 1967.

25. Clinton J. Dawes and Ernest L. Rkamstine. An ultrastructural study of the giant green algal coenocyte, Caulerpa prolifera. *Journal of Phycology*, 3(3):117–126, September 1967.

26. Fernando Pérez, Brian E. Granger, and John D. Hunter. Python: An Ecosystem for Scientific Computing. *Computing in Science & Engineering*, 13(2):13–21, 2011.

27. Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C. J. Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, and Paul van Mulbregt. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods*, 17(3):261–272, March 2020. Number: 3 Publisher: Nature Publishing Group.

28. Wes McKinney. Data Structures for Statistical Computing in Python. In Stéfan van der Walt and Jarrod Millman, editors, *Proceedings of the 9th Python in Science Conference*, pages 51 – 56, 2010.

29. Fernando Pérez and Brian E. Granger. IPython: A System for Interactive Scientific Computing. *Comput. Sci. Eng.*, 9(3):21–29, 2007.

Acknowledgments

We thank A. Kornfeld for the computer-aided design of the 3d-printed apparatus, as well as Caltech Library Techlab and staff for 3D Printers access, materials and support. We are grateful to L. Michaeli, A.I. Flamholz, and G. Manella for helpful discussions and critical comments on the manuscript. E.A. is thankful for fruitful discussions at the SRBR2022 and EBRS2022 meetings.

Funding: The laboratory of E.M.M. is supported by the Howard Hughes Medical Institute. E.A. has been awarded the Zuckerman Israeli Postdoctoral Scholar, Zuckerman STEM Leadership Program, and the Biology and Biological Engineering Divisional Fellowship, Caltech. T.J.B.L. has been awarded the Summer Undergraduate Research Fellowship (SURF), Caltech.

Author contributions: Conceptualization: E.M.M proposed studying morphogenesis in *Caulerpa*; E.A. designed the study; Methodology: E.A. designed the experimental system and analysis; Investigation: E.A. performed the measurements; E.A. and T.J.B.L. performed computational analysis; Visualization: E.A. and T.J.B.L.; Funding acquisition: E.M.M.; Writing – original draft: E.A.; Writing – review & editing: E.A. and E.M.M.; All authors discussed and commented on the manuscript.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: Analyzed data are available in the main text. Raw data, code and materials are available upon request from E.A. and E.M.M. All programming and computer aided analysis has been done using open-source projects, primarily tools from the Scientific Python ecosystem. Distributed image processing was conducted in the Resnick High Performance Computing Center, a facility supported by Resnick Sustainability Institute at the California Institute of Technology.

Supplementary materials

Materials and Methods
Movie S1

Supplementary materials

Materials and Methods

To quantitatively study the green wave dynamics in *Caulerpa brachypus*, we have developed an affordable and scalable experimental system for the algal culture, propagation and imaging, summarized in Fig. 1A. A Raspberry-Pi-Zero-W is mounted on a custom designed 3d-printed apparatus, such that its camera module is focused at an 8-well-culture-dish. Using multiple replicas, we concurrently track over weeks the morphogenesis of tens of samples, while tracing at resolution of tens of seconds the variation of the green coverage. The same computers additionally control LED strips, assigning distinct time-dependent illumination per culture dish that provides both for photosynthesis and imaging. Samples regenerating over weeks can provide tens of segments for the next experiment; hence the experiments provide living material for their follow-ups in a cyclic manner. A time-lapse video of samples regenerating under 12hL-12hD, 24h Light-Dark cycles, is presented in movie S1.

Towards reproducibility, controlling for the variability in initial conditions of the living samples, we use the regenerative properties of *Caulerpa brachypus*. Individual algal samples are cut in segments of 1-3 centimeters in length; within 1 to 2 months a regenerating sample can provide for about 20 new segments. This approach contributes to the homogeneity of the observed individuals, both in their genetic background, as well as the developmental state and geometry set as initial conditions. A visual summary of the experimental setup is presented in Fig. 1A.

The nature of the data presents both conceptual and computational challenges. Conceptually it is desired to identify a coarse-graining procedure, to reduce the high-dimensionality of the spatio-temporal dynamics of the green waves, and at the same time preserve a minimal set of essential degrees of freedom. Images are taken every 150 seconds over weeks. This amounts to datasets of about 400 GiB/month/dish of RGB (Red-Green-Blue) images. Running several experiments concurrently leads to high-throughput data generation. It is therefore challenging for human detailed inspection, introducing computational challenges: for computer vision and for computational resources.

In this report, our dynamic observable is the apparent green area. Its time series are inferred from applying a threshold to the blue channel of the RGB images. Chlorophyll absorbs strongly in the blue. Therefore regions rich in Chlorophyll are expected to show relative darker values in the Blue channel. An example is summarized in Fig. 1B–E, where the time series correspond to a region of interest tracking a single leaf-like organ. Everywhere else in this report the time series have been inferred from regions of interest corresponding to wells; hence these represent the apparent fraction of green area, where the area of the well is the normalization factor.

Caulerpa brachypus culture and propagation

Caulerpa brachypus algal samples in this report were all derived from a single origin (shipped from Inland Aquatics, Indiana).

As culture medium we use ErdsSchreiber's Medium (UTEX Culture Collection of Algae at The University of Texas at Austin). It is a variation over the one in previous reports on *Caulerpa* (23, 24, 25).

Prior to cutting, samples were triple washed using filtered artificial sea water (35‰ Instant Ocean in deionized water; filtered using a Nalgene™ Rapid-Flow™ with 0.1μm PES Membrane filters, Thermo Fisher Scientific Inc.). Algal segments were cut using a surgical blade, and let to regenerate in 8-well-dishes (Non-treated Nunc™ Rectangular Dishes, Thermo Fisher Scientific Inc.), each well filled 7ml of the culture medium.

Illumination and imaging setup

Custom made apparati were 3D-printed using translucent neutral color Polylactic Acid (PLA and 3D Printers access courtesy of Caltech Library Techlab). These were designed to hold the culture dish, as well as the LEDs, Raspberry-Pi and camera module.

Raspberry-Pi-Zero-W and its 8MP Camera v2 module (Adafruit Industries) were used for time-lapse image acquisition (using the RPi-Cam-Web-Interface). Exposure time and gain were automatically set by the software to allow consistent imaging at various illumination intensities. The Raspberry-Pi also controlled the illumination protocols, allowing the assignment of time-dependent photon-flux per culture dish. This was done by pulse-width modulation (using the Python interface of the pigpio library) of LED strips (5000K cool-white SMD5050 12V, L1012V-502-1630 from HitLights), two triplets per dish. Reference illumination intensity is estimated at $I_L \approx 4.5\mu\text{mol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$, measured at the culture medium level covered by the culture dish lid (photodiode power sensor connected to an optical power meter, S120C and PM100D from Thorlabs). Under these conditions, the measured illumination spectrum shows two main modes, one at 545nm, half-max extends 505nm-614nm, and the other at 440nm, half-max extends 428nm-449nm (optic spectrometer USB2000+ from OceanOptics).

The experiments were conducted in incubators (DT2-MP-47L from Tritech Research, Inc., and VWR 2015 from Sheldon Manufacturing Inc.) at temperatures maintained within 22.5°C to 24.5°C.

Spatial coarse-graining, post-processing and inference of power spectra

Images have been split into regions of interest, each corresponding to a well in the 8-well-dish. For each region of interest, time series correspond to the fraction of pixels whose value does not exceed a threshold of 70 out of 255 in the blue channel.

Post-processing of the time series consists of outlier detection and replacement. Outliers have been defined as those exceeding 75% of the well, or lying outside the 1.5 x Interquartile-Range of a rolling window spanning 1h40m. Where there are no more than three consecutive outliers, these have been replaced by linear interpolation.

Power spectra have been inferred based on 35 day intervals, starting two days after transitioning from 12hL-12hD to the new driving illumination protocol. Following the above post-

processing procedure for outlier detection and replacement, time series consisting of more than 2% unreplaced outliers within these intervals were excluded from this report.

Power spectra have been estimated by Welch's method (*16*), applying Hann taper to segments of 8.5 days (4896 time-points), 50% overlap between segments, zero-padded for interpolated frequency resolution of $1/18 \times (24h)^{-1}$, without detrending. Any remaining detected outliers were replaced by forward and backwards propagation of nearest valid data.

To estimate dominant frequencies, local maxima in the power spectra are detected by a two step process: first, local peaks are filtered by requiring a prominence of at least twice the level of their baseline; second, these are refined by a local fit to a parabola on logarithmic scale, applied to 5 points centered at the peaks detected by the first step.

The trends in Fig. 4E have been estimated by a rolling average, using gaussian weighting window of σ corresponding to 2 days 00h55m45s, and spanning 4σ.

All programming and computer aided analysis has been done using open-source projects, primarily tools from the Scientific Python ecosystem (*26*): SciPy (*27*), pandas (*28*), IPython and JupyterLab (*29*); image processing has been distributed using dask and xarray; visualization has been done using HoloViz and Matplotlib.

Movie S1. Time-lapse imaging of algal samples regenerating under 12hL-12hD, T=24h cycles. One month of regeneration under 12h Light - 12h Dark; 3 seconds correspond to 24h; 3 wells, 3cm x 4cm each. The waves of greenness show synchronization within individual cells, and among them.