IMD-mediated innate immune priming increases Drosophila survival and reduces pathogen transmission

Arun Prakash[1,4]*, Florence Fenner[1], Biswajit Shit[2], Tiina S. Salminen[3], Katy M. Monteith[1], Imroze Khan[2], Pedro F. Vale[1]*

[1]Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh, UK

[2]Ashoka University, Plot No. 2, Rajiv Gandhi Education City, P.O. Rai, Sonepat, Haryana-131029, India

[3]Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

[4]Current address: Vanderbilt University, 465 21st Ave S, MRB-III, Biological Sciences, Nashville, TN 37212, USA

* Pedro F. Vale,

Email: pedro.vale@ed.ac.uk

Author Contributions: AP and PFV conceived and designed the experiments. AP, FF, BS, KMM, TSS performed the experiments. AP analysed the data with help from FF and PFV. AP and PFV wrote the paper with additional inputs from IK, TSS, FF, BS & KMM.

Competing Interest Statement: The authors declare no competing interests.

Keywords: antimicrobial peptides, immune priming, immune regulation, pathogen transmission, pathogen shedding.

Abstract

Invertebrates lack the immune machinery underlying vertebrate-like acquired immunity. However, in many insects past infection by the same pathogen can 'prime' the immune response, resulting in improved survival upon reinfection. Here, we investigated the generality, specificity and mechanistic basis of innate immune priming in the fruit fly *Drosophila melanogaster* when infected with the gram-negative bacterial pathogen *Providencia rettgeri*. We find that priming in response to *P. rettgeri* infection is a long-lasting and pathogen-specific response. We further explore the epidemiological consequences of immune priming and find it has the potential to curtail pathogen transmission by reducing pathogen shedding and spread. The enhanced survival of individuals previously exposed to a non-lethal bacterial inoculum coincided with a transient decrease in bacterial loads, and we provide strong evidence that the effect of priming requires the IMD-responsive antimicrobial-peptide *Diptericin-B* in the fat body. Further, we show that while *Diptericin B* is the main effector of bacterial clearance, it is not sufficient for immune priming, which requires regulation of IMD by peptidoglycan recognition proteins. This work underscores the plasticity and complexity of invertebrate responses to infection, providing novel experimental evidence for the effects of innate immune priming on population-level epidemiological outcomes.

Introduction

Immunisation using attenuated or inactivated pathogens is one of the most successful public health practices to reduce the incidence of infectious diseases(Pollard and Bijker, 2021). Immunisation works because humans and other vertebrate animals have evolved an acquired immune response capable of specific immune memory, which ensures a strong, precise, and effective response to a secondary infection(Boehm and Swann, 2014). Insects possess a robust innate immune response to pathogens which includes both cellular and humoral components (Hoffmann, 2003; Hultmark, 2003; Buchon et al., 2014), but lack vertebrate-like specialized immune cells responsible for acquired immunity. These differences in immune physiology resulted in the long-standing assumption that invertebrates should not be capable of immune 'memory', though this view was clearly at odds with empirical evidence from several invertebrate host-pathogen systems(Lanz-Mendoza et al., 2024; Milutinović and Kurtz, 2016; Prakash and Khan, 2022). There is now substantial evidence that insects also possess a form of "immune priming", where low doses of an infectious pathogen, or even an inactivated pathogen, can lead to increased survival upon reinfection (Kurtz and Franz, 2003; Sadd and Schmid-Hempel, 2006; Lanz-Mendoza et al., 2024; Contreras-Garduño et al., 2016; Milutinović and Kurtz, 2016).

There are many ways in which invertebrates may enhance their immune responses upon reinfection(Arch et al., 2022; Gomes et al., 2022; Prakash and Khan, 2022). In *Drosophila,* the priming response during infection with the gram-positive bacterial pathogen *Streptococcus pneumoniae* was shown to be dependent on haemocytes and phagocytosis, while the Toll-pathway - the main pathway involved in clearance of gram-positive bacteria - was shown to be insufficient for successful priming (Pham et al., 2007). Increased phagocytic activity in primed individuals has also been shown to play a key role in priming during *Pseudomonas aeruginosa* infection in Drosophila(Apidianakis et al., 2005), while in the woodlouse, prior exposure to heat-killed bacteria led to increased phagocytosis by haemocytes upon reinfection (Roth and Kurtz, 2009). In other *Drosophila* work, *PGRP-LB* (peptidoglycan recognition protein *LB*, a negative regulator of the Immune deficiency (IMD)-pathway) was identified as a key mediator of transgenerational immune priming against infection with parasitoid wasps (*Leptopilina heterotoma* and *Leptopilina victoriae*). Here, downregulation of *PGRP-LB* was necessary to increase haemocyte proliferation, required for wasp encapsulation by haemocytes in the offspring (Bozler et al., 2020). By contrast, in response to infection with *Drosophila C virus* (DCV)*, Drosophila* progeny can produce a DCV-specific priming response by inheriting viral cDNA from the infected adult flies which is a partial copy of the virus genome (Mondotte et al., 2018, 2020). The benefits of priming are not always associated with increased pathogen clearance, as shown during Enterococcus faecalis infection, where protection was explained by increases tolerance of infection, not increased clearance (Cabrera et al., 2023).

Beyond its underlying mechanisms, an important but understudied aspect of immune priming is its potential consequences for pathogen transmission (Gomes et al., 2022; Tate, 2017, 2016; Tidbury et al., 2012). Epidemiological modelling predicts that priming should affect the

likelihood of pathogen persistence, destabilise host–pathogen population dynamics, and that
these effects depend on the degree of protection conferred by priming (Tidbury et al., 2012).
Further theoretical work suggests that priming can either increase or decrease infection
prevalence depending on the extent to which it affects the pathogen's colonization success
and the host's ability to clear or tolerate the infection (Tate, 2017). While primed individuals
may live longer, thus extending the infectious period, their pathogen burden may be lower,
which could lead to lower pathogen shedding and less severe epidemics. Immune priming is
therefore expected to have a significant impact on the outcome of pathogen transmission by
directly modifying important epidemiological parameters, but the strength and direction of
these effects is not intuitive to predict.
100
Here, we focus on immune priming in *Drosophila* when infected with the gram-negative
bacterial pathogen *Providencia rettgeri* to investigate the occurrence, generality, duration,
and mechanistic basis of immune priming during systemic infection. Furthermore, motivated
by the theoretical predictions about the role of immune priming on epidemiological dynamics,
we also designed transmission experiments to enable us to test how immune priming could
affect each of these behavioural and immunological components of pathogen spread.
107
108

Results

Immune priming in *Drosophila* is a long-lasting response, showing sex-specific effects in different genetic backgrounds

We first examined if the length of time between the initial non-lethal exposure with heat-killed *P. rettgeri* and the secondary pathogenic challenge with live *P. rettgeri* affects the extent of priming. To address this, we exposed w^{1118} male and female control flies to live *P. rettgeri* 18-hours, 48-hours, 96-hours, 1-week or 2-weeks following the initial exposure to heat-killed bacteria. Male w^{1118} flies showed increased survival after initial priming for time points 18-hours, 48-hours and 96-hours, and still showed a significant, albeit reduced, priming response 1-week and 2-weeks after the initial exposure (**Fig**. 1 A-E). Female flies did not show a priming response when infected 18-hours following the initial challenge, but the priming response increased with 48-hours and 96-hours priming intervals before completely disappearing after a week time-interval (including 2-weeks) (**Fig**. 1A-E, **Table**-S1).

Figure 1: The benefit of priming is long-lasting. Survival curves of w^{1118} females and males with primed (exposed to heat-killed *P. rettgeri* in the first exposure) and unprimed (exposed to a sterile solution in the first exposure) treatments challenged with live *P. rettgeri* pathogen after (A) 18-hours (B) 48-hours and (C) 96-hours (D) 168-hours/1-week and (E) 336-hours/2-weeks post priming that is, initial non-lethal exposure to heat-killed *P. rettgeri* (n=7-9 vial with 10-15 flies in each vial/sex/treatment/timepoint).

Next, we asked if priming occurred in two other commonly used *Drosophila* lines, Canton-S
and Oregon-R (Ore-R) as observed with w^{1118}. Given the widespread effects of the
endosymbiont *Wolbachia* on *Drosophila* immunity (Brownlie and Johnson, 2009; Teixeira et
al., 2008; Vanika Gupta et al., 2017) we also tested whether the presence of *Wolbachia* had
any effect on immune priming by comparing the priming response of Oregon-R (OreR), the
line originally infected with *Wolbachia* strain wMel, henceforth OreR^{Wol+} and a *Wolbachia*-free
line OreR^{Wol-} that was derived from OreR^{Wol+} by antibiotic treatment (Vanika Gupta et al.,
2017). Females and males of the four *Drosophila* lines were treated first with heat-killed *P.
rettgeri*, followed by infection with live *P. rettgeri* 96-hours after the first treatment. Since the
96-hour time gap between priming and live *P. rettgeri* treatments showed maximum priming
response (difference in survival) for both males and females of the w^{1118} line (Fig. 1C), we
kept the 96-hours timepoint as the consistent time-gap between priming and live infection
throughout the study (**Fig**. 2C). Canton-S flies showed increased survival following priming
and we observed this survival benefit across both sexes, although stronger in females (**Fig**.
2B, **Table** S2B). Oregon-R males also exhibited increased survival following priming, but
priming had no significant effect on the females (**Fig**. 2C, **Table**-S2). We found that the
presence of *Wolbachia* significantly improved overall survival of both males and females
(**Fig**. 2C and D, **Table**-S3). However, the immune priming observed in males in absence of
Wolbachia (OreR^{Wol-}) was no longer present in flies carrying *Wolbachia* (OreR^{Wol+}) (**Fig.** 2D,
Table-S3).

**Figure 2: Survival curves of females and males of the commonly used *Drosophila*
lines either primed-challenged or unprimed-challenged with *P. rettgeri*.** (A) w^{1118} (B)
Canton-S (C) Oregon-R carrying endosymbiont *Wolbachia* (D) Oregon-R cleared of
Wolbachia. (n=7-9 vial with 10-15 flies in each vial/sex/treatment/line).

The strength of immune priming is bacteria species-specific

A notable feature of immune priming is its specificity (Roth et al., 2009; Khan et al., 2017), which suggests an added level of functional complexity of insect immunity. We tested whether *D. melanogaster* also shows specific immune priming against *Providencia rettgeri (Pr)*. We used males of the w^{1118} line, as males were shown to have stronger priming response than females and two pathogens, one closely related species of *P. rettgeri* i.e., *Providencia burhodogranariea (Pb)* and another less closely related gram-negative bacteria i.e., *Pseudomonas entomophila (Pe)*. Male flies primed and challenged homologously (*Pr – Pr*) species showed significantly improved survival rates compared with the heterologous treatment combinations (primed with *Pr* followed by challenge with other bacteria) and corresponding control (primed with PBS – challenge with live bacteria) (**Fig**. 3A-C, **Table** S4A-B).

Figure 3: Bacterial species-specific immune priming response in w^{1118} males. Two closely related species of Providencia bacteria *Providencia rettgeri* (Pr) and *Providencia burhodogranariea* (Pb) and another gram-negative bacteria *Pseudomonas entomophila* were used for testing the immune priming specificity in *Drosophila*. (A) survival curves (B) hazard ratio of unprimed males with 1x PBS followed by live *P. rettgeri* against males with primed *P. rettgeri* followed by live infection with other bacterial species *Pb* and *Pe*. (C). hazard ratio of primed males with heat-killed *P. rettgeri* followed by live *P. rettgeri* against males with primed *P. rettgeri* followed by live infection with other bacterial species *Pb* and *Pe*.

Primed male w^{1118} flies exhibit a transient reduction in bacterial load

Previous studies have shown that the priming response to fungi and gram-positive bacteria can be explained by increased clearance of bacterial pathogens in primed individuals (Pham et al., 2007; Khan et al., 2019). We therefore examined whether the increased survival

following a prior challenge we observed was a result of greater bacterial clearance in the
primed individuals, or if the primed flies were simply better able to tolerate the bacterial
infection (Cabrera et al., 2023). To investigate this, we repeated the priming experiment with
w^{1118} as it showed the strongest priming phenotype in both sexes (**Fig**. 1C and **Fig**. 4A), and
measured the bacterial load at 24-hours and 72-hours post-infection for both primed and
unprimed female and male flies. Again, we observed a clear priming effect in both sexes
(**Fig**. 4A, **Table**-S5). In primed males the systemic infection resulted in higher survival and
also in decreased bacterial loads at 24-hours post-infection when compared to unprimed
individuals (**Fig**. 4, **Table**-S5 for survival, **Table**-S6 for load). However, by 72-hours post-
infection, bacterial loads had dropped in both primed and unprimed male flies and there was
no detectable effect of priming on the bacterial loads (**Fig**. 4B, **Table**-S6).
201

202
Figure 4: Bacterial loads of primed and unprimed w^{1118} flies. (A). Survival curves of
primed and unprimed control w^{1118} flies (B). Internal bacterial load (n=20-22 vial with 5-7
flies in each vial/sex/treatment) after (i) 24-hours and (ii) 72-hours following *P. rettgeri*
infection. Different letters in panel-B denotes primed and unprimed individuals are
significantly different, tested using Tukey's HSD pairwise comparisons, analysed separately
for each timepoint and sex combination. The error bars in panel B represent standard error.
209
210
Priming following oral infection reduces bacterial shedding and transmission in males
Apart from increasing survival during infections, immune priming may also have
consequences to pathogen spread and transmission (Tate, 2016). Despite having enhanced
survival, the primed individuals may also extend the infectious period or their pathogen

burden may be lower, which would lead to less severe epidemics (Tate, 2016). To
investigate how immune priming affects epidemiological parameters we tested whether
primed individuals have reduced bacterial shedding and spread. Given the oral-faecal nature
of bacterial transmission and that our previous results all related to systemic infections, we
first established that a survival benefit fo priming also occurs under oral infection. Following
an initial oral exposure to a heat-killed *P. rettgeri* culture, after a 72-hour period we exposed
female and male *w*[1118] flies orally to a lethal dose of live *P. rettgeri*. Primed *w*[1118] males, but
not females, showed increased survival decreased internal bacterial loads after priming via
the oral route of infection. (**Fig.** 5A and 5B, **Table**-S7 and S8).

225
Figure 5: Priming occurs in males but not females during oral infections. (A) Survival
curves of male and female wildtype (*w*[1118]) flies with (i) systemic (ii) oral priming where flies
initially exposed to heat-killed *P. rettgeri* or unprimed flies exposed to a sterile solution in the
first exposure, this is followed by live *P. rettgeri* pathogen after 72-hour time gap [systemic
infection dose = 0.75 OD (~45 cells/fly) and oral *P. rettgeri* infection dose = 25 OD_{600}]. (n= 7-
9 vials/sex/treatment/infection route) (B). Mean bacterial load measured as colony forming
units at 24 hours following (i) systemic and (ii) oral priming, followed by live *P. rettgeri*
infection for male and female *w*[1118]. Different letters in panel-B denotes primed and unprimed
individuals are significantly different, tested using Tukey's HSD pairwise comparisons.

235
236

Given this priming effect on bacterial loads (Fig 5B) we hypothesised that priming could
directly impact the amount of bacterial shedding, and therefore have a direct impact on
pathogen transmission. We measured the shedding of single flies 4-hours after infection by
live *P. rettgeri* exposure. We found that male flies previously primed with a heat-killed

bacterial inoculum shed less bacteria than unprimed flies. However, both primed and
unprimed females showed increased bacterial shedding following oral *P. rettgeri* infection
(**Fig.** 6A, **Table**-S10). These effects on bacterial shedding are likely to have a drirect effect
on the spread of infection in groups of individuals. In transmission assays, primed donor
males also spread very little pathogen upon re-infection with oral *P. rettgeri*, compared to the
unprimed male treatment where successful transmission was detected recipient flies (**Fig.**
6B, **Table-**S10). However, both primed and unprimed females showed equivalent levels of
bacterial spread following oral *P. rettgeri* infection (**Fig.** 6B, **Table-**S10).
249

250
Figure 6: The effects of priming on bacterial shedding and transmission. (A). Bacterial
shedding after 4-hours following oral priming (initial heat-killed exposure) and infection with
live *P. rettgeri* (OD_{600}=25) for males and females (n=15 individual flies per treatment). (B)
percentage transmission of measurable bacterial loads for male and female *w*[1118] flies
(recipient-flies) over the first 4-hours following exposure with infected donor flies (n=6
independent transmission assays). Different letters denote primed and unprimed individuals
are significantly different, tested using Tukey's HSD pairwise comparisons.

258
259
While multiple traits, including host activity levels and contact rates may influence pathogen transmission (Siva-Jothy and Vale, 2021; White et al., 2020), we did not find any difference in the locomotor activity of primed and unprimed flies (Fig 7). Our results therefore provide evidence that, in male flies, immune priming reduces pathogen transmission by directly decreasing bacterial shedding from infectious flies.
265

Figure 7: The effect of priming on fly locomotor activity. Mean ±SE total locomotor activity for males and females (n= 52 individual flies per treatment), during first 72-hours following systemic and oral priming and infection. (A) average total locomotor activity (B) average awake activity and (C) proportion of flies spent sleeping.

The *IMD*-pathway, but not the Toll pathway, is required for immune priming during *P. rettgeri* infections

In fruit flies, the production of AMPs during antibacterial immunity is mediated by the Immune deficiency (IMD) and Toll pathways. In both pathways, pathogens are recognised by

peptidoglycan receptors (PGRPs), initiating a signalling cascade that culminates in the
activation of the NF-κB-like transcription factors (*Dorsal* in Toll or *Relish* in IMD), resulting in
the upregulation of AMP genes. The Toll-pathway generally recognises LYS-type
peptidoglycan found in gram-positive bacteria and fungi. The IMD-pathway recognises DAP-
type peptidoglycan found in gram-negative bacteria and produces AMPs such as *Diptericins,*
Attacins and *Drosocin* among others. AMPs work with a high degree of specificity, so that
only a small subset of the total AMP repertoire provides the most effective protection against
specific pathogens (Hanson et al., 2019),although this specificity has been shown to be
greatly reduced during aging (Shit et al., 2022).
285
The inducible AMPs regulated by the IMD signalling pathway play a crucial role in resisting
gram-negative bacterial infection such as *P. rettgeri* (Lemaitre and Hoffmann, 2007).
Therefore, we wanted to investigate whether the IMD signalling pathway and IMD-responsive
AMPs contribute to immune priming. To address this, we used several transgenic fly lines
(CRISPR knockouts and UAS-RNAi knockdowns) with functional absence of or knockdown
of different regulatory and effector components of the IMD-signalling pathway. First, we used
a *Relish* loss-of-function mutant Rel^{E20}, a key regulator of the IMD immune response. As
expected, we found that *Relish* mutants did not show any bacterial clearance, died at faster
rate, and therefore did not show any benefit of a priming treatment (**Fig**. 8A-ii for survival and
Fig. 8B-ii for bacterial load, **Table**-S11 for survival and **Table**-S12 for bacterial load). Loss-
of—function of *spätzle (spz),* a key regulator in the *Toll* pathway, showed enhanced survival
following initial heat-killed exposure, and their mortality rates were the same as in controls
(w^{1118}) (**Fig**. 8A-iii, **Table**-S11) indicating that Toll-pathway does not contribute to priming
during *P. rettgeri* infection (**Fig**. 6A-iii, **Table**-S11).
300

Figure 8. Priming requires the IMD but not the Toll pathway. (A): Survival curves for control flies and flies lacking different innate immune pathway components in females and males (i) control *iso-w[1118]* (ii) *Rel[E20]*- IMD-pathway transcription factor (iii) *Spz*- Toll pathway regulator in both males and females (B) (i-iii) bacterial load measured after 24-hours post-secondary pathogenic exposure (n=7-9 vial/sex/treatment/transgenic lines). Different letters in panel-B indicate that primed and unprimed individuals are significantly different, tested using Tukey's HSD pairwise comparisons, analysed separately for each timepoint and sex combination. The error bars in panel B represent standard error.

Expression of IMD-regulated *Diptericin B* in the fat body is required for priming
Given the important role of the IMD pathway for the priming response (Fig. 5A), next, we tested whether mutants with defective IMD signalling or unable to produce specific antimicrobial peptides were capable of immune priming. We first used a ΔAMP transgenic line, which lacks most of the known *Drosophila* AMPs (10 AMPs in total). ΔAMPs flies are extremely susceptible to the majority of microbial pathogens, including gram-negative bacteria (Hanson et al., 2019), and we confirmed that both primed and unprimed ΔAMP flies succumb to death at a similar rate, and both primed and unprimed ΔAMP flies also exhibited elevated bacterial loads (measured 24-hours after the secondary pathogenic exposure) (**Fig. 9A-i** for survival, **Fig**. 9B-i for bacterial load, **Table**-S11 and S12, relative to control w[1118] flies – compare with previous figure Fig. 8B-i), showing that AMPs are needed for the priming response against *P. rettgeri*. To investigate which AMPs are required for priming against *P.*

rettgeri infection, we used a Group-B transgenic line, lacking major IMD regulated AMPs
(including *Diptericins* and *Attacins* and *Drosocin*) but have all upstream IMD signalling intact.
Primed Group-B flies showed mortality similar to unprimed Group-B flies (**Fig**. 9A-ii, **Table**
S11) and exhibited increased bacterial loads across both the sexes irrespective of being
primed or not (**Fig**. 9B-ii, **Table** S12). Thus, despite being able to produce other AMPs,
removal of IMD-regulated AMPs completely eliminated the priming effect, indicating that
AMPs regulated by the IMD-pathway are required for immune priming against *P. rettgeri*
infection.
332

333
334
Figure 9: Priming requires specific AMP expression. (A). Survival curves for
CRISPR/Cas9 AMP mutants (i) *ΔAMP* fly line: lacking all known 10 fly AMPs – (ii) *Group-B:*
lacking major IMD pathway AMPs (iii) *Dpt* knockout and (iv) *ΔAMP^{+Dpt}*: lacking all AMPs
except *Dpt*. (B) internal bacterial load quantified after 24-hours post-secondary exposure
(n=7-9 vials with 10-15 flies in each vial/sex/treatment/transgenic lines). Different letters in
panel-B denotes primed and unprimed individuals are significantly different, tested using

Tukey's HSD pairwise comparisons, analysed separately for for each fly line and sex
combination. The error bars in panel B represent standard error.
343
344

Since *Diptericins* have been shown previously to play a key role in defense against *P.*
rettgeri, we then used a *Dpt* mutant (lacking *Diptericin-A* and *B*) and *AMPs*$^{+Dpt}$ transgenic fly
lines (flies lacking all known AMPs except *Diptericin*) to test whether *Diptericins* are required
and sufficient for priming in both females and males. The survival benefit of priming
disappeared in flies lacking *Dpt* across both females and males and both primed and
unprimed *Dpt* mutants exhibited increased bacterial load (**Fig**. 9B-iii, **Table** S12). However,
the priming response was recovered completely in male flies that lacked other AMPs but
possessed functional *Diptericins* (*AMPs*$^{+Dpt}$). However, the same effect was not seen in
females (**Fig**. 9A-iv for survival, **Fig**. 9B-iv for bacterial load; **Table** S11 for survival, **Table**
S12 for bacterial load).
355

In response to gram-positive *Streptococcus pneumoniae* infection, previous work described
the role of haemocytes in immune priming through increased phagocytosis (Pham et al.,
2007). Subsequent work has also shown that reactive oxygen species (ROS) burst from
haemocytes is important for immune priming during *Enterococcus faecalis* infection
(Chakrabarti and Visweswariah, 2020). Since our results pointed to a *Diptericins* being
required for immune priming against *P. rettgeri*, we wanted to determine if *Diptericin*
expression in either the fat body or haemocytes was more important for immune priming.
Using-tissue specific *Diptericin-B* UAS-RNAi knockdown, we found that male flies with
knocked-down *DptB* in fat bodies no longer showed immune priming compared to the
background or control iso-w^{1118}, while knocking down *DptB* in haemocytes resulted in a
smaller but still significant increase in survival following an initial exposure (**Fig**. 10, **Table**-
S13). This would therefore support that immune priming requires *DptB* expression in the fat
body, while haemocyte-derived *DptB* is not important for priming. It is worth noting that it is
possible that haemocytes contribute to immune priming through phagocytosis or
melanisation, or via cross-talk with the IMD pathway, and this remains a question for future
research.
372
373

Figure 10: Survival curves tissue-specific Dpt knockdowns. (i) control *iso-w[1118]* and *UAS-RNAi* mutants - (ii) *FB>DptB* RNAi knock down of *DptB* in fat body and (iii) *FB>DptB* RNAi knock down of *DptB* in haemocytes. (n=~50 flies/sex/treatment/fly lines).

378
379
Priming is not an outcome of constant *Dpt* upregulation
As our results indicated that AMPs, especially *Diptericins* play a key role in *Drosophila*
priming against *P. rettgeri*, we wanted to test if the priming effect we had observed was a
result of constant upregulation of AMPs after initial heat-killed exposure allowing rapid
bacterial clearance during the secondary exposure, or if AMP expression returned to a
baseline level within the 96-hours between priming and the live infection. To do this, we
measured *Dpt* gene expression at 18-hours and 72-hours following initial heat-killed
exposure, and then 12-hours, 24-hours and 72-hours following secondary live *P. rettgeri*
infection. *DptB* expression increased 18-hours after initial heat-killed exposure but returned
to baseline levels by 72-hours (**Fig.** 11A and 11B, **Table**-S14) indicating that flies did mount
an immune response to the heat killed bacteria, but that this was resolved by the time they
were infected with the live bacteria at 96-hours.
392

393
Figure 11: Antimicrobial gene expression following priming (A) Mean±SE (standard
error) of *Diptericin*-AMPs expression at different time points (hours) for females and males
wild-type w[1118] flies with treated with - naïve or unhandled flies; flies exposed to initial heat-
killed *P. rettgeri*; primed flies that are initially exposed to heat-killed P. rettgeri followed by
challenge with *P. rettgeri*; unprimed flies that receive 1xPBS during primary exposure
followed by live *P. rettgeri* during secondary exposure [n=15-21 flies (3 flies pooled)
/sex/treatment/timepoint]. (B). *Diptericin (A+B), Attacin-C* and *Drosocin* AMPs expression 12-
hours and 24-hours after exposure to live *P. rettgeri* in w[1118] female flies. Asterisks '*'
indicates that primed and unprimed individuals are significantly different (p<0.05). The error
bars represent standard error.

Further, 72-hours following a lethal secondary live *P. rettgeri* infection, we found that primed

w^{1118} females and males showed increased *DptB* levels compared to unprimed individuals

(**Fig.** 12A and 12B). The increased *DptB* expression also resulted in increased bacterial

clearance in males (see **Fig**. 4Bi and 8Bi). In case of females, despite higher *Dpt* levels at

72-hours following secondary live infection in primed flies, we did not detect any difference in

bacterial clearance between primed and unprimed flies (see **Fig**. 4Bi and 8Bi). We also

tested the expression pattern of other IMD-responsive AMPs such as *Attacin-C* and *Drosocin*

in females. Overall, we found that the expression of *Attacin-C* and *Drosocin* was not different

between primed and unprimed females (**Fig.** 11C for *AttC and Dro*, **Table**-S14).

Figure 12: (A). Survival curves for males and females of Loss-of-function in *PGRPs*

(peptidoglycan recognition proteins) of IMD **(i)** *PGRP-LB* **(ii)** *PGRP-LC* and **(iii)** *PGRP-*

LE **(B).** internal bacterial load quantified after 24-hours post-secondary exposure in

female and male flies (n=7-9 vials with 10-15 flies in each vial/sex/treatment/ fly lines).

Different letters in panel-B denotes primed and unprimed individuals are significantly

different, tested using Tukey's HSD pairwise comparisons for each fly line and sex

combination. The error bars represent standard error.

Regulation of IMD by PGRPs is required for immune priming

Following gram-negative bacterial infection, DAP-type peptidoglycans from gram-negative bacteria are recognised by the peptidoglycan receptors *PGRP- LC* (a transmembrane receptor) and *PGRP-LE* (a secreted isoform and an intracellular isoform), which then activate the IMD intracellular signalling cascade (Lemaitre and Hoffmann, 2007; Valanne et al., 2022). Immune regulation is achieved by several negative regulators, including *PGRP-LB*, a secreted peptidoglycan with amidase activity, that break down peptidoglycans into smaller, less immunogenetic fragments(Lemaitre and Hoffmann, 2007; Valanne et al., 2022). *PGRP-LB* has also been shown to be important for transgenerational immune priming in *Drosophila* against parasitoid wasp infection (Bozler et al., 2020). Given the role of DptB in the observed priming benefit, we therefore decided to investigate the role of *PGRPs* in the immune priming we observed during *P. rettgeri* infection.

To address this, we used fly lines with loss-of-function in *PGRP-LB*, *LC* and *LE*. Regardless of which *PGRP* was disrupted, we observed that flies were no longer able to increase their survival following an initial exposure (**Fig**. 12A, **Table**-S15). However, unlike similar outcomes with *Relish* or *ΔAMP*, here the lack of priming was not driven by an inability to clear bacterial loads, as microbe loads in *PGRP* mutants were ~100-fold lower than in *Relish* or AMP mutants (**Fig**. 12B, **Table**-S16; also compare **Figs**. 8B-ii and 9B-i. Further, Dpt expression compared to the w^{1118} control was either similar (PGRP-LC, PGRP-LE) or higher (PGRP-LB), as expected, in both primed and unprimed flies (**Fig.** 13, **Table**-S17). This suggests that *Diptericin* expression is required, but not sufficient, for successful priming, which requires adequate regulation by *PGRPs*. Which specific molecular signal modifies *PGRP-mediated* regulation of *Diptericin* expression following an initial priming challenge is unclear, but must lie upstream of the IMD pathway and the fat body.

455 **Figure 13: *Diptericin*-AMPs expression 24-hours after exposure to live *P. rettgeri* in**
456 **male and female control *w^1118* flies and flies with loss-of-function in different *PGRPs*,**
457 ***PGRP-LB, PGRP-LC & -LE*.**
458
459

Discussion

The observation of immune priming in invertebrates underlines the importance of whole
organism research in immunity in lieu of a purely mechanistic approach to immunology,
highlighting that the same phenomenology can originate in very different mechanisms (Arch
et al., 2022; Little et al., 2005). There is now substantial evidence of diverse priming
responses occurring within the same invertebrate life stage (larvae or adults), across life
stages (from larvae to adults), or even between generations (parents to offspring) (Arch et
al., 2022; Contreras-Garduño et al., 2016; Khan et al., 2016; Sheehan et al., 2020). The
survival benefit of priming has been observed in a range of arthropod taxa, including
Dipterans: fruit flies (Pham et al., 2007), mosquitoes (Ramirez et al., 2015); Coleopterans:
flour beetles (Khan et al., 2016; Moret and Siva-Jothy, 2003); Lepidopterans: the greater
wax-moth (Fallon et al., 2011); Hymenopterans: bumblebee (Sadd and Schmid-Hempel,
2006); Crustaceans: water fleas (Little et al., 2003) and Arachnids: spiders and scorpions
(Gálvez et al., 2020). Thus, substantial experimental evidence in both lab adapted and wild-
caught arthropods suggests that immune priming is a widespread phenomenon, and is
predicted to have a profound impact on the outcome of host–pathogen interactions, including
infection severity and pathogen transmission (Tate and Rudolf, 2012; Tidbury et al., 2012;
Tate, 2016; Gomes et al., 2022).

In the present work, we investigated the occurrence, generality, specificity and mechanistic
basis of immune priming in fruit flies when infected with the gram-negative pathogen
Providencia rettgeri. We present evidence that priming with an initial non-lethal bacterial
inoculum results increased survival after a secondary lethal challenge with the same live
bacterial pathogen. This protective response may last at least two weeks after the initial
exposure, is particularly strong in male flies, and occurs in several genetic backgrounds. We
show that the increased survival of primed individuals coincides with a transient decrease in
bacterial loads, and that this is likely driven by the expression of the IMD-responsive AMP
Diptericin-B in the fat body. Further, we show that while *Diptericin* is required as the effector
of bacterial clearance, it is not sufficient for immune priming, which requires the regulation by
at least three PGRPs (*PGRP-LB, PGRP-LC*, and *PGRP-LE*). Therefore, despite having an
intact IMD signalling cascade, and being able to express *Diptericins*, flies lacking any one of
PGRP-LB, LC or *LE* were not capable of increasing survival following an initial sublethal
challenge. Together, our data indicates that PGRPs are necessary for regulating immune
priming against *P. rettgeri*.

These results are largely consistent with other work showing that the pathways that signal
pathogen clearance may not be the same that underlie the signalling of the priming
response. For example, Cabrera et al   investigated priming with the Gram-positive *E.*

faecalis and found that while the Toll pathway was required for responding to a single infection, Toll signalling was dispensable for immune priming (Cabrera et al., 2023). By contrast, IMD-deficient flies were still able to clear single *E. faecalis* infections (due a functional Toll response) but were no longer able to increase survival following an initial exposure. That is, in support of our results here, the IMD played a distinct role in signalling priming that was independent of its role in clearance. Further, previous work found that the Toll pathway is required (though not sufficient) for priming in response to gram-positive bacterial *Streptococcus pneumoniae* infection (Pham et al., 2007) and here we find that the Toll pathway is not required for a functional immune priming response.

In response to infection, the expression of several AMP genes in *Drosophila* increases and then drops once the infection threat is resolved or controlled. Previous studies have shown that this occurs within a period of few hours during bacterial infections (Lemaitre et al., 1997), and that in many cases, this increased AMP expression underlies the immune priming response (Arch et al., 2022; Wen et al., 2019) . In other host-pathogen systems, genome wide insect transcriptome studies have identified upregulation of several AMPs in primed individuals, for example, *Attacins, Defensins and Coleoptericins* in flour beetles (Ferro et al., 2019; Greenwood et al., 2017), *Cecropin, Attacin, Gloverin, Moricin* and *Lysozyme* in silkworms (Yi et al., 2019), *Gallerimycin and Galiomicin* in wax-moths (Bergin et al., 2006) and finally, *Cecropin* in tobacco moths and fruit flies (Roesel et al., 2020; Chakrabarti and Visweswariah, 2020). However, there are also examples where innate immune priming involves the downregulation of AMP expression, such as priming with the gram-positive *E. faecalis* (Cabrera et al., 2023).

One key aspect of our results is that immune priming was not the result of continued upregulation of *Diptericin* following the initial exposure to heat-killed bacteria. Instead, we observed that the initial immune response to heat-killed bacteria had already been resolved at 72-hours, before the secondary exposure to a lethal infection at 96-hours (see **Fig**. 11A and B). This second up-regulation of AMP expression was at least 10-times higher than the response to heat-killed bacteria, and was initially similar between both primed and unprimed flies (exposed to a sterile solution in the first exposure). It was only at 72-hours following the second lethal exposure that we observed significantly higher expression of *Diptericin* in primed flies. This difference appears to arise because unprimed flies show a faster resolution of the immune response compared to primed flies; that is, the expression of *Diptericin* shows a faster decline between 24-hours and 72-hours post-exposure in unprimed compared to primed flies (see **Fig**. 11A and B). These patterns of gene expression provide a partial explanation for the increased survival following priming, but they do not explain why primed individuals had lower bacterial loads at 24-hours after exposure to the second lethal infection, as at this timepoint we did not detect any differences in *Diptericin* expression between primed and unprimed flies. It is also unclear why primed females showed increased expression of *Diptericin* after 24-hours, but do not show any reduction in bacterial loads at this timepoint following priming, in the way male flies did.

The sex differences we observed in priming reflect a larger pattern of sexual dimorphism in
immunity present in most including *Drosophila* (Belmonte et al., 2020; Klein and Flanagan,
2016; Simon, 2005). Here we found that males showed better bacterial clearance after initial
exposure to heat-killed *P. rettgeri* which enabled them to experience enhanced survival
compared to females, who exhibited higher bacterial loads and greater mortality. One
possibility for the observed sex-differences could relate to different nutritional demands and
metabolic activities in female *Drosophila*. For instance, female fruit flies are often able to
reallocate resources in accordance with their reproductive demands, as observed in terminal
investment during infection (Camus et al., 2019; Martínez et al., 2020; Hudson et al., 2020).
Studies from other insects suggest that inducing priming responses can directly reduce the
reproductive fitness in mosquitoes (Contreras-Garduño et al., 2014), wax moth (Trauer and
Hilker, 2013), and mealworm beetles (Zanchi et al., 2011). There are therefore potential
trade-offs between investment in reproductive effort and investment in stronger immune
responses following priming. Given that all females used in this study were mated, it is
possible that such trade-offs forced a reallocation of resources towards reproduction, thereby
reducing the observed magnitude of the priming response in females. Future studies may
consider comparing priming responses in females with different reproductive states in order
to test whether immune priming is costlier for female *Drosophila*.

A subsidiary finding of this work was the effect of the endosymbiont *Wolbachia* on immune
priming. How endosymbionts that are widespread among insects are likely to influence
innate immune priming is a topic of considerable interest (Prigot-Maurice et al., 2022). There
is abundant evidence that *Drosophila* carrying the endosymbiont *Wolbachia* are better able
to survive infections, especially viral infections (Hedges et al., 2008; Teixeira et al., 2008;
Martinez et al., 2014; Vanika Gupta et al., 2017). In this case, we observed that the priming
response that was present in male flies cleared of *Wolbachia* disappeared in males carrying
the endosymbiont. This effect is unlikely due to a direct effect of *Wolbachia* on the ability to
clear *P. rettgeri* in primed flies, as previous work has found that *Wolbachia* had no effect on
the ability to suppress *P. rettgeri* during systemic infection (Rottschaefer and Lazzaro, 2012).
Indeed, if priming is the result of upregulation of AMP expression, this may suggest that
Wolbachia may be actively suppressing the expression of AMPs, thereby reducing the
beneficial effects of priming. However, this hypothesis would contradict work showing that
some *Wolbachia* strains upregulate the host's immune response and result in a reduction of
pathogen growth (Rancès et al., 2012; Chrostek et al., 2013). Further rigorous experimental
work is therefore required to fully understand this effect of *Wolbachia* on immune priming.

Finally, it is important to consider the implications of immune priming for disease ecology and
epidemiology, and particularly how it may affect pathogen transmission. If priming acts by
improving bacterial clearance via increased AMP expression, as observed in the present
work, we predict that priming is likely to reduce pathogen shedding at the individual level,
resulting in reduced disease transmission at the population level. Epidemiological models
that have incorporated priming predict that primed individuals with enhanced survival
following an initial sub-lethal pathogenic exposure are less likely to become infectious upon
re-infections (Tate, 2016). It remains unclear whether immune priming reduces pathogen

transmissibility, varies the infectious period, or alters infection-induced behavioural changes
in the host (Tate, 2016). An additional level of complexity is that there is likely to be
substantial within-population genetic variation in how priming affects each of these
components of pathogen transmission. Insects offer a powerful system to investigate these
effects, because they rely on an innate immune system that can induce an easily
measurable priming response, but further, insect are also important vectors of many
infectious diseases. Immune priming has thus emerged as a provocative idea to reduce the
vectorial capacity of insect vectors, thereby reducing the transmission of vector-borne
pathogens (Gomes et al., 2022). A better understanding of how immune priming contributes
to host heterogeneity in disease outcomes would aid our understanding of the causes and
consequences of variation in infectious disease dynamics.
598
599

600 ## Materials and Methods

601
Fly strains and maintenance
Several *D. melanogaster* strains: w^{1118} (Vienna *Drosophila* Resource Center)*, iso-w^{1118}*
(Bloomington *Drosophila* Stock Center)*, Canton-S, Oregon-R *Wolbachia^{+ve} and Oregon-R
Wolbachia^{-ve} (Vanika Gupta et al., 2017). Transgenic flies included immune mutants *Rel^{E20}*
(*relish - IMD* pathway regulator (Hedengren et al., 1999)) and *spz (spätzle – Toll* pathway
regulator (Levashina et al., 1998), and were a gift from the Saleh lab (Pasteur, Paris). We also
used the following CRISPR/Cas9 deletion lines, originally gifted by the Lematire lab (EPFL,
Lausanne) (Hanson et al., 2019): (a) ΔAMPs - flies lacking 10 fly AMPs, (b) Group-B - flies
lacking major IMD regulated AMPs including *Attacins* (*AttC^{Mi}; AttD^{SK1}), Drosocin (Dro^{SK4}) and
Diptericins (Dpt^{SK12}), (c) *Dpt^{SK12}* – flies lacking *Diptericins (DptA and DptB),* and (d)
ΔAMPs^{+Dpt} – flies lacking 10 known AMPs except *Diptericins*. All the CRISPR/Cas9 mutants
were generated previously from the *iso-w^{1118}* genetic background using CRISPR∕Cas9 gene
editing technology to induce null mutations in the selected genes (Hanson et al., 2019).
615
We also used the binary GAL4/UAS system for tissue specific silencing of target genes with
the RNA interference (RNAi) method. *Diptericin-B (DptB)* (Bloomington stock# *28975*) was
tissue-specifically knocked down in the fat body (*w$^{1118-iso}$,Fb-Gal4i+(P{fat})* and in
haemocytes [*w$^{1118-iso}$,Hmldelta- Gal4; He-Gal4* – a combination of two haemocyte GAL4
drivers, the Hml-GAL4.Δ (Sinenko and Mathey-Prevot, 2004) and He-GAL4.Z (Zettervall et
al., 2004)]. All fly stocks and experimental flies were maintained at 25°C ±1°C on a 12:12
hour light: dark cycle in vials containing 7ml of standard cornmeal fly medium (Lewis, 2014).
For the experiments, we controlled larval density by placing 10 females and 5 males in each
vial and the females were allowed to lay eggs for 48-hours. Fourteen days later, the eclosing
males and females were sorted and collected and separated into group of 25 flies in each
vial. Three-day old, mated individuals were used in all experiments.
627
Systemic immune priming and infection assays
P. rettgeri was grown at 37°C in 10ml Luria broth (Sigma Ltd) overnight to reach optical
density OD_{600}=0.95 (measured at 600nm in a Cell Density Meter, Fisherbrand™). The

culture was centrifuged at 5000 rpm for 5 min at 4°C, and the supernatant was removed, and
the final OD was adjusted to 0.1 and 0.2 by using sterile 1xPBS (Phosphate buffer saline).
To obtain heat-killed bacteria the dilution was incubated at 90°C for 20-30 mins (Khan et al.,
2016). To ensure all bacteria were dead in the heat-treated culture, it was plated and no
growth was confirmed. To prime individuals, 3-day old adults were pricked with a 0.14-mm
pin (Fine Science Tools) dipped in either heat-killed bacteria for the primed treatment or in
1xPBS solution for the unprimed treatment (exposed to a sterile solution in the first
exposure) in the mesopleuron region (the area situated under the wing and to the left of the
pleural suture) (Prakash et al., 2021). Following this initial priming treatment, the individuals
were pricked using OD_{600}=0.1 live *P. rettgeri* bacteria (resulting in approximately 70 bacterial
cells/fly). To test whether male and female adult flies show priming (measured as enhanced
survival) with increasing time intervals between the initial heat-killed exposure and later
challenge with live *P. rettgeri*, we tested several time points between the two challenges 18-
hours, 48-hours, 96-hours, 1-week and 2-weeks (See **Fig**. S1 for experimental design; n=9-
13 vial treatment/sex/ fly line). We used a split vial experimental design to obtain replicate
matched data for both survival and bacterial load see (Prakash et al., 2023) for details.
Briefly after infection each vial containing about 25 flies (of each treatment, sex and fly line
combination) were divided into 2 vials for measuring (i). survival following infection (see **Fig**.
S1i; 13-17 flies/combination) and (ii). internal bacterial load (see **Fig**. S1ii).
650

Bacterial load quantification
To test whether the host's ability to supress bacterial growth varies across primed and
unprimed individuals, we quantified bacterial load as colony forming units (CFUs) at 24-hours
after *P. rettgeri* infection for controls and transgenic flies in both sexes. Flies were surface
sterilized in groups of 3-5 flies per vial in 70% ethanol for 30s and washed twice with distilled
water before homogenising flies individually using micro pestles. We immediately performed
serial dilutions of the homogenate with 1xPBS and plated them on LB agar plates and
cultured at 29°C overnight. The following day, we counted the CFUs manually (Siva-Jothy et
al., 2018) (n=9-13 vial/sex/treatment/fly mutants).
660

Gene expression quantification
The expression of *Diptericins* was quantified by qRT-PCR. In parallel with the survival
experiment, we randomly selected a subset of control w[1118] individuals (both males and
females) for RNA extraction, we included 15 flies [5-7 replicates of 3 flies pooled together for
each treatment (primed and unprimed) for both males and females]. We randomly removed
selected flies (3 flies per vial) at different time points post exposure to *P. rettgeri* (18-hours
and 72-hours post-priming, 12-hours and 24-hours post-challenge). We then homogenised
pools of three in 80µl of TRIzol reagent (Invitrogen, Life Technologies). Homogenates were
kept frozen at -70°C until RNA extraction. We performed mRNA extractions using the
standard phenol-chloroform method and included a DNase treatment (Ambion, Life
Technologies).
672

673 We confirmed the purity of eluted RNA using a Nanodrop 1000 Spectrophotometer (version
674 3.8.1) before going ahead with reverse transcription (RT). The cDNA was synthesized from
675 2µl of the eluted RNA using M-MLV reverse transcriptase (Promega) and random hexamer

primers, and then diluted 1 : 1 in nuclease free water. We then performed quantitative RT-PCR (qRT-PCR) on an Applied Biosystems StepOnePlus machine using Fast SYBR Green Master Mix (Invitrogen) using a 10μl reaction containing 1.5L of 1:1 diluted cDNA, 5μl of Fast SYBR Green Master Mix an 3.5μl of a primer stock containing both forward and reverse primer at 1μM suspended in nuclease free water (final reaction concentration of each primer 0.35μM). For each cDNA sample, we performed two technical replicates for each set of primers and the average threshold cycle (Ct) was used for analysis. We obtained the *AMP* primers from Sigma-Aldrich Ltd; *Dpt*_Forward: 5' GACGCCACGAGATTGGACTG 3', Dpt_Reverse: 5' CCCACTTTCCAGCTCGGTTC 3', *AttC*_Forward: TGCCCGATTGGACCTAAGC, *AttC*_Reverse: GCGTATGGGTTTTGGTCAGTTC, *Dro*_Forward: ACTGGCCATCGAGGATCACC, *Dro*_Reverse: TCTCCGCGGTATGCACACAT. We used *RpL*49 as endogenous reference gene, *RpL*49_Forward: 5' ATGCTAAGCTGTCGCACAAATG 3', *RpL*49_Reverse: 5' GTTCGATCCGTAACCGATGT 3'. We optimised the annealing temperature (T▯) and the efficiency (Eff) of the *Dpt* primer pair was calculated by 10-fold serial dilution of a target template (each dilution was assayed in duplicate); *Dpt*: T▯= 59 °C, Eff= 102%; *AttC*: T▯= 60 °C, Eff= 94%; *Dro*: T▯= 61 °C, Eff= 104%.

Oral priming and infection

For oral priming and live infection, we adjusted the final concentration to OD_{600}=25 (Prakash et al., 2022; Siva-Jothy et al., 2018). We initially prepared vials for oral priming by pipetting 350-400 μl of standard agar [see (Siva-Jothy et al., 2018)] onto lid of a 7ml tubes (bijou vials) and allowed it to dry. Simultaneously, we starved the experimental flies on 12ml agar vials for 4 hours. Once the agar on the lids dried, we placed a small filter disc (Whattmann-10) in the lid and pipetted 80μl of heat-killed bacterial culture (primed treatment) or 5% sucrose solution (unprimed control treatment) directly onto the filter disc. Once the agar dried, we orally exposed flies (heat-killed only, primed and unprimed treatment) by adding approximately 10 flies per vial for 18-hours and then transferred the flies onto fresh Lewis food vials. After 3-days, we again prepared the bijou vials and once the agar dried, this time we added 80μl of live bacterial culture (OD_{600}=25) and exposed flies to live *P. rettgeri* for 18-hours. We then transferred flies onto fresh food vials and observed survival after oral exposure to *P. rettgeri* every 12 hours for the following 8 days.

Measuring locomotor activity

We measured the locomotor activity of single flies (n=52 flies for each for each sex and treatment combination) during three continuous days using a *Drosophila* Activity Monitor – DAM (v2 and v5) System (Pfeiffenberger et al., 2010), in an insect incubator maintained at 25°C ± 1°C in a 12 D: 12 L cycle. We then processed the raw activity data using the DAM System File Scan Software (Pfeiffenberger et al., 2010). We analysed fly activity data using three metrics (Anderson et al., 2022; V. Gupta et al., 2017; Siva-Jothy and Vale, 2019; Vale and Jardine, 2015): total activity, the average activity during 5-min activity bouts, and proportion of 5-min bouts with zero activity (which has been defined as sleep in *Drosophila* (Shaw et al., 2000)) (**Fig.** S2-III)

Measuring bacterial shedding

We measured the bacterial shedding of single flies (n = 8-12 flies per treatment and sex
combination) at a single time point, 4-hours following overnight oral bacterial exposure. We
chose this timepoint as in other work we have found that most faecal shedding of bacterial
pathogens occurs within the first 4 hours, and steadily decreases by 8 hours following
overnight oral infection. Following oral priming to either heat-killed bacterial culture (primed
treatment) or 5% sucrose solution (unprimed treatment), flies were exposed to an oral
infection with live infection with 80µl of live *P. rettgeri* culture (OD$_{600}$=25). Following 18 hours
of oral exposure, flies were placed individually into 1.5ml Eppendorf tubes with approximately
50µl of Lewis medium in the bottom of the tube for 4-hours. After 4-hours, flies were removed
from the tube and the remaining content of each tube was washed with 50µl of 1xPBS buffer
by vortexing thoroughly for at least 5 secs. We then plated these samples on a LB agar
plates, incubated them at 29°C and counted the colonies manually after 18-hours.
733
Measuring bacterial transmission
We measured transmission in groups of flies, by collecting age-matched donor and recipient
w^{1118} flies, separately for each sex. To test the effect of priming on the ability of flies to
transmit *P.rettgeri*, we orally exposed 3-day old w^{1118} flies (donor flies) with either 5%
sucrose and heat-killed bacteria (primed) or 5% sucrose only (unprimed). After 96-hours the
donor flies were exposed to OD$_{600}$=25 of *P. rettgeri* (*see oral infection section above*). The
infected donors were marked by cutting the corner of a fly wing. We then placed one donor
and five uninfected recipient flies in 7ml bijou vials with a small amount of Lewis food on the
lid for each treatment (heat-killed bacteria only and 5% sucrose exposed without live
infection) and sex-combination. After 4-hours exposure we surface-sterilised and
homogenized the flies and plated the homogenate to measure the presence or absence of *P.*
rettgeri infection inside each recipient fly, as an indication of successful transmission of
bacteria from the donors to the recipient flies. We also set up and plated flies in groups with
no infection, to confirm that our measures of *P. rettgeri* prevalence reflected successful
transmission from the donor flies.
749
Data analysis
All statistical analyses and graphics were carried out and produced in R (version 4.2.2) using
the ggplot2, coxhz and lme4 packages (Wickham, 2016; Therneau, 2015; Bates et al., 2014),
and all raw data and code is available at 10.5281/zenodo.7624084 (Prakash and Vale,
2023). We analysed the survival data following systemic *P. rettgeri* infection with a mixed
effects Cox model using the R package 'coxme' (Therneau, 2015) for different treatment
groups (that is, primed and unprimed) across both the sexes and fly lines (controls and
transgenic lines). We specified the model as: survival ~ treatment * sex * (1|vials/block), with
'treatment' and 'sex' and their interactions as fixed effects, and 'vials' nested within each
'block' as a random effect for control and transgenic lines. We used ANOVA to test the
impact of each fixed effect in the 'coxme' model. We analysed the bacterial load, measured
as log$_{10}$ bacterial colony-forming units (CFUs) at 24-hours following *P. rettgeri* infection. As
bacterial load data was non-normally distributed, we log-transformed the data and analysed
using a non-parametric one-way ANOVA (Kruskal-Wallis test) to test whether the 'treatment'
groups that is, primed and unprimed individuals significantly differed in internal bacterial load
for males and females of each fly lines (control and transgenic lines).

766
We analysed the gene expression data by calculating the ΔΔCT value (Livak and
Schmittgen, 2001). We used *RpL32* as a reference gene as it was expressing steadily in our
treatment and control conditions. We calculated fold change in gene expression relative to
the uninfected controls to calculate ΔΔCT and used ANOVA to test whether AMP expression
differed significantly between primed and unprimed treatment for males and females. Data
for bacterial shedding and transmission were non-normally distributed (tested using Shapiro-
Wilks test for normality) hence we performed non-parametric Wilcoxon Kruskal-Wallis tests
for bacterial shedding and transmission data.
775

Acknowledgments

We thank Bruno Lemaitre's lab for generating and generously sharing the CRISPR/Cas9
AMP transgenic lines. We thank Emily Robertshaw, Srijan Seal, Saubhik Sarkar and Pavan
Thunga for laboratory assistance. We thank Sveta Chakrabarti and Ashworth fly group
members for helpful discussion. Finally, we thank Angela Reid, Lucinda Rowe, James King
and Alison Fulton for help with media preparation. Figures *SI-1* & *SI-2*, were prepared using
Biorender platform. We acknowledge funding and support from the Branco Weiss fellowship
and a Chancellor's Fellowship to PFV; a Darwin Trust PhD studentship to AP from the
School of Biological Sciences, The University of Edinburgh, United Kingdom and the Ashoka
University, India. For the purpose of open access, the author has applied a Creative
Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising
from this submission.

References

Anderson L, Camus MF, Monteith KM, Salminen TS, Vale PF. 2022. Variation in
mitochondrial DNA affects locomotor activity and sleep in Drosophila
melanogaster. *Heredity* in press. doi:10.1101/2021.10.30.464953
Apidianakis Y, Mindrinos MN, Xiao W, Lau GW, Baldini RL, Davis RW, Rahme LG.
2005. Profiling early infection responses: Pseudomonas aeruginosa eludes host
defenses by suppressing antimicrobial peptide gene expression. *Proceedings of*
the National Academy of Sciences **102**:2573–2578. doi:10.1073/pnas.0409588102
Arch M, Vidal M, Koiffman R, Melkie ST, Cardona P-J. 2022. Drosophila melanogaster
as a model to study innate immune memory. *Frontiers in Microbiology* **13**.
Bates D, Mächler M, Bolker B, Walker S. 2014. Fitting Linear Mixed-Effects Models
using lme4. *arXiv:14065823 [stat]*.
Belmonte RL, Corbally M-K, Duneau DF, Regan JC. 2020. Sexual Dimorphisms in
Innate Immunity and Responses to Infection in Drosophila melanogaster. *Front*
Immunol **10**:3075. doi:10.3389/fimmu.2019.03075
Bergin D, Murphy L, Keenan J, Clynes M, Kavanagh K. 2006. Pre-exposure to yeast
protects larvae of Galleria mellonella from a subsequent lethal infection by
Candida albicans and is mediated by the increased expression of antimicrobial
peptides. *Microbes and Infection* **8**:2105–2112. doi:10.1016/j.micinf.2006.03.005

Boehm T, Swann JB. 2014. Origin and Evolution of Adaptive Immunity. *Annual Review of Animal Biosciences* **2**:259–283. doi:10.1146/annurev-animal-022513-114201

Bozler J, Kacsoh BZ, Bosco G. 2020. Maternal Priming of Offspring Immune System in *Drosophila*. *G3 Genes|Genomes|Genetics* **10**:165–175. doi:10.1534/g3.119.400852

Brownlie JC, Johnson KN. 2009. Symbiont-mediated protection in insect hosts. *Trends in Microbiology* **17**:348–354. doi:10.1016/j.tim.2009.05.005

Buchon N, Silverman N, Cherry S. 2014. Immunity in Drosophila melanogaster — from microbial recognition to whole-organism physiology. *Nat Rev Immunol* **14**:796–810. doi:10.1038/nri3763

Cabrera K, Hoard DS, Gibson O, Martinez DI, Wunderlich Z. 2023. Drosophila immune priming to Enterococcus faecalis relies on immune tolerance rather than resistance. *PLoS Pathog* **19**:e1011567. doi:10.1371/journal.ppat.1011567

Camus MF, Piper MD, Reuter M. 2019. Sex-specific transcriptomic responses to changes in the nutritional environment. *eLife* **8**:e47262. doi:10.7554/eLife.47262

Chakrabarti S, Visweswariah SS. 2020. Intramacrophage ROS Primes the Innate Immune System via JAK/STAT and Toll Activation. *Cell Reports* **33**:108368. doi:10.1016/j.celrep.2020.108368

Chrostek E, Marialva MSP, Esteves SS, Weinert LA, Martinez J, Jiggins FM, Teixeira L. 2013. Wolbachia Variants Induce Differential Protection to Viruses in Drosophila melanogaster: A Phenotypic and Phylogenomic Analysis. *PLOS Genetics* **9**:e1003896. doi:10.1371/journal.pgen.1003896

Contreras-Garduño J, Lanz-Mendoza H, Franco B, Nava A, Pedraza-Reyes M, Canales-Lazcano J. 2016. Insect immune priming: ecology and experimental evidences. *Ecological Entomology* **41**:351–366. doi:10.1111/een.12300

Contreras-Garduño J, Rodríguez MC, Rodríguez MH, Alvarado-Delgado A, Lanz-Mendoza H. 2014. Cost of immune priming within generations: trade-off between infection and reproduction. *Microbes and Infection* **16**:261–267. doi:10.1016/j.micinf.2013.11.010

Fallon JP, Troy N, Kavanagh K. 2011. Pre-exposure of Galleria mellonella larvae to different doses of Aspergillus fumigatus conidia causes differential activation of cellular and humoral immune responses. *Virulence* **2**:413–421. doi:10.4161/viru.2.5.17811

Ferro K, Peuß R, Yang W, Rosenstiel P, Schulenburg H, Kurtz J. 2019. Experimental evolution of immunological specificity. *Proc Natl Acad Sci USA* **116**:20598–20604. doi:10.1073/pnas.1904828116

Gálvez D, Añino Y, Vega C, Bonilla E. 2020. Immune priming against bacteria in spiders and scorpions? *PeerJ*. doi:http://doi.org/10.7717/peerj.9285

Gomes FM, Silva M, Molina-Cruz A, Barillas-Mury C. 2022. Molecular mechanisms of insect immune memory and pathogen transmission. *PLOS Pathogens* **18**:e1010939. doi:10.1371/journal.ppat.1010939

Greenwood JM, Milutinović B, Peuß R, Behrens S, Esser D, Rosenstiel P, Schulenburg H, Kurtz J. 2017. Oral immune priming with Bacillus thuringiensis induces a shift in the gene expression of Tribolium castaneum larvae. *BMC Genomics* **18**:329. doi:10.1186/s12864-017-3705-7

Gupta V., Stewart CO, Rund SSC, Monteith K, Vale PF. 2017. Costs and benefits of
sublethal Drosophila C virus infection. *Journal of Evolutionary Biology* **30**:1325–
1335. doi:10.1111/jeb.13096
Gupta Vanika, Vasanthakrishnan RB, Siva-Jothy J, Monteith KM, Brown SP, Vale PF.
2017. The route of infection determines Wolbachia antibacterial protection in
Drosophila. *Proc R Soc B* **284**:20170809. doi:10.1098/rspb.2017.0809
Hanson MA, Dostálová A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. 2019. Synergy
and remarkable specificity of antimicrobial peptides in vivo using a systematic
knockout approach. *eLife* **8**:e44341. doi:10.7554/eLife.44341
Hedengren M, BengtÅsling, Dushay MS, Ando I, Ekengren S, Wihlborg M, Hultmark D.
1999. Relish, a Central Factor in the Control of Humoral but Not Cellular
Immunity in Drosophila. *Molecular Cell* **4**:827–837. doi:10.1016/S1097-
2765(00)80392-5
Hedges LM, Brownlie JC, O'Neill SL, Johnson KN. 2008. Wolbachia and Virus
Protection in Insects. *Science* **322**:702–702. doi:10.1126/science.1162418
Hoffmann JA. 2003. The immune response of Drosophila. *Nature* **426**:33–38.
doi:10.1038/nature02021
Hudson AL, Moatt JP, Vale PF. 2020. Terminal investment strategies following infection
are dependent on diet. *Journal of Evolutionary Biology* **33**:309–317.
doi:10.1111/jeb.13566
Hultmark D. 2003. Drosophila immunity: paths and patterns. *Current Opinion in
Immunology* **15**:12–19. doi:10.1016/S0952-7915(02)00005-5
Khan I, Prakash A, Agashe D. 2019. Pathogen susceptibility and fitness costs explain
variation in immune priming across natural populations of flour beetles. *J Anim
Ecol* **88**:1332–1342. doi:10.1111/1365-2656.13030
Khan I, Prakash A, Agashe D. 2016. Divergent immune priming responses across flour
beetle life stages and populations. *Ecol Evol* **6**:7847–7855. doi:10.1002/ece3.2532
Klein SL, Flanagan KL. 2016. Sex differences in immune responses. *Nat Rev Immunol*
16:626–638. doi:10.1038/nri.2016.90
Kurtz J, Franz K. 2003. Evidence for memory in invertebrate immunity. *Nature* **425**:37–
38. doi:10.1038/425037a
Lanz-Mendoza H, Gálvez D, Contreras-Garduño J. 2024. The plasticity of immune
memory in invertebrates. *Journal of Experimental Biology* **227**:jeb246158.
doi:10.1242/jeb.246158
Lemaitre B, Hoffmann J. 2007. The Host Defense of Drosophila melanogaster. *Annu Rev
Immunol* **25**:697–743. doi:10.1146/annurev.immunol.25.022106.141615
Lemaitre B, Reichhart J-M, Hoffmann JA. 1997. Drosophila host defense: Differential
induction of antimicrobial peptide genes after infection by various classes of
microorganisms. *Proceedings of the National Academy of Sciences* **94**:14614–
14619. doi:10.1073/pnas.94.26.14614
Levashina EA, Ohresser S, Lemaitre B, Imler JL. 1998. Two distinct pathways can
control expression of the gene encoding the Drosophila antimicrobial peptide
metchnikowin. *J Mol Biol* **278**:515–527. doi:10.1006/jmbi.1998.1705
Lewis E. 2014. A new standard food medium. 1960 Drosophila Information Service. in.
Cold Spring Harb Protoc **2014**:pdb.rec081414. doi:10.1101/pdb.rec081414

Little TJ, Hultmark D, Read AF. 2005. Invertebrate immunity and the limits of
mechanistic immunology. *Nat Immunol* **6**:651–654. doi:10.1038/ni1219
Little TJ, O'Connor B, Colegrave N, Watt K, Read AF. 2003. Maternal Transfer of
Strain-Specific Immunity in an Invertebrate. *Current Biology* **13**:489–492.
doi:10.1016/S0960-9822(03)00163-5
Livak KJ, Schmittgen TD. 2001. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**:402–408.
doi:10.1006/meth.2001.1262
Martínez BA, Hoyle RG, Yeudall S, Granade ME, Harris TE, Castle JD, Leitinger N,
Bland ML. 2020. Innate immune signaling in Drosophila shifts anabolic lipid
metabolism from triglyceride storage to phospholipid synthesis to support
immune function. *PLoS Genet* **16**:e1009192. doi:10.1371/journal.pgen.1009192
Martinez J, Longdon B, Bauer S, Chan Y-S, Miller WJ, Bourtzis K, Teixeira L, Jiggins
FM. 2014. Symbionts Commonly Provide Broad Spectrum Resistance to Viruses
in Insects: A Comparative Analysis of Wolbachia Strains. *PLOS Pathogens*
10:e1004369. doi:10.1371/journal.ppat.1004369
Milutinović B, Kurtz J. 2016. Immune memory in invertebrates. *Seminars in Immunology*
28:328–342. doi:10.1016/j.smim.2016.05.004
Mondotte JA, Gausson V, Frangeul L, Blanc H, Lambrechts L, Saleh M-C. 2018.
Immune priming and clearance of orally acquired RNA viruses in Drosophila. *Nat*
Microbiol **3**:1394–1403. doi:10.1038/s41564-018-0265-9
Mondotte JA, Gausson V, Frangeul L, Suzuki Y, Vazeille M, Mongelli V, Blanc H,
Failloux A-B, Saleh M-C. 2020. Evidence For Long-Lasting Transgenerational
Antiviral Immunity in Insects. *Cell Reports* **33**:108506.
doi:10.1016/j.celrep.2020.108506
Moret Y, Siva-Jothy MT. 2003. Adaptive innate immunity? Responsive-mode
prophylaxis in the mealworm beetle, Tenebrio molitor. *Proceedings of the Royal*
Society of London Series B: Biological Sciences **270**:2475–2480.
doi:10.1098/rspb.2003.2511
Pfeiffenberger C, Lear BC, Keegan KP, Allada R. 2010. Processing Sleep Data Created
with the Drosophila Activity Monitoring (DAM) System. *Cold Spring Harb*
Protoc **2010**:pdb.prot5520. doi:10.1101/pdb.prot5520
Pham LN, Dionne MS, Shirasu-Hiza M, Schneider DS. 2007. A Specific Primed Immune
Response in Drosophila Is Dependent on Phagocytes. *PLoS Pathog* **3**:e26.
doi:10.1371/journal.ppat.0030026
Pollard AJ, Bijker EM. 2021. A guide to vaccinology: from basic principles to new
developments. *Nat Rev Immunol* **21**:83–100. doi:10.1038/s41577-020-00479-7
Prakash A, Khan I. 2022. Why do insects evolve immune priming? A search for
crossroads. *Dev Comp Immunol* **126**:104246. doi:10.1016/j.dci.2021.104246
Prakash A, Monteith KM, Bonnet M, Vale PF. 2023. Duox and Jak/Stat signalling
influence disease tolerance in Drosophila during Pseudomonas entomophila
infection. *Developmental & Comparative Immunology* 104756.
doi:10.1016/j.dci.2023.104756

Prakash A, Monteith KM, Vale PF. 2022. Mechanisms of damage prevention, signalling,
and repair impact disease tolerance. *Proceedings of the Royal Society B*
2021.10.03.462916. doi:10.1101/2021.10.03.462916
Prakash A, Monteith KM, Vale PF. 2021. Negative regulation of IMD contributes to
disease tolerance during systemic bacterial infection in Drosophila. *bioRxiv*
2021.09.23.461574. doi:10.1101/2021.09.23.461574
Prakash A, Vale P. 2023. Data and code for "The immune regulation and epidemiological
consequences of specific immune priming in Drosophila."
doi:10.5281/zenodo.7624085
Prigot-Maurice C, Beltran-Bech S, Braquart-Varnier C. 2022. Why and how do
protective symbionts impact immune priming with pathogens in invertebrates?
Developmental & Comparative Immunology **126**:104245.
doi:10.1016/j.dci.2021.104245
Ramirez JL, de Almeida Oliveira G, Calvo E, Dalli J, Colas RA, Serhan CN, Ribeiro JM,
Barillas-Mury C. 2015. A mosquito lipoxin/lipocalin complex mediates innate
immune priming in Anopheles gambiae. *Nat Commun* **6**:7403.
doi:10.1038/ncomms8403
Rancès E, Ye YH, Woolfit M, McGraw EA, O'Neill SL. 2012. The Relative Importance
of Innate Immune Priming in Wolbachia-Mediated Dengue Interference. *PLOS*
Pathogens **8**:e1002548. doi:10.1371/journal.ppat.1002548
Roesel CL, Rosengaus RB, Smith W, Vollmer SV. 2020. Transcriptomics reveals
specific molecular mechanisms underlying transgenerational immunity in
Manduca sexta. *Ecology and Evolution* **10**:11251–11261. doi:10.1002/ece3.6764
Roth O, Kurtz J. 2009. Phagocytosis mediates specificity in the immune defence of an
invertebrate, the woodlouse Porcellio scaber (Crustacea: Isopoda). *Developmental*
& Comparative Immunology **33**:1151–1155. doi:10.1016/j.dci.2009.04.005
Rottschaefer SM, Lazzaro BP. 2012. No Effect of Wolbachia on Resistance to
Intracellular Infection by Pathogenic Bacteria in Drosophila melanogaster. *PLOS*
ONE **7**:e40500. doi:10.1371/journal.pone.0040500
Sadd BM, Schmid-Hempel P. 2006. Insect Immunity Shows Specificity in Protection
upon Secondary Pathogen Exposure. *Current Biology* **16**:1206–1210.
doi:10.1016/j.cub.2006.04.047
Shaw PJ, Cirelli C, Greenspan RJ, Tononi G. 2000. Correlates of sleep and waking in
Drosophila melanogaster. *Science* **287**:1834–7.
doi:10.1126/science.287.5459.1834
Sheehan G, Farrell G, Kavanagh K. 2020. Immune priming: the secret weapon of the
insect world. *Virulence* **11**:238–246. doi:10.1080/21505594.2020.1731137
Shit B, Prakash A, Sarkar S, Vale PF, Khan I. 2022. Ageing leads to reduced specificity
of antimicrobial peptide responses in Drosophila melanogaster. *Proceedings of*
the Royal Society B 2022.06.25.497570. doi:10.1101/2022.06.25.497570
Simon V. 2005. Wanted: women in clinical trials. *Science* **308**:1517–1518.
Sinenko SA, Mathey-Prevot B. 2004. Increased expression of Drosophila tetraspanin,
Tsp68C, suppresses the abnormal proliferation of ytr-deficient and Ras/Raf-
activated hemocytes. *Oncogene* **23**:9120–9128. doi:10.1038/sj.onc.1208156

Siva-Jothy JA, Prakash A, Vasanthakrishnan RB, Monteith KM, Vale PF. 2018. Oral
Bacterial Infection and Shedding in Drosophila melanogaster. *JoVE* 57676.
doi:10.3791/57676
Siva-Jothy JA, Vale PF. 2021. Dissecting genetic and sex-specific sources of host
heterogeneity in pathogen shedding and spread. *PLOS Pathogens* **17**:e1009196.
doi:10.1371/journal.ppat.1009196
Siva-Jothy JA, Vale PF. 2019. Viral infection causes sex-specific changes in fruit fly
social aggregation behaviour. *Biology Letters* **15**:20190344.
doi:10.1098/rsbl.2019.0344
Tate AT. 2017. A general model for the influence of immune priming on disease
prevalence. *Oikos* **126**:350–360. doi:10.1111/oik.03274
Tate AT. 2016. The Interaction of Immune Priming with Different Modes of Disease
Transmission. *Frontiers in Microbiology* **7**:1102. doi:10.3389/fmicb.2016.01102
Tate AT, Rudolf VHW. 2012. Impact of life stage specific immune priming on
invertebrate disease dynamics. *Oikos* **121**:1083–1092. doi:10.1111/j.1600-
0706.2011.19725.x
Teixeira L, Ferreira Á, Ashburner M. 2008. The Bacterial Symbiont Wolbachia Induces
Resistance to RNA Viral Infections in Drosophila melanogaster. *PLOS Biology*
6:e1000002. doi:10.1371/journal.pbio.1000002
Therneau T. 2015. Mixed Effects Cox ModelsMixed Effects Cox Models. CRAN
repository.
Tidbury HJ, Best A, Boots M. 2012. The epidemiological consequences of immune
priming. *Proceedings of the Royal Society B: Biological Sciences* **279**:4505–4512.
doi:10.1098/rspb.2012.1841
Trauer U, Hilker M. 2013. Parental Legacy in Insects: Variation of Transgenerational
Immune Priming during Offspring Development. *PLOS ONE* **8**:e63392.
doi:10.1371/journal.pone.0063392
Valanne S, Vesala L, Maasdorp MK, Salminen TS, Rämet M. 2022. The Drosophila Toll
Pathway in Innate Immunity: from the Core Pathway toward Effector Functions.
The Journal of Immunology **209**:1817–1825. doi:10.4049/jimmunol.2200476
Vale PF, Jardine MD. 2015. Sex-specific behavioural symptoms of viral gut infection and
Wolbachia in Drosophila melanogaster. *Journal of Insect Physiology* **82**:28–32.
doi:10.1016/j.jinsphys.2015.08.005
Wen Y, He Z, Xu T, Jiao Y, Liu X, Wang Y-F, Yu X-Q. 2019. Ingestion of killed
bacteria activates antimicrobial peptide genes in Drosophila melanogaster and
protects flies from septic infection. *Developmental & Comparative Immunology*
95:10–18. doi:10.1016/j.dci.2019.02.001
White LA, Siva-Jothy JA, Craft ME, Vale PF. 2020. Genotype and sex-based host
variation in behaviour and susceptibility drives population disease dynamics.
Proceedings of the Royal Society B: Biological Sciences **287**:20201653.
doi:10.1098/rspb.2020.1653
Wickham H. 2016. Ggplot2: elegant graphics for dataanalysis.
Yi Y, Xu H, Li M, Wu G. 2019. RNA-seq profiles of putative genes involved in specific
immune priming in Bombyx mori haemocytes. *Infection, Genetics and Evolution*
74:103921. doi:10.1016/j.meegid.2019.103921

Zanchi C, Troussard J-P, Martinaud G, Moreau J, Moret Y. 2011. Differential expression
and costs between maternally and paternally derived immune priming for
offspring in an insect. *Journal of Animal Ecology* **80**:1174–1183.
doi:10.1111/j.1365-2656.2011.01872.x
Zettervall C-J, Anderl I, Williams MJ, Palmer R, Kurucz E, Ando I, Hultmark D. 2004. A
directed screen for genes involved in Drosophila blood cell activation. *PNAS*
101:14192–14197. doi:10.1073/pnas.0403789101

1037