1   **A Novel Memory Type in *C. elegans* Exhibits Post-Training Consolidation**
2

3   Daniel M. Merritt*[1,2], Alexandra Udachina[3,4], Ninon Freidel[3,4], Sylvia M. T. Almeida[5], Yan
4   Ming Anson Lau[5,6] and Derek van der Kooy[1,5]
5

Affiliations:

[1] Institute for Medical Science, University of Toronto, Toronto, Canada
[2] Current address: Department of Biological Sciences, Columbia University, New York, USA
[3] Human Biology Program, University of Toronto, Toronto, Canada
[4] Department of Psychology, University of Toronto, Toronto, Canada
[5] Department of Molecular Genetics, University of Toronto, Toronto, Canada
[6] Current address: Institute of Biomedical Engineering, University of Toronto, Toronto, Canada
13
14 * Correspondence to: dmerritt@gmail.com
15

Abstract

Memories are often categorized into types, reflecting their behavioural, anatomical and

molecular diversity: these classifications both aid understanding of the differences among

varieties of memory and help delineate the unifying cross-species principles underlying them. In

the nematode worm *Caenorhabditis elegans,* we find that an associative memory of the pairing

of the normally attractive odorant benzaldehyde and starvation depends on *de novo* transcription

and translation, is independent of CREB, and is produced by massed training: a pattern which

does not correspond to any of the well-characterized molecular categories of invertebrate

memory. Further, as has been shown for many memories in vertebrates, but not previously in

nematodes, we find that formation of this memory continues after removal of the stimuli initially

causing it, and that it is labile to disruption through protein synthesis inhibition following

training, but that inhibition of proteasomal activity does not extend the duration of the memory.

Previous findings have implicated insulin pathway signalling as a key component of this

benzaldehyde/starvation memory, however we find that the transcriptional activity required for

the memory is likely to be independent of the transcription factors that function at the terminus

of this pathway. These findings better characterize this model associative memory in relation to

other invertebrate memory types and identify ways in which it both shares their traits and differs

from them.

Introduction

Since at least the era of William James, researchers have understood the importance of

systematically classifying the diverse phenomena of memory to understand how they function

(James, 1890). Memories have been categorized along many different axes, including conceptual

structure (non-associative and associative memory), stimuli and paradigm (fear conditioned

memory, conditioned taste avoidance memory), cognitive function (episodic memory, semantic

memory, implicit memory, working memory), and by anatomical location (hippocampal

memory, cerebellar memory, etc.).

Within neuroscience, one of the most influential ways of categorizing memories has been

to divide them by the dissociable molecular mechanisms underlying them (and the corresponding

perturbations disrupting these mechanisms). These molecular mechanisms have proven to be at

least loosely correlated with the temporal sequence by which the memory develops and appear to

be widely conserved, particularly in invertebrates. In the fruit fly *Drosophila melanogaster*,

memories have been divided into short-term memory (STM), anesthesia-sensitive memory

(ASM), anesthesia-resistant memory (ARM), and long-term memory (LTM) (Tully et al., 1994),

with broadly similar divisions in *Aplysia* (Hawkins et al., 2006) (for a review of similar

delineations in mammals, see Hernandez & Abel (2008)). Across all animals, most of the longest

term memories require protein synthesis, as well as repeated training sessions (spaced training)

rather than single training blocks (massed training) to form (Smolen et al., 2016), although there

are prominent exceptions (Garcia et al., 1955).

Memory in the nematode worm *Caenorhabditis elegans* has been the subject of extensive

research, and a broad assortment of genes have been identified, organized into several signalling

cascades, which are required for various learning modalities. How these modalities relate to the

categories of memory studied in other model invertebrates has not been clearly worked out:

although types of memory are often treated as if they are broadly generalizable categories (at

least within invertebrates), evidence for this remains lacking.

	Memory Type			
Memory Characteristic	**STM**	**ASM**	**ARM**	**LTM**
Protein Synthesis-Dependent	No (Tully et al., 1994)	No (Tully et al., 1994)	No (Tully et al., 1994)	Yes (Tully et al., 1994)
Spaced Training Required	No (Davis, 2011)	No (Tully et al., 1994)	No (Tully et al., 1994)	Yes (Tully et al., 1994)
Blocked by Cold Shock	Yes (Saitoe et al., 2005)	Yes (Tully et al., 1994)	No (Tully et al., 1994)	No (Xia et al., 1998)
Duration	< 2 h (Tully et al., 1990)	~ 7 h (Tully et al., 1994)	2-4 days (Tully et al., 1994)	> 7 days (Tully et al., 1994)
Genes Involved	*dunce, rutabaga* (Tully & Quinn, 1985)	*amnesiac* (Tully et al., 1994)	*radish* (Folkers et al., 1993)	CREB (Yin et al., 1994)

Table 1 Characteristics of memory types in *Drosophila melanogaster*.

To better understand how these categories of memories which were primarily developed

from research in fruit flies and *Aplysia* apply in *C. elegans*, we characterized a model aversive

memory in which the smell of benzaldehyde, which is normally attractive to the worm, becomes

temporarily aversive following paired presentation of the odour with starvation (Nuttley et al.,

2002). We try to orient this memory within the categories established in the *Drosophila* literature

(**Table 1**) to understand the general principles, if any, underlying molecular memory types across

animals.

Materials and Methods

Strains and General Methods

All experiments were performed using wild type N2 *Caenorhabditis elegans* unless

otherwise stated. Worms were grown using standard techniques at 20°C on nematode growth

medium (NGM) agar plates seeded with *Escherichia coli* OP50. All behavioural experiments

were done on synchronized populations of worms 52 h after release from L1 arrest and were

performed in an environmentally controlled room at 20 °C and less than 25% humidity. Worms

used in behavioural tests did not experience starvation between release from L1 arrest and

initiation of training. The *C. elegans* strains N2, CF1038 *daf-16(mu86)*, RB759 *akt-1(ok525),*

YT17 *crh-1(tz2)* and VC3149 *crh-2(gk3293)* were provided by the CGC, which is funded by

NIH Office of Research Infrastructure Programs (P40 OD010440). The 6x outcrossed CQ528

pqm-1(ok485) strain was a generous gift from Coleen Murphy. The double CREB mutant

UT1343 *crh-1(tz2); crh-2(gk3293)* was created using standard methods.

The *akt-1(mm200)* allele was generated by ethyl methanesulfonate mutagenesis (Brenner,

1974), and isolated from a mixed population following repeated selection for learning mutants in

an approach similar to that of Colbert & Bargmann (1995). *mm200* contains a single base pair

change resulting in an p.L199F substitution, and the originally isolated strain was outcrossed 4x

to give UT1306 *akt-1(mm200)*. The *akt-1* rescue strain was created by microinjection of the

fosmid WRM065aH03 into UT1306 to give UT1309.

Statistical Analysis

No statistical test was performed to predetermine sample sizes, which were set at 9 test

plates for all experiments. The mean chemotaxis index (C.I.) and the standard error of the mean

(SEM) were calculated using Excel. In most experiments with multiple independent variables,

comparisons between C.I. were made by two- or three-way ANOVA (as appropriate) performed

in GraphPad Prism, which were then followed by t-tests using Tukey's multiple comparison test.

In experiments (**Fig. 2B**, **Supplemental Fig. 1**) where a full factorial design was impractical,

precluding the use of a two-way ANOVA, t-tests alone were performed to test specific predicted

effects, and the results adjusted for multiple comparison using Bonferroni correction. Differences

were considered significant when the adjusted $p<0.05$. All figures depict data from three

replicates performed on three different days. All statistical data are provided in **Supplemental**

Table 1.

CREB Phylogenetic Tree

The phylogenetic tree of CREB proteins shown in **Fig. 3A** was produced with T-Coffee

sequence alignment (Notredame et al., 2000), Gblocks curation (Castresana, 2000), PhyML

phylogeny reconstruction (Guindon & Gascuel, 2003) and TreeDyn rendering (Chevenet et al.,

2006) using http://www.phylogeny.fr (Dereeper et al., 2008). Sequence labels were edited

following tree generation for clarity.

Protein isoforms used for analysis were *C. elegans* CRH-1a (NP_001022859.1), CRH-2a

(NP_740985.2), *Drosophila melanogaster* CREBA-PA (NP_524087.3), *Drosophila*

melanogaster CREBB-PF (NP_996504.1), *Aplysia californica* CREB1 (XP_012939791.1),

Aplysia californica CREB1 (NP_001191630.1), *Mus musculus* CREB1A (NP_598589.2), *Mus*

musculus ATF-2 isoform 1 (NP_001020264.1) and *Mus musculus* CREB3 (NP_001369747.1).

Behavioural Tests

Benzaldehyde/starvation experiments were conducted as per Nuttley et al. (2002).

Briefly, learning was tested by training 1000-1500 worms for 1 hour on 6 mL 10 cm parafilm-

sealed NGM agar plates without bacteria, with either 2 µL of 100% benzaldehyde placed on a

small parafilm square on the inside centre of the lid (trained condition), or a parafilm square

without benzaldehyde placed on the inside centre of the lid (naïve condition). Following training,

worms were rinsed off the plate with M9 and divided into three approximately equal groups,

which were then moved to the centre of three fresh 10 cm NGM testing plates, each with 1 µL of

1% benzaldehyde (Bnz) in ethanol on one side, and a second spot of 1 µL 100% ethanol (EtOH)

on the other. 1 µL of 1 M sodium azide was placed on top of each odorant to paralyze the worms

upon reaching it. The number of worms within 2 cm of each point of odorant, and elsewhere on

the plate, was counted after 1 hour, with any worms that remained where they were placed in the

centre of the plate and appeared dead or injured discounted (usually < 2%), and the chemotaxis

index calculated using the equation C.I.=((#Bnz)-(#EtOH))/(#Total Worms).

Cycloheximide Treatment

126   Experiments utilizing cycloheximide (Bioshop) to disrupt protein synthesis were

127   performed using 1 mL of 6 mg/mL cycloheximide dissolved in water. This was added to 5 mL of

128   NGM agar, which had been melted and allowed to cool to below 60 °C, for a final concentration

129   of 1 mg/mL in the training plates and/or testing plates unless otherwise stated.

130   For cycloheximide experiments which employed a gap between training and testing, after

131   being washed off training plates, worms were suspended in 3 mL of M9 buffer, with or without 1

132   mg/mL of cycloheximide, and agitated on a rocker for 1 hour.

133   *Bortezomib Treatment*

134   Worms were treated with 60 µg/mL bortezomib (Cell Signaling Technology) for 5 hours

prior to training, during training, and during a 1 hour gap between training and testing. Prior to

training, bortezomib was added to HT115 *E. coli* which had been grown overnight (to saturation)

at 37°C in agitated LB media, and then concentrated 3 times by centrifugation followed by

resuspension of the pellet in an appropriate volume of M9. During training and the forgetting

gap, bortezomib was added to M9 buffer along with 0.006% benzaldehyde, or M9 buffer alone,

respectively. Naïve controls were treated with bortezomib in M9 buffer alone during both

training and the forgetting gap. Tubes containing worms were agitated on a rocker throughout

the experiment. Bortezomib was prepared as a concentrated stock solution at 300 µg/mL in M9.

α-amanitin/actinomycin D Treatment

Worms were treated with 100 µg/ml α-amanitin (Sigma) or 200 µg/ml actinomycin D

(Sigma) for 4 hours prior to, and subsequently during, training. Prior to training, drugs were

added to HT115 *E. coli* which had been grown overnight at 37 °C in agitated LB media before

being concentrated 3 times by centrifugation followed by resuspension of the pellet in an

appropriate volume of M9. During training, drugs were combined in M9 buffer along with

0.006% benzaldehyde, or M9 buffer alone for naïve controls. Tubes containing worms were kept

agitated on a rocker throughout the experiment. α-amanitin was prepared as a concentrated stock

solution at 1 mg/mL in M9, while actinomycin D was dissolved at 222 µg/ml in M9.

Cold Shock

Worms were cold shocked in 3 mL ice-cold M9 buffer for 6 minutes immediately before

and/or 500 µL ice-cold M9 buffer immediately after training, as indicated. Controls conditions

which were not cold shocked were allowed to settle in 20 °C M9 buffer for 6 minutes, before

and/or after training. Training took place at 20 °C in M9 buffer containing 0.006%

benzaldehyde, or in M9 buffer alone for naïve controls. Tubes containing worms were agitated

on a rocker during training.

Heat Shock

Worms were transferred to NGM agar plates with a lawn of *E. coli* OP50 which had been

pre-warmed to 37 °C for 2 hours, and subsequently heat shocked at this temperature for 45

minutes in an air incubator immediately prior to training.

Results

Benzaldehyde/Starvation Associative Memories are Protein Synthesis-Dependent

We first attempted to determine whether a model associative memory in which benzaldehyde is paired with starvation, resulting in subsequent aversion to benzaldehyde, is dependent on protein synthesis. Application of the chemical protein synthesis inhibitor cycloheximide during both initial training to avoid benzaldehyde, and during subsequent testing for benzaldehyde preference, revealed a strong effect of protein synthesis inhibition in preventing the memory, with only mild effects on naïve approach to benzaldehyde (**Fig. 1A**). Decreasing the concentration of cycloheximide resulted in a modestly weaker learning deficit while fully eliminating the naïve approach deficit (**Supplemental Fig. 1**).

To determine when protein synthesis inhibition was acting to inhibit the memory, we applied cycloheximide selectively during training, during testing, and during both training and testing, and evaluated the strength of the resultant memory. To our surprise, cycloheximide was capable of partially inhibiting the memory during either training or testing, with the strongest inhibition resulting when it was present during both (**Fig. 1B**). This suggested that protein synthesis was required even after removal of the training stimuli for complete learning to take place.

Cycloheximide Inhibits Consolidation Rather than Recall

We reasoned that the inhibition of memory we observed when cycloheximide was given during testing could be due to impaired chemotaxis during testing, impaired memory consolidation following training, or impaired memory retrieval. Impaired chemotaxis was inconsistent with the larger absolute chemotaxis indices we observed in the group receiving cycloheximide during both training and testing compared to those seen in the no drug group. To

Figure 1 Benzaldehyde/Starvation Memory is Translation- and Transcription-Dependent. **A)** Chemotaxis of wild type animals to a point of benzaldehyde after benzaldehyde/starvation training, with and without cycloheximide in training and testing plates. **B)** Chemotaxis of wild type animals to a point of benzaldehyde after benzaldehyde/starvation training, with cycloheximide absent, present during training, during testing, and during training and testing. **C)** Chemotaxis of wild type animals to a point of benzaldehyde after benzaldehyde/starvation training and a 1 hour consolidation gap, with cycloheximide absent, present during training, present during the gap, present during testing or present during training and the gap. **D)** Chemotaxis of wild type animals to a point of benzaldehyde following benzaldehyde/starvation training in α-amanitin or actinomycin D, after 4h pre-exposure to the drug. **E)** Chemotaxis of wild type animals to a point of benzaldehyde after benzaldehyde/starvation training following 5 hours of proteasome inhibition by bortezomib, with and without a 1 h forgetting period after training.

distinguish between the remaining two possibilities, we added a 1 h post-training gap between

training and testing for consolidation to occur in, reasoning if the observed memory inhibition

was due to impaired consolidation, cycloheximide given during this gap and during training (but

not during testing) would inhibit consolidation, resulting in a memory deficit, while

cycloheximide given during testing alone would have no effect. Conversely, if the memory

inhibition was caused by impaired recall, cycloheximide during testing alone should retain its

inhibitory effects. We find that cycloheximide during testing has no effect on memory given a 1

h post-training gap for consolidation to occur in, suggesting that it impairs memory consolidation

and that the effect of the drug during recall is in fact due to an extended period of memory

formation that begins during training and continues after its end (**Fig. 1C**).

Benzaldehyde/Starvation Memories are Transcription-Dependent

We next wondered whether benzaldehyde/starvation associative memories in *C. elegans*

were dependent on *de novo* transcription, in addition to translation. We employed the

transcriptional inhibitors α-amanitin and actinomycin D to inhibit transcription during training,

and find that actinomycin D can inhibit the memory to a comparable extent as cycloheximide. α-

amanitin was found to also inhibit memory to a lesser extent, but this was not statistically

significant after adjustment for multiple testing (**Fig. 1D**).

Proteasome Inhibition does not Extend Benzaldehyde/Starvation Memory Duration

Our finding that inhibition of protein translation impairs benzaldehyde/starvation

memory led us to wonder if inhibition of the proteasome might, conversely, prolong the duration

of the memory by extending the lifetime of the relevant translated proteins, inhibiting an

endogenous memory decay process. To test this hypothesis, we exposed worms to the small

molecule proteasome inhibitor bortezomib for 5 hours prior to training, during training, and then

during a 1 hour gap between training and testing: 6 hours of exposure to bortezomib at this

concentration has previously been shown to be sufficient to inhibit the *C. elegans* proteasome

(Melo & Ruvkun, 2012). To maximize our ability to see an increase in memory retention, we

trained worms in liquid, exploiting an earlier chance observation that duration of the memory is

shorter when worms are trained in liquid media than on agar plates.

We find that bortezomib-mediated inhibition of the proteasome does not result in

extension of the benzaldehyde/starvation memory (**Fig. 1E)**, suggesting that forgetting of this

memory in the worm was not mediated by the proteasome.

Heat Shock Impairs Benzaldehyde/Starvation Memory

Inhibition of protein synthesis by heat shock has been demonstrated in yeast (Lindquist,

1981), *Drosophila* (Lindquist, 1981), mammalian cell culture (Shalgi et al., 2013) and *C. elegans*

(Snutch & Baillie, 1983), and has been suggested to be mediated by pausing of translation

elongation (Shalgi et al., 2013)*.* Although not often used for classification of memories in flies

and *Aplysia sp.*, heat shock has been used to impair memory in *C. elegans* (Beck & Rankin,

1995; Rose & Rankin, 2001), presumably through inhibition of protein synthesis. We find that

heat shock prior to training partially blocks the benzaldehyde/starvation memory, with no effect

on naïve approach to benzaldehyde (**Fig. 2A**).

Benzaldehyde/Starvation Memory is Resistant to Cold Shock

In *Drosophila* (Quinn & Dudai, 1976), the slug *Limax flavus* (Yamada et al., 1992) and

the snail *Lymnaea stagnalis* (Sangha et al., 2003), post-training cold shock is capable of

disrupting the consolidation of one type of memory into another, and this effect has been

Figure 2 Heat Shock, but not Cold Shock, Inhibits the Benzaldehyde/Starvation Memory. **A)** Chemotaxis of wild type animals to a point of benzaldehyde after benzaldehyde/starvation training, with and without heat shock immediately before training. **B)** Chemotaxis of wild type animals to a point of benzaldehyde without cold shock, and with cold shock given before, after, and before and after training.

suggested to be mediated by cooling-induced disruption of protein synthesis (Takahashi et al.,

2013). In *Drosophila*, anesthesia-resistant memory is behaviourally differentiated from

anesthesia-sensitive memory by resistance to disruption by cold shock. While some memories in

C. elegans can be disrupted by cold shock (Morrison & van der Kooy, 1997; Nishijima &

Maruyama, 2017), long-term tap habituation and long-term 1-nonanol/food appetitive learning

have been reported to be cold shock-resistant (Aamodt, 2006, p. 49; Nishijima & Maruyama,

2017). To determine whether the benzaldehyde/starvation model associative memory is sensitive

to cold shock, we first evaluated various durations of cold shock to determine the maximum

duration we could subject worms to ice-cold M9 for without severely hindering subsequent

chemotaxis, and found that 6 minutes of cold shock resulted in only a minor deficit in

chemotaxis (**Supplemental Fig. 2**). This duration of cold shock is far longer than that required to

disrupt memory in analogous worm paradigms (Nishijima & Maruyama, 2017), suggesting that it

should be sufficient to reveal cold shock disruption if possible in our paradigm.

We previously determined that cold shock of appetitive food/salt memories in *C. elegans*

had distinct effects when performed before and after training (Morrison & van der Kooy, 1997).

Therefore, to determine whether this benzaldehyde/starvation memory was similarly sensitive to

cold shock, we tested the effects of cold shock before, after, and both before and after training.

We find that, contrary to food/salt memory but consistent with long-term tap habituation and 1-

nonanol/food memory, the benzaldehyde/starvation memory was cold shock-resistant under all

tested conditions (**Fig. 2B**).

CREB is Dispensable for the Benzaldehyde/Starvation Memory

Our finding that transcription and translation are required for formation of the

benzaldehyde/starvation memory prompted us to investigate candidate transcription factors

which might mediate memory formation. The basic region/leucine zipper (bZIP) transcription

factor CREB1 has been shown to be necessary for long-term memories in diverse organisms,

including *Aplysia* (Dash et al., 1990; Kaang et al., 1993), *Drosophila* (Yin et al., 1994), mice

(Bourtchuladze et al., 1994) and in some paradigms *C. elegans* (Amano & Maruyama, 2011;

Dahiya et al., 2019; Timbers & Rankin, 2011).

In most animals, CREB proteins exist as a diversified family, with the majority of

members not having any role in memory. While the *C. elegans* genome contains two genes

encoding for CREB family members, *crh-1* and *crh-2,* only the protein product of *crh-1* clusters

with known positive regulators of memory function in other animals (**Fig. 3A**). To determine

whether the benzaldehyde/starvation memory requires CREB, we tested a strain carrying the *crh-*

1(tz2) allele. We find that *crh-1(tz2)* mutants exhibit slightly impaired naïve approach to

odorants, as has been reported previously (Dahiya et al., 2019), but see no evidence of a

benzaldehyde/starvation learning deficit (**Fig. 3B**), consistent with our previous observations on

these mutants in more complex memory paradigms (Merritt et al., 2019). Since the *crh-1(tz2)*

allele deletes most of the bZIP domain of *C. elegans* CREB1 **(Fig. 3C)**, and the allele results in

the elimination of CREB as detected by anti-phospho-CREB antibody (Kimura et al., 2002), it is

a presumptive null, and we therefore conclude that the benzaldehyde/starvation memory is CRH-

1-independent.

While homology and behavioural evidence suggested that CRH-1 was the most likely

CREB family candidate involved in this memory, it remained possible that the CRH-2 was the

relevant CREB protein. We therefore tested a strain carrying the *crh-2(gk3293)* allele, which

eliminates the entire CRH-2 bZIP domain and is therefore a probable null, but found no evidence

of a learning deficit. A double mutant strain carrying mutations in both *crh-1* and *crh-2* shows no

additive phenotype over the *crh-1* mutant strain, suggesting that our failure to observe a learning

deficit was not due to genetic redundancy **(Fig. 3D)**.

DAF-16 Regulated Transcriptional Targets May Constitute the Necessary Translated Proteins

Previous work from our group and others (Cheng et al., 2022; C. H. A. Lin et al., 2010;

Tomioka et al., 2006) has described a key role for insulin signalling in olfactory and gustatory

learning in *C. elegans*. The extensive transcriptional changes known to take place downstream of

insulin signalling (Murphy, 2006) suggested a straightforward explanation: could the

requirement we observed for transcription in the benzaldehyde/starvation learning paradigm

result from insulin signalling-mediated changes in gene expression?

Canonical regulation of transcription through the insulin signalling pathway is mediated

by two downstream transcription factors with opposing regulatory effects: DAF-16

Figure 3 CREB is Dispensable for Benzaldehyde/Starvation Memory. **A)** Phylogenetic tree of CREB family members, with proteins implicated as positive regulators of learning highlighted in pink. **B)** Chemotaxis of wild type and *crh-1* mutant animals to a point of benzaldehyde after benzaldehyde/starvation training. **C)** Structure of *C. elegans* CRH-1 and CRH-2 proteins, with deletions tested indicated above. **D)** Chemotaxis of wild type, *crh-2* mutant and *crh-1;crh-2* double mutant animals to a point of benzaldehyde after benzaldehyde/starvation training.

(Schuster et al., 2010) and PQM-1 (Tepper et al., 2013). DAF-16 is excluded from the nucleus

by phosphorylation by the kinase AKT-1 when the insulin signalling pathway is active (Paradis

& Ruvkun, 1998), but in the absence of upstream signalling enters the nucleus to

transcriptionally activate target genes (class I genes). PQM-1 is excluded from the nucleus by

nuclear DAF-16, and so only in the presence of insulin signalling is it able to enter the nucleus to

transcriptionally activate its targets (class II genes) (Tepper et al., 2013).

We first sought to clarify whether the entire canonical insulin signalling pathway was

required for benzaldehyde/starvation associative learning. Previous work has shown a

requirement for the sole insulin receptor *daf-2*, its ligand *ins-1* and the downstream kinase *age-1*

in this paradigm (C. H. A. Lin et al., 2010). Since in the canonical pathway the final step before

transcription factor involvement is phosphorylation of the serine/threonine kinase AKT-1, we

first tested mutants in *akt-1* to confirm the necessity of the pathway up to the point of

transcriptional regulation. We find that *akt-1* mutant animals exhibit a severe abrogation of

learning (**Fig. 4A**) which can be rescued by wild type *akt-1* (**Supplemental Fig. 3**). This

confirmed that the requirement for the insulin signalling pathway extends at least as far in the

pathway as *akt-1*.

Since AKT-1 activity results in the nuclear localization of PQM-1 (via nuclear exclusion

of DAF-16), we reasoned that PQM-1 was the downstream transcription factor most consistent

with our observed effects of transcriptional inhibition. We therefore tested *pqm-1* loss of function

mutant animals (Tepper et al., 2013) for learning deficits, but to our surprise found that this

strain exhibited only an extremely weak learning deficit (**Fig. 4B**). *daf-16* null animals (K. Lin et

al., 1997) were also capable of learning (**Fig. 4C**), however they showed a mild deficit in naïve

approach to benzaldehyde similar to that previously reported in this strain (Tomioka et al., 2006).

This impaired approach made it difficult to conclusively rule out a simultaneous impairment of

learning comparable to *akt-1* mutants, since the loss of trained aversion to benzaldehyde we

observed could be due to either a genuine failure to learn or a movement disorder.

Discussion

Protein Synthesis-Dependent Consolidation Continues After Training

Our data suggest that withdrawal of protein synthesis inhibition after completion of

training allows for partial formation of the benzaldehyde/starvation memory. When training is

immediately followed by testing, inhibition of protein synthesis during either phase results in

significant abrogation of memory, and inhibition of protein synthesis during both results in

nearly total loss of memory. When a gap period is interspersed between training and testing

phases, however, preventing protein synthesis during testing or training individually no longer

impairs memory: memory formation is only blocked, and even then not fully, when protein

synthesis is inhibited during both training and the gap period. This pattern of results suggests that

there is a wide temporal window for protein synthesis-dependent memory formation to occur in,

and further, that there is no necessary requirement for protein synthesis during memory recall.

Post-training consolidation of memory has been extensively studied in mammals, flies

and *Aplysia*, primarily in the context of long-term memories. Consolidation in these organisms

can be disrupted by protein synthesis inhibition, with a critical period of a few hours post-

training generally constituting the window during which protein synthesis inhibitors are effective

(Flexner et al., 1965; Goelet et al., 1986; Wu et al., 2017) except during episodes of subsequent

reconsolidation (Nader et al., 2000). Our data support a similar mechanism in formation of the

benzaldehyde/starvation memory, whereby external stimuli set in motion memory formation

processes which continue after their removal.

The Identity of Gene Products Requiring Translation Remains Unclear

Many genes have been implicated in *C. elegans* memory, and in principle translation of

any of them might be required during learning. Our chemical inhibition data suggest that *de novo*

transcription and translation of previously inactive genes is required during learning. This

constrains the range of possible genes to those which are not normally transcribed and translated

in the relevant cell type, and to protein coding genes specifically (ruling out, for example, *odr-1-*

targeting endo-siRNAs (Juang et al., 2013)).

The key role played by insulin signalling during benzaldehyde/starvation associative

learning, and its well-studied termination in DAF-16- and PQM-1-mediated transcriptional

changes, make it an appealing mechanism to explain our drug inhibition effects, however the

data we present here suggest it is unlikely to be the causative mechanism. Our present findings,

in conjunction with the previous identification of *ins-1*, *daf-2* and *age-1* as required genes (C. H.

A. Lin et al., 2010), support the necessity of components of the insulin signalling pathway as far

downstream as AKT-1, but *pqm-1* mutants exhibit only extremely small learning deficits relative

to those caused by transcriptional or translational inhibition. *daf-16* mutant animals are more

ambiguous because of a fairly severe naïve approach deficit, however their learning deficit is still

weaker than that caused by our chemical manipulations. A requirement for *daf-16*-mediated

transcription is also mechanistically inconsistent with a requirement for transcription, since *akt-1*

mutations result in the constitutive nuclear localization of DAF-16, but result in abrogation of

learning. These findings suggest an insulin signalling-independent role of transcription in the

benzaldehyde/starvation memory, and further, that the role of the insulin signalling pathway in

learning is not mediated by its canonical transcriptional regulators DAF-16 and PQM-1.

Evidence for DAF-16-independent roles of the insulin signalling pathway in learning has

previously been identified in the worm (Tomioka et al., 2006), and highly paradigm-specific

roles for various components of the pathway have been suggested (Nagashima et al., 2019).

Research in other learning modalities in *C. elegans* has shown that regulation of

experience-dependent plasticity in synaptic size, thought to be a mechanism of learning, requires

translation of the ARP2/3 complex, and that this is negatively regulated by the pro-forgetting

protein MSI-1 (Hadziselimovic et al., 2014). This represents a second potential mechanism by

which transcriptional and translational inhibitors may impair memory in *C. elegans*.

Our finding that inhibition of the proteasome does not extend memory duration is

consistent with both a mechanism in which protein synthesis is necessary for formation of the

memory engram but does not itself constitute it, or with a proteasome-independent mechanism of

engram decay (for example, free cellular proteases). Genes mediating active processes of

forgetting in *C. elegans* have been identified (Arai et al., 2022; Bai et al., 2022; Hadziselimovic

et al., 2014; Inoue et al., 2013; Teo et al., 2022), however none have encoded proteins involved

in protein decay, consistent with our findings.

Benzaldehyde/Starvation Memory Exhibits Characteristics of Several Invertebrate Memory

Types

In fruit flies and *Aplysia*, memories produced by a single training session (massed

training) are generally independent of transcription, translation and the transcription factor

CREB, while repeated training sessions (spaced training) give rise to long-term memories

dependent on transcription, translation and CREB (**Table 1**). The *C. elegans* memory we

describe here does not clearly correspond to any of these: it depends on transcription and

translation, is independent of CREB, and is produced through a single massed training session.

Two very different interpretations of these results can be imagined. It could be that the

memory described here is a genuinely independent class of memory with different behavioural

and molecular requirements to those of other well-studied invertebrate memories. Alternatively,

it may be that the behavioural manifestation of this memory is mediated by an overlapping suite

of memory classes which independently correspond to the memory types seen in flies and

Aplysia, but which in conjunction manifest as a blended memory mechanism. However, since

this latter interpretation would at the very least require postulating a protein synthesis-dependent

memory produced by massed training as one of these constituent components, the requirement

for a novel class of memory cannot be fully avoided, and so we favour the former interpretation

as more parsimonious.

Massed training protocols in the worm have been reported to be either independent

(Amano & Maruyama, 2011; Nishijima & Maruyama, 2017) or dependent (Stein & Murphy,

2014) on *de novo* protein translation, depending on the protocol. While it is conceptually

possible that some apparently massed protocols involve cryptic spaced training, it is difficult to

see ways by which this could occur in ours: food is uniformly absent on the agar plate (or in

liquid media) during training, and the worms are exposed to the smell through the gas phase,

which is ubiquitously present. Our findings, in conjunction with a similar report of protein

synthesis-dependent massed learning using a different paradigm (Stein & Murphy, 2014),

challenge the view that protein synthesis-dependent memory necessarily requires spaced

training: in *C. elegans,* requirement for spaced training appears instead to depend on the exact

conditioned and/or unconditioned stimuli employed.

None of the manipulations we describe here fully prevent memory. It is possible that the

small fraction of retained memory in our experiments represents a (potentially short-term)

memory process which is independent of translation and transcription. Since experiments in a

similar paradigm found that only 50% of protein synthesis was inhibited by a concentration of

cycloheximide three times greater than the one we use (Amano & Maruyama, 2011), however, at

least for our experiments utilizing cycloheximide, limited drug efficacy is a more likely

explanation.

Learning from your mistakes is a valuable skill for all animals, and for most animals

there are many different mistakes which might profitably be learned from. Our findings suggest

that memory types across invertebrates, and even different memories within a single species,

may exhibit fewer unifying principles than is commonly appreciated, at least as revealed by the

systems of classification which have been influential in flies and *Aplysia*. This both highlights

the importance of considering the specific paradigm and species in question when reasoning

about the mechanisms of memory processes, and draws attention to the diversity of ways in

which evolution has provided mechanisms to change behaviour in response to a dangerous

world.

Acknowledgements

442   Some strains were provided by the CGC, which is funded by NIH Office of Research
443   Infrastructure Programs (P40 OD010440). We thank Coleen Murphy for the generous gift
444   of the 6x outcrossed *pqm-1* strain. We thank Sebastien Santini (CNRS/AMU IGS
445   UMR7256) and the PACA Bioinfo platform for the availability and management of the
446   phylogeny.fr website used for the analysis depicted in **Fig. 3A**.

Funding

448   NSERC RGPIN Grant 8319
449   NSERC CREATE in Manufacturing, Materials, Mimetics

Author contributions

DMM: Conceptualization, Data Curation, Formal Analysis, Project Administration,
Supervision, Visualization, Writing – Original Draft Preparation, Writing – Review and
Editing.
A.U.: Data Curation, Formal Analysis, Investigation, Methodology, Conceptualization,
Writing – Review and Editing.
N.F.: Data Curation, Investigation, Conceptualization, Writing – Review and Editing.
S.M.T.A.: Formal Analysis, Investigation, Conceptualization, Writing – Review and
Editing.
Y.M.A.L.: Investigation, Validation, Writing – Review and Editing.
D.v.d.K.: Conceptualization, Project Administration, Supervision, Funding Acquisition,
Writing – Review and Editing.

Competing interests

463   Authors declare no competing interests.

Data and materials availability

465   All data is available in the main text or the supplementary materials.

References

467   Aamodt, E. (Ed.). (2006). *The neurobiology of* C. elegans. Elsevier.

468   Amano, H., & Maruyama, I. N. (2011). Aversive olfactory learning and associative long-term

469   memory in *Caenorhabditis elegans*. *Learning & Memory*, *18*(10), 654–665.

470 Arai, M., Kurokawa, I., Arakane, H., Kitazono, T., & Ishihara, T. (2022). Regulation of

471    Diacylglycerol Content in Olfactory Neurons Determines Forgetting or Retrieval of

472    Olfactory Memory in *Caenorhabditis elegans*. *Journal of Neuroscience*, *42*(43), 8039–

473    8053.

474 Bai, H., Huang, H., Zhao, N., Gu, H., Li, Y., Zou, W., Wu, T., & Huang, X. (2022). Small G

475    Protein RAC-2 Regulates Forgetting via the JNK-1 Signaling Pathway in *Caenorhabditis*

476    *elegans*. *European Journal of Neuroscience*, *56*(12), 6162–6173.

477 Beck, C. D., & Rankin, C. H. (1995). Heat shock disrupts long-term memory consolidation in

478    *Caenorhabditis elegans*. *Learning & Memory*, *2*(3–4), 161–177.

479 Bourtchuladze, R., Frenguelli, B., Blendy, J., Cioffi, D., Schutz, G., & Silva, A. J. (1994).

480    Deficient long-term memory in mice with a targeted mutation of the cAMP-responsive

481    element-binding protein. *Cell*, *79*(1), 59–68.

482 Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics*, *77*, 71–94.

483 Castresana, J. (2000). Selection of Conserved Blocks from Multiple Alignments for Their Use in

484    Phylogenetic Analysis. *Molecular Biology and Evolution*, *17*(4), 540–552.

485 Cheng, D., Lee, J. S., Brown, M., Ebert, M. S., McGrath, P. T., Tomioka, M., Iino, Y., &

486    Bargmann, C. I. (2022). Insulin/IGF signaling regulates presynaptic glutamate release in

487    aversive olfactory learning. *Cell Reports*, *41*(8), 111685.

488 Chevenet, F., Brun, C., Bañuls, A.-L., Jacq, B., & Christen, R. (2006). TreeDyn: Towards

489    dynamic graphics and annotations for analyses of trees. *BMC Bioinformatics*, *7*(1), 439.

490 Colbert, H. A., & Bargmann, C. I. (1995). Odorant-specific adaptation pathways generate

491    olfactory plasticity in *C. elegans*. *Neuron*, *14*, 803–812.

Dahiya, Y., Rose, S., Thapliyal, S., Bhardwaj, S., Prasad, M., & Babu, K. (2019). Differential

Regulation of Innate and Learned Behavior by *Creb1/Crh-1* in *Caenorhabditis elegans*.

The Journal of Neuroscience, *39*(40), 7934–7946.

Dash, P. K., Hochner, B., & Kandel, E. R. (1990). Injection of the cAMP-responsive element

into the nucleus of *Aplysia* sensory neurons blocks long-term facilitation. *Nature*, *345*,

718–721.

Davis, R. L. (2011). Traces of *Drosophila* Memory. *Neuron*, *70*(1), 8–19.

Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., Dufayard, J.-F.,

Guindon, S., Lefort, V., Lescot, M., Claverie, J.-M., & Gascuel, O. (2008). Phylogeny.fr:

Robust phylogenetic analysis for the non-specialist. *Nucleic Acids Research*, *36*(Web

Server issue), W465-469.

Flexner, L. B., Flexner, J. B., & Stellar, E. (1965). Memory and cerebral protein synthesis in

mice as affected by graded amounts of puromycin. *Experimental Neurology*, *13*(3), 264–

272.

Folkers, E., Drain, P., & Quinn, W. G. (1993). Radish, a *Drosophila* mutant deficient in

consolidated memory. *Proceedings of the National Academy of Sciences of the United

States of America*, *90*(17), 8123–8127.

Garcia, J., Kimeldorf, D. J., & Koelling, R. A. (1955). *Conditioned aversion to saccharin

resulting from exposure to gamma radiation*. *122*(3160), 157–158.

Goelet, P., Castellucci, V. F., Schacher, S., & Kandel, E. R. (1986). The long and the short of

long–term memory—A molecular framework. *Nature*, *322*(6078), 419–422.

Guindon, S., & Gascuel, O. (2003). A Simple, Fast, and Accurate Algorithm to Estimate Large

Phylogenies by Maximum Likelihood. *Systematic Biology*, *52*(5), 696–704.

Hadziselimovic, N., Vukojevic, V., Peter, F., Milnik, A., Fastenrath, M., Fenyves, B. G., Hieber,

P., Demougin, P., Vogler, C., de Quervain, D. J.-F., Papassotiropoulos, A., & Stetak, A.

(2014). Forgetting is regulated via Musashi-mediated translational control of the Arp2/3

complex. *Cell*, *156*(6), 1153–1166.

Hawkins, R. D., Kandel, E. R., & Bailey, C. H. (2006). Molecular Mechanisms of Memory

Storage in *Aplysia*. *The Biological Bulletin*, *210*(3), 174–191.

Hernandez, P. J., & Abel, T. (2008). The role of protein synthesis in memory consolidation:

Progress amid decades of debate. *Neurobiology of Learning and Memory*, *89*(3), 293–

311.

Inoue, A., Sawatari, E., Hisamoto, N., Kitazono, T., Teramoto, T., Fujiwara, M., Matsumoto, K.,

& Ishihara, T. (2013). Forgetting in *C. elegans* is accelerated by neuronal communication

via the TIR-1/JNK-1 pathway. *Cell Reports*, *3*(3), 808–819.

James, W. (1890). The principles of psychology, Vol I. In *New York Holt* (Vol. 1, Issue 2).

Juang, B.-T., Gu, C., Starnes, L., Palladino, F., Goga, A., Kennedy, S., & L'etoile, N. D. (2013).

Endogenous Nuclear RNAi Mediates Behavioral Adaptation to Odor. *Cell*, *154*(5), 1010–

1022.

Kaang, B.-K., Kandel, E. R., & Grant, S. G. N. (1993). Activation of cAMP-Responsive genes

by stimuli that produce long-term facilitation in Aplysia sensory neurons. *Neuron*, *10*(3),

427–435.

Kimura, Y., Corcoran, E. E., Eto, K., Gengyo-Ando, K., Muramatsu, M.-A., Kobayashi, R.,

Freedman, J. H., Mitani, S., Hagiwara, M., Means, A. R., & Tokumitsu, H. (2002). A

CaMK cascade activates CRE-mediated transcription in neurons of *Caenorhabditis*

elegans. *EMBO Reports*, *3*(10), 962–966.

Lin, C. H. A., Tomioka, M., Pereira, S., Sellings, L., Iino, Y., & van der Kooy, D. (2010). Insulin

signaling plays a dual role in *Caenorhabditis elegans* memory acquisition and memory

retrieval. *The Journal of Neuroscience*, *30*, 8001–8011.

Lin, K., Dorman, J. B., Rodan, A., & Kenyon, C. (1997). *daf-16*: An HNF-3/forkhead Family

Member That Can Function to Double the Life-Span of *Caenorhabditis elegans*. *Science*,

278(5341), 1319–1322.

Lindquist, S. (1981). Regulation of protein synthesis during heat shock. *Nature*, *293*(5830), 311–

314.

Melo, J. A., & Ruvkun, G. (2012). Inactivation of Conserved *C. elegans* Genes Engages

Pathogen- and Xenobiotic-Associated Defenses. *Cell*, *149*(2), 452–466.

Merritt, D. M., Melkis, J. G., Kwok, B., Tran, C., & van der Kooy, D. (2019). Analysis of

Mutants Suggests Kamin Blocking in *C. elegans* is Due to Interference with Memory

Recall Rather than Storage. *Scientific Reports*, *9*(1), 2371.

Morrison, G. E., & van der Kooy, D. (1997). Cold shock before associative conditioning blocks

memory retrieval, but cold shock after conditioning blocks memory retention in

Caenorhabditis elegans. *Behavioral Neuroscience*, *111*(3), 564–578.

Murphy, C. T. (2006). The search for DAF-16/FOXO transcriptional targets: Approaches and

discoveries. *Experimental Gerontology*, *41*(10), 910–921.

Nader, K., Schafe, G. E., & Le Doux, J. E. (2000). Fear memories require protein synthesis in the

amygdala for reconsolidation after retrieval. *Nature*, *406*(6797), 722–726.

Nagashima, T., Iino, Y., & Tomioka, M. (2019). DAF-16/FOXO promotes taste avoidance

learning independently of axonal insulin-like signaling. *PLOS Genetics*, *15*(7), e1008297.

Nishijima, S., & Maruyama, I. N. (2017). Appetitive Olfactory Learning and Long-Term

Associative Memory in *Caenorhabditis elegans*. *Frontiers in Behavioral Neuroscience*,

11.

Notredame, C., Higgins, D. G., & Heringa, J. (2000). T-coffee: A novel method for fast and

accurate multiple sequence alignment. *Journal of Molecular Biology*, *302*(1), 205–217.

Nuttley, W. M., Atkinson-Leadbeater, K. P., & van der Kooy, D. (2002). Serotonin mediates

food-odor associative learning in the nematode *Caenorhabditis elegans*. *Proceedings of*

the National Academy of Sciences of the United States of America, *99*(19), 12449–12454.

Paradis, S., & Ruvkun, G. (1998). *Caenorhabditis elegans* Akt/PKB transduces insulin receptor-

like signals from AGE-1 PI3 kinase to the DAF-16 transcription factor. *Genes &*

Development, *12*(16), 2488–2498.

Quinn, W. G., & Dudai, Y. (1976). Memory phases in Drosophila. *Nature*, *262*(5569), 576–577.

Rose, J. K., & Rankin, C. H. (2001). Analyses of Habituation in *Caenorhabditis elegans*.

Learning & Memory, *8*(2), 63–69.

Saitoe, M., Horiuchi, J., Tamura, T., & Ito, N. (2005). *Drosophila* as a novel animal model for

studying the genetics of age-related memory impairment. *Reviews in the Neurosciences*,

16(2), 137–149.

Sangha, S., Morrow, R., Smyth, K., Cooke, R., & Lukowiak, K. (2003). Cooling blocks ITM and

LTM formation and preserves memory. *Neurobiology of Learning and Memory*, *80*(2),

130–139.

Schuster, E., McElwee, J. J., Tullet, J. M. A., Doonan, R., Matthijssens, F., Reece-Hoyes, J. S.,

Hope, I. A., Vanfleteren, J. R., Thornton, J. M., & Gems, D. (2010). DamID in *C. elegans*

reveals longevity-associated targets of DAF-16/FoxO. *Molecular Systems Biology*, *6*(1),

399.

Shalgi, R., Hurt, J. A., Krykbaeva, I., Taipale, M., Lindquist, S., & Burge, C. B. (2013).

Widespread regulation of translation by elongation pausing in heat shock. *Molecular*

Cell, *49*(3), 439–452.

Smolen, P., Zhang, Y., & Byrne, J. H. (2016). The right time to learn: Mechanisms and

optimization of spaced learning. *Nature Reviews Neuroscience*, *17*(2), Article 2.

Snutch, T. P., & Baillie, D. L. (1983). Alterations in the pattern of gene expression following

heat shock in the nematode *Caenorhabditis elegans*. *Canadian Journal of Biochemistry*

and Cell Biology, *61*(6), 480–487.

Stein, G. M., & Murphy, C. T. (2014). *C. elegans* positive olfactory associative memory is a

molecularly conserved behavioral paradigm. *Neurobiology of Learning and Memory*,

115, 86–94.

Takahashi, T., Takigami, S., Sunada, H., Lukowiak, K., & Sakakibara, M. (2013). Critical Period

of Memory Enhancement during Taste Avoidance Conditioning in *Lymnaea stagnalis*.

PLOS ONE, *8*(10), e75276.

Teo, J. H.-M., Kurokawa, I., Onishi, Y., Sato, N., Kitazono, T., Tokunaga, T., Fujiwara, M., &

Ishihara, T. (2022). Behavioral Forgetting of Olfactory Learning Is Mediated by

Interneuron-Regulated Network Plasticity in *Caenorhabditis elegans*. *ENeuro*, *9*(4).

Tepper, R. G., Ashraf, J., Kaletsky, R., Kleemann, G., Murphy, C. T., & Bussemaker, H. J.

(2013). PQM-1 Complements DAF-16 as a Key Transcriptional Regulator of DAF-2-

Mediated Development and Longevity. *Cell*, *154*(3), 676–690.

Timbers, T. A., & Rankin, C. H. (2011). Tap withdrawal circuit interneurons require CREB for

long-term habituation in *Caenorhabditis elegans. Behavioral Neuroscience*, *125*(4), 560–

566.

Tomioka, M., Adachi, T., Suzuki, H., Kunitomo, H., Schafer, W. R., & Iino, Y. (2006). The

Insulin/PI 3-Kinase Pathway Regulates Salt Chemotaxis Learning in *Caenorhabditis*

elegans. *Neuron*, *51*, 613–625.

Tully, T., Boynton, S., Brandes, C., Dura, J. M., Mihalek, R., Preat, T., & Villella, A. (1990).

Genetic Dissection of Memory Formation in Drosophila melanogaster. *Cold Spring*

Harbor Symposia on Quantitative Biology, *55*, 203–211.

Tully, T., Preat, T., Boynton, S. C., & Del Vecchio, M. (1994). Genetic dissection of

consolidated memory in *Drosophila*. *Cell*, *79*(1), 35–47.

Tully, T., & Quinn, W. G. (1985). Classical conditioning and retention in normal and mutant

Drosophila melanogaster. *Journal of Comparative Physiology A*, *157*, 263–277.

Wu, J.-K., Tai, C.-Y., Feng, K.-L., Chen, S.-L., Chen, C.-C., & Chiang, A.-S. (2017). Long-term

memory requires sequential protein synthesis in three subsets of mushroom body output

neurons in *Drosophila*. *Scientific Reports*, *7*(1), 7112.

Xia, S.-Z., Feng, C.-H., & Guo, A.-K. (1998). Temporary Amnesia Induced by Cold Anesthesia

and Hypoxia in *Drosophila*. *Physiology & Behavior*, *65*(4), 617–623.

Yamada, A., Sekiguchi, T., Suzuki, H., & Mizukami, A. (1992). Behavioral analysis of internal

memory states using cooling-induced retrograde amnesia in *Limax flavus*. *Journal of*

Neuroscience, *12*(3), 729–735.

Yin, J. C., Wallach, J. S., Del Vecchio, M., Wilder, E. L., Zhou, H., Quinn, W. G., & Tully, T.

(1994). Induction of a dominant negative CREB transgene specifically blocks long-term

memory in *Drosophila*. *Cell*, *79*(1), 49–58.