1 **Sensory prediction error drives subconscious motor learning outside of the laboratory**

2 Scott T. Albert[1], Emily Blaum[2], Daniel H. Blustein[3]
3 *[1]Neuroscience Center, UNC Chapel Hill, NC, USA*
4 *[2]Neuroscience Program, Rhodes College, Memphis, TN, USA*
5 *[3]Department of Psychology, Acadia University, Wolfville, NS, Canada*
6
Corresponding Author:
Daniel Blustein
18 University Ave, Wolfville, NS B4P 2R6, Canada
Phone: +1 902 585 1220
blustein.neuro@gmail.com

Author contributions:

Wrote main manuscript text (STA, DHB), prepared figures 1 and 4 (STA), prepared figures 2 and

3 (DHB, STA), collected the data (EB), analyzed the data (DHB), and supervised student

researchers (DHB). All authors contributed to the study design and reviewed the manuscript.

Abstract

Sensorimotor adaptation is supported by at least two parallel learning systems: an intentionally

controlled explicit strategy, and an involuntary implicit learning system. To investigate the error

sources that drive these two systems, past work has focused on constrained reaching, or finger

movements in laboratory environments: studies that have shown subconscious learning systems to

be driven in part by sensory prediction error (SPE), i.e., the mismatch between the realized and

expected outcome of an action. Yet while SPE-driven adaptation has become central to qualitative

and quantitative models about the brain, little to no work has been done to understand whether this

error class meaningfully contributes to more complex, skill-based motor actions we perform in our

daily life. To fill this gap, we designed a ball rolling task to explore whether SPEs can drive implicit

motor adaptation outside the laboratory, where motor skills require (1) multi-segmented whole-

body movements that (2) impart physical motion on external objects. After applying a visual shift,

participants rapidly adapted their rolling angles to reduce the error between the ball and target. We

removed all visual feedback and told participants to aim their throw directly toward the primary

target, revealing an unintentional 5.06° implicit adjustment to reach angles that decayed over time.

To determine whether this implicit adaptation was driven by SPE, we tested participants in a

second paradigm similar to Mazzoni and Krakauer (2006): participants were given a second aiming

target that would 'solve' the visual shift. Remarkably, after rapidly reducing ball rolling error to

zero (due to enhancements in strategic aiming), the additional aiming target caused rolling angles

to drift beyond the primary target by 3.15°. This involuntary overcompensation, which worsened

task performance, is a hallmark of SPE-driven implicit learning. These results show that SPE-

driven implicit processes, previously observed within simplified finger or planar reaching

movements, actively contribute to motor adaptation in naturalistic skill-based tasks involving

complex multi-segmented body movements that impart motion on physical objects.

Keywords: motor learning; sensory prediction error; visuomotor rotation; motor adaptation

New and Noteworthy: Implicit and explicit learning systems have been detected within simple

finger or reaching movements inside the laboratory. But how these systems impact movements

during complex whole-body, skill-based tasks has not been established. Here we demonstrate that

sensory prediction errors significantly impact how a person updates their movements, replicating

findings from the laboratory in an unconstrained ball-rolling task. This real-world validation is an

important step towards explaining how subconscious learning helps humans execute common

motor skills in dynamic environments.

Introduction

Current theories posit that implicit, i.e., subconscious, motor learning is driven in part by sensory prediction error (SPE): a mismatch between a movement's outcome versus the expected result[1–7]. The notion that errors are a critical implicit learning substrate is central to both descriptive and computational motor control models[1,3,8–10]. The visuomotor rotation (VMR) paradigm has been applied extensively to examine error-based adaptation in simplified motor tasks where participants perform a planar reaching or pointing movement to advance a cursor towards a virtual target (Fig 1, Inset). For example, SPE-driven learning was initially detected by Mazzoni and Krakauer[2], in an experimental condition where the arm was splinted to a tripod, and the movement of a single index finger was tracked. While these constraints on task complexity allow precise error manipulation[2,7], the SPE-learning mechanisms detected in these contexts may not generalize to skill learning in natural environments, where individuals must coordinate multiple body segments to impart a desired motion on an external object. In other words, do SPEs contribute to our motor control and learning within full-body actions we perform in day-to-day life, like soccer or bowling?

Figure 1. *Unconstrained motor task setup.* Participants wearing goggles with or without a visual shift rolled a tennis ball to a small target 6.1 m away on sequential trials. **Inset**. Traditional VMR experiment (adapted from[46])

In the rare cases where sensorimotor adaptation has been examined in skill-based settings[11–14], reliable assays to separate implicit learning from explicit (i.e., intentional) strategies have not been applied; these studies have either relied on the tenuous assumption that implicit learning alone persists once a perturbation has been removed, or have

attempted to reduce explicit learning by providing aiming instructions, an intervention that could

cause the opposite effect: priming explicit adjustments[13]. Furthermore, no attempts have been

made to dissociate between error sources that could drive the implicit system, in particular, SPEs

versus target (i.e., performance) errors[1,3,7,10,15] which co-occur unless specialized conditions are

used to isolate the SPE component.

Here we aimed to fill this gap using a task designed to detect SPE learning in a more naturalistic

skill-based task that requires complex multi-segmented body movements. By eliminating target

errors and isolating SPEs, we observed an obligatory overcompensation in motor actions,

demonstrating that subconscious adjustments to everyday movements are driven at least in part by

SPEs.

Results

We trained participants to roll a tennis ball (D = 0.425° at target displacement) towards a small

target (D=0.85°) 6.1m away; the ball arrived at the target about 750ms after release (Fig. 1). Unlike

constrained motor tasks where motion is restricted to one or two joints[2,7,16], our paradigm

introduced an array of additional control variables: wrist, elbow, and shoulder angle, bend at the

knees and hip, orientation of the trunk, release timing, grip, and arm speed. Further, to internalize

sensory error, the brain needed to map visual error between the ball and target onto the participant's

temporally separated motion (conditions known to attenuate implicit learning[8,17,18]).

Despite these complex elements, participants achieved excellent performance, hitting the target

with high precision (Fig. 2A,B, first 40 trials; 0.095° average error [mean across all 27 participants

of median error on last 10 trials of initial baseline trials]). Next, we rotated the participant's visual

scene with prism glasses. The glasses, which occluded the view of the body and throwing arm,

caused participants to perceive the target offset by 30° in space from its true position. Past the

target, balls disappeared underneath a black curtain, which along with black flooring, reduced

spatial reference cues. In response, participants committed throwing errors (Fig. 2A, blue rotation

period) which induced rapid adaptation; participants adjusted their movement angle to mitigate the

discrepancy between ball and target, reaching asymptotic performance[19] by approximately their

13th trial with -1.29° error persisting over the last 3 trials of the period (Fig. 2A,C).

Figure 2. *Implicit and explicit components of unconstrained movements*. **A.** Trial-by-trial lateral error during ball rolling relative to the perceived target location with no aiming target provided (n = 27). Black-outlined datapoints represent trials with no visual shift applied. Blue-outlined points represent trials with a 30° visual shift applied. After the first two visually shifted trials (blue squares), participants were

reminded of the goal to hit the target. Black-filled datapoints indicate the absence of endpoint visual error feedback. **B.** Trial-by-trial error for participants given an aiming target to minimize error (n = 27). Red-outlined datapoints indicate trials with a 30° visual shift applied. After the first two visually shifted trials (red squares), the aiming target was added and described. **C.** Mean residual error on the last three trials of the visual shift block drifted beyond the target for the aiming-target group (red boxplot). Color-coded lines connect each individual participant's results for both conditions: blue lines represent participants who completed the no-aiming-target condition first; red lines represent participants who completed the aiming-target condition first. **D.** Statistically significant differences between aiming-target and no-aiming-target groups in total adaptation and explicit adaptation. Error bars represent standard error and asterisks indicate results of Bonferroni-corrected p-values from paired t-tests [* p=0.012; ** p=0.0005].

Was this rapid motor adaptation solely dependent on explicit changes in throwing strategies, e.g.,

intended aiming angle, or was some adaptation due to subconscious implicit learning? To measure

implicit and explicit learning components, we instructed participants to continue making throws

over 20 trials, but without error feedback. On these trials, participants could view the initial target

location, but the room was darkened once the ball left the hand. For the initial 10 no-feedback

trials, the rotation was maintained. Then, participants switched to the unshifted glasses for an

additional 10 no-feedback trials. For these trials, participants were told that the glasses had been

switched, and that they should aim towards the initial target location. We gave these instructions

to match the conditions used to measure implicit learning via aftereffect trials in modern VMR

experiments[1,8,20]. Throwing angles decreased from 28.71° (relative to perceived target location) at

the end of the adaptation period, to approximately 5.06° on the initial 3 unshifted no-feedback

trials (Fig. 2A, implicit aftereffect across Trials 11-13 of the no-feedback period; Trials 1-10 were

not used as vision remained shifted and no aiming instruction had been provided). This immediate

voluntary change in angle implied that participants had explicitly aimed about 23.7° away from

the perceived visual target (explicit strategy was estimated by subtracting implicit learning from

total adaptation during the shifted vision block). An additional 5° of subconscious implicit learning

decayed after a return to unshifted vision, both in the absence of feedback (one sample Bonferroni-

corrected t-test confirming the slope of error over these trials to be significantly different from

zero, $t(26) = -1.54$, $p = 0.0097$) and over the ensuing washout period ($t(26) = -2.74$, $p < 0.0001$).

In sum, motor behavior in our task showed a clear correspondence to lab-based VMR studies: (1)

Participants adapted to the rotation, achieving steady-state performance that never completely

counteracted the rotation[3,8,21]. (2) Learning was supported by two parallel systems[2,7,22,23]: an

explicit strategy that participants could willfully disengage, and an implicit correction that

persisted despite the participant's conscious intentions. But which error sources had engaged the

implicit learning system? Implicit learning could have been caused by SPEs (error between

predicted and realized motion), task errors (error between ball and target)[1,3,10,15], or a model-free

reinforcement of successful motor actions[24].

To test whether SPEs contributed to adaptation we tested subjects in a second paradigm similar to

Mazzoni and Krakauer (2006) (Fig. 2B). Here, after performing two ball rolls with rotated vision,

participants were told they could aim towards an assistive target in order to move the ball to the

desired goal target. Subjects in this aiming-target condition adopted this strategy quickly, reaching

steady-state performance in about 10 trials (see Methods[19]). Remarkably, however, despite rapidly

achieving zero error (i.e., hitting the target's center), the ball angle rapidly surpassed the target's

location (horizontal dashed line in Fig. 2B), overcompensating for the imposed rotation. By the

last 3 trials of the period, the ball's path deviated by 3.15° *beyond* the task-related target (Fig. 2C),

exceeding on average the target's extent (radius = 0.425°, $t(26) = 2.89$, $p = 0.0077$, 95% CI lower

bound = 1.21°). This involuntary overcompensation in reach angle is a distinct hallmark of SPE-

driven implicit learning, indicating that SPE learning mechanisms detected in simplified arm and

hand movements studied in the laboratory, also contribute to subconscious adaptation in skill-

based tasks involving multi-segmented full-body motion and object manipulation.

To examine why this involuntary overcompensation occurred only when a secondary aiming target

was provided, we compared implicit and explicit learning measures across the aiming-target and

no-aiming-target conditions. As expected, the increase in total adaptation observed in the aiming-

target condition (main effect of aiming target on total adaptation with a multiple linear regression;

$F(2,51) = 10.00$, $p = 0.00022$ for total adaptation response variable; $\beta_{aiming\text{-}target} = 4.51$, $t(24) = 4.19$,

$p = 0.00011$), was supported by an increase in explicit strategy (Fig. 2D; no aiming target: 23.64°;

aiming target: 27.85°; $t(52) = -3.00$, $p = 0.0123$; follow up paired t-test with Bonferroni correction,

see Methods); thus, participants utilized the aiming target and instructions to increase their explicit

strategy. On the contrary, the aftereffects observed immediately upon returning to unshifted vision,

indicating implicit adaptation, did not show a statistically significant difference between groups

(5.06° (no aiming target), 5.30° (aiming target) $(F(2,51) = 0.89$, $p = 0.42$ for implicit adaptation

response variable). This indicated that the SPEs related to the goal target (no-aiming-group) and

aiming-target (aiming-group) both drove implicit learning in a similar manner, suggesting that

implicit learning was largely immune to any change in explicit strategy in our experiment. This

was further supported at the individual participant level; we did not observe that individuals with

greater implicit aftereffects exhibited greater total adaptation, and hence overcompensation.

Specifically, we observed only weak correlations between total adaptation and implicit adaptation

across the no-aiming-target condition $(r(25) = -0.25$, $p = 0.20)$ and the aiming-target condition

$(r(25) = 0.098$, $p = 0.63)$. In sum, overcompensation appeared to be due to an overactive explicit

process that did not appropriately counterbalance obligatory SPE-driven learning in the implicit

learning system.

This overcompensation pattern appeared largely insensitive to the order in which each condition

was experienced (i.e., all participants completed the aiming-target and no-aiming-target periods,

but with their order counterbalanced). We did not observe a statistically significant increase in

overcompensation when aiming targets were provided during the second learning period (4.31°,

Fig. 3D) than the first learning period (2.06°, Fig. 3B) (t-test: $t(25) = -1.20$, $p = 0.24$). Similarly,

during the no-aiming-target condition, we also did not detect a statistically significant change in

residual error between the first (Fig. 3A, -2.05°) and second (Fig. 3C, -0.59°) learning periods

(independent t-test: $t(25) = -1.33$, $p = 0.20$). On the contrary, initial learning rates were highly

sensitive to the block order; both the aiming-target and no-aiming target conditions exhibited an

increased learning rate when these periods were experienced during the second block of the

experiment, likely owing to a savings in explicit strategy[25–27] upon re-exposure to the rotation

[mean error on the first two shifted vision trials on Block 1 (19.36°, Fig. 3A,B) exceeded that in

Block 2 (-4.71°, Fig. 3C,D); paired t-test ($t(26)=-8.09$, $p<0.0001$]. These increases in overall

adaptation during the early phase of adaptation, but not the later steady-state period suggests that

participants downregulated strategic aiming over time, likely to mitigate the obligatory SPE-driven

increases in the implicit process[22].

Figure 3. *Trial-by-trial error across all block order and strategy condition pairs.* **A.** Data plotted for participants completing their first block without an aiming target (n = 13). Plotting conventions as described in Figure 2. **B.** Data from Block 1 of participants using an aiming target (n = 14). **C.** Data from Block 2 of participants without an aiming target who first used an aiming target, the same participants as in panel B (n = 14). **D.** Data from Block 2 participants with an aiming target who first had no aiming target, the same participants as in panel A (n = 13).

Discussion

Error-based motor adaptation has been studied extensively in laboratory experiments that constrain motor tasks to the planar motion of a finger, hand, or forelimb. Little is known about whether this vast body of work applies to more complex skill-based motor actions we perform in day-to-day life involving coordination across multiple segments across our entire body. Here, we observed

the first direct evidence that (1) subconscious error-driven learning processes make substantial

contributions to more sophisticated motor actions, and (2) that this learning is in part mediated by

SPEs not only in our body's motion, but in the motion of external objects (i.e., the ball in our task).

Admittedly, using prism glasses to shift the visual world is not naturalistic. These findings simply

represent a step towards more rigorous study of implicit and explicit learning processes "in the

wild". To this end, our study differs from past skill-based learning investigations[11–14] in notable

ways. Most importantly, earlier reports[2,7,17,22,23] have inferred implicit (or procedural) adaptation,

under the assumption that the aftereffect during the washout period is entirely driven by implicit

processes. This is problematic, as there is no guarantee that participants will stop using explicit

strategies, especially when errors are not removed during the washout period which can continue

to guide aiming strategies.

Furthermore, there has been no attempt to delineate the potential error sources which could drive

subconscious adaptation in a skill-based setting. That is, SPEs and target-errors always co-occur

with one another (i.e., the discrepancy between the hand/object and target creates a task error, and

a discrepancy between the hand/object and predicted path creates the SPE) and are both known to

drive implicit adaptation[1,3,7,10,15]. On the other hand, implicit recalibration may occur through an

error-free process such as reinforcement learning. That is, strategies could initially modulate the

motor plan, and then be transferred to the implicit system through the development of a use-

dependent bias[24,28]. In sum, special experimental conditions must be used to isolate adaptation that

can only be attributed to SPEs, which have rarely been applied even in lab-based settings[1,2,7].

Here we solved both these issues by adopting the standard best practices in measuring the implicit

contribution to adaptation: (1) instructing participants to not use strategy (i.e., aim directly to the

target)[3,8,25], and (2) removing all visual feedback that could be used to inform them about the

success of their movements[29,30] (in our case, by darkening the room during the no-aiming period).

Second, to ensure that this implicit aftereffect was at least in part driven by SPEs, we provided a

secondary aiming landmark[2,7] that helped participants eliminate the target error; adaptation beyond

this point must be driven by an SPE, because there is no remaining discrepancy between the ball

and the primary target.

The result was a clear implicit aftereffect that exhibited decay properties resembling those studied

in lab-based settings. That is, even in the absence of error, the act of moving and the passage of

time alone resulted in the partial loss of this memory trial-to-trial[1,29,31–34]. Further, we could

attribute this implicit aftereffect at least in part to SPEs due to the striking overcompensation

beyond the primary target (Fig. 2B, overcompensation beyond the target). It is rather remarkable

that this distinct SPE-driven phenotype is preserved across motor actions with stark differences in

complexity; Mazzoni and Krakauer (2006) studied the movement of an index finger with their arm

immobilized via splinting to a tripod, whereas the ball's path in our task was due to a complicated

whole-body coordination that imparted motion on a physical object. In our view, this speaks to the

strength and ubiquity of SPE-learning, which we speculate fine tunes our motor behaviors in

everyday life.

There are, however, some differences in the magnitude of this aftereffect between our study and

that observed by Mazzoni and Krakauer[2]. Namely, the 5° implicit aftereffect we detected was

smaller than that observed in their task, which reached 20-30°. We suspect that the aftereffects we

report in Fig. 2 largely underapproximate the total implicit learning that was achieved during the

adaptation period. First, these aftereffects were measured after a substantial break in time (about 1

minute) during which new instructions to aim toward the target were given and the prism goggles

were switched to remove the visual shift. Past studies have shown that such breaks in time lead to

decay in temporally volatile implicit learning systems[3,20,31,35]. Second, the implicit aftereffect was

measured after participants had already completed 10 trials without visual feedback, prior to the

removal of the shifted prism goggles (see the period between "remove FB" and "goggles off" in

Figs. 2A,B). Multiple studies have demonstrated that moving without error feedback also results

in a sizeable decay in implicit learning[3,8,29,36]. Thus, we strongly suspect that implicit learning is

underapproximated by the aftereffect measures in Fig. 2. This is an important limitation of our

analysis that could be improved in future work by measuring aftereffects on the very initial no-

feedback trials while also minimizing any delay between the adaptation and aftereffect periods.

Apart from these potential bias sources in our aftereffect-based implicit measures, there are critical

experimental differences between our work and that of Mazzoni and Krakauer, that could greatly

suppress the implicit learning achieved in our ball rolling task. The first factor is temporal delay;

given the ~6.1m displacement between participant and target, the ball arrived approximately 0.75

sec after the motor action had completed. These conditions are known to strongly attenuate implicit

learning[8,17,18], likely due to the misalignment between motor and error signals in the cerebellar

cortex[37–39]. Future studies could boost implicit learning by (1) decreasing the distance between the

target and participant, (2) requiring faster rolling speeds, (3) or using virtual reality to reduce error

latency. The second factor is target error, which is also known to increase implicit learning[1,3,10,40].

In Mazzoni and Krakauer (2006), learning occurred slowly without a secondary target (see Fig.

4B, no aiming-target), leading to sustained target errors that were at least 10° or larger throughout

the adaptation period. On the other hand, no-aiming-target learning in our task was rapid, reducing

the target error to only about 2° within 10 trials or so (Fig. 4A, no-aiming-target). This rapid

learning was likely due to strategy use, which was high in our task, about 25° (80% of the total

perturbation size). We hypothesize that large explicit strategies deprived the implicit system of

target errors between the ball and the task-related target, which would have provided a learning

substrate that enhanced the implicit aftereffect. These ideas could be further investigated in future

studies by reducing explicit strategies (keeping target errors large) by adding more training

targets[41], or limiting movement preparation time[3,8,25,26,42,43].

Reduction in implicit adaptation in our experiment could also be inferred based on the time course

of overcompensation detected during the aiming-target condition (Fig. 4A, red). Whereas motor

behaviors drifted further and further beyond the primary target in Mazzoni and Krakauer (see Fig.

4B, aiming-target), throwing angles in our task (Fig. 4A, aiming-target) remained fixed at nearly

a constant 3° overcompensation across the entire adaptation period. This attenuation in drift in our

task is entirely consistent with a reduction in the total extent of implicit learning. Indeed, Taylor

and Ivry (2011) demonstrated that the extent of overcompensation in an aiming-target condition is

directly linked to the extent of implicit learning (Fig. 4C). By providing an aiming target during

the entire movement (full-aiming-target), removing it after movement start (partial-aiming-target),

or never showing the aiming target during adaptation (no-aiming-target), Taylor and Ivry showed

that the extent of overcompensation could be modulated: the more salient the presence of an aiming

target, the greater the drift. These variations in overcompensations were directly linked to changes

in implicit aftereffects measured following the adaptation period (not shown in Fig. 4C): the more

salient the aiming target during learning, the greater the implicit aftereffect. This again is consistent

with past work indicating that SPEs are reinforced by visual target error[1,3,10,40].

Figure 4. *Implicit overcompensation occurs in several experimental contexts.* **A.** Trial-by-trial error data from Figure 2A,B with actual target error appearing on the y-axis. Providing an aiming target (in red) caused the angular ball error to drift beyond the target (i.e., past zero), thus overcompensating for the visual rotation. **B.** Results from Mazzoni and Krakauer (2006), an analogous lab-based experiment, which have been widely cited as evidence for SPE learning. Participants were either given no instruction (blue) or were instructed to use an aiming target to "solve" the rotation and reduce error (red). **C.** Results from Taylor & Ivry (2011), a modified version of the experiment from B. Participants were either given an aiming target during the entire movement (red), an aiming target that disappeared when the movement began (black), or taught how re-aim their movement to counter the rotation without an additional target (blue). Note the graded levels of overcompensation that correspond to the salience of the visual aiming target.

Another possible force that attenuates potential drift is a change in explicit strategy. That is, aim

reporting protocols have demonstrated late reductions in explicit strategy during training, which

presumably functions to prevent the overcompensation that can result from continued SPE-driven

implicit learning[22,23]. Similarly, Taylor and Ivry (2011) demonstrated that drift beyond the target

can be eliminated with long exposure to the rotation (Fig. 4C, full-aiming-target). The notion that

the explicit system is downregulated over time to counteract SPE-driven implicit learning would

be consistent with the order effects we observed in Fig. 3: namely, that initial adaptation rates were

greater during the second exposure to the rotation due to savings in explicit strategy, but no change

in the terminal residual error could be detected. This hypothesis could be examined in future work

by measuring implicit and explicit learning at various points throughout adaptation. Indeed, both

our experiment and the other studies illustrated in Fig. 4 provide very limited insight into the time

course of implicit learning and explicit strategy, because these are only measured after the rotation

period had ended. Measuring the time course of implicit and explicit compensation as they evolve

is critically needed to inform our qualitative and quantitative models of SPE-driven adaptation. It

is also needed to assess potential limitations in our assumption that implicit and explicit learning

are 'perfectly additive' processes[20], which may be violated under certain circumstances.

In sum, our work demonstrates that SPE-driven implicit adaptation is not a theoretical construct

observable only for constrained motor actions studied in the laboratory. Rather, SPEs are potent

drivers of adaptation even during movements that require sophisticated coordination across our

entire body as we execute common motor skills. Substantial work remains to better elucidate how

SPE-driven implicit learning can be strengthened or attenuated (e.g., via target errors), how

implicit processes are coordinated with explicit strategy, and how these two systems evolve over

time. Another important future direction will be the application of motion capture to more deeply

investigate whether markers of implicit and explicit adaptation can be differentiated across distinct

movement components. Initial motion capture efforts have identified a connection between motor

learning and whole-body motor adjustments, along with correlations between kinematic variability

and learning rates[11]. It may be that implicit and explicit processes are differentially expressed

across distinct motor control variables (e.g., trunk orientation, ball release timing and variability,

wrist and elbow joint angles).

Methods

Ethics statement. Human participant research was completed with the approval and oversight of

the Rhodes College Institutional Review Board.

Subject recruitment. Thirty study participants were recruited by word of mouth, and each given a

$10 gift card for their participation (mean age = 20yrs; age range – 18 – 21; 21 female; 29 right

handed but all completed ball throws with right). Data from three participants were omitted

because of technical issues that resulted in missing video files that contained motor error data.

Written informed consent was obtained from each participant before completing the experiment.

Experimental setup. Participants stood behind a line marked on the floor facing a small target

(D=0.85°) on the ground 6.1m away. They were informed about the task parameters: to use an

underhand motion to roll a tennis ball so that it moved through the target. The experiment took

place in a black box theater with black flooring and a black curtain behind the target area under

which balls rolled out of view. Participants started by wearing goggles with clear lenses that

allowed for unperturbed vision (Fork in the Road Vision Rehabilitation Services LLC, Madison,

WI). During shifted vision blocks, participants wore Perception (Distortion) goggles

(WinginItProducts, Pleasant Prairie, WI) that caused the participants to perceive visual targets

offset in space by 30° from their true position. Both types of goggles occluded the view of the

body and throwing arm and participants were asked to release the tennis ball below their waist

level so the hand was never visible.

An infrared video camera (Prime Color, Optitrack) was mounted directly above the target (~7

meters above attached to a theater lighting grid). The camera was pointed at the target, capturing

several meters of linear space on both sides of the target. It recorded the tennis ball rolling along

the floor so that rolling error could be computed (see Data analysis). The camera was attached to

a computer which displayed and recorded the video in Motive software (Optitrack). A single

eStrobe (Optitrack) mounted above the target (next to the camera) provided the only illumination

in the room and could be toggled on and off with a single button press in the Motive software. The

camera system did not record any motion tracking of participant kinematics.

A researcher kneeled next to the participant and handed tennis balls to the participant for each

subsequent roll. Another researcher stood behind the curtain and collected tennis balls to make

sure they did not roll back onto the visible floor space. A third researcher monitored the video

recording and controlled the lights during no feedback trials. For some participants, only two

researchers were present and the tennis ball wrangling and computer monitoring were done by the

same individual.

Data collection. Each participant completed two blocks of 192 tennis ball rolls. In the 'No

Instructions' block, participants were not given a strategy to counteract the visual perturbation. In

the 'Aiming Target' block, participants were given a secondary target to use that would minimize

the error resulting from the visual perturbation. The order of the blocks was randomly assigned to

the participants (after the three excluded participants, 13 participants completed 'No Instructions'

first and 14 participants completed 'Aiming Target' first).

Participants first completed 40 baseline movement trials, rolling the ball to the target without

shifted vision. Then participants switched to goggles that shifted their vision by 30°. To switch the

goggles, participants first closed their eyes. The researcher then assisted with doffing the current

goggles and donning the new goggles. With eyes still closed, participants were then guided through

a series of in-place rotations in order to reset the relative orientation between the target and the

body. Then participants opened their eyes.

In the 'No Instructions' block, participants then completed two trials with shifted vision after

which the researcher said: "Try to hit the target.". Participants then completed another 70 trials

with the shifted visual feedback.

In the 'Aiming Target' block, participants completed two trials with shifted vision after which

the researcher placed an aiming target, matching the size and appearance of the original target,

30° to the right of the original target. The researcher then explained to the participant: "You just

made two large errors because we imposed a rotation that pushes you 30 degrees to the left. You

can reduce this error by aiming at the new target to your right."

After 72 total trials with the shifted goggles, participants then completed 10 trials with the same

visual shift but with no endpoint error feedback. Participants were made aware before these trials

that the light would turn off immediately after they released the ball. The researcher controlling

the computer manually shut off the eStrobe light as soon as the tennis ball first hit the floor at the

start of the participant's roll, making the room completely dark. Since the camera was imaging in

the infrared spectrum, the tennis ball could still be viewed from the computer. As soon as the tennis

ball passed the target and went out of view underneath the curtain, the researcher turned on the

light.

After 10 trials with no endpoint visual feedback, the goggles were switched back to the clear lens

goggles and participants were told: "I would like you to aim for the original target again."

Participants completed 10 additional trials without endpoint visual feedback with no vision shift.

Then participants completed 60 trials with no visual shift with endpoint visual feedback (i.e. the

lights stayed on).

Data analysis. Video recordings were analyzed using Vernier Video Analysis software (Vernier

Science Education, Beaverton, OR). Small pieces of tape on the ground set at 1m apart and visible

in the video frame but not visible to participants were used to calibrate position measurements in

the software. When the tennis ball was in line with the target, its position was marked manually

with a mouse click in the software for each trial. The lateral error between the target and tennis

ball was computed for the entire set of trials using the marked positions on the video frames.

Data were loaded into MATLAB software and a custom script was used to calculate descriptive

statistics and identify the steady-state trials during the visual rotation block after which learning

had stabilized[19]. Total Adaptation was calculated as the sum of the visual shift (30°) and the mean

error over the last three trials of the shifted vision with feedback trials [i.e. how much movements

were adapted under the visual shift]. Implicit adaptation was calculated as the mean error over the

first three trials immediately after switching to unshifted vision (after the shifted vision block).

Explicit adaptation was computed as the difference between Total Adaptation and Implicit

Adaptation. Graphical plots were generated in R and MATLAB supported by publicly available

scripts[44,45].

A multiple linear regression was run in RStudio (v2022.07.1, R version 4.1.1) with Total

Adaptation and Implicit Adaptation as the outcome variables and Block Number and Strategy

Condition (i.e. with or without aiming target) as the predictor variables. Explicit Adaptation was

not included in the regression since it is computed from the included outcome variables and thus

correlated with them. Secondary inferential statistical analyses were conducted to further explain

the results. Bonferroni-corrected p-values from t-tests were used to describe pairwise comparisons

that appear in Fig. 2C,D. The same analysis was run on pairwise comparisons with data separated

into the four different blocks (Fig. 3). The mean tennis ball error during the last three trials of the

rotated vision trials with feedback was analyzed using a one sample t-test to determine if the ball's

error was beyond the physical extent of the target (i.e. including the radius of the target and ball).

To investigate savings (and by extension, the influence of an explicit strategy), a paired t-test was

run comparing initial error on the first two visual shift trials between Block 1 and Block 2. To

investigate individual correlations between total and implicit adaptation rates, Pearson's

correlation coefficients were computed for no-aiming-target and aiming-target data.

Data and code. All data, analysis code and plotting scripts are available here:

https://osf.io/8kf59/?view_only=cfeb2f109d304994bbd3a5f5a44c11ae

Acknowledgments

400 We thank Jenna Floyd, Parker Hill, Erin Kuylenstierna, and Brittney Stone for assistance with data

401 collection. We thank Dr. Shubho Banerjee for helpful data analysis suggestions. Funding provided

402 by a Harrison McCain Emerging Scholar Award (Blustein).

References

1. Tsay, J. S., Haith, A. M., Ivry, R. B. & Kim, H. E. Interactions between sensory prediction

error and task error during implicit motor learning. *PLoS Comput Biol* **18**, e1010005 (2022).

2. Mazzoni, P. & Krakauer, J. W. An Implicit Plan Overrides an Explicit Strategy during

Visuomotor Adaptation. *J Neurosci* **26**, 3642–3645 (2006).

3. Albert, S. T. *et al.* Competition between parallel sensorimotor learning systems. *eLife* **11**,

e65361 (2022).

4. Wong, A. L. & Shelhamer, M. Using prediction errors to drive saccade adaptation: the

implicit double-step task. *Exp Brain Res* **222**, 55–64 (2012).

5. Tseng, Y., Diedrichsen, J., Krakauer, J. W., Shadmehr, R. & Bastian, A. J. Sensory

Prediction Errors Drive Cerebellum-Dependent Adaptation of Reaching. *J Neurophysiol* **98**,

54–62 (2007).

6. Shadmehr, R., Smith, M. A. & Krakauer, J. W. Error Correction, Sensory Prediction, and

Adaptation in Motor Control. *Annu Rev Neurosci* **33**, 89–108 (2010).

7. Taylor, J. A. & Ivry, R. B. Flexible Cognitive Strategies during Motor Learning. *PLoS

Comput Biol* **7**, e1001096 (2011).

8. Albert, S. T. *et al.* An implicit memory of errors limits human sensorimotor adaptation. *Nat

Hum Behav* **5**, 920–934 (2021).

9. Herzfeld, D. J., Vaswani, P. A., Marko, M. K. & Shadmehr, R. A memory of errors in

sensorimotor learning. *Science* **345**, 1349–1353 (2014).

10. Miyamoto, Y. R., Wang, S. & Smith, M. A. Implicit adaptation compensates for erratic

explicit strategy in human motor learning. *Nat Neurosci* **23**, 443–455 (2020).

11. Haar, S., van Assel, C. M. & Faisal, A. A. Motor learning in real-world pool billiards. *Sci

Rep* **10**, 20046 (2020).

12. van der Kamp, J., Steenbergen, B. & Masters, R. S. W. Explicit and implicit motor learning

in children with unilateral cerebral palsy. *Disabil Rehabil* **40**, 2790–2797 (2018).

13. Leukel, C., Gollhofer, A. & Taube, W. In Experts, underlying processes that drive

visuomotor adaptation are different than in Novices. *Front Hum Neurosci* **9**, (2015).

14. Martin, T. A., Keating, J. G., Goodkin, H. P., Bastian, A. J. & Thach, W. T. Throwing while

looking through prisms: I. Focal olivocerebellar lesions impair adaptation. *Brain* **119**, 1183–

1198 (1996).

15. Leow, L.-A., Marinovic, W., de Rugy, A. & Carroll, T. J. Task errors contribute to implicit

aftereffects in sensorimotor adaptation. *Eur J Neurosci* **48**, 3397–3409 (2018).

16. Ruttle, J. E., 't Hart, B. M. & Henriques, D. Y. P. Implicit motor learning within three trials.

Sci Rep **11**, 1627 (2021).

17. Brudner, S. N., Kethidi, N., Graeupner, D., Ivry, R. B. & Taylor, J. A. Delayed feedback

during sensorimotor learning selectively disrupts adaptation but not strategy use. *J*

Neurophysiol **115**, 1499–1511 (2016).

18. Schween, R. & Hegele, M. Feedback delay attenuates implicit but facilitates explicit

adjustments to a visuomotor rotation. *Neurobiol Learn Mem* **140**, 124–133 (2017).

19. Blustein, D., Shehata, A., Englehart, K. & Sensinger, J. Conventional analysis of trial-by-

trial adaptation is biased: Empirical and theoretical support using a Bayesian estimator. *PLoS*

Comput Biol **14**, e1006501 (2018).

20. Maresch, J., Werner, S. & Donchin, O. Methods matter: Your measures of explicit and

implicit processes in visuomotor adaptation affect your results. *Eur J Neurosci* **53**, 504–518

(2021).

21. Vaswani, P. A. *et al.* Persistent Residual Errors in Motor Adaptation Tasks: Reversion to

Baseline and Exploratory Escape. *J Neurosci* **35**, 6969–6977 (2015).

22. Taylor, J. A., Krakauer, J. W. & Ivry, R. B. Explicit and Implicit Contributions to Learning

in a Sensorimotor Adaptation Task. *J. Neurosci.* **34**, 3023–3032 (2014).

23. McDougle, S. D., Bond, K. M. & Taylor, J. A. Explicit and Implicit Processes Constitute the

Fast and Slow Processes of Sensorimotor Learning. *J Neurosci* **35**, 9568–9579 (2015).

24. Huang, V. S., Haith, A., Mazzoni, P. & Krakauer, J. W. Rethinking Motor Learning and

Savings in Adaptation Paradigms: Model-Free Memory for Successful Actions Combines

with Internal Models. *Neuron* **70**, 787–801 (2011).

25. Huberdeau, D. M., Krakauer, J. W. & Haith, A. M. Practice induces a qualitative change in

the memory representation for visuomotor learning. *J Neurophysiol* **122**, 1050–1059 (2019).

26. Haith, A. M., Huberdeau, D. M. & Krakauer, J. W. The Influence of Movement Preparation

Time on the Expression of Visuomotor Learning and Savings. *J Neurosci* **35**, 5109–5117

(2015).

27. Morehead, J. R., Qasim, S. E., Crossley, M. J. & Ivry, R. Savings upon Re-Aiming in

Visuomotor Adaptation. *J Neurosci* **35**, 14386–14396 (2015).

28. Diedrichsen, J., White, O., Newman, D. & Lally, N. Use-Dependent and Error-Based

Learning of Motor Behaviors. *J Neurosci* **30**, 5159–5166 (2010).

29. Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R. & Ivry, R. B. Invariant errors reveal

limitations in motor correction rather than constraints on error sensitivity. *Commun Biol* **1**,

19 (2018).

30. Tsay, J. S., Kim, H., Haith, A. M. & Ivry, R. B. Understanding implicit sensorimotor

adaptation as a process of proprioceptive re-alignment. *eLife* **11**, e76639 (2022).

31. Neville, K.-M. & Cressman, E. K. The influence of awareness on explicit and implicit

contributions to visuomotor adaptation over time. *Exp Brain Res* **236**, 2047–2059 (2018).

32. Hadjiosif, A. M. & Smith, M. A. Flexible Control of Safety Margins for Action Based on

Environmental Variability. *J Neurosci* **35**, 9106–9121 (2015).

33. Morehead, J. R. & Xivry, J.-J. O. de. A Synthesis of the Many Errors and Learning Processes

of Visuomotor Adaptation. *bioRxiv* doi:10.1101/2021.03.14.435278 (2021).

34. Orozco, S. P., Albert, S. T. & Shadmehr, R. Adaptive control of movement deceleration

during saccades. *PLoS Comp Biol* **17**, e1009176 (2021).

35. Hadjiosif, A. M., Morehead, J. R. & Smith, M. A. A double dissociation between savings

and long-term memory in motor learning. *PLoS Biol* **21**, e3001799 (2023).

36. Alhussein, L., Hosseini, E. A., Nguyen, K. P., Smith, M. A. & Joiner, W. M. Dissociating

effects of error size, training duration, and amount of adaptation on the ability to retain motor

memories. *J Neurophysiol* **122**, 2027–2042 (2019).

37. Ekerot, C.-F. & Kano, M. Stimulation parameters influencing climbing fibre induced long-

term depression of parallel fibre synapses. *Neurosci Res* **6**, 264–268 (1989).

38. Suvrathan, A., Payne, H. L. & Raymond, J. L. Timing Rules for Synaptic Plasticity Matched

to Behavioral Function. *Neuron* **92**, 959–967 (2016).

39. Herzfeld, D. J., Kojima, Y., Soetedjo, R. & Shadmehr, R. Encoding of error and learning to

correct that error by the Purkinje cells of the cerebellum. *Nat Neurosci* **21**, 736–743 (2018).

40. Kim, H. E., Parvin, D. E. & Ivry, R. B. The influence of task outcome on implicit motor

learning. *eLife* **8**, e39882 (2019).

41. Tsay, J. S. *et al.* Predictors of sensorimotor adaption: insights from over 100,000 reaches.

bioRxiv doi:10.1101/2023.01.18.524634 (2023).

42. McDougle, S. D. & Taylor, J. A. Dissociable cognitive strategies for sensorimotor learning.

Nat Commun **10**, 40 (2019).

43. Fernandez-Ruiz, J., Wong, W., Armstrong, I. T. & Flanagan, J. R. Relation between reaction

time and reach errors during visuomotor adaptation. *Behav Brain Res* **219**, 8–14 (2011).

44. Campbell, R. sigstar. *GitHub* https://github.com/raacampbell/sigstar, (2023).

45. Fachada, N. & Rosa, A. micompm: A MATLAB/Octave toolbox for multivariate

independent comparison of observations. *J Open Source Softw* **3**, (2018).

46. Albert, S. T. *et al.* Postural control of arm and fingers through integration of movement

commands. *eLife* **9**, e52507 (2020).