The song does not remain the same: daily singing of adult songbirds

prevents passive changes in song structure independently of auditory

feedback

Daisuke Mizuguchi[a], Miguel Sánchez-Valpuesta[a], Yunbok Kim[a], Ednei B. dos Santos [a], HiJee

Kang[a], Chihiro Mori[b, †], Kazuhiro Wada[c], and Satoshi Kojima[a,*]

[a] Sensory and Motor Systems Research Group, Korea Brain Research Institute, 61,

Cheomdan-ro, Dong-gu, Deagu, 41062, Republic of Korea

[b] Department of Life Sciences, Graduate School of Arts and Sciences, University of Tokyo,

Tokyo, Japan

[c] Department of Biological Sciences, Faculty of Science, Hokkaido University, Sapporo, 060-

0810, Hokkaido, Japan

* Correspondence: Satoshi Kojima, satoshikojima.sk@gmail.com

[†] Current affiliation: Faculty of Pharmaceutical Sciences, Department of Life and Health

Sciences, Teikyo University

Author Contributions: D.M. and S.K. designed and performed the research; D.M., M.S.V.,

and E.B.D.S., analyzed the data; and D.M., M.S.V., E.B.D., Y.K., H.K., C.M., K.W., and S.K.

wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: zebra finch | song | maintenance | vocal training | auditory feedback

Abstract

Many songbirds learn to produce songs through vocal practice early in life and continue to sing numerous renditions of their learned songs daily throughout their lifetime. While it is well-known that adult songbirds sing as part of a mating ritual, the functions of singing behavior outside of reproductive contexts are not fully understood. Here, we demonstrate that adult singing outside of reproductive contexts serves to prevent passive changes in song performance. We suppressed the daily singing behavior of adult zebra finches produced in the solo context for two weeks using a reversible behavioral manipulation and examined the effect on song performance. Our results indicated that suppressing daily singing significantly decreased the pitch of song elements and both the amplitude and duration of song motifs. These findings suggest that adult song is not acoustically stable without singing, and that adult birds maintain their song performance by daily singing. Moreover, we found that the changes in song structure caused by singing suppression were substantially recovered within two weeks of free singing, even in deafened birds. Thus, the recovery of song performance does not necessarily require auditory feedback but is predominantly caused by singing behavior per se (i.e., the physical act of singing). Finally, unlike the auditory feedback-dependent song plasticity reported previously, the passive song changes caused by singing suppression were not significantly dependent on age. Taken together, our findings demonstrate that adult songbirds maintain song performance by preventing passive song changes through the physical act of daily singing throughout their life. Such daily singing likely functions as vocal training to maintain the neural and/or muscular system in optimal conditions for song performance in reproductive contexts, similar to how human singers and athletes practice daily to maintain their performance.

Introduction

Singing behavior is a highly complex motor skill that has been extensively studied across many taxa, including birds, mice, whales, and primates including humans[1]. In songbirds, the primary function of singing is mate attraction and territorial defense in reproductive contexts[2]. Singing behavior in reproductive contexts is observed in the presence of apparent recipients, and thus the purpose of singing can be easily determined by the response of the recipients. In contrast, the functions of singing outside reproductive contexts, produced even when no apparent recipients are around, are much less understood. Although this type of singing behavior has been suggested to play important roles in vocal learning and song structure maintenance[3–5], the exact function of singing outside reproductive contexts remains controversial[6–9].

The zebra finch (*Taeniopygia guttata*) is a non-territorial songbird commonly used for neuroethological studies of birdsong. Male zebra finches learn to produce a song during a limited period in early life (closed-ended learner[10]). As adults (>90 days post-hatch [dph]) they repeatedly produce thousands of renditions of the learned song daily throughout life, even when isolated from other birds (i.e., outside reproductive contexts)[11–13]. This daily singing in

the solo context, generally called "undirected singing", has been shown to have a function to refine and maintain song structure: through undirected singing, birds evaluate the song structure using auditory feedback and refine it if necessary[3,4,14–17] (although other functions of undirected song have been suggested[7–9]). This auditory feedback-dependent function of undirected singing, however, appears to be limited to young adult birds, as the removal of auditory feedback causes substantial changes in song structure in relatively young birds but not in old birds[16,17]. Nevertheless, old birds still sing many renditions of undirected songs every day[12,18], raising the possibility that undirected singing serves an unknown function in addition to the auditory feedback-dependent refinement of song structure. To explore novel functions of daily undirected singing, we suppressed undirected singing of adult zebra finches for two weeks (Fig. 1A) using a method to physically interfere with their singing posture[19] and examined how their song performance was affected.

Results

We compared undirected songs between one day before (pre1d) and one day after (post1d) the singing suppression (SS) period (Fig. 1A) in young adult birds (96-129 dph), and found small but significant changes in song structure following the SS period: Although the overall song acoustic structure was maintained (Fig. 1B), the pitch of harmonic syllables significantly decreased compared control birds, which were allowed to freely sing for 2 weeks (post1d in Figs. 1C, E; n = 10 SS-treated birds and 9 control birds; $p = 0.002$, two-way ANOVA and post-hoc Bonferroni's multiple comparisons test, see Table 1 for the results of two-way ANOVA). Also, significant decreases were observed in both the root mean square (RMS) amplitude and mean duration of song motifs (a stereotyped sequence of syllables) (post1d in Figs. 1D, F, G; $p = 0.029$ for motif amplitude; $p = 0.003$ for motif duration). The decrease in motif duration was predominantly attributable to the decrease in syllable duration rather than in silent gap duration (Fig. 1H; $p = 0.0024$ and $W = 75$ for syllable duration and $p = 0.57$ and $W = 31$ for gap duration, Wilcoxon signed-rank test). Moreover, those syllable and motif structures tended to return to their pre-SS levels after 2 weeks of free singing following the SS period, suggesting reversible effects of SS on song structure (post2w in Fig. 1E-G; $p = 0.11$ and 0.25 for syllable pitch and motif duration, respectively; the results of motif amplitude were barely significant [$p = 0.049$]). These results provide evidence that neither the syllable nor motif structures of adult songs are stable without daily undirected singing, and suggest that undirected singing has a function to prevent passive changes in song performance to maintain it in the optimal condition.

Undirected singing has been previously shown to function in refinement and maintenance of song structure by providing auditory feedback, through which the birds evaluate their own song structure and correct vocal errors[3,4,14,15]. We wondered whether similar, auditory feedback-dependent mechanisms mediate the recoveries of syllable pitch and motif structure after SS. To examine this, we deafened birds to remove auditory feedback immediately before the SS period and examined whether their song recovery following SS

was affected (Fig. 2*A*). Despite the lack of auditory feedback, the deafened birds showed
reversible changes in syllable pitch and motif amplitude similar to those observed in intact-
hearing birds (Fig. 2*B-C*; n = 6 SS-treated and deafened birds and 9 control [freely singing
and intact hearing] birds; for syllable pitch, *p* = 0.013 at post1d and >0.99 at post2w; for motif
amplitude, *p* = 0.0001 at post1d and >0.99 at post2w; see Table 1 for the results of two-way
ANOVA). A similar trend was observed for motif duration as well, although the results were
not significantly different (Fig. 2*D*; *p* = 0.8164 at post1d and 0.2197 at post2w). Also, there
was a similar trend for the decreases in syllable duration and silent gap duration as well (Fig.
2*E*), although neither decrease was statistically significant (*p* = 0.44 and *W* = 15 for syllable
duration; *p* = 0.69 and *W* = 8 for gap duration). Thus, the recovery of song performance after
SS does not necessarily require auditory feedback, and is likely to be caused mainly by
singing behavior per se (i.e., the physical act of singing). These results suggest an auditory
feedback-independent function of undirected singing to maintain song performance.

In zebra finches, the auditory feedback-dependent function of undirected singing to
maintain song structure strongly depends on age: the removal of auditory feedback causes
substantial changes in song structure in young adult birds (< ~200 dph) but not in older
birds[16,17]. Given that old birds still sing many renditions of undirected songs every day
(although the number of undirect songs decreases with age)[12,18], it is possible that passive
changes in song structure can occur independently of age and that those changes are
prevented by undirected singing throughout life. To test this possibility, we examined the
effect of SS on song structure in relatively old birds with intact hearing (396–809 dph, n = 6
birds) and combined their data with the data of young adult birds shown in Fig. 1*E-G*, to ask
whether SS-induced song changes depend on age. In contrast to auditory feedback-
dependent song plasticity previously reported[16,17], we found no significant effects of age on
SS-induced song changes (Fig. 3; *r* = -0.018, *p* = 0.94 for syllable pitch; *r* = 0.022, *p* = 0.93 for
motif amplitude; *r* = 0.17, *p* = 0.49 for motif duration). Thus, passive changes in adult song
structure can occur regardless of age, suggesting that birds produce daily undirected songs to
prevent passive song changes throughout life.

Because our procedure of SS is likely to induce substantial physical stress in the
birds, we cannot exclude the possibility that the passive song changes observed after SS
period were predominantly caused by the physical stress induced by the SS procedure rather
than the lack of undirected singing. To examine this possibility, we measured plasma
corticosterone (CORT) levels in a different group of birds before and during the SS period
(see Methods for more detail). We found that many birds increased CORT levels on the first
day of the SS period to levels similar to those induced by restraint stress reported in previous
studies[20,21] and subsequently decreased toward the pre-SS levels (Fig. 4*A*; *n* = 10 birds). Also,
the increases in CORT levels on the first day of SS period were not significantly correlated
with SS-induced changes in the song features that we examined (syllable pitch, motif
amplitude, or motif duration) (Fig. 4*B*; *n* = 10 birds). These results support the notion that SS-
induced song changes are mainly attributed to the lack of undirected singing rather than the

physical stress caused by SS procedure. Moreover, although many birds that received SS
exhibited weight loss, an indirect indicator of stress level, no significant correlations were
found between the weight changes and the song changes (Fig. 4C; the song data in Figs 1 &
2 were replotted to be compared with weight changes), further supporting the independence
of SS-induced song changes from physical stress. These results are also consistent with a
previous study that the pitch and amplitude of undirected songs are significantly affected by
food restriction only when food is restricted in early life but not in adulthood[22]. In summary, the
song changes induced by SS are most likely caused by the lack of singing behavior, rather
than stress.
153
Discussion
In summary, we found that suppression of undirected singing in adult zebra finches
changes both syllable and motif structures of songs, which can be recovered within a short
period of free singing. This is the first study in a closed-ended learner showing that the adult
song can substantially change over time in the absence of singing behavior, revealing a novel
function of undirected singing to prevent passive song changes. Moreover, the recovery of
song performance following the SS period does not necessarily require auditory feedback,
highlighting the importance of the physical act of singing in the maintenance of adult song
performance. Finally, in contrast to song plasticity reported in previous studies[16,17], SS-
induced song changes do not depend on age, providing a plausible explanation for why adult
birds sing many renditions of the undirected song throughout life even in old age. Taken
together, our findings provide evidence that zebra finches maintain their song performance by
preventing passive song changes through the physical act of daily singing throughout life.
Because male zebra finches produce courtship songs with the structure that is virtually the
same as that of undirected song[10], it is likely that they maintain the song production system in
the optimal condition through undirected singing to prepare for the best song performance in
reproductive contexts.
We previously showed that temporary suppression of undirected singing (for 5 or 10
h) in adult zebra finches induces intense singing immediately after the SS period, indicating
that intrinsic motivation for undirected singing increases while singing is suppressed[23]. This
SS-induced increase in singing motivation and resulting intense singing can be explained by
the song maintenance function of undirected singing that we found in the present study. It is
reasonable to assume that birds produce intense singing after SS to compensate for the loss
of vocal practice during SS and to quickly re-optimize the vocal system and song structure for
future courtship activity. Although it remains unclear whether adult song structure actually
deteriorates during such short SS periods and recovers through subsequent intense singing,
substantial song deterioration during non-singing periods and improvement during
subsequent singing are observed in juvenile birds producing immature songs[11,13]. Moreover,
this idea of vocal re-optimization through intense singing after SS is consistent with a
hypothesized role of the "dawn chorus", intense singing behavior at dawn widely observed in

many songbird species[24,25]. Because dawn chorus occurs after a long break of daytime
singing by the night and is accompanied by significant improvement in song structure[26,27], it
has been hypothesized that through intense dawn singing, birds warm up their vocal system
to prepare for future courtship and territorial defense activities just as human athletes and
singers do before their performance[26,27]. Our results in the present and previous studies
provide empirical support for this warm-up hypothesis of dawn chorus, advancing our
understanding of the functional roles of non-reproductive singing in wild birds as well as in
captive birds.
While our findings suggest that the limited singing behavior of adult birds results in
substantial changes in the song performance, the physiological mechanisms underlying such
passive song changes should be further studied. It is presumably caused, at least in part, by
weakening (atrophy) of the syringeal muscles responsible for song production, similar to the
atrophy of skeletal muscles that is generally known to be caused by a lack of physical activity.
Comparing the syringeal muscle volumes or fiber sizes before and after SS would be
necessary to examine any possible muscle atrophy. Given the decreasing effect of SS on
motif and syllable duration, it is also possible that the neural circuitry that generates song
structure passively drifts without daily undirected singing. Consistent with this idea,
substantial changes in singing-related neural activity in the song motor pathway occur
primarily during the nighttime (when birds do not sing), and less frequently during the
daytime[28,29]. Also, changes in motif duration are observed even in birds that freely sing when
the central neural circuits are manipulated[30,31], further supporting the possibility that SS-
induced song changes are attributable, at least in part, to passive changes in neural circuits.
Comparing singing-related neural activity patterns before and after SS would be necessary to
examine this possibility. Regardless of the mechanisms of passive song changes induced by
the lack of daily undirected singing, our findings open the possibility that the physical activity-
dependent and hearing-independent mechanisms for maintaining vocal performance are
commonly used in many other bird species and other animal taxa, including non-vocal
learning birds and mammals.
212
Materials and Methods
Subjects. Adult male zebra finches (98–809 dph) were bred in our colony at Korea Brain
Research Institute and individually housed in sound-attenuating chambers (MC-050,
Muromachi Kikai, or custom-made ones) on a 14:10-h light:dark cycle throughout the
experiments. Their care and treatment were reviewed and approved by the Animal Care and
Use Committee at Korea Brain Research Institute.
219
Song recording. Songs were recorded using a microphone (PRO35, Audio-Technica)
positioned 5~10 cm above the cage (20 x 20 x 20 cm) and a custom-made recording program
at a 32-kHz sampling rate with 16-bit resolution. To minimize the influence of the bird's
position on the amplitude of song recording, no perch was placed in the cage and the food

was provided in a petri dish placed on the bottom of the cage so as to have the bird sing at
the cage bottom (zebra finches rarely sing while holding on to the side walls or ceiling of the
cage). Song recording was triggered when the program detected 4 or 5 successive song
elements, each of which reached predefined thresholds of amplitude and duration, and ended
when a silent period lasted >0.5 sec. Our recording system thus captured mostly the song
bouts, although short calls (and cage noises) were occasionally recorded. We counted the
number of song bouts daily to monitor singing activity (Supplementary Tables 1-2). Birds that
produced songs containing at least one harmonic syllable with sufficient singing rates (>200
undirected song bouts per day) were used for experimentation.

Singing suppression. Spontaneous undirected singing was suppressed for two weeks using
a previously published method[19]. Briefly, a custom-made weight was attached to the neck of
the bird each morning to interfere with his singing posture to prevent him from singing (the
weight shifted the posture toward an inferior position). The weight (20.1-32.6 g) was adjusted
daily for each bird so that the procedure mostly suppressed undirected singing but did not
severely affect the daily behaviors of the bird, such as eating, drinking, preening, and calling
(body weight and health conditions were carefully monitored every day). Each night, the
weight was removed when the light was turned off as no singing behavior was observed
during the night. With this method, the daily singing amount (the number of song bouts per
day) decreased to less than 10% of that of pre-SS (see Supplementary Tables 1-2). Birds in
the control group were allowed to sing freely without a weight all time.

Song analysis. To examine the effects of SS on song performance, we compared the
acoustic and temporal structure of songs recorded 1 day before SS (pre1d) with songs
recorded 1 day and 2 weeks after SS (post1d and post2w, respectively; Fig. 1*A*). Adult zebra
finch's songs consist of stereotyped sequences of several syllables, which are termed
"motifs," in which syllables are separated by silent gaps (see Fig. 1*B*). To test the effect of SS
on the pitch of individual syllables, >30 motifs from pre1d, post1d, and post2w songs were
randomly selected for each bird, and syllables containing clear harmonic structure were
extracted. For each syllable rendition, the pitch trajectory was obtained from a segment of a
sound waveform described in a previous study[31,32]. Briefly, spectrograms were calculated
using a Gaussian-windowed short-time Fourier transform (σ = 1 ms) sampled at 8 kHz, and a
pitch trajectory was obtained by calculating the fundamental frequency in individual time bins.
The fundamental frequency was calculated by parabolic interpolation. For each syllable, all
renditions of pitch trajectories were aligned by the syllable onsets, which were determined
based on amplitude threshold crossings, and the mean pitch values over the flat portions of
pitch trajectories were averaged across all renditions. All the birds examined in the present
study had at least one syllable with a clear harmonic structure in their song motif; for the birds
that had more than one harmonic syllable, we calculated the effects (percent changes) of SS

on the pitch of individual syllables and averaged them across the syllables for each bird (the

mean number of syllables examined per bird = 1.76; max = 3).

For the amplitude measurements, 12-20 motifs with low background noise levels from

pre1d, post1d, and post2w songs were randomly selected for each bird. We then measured

the RMS amplitude values for individual motifs and averaged them. Since we were only

interested in SS-induced changes in the motif amplitude, we did not calculate the absolute

amplitude and instead obtained percent changes relative to the amplitude of pre1d song.

Because our cage setting limited birds' singing sites to the bottom of the cage (see the

description of song recording above), the distance between a singing bird and the microphone

did not dramatically changes, and therefore changes in the RMS amplitude across different

time points are attributed mainly to changes in song volume rather than changes in singing

sites. One control bird was excluded from the amplitude analysis since the amplitude gain

was not consistent over the song recording due to the malfunction of the recording system.

Durations of song motifs, syllables, and silent gaps were measured by thresholding

amplitude envelopes that were normalized to their peak amplitudes (the same thresholds

were used for all the time points for each bird). The obtained song segmentations were

visually inspected and manually modified if necessary, so that syllable/gap segmentation

patterns were identical across motif renditions. Percent changes in syllable duration or in

silent gap duration were calculated for individual syllables or gaps, and then averaged across

different syllables or gaps for each bird.

Deafening. For experiments to test whether auditory feedback was related to the song

change after the SS period, we deafened young adult birds immediately before SS by bilateral

cochlea removal following a previously published method[5,18] and examined whether song

recovery following SS was affected. SS-induced changes in syllable pitch, motif amplitude

and motif duration were analyzed in the same way as normal-hearing birds. One bird was

excluded from the analysis because the song structure was severely deteriorated by

deafening.

Corticosterone analysis. To measure the stress levels of SS-treated birds, blood samples

were collected (<75 µl for each sampling) at 5-time points before and during the SS period

(pre4d, post1d, post5d, post9d, and post13d) by puncturing the brachial wing vein with a

sterile needle (23G) 1 hour after the weight attachment. The blood samples were then

centrifuged at 3500 rpm for 20 min at room temperature (approximately 20 °C). The

separated serums were stored at -20 °C until the assay. Corticosterone (CORT) levels were

measured using an Enzyme Immunoassay kits (Enzo Life Science). Two separate 96-well

plates were used in this study, and a separate standard curve was run on each plate. All

serum samples (1:20 dilution) and standards were run in duplicates. Plates were read on

Absorbance 96 plate reader (Enzo Life Sciences). CORT levels were determined from 4-PL

regression of 5 standards ranging from 32 to 20,000 pg/ml. We calculated intra-assay

variability as the average %CV between duplicates for each assay and inter-assay variability as %CV among CORT concentrations calculated from each standard curve for 5 corticosterone standards run in each assay. Average intra- and inter-assay variability was 18.0% and 4.8%, respectively.

Statistics. Statistical tests were performed using R 3.6.2 (R Core Team and R Foundation for Statistical Computing) or MATLAB (MathWorks). The effects of SS on syllable pitch and motif amplitude and duration were examined using a two-way ANOVA and post-hoc Bonferroni's multiple comparisons test. The effects of SS on syllable duration and silent gap duration were examined using a Wilcoxon signed-rank test. The relationships of song changes with age, CORT changes, and weight changes were examined using Pearson's correlation coefficients.

Data availability
All data are included in the article and supporting information.

Acknowledgments and funding sources
This work was supported by grants from Korea Brain Research Institute basic research program through Korea Brain Research Institute funded by Ministry of Science and ICT (Grant 23-BR-01-01 to S.K.). We thank Dr. Ji Young Mun, and Dr. Min Kyo Jung at Korea Brain Research Institute and Dr. Ryosuke O. Tachibana at Tokyo University for their valuable comments and discussion.

References

1. Petkov, C. I. & Jarvis, E. D. Birds, primates, and spoken language origins: behavioral phenotypes and neurobiological substrates. *Front Evol Neurosci* **4**, 12 (2012).

2. Catchpole, C. K. & Slater, P. J. B. *Bird song: Biological themes and variations, 2nd ed*. xi, 335 (Cambridge University Press, 2008). doi:10.1017/CBO9780511754791.

3. Leonardo, A. & Konishi, M. Decrystallization of adult birdsong by perturbation of auditory feedback. *Nature* **399**, 466–470 (1999).

4. Tumer, E. C. & Brainard, M. S. Performance variability enables adaptive plasticity of 'crystallized' adult birdsong. *Nature* **450**, 1240–1244 (2007).

5. Konishi, M. The role of auditory feedback in the control of vocalization in the white-crowned sparrow. *Z Tierpsychol* **22**, 770–783 (1965).

6. Riters, L. V., Spool, J. A., Merullo, D. P. & Hahn, A. H. Song practice as a rewarding form of play in songbirds. *Behav. Processes* (2017) doi:10.1016/j.beproc.2017.10.002.

7. Dunn, A. M. & Zann, R. A. Undirected Song Encourages the Breeding Female Zebra

Finch to Remain in the Nest. *Ethology* **102**, 540–548 (1996).

8. Bolund, E., Schielzeth, H. & Forstmeier, W. Singing activity stimulates partner

reproductive investment rather than increasing paternity success in zebra finches.

Behavioral Ecology and Sociobiology **66**, 975–984 (2012).

9. Loning, H., Verkade, L., Griffith, S. C. & Naguib, M. The social role of song in wild zebra

finches. *Current Biology* **33**, 372-380.e3 (2023).

10. Zann, R. A. *The zebra finch: A synthesis of field and laboratory studies.* (Oxford

University Press, 1996).

11. Ohgushi, E., Mori, C. & Wada, K. Diurnal oscillation of vocal development associated with

clustered singing by juvenile songbirds. *Journal of Experimental Biology* **218**, 2260–2268

(2015).

12. Johnson, F., Soderstrom, K. & Whitney, O. Quantifying song bout production during

zebra finch sensory-motor learning suggests a sensitive period for vocal practice.

Behavioural Brain Research **131**, 57–65 (2002).

13. Derégnaucourt, S., Mitra, P. P., Fehér, O., Pytte, C. & Tchernichovski, O. How sleep

affects the developmental learning of bird song. *Nature* **433**, 710–716 (2005).

14. Brainard, M. S. & Doupe, A. J. Auditory feedback in learning and maintenance of vocal

behaviour. *Nat Rev Neurosci* **1**, 31–40 (2000).

15. Andalman, A. S. & Fee, M. S. A basal ganglia-forebrain circuit in the songbird biases

motor output to avoid vocal errors. *Proc. Natl. Acad. Sci. U.S.A* **106**, 12518–12523

(2009).

16. Brainard, M. S. & Doupe, A. J. Postlearning consolidation of birdsong: stabilizing effects

of age and anterior forebrain lesions. *J. Neurosci* **21**, 2501–2517 (2001).

17. Lombardino, A. J. & Nottebohm, F. Age at deafening affects the stability of learned song

in adult male zebra finches. *J. Neurosci* **20**, 5054–5064 (2000).

18. Mori, C. & Wada, K. Audition-independent vocal crystallization associated with intrinsic

developmental gene expression dynamics. *J. Neurosci.* **35**, 878–889 (2015).

19. Hayase, S. *et al.* Vocal practice regulates singing activity–dependent genes underlying

age-independent vocal learning in songbirds. *PLOS Biology* **16**, e2006537 (2018).

20. Wada, H. *et al.* Adrenocortical responses in zebra finches (Taeniopygia guttata):

individual variation, repeatability, and relationship to phenotypic quality. *Horm Behav* **53**,

472–480 (2008).

21. Rensel, M. A. & Schlinger, B. A. The stressed brain: regional and stress-related

corticosterone and stress-regulated gene expression in the adult zebra finch

(Taeniopygia guttata). *J Neuroendocrinol* **32**, e12852 (2020).

22. Mishra, I. *et al.* Developmental effects of daily food availability times on song behaviour

and neuronal plasticity of song-control system in male zebra finches. *Behav. Brain Res.*

382, 112497 (2020).

23. Kim, Y., Kwon, S., Rajan, R., Mori, C. & Kojima, S. Intrinsic motivation for singing in

songbirds is enhanced by temporary singing suppression and regulated by dopamine. *Sci*

Rep **11**, 20350 (2021).

24. Gil, D. & Llusia, D. The Bird Dawn Chorus Revisited. in *Coding Strategies in Vertebrate*

Acoustic Communication (eds. Aubin, T. & Mathevon, N.) 45–90 (Springer International

Publishing, 2020). doi:10.1007/978-3-030-39200-0_3.

25. Staicer, C. A., Spector, D. A. & Horn, A. G. *24. The Dawn Chorus and Other Diel*

Patterns in Acoustic Signaling. Ecology and Evolution of Acoustic Communication in

Birds 426–453 (Cornell University Press, 1996).

26. Dinh, J. P., Peters, S. & Nowicki, S. Song performance improves with continued singing

across the morning in a songbird. *Animal Behaviour* **167**, 127–137 (2020).

27. Schraft, H. A., Medina, O. J., McClure, J., Pereira, D. A. & Logue, D. M. Within-day

improvement in a behavioural display: wild birds 'warm up'. *Animal Behaviour* **124**, 167–

174 (2017).

28. Liberti, W. A. *et al.* Unstable neurons underlie a stable learned behavior. *Nat. Neurosci.*

(2016) doi:10.1038/nn.4405.

29. Rauske, P. L., Chi, Z., Dave, A. S. & Margoliash, D. Neuronal Stability and Drift across

Periods of Sleep: Premotor Activity Patterns in a Vocal Control Nucleus of Adult Zebra

Finches. *J. Neurosci.* **30**, 2783–2794 (2010).

30. Long, M. A. & Fee, M. S. Using temperature to analyse temporal dynamics in the

songbird motor pathway. *Nature* **456**, 189–194 (2008).

31. Kojima, S., Kao, M. H., Doupe, A. J. & Brainard, M. S. The Avian Basal Ganglia Are a

Source of Rapid Behavioral Variation That Enables Vocal Motor Exploration. *J. Neurosci.*

38, 9635–9647 (2018).

32. Charlesworth, J. D., Tumer, E. C., Warren, T. L. & Brainard, M. S. Learning the

microstructure of successful behavior. *Nat. Neurosci.* **14**, 373–380 (2011).

Figure 1. Changes in song performance by suppression of daily undirected singing. (A) Schematic diagram of the experimental procedure. Undirected singing was suppressed for 2 weeks (SS) and songs recorded one day before SS (pre1d) and one day and 2 weeks after SS (post1d and post2w, respectively) were analyzed. (B) Example spectrograms of undirected song motifs at pre1d and at post1d, showing that the overall song structure was maintained. (C) Pitch trajectories of the rectangle regions shown in *B*. Thin lines indicate pitch trajectories from a single syllable rendition; thick lines indicate mean trajectories across all renditions examined; black and magenta lines indicate pre1d and post1d songs, respectively. (D) Overlaid amplitude envelopes of song motifs obtained from the bird shown in *B*. Conversions are as in *C*. (E-G) Percent changes in syllable pitch (E), motif amplitude (F), and motif duration (G) of SS-treated (red lines) and control (blue lines) birds at post1d and post2w relative to pre1d songs. Circles indicate the data of individual birds and lines indicate the mean and SEM across birds. All the song features significantly decreased at post1d in SS birds compared to those in control birds and substantially recovered at post2w (**p < 0.01, *p < 0.05). (H) Percent changes in syllable and silent gap duration of SS-treated birds at post1d relative to pre1d data. Syllable duration but not gap duration significantly changed by SS procedure (**p < 0.01).

Figure 2. Auditory feedback-independent recovery in song performance after SS through

daily undirected singing. (A) Schematic diagram of the experimental procedure. Birds were

deafened immediately before SS. (B-D) Percent changes in syllable pitch (B), motif amplitude

(C), and motif duration (D) in deafened and SS-treated birds (red lines) and control (normal-

hearing and free-singing) birds (blue lines). Conversions are as in Figs. 1*E-G*. ***$p < 0.001$, *p

< 0.05. (E) Percent changes in syllable and silent gap duration of experimental birds at post1d

relative to pre1d data.

Figure 3. Age-independent changes of the song performance by SS. (A-C) Percent changes
in syllable pitch (A), motif amplitude (B), and motif duration (C) at post1d are plotted against
the ages of birds. There were no significant effects of age on the changes in the song
features.

Figure 4. SS-induced changes in song performance are not well explained by stress or bodyweight loss. (A) Percent changes in plasma CORT levels caused by SS treatment; thin lines indicate the data of individual birds and thick lines indicate the mean (±SEM) across the birds. (B) Percent changes in syllable pitch (left), motif amplitude (middle), and motif duration (right), plotted against those in CORT levels measured on the 1st day of SS. (C) Percent changes in syllable pitch (left), motif amplitude (middle), and motif duration (right), plotted against bodyweight changes (the song data in Figs 1 & 2 were replotted). Circles and triangles indicate SS birds with normal hearing and those with hearing impaired (deafened), respectively.

454 **Table 1.** The results of a two-way repeated measures ANOVA analysis for the data in Fig.

455 1*E-G* (SS alone) and those in Fig. 2*B-D* (SS and deafening).

	Treatment		Time		Interaction	
	F_{df}	p	F_{df}	p	F_{df}	p
SS alone						
Syllable pitch	$F_{1,17} = 10.64$	**0.0046****	$F_{2,33} = 3.287$	**0.0499***	$F_{2,33} = 3.441$	**0.0439***
Motif amplitude	$F_{1,55} = 0.003$	0.9552	$F_{2,55} = 7.314$	**0.0015****	$F_{2,55} = 6.658$	**0.0026****
Motif duration	$F_{1,19} = 5.919$	**0.0250***	$F_{2,36} = 2.308$	0.114	$F_{2,36} = 4.204$	**0.0229***
SS & deafened						
Syllable pitch	$F_{1,13} = 1.681$	0.2173	$F_{2,26} = 5.061$	**0.0139***	$F_{2,26} = 4.601$	**0.0195***
Motif amplitude	$F_{1,13} = 3.820$	0.0725	$F_{2,26} = 11.24$	**0.0003*****	$F_{2,26} = 11.42$	**0.0003*****
Motif duration	$F_{1,13} = 0.2001$	0.662	$F_{2,26} = 2.210$	0.1299	$F_{2,26} = 2.101$	0.1426