Thermal and alkaline pre-treatments of inoculum halt methanogenesis and enables cheese whey valorization by batch acidogenic fermentation

Maria Paula Giulianetti de Almeida[1,2,3], Camille Mondini[2], Guillaume Bruant[4], Julien Tremblay[4], David G. Weissbrodt[2*] and Gustavo Mockaitis[1*]

[1] Interdisciplinary Research Group on Biotechnology Applied to the Agriculture and the Environment, School of Agricultural Engineering, University of Campinas (GBMA/FEAGRI/UNICAMP) - , Campinas, SP, Brazil

[2] Department of Biotechnology, Delft University of Technology, Delft, The Netherlands

[3] Interinstitutional Graduate Program in Bioenergy (USP/UNICAMP/UNESP) – 330 Cora Coralina Street, Campinas/SP, CEP 13.083-896, Brazil.

[4] National Research Council Canada, Energy, Mining and Environment research centre, Genomics and microbiomes group, 6100 Royalmount Avenue, H4P 2R2, Montreal, QC, Canada

* Shared senior authorship and correspondence: Gustavo Mockaitis (gusmock@unicamp.br) and David G. Weissbrodt (D.G.Weissbrodt@tudelft.nl)

Abstract

Carboxylates like volatile fatty acids (VFAs) can be produced by acidogenic fermentation (AF) of

dairy wastes like cheese whey, a massive residue produced at 160.67 million m^3 of which 42% are

not valorized and impact the environment. In mixed-culture fermentations, selection pressures are

needed to favor AF and halt methanogenesis. Inoculum pre-treatment was studied here as selective

pressure for AF demineralized cheese whey in batch processes. Alkaline (NaOH, pH 8.0, 6 h) and

thermal (90°C for 5 min, ice-bath until 23°C) pre-treatments, were tested together with batch

operations run at initial pH 7.0 and 9.0, food-to-microorganism (F/M) ratios of 0.5 to 4.0 g COD g^{-1}

VS, and under pressurized and non-pressurized headspace, in experiments duplicated in two

institutes. Acetic acid was highly produced (1.36 and 1.40 g COD_{AcOH} L^{-1}) at the expense of

methanogenesis by combining a thermal pre-treatment of inoculum with a non-pressurized batch

operation started at pH 9.0. Microbial communities comprised of VFAs and alcohol producers, such

as *Clostridium*, *Fonticella*, and *Intestinimonas*, and fermenters such as *Longilinea* and *Leptolinea*.

Communities also presented the lipid-accumulating and bulk and foaming *Candidatus Microthrix*

and the metanogenic *Methanosaeta* regardless of no methane production. An F/M ratio of 0.5 g

COD g^{-1} VS led to the best VFA production of 1,769.38 mg L^{-1}. Overall, inoculum thermal pre-

treatment, initial pH 9.0, and non-pressurized headspace acted as a selective pressure for halting

methanogen and producing VFAs, valorizing cheese whey via batch acidogenic fermentation.

Keywords: acidogenic fermentation, physicochemical pre-treatments, alkaline, thermal, volatile

fatty acids, cheese whey

1 Introduction

Anaerobic digestion (AD) has been used from the 19[th] century onwards to obtain biogas as an energy carrier (Kigozi et al., 2014). Since then, AD processes are being constantly improved (Abbasi et al., 2012; van Lier et al., 2001) to process various waste streams (Ostrem et al., 2004) and address energy production, environmental burdens, and circular economy. The primary goal remains to reduce organic matter and remove nutrients from agro-industrial, food, and municipal solid wastes (Chen et al., 2008; Goud and Mohan, 2012; Wonglertarak and Wichitsathian, 2014).

Cheese whey (CW) is a by-product of the dairy industry with a high organic load (i.e., 50 to 80 g O_2 L^{-1}, in terms of chemical oxygen demand (COD))(Saddoud et al., 2007). It has an annual production of 160.67 million m^3 where 58% of the production is absorbed by various industries (e.g., food, nutrition, cosmetics, and pharmaceutical). However, a staggering amount of 66.5 million m^3 year $^{-1}$ of CW (Food and Agriculture Organization of the United Nations, n.d.; Tsakali et al., 2010; USDA - Foreign Agricultural Service, 2022) is currently transformed into low added-value products such as animal feed, fertilizer, or being discharged into water bodies leading to eutrophication processes (Smithers, 2008; Tsakali et al., 2010). Hence, cheese whey can be an excellent substrate for AF processes.

Acidogenic fermentation focuses on the valorization of organic matter via the carboxylate platform, by combining the inhibition of methanogenesis and production of VFAs at high yields, which are of great economic interest due to their potential industrial applications (e.g., biofuels, biopolymers, and chemicals) (Angenent et al., 2016; Kleerebezem et al., 2015; Mulders et al., 2020; Rombouts et al., 2019).

Selective pressures (e.g., pH, physicochemical pre-treatments, reactor headspace pressure, and F/M ratio) can dictate microbial diversity and dynamics, interactions, energy requirements, and

preferred metabolisms (Pretorius, 1987). Most studies in AD and AF processes focus on the pre-

treatment of the substrate and waste-activated sludge (Sarkar et al., 2021). The present study

focused on various selective pressure mechanisms for enhancing VFA production and halting

methanogenesis in AF processes. We also investigated the effects of alkaline and thermal inoculum

pre-treatments, variations in the initial pH in headspace pressure, and different F/M ratios on the

acidogenic fermentation of cheese whey.

Alkaline pre-treatments have been shown to improve AD processes by increasing sludge

solubilization and enhancing methane production (Chen et al., 2007; Li et al., 2012; López-Torres

and Llórens, 2016; Navia and Vidal, 2002; Wonglertarak and Wichitsathian, 2014), while thermal

pre-treatments have been used to inhibit methanogenesis (Alibardi et al., 2012; Ramos-Suarez et

al., 2021; Sarkar et al., 2017).

However, thermal pre-treatments, that can be performed at mesophilic or hyperthermophilic

temperatures (i.e., 30°C to 180°C) are time-consuming (up to several hours), which would increase

AF processes' overall costs (Corti and Lombardi, 2007). Contradictory results have also been

observed regarding the effects of such pre-treatments on methanogenesis, with an inhibition

observed after alkaline pre-treatment (Yuan et al., 2006) and an enhancement observed after

thermal pre-treatment (Zhang et al., 2015). The lack of consensus on the best conditions for

inoculum pre-treatment and their influence on AF processes still needs to be elucidated. To date,

studies on thermal pre-treatments are mostly performed on waste-activated sludge. Their use as a

selective pressure mechanism for VFA production remains scarce.

Variations in the initial pH can positively influence VFA production (Dareioti et al., 2014). In

addition, uncontrolled pH approaches can reduce AF process costs since there is no further need for

chemical utilization for stabilizing pH (Sarkar et al., 2021).

Headspace gas composition and pressure play a role in product formation and metabolic

pathways preferentially used. However, most studies still focus on hydrogen production (Darvekar

et al., 2019). According to Zhou (2018) and Sarkar (2017), low hydrogen pressure favors VFA

formation. Finally, the F/M ratio, which is inoculum and substrate-dependent, is a parameter that

impacts acidogenesis, with lower F/M ratios being beneficial to VFAs production (Pang et al.,

2019; Shah et al., 2015).

In this work, we aimed at (*i*) identifying how abiotic factors (i.e., pH, inoculum pre-treatment,

headspace pressure, and F/M ratio) influence the products spectra of cheese whey via AF, (*ii*)

validating thermal pre-treatment efficiency for halting methanogenesis, and (*iii*) identifying the

parameters that increase acetate level of production for other biological processes (e.g., microalgal

photoorganoheterotrophic biomass production).

2   Materials & Methods

98   Figure 1 depicts the overall methodology, involving experiments performed at the University of

99   Campinas (UNICAMP, Brazil) and reproduced at Delft University of Technology (TU Delft, The

100   Netherlands). Substrate and inoculum preparation and inoculum pre-treatment are common to all

101   experiments apart from a few modifications described hereafter.

Effect of Inoculum pre-treatment, pH and headspace pressure

University of Campinas, Brazil

Effect of pre-treatment efficiency

Effect of F/M ratio for optimal VFAs production

Delft University of Technology, The Netherlands

Figure 1 – Overall experiment methodology. The inoculum went through alkaline (A_{PT}) and thermal
(T_{PT}) pre-treatments. The initial pH at the start of the digestions was set at 7.0 and 9.0 and
headspace assays were carried in pressurized (P) and non-pressurized (NP) batches. Analyses
encompassed gas and VFA profiles, total organic carbon (TOC), total nitrogen (TN), COD, solid
series, carbohydrates, and amplicon sequencing (16S rRNA gene).

2.1 Demineralized cheese whey media

The chosen carbon source for all experiments consisted of 40% demineralized whey powder

(WPC40), which was the closest substrate to raw cheese whey. The substrate medium for

experiments held at UNICAMP was composed of (per L): WPC40 (Pic-Nic, Brazil) (4 g)

(Mockaitis et al., 2006) and the following reagents: $NaHCO_3$ (1 g), NaCl (250 μg), $MgCl_2.6H_2O$ (7

μg) and $CaCl_2 \cdot 2H_2O$ (9.5 μg) (Torres, 1992). The medium was prepared fresh before experiments

and its initial pH was approximately 7.0. No pH corrections were made after medium preparation.

The sludge used as inoculum was obtained from a UASB reactor from a poultry abattoir in Brazil

(25°05'10.1" S 47°58'49.7" W). National System for the Management of Genetic Heritage and

Associated Traditional Knowledge (SisGen) register number is AE43468.

For the experiments held at TU Delft, the medium was composed of (per L): WPC40 (Subo

International, The Netherlands) (4 g) amended with 0.3 mL of a micronutrients solution per g of

substrate COD. The micronutrients solution was composed of $FeCl_3 \cdot 6H_2O$ (2 g L^{-1}), $CoCl_2 \cdot 6H_2O$ (2

g L^{-1}), $MnCl_2 \cdot 4H_2O$ (0.5 g L^{-1}), $CuCl_2 \cdot 2H_2O$ (32 mg L^{-1}), $ZnCl_2$ (50 mg L^{-1}), $HBO3$ (50 mg L^{-1}),

$(NH_4)6Mo_7O_2 \cdot 4H_2O$ (90 mg L^{-1}), $Na_2SeO_3 \cdot 5H_20$ (100 mg L^{-1}), $NiCl_2 \cdot 6H_2O$ (6 mg L^{-1}), EDTA (1 g

L^{-1}), HCl 36% (1 mL L^{-1}), Resazurine (0.5 g L^{-1}) and yeast extract (2 g L^{-1}). Sludge used as

inoculum was obtained from an anaerobic digester at WWTP Harnashpolder in The Netherlands

(52°00'48.8"N 4°19'02.5"E). It is valid to notice that 1 g L^{-1} of WPC40 corresponds to 1 g O_2 L^{-1} in

COD terms.

2.2 Mechanical homogenization, thermal and alkaline pre-treatments of inocula

The anaerobic granular sludge used as inoculum at UNICAMP was mechanically homogenized

by using a blender (Britannia, Brazil) to decrease the size of bioaggregates while increasing their

superficial area. The anaerobic granular sludge used as inoculum in TU Delft was used as is.

Thermal pre-treatment, which was performed both at UNICAMP and TU Delft, was adapted

from Mockaitis (2020). Briefly, the sludge was heated at 90°C in a water bath for 20 minutes under

constant stirring. Heating was then halted by decreasing the sludge temperature to 23°C in an ice

bath.

Alkaline pre-treatment consisted in increasing the sludge pH up to 8.0 using 1 M NaOH under

continuous mixing, and controlling it for the next 6 hours, with a correction to 8.0 if needed. This

pre-treatment was only performed at UNICAMP.

2.3   Effect of thermal and alkaline pre-treatments of inocula, initial batch pH, and headspace

pressure on acidogenic fermentation

Batch experiments were conducted at UNICAMP. They assessed the effects of three factors

evaluated at two levels in WPC40 acidogenic fermentation, namely: (*i*) inoculum pre-treatment

(alkaline (A_{PT}) and thermal (T_{PT})) to enhance sludge biodegradability and select for alkaline-tolerant

acetoclastic microorganisms, and to inactivate methanogenic microorganisms, respectively; (*ii*)

variation in the initial pH of digestion process (7.0 and 9.0) to further select microorganisms

adaptable to neutral and alkaline pH environments, and (*iii*) gas headspace pressure (pressurized (P)

and non-pressurized (NP)) to investigate the influence on AF products spectra. Table 1 shows the

two-level factorial experimental design with 3 factors (2^3 designs) applied to this setup.

Table 1 – The 23 experimental designs for phase one. The (±1) values indicate the upper and lower
levels of the investigated factors. The design aimed at testing the main effects (referred to as a, b
and c), the 2-factor interactions effects (ab, ac, and bc), and the 3- factors interaction effect (abc).
Two control experiments were included.

Experiment	Factors		
	Pre-treatment (binomial)	pH (continuous)	Headspace (continuous)
A7P (*ac*)	Alkaline (+1)	7.0 (-1)	P (+1)
A9P (*abc*)	Alkaline (+1)	9.0 (+1)	P (+1)
A7NP (*a*)	Alkaline (+1)	7.0 (-1)	NP (-1)
A9NP (*ab*)	Alkaline (+1)	9.0 (+1)	NP (-1)
T7P (*c*)	Thermal (-1)	7.0 (-1)	P (+1)
T9P (*bc*)	Thermal (-1)	9.0 (+1)	P (+1)
T7NP (*1*)	Thermal (-1)	7.0 (-1)	NP (-1)
T9NP (*b*)	Thermal (-1)	9.0 (+1)	NP (-1)
Control (*P*)*	______	8.25	P (+1)
Control (*NP*)*	______	8.25	NP (-1)

Experiments (Figure 1) were performed in 1 L Duran flasks (total volume of 1,130 mL) with an
initial working volume of 730 mL and an initial headspace volume of 398 mL. Bottles were
inoculated at a concentration of 6.6 g total volatile solids (TVS) L^{-1}, as depicted in Mockaitis et al.
(2020). Batch reactors were continuously agitated at 50 rpm (Orbital Incubator Marconi – MA420)
at a mesophilic temperature of 35°C for 30 days.

2.3.1   Batches headspace pressure assays

The influence of headspace pressure (pressurized and non-pressurized batches) on VFAs
production was investigated according to Peixoto et al., (2011). Manometric pressure was
measured, then, 3 mL samples of gas were collected with a syringe containing a pressure lock
(Thermo Fisher). Each sample corresponded to a batch condition.

In pressurized headspace assays, butyl rubber stoppers of each flask were covered with silicone

sealant after manometric measurement and gas sample collection, allowing gas accumulation in the

batch headspace.

In non-pressurized experiments, the headspace of each flask was punctured with a needle after

pressure manometric reading and gas sampling, allowing the remaining gas to be released until

reaching the atmospheric pressure value. After this step, butyl rubber stoppers were also covered

with silicone sealant avoiding any gas release into the atmosphere.

Once collected, gas samples were analyzed with a gas chromatograph (GC) equipped with a

thermal conductivity detector GC-TCD (Construmac, Brazil) with hydrogen as a carrier gas.

The volumetric production of biogas and its constituents were inferred through equation 1 as a

punctual function from sampling timestep $t_0 = 0$ to t for non-pressurized assays.

$$\Gamma_n = \frac{T^{STP}}{T \cdot P_{STP}} \cdot \sum_{i=1}^{n} V_{i-1}^{H} \left(P_i \cdot {}^{A}x_i \ - \ P_{atm} \cdot {}^{A}x_{i-1} \right) \qquad \text{Equation 1}$$

Where Γ_n is the volumetric production of a gas of interest (N_2, H_2, CH_4 or CO_2) in standard

temperature and pressure (STP) equivalents (L), at a given time n; P_i is the measured pressure at

sampling time (atm); V_{i-1}^{H} is the headspace volume before sampling (L); T is system's temperature

(K), T^{STP} is the standard temperature (273 K), P^{STP} is standard pressure (1 atm), ${}^{A}x_i$ is the molar

fraction of the gas of interest at sampling time and ${}^{A}x_{i-1}$ is the molar fraction of the gas of interest

before sampling.

2.3.2 VFAs measurements

VFAs, alcohols and carbohydrates were quantified by high performance liquid chromatography

(HPLC) based on Penteado et al.,(2012). The VFAs and alcohol profile consisted of the following:

lactic acid, formic acid, acetic acid, propionic acid, butyric acid, iso-butyric acid, iso-valeric acid

and ethanol.

The chromatograph was equipped with two LC-20AD pumps, one DGU-20A3R degasser, a

SIL-20AHT autosampler, a CTO-20A column oven, a SPD-20 UV detector with readings at 210

nm (Shimadzu, Japan), an Aminex HPX-87H 300 x 7.8 mm column (BioRad, USA), a RID-10A

index refraction detector and a CBM-20A controller (Shimadzu, Japan).

Two mL of mixed liquors were centrifuged at 16,025 xg for 5 min.

Forty µL of H_2SO_4 (2M) was added to 1 mL of the supernatant to acidify the sample for ideal

analyte separation. The samples were then filtered through regenerated cellulose (RC) syringe

filters (RC 0.20 µm) (GVS, Italy), and transferred to 1.5 mL pre-washed vials (H_2SO_4 at 2 mol L^{-1}

to avoid any contamination) before HPLC analyses. The HPLC run time was 60 minutes per sample

with a constant column oven temperature of 43°C and 0.005 M H_2SO_4 mobile phase at a flux rate of

0.5 mL min $^{-1}$.

2.4   Inoculum thermal pre-treatment and non-pressurized headspace efficiency

Inoculum thermal pre-treatment and non-pressurized headspace assays were reproduced to

confirm the efficiency of such imposed conditions in halting methanogenesis while enhancing

VFAs production. Each factor was tested at one level, based on the effects identified at UNICAMP

(section 2.3): inoculum thermal pre-treatment, initial pH of 9.0, and non-pressurized headspace. A

batch with no inoculum pre-treatment, initial pH of 9.0, and non-pressurized headspace acted as a

control. The experimental design with three factors applied to this set-up is shown in Table 2.

Similar to the first experiments conducted, the initial working volume of the 1 L Duran flasks

was 750 mL with an initial headspace volume of 398 mL, for a total working volume of 1,148 mL.

The inoculum concentration was identical at 6.6 g TVS L^{-1} as depicted in Mockaitis et al., (2020).

Batch reactors were incubated for 10 days at a mesophilic temperature of 35°C in a Certomat® BS1

incubator (Sartorius Stedim Biotech, Germany) under continuous agitation (145 rpm). The

headspace was released with the aid of Cole-Palmer stopcocks with Luer connection, a 1-way male

lock (Cole-Parmer, USA). Experiments were performed in triplicate and lasted 12 days.

2.4.1 VFAs measurements

VFAs and carbohydrates were quantified by HPLC. One mL of mixed liquors was

centrifuged at 5,073 x g for 5 minutes and supernatants were filtered using a 0.45 μm syringe filters.

Filtered samples were then transferred to 300 μL Waters total recovery vials, from which 10 μL

were injected with a 2707 Waters HPLC autosampler at 15°C The chromatograph was equipped

with a Waters M515 HPLC (Waters, USA) pump, a Waters 2414 refractive index (RI) detector,

with 1,024 of sensitivity and a Waters 2489 UV/Visible detector at 210 nm. The column was a

BioRad HPX-87H (300 x 7.8 mm) with a BioRad Cation-H refill cartridge (30 x 4.6 mm) guard

column (BioRad, USA) and the column oven was built in-house. The flow rate of the pump was 0.6

mL min^{-1} and the temperature of the column was set at 59°C. The RI detector was operated at

30 °C. The mobile phase was 1.5 mmol L^{-1} phosphoric acid diluted in ultrapure water (MilliQ,

Merck Millipore). The VFAs measured were acetic acid, butyric acid, formic acid, caproic acid,

propionic acid, valeric acid, iso-butyric acid, iso-caproic acid, and iso-valeric acid.

2.5   Ideal F/M ratio for optimal VFAs production with thermally pre-treated inocula

F/M ratio is a measurement used to determine the amount of substrate needed for the number of

microorganisms present in a system. It is an important parameter to evaluate in an approach aiming

at maximizing VFA production. To determine the best FM ratio for VFA production when working

with a thermal pre-treated sludge at non-pressurized headspace, four different F/M ratios were

tested: 0.5, 1.0, 2.0, and 4.0,g_{COD} g_{VS} $^{-1}$. These ratios were obtained by dividing the COD of the

substrate by the VS of the sludge. The initial pH of all experiments was 9.0. In section 2.4, both

control and thermal batches presented an F/M ratio of 0.5 g $_{COD}$ g $_{VS}$ $^{-1}$. g $_{COD}$ g $_{VS}$ $^{-1}$ acted as control.

Assays were conducted in duplicates for 14 days. The experimental design is depicted in Table 2.

Table 2 – Experimental design for the F/M ratio experiments. All experiments were done in
duplicate.

Experiment	Factors		
	Pre-treatment	F/M ratio (continuous)	Headspace (continuous)
FM0.5	Thermal	0.5	NP
FM0.5	Thermal	0.5	NP
FM1.0	Thermal	1.0	NP
FM1.0	Thermal	1.0	NP
FM2.0	Thermal	2.0	NP
FM2.0	Thermal	2.0	NP
FM4.0	Thermal	4.0	NP
FM4.0	Thermal	4.0	NP

2.5.1   Biogas measurements

The presence or absence of CH_4 in the gas was detected by injecting 10 mL samples of gas

from the headspace of the bottles in a GC (Agilenttech 7890 A, Agilent Technologies Inc., USA)

equipped with an HP-PLOT Moleseive GC column (Agilent 19095P-MS6, Agilent Technologies

Inc., USA) of 60 m x 0.53 mm x 200 μm and a thermal conductivity detector (TCD). The carrier

gas was helium (14.8 psi, 23 mL min^{-1}) and the operating temperature was 200 ℃.

2.6   Physicochemical analyses

Mixed liquors and gas phases of the experimental flasks were sampled for physicochemical

analyses in both original and replicated experiments. For the experiments held at UNICAMP, every

15 days, 150 mL of mixed liquors were collected and used in the following analyses and respective

code methods: solid series (2540 B-F; 50 mL), COD (5220 B; 1 mL), sulfate (4500-SO_4^{2-} E; 1 mL),

sulfide (4500-S^2 D, 1 mL), total organic carbon (5310-TOC; 15 mL) (APHA, 2005) and total

nitrogen (TN, ASTM D8083; 15 mL) (ASTM International, 2016). Additional analyses were

performed consisting of pH measurements (4500 H) (Federation 2005), headspace assays (CH_4, N_2,

and CO_2, 3 mL of gas phase), HPLC (2 mL) for VFAs, sugars and alcohols characterization, total

carbohydrates measurements (Dubois et al., 1956) (1 mL), total alkalinity and VFAs measurements

(Dilallo and Albertison, 1961; Ripley, L.E., Boyle, W.C. and Converse, 1986) (60 mL). The

remainder sludge was discharged. On alternate days, these analyses were also performed.

For the replicated experiments performed at TU Delft, 100 mL of mixed liquors were collected

on the first and last days of the process and were used in the following analyses in the same volume

as mentioned above: solid series, COD, and TOC/TN. Other analyses were also performed,

consisting of pH measurements, biogas composition determination, HPLC for VFAs and sugars

characterization, and total carbohydrates measurements. In addition, on alternate days 12 mL of

mixed liquors were sampled for pH, HPLC, COD, TOC, and TN measurements.

2.7   16S rRNA gene amplicon sequencing

2.7.1   Sequencing experiments

Microbial community analyses were performed through 16S rRNA gene amplicon sequencing

on (*i*) the initial and final samples from inoculum thermal pre-treatment efficiency assays and (*ii*)

the final samples of each F/M ratio experiment.

0.5 mL samples of mixed liquors were collected from each condition in 1.5 mL Eppendorf

tubes completed with demineralized water for initial washing. Samples were centrifuged at 10,000 x

g, 4 ºC for 3 minutes, and the supernatant was discarded. DNA was extracted using the Qiagen

DNeasy® UltraClean® Microbial Kit (Qiagen, Germany) according to the manufacturer's

instructions. The quality and concentrations of the DNA extracts were assessed using a Qubit

fluorometer and the Qubit dsDNA HS assay kit (Thermo Fischer Scientific, USA) according to the

manufacturer's instructions.

For each sample, a minimum of 20 μL of DNA extracts with concentrations ranging from 13.9

and 95 ng μL^{-1} were sent to Novogene (Beijing, China) for 16S rRNA gene amplicon sequencing.

All sequencing experiments (i.e 16S rRNA sequence amplifications, libraries preparations, and

sequencing runs) were performed at Novogene, according to their protocols. The targeted

hypervariable regions V3-V4 of the 16S rRNA gene were amplified by PCR with the pair of

barcoded   primers   341F   (CCTAYGGGRBGCASCAG)   and   806R

(GGACTACNNGGGTATCTAAT) (Yu et al., 2005). This set of primers is known to cover both

bacteria and archaea with high specificity, except for *Planctomycetes* (Weissbrodt et al., 2020).

2.7.2 Sequences analyses

All sequencing data received from Novogene were analyzed using AmpliconTagger (Julien

Tremblay, 2019). Briefly, raw reads were scanned for sequencing adapters and PhiX spike in

sequences. The remaining paired-end reads were processed to remove primer sequences (pTrimmer

v1.3.3 – (Zhang et al., 2019) and discard reads having average quality Phred score lower than 20.

The remaining sequences were processed for generating Amplicon Sequence Variants (ASVs)

(DADA2 v1.12.1) (Callahan 2016) using the following key parameters: filterAndTrim (maxEE =2;

truncQ = 0; maxN = 0; minQ = 0) and learnErrors (nbases=1e8) functions for both forward and

reverse filtered reads. Reads were then merged using the mergePairs (minOverlap = 10;

maxMismatch = 0) function. Chimeras were removed with DADA2's internal

removeBimeraDeNovo (method = 'consensus') method followed by UCHIME reference (Rognes et

al., 2016). ASVs were assigned a taxonomic lineage with the RDP classifier (Wang et al., 2007)

using training sets containing the complete Silva release 138 database (Quast et al., 2013)

supplemented with a customized set of mitochondria and plastid sequences. The RDP classifier

gave a score of (0 to 1) to each taxonomic depth of each ASV. Any taxonomic depth having a score

of ≥ 0.5 was kept reconstructing the final lineage. Taxonomic lineages were combined with the

cluster abundance matrix obtained above to generate a raw ASV table from which the bacterial or

fungi ASV tables were generated. Five hundred 1,000 reads rarefactions were then performed on

these ASV tables and the average number of reads of each ASV of each sample was computed to

obtain consensus rarefied ASV tables. Alpha diversity metrics (RTK v 0.93.2 – ((Saary et al., 2017)

and taxonomic summaries microbiomeutils v 0.9.3 – (Julien Tremblay, 2020)) were then computed

using the consensus rarefied ASV tables. Figures were generated in R using the ggplot2 package.

3 Results and Discussion

3.1 Effects of inoculum thermal and alkaline pre-treatments and initial pH on the anaerobic

digestion process

3.1.1   Inoculum thermal pre-treatment efficiently halted methanogenesis

As demonstrated in Figure 2, inoculum thermal pre-treatment successfully inhibited methanogenesis in all tested conditions (i.e., pressurized and non-pressurized headspace assays with initial pH of 7.0 and 9.0).

Inoculum alkaline pre-treatment halted methanogenesis in both pressurized and non-pressurized assays but only with initial pH 7.0, presenting negligible to no levels of total methane production of 0.0 and 0.54 mL, respectively. Initial pH 9.0 presented a total methane production of about 52.29 mL for pressurized assays and 0.67 mL for non-pressurized ones.

As a strategy to maximize VFA production, inoculum thermal pre-treatment appears to be the best selective pressure for impairing methanogens in batch processes. It is of utmost importance to confirm whether this selective pressure is also successful to halt methanogenesis considering different types of inoculum (e.g., waste activated sludge and secondary sludge), reactors (e.g., sequence- batch reactors, and continuous-stirred tank reactors) and configurations (e.g., organic loading rate, hydraulic retention time, coupled reactors). If so, then inoculum thermal pre-treatment can be implemented in full-scale AF processes, decreasing their overall costs (Corti and Lombardi, 2007).

Some studies have used alkaline pre-treatment of inoculum to increase the biodegradability of waste-activated sludge (Wonglertarak and Wichitsathian, 2014), and to stabilize the sludge originated from aerobic processes while treating the mineralization of its remaining organic compounds (Navia and Vidal, 2002). Sludge alkaline pre-treatment increased the COD: COD_{Total}

ratio (Navia and Vidal, 2002) VS while improving volatile solids (VS) reduction (Appels et al.,

2011; Lin et al., 1997). Hence, their methane production would also increase.

Noteworthily, the methane production levels observed in our experiments over 30 days (i.e.,

A_{PT}7NP, 0.54 mL; A_{PT}7P, 0.00 mL; A_{PT}9NP 0.67 mL, and A_{PT}9P, 52.29 mL), did not agree with

the findings of Wonglertarak and Wichitsathian (2014). Their methane production was about 2,

526.60 mL in a thermophilic reactor with alkaline pre-treated sludge and 2,173.8 mL in a

mesophilic reactor with alkaline pre-treated sludge. Our findings also contradicted the ones found

by Lin et al. (1997) which was an average CH_4 production rate of 222.25 L m^3 d^{-1}.

This could be explained using different methodologies in each work such as maintaining the

reactor with a constant alkaline pH after treating the sludge, different reactors configurations (e.g.,

single-stage high-rate reactor and continuous stirred tank reactor and batch reactor), hydraulic

retention times, organic loading rates as well as the type of inoculum used (i.e., WAS and UASB

granular sludge).

For instance, we used 1 M NaOH to treat the inoculum, correcting the pH to 8.0 hourly for 6

hours. However, after pre-treating the inoculum, batch reactors were assembled with initial pH of

7.0 and 9.0, with the final pH from both pressurized and non-pressurized headspaces, ranging from

4.9 and 5.1, respectively. Lin et al. (1997), treated secondary sludge with NaOH ranging from 0.02

to 0.2 M, which pH increased from 7.21 to 13.2 in 24 h hydrolysis and 0.2 M NaOH. Alternatively,

Wonglertarak and Wichitsathian's (2014) alkaline inoculum pre-treatment consisted of treating

waste-activated sludge with pH varying from 8.0 to14.0 and varying NaOH concentration from 8 to

31,000 g m^{-3} wet sludge.

Another result of interest regarding inoculum alkaline pre-treatment was the correlation

between acetate and methane production. While there was no methane production in inoculum

alkaline treatment with initial pH 7.0 in both non-pressurized and pressurized headspace batches,

acetate production differed 185.6% from non-pressurized headspace assay to pressurized headspace

(i.e., $A_{PT}7NP$ 1,056.04 mg COD_{AcOH} L^{-1} and $A_{PT}7P$ 369.82 mg COD_{AcOH} L^{-1}). So, headspace

pressure can favor acetate production in acetoclastic methanogens.

The detected levels of CO_2 in both $A_{PT}7NP$ and $A_{PT}7P$ (i.e., 1.84 mL and 60.78 mL,

respectively) were low as was the average of the final pH for both conditions (i.e., 4.8). Hence,

instead of methanogenesis being inhibited due to disturbance in the buffer system (Casallas-Ojeda

et al., 2020), we can infer that most CO_2 was dissolved in the liquid phase, decreasing the pH

(Deublein and Steinhauser, 2011). Figure 2 shows the methane and carbon dioxide production in

alkaline and thermal batches with both pressurized and non-pressurized headspaces.

Figure 2. Methane and carbon dioxide gas production profiles for control (C), inoculum alkaline (A_{PT}), and thermal (T_{PT}) pre-treatment with initial pH

of 7.0 and 9.0 during 30 days. Non-pressurized (NP) and Pressurized (P) headspace assays are also represented. In the figure: ■ CH_4NP, ☐ CH_4P,

CO_2NP and ⊙ CO_2P

3.1.2 Initial pH, headspace pressure, and inoculum pre-treatments influenced VFAs product

spectra.

Cheese whey has a natural tendency to acidify and since pH was not controlled during the

experiments (pressurized and non-pressurized batches displayed an average pH of 4.96 and 5.26,

respectively), it is likely that hydrolysis was not a limiting step in the AF process.

An additional hypothesis to justify the different production of VFAs and alcohol in all batches

was that together with uncontrolled pH, the headspace internal pressure variation from pressurized

and non-pressurized batches played a key role on the spectra and quantity of VFAs produced. In

pressurized batches, the pressure would contribute to the gas phase being accumulated within the

liquid phase, which influenced both gas and VFA spectra. In non-pressurized batches, due to

headspace release, gas and VFAs production restarted daily, after each sampling.

Overall, non-pressurized batches produced more VFAs than pressurized ones. Special

highlights can be given to $A_{PT}9$ NP (21,360.54 mg $COD_{compound}$ L^{-1}), $T_{PT}9$ NP (4,108.71 mg

$COD_{compound}$ L^{-1}), and $T_{PT}7$ NP (3,414.15 mg $COD_{compound}$ L^{-1}), when compared to their counterparts

($A_{PT}9$ P 1,940.14 mg $COD_{compound}$ L^{-1}, $T_{PT}9$ P 2,151.99 mg $COD_{compound}$ L^{-1} and $T_{PT}7$ P 2,140.62 mg

$COD_{compound}$ L^{-1}). Both pressurized and non-pressurized controls and $A_{PT}7$ did not present a

substantial difference with an increase of approximately 2.9% in VFAs produced in non-pressurized

batches compared to pressurized batches as seen in table 3.

It is important to stress that although $A_{PT}9$ NP, $T_{PT}9$ NP, and $T_{PT}7$ NP produced greater

quantities of metabolites, they did not produce a great variety of VFAs and alcohol. $A_{PT}9$ NP

produced 7,677.30 mg COD_{IBA} L^{-1} of iso-butyric acid and 13,683.24 mg COD_{EtOH} L^{-1} of ethanol.

These metabolites production together with the absence of acetic acid production are a clear result

of a metabolic shift that was favored due to the imposed selective pressures in this batch (i.e.,

alkaline inoculum pre-treatment, initial pH of 9.0, and non-pressurized headspace).

This was in opposition to all other conditions tested. It is likely that in such operational

conditions, acetate served as the substrate for fermentative bacteria to form iso-butyric acid and

ethanol, as already described by Thatikayala et al. (2021) and Liu et al. (2022). Both chemicals are

of great industrial interest (i.e., pharmaceutical, feed, chemical, biofuels). Optimization of AF

processes towards the production of either ethanol or iso-butyric acid or both can be an additional

approach for cheese whey valorization.

$T_{PT}9$ NP and $T_{PT}7$ NP displayed a production of acetic acid of 1,479.13 and 1,558.17 mg

COD_{AcOH} L^{-1}, respectively. As shown in figure 2, there was no production of methane. Acetic acid

was consumed in $A_{PT}9NP$, whereas in $A_{PT}7NP$ its production was slightly lower than both $T_{PT}9$ NP

and $T_{PT}7$ NP with 1,056.04 mg COD_{AcOH} L^{-1}.

Interestingly, CNP also presented a higher acetic acid production of 1,109.13 mg COD_{AcOH} L^{-1},

which shows that headspace pressure is a more suitable selective pressure parameter than inoculum

pre-treatment for acetic acid production. The produced acetic acid can thus be utilized as an organic

carbon source for higher added-value biological processes (i.e., microalgal

photoorganoheterotrophic biomass production).

The imposed parameters (*i.e.*, inoculum pre-treatments, initial pH, and headspace pressure) also

acted as selective pressures in the batch experiments modifying the microbial fermentation end-

products. In general, thermal pre-treatment, non-pressurized headspace, and initial pH 9.0 displayed

a trend for higher VFA production. Nevertheless, it is not possible to affirm what the best

combination of parameters would be since there was not a pattern of production observed for all

VFAs in this given setup.

However, the combination of parameters can be suitable for the production of specific volatile

acids or ethanol, as seen in $T_{PT}7NP$ and $T_{PT}9NP$ for the production of acetic (1,479.13 and 1,558.17

mg COD_{AcOH} L^{-1}, respectively) and butyric acids (1,207.46 and 1,425.36 mg COD_{BTA} L^{-1}), and

$A_{PT}9NP$ for iso-butyric acid and ethanol (7,677.30 mg COD_{IBA} L^{-1} and 13,683.24 mg COD_{EtOH} L^{-1}).

Non-pressurized batches presented a global production of 3 times more VFAs than pressurized

batches. Besides common VFAs, pressurized headspace batches produced formic and lactic acids

but in negligible quantities as shown in table 3.

Table 3. VFAs and alcohols product spectra of pressurized (P) and non-pressurized (NP) headspace batches with alkaline (A_{PT}) and thermal (T_{PT}) pre-treatments and initial pH of 7.0 and 9.0. Experiments lasted for 30 days.

	Time	Batch operational conditions (mg COD compound L-1)									
		Control		APT7		APT9		TPT7		TPT9	
		P	NP	P	NP	P	NP	P	NP	P	NP
Acetic acid	t0	53.88	154.34	97.50	0.00	0.00	180.11	183.94	314.61	154.56	308.26
	tf	267.62	1,109.13	369.82	1,056.04	280.22	0.00	377.39	1,479.13	417.91	1,558.17
Butyric acid	t0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	tf	929.83	317.63	662.53	177.73	773.31	0.00	881.05	1,207.46	821.59	1,425.36
Iso-butyric acid	t0	0.00	0.00	0.00	0.00	0.00	0.00	536.00	10.84	537.85	0.00
	tf	108.02	0.00	7.47	113.92	78.36	7,677.30	57.69	29.77	182.37	85.14
Ethanol	t0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	tf	494.93	0.00	489.74	0.00	598.57	13,683.24	621.32	302.99	479.02	303.51
Formic acid	t0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	tf	0.33	0.00	0.50	0.00	0.46	0.00	0.52	0.00	0.44	0.00
Lactic acid	t0	0.00	0.00	0.00	0.00	94.07	0.00	37.61	1,524.34	38.72	1,137.60
	tf	0.50	0.00	0.42	0.00	0.00	0.00	0.22	0.00	0.00	0.00
Propionic acid	t0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	tf	69.37	189.39	34.78	130.51	65.84	0.00	15.53	91.24	20.04	111.16
Valeric acid	t0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	tf	0.00	0.00	0.00	0.00	0.00	0.00	44.12	0.00	47.95	0.00
Iso-valeric acid	t0	0.00	0.00	0.00	0.00	0.00	0.00	270.08	0.00	260.84	0.00
	tf	45.40	194.40	45.39	179.11	143.39	0.00	142.77	303.56	182.67	625.38

Initial WPC40 carbohydrate concentration was approximately 260 $mg_{glucose}$ L^{-1}, stabilizing at

65 $mg_{glucose}$ L^{-1} on the fourth day. The acidogenesis rate of 1.9 d^{-1}, was close to the typical rate of 2

d^{-1} observed in similar studies(van Lier et al., 2008). COD removal and volatile solids (VS) showed

similar patterns, with sCOD decreasing from 4.2 to 2.4 g L^{-1} and VS decreasing from 4.92 to 4.74 g

L^{-1}. This was likely due to biomass production and organic matter oxidation. Control batches

showed a decrease from 4.92 to 4.74 g L^{-1}, corroborating with COD removal results. Cheese whey

alkalinity varied by 6% in all batches (1.53 to 1.45 $CaCO_3$ L^{-1}). Acidification of whey might thus

have led to the production of VFAs, as seen in both control and thermal pre-treatment pH 9.0. Table

4 shows the results of the main physicochemical analyses.

Table 4 – Initial and final results of the main physicochemical analyses performed in the AF experiment for 30 days. Batches consisted of control,

alkaline and thermal pre-treatments with initial pH 7.0 and 9.0 ($A_{PT}7$, $A_{PT}9$, $T_{PT}7$, and $T_{PT}9$) with pressurized (P) and non-pressurized (NP) headspaces.

Analyses considered are: chemical oxygen demand (COD), total organic carbon (TOC), total nitrogen (TN), total solids (TS), total volatile solids

(TVS), and pH.

Analyses (mg L-1) * × •	Time	Batch operational conditions									
		Control		APT7		APT9		TPT7		TPT9	
		P	NP	P	NP	P	NP	P	NP	P	NP
CH4•	t0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	tf	11,086.37	5.69	0.00	0.54	52.29	0.67	0.00	0.00	0.00	0.00
COD*	t0	8,368.63	8,634.33	7,336.90	11,174.50	8,583.00	15,294.50	9,325.30	8,758.00	10,000.00	9,736.00
	tf	7,391.10	11,053.99	7,694.78	10,658.81	7,610.05	10,584.72	8,859.74	14,289.56	8,779.31	13,844.98
TOC	t0	2,005.67	1,695.33	3,280.00	1,326.00	2,890.00	1,636.00	3,467.00	2,230.00	3,691.00	2,237.00
	tf	2,284.00	1,181.00	2,229.00	1,213.00	2,385.00	1,176.00	3,133.00	1,842.00	3,266.00	1,784.00
TN	t0	333.40	617.99	309.80	69.25	306.50	100.30	468.30	219.20	463.50	216.10
	tf	279.33	174.97	268.20	157.40	403.60	136.80	544.90	317.60	484.10	284.20
C: N ratio	t0	6.02:1	2.7:1	10.59:1	19.15:1	9.43:1	16.31:1	7.40:1	10.17:1	7.96:1	10.35:1
	tf	8.18:1	6.75:1	8.31:1	7.71:1	5.91:1	8.60:1	5.75:1	5.80:1	6.75:1	6.28:1
TS	t0	0.00516	5.72	0.00478	6.69	0.00539	6.71	0.00729	7.68	0.00698	8.52
	tf	0.00323	5.50	0.00344	5.08	0.00330	4.69	0.00376	6.13	0.00401	5.64
TVS	t0	0.00453	4.98	0.00421	5.87	0.00468	5.84	0.00578	6.81	0.00600	7.40
	tf	0.00264	4.66	0.00278	4.23	0.00260	3.99	0.00311	5.10	0.00324	4.58
pH	t0	8.25	8.25	7.00	7.00	9.00	9.00	7.00	7.00	9.00	9.00
	tf	4.76	5.14	4.65	4.99	5.00	5.01	4.87	5.11	5.06	5.19

3.1.3 Inoculum thermal pre-treatment, initial pH, and non-pressurized headspace acted as selective pressure mechanisms on microbial communities during cheese whey acidogenic fermentation

In the replicate experiment performed in TU Delft, both control and inoculum thermal pre-treated microbial communities' composition and evolution were determined through 16S rRNA gene amplicon sequencing (Figures 3 and 4). Initial microbial communities were highly similar, with few quantitative and only minor qualitative differences being observed. They both shared 7 of their 8 main detected phyla, 10 of their 11 main detected classes and 18 of their 19 main detected orders, as well as their 10 and 5 most important detected families and genera, respectively

As shown in figures 3 and 4, the common genera included fermenters and VFAs producers (such as *Sedimentibacter* (Lin et al., 2019), *Longilinea*, *Leptolinea* (Dai et al., 2016), and *Christensenellaceae* R-7 group (Y. Gao et al., 2022), the lipid-accumulating and bulk and foaming *Candidatus Microthrix* (Sheik et al., 2016), microorganisms involved in sulfur and nitrogen metabolisms (*Magnetovibrio* (Bazylinski et al., 2013), *Sulfurovum* and *Sulfurimonas* (Li et al., 2018), as well as the acetoclastic methanogen *Methanosaeta* (Stams et al., 2019).

The only noticeable difference consisted of *Cyanobacteria* being detected in the inoculum thermal pre-treated population (Figure 4). Cyanobacteria are aerobic photosynthetic microorganisms (Stal and Moezelaar, 1997). However, they display mechanisms that enable them to survive in unfavourable conditions such as in hypoxic or anaerobic (Li et al., 2018) environments. Although chlorophyll (Chl) deficiency can be caused by low levels of oxygen, these microorganisms can regulate Chl production by activating oxygen-independent oxidases or by inducing the transcription of genes that encodes enzymes that work in microoxic conditions (Fujita et al., 2015). In the given scenario, facultative anoxygenic photosynthesis uses sulfide as the electron donor with photosystem I-driven photoassimilation (Hamilton et al., 2018).

Control and inoculum thermal pre-treated microbial communities evolved differently along the

process. The final populations presented highly similar compositions at the phylum, class, order,

and family levels, sharing all or almost all the main detected taxons with only quantitative

differences being perceived. Noticeable qualitative and quantitative differences were however

observed at the genus level (Figures 3 and 4). Figure 5 depicts the beta diversity analysis of the

samples.

In the control final microbial community, an increase was observed in fermenters, such as

Longilinea and *Leptolinea* (Dai et al., 2016), and in microorganisms involved in sulfur and nitrogen

metabolisms, such as *Magnetovibrio* (Bazylinski et al., 2013), *Sulfurovum* and *Sulfurimonas* (Li et

al., 2018). *Arcobacter* (S. Gao et al., 2022), a sulfide oxidizer and denitrifier, also appeared and

became the second genus of importance, while the two VFAs producers *Sedimentibacter* (Lin et al.,

2019) and *Christensenelaceae* R-7 group (Y. Gao et al., 2022) strongly decreased.

In opposition, in the inoculum thermal pre-treated final microbial community, an increase was

observed in VFAs and alcohols producers, such as *Clostridium* (Mockaitis et al., 2022), *Fonticella*

(Joshi et al., 2021), and *Intestinimonas* (Kang et al., 2021), while fermenters such as *Longilinea* and

Leptolinea strongly decreased. The lipid-accumulating and bulk and foaming *Candidatus*

Microthrix (Sheik et al., 2016) and the *Cyanobacteria Synechocystis* (Fujita et al., 2015) increased,

and *Magnetovibrio* (Bazylinski et al., 2013), *Sulfurovum* and *Sulfurimonas* (Li et al., 2018),

involved in sulfur and nitrogen metabolisms, slightly decreased.

Those results could explain the differences in performance observed in both processes. It is

likely that methanogenesis was halted in the inoculum thermal pre-treated assay by the strong

decrease observed in microorganisms such as *Longilinea* and *Leptolinea*, which are fermenters

known to provide substrates to methanogens (Yamada et al., 2007, 2006), concomitantly with the

strong increase in VFAs and alcohols producers.

The decrease, even if limited, observed in *Methanosaeta* tends to confirm that methanogens were

inhibited. The presence of *Methanosaeta* could be explained by several hypotheses: (*i*) samples

detected were dead biomass or spores, (*ii*) inoculum pre-treatments were sufficient to inhibit or

diverge the metabolic pathway for methane formation, or (*iii*) they could have been outcompeted by

acetate fermenters (Callbeck et al., 2019; Miñana-Galbis et al., 2002; Nuppunen-Puputti et al.,

2018; Shi et al., 2020). Further -omics studies can clarify this finding. The presence of denitrifiers

could be a response to batches oxygen limitation since they are aerobic facultative (Skiba, 2008).

They can also compete for nitrate content in cheese whey (Oliveira et al., 1995).

Initial WPC40 carbohydrate concentration was approximately 260 $mg_{glucose}$ L^{-1}, stabilizing at

65 $mg_{glucose}$ L^{-1} on the fourth day. The acidogenesis rate of 1.9 d^{-1}, was close to the typical rate of 2

d^{-1} observed in similar studies(van Lier et al., 2008). COD removal and volatile solids (VS) showed

similar patterns, with sCOD decreasing from 4.2 to 2.4 g L^{-1} and VS decreasing from 4.92 to 4.74 g

L^{-1}. This was likely due to biomass production and organic matter oxidation. Control batches

showed a decrease from 4.92 to 4.74 g L^{-1}, corroborating with COD removal results. Cheese whey

alkalinity varied by 6% in all batches (1.53 to 1.45 $CaCO_3$ L^{-1}). Acidification of whey might thus

have led to the production of VFAs, as seen in both control and thermal pre-treatment pH 9.0.

Table 5 – VFAs spectra of replicated control and thermal pre-treatment (pH 9.0) acidogenic
fermentation experiment compared to the original experiment for 10 days. Replicate results are
expressed as a means average of the triplicate assays. (--) The product was not evaluated.

| | | VFAs Production (mg COD compound L^{-1}) | | | |
| | | UNICAMP | | TU Delft | |
Products	Time	C	T	C	T
Acetic acid	t0	154.34	308.26	0.00	19.28
	tf	1,037.34	1,361.07	319.13	1,407.90
Butyric acid	t0	0.00	0.00	0.00	0.00
	tf	187.29	1121.49	5.35	106.91
Iso-butyric acid	t0	0.83	0.00	16.04	149.67
	tf	0.00	0.00	16.04	58.80
Caproic acid	t0	0.00	0.00	290.94	179.70
	tf	0.00	0.00	479.20	2,587.25
Iso-caproic acid	t0	--	--	0.00	0.00
	tf	--	--	0.00	42.79
Ethanol	t0	0.00	0.00	--	--
	tf	0.00	155.07	--	--
Formic acid	t0	0.00	0.00	4.30	4.30
	tf	0.00	0.00	0	0.00
Lactic acid	t0	0.00	1,137.59	--	--
	tf	0.00	0.00	--	--
Malic acid	t0	0.00	24.74	--	--
	tf	0.00	0.00	--	--
Propionic acid	t0	0.00	0.00	11.19	11.19
	tf	161.48	55.21	85.76	55.93
Valeric acid	t0	0.00	0.00	0.00	6.94
	tf	0.00	0.00	20.83	27.78
Iso-valeric acid	t0	0.00	0.00	0.00	20.83
	tf	46.54	235.89	20.83	83.34

Figure 3 – Microbial communities' compositions and evolution. The 20 most abundant archaeal and bacterial genera identified through 16S rRNA gene amplicon sequencing. A microbial population from the F/M ratios experiments (FM). B, microbial populations from the control and thermal pre-treated inoculum batches ($T_{PT}9$ NP).

Figure 4 – ASV abundance. The 50 most abundant archaeal and bacterial ASVs detected by the AmpliconTagger pipeline. FM: F/M ratios experiments; $T_{PT}NP9$: thermal pre-treated inoculum batches.

3.2 Impact of F/M ratio on VFAs production during acidogenic fermentation of cheese whey

3.2.1 Lower F/M ratio is more efficient for VFAs production

Four different F/M ratios of WPC40 were evaluated in an experiment performed with a thermal

pre-treated inoculum at initial pH of 9.0 and under non-pressurized conditions. Ratios were F/M

0.5, F/M 1.0, F/M 2.0, and F/M 4.0, corresponding to 0.5, 1.0, 2.0, and 4.0 g COD g VSS^{-1},

respectively.

The F/M 0.5 ratio acted as a positive control since the thermal batch in the replicate experiment

presented the same F/M ratio of 0.5 g COD g VSS^{-1}. About 95% of carbohydrates were consumed

by the third day, while substrate acidification started in the first couple of days F/M 0.5 (1,056 mg

$COD_{compound}$ L^{-1} d^{-1}), F/M 1.0 (789 mg $COD_{compound}$ L^{-1} d^{-1}), F/M 0.2 (486 mg $COD_{compound}$ L^{-1} d^{-1}nd

F/M 4.0 (390 mg $COD_{compound}$ L^{-1} d^{-1}).

As shown in Table 6, F/M 0.5 ratio showed the best VFA production (3,317.88 mg $COD_{compound}$ L^{-1}

), whereas the F/M ratio 2.0 (345.93 mg L^{-1}) had the lowest. Non-pressurized thermal pre-treatment

batch in the replicate experiment had a VFAs production of 2,568.26 mg L^{-1}. There was no

production of methane in all samples which corroboratto the results observed in previous

experiments.

Table 6 – VFAs spectra of F/M ratios experiments. F/M 0.5, F/M 1.0, F/M 2.0, and F/M 4.0: 0.5,
1.0, 2.0 and 4.0 g COD g VSS^{-1}, respectively. F/M 0.5 acted as a positive control as it has the same
0.5 g COD g VSS^{-1} ratio as the replicated thermal pre-treatment batch in section 3.2. The
experiment was performed in triplicate and lasted 10 days.

VFAs Production (mg CODcompound L-1)					
Products	Time	F/M 0.5	F/M 1.0	F/M 2.0	F/M 4.0
Acetic acid	t0	34.02	9.19	0.00	0.00
	tf	900.48	956.51	493.42	865.84
Butyric acid	t0	0.00	0.00	0.00	0.00
	tf	349.75	25.26	34.64	36.48
Iso-butyric acid	t0	1,707.03	78.98	60.70	25.58
	tf	43.22	38.89	19.32	12.99
Caproic acid	t0	154.28	30.42	0.00	0.00
	tf	1,823.41	1,042.24	3,620.88	194.20
Iso-caproic acid	t0	0.00	0.00	0.00	0.00
	tf	0.00	0.00	0.00	0.00
Formic acid	t0	1.85	2.21	2.89	3.33
	tf	0.00	0.00	0.30	0.00
Propionic acid	t0	5.26	0.00	0.00	0.00
	tf	66.33	34.40	7.27	23.27
Valeric acid	t0	9.79	0.00	0.00	0.00
	tf	45.21	0.00	27.19	4.90
Iso-valeric acid	t0	32.61	17.61	0.00	0.00
	tf	89.48	58.23	40.31	27.40
Total production		3,317.88	2,155.53	4,211.46	1,165.08

3.2.2   F/M ratios affected microbial communities' evolution

Microbial communities evolved differently during the F/M experiments. As shown in Figure 5,

increasing F/M ratios significantly impacted the microbial populations' evolution, with the

strongest impact observed at the highest ratio of 4.0. Under such conditions, the microbial

population was ultra-dominated by a single genus, *Clostridium* (Figures 3 and 4). At lower F/M

ratios, the community was mainly composed of the same four genera, which altogether represented

84.3%, 77.6%, and 74.7% of the F/M-0.5, F/M-1, and F/M-2 populations, respectively (Figure 3

and 4). Those four genera were *Gemmobacter*, a poly-hydroxybutyrate-accumulating and

denitrifying bactéria (Wang et al., 2019), *Clostridium*, a known fermenter and VFAs producer

(Mockaitis et al., 2022), *Lachnoclostridium*, a known butyrate producer (Detman et al., 2021), and

Rhodopseudomonas, a highly metabolically versatile bacteria capable of hydrogen production

through nitrogen fixation or poly-hydroxybutyrate production (Ranaivoarisoa et al., 2019). The

operational conditions applied in the F/M-0.5 experiment were thus apparently those that favor the

best microbial combination to produce the highest level of VFAs. Further characterizations (i.e.,

metagenomics and/or metatranscriptomics analyses) have to be performed to better understand why

such close microbial populations could generate different amounts of VFAs.

Figure 5 – Beta diversity analysis of the microbial populations. Principal Coordinates Analysis
computed from Bray-Curtis dissimilarity distance matrix. FM: F/M ratios experiments; $T_{PT}NP9$:
thermal pre-treated inoculum batches.

4   Conclusions

AF is a successful alternative for producing VFAs of economic interest out of high-strength residues such as cheese whey. Inoculum pre-treatments can play as selective pressures on microbial communities, selecting microorganisms that can thrive in imposed conditions. The main conclusions from this work are the following:

1. Thermal pre-treatment of inocula was efficient to halt methanogenesis regardless of headspace pressure, pH, and F/M ratios.

2. Contrary to the literature, alkaline pre-treatment did not improve methanogenesis. However, non-pressurized headspace alkaline pre-treatment pH 9.0 did produce significant quantities of iso-butyric acid (7,677.30 mg COD_{IBA} L^{-1}) and ethanol (13,693.24 mg COD_{EtOH} L^{-1}). Further studies on the mechanisms of this pre-treatment are necessary to optimize the process.

3. *Methanosaeta* was present in the replicate of the control (pH 8.25) and thermal (pH 9.0) batches, despite the absence of methane production. Thermal pre-treatment could have either inactivated or destroyed this microorganism, which in its turn enabled the detection of *Methanosaeta* in the 16S rRNA gene amplicon sequencing analysis.

4. A low F/M ratio of 0.5 g COD g^{-1} VS selected a microbial community producing a high level of VFAs by AF.

5. The setup of parameters used as selective pressure during cheese whey AF must vary according to the desired end product.

6. Headspace pressure directs the metabolic pathways for VFAs and alcohol production in AF. Pressurized headspace batches had a more di while non-pressurized batches produced higher amounts of VFAs. The exception would be iso-butyric acid and ethanol produced in non-pressurized alkaline pH 9.0 batches.

7. Another important parameter was the Initial pH of 9.0. Regardless of the headspace
pressure and inoculum pre-treatment, an initial pH of 9.0 produced the greatest number of
products. The drastic pH drop influences the redox potential of the medium facilitating the
uptake of compounds while setting the grounds for ecologic relationships within the
microbial community.

This work aimed to verify if imposed parameters (i.e., headspace pressure, inoculum thermal
and alkaline pre-treatment, and initial pH) in cheese whey AF would work as selective pressures on
the inoculum microbial community. Our objectives were met when we managed to halt
methanogenesis and increase acetic acid, iso-butyric acid, and ethanol. Acetate is an organic carbon
source that can be used for different biological conversions (e.g., microalgal
photoorganoheterotrophic processes), whereas iso-butyric and ethanol are of interest of
pharmaceutical, feed, chemical, and biofuels industries.

Still, there is plenty to be understood, especially regarding the influence of these parameters
on metabolic pathways and microbial interaction. The fact that thermal pre-treatment time and
temperature were optimized and methanogenesis was halted, despite the presence of methanogens.
Understanding the mechanism behind this can overcome competition for substrate.

Thermal inoculum pre-treatment can be used in short fermentation processes. Nevertheless,
the interaction among the factors with different substrates, inoculum and reactor configurations
must be studied to certify thermal pre-treatment efficiency. Thus, it is necessary to evaluate other
configurations for process implementation and optimization.

Understanding the VFAs production and consumption mechanisms is crucial for driving the
spectrum of fermentation products to desired compounds. Future 16S rRNA metagenomics analyses
will be fundamental to decrease these knowledge gaps.

5 Acknowledgements

This work was funded in major CAPES PDS scholarship (CAPES PDS 88882.435082/2019-01) and CNPq (CNPq 166460/2017-6). The work at the TU Delft was funded by the start-up grant of the Department of Biotechnology of the TU Delft (Prof. Dr. David Weissbrodt, PI). The lab works at UNICAMP benefited from the assistance of Vítor Augusto de Oliveira, Juliana Martins Valença, Giovani Archanjo Brotto and Rosa Helena Aguiar while the lab works in the TU Delft benefited from the assistance of Cor Ras, Johan Knoll, Marcel Langerveld, Ben Abbas and Armand Middeldorp.

6 Conflict of interest statement

The authors share no conflict of interest.

7 Authors' contributions

M.P.G. d A conceptualized the experiment and wrote the manuscript with direct core inputs by D.G.W and G.M. M.P.G.dA performed the analysis for the acidogenic fermentation in Brazil and in The Netherlands, the latter with the assistance of C.M. The analysis and conceptualisation the F/M experiments were performed by C.M. The experiment was designed by MPGdA, DGW and GM by confronting ideas, concepts and solutions to technological, outcomes on microbial communities genomics were provided by G.B and J.T. All authors read, edited, and provided critical feedback to the manuscript.

8 References

Abbasi, T., Tauseef, S.M., Abbasi, S.A., 2012. Biogas energy, in: Abbasi, Tasneem, Tauseef, S.M., Abbasi, S.A. (Ed.), Biogas Energy. Springer, pp. 1–169. https://doi.org/10.1007/978-1-4614-1040-9

Alibardi, L., Favaro, L., Lavagnolo, M.C., Basaglia, M., Casella, S., 2012. Effects of heat treatment
on microbial communities of granular sludge for biological hydrogen production. Water
Science and Technology 66, 1483–1490. https://doi.org/10.2166/wst.2012.336

Angenent, L.T., Richter, H., Buckel, W., Spirito, C.M., Steinbusch, K.J., Strik, D.P.B.T.B.,
Grootscholten, T.I.M., Buisman, C.J.N., Hamelers, H.V.M., 2016. Chain elongation with
reactor microbiomes: open- culture biotechnology to produce biochemicals Extensive review
Chain elongation with reactor microbiomes□: open-culture biotechnology to produce
biochemicals. Environ Sci Technol 50, 2796–2810. https://doi.org/10.1021/acs.est.5b04847

APHA, 2005. Standard Methods for the Examination of Water and Wastewater, 21st ed. American
Public Health Association, Washington, DC.

Appels, L., Lauwers, J., Degrve, J., Helsen, L., Lievens, B., Willems, K., van Impe, J., Dewil, R.,
2011. Anaerobic digestion in global bio-energy production: Potential and research challenges.
Renewable and Sustainable Energy Reviews 15, 4295–4301.
https://doi.org/10.1016/j.rser.2011.07.121

ASTM International, 2016. ASTM D8083-16 - Standard Test Method for Total Nitrogen, and Total
Kjeldahl Nitrogen (TKN) by Calculation, in Water by High Temperature Catalytic Combustion
and Chemiluminescence Detection. ASTM Standard 1–9.

Bazylinski, D.A., Williams, T.J., Lefèvre, C.T., Trubitsyn, D., Fang, J., Beveridge, T.J., Moskowitz,
B.M., Ward, B., Schübbe, S., Dubbels, B.L., Simpson, B., 2013. *Magnetovibrio blakemorei*
gen. nov., sp. nov., a magnetotactic bacterium (Alphaproteobacteria□*: Rhodospirillaceae*)
isolated from a salt marsh. Int J Syst Evol Microbiol 63, 1824–1833.
https://doi.org/10.1099/ijs.0.044453-0

Callbeck, C.M., Pelzer, C., Lavik, G., Ferdelman, T.G., Graf, J.S., Vekeman, B., Schunck, H.,
Littmann, S., Fuchs, B.M., Hach, P.F., Kalvelage, T., Schmitz, R.A., Kuypers, M.M.M., 2019.
Arcobacter peruensis sp. nov., a chemolithoheterotroph isolated from sulfide-and organic-rich
coastal waters off Peru. Appl Environ Microbiol 85, 1–17.
https://doi.org/10.1128/AEM.01344-19

Casallas-Ojeda, M.R., Marmolejo-Rebellón, L.F., Torres-Lozada, P., 2020. Evaluation of
simultaneous incidence of head space and temperature on biochemical methane potential in
food waste. Cogent Eng 7, 1729514. https://doi.org/10.1080/23311916.2020.1729514

Chen, Y., Cheng, J.J., Creamer, K.S., 2008. Inhibition of anaerobic digestion process. Bioresource
Technologyr 99, 4.

Chen, Y., Jiang, S., Yuan, H., Zhou, Q., Gu, G., 2007. Hydrolysis and acidification of waste
activated   sludge   at   different   pHs.   Water   Res   41,   683–689.
https://doi.org/10.1016/j.watres.2006.07.030

Corti, A., Lombardi, L., 2007. Anaerobic co-digestion of source selected organic waste and sewage
sludge, in: 11th International Symposium on Waste Management and Sustainable Landfilling.
Sardinia, pp. 1–5.

Dai, Y., Yan, Z., Jia, L., Zhang, S., Gao, L., Wei, X., Mei, Z., Liu, X., 2016. The composition,
localization and function of low-temperature-adapted microbial communities involved in
methanogenic degradations of cellulose and chitin from Qinghai-Tibetan Plateau wetland soils.
J Appl Microbiol 121, 163–176. https://doi.org/10.1111/jam.13164

Dareioti, M.A., Vavouraki, A.I., Kornaros, M., 2014. Effect of pH on the anaerobic acidogenesis of
agroindustrial wastewaters for maximization of bio-hydrogen production: A lab-scale
evaluation   using   batch   tests.   Bioresour   Technol   162,   218–227.
https://doi.org/10.1016/j.biortech.2014.03.149

Darvekar, P., Liang, C., Karim, M.N., Holtzapple, M.T., 2019. Effect of headspace gas composition
on carboxylates production in open-culture fermentation of corn stover. Biomass Bioenergy
126, 57–61. https://doi.org/10.1016/j.biombioe.2019.04.019

Detman, A., Laubitz, D., Chojnacka, A., Kiela, P.R., Salamon, A., Barberán, A., Chen, Y., Yang,
F., Błaszczyk, M.K., Sikora, A., 2021. Dynamics of dark fermentation microbial communities
in   the   light   of   lactate   and   butyrate   production.   Microbiome   9,   158.
https://doi.org/10.1186/s40168-021-01105-x

Deublein, D., Steinhauser, A. (Eds.), 2011. Biogas from renewable resources: An introduction, 2nd
ed. Wiley-VCH Verlag GmbH & Co. KGaA.

Dilallo, R., Albertison, O.E., 1961. Volatile Acids by Direct Titration. J Water Pollut Control Fed
33, 356–365.

Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A., Smith, F., 1956. Colorimetric Method for
Determination of Sugars and Related Substances. Anal Chem 28, 350–356.

Food and Agriculture Organization of the United Nations, n.d. FAOSTAT - Livestock processed
[WWW Document]. URL http://www.fao.org/faostat/en/?#data/QP (accessed 6.20.20).

Fujita, Y., Tsujimoto, R., Aoki, R., 2015. Evolutionary Aspects and Regulation of Tetrapyrrole
Biosynthesis in Cyanobacteria under Aerobic and Anaerobic Environments. Life 5, 1172–
1203. https://doi.org/10.3390/life5021172

Gao, S., Li, Z., Hou, Y., Wang, A., Liu, Q., Huang, C., 2022. Effects of different carbon sources on
the efficiency of sulfur-oxidizing denitrifying microorganisms. Environ Res 204, 111946.
https://doi.org/10.1016/j.envres.2021.111946

Gao, Y., Guo, L., Jin, C., Zhao, Y., Gao, M., She, Z., Wang, G., 2022. Metagenomics and network
analysis elucidating the coordination between fermentative bacteria and microalgae in a novel
bacterial-algal coupling reactor (BACR) for mariculture wastewater treatment. Water Res 215,
118256. https://doi.org/10.1016/j.watres.2022.118256

Goud, R.K., Mohan, S.V., 2012. Acidic and alkaline shock pretreatment to enrich acidogenic
biohydrogen producing mixed culture: long term synergetic evaluation of microbial inventory,
dehydrogenase   activity   and   bio-electro   kinetics.   RSC   Adv   2,   6336–6353.
https://doi.org/10.1039/c2ra20526b

Hamilton, T.L., Klatt, J.M., de Beer, D., Macalady, J., 2018. Cyanobacterial photosynthesis under
sulfidic conditions: insights from the isolate *Leptolyngbya* sp. strain hensonii. ISME J 12, 568–
584. https://doi.org/https://doi.org/10.1038/ismej.2017.193

Joshi, S., Robles, A., Aguiar, S., Delgado, A.G., 2021. The occurrence and ecology of microbial
chain   elongation   of   carboxylates   in   soils.   ISME   J   15,   1907–1918.
https://doi.org/10.1038/s41396-021-00893-2

Julien Tremblay, 2020. microbiomeutils 0.9.4: Microbiome utilities [WWW Document]. URL
https://github.com/jtremblay/microbiomeutils

Julien Tremblay, 2019. AmpliconTagger Pipeline Databases [WWW Document].

Kang, D., Saha, S., Kurade, M.B., Basak, B., Ha, G.-S., Jeon, B.-H., Lee, S.S., Kim, J.R., 2021.
Dual-stage pulse-feed operation enhanced methanation of lipidic waste during co-digestion
using acclimatized consortia. Renewable and Sustainable Energy Reviews 145, 111096.
https://doi.org/10.1016/j.rser.2021.111096

Kigozi, R., Muzenda, E., Aboyade, A.O., 2014. Biogas technology: Current trends, opportunities
and challenges, in: 6th Int'l Conf. on Green Technology, Renewable Energy & Environmental
Engg. (ICGTREEE'20114). Cape Town, pp. 311–317.

Kleerebezem, R., Joosse, B., Rozendal, R., van Loosdrecht, M.C.M., 2015. Anaerobic digestion
without biogas? Rev Environ Sci Biotechnol 14, 787–801. https://doi.org/10.1007/s11157-015-
9374-6

Li, H., Li, C., Liu, W., Zou, S., 2012. Optimized alkaline pretreatment of sludge before anaerobic
digestion. Bioresour Technol 123, 189–194. https://doi.org/10.1016/j.biortech.2012.08.017

Li, Y., Tang, K., Zhang, L., Zhao, Z., Xie, X., Chen, C.-T.A., Wang, D., Jiao, N., Zhang, Y., 2018.
Coupled Carbon, Sulfur, and Nitrogen Cycles mediated by microorganisms in the water
column of a shallow-water hydrothermal ecosystem. Front Microbiol 9.
https://doi.org/10.3389/fmicb.2018.02718

Lin, J.-G., Chang, C.-N., Chang, S.-C., 1997. Enhancement of anaerobic digestion of waste
activated sludge by alkaline solubilization. Bioresour Technol 62, 85–90.

Lin, X.-Q., Li, Z.-L., Liang, B., Zhai, H.-L., Cai, W.-W., Nan, J., Wang, A.-J., 2019. Accelerated
microbial reductive dechlorination of 2,4,6-trichlorophenol by weak electrical stimulation.
Water Res 162, 236–245. https://doi.org/10.1016/j.watres.2019.06.068

Liu, Y., Zhang, Z., Jiang, W., Gu, Y., 2022. Protein acetylation-mediated cross regulation of acetic
acid and ethanol synthesis in the gas-fermenting *Clostridium ljungdahlii*. Journal of Biological
Chemistry 298. https://doi.org/10.1016/j.jbc.2021.101538

López-Torres, M., Llórens, Ma. del C.E., 2016. Effect of alkaline pretreatment on anaerobic
digestion of olive mill solid waste. Waste Management 58, 160–168.
https://doi.org/10.1016/j.wasman.2016.08.008

Miñana-Galbis, D., Farfán, M., Lorén, J.G., Fusté, M.C., 2002. Biochemical identification and
numerical taxonomy of *Aeromonas* spp. isolated from environmental and clinical samples in
Spain. J Appl Microbiol 93, 420–430. https://doi.org/10.1046/j.1365-2672.2002.01711.x

Mockaitis, G., Bruant, G., Foresti, E., Zaiat, M., Guiot, S.R., 2022. Physicochemical pretreatment
selects microbial communities to produce alcohols through metabolism of volatile fatty acids.
Biomass Convers Biorefin. https://doi.org/10.1007/s13399-022-02383-7

Mockaitis, G., Bruant, G., Guiot, S.R., Peixoto, G., Foresti, E., Zaiat, M., 2020a. Acidic and thermal
pre-treatments for anaerobic digestion inoculum to improve hydrogen and volatile fatty acid
production using xylose as the substrate. Renew Energy 1388–1398.

Mockaitis, G., Ratusznei, S.M., Rodrigues, J.A.D., Zaiat, M., Foresti, E., 2006. Anaerobic whey
treatment by a stirred sequencing batch reactor (ASBR): effects of organic loading and
supplemented alkalinity. J Environ Manage 79, 198–206.
https://doi.org/10.1016/j.jenvman.2005.07.001

Mulders, M., Tamis, J., Abbas, B., Sousa, J., Dijkman, H., Rozendal, R., Kleerebezem, R., 2020.
Pilot-scale polyhydroxyalkanoate production from organic waste: Process characteristics at
high pH and high ammonium concentration. Journal of Environmental Engineering 146,
4020049. https://doi.org/10.1061/(ASCE)EE.1943-7870.0001719

Navia, R., Vidal, G., 2002. Alkaline pretreatment of kraft mill sludge to improve its anaerobic
digestion alkaline pretreatment of kraft mill sludge to improve its anaerobic digestion. Bull.
Environ. Contam. Toxicol. 69, 869–876. https://doi.org/10.1007/s00128-002-0140-4

Nuppunen-Puputti, M., Purkamo, L., Kietäväinen, R., Nyyssönen, M., Itävaara, M., Ahonen, L.,
Kukkonen, I., Bomberg, M., 2018. Rare biosphere archaea assimilate acetate in Precambrian
terrestrial subsurface at 2.2 km depth. Geosciences (Switzerland) 8, 1–20.
https://doi.org/10.3390/geosciences8110418

Oliveira, C.P., Gloria, M.B.A., Barbour, J.F., Scanlan, R.A., 1995. Nitrate, nitrite, and volatile
nitrosamines in whey-containing food products. J Agric Food Chem 43, 967–969.
https://doi.org/10.1021/jf00052a023

Ostrem, K.M., Millrath, K., Themelis, N.J., 2004. Combining anaerobic digestion and waste-to-
energy, in: Proceedings of 12TH Annual North American Waste to Energy Conference,
NAWTEC12. pp. 265–271. https://doi.org/10.1115/nawtec12-2231

Pang, N., Gu, X., Kirchhoff, H., Lei, H., Roje, S., 2019. Exploiting mixotrophic for improving
productivities of biomass and co-products of microalgae. Renewable and Sustainable Energy
Reviews 112, 450–460. https://doi.org/10.1016/j.rser.2019.06.001

Peixoto, G., Saavedra, N.K., Varesche, M.B.A., Zaiat, M., 2011. Hydrogen production from soft-
drink wastewater in an upflow anaerobic packed-bed reactor. Int J Hydrogen Energy 36, 8953–
8966. https://doi.org/10.1016/j.ijhydene.2011.05.014

Penteado, E.D., Adorno, M.A.T., Zaiat, M., 2012. Simple and accurated method for determination
of fatty acids, alcohols andcarbohydrates by HPLC with UV/DAD and RID detectors, in: 38th
International Symposium on High Performance Liquid Phase Separation. Anaheim, California,
USA.

Pretorius, W.A., 1987. A conceptual basis for microbial selection in biological wastewater
treatment. Water Res 21, 891–894.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.,
2013. The SILVA ribosomal RNA gene database project: Improved data processing and web-
based tools. Nucleic Acids Res 41, D590–D596. https://doi.org/10.1093/nar/gks1219

Ramos-Suarez, M., Zhang, Y., Outram, V., 2021. Current perspectives on acidogenic fermentation
to produce volatile fatty acids from waste. Rev Environ Sci Biotechnol 20, 439–478.
https://doi.org/10.1007/s11157-021-09566-0

Ranaivoarisoa, T.O., Singh, R., Rengasamy, K., Guzman, M.S., Bose, A., 2019. Towards
sustainable bioplastic production using the photoautotrophic bacterium *Rhodopseudomonas*
palustris TIE-1. J Ind Microbiol Biotechnol 46, 1401–1417. https://doi.org/10.1007/s10295-
019-02165-7

Ripley, L.E., Boyle, W.C. and Converse, J.C., 1986. Improved alkalimetric monitoring for
anaerobic digestion of high-strength wastes. Journal WPCF 48, 406–411.

Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: A versatile open-
source tool for metagenomics. PeerJ 2016. https://doi.org/10.7717/peerj.2584

Rombouts, J.L., Mos, G., Weissbrodt, D.G., Kleerebezem, R., van Loosdrecht, M.C.M., 2019.
Diversity and metabolism of xylose and glucose fermenting microbial communities in
sequencing batch or continuous culturing. FEMS Microbiol Ecol 95, 1–12.
https://doi.org/10.1093/femsec/fiy233

Saary, P., Forslund, K., Hildebrand, F., 2017. RTK: efficient rarefaction analysis of large datasets.
Bioinformatics 33, 2594–2595. https://doi.org/10.1093/bioinformatics/xxxxx

Saddoud, A., Hassaïri, I., Sayadi, S., 2007. Anaerobic membrane reactor with phase separation for
the treatment of cheese whey. Bioresour Technol 98, 2102–2108.
https://doi.org/10.1016/j.biortech.2006.08.013

Sarkar, O., Butti, S.K., Venkata Mohan, S., 2017. Acidogenesis driven by hydrogen partial pressure
towards bioethanol production through fatty acids reduction. Energy 118, 425–434.
https://doi.org/10.1016/j.energy.2016.12.017

Sarkar, O., Rova, U., Christakopoulos, P., Matsakas, L., 2021. Influence of initial uncontrolled pH
on acidogenic fermentation of brewery spent grains to biohydrogen and volatile fatty acids
production: Optimization and scale-up. Bioresour Technol 319.
https://doi.org/10.1016/j.biortech.2020.124233

Shah, F.A., Mahmood, Q., Rashid, N., Pervez, A., Raja, I.A., Shah, M.M., 2015. Co-Digestion,
pretreatment and digester design for enhanced methanogenesis. Renewable and Sustainable
Energy Reviews 42, 627–642. https://doi.org/10.1016/j.rser.2014.10.053

Sheik, A.R., Muller, E. el, Audinot, J.-N., Lebrun, L.A., Grysan, P., Guignard, C., Wilmes, P.,
2016. *In situ* phenotypic heterogeneity among single cells of the filamentous bacterium
Candidatus Microthrix parvicella. ISME J 10, 1274–1279.
https://doi.org/10.1038/ismej.2015.181

Shi, L.L., Da, Y.Y., Zheng, W.T., Chen, G.Q., Li, Z.J., 2020. Production of polyhydroxyalkanoate
from acetate by metabolically engineered *Aeromonas hydrophilia*. J Biosci Bioeng 130, 290–
294. https://doi.org/10.1016/j.jbiosc.2020.05.003

Skiba, U., 2008. Denitrification, in: Jorgensen, S.E., Fath, B.D. (Eds.), Encyclopedia of Ecology.
Oxford, Elsevier, pp. 866–871.

Smithers, G.W., 2008. Whey and whey proteins-From "Gutter-to-Gold." Int Dairy J 18, 695–704.
https://doi.org/10.1016/j.idairyj.2008.03.008

Stal, L.J., Moezelaar, R., 1997. Fermentation in cyanobacteria. FEMS Microbiol Rev 21, 179–211.

Stams, A.J.M., Teusink, B., Sousa, D.Z., 2019. Ecophysiology of Acetoclastic Methanogens, in:
Biogenesis of Hydrocarbons. Springer International Publishing, Cham, pp. 1–14.
https://doi.org/10.1007/978-3-319-53114-4_21-1

Thatikayala, D., Pant, D., Min, B., 2021. MnO_2/reduced graphene oxide nanohybrids as a cathode
catalyst for the microbial reduction of CO_2 to acetate and isobutyric acid. Sustainable Energy
Technologies and Assessments 45. https://doi.org/10.1016/j.seta.2021.101114

Torres, P.L., 1992. Desempenho de um reator anaerobio de manta de lodo (uasb) de bancada no
tratamento de substrato sintetico simulando esgotos sanitarios (Doctorate). Universidade de
Sãp Paulo, São Carlos.

Tsakali, E., Petrotos, K., d'Allessandro, A., Goulas, P., 2010. A review on whey composition and
the methods used for its utilization for food and pharmaceutical products, in: Centre, C.R.
(Ed.), 6[th] International Conference on Simulation and Modelling in the Food and Bio-Industry.
Braganca, Portugal, pp. 195–201.

USDA - Foreign Agricultural Service, 2022. Dairy: World markets and trade.

van Lier, J.B., Mahmoud, N., Zeeman, G., 2008. Anaerobic wastewater treatment, in: Henze, M.,
Loosdrecht, M.C.M. van, Ekama, G.A., Brdjanovic, D. (Eds.), Biological Wastewater
Treatment: Principles, Modelling and Design. IWA Publishing, London, pp. 401–442.
https://doi.org/10.1021/es00154a002

van Lier, J.B., Tilche, A., Ahring, B.K., Macarie, H., Moletta, R., Dohanyos, M., Pol, L.W.H.,
Lens, P., Verstraete, W., 2001. New perspectives in anaerobic digestion. Water Science and
Technology 43, 1–18.

Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naïve Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. Appl Environ Microbiol 73,
5261–5267. https://doi.org/10.1128/AEM.00062-07

Wang, S., Zhao, J., Ding, X., Zhao, R., Huang, T., Lan, L., Naim Bin Nasry, A. al, Liu, S., 2019.
Effect of starvation time on NO and N$_2$O production during heterotrophic denitrification with
nitrite   and   glucose   shock   loading.   Process   Biochemistry   86,   108–116.
https://doi.org/10.1016/j.procbio.2019.07.023

Weissbrodt, D.G., Wells, G.F., Laureni, M., Agrawal, S., Goel, R., Russo, G., Men, Y., Johnson,
D.R., Christensson, M., Lackner, S., Joss, A., Lund Nielsen, J., Bürgmann, H., Morgenroth, E.,
Nielsen, L., 2020. Systems microbiology and Engineering of aerobic-anaerobic ammonium
oxidation. ChemRxiv preprint 1–34. https://doi.org/10.26434/chemrxiv.12243077.v1

Wonglertarak, W., Wichitsathian, B., 2014. Alkaline Pretreatment of Waste Activated Sludge in
Anaerobic   Digestion.   Journal   of   Clean   Energy   Technologies   118–121.
https://doi.org/10.7763/jocet.2014.v2.104

Yamada, T., Imachi, H., Ohashi, A., Harada, H., Hanada, S., Kamagata, Y., Sekiguchi, Y., 2007.
Bellilinea caldifistulae gen. nov., sp. nov. and *Longilinea arvoryzae* gen. nov., sp. nov., strictly
anaerobic, filamentous bacteria of the phylum *Chloroflexi* isolated from methanogenic
propionate-degrading   consortia.   Int   J   Syst   Evol   Microbiol   57,   2299–2306.
https://doi.org/10.1099/ijs.0.65098-0

Yamada, T., Sekiguchi, Y., Hanada, S., Imachi, H., Ohashi, A., Harada, H., Kamagata, Y., 2006.
Anaerolinea thermolimosa sp. nov., *Levilinea saccharolytica* gen. nov., sp. nov. and
Leptolinea tardivitalis gen. nov., sp. nov., novel filamentous anaerobes, and description of the
new classes *Anaerolineae* classis nov. and *Caldilineae* classis nov. in the bacterial phylum
Chloroflexi. Int J Syst Evol Microbiol 56, 1331–1340. https://doi.org/10.1099/ijs.0.64169-0

Yu, Y., Lee, C., Kim, J., Hwang, S., 2005. Group-specific primer and probe sets to detect
methanogenic communities using quantitative Real-Time Polymerase Chain Reaction.
Biotechnol Bioeng 89, 670–679. https://doi.org/10.1002/bit.20347

Yuan, H., Chen, Y., Zhang, H., Jiang, S., Zhou, Q., Gu, G., 2006. Improved bioproduction of short-

chain fatty acids (SCFAs) from excess sludge under alkaline conditions. Environ Sci Technol

40, 2025–2029. https://doi.org/10.1021/es052252b

Zhang, S., Guo, H., Du, L., Liang, J., Lu, X., Li, N., Zhang, K., 2015. Influence of NaOH and

thermal pretreatment on dewatered activated sludge solubilisation and subsequent anaerobic

digestion: Focused on high-solid state. Bioresour Technol 185, 171–177.

Zhang, X., Shao, Y., Tian, J., Liao, Y., Li, P., Zhang, Y., Chen, J., Li, Z., 2019. PTrimmer: An

efficient tool to trim primers of multiplex deep sequencing data. BMC Bioinformatics 20.

https://doi.org/10.1186/s12859-019-2854-x

Zhou, M., Yan, B., Wong, J.W.C., Zhang, Y., 2018. Enhanced volatile fatty acids production from

anaerobic fermentation of food waste: A mini-review focusing on acidogenic metabolic

pathways. Bioresource Biotechnology 248, 68–78.

https://doi.org/https://doi.org/10.1016/j.biortech.2017.06.121