Vitamin D induces SIRT1 activation through K610 deacetylation in colon cancer

José Manuel García-Martínez[1], Ana Chocarro-Calvo[1], Javier Martínez-Useros[1,2], María Jesús
Fernández-Aceñero[3], M. Carmen Fiuza[4], Jose Cáceres-Rentero[1], Antonio De la Vieja[5,6], Antonio
Barbáchano[6,7,8], Alberto Muñoz[6,7,8], María Jesús Larriba[6,7,8] and Custodia García-Jiménez[1]*

[1] Area of Physiology, Faculty Health Sciences, University Rey Juan Carlos, 28922 Alcorcón,
Madrid, Spain.
9
10 [2] Translational Oncology Division, OncoHealth Institute, Health Research Institute-University
11 Hospital Fundación Jiménez Díaz-Universidad Autónoma de Madrid, 28040 Madrid, Spain.
12
13 [3] Department of Surgical Pathology, Hospital Clínico San Carlos, 28040 Madrid, Spain.
14
15 [4] Department of Surgery, University Hospital Fundación Alcorcón-Universidad Rey Juan Carlos,
16 28922 Alcorcón, Madrid, Spain.
17
18 [5] Unidad de Tumores Endocrinos (UFIEC), Instituto de Salud Carlos III, 28220 Majadahonda,
19 Madrid, Spain.
20
21 [6] CIBER de Cáncer, Instituto de Salud Carlos III, Madrid, Spain.
22
23 [7] Instituto de Investigaciones Biomédicas "Alberto Sols", Consejo Superior de Investigaciones
24 Científicas, Universidad Autónoma de Madrid, 28029 Madrid, Spain.
25
26 [8] Instituto de Investigación Sanitaria del Hospital Universitario La Paz, 28046 Madrid, Spain.
27
28

29 Running title: Vitamin D activates SIRT1 through deacetylation
30
Keywords: colorectal cancer; vitamin D; VDR; SIRT1; acetylation

* Correspondence: custodia.garcia@urjc.es; Tel.: +(34) 91 488 86 18

ABSTRACT

Posttranslational modifications of epigenetic modifiers provide a flexible and timely

mechanism for rapid adaptations to the dynamic environment of cancer cells. SIRT1 is an NAD^+-

dependent epigenetic modifier whose activity is classically associated with healthy aging and

longevity, but its function in cancer is not well understood. Here, we reveal that 1α,25-

dihydroxyvitamin D_3 (1,25(OH)$_2$D$_3$, calcitriol), the active metabolite of vitamin D (VD), promotes

SIRT1 activation through auto-deacetylation in human colon carcinoma cells, and identify lysine 610

as an essential driver of SIRT1 activity. Remarkably, our data show that the post-translational control

of SIRT1 activity mediates the antiproliferative action of 1,25(OH)$_2$D$_3$. This effect is reproduced by

the SIRT1 activator SRT1720, suggesting that SIRT1 activators may offer new therapeutic possibilities

for colon cancer patients who are VD deficient or unresponsive. Moreover, this might be extrapolated

to inflammation and other VD deficiency-associated and highly prevalent diseases in which SIRT1

plays a prominent role.

INTRODUCTION

Over the last decades the study of genetic lesions in tumour cells provided key insights into the deregulated signalling pathways that promote cell proliferation and invasiveness and suppress senescence. In addition, increasing evidence from epidemiological studies suggests a significant impact of non-genetic modifiable factors on the oncogenic process by affecting cell epigenetics and thus, gene expression and phenotype. The key metabolic sensor SIRT1 acts an NAD^+-dependent deacetylase and a critical post-translational modifier and epigenetic regulator. As such, SIRT1 promotes efficient energy utilisation and cellular defences in response to environmental challenges, whereas its dysregulation accelerates age-related diseases such as diabetes and cancer (Bonkowski and Sinclair, 2016).

Colorectal cancer (CRC) is the second most diagnosed malignancy in women, the third in men, and a ~~major~~ leading cause of cancer-related deaths worldwide, with an incidence estimated to increase by 60% by 2030 (Bray et al., 2018; Cancer Genome Atlas Network, 2012). Many observational/epidemiological studies suggest that vitamin D (VD) deficiency is a risk factor for developing and dying of cancer, particularly for CRC (Feldman et al., 2014; Grant et al., 2022; Kim et al., 2022, 2021), albeit data from supplementation studies in the human population are controversial. Confirmation of a clinically relevant anti-CRC effect of VD in well-designed prospective randomized studies is still pending (Chandler et al., 2020; Henn et al., 2022; Manson et al., 2020; Muñoz and Grant, 2022; Song et al., 2021)

VD in humans has two origins: synthesis in the skin by the action of solar ultraviolet radiation on 7-dehydrocholesterol and the diet. The active VD metabolite is $1\alpha,25$-dihydroxyvitamin D_3 ($1,25(OH)_2D_3$, calcitriol), which results from two consecutive hydroxylations of VD, the first in the liver and the second in the kidney or in many epithelial and immune cell types (Bikle and Christakos, 2020; Feldman et al., 2014). VD actions are mediated by $1,25(OH)_2D_3$ binding to a member of the nuclear receptor superfamily of transcription factors,

the VD receptor (VDR). Upon ligand binding, VDR regulates the expression of hundreds of target genes, many of which are involved in calcium and phosphate homeostasis, bone biology, immune response, metabolism, detoxification, and cell survival, proliferation and differentiation (Bikle and Christakos, 2020; Carlberg and Muñoz, 2022; Carlberg and Velleuer, 2022; Feldman et al., 2014; Fernández-Barral et al., 2020).

Consistent with a protective role of VD against CRC, the colonic epithelium expresses high levels of VDR. Many studies in CRC cells and associated fibroblasts and in experimental animals have shown a large series of VDR-mediated VD anti-CRC effects (Carlberg and Muñoz, 2022; Carlberg and Velleuer, 2022; Feldman et al., 2014; Fernández-Barral et al., 2020; Ferrer-Mayorga et al., 2019) In support, germline deletion of *Vdr* in mice with constitutively active Wnt/β-catenin signalling, as a main driver of CRC, results in increased intestinal tumour load (Larriba et al., 2011; Zheng et al., 2012).

Notably, certain evidence links SIRT1 and VD. First, decreased SIRT1 activity has been linked to the pathogenesis of CRC (Ren et al., 2017; Strycharz et al., 2018). However, using SIRT1 level as a marker of malignancy is controversial since both increased or decreased SIRT1 tumour expression have been described in different studies (Carafa et al., 2019; Chen et al., 2014; Firestein et al., 2008), which points to a discrepancy between SIRT1 level and activity. Second, SIRT1 deacetylates the VDR to enhance its activity in kidney and bone cells (Sabir et al., 2017) and, consequently, decreased SIRT1 activity may drive VD insensitivity. Third, promoter-bound VDR increases *SIRT1* gene expression in kidney and liver cells (Yuan et al., 2022) and thus, VD deficiency may decrease SIRT1 levels leading to reduced activity. However, whether VD alters SIRT1 activity *per se* and/or SIRT1 protein expression in CRC remains unclear.

Here we reveal that $1,25(OH)_2D_3$ induces specific SIRT1 activity at multiple levels in human CRC cells. Notably, liganded VDR post-translationally increases SIRT1 activity through auto-deacetylation. Mechanistically, SIRT1 K610 is identified as a critical player for

$1,25(OH)_2D_3$-enhanced VDR-SIRT1 interaction, SIRT1 auto-deacetylation and activation. This

occurs independently of increased SIRT1 RNA and protein expression and elevated cofactor

(NAD^+) levels. Discrepant SIRT1 level and activity may explain the controversial role of SIRT1

as tumor suppressor or tumor promoter in CRC. Our results provide a new mechanistic insight

into the association of VD deficiency with CRC and suggest potential therapeutic benefit for

SIRT1 activators in SIRT1-positive CRC.

RESULTS

$1,25(OH)_2D_3$ increases SIRT1 level via the VDR in CRC cells

The question of whether VD governs SIRT1 activity in CRC was first approached in two

well characterized CRC cell lines, HCT 116 and HT-29. Immunofluorescence analyses (Fig.

1A) revealed that SIRT1 protein level doubled in response to $1,25(OH)_2D_3$, a result confirmed

by Western blotting of nuclear extracts (Fig. 1B). The increased expression of SIRT1 protein by

$1,25(OH)_2D_3$ could be explained by a 2-fold upregulation of SIRT1 RNA revealed by RT-qPCR

(Fig.1C). The induction of *CYP24A1* (Fig. 1D), a major VDR target gene in many cell types

served as control for $1,25(OH)_2D_3$ activity.

VDR expression is the main determinant of cell responsiveness to $1,25(OH)_2D_3$ and is

downregulated in a proportion of patients with advanced CRC, implying that these patients

would probably not benefit from the anticancer effects of $1,25(OH)_2D_3$ (Ferrer-Mayorga et al.,

2017; Larriba and Muñoz, 2005; Pálmer et al., 2004; Peña et al., 2005). To model CRCs patients

with low VDR tumour level or with VD deficiency, we used an HCT 116-derived cell line stably

depleted of VDR using shRNA (hereafter named ShVDR) (Larriba et al., 2011). Induction of

SIRT1 gene expression by $1,25(OH)_2D_3$ was confirmed by RT-qPCR analysis in ShControl but

not in ShVDR cells, although basal SIRT1 levels were similar (Fig. 1E). At the protein level,

confocal imaging showed reduced SIRT1 protein in ShVDR as compared to ShControl cells

(Fig. 1F), suggesting that VDR contributes to maintaining basal SIRT1 protein expression.

Western blot analyses on nuclear extracts confirmed that ShControl cells responded to

$1,25(OH)_2D_3$ with increased SIRT1 protein levels in contrast to the unresponsiveness of ShVDR

cells (Fig. 1G). Moreover, the upregulating effect of $1,25(OH)_2D_3$ was specific for SIRT1, since

the expression of the other nuclear sirtuin, SIRT7, was unaffected (Fig. 1H). Together, these data

indicated that $1,25(OH)_2D_3$ specifically increased SIRT1 mRNA and protein levels in CRC cell

lines via VDR. This is important because SIRT1 depletion and/or sirtuin activity decay might

represent an important neglected mediator of VD deficiency and a target for intervention in CRC.

Whether VD controls sirtuin activity independently of sirtuin level in CRC is a critical issue that

deserved further exploration.

$1,25(OH)_2D_3$ induces SIRT1 deacetylase activity in CRC cells

Evaluation of global sirtuin activity in the nuclei of CRC cells revealed that treatment

with $1,25(OH)_2D_3$ increased NAD^+-dependent deacetylase activity 1.5-fold as compared to

control untreated cells (Fig. 2A). NAD^+ is a cofactor required for sirtuin activity. Metabolic

control of the level of NAD^+ represents an important node for sirtuin regulation and could explain

the induction of general sirtuin activity by $1,25(OH)_2D_3$. Indeed, nuclear NAD^+ level doubled in

response to $1,25(OH)_2D_3$ in human CRC cells (Fig. 2B). However, whether $1,25(OH)_2D_3$ could

specifically increase SIRT1 activity remained unknown.

Liganded VDR interacts with SIRT1 in kidney and bone cells (Sabir et al., 2017)

suggesting that $1,25(OH)_2D_3$ might alter SIRT1 expression or activity via post-transcriptional

mechanisms. Since protein-protein interactions regulate stability and activity of proteins, the

ability of $1,25(OH)_2D_3$ to promote VDR-SIRT1 interaction in CRC cells was evaluated by co-

immunoprecipitation assays. The results revealed that $1,25(OH)_2D_3$ potently (4-fold) enhanced

VDR/SIRT1 complex formation (Fig. 2C). To test if this interaction alters SIRT1 activity, the

acetylation status of two specific SIRT1 substrates (FOXO3a and H3K9) was evaluated as a read

out of SIRT1 deacetylase activity in response to $1,25(OH)_2D_3$. Acetyl-lysine

immunoprecipitation followed by Western blotting demonstrated efficient deacetylation of

FOXO3a in response to $1,25(OH)_2D_3$ (Fig. 2D). Likewise, the level of acetylated histone H3K9

(Ace H3K9) dramatically decreased in response to $1,25(OH)_2D_3$ (Fig. 2E), indicating specific

activation of SIRT1 by $1,25(OH)_2D_3$. The effect of $1,25(OH)_2D_3$ was mediated by VDR since

ShVDR cells did not show Ace H3K9 deacetylation in response to $1,25(OH)_2D_3$, whereas

ShControl cells maintained the response (Fig. 2F). Importantly, the specific SIRT1 activator

SRT1720 reduced Ace H3K9 levels in both ShControl and ShVDR cells, thus mimicking

$1,25(OH)_2D_3$ action independently of VDR (Fig. 2F).

The ability of $1,25(OH)_2D_3$ to increase protein deacetylation mediated by SIRT1 could

be driven either by elevated SIRT1 level or by increased SIRT1 activity. To distinguish between

these two possibilities, we performed an *in vitro* deacetylase assay using equivalent amounts of

immunoprecipitated SIRT1 from cells treated or not with $1,25(OH)_2D_3$. SIRT1-specific

deacetylase activity was enhanced 3-fold in cells treated with $1,25(OH)_2D_3$ despite similar levels

of immunoprecipitated SIRT1 (Fig. 2G).

Together, these results indicated that $1,25(OH)_2D_3$ induces nuclear SIRT1 activity. This

implies that VD deficiency and/or loss of tumour VDR expression (both observed in a proportion

of CRC patients) (Evans et al., 1998; Ferrer-Mayorga et al., 2019; Giardina et al., 2015; Larriba

et al., 2013, 2009; Pálmer et al., 2004; Peña et al., 2005) may negatively impact SIRT1

deacetylase activity and prompted us to examine samples of CRC patients.

CRC biopsies exhibit discrepancies between SIRT1 protein levels and deacetylase activity

Changes in SIRT1 RNA and protein expression and enzymatic activity were analysed in

samples of CRC patients. First, we compared available data (TCGA-COAD database) from 160

pairs of human colon adenocarcinoma samples matched with their normal adjacent tissue. TNM

plot analysis revealed that both *VDR* and *SIRT1* gene expression decreased from normal to

tumour colonic tissue in a highly significant fashion ($P<0.001$) (Fig. 3A-B).

Immunohistochemical analysis of human colon tumor tissue microarrays (TMAs) from

the Biobank of Hospital Clínico San Carlos (HCSC, Madrid) revealed a significant ($P=0.017$)

positive association between the levels of VDR and SIRT1 proteins (Fig. 3C). An additional

TMA containing healthy colon samples was used to compare tumor and nontumor tissues. The

level of VDR protein was usually lower in tumour samples (Fig. 3D), while SIRT1 protein

expression showed high variability between tumors and high cellular intratumor heterogeneity

(Fig. 3D-E). Globally, TMA data suggested that SIRT1 protein levels in tumor samples (Fig.

3E), are subjected to high variability, which hampers statistical significance even though for

SIRT1 gene expression, differences were statistically significant as shown in Figure 3B.

Global acetylation also showed high cellular intratumor heterogeneity and variability

between samples without statistically significant differences between tumor and nontumor

samples (data not shown), showing that is not a valid, specific read out of SIRT1 activity. As an

alternative approach, the levels of SIRT1 protein and its substrate Ace H3K9 were compared

first in intestinal cell lines and then in human tumours. Surprisingly, SIRT1 protein level

increased in CRC cells as compared to normal HIEC6 cells (Fig. 3F). However, increased SIRT1

levels coincided with elevated Ace H3K9 (Fig. 3F), indicating SIRT1 inactivation in CRC cells.

The results in cell lines encouraged us to evaluate the potential discordance of SIRT1 and Ace

H3K9 levels in fresh-frozen samples from paired tumor and nontumor colonic tissue of patients

from Hospital Universitario Fundación Alcorcón (HUFA). All tissues were evaluated to have at

least 85% of tumor cells by immunohistochemistry (IHC). Western blotting allowed

simultaneous detection of SIRT1 and Ace H3K9. Notably, whereas the level of SIRT1 protein

variably increased or decreased from nontumor to tumor tissues in different patients, acetylation

of its specific substrate H3K9 consistently increased in tumor samples (Fig. 3G), reflecting SIRT1 inactivation as found in carcinoma cell lines (Fig. 3F). *In vitro* deacetylation assays performed on SIRT1 immunoprecipitated from SIRT1-positive patient samples confirmed a common decrease of SIRT1 activity in colon tumor tissue (Fig. 3H). Western blotting illustrated the variability of SIRT1 level among tumors and its correlation with VDR level (Fig. 3I), while Ace H3K9 levels are generally high, but variable. Importantly, in vitro deacetylation assays on SIRT1 immunoprecipitated from the same samples (Fig. 3J) confirmed Ace H3K9 as a *bona fide* read out of SIRT1 inactivation. The discrepancy between the levels of SIRT1 protein expression and activity in tumor samples may solve the controversy previously reported regarding the use of SIRT1 as a tumor biomarker and poses the question of how $1,25(OH)_2D_3$ specifically enhances SIRT1 deacetylase activity.

$1,25(OH)_2D_3$ induction of SIRT1 activity is mediated through auto deacetylation

Since SIRT1 catalyses its auto deacetylation as a mechanism to increase its deacetylase activity towards other substrates (Fang et al., 2017) and $1,25(OH)_2D_3$ facilitates VDR-SIRT1 interaction, we explored the possibility that liganded VDR increases SIRT1 activity by facilitating SIRT1 auto deacetylation. To evaluate SIRT1 acetylation status upon exposure to $1,25(OH)_2D_3$, nuclear extracts from CRC cells challenged or not with $1,25(OH)_2D_3$ were immunoprecipitated using anti-acetyl-lysine antibodies and subjected to Western blotting. $1,25(OH)_2D_3$ greatly reduced the nuclear pool of acetylated SIRT1 (Fig. 4A). Remarkably, the specific SIRT1 activator SRT1720 also drove SIRT1 deacetylation in HCT 116 CRC cells (Fig. 4B). Consistently, ShVDR cells did not reduce acetylated SIRT1 in response to $1,25(OH)_2D_3$ but retain their response to SRT1720 (Fig 4C). These results indicated that $1,25(OH)_2D_3$ activation of SIRT1 is mediated through SIRT1 deacetylation.

Next, we sought to get further insight into the mechanism of regulation of SIRT1 activity

by 1,25(OH)$_2$D$_3$. As the acetyltransferase EP300 acetylates and inactivates SIRT2 (Han et al.,

2008) and is critical for CRC cell signalling (Chocarro-Calvo et al., 2013; Gutiérrez-Salmerón

et al., 2020), we used the ASEB engine (Wang et al., 2012) to search for SIRT1 residues

potentially acetylated by EP300. This approach identified K610, a lysine conserved from mouse

to human, as a potential EP300 target (Fig. 4D). The role of K610 in the regulation of SIRT1

activity was assessed by generating a non-acetylable K610R mutant, which should be resistant

to inactivation through acetylation. We compared the enzymatic activity of K610R SIRT1 with

that of wild type (WT) and inactive H363Y SIRT1 (Vaziri et al., 2001). Nuclear extracts from

cells transfected with MYC-tagged SIRT1 WT or mutants were immunoprecipitated using anti-

acetyl-lysine antibodies. Western blotting using an anti-MYC antibody revealed that exogenous

WT SIRT1 was acetylated (Fig. 4E), as it was endogenous SIRT1 (Fig 4A). Of note, catalytically

inactive H363Y SIRT1 was highly acetylated, in striking contrast with the lack of acetylation of

the K610R SIRT1 mutant (Fig. 4E), This highlights the importance of K610 for the regulation

of SIRT1 activity.

Since interaction with VDR was important for SIRT1 deacetylation activity, the ability

of each mutant to interact with VDR was compared in immunoprecipitation assays. The results

revealed that the K610R SIRT1 mutation facilitated the interaction with VDR, whereas the

inactive H363Y SIRT1 showed low interaction with VDR (Fig 4F). This suggested that

acetylation of SIRT1 might interfere with or destabilize its interaction with VDR, and that the

ability of 1,25(OH)$_2$D$_3$ to favour VDR-SIRT1 interaction may allow SIRT1 auto deacetylation,

and thus enhance its activity. The activity of mutant SIRT1 enzymes was first inferred from the

level of Ace H3K9, which was lower in cells expressing K610R SIRT1 (Fig 4G). *In vitro* NAD$^+$-

dependent deacetylase assays were next used to compare the activity of WT and mutant SIRT1.

Using equivalent amounts of immunoprecipitated MYC-tagged enzymes, we found a 10-fold

higher deacetylase activity for the K610R mutant than for WT SIRT1, whereas as expected the

H363Y inactive mutant exhibited very low activity as expected (Fig. 4H).

SIRT1 activation mimics the antiproliferative effects of 1,25(OH)₂D₃ in CRC cells

unresponsive to VD

Collectively, our data indicated that the acetylation status of SIRT1 K610 is critically controlled

and governs SIRT1 activity in the nuclei of CRC cells. By promoting VDR-SIRT1 interaction,

$1,25(OH)_2D_3$ facilitates SIRT1 auto deacetylation and enhances SIRT1 activity. These results

suggest that SIRT1 is an effector of $1,25(OH)_2D_3$ and prompted us to ask whether the

pharmacological activation of SIRT1 would rescue CRC cells that are unresponsive to

$1,25(OH)_2D_3$.

The anticancer effects of $1,25(OH)_2D_3$ are exerted at least partially through the inhibition

of tumor cell proliferation (Carlberg and Muñoz, 2022; Muñoz and Grant, 2022). $1,25(OH)_2D_3$

strongly reduced the percentage of CRC cells in S-G$_2$M cell-cycle phases (Fig. 5A) and

consequently, it lengthened the cell cycle and reduced CRC cell proliferation. We aimed to

analyse the effect of the modulation of SIRT1 activity on the proliferation of CRC cells and on

the antiproliferative action of 1,25(OH)2D3. Inhibition of general sirtuin activity with

Nicotinamide (NAA) abolished the antiproliferative effects of $1,25(OH)_2D_3$ on CRC cells (Fig.

5B). Moreover, activation of SIRT1 by the small molecule SRT1720 mimicked the response to

$1,25(OH)_2D_3$ and efficiently reduced the percentage of cells in S-G$_2$M cell cycle phases (Fig.

5C). These data indicate that induction of SIRT1 activity mediates $1,25(OH)_2D_3$-driven cell cycle

lengthening in CRC cells.

As expected, in contrast with ShControl cells, ShVDR CRC cells did not extend the cell

cycle in response to $1,25(OH)_2D_3$ (Fig. 5D). Accordingly, an early and strong reduction in the

proliferation rate in response to $1,25(OH)_2D_3$ was noted in ShControl cells but it was almost

absent in ShVDR cells (Fig. 5E). Specific SIRT1 activation with SRT1720 reduced the

proliferation of ShControl cells at a comparable magnitude to $1,25(OH)_2D_3$ and, importantly,

also reduced proliferation of $1,25(OH)_2D_3$-unresponsive ShVDR cells (Fig.5E).

Collectively, these results show that SIRT1 acts downstream of $1,25(OH)_2D_3$-activated

VDR in the inhibition of CRC cell proliferation. They also suggest that activation of SIRT1 in

CRC patients with vitamin D deficiency or with VD unresponsive tumors (conditions frequently

observed in advanced CRC), could be a therapeutic strategy to effectively antagonize cell

proliferation.

DISCUSSION

Here we reveal that $1,25(OH)_2D_3$ activates the epigenetic regulator SIRT1 by acting at

multiple levels. Ligand-activated VDR post-translationally upregulates SIRT1 activity by

facilitating its auto deacetylation. Mechanistically, SIRT1 K610 is identified as essential to

control VDR-SIRT1 interaction and SIRT1 acetylation status and activity. Notably, both

SRT1720 and $1,25(OH)_2D_3$ activate SIRT1 through auto deacetylation. Modulation of SIRT1

activity by $1,25(OH)_2D_3$ through deacetylation occurs in addition to increased mRNA and

protein levels as well as to an elevation of cofactor (NAD^+) availability. Thus, $1,25(OH)_2D_3$

activates SIRT1 acting at multiple levels, from gene expression to activity, and these findings

broaden our understanding of how VD deficiency is associated with CRC.

We also show that SIRT1 deacetylase activity mediates the anti-proliferative action of

$1,25(OH)_2D_3$ in CRC, which is in line with the recent proposal that SIRT1 activity opposes

several cancer cell hallmarks (Yousafzai et al., 2021). In cases of VD deficiency and/or

unresponsiveness such as advanced colorectal tumor lacking VDR expression due to

transcriptional repression by SNAIL1/2 (Pálmer et al., 2004; Larriba et al., 2009), the activation

of SIRT1 by compounds other than $1,25(OH)_2D_3$ may be especially relevant. Clinical trials that

explore SIRT1 activators in CRC are rare (https://clinicaltrials.gov). Some trials are designed to

study doses and pharmacokinetics of resveratrol, a known sirtuin activator that has many other

targets (Patel et al., 2011). Currently there are not known clinical trials using SRT1720 or

alternative SIRT1 specific activators.

Consideration of SIRT1 protein level as a prognostic marker for cancer has received

much attention, but conflicting results reporting both, tumor SIRT1 increases and decreases

raised controversy (Ren et al., 2017; Wu et al., 2017). Accordingly, some studies propose SIRT1

either as tumour suppressor or as tumor promoter in CRC (Carafa et al., 2019; Ren et al., 2017).

Importantly, SIRT1 protein level might not always reflect SIRT1 activity, and some tumors could

possibly increase SIRT1 protein expression to counteract inactivation. Our results, which

indicate the existence of discrepancies between SIRT1 levels and activity in human colorectal

tumors, may help to resolve the controversy. Interestingly, alternative posttranslational SIRT1

modifications may have a similar outcome. For example, proteolytic cleavage of SIRT1 induced

by TNFα in osteoarthritis renders accumulation of a catalytically inactive 75 kDa resistant

fragment (Oppenheimer et al., 2012) that could be detected by immunohistochemistry. Since

inflammation (in general) and especially TNFα is an important mediator of CRC, understanding

if this fragment is generated in CRC patients and whether its presence would also lead to

decreased SIRT1 activity in samples strongly positive for SIRT1 remains as an interesting

unexplored issue.

The ability of $1,25(OH)_2D_3$ to activate SIRT1 may have further major implications for

physiological and pathological processes other than CRC since SIRT1 target proteins control

several processes such as mitochondria physiology, apoptosis, and inflammation (Strycharz et

al., 2018). Thus, $1,25(OH)_2D_3$ may not only regulate gene expression via direct transcriptional

regulation of VDR-bound targets, but it can secondarily impact a wide range of events and

processes via the control of SIRT1 activity. This may explain how $1,25(OH)_2D_3$ regulates the

activity of FOXO3a, an important modulator of cell proliferation and the immune response that is targeted by SIRT1 (An et al., 2010). SIRT1 also modulates the activity of the tumour suppressor p53, whose mutation is a relatively early event in a high proportion of CRCs (Polidoro et al., 2013; Strycharz et al., 2018). In this way, SIRT1 activation greatly broaden the known vitamin D regulatory action, albeit whether $1,25(OH)_2D_3$ directs SIRT1 activity towards specific substrates remains to be elucidated. The accumulating evidence for SIRT1 involvement in chronic inflammation (Mendes et al., 2017) suggests a potential benefit of SIRT1 activation by $1,25(OH)_2D_3$ in chronic inflammation-linked CRC and other diseases. Moreover, the mechanism revealed here for SIRT1 activation by $1,25(OH)_2D_3$ may be relevant for diseases associated with VD deficiency beyond CRC, including autoimmune disorders and diabetes.

In conclusion, the multilevel activation of SIRT1 deacetylase by $1,25(OH)_2D_3$ broadens the range of known $1,25(OH)_2D_3$ targets and positions SIRT1 as an important mediator of the protective action of VD against CRC and potentially other neoplasias and non-tumoral diseases. These data support SIRT1 activators as useful agents in situations of dysfunction of the VD system.

MATERIALS AND METHODS

All materials and resources can be found in Table 1.

Experimental model and subject details

Human samples

Patient samples were derived from surgical removal at Fundación Jimenez Diaz University Hospital, General and Digestive Tract Surgery Department: 95 patient samples diagnosed with stage II CRC. The Institutional Review Board (IRB) of the Fundación Jimenez Diaz Hospital, reviewed and approved the study, granting approval on December 9, 2014, act number 17/14.

Samples from seven healthy individuals were obtained from Hospital Clínico San Carlos

(HCSC). IRB-HCSC act number 21/498-E granting approval on June 25th, 2021. Paired fresh-

frozen samples from tumor and nontumor adjacent tissue were collected from 18 patients at

Hospital Universitario Fundación Alcorcón (HUFA). IRB-HUFA, act number 17/68 granting

approval on June 8th, 2017. Colon cancer tissues were collected using protocols approved by the

corresponding Ethics Committees and following the legislation. All patients gave written

informed consent for the use of their biological samples for research purposes. Fundamental

ethical principles and rights promoted by Spain (LOPD 15/1999) and the European Union EU

(2000/C364/01) were followed. All patients' data were processed according to the Declaration

of Helsinki (last revision 2013) and Spanish National Biomedical Research Law (14/2007, July

3).

Colorectal cell panels

Human colorectal adenocarcinoma HT-29 and HCT 116 and HCT 116 derived ShControl and

ShVDR cells were cultured under standard conditions, in 5%CO2 at 37ºC with DMEM

containing 25 mM glucose (unless specifically indicated) and 40 mM LiCl to mimic Wnt

signalling and supplemented with 10% foetal bovine serum (FBS) and 1% penicillin-

streptomycin. Cells were treated as indicated for 24h. HCT 116 ShControl and ShVDR cell lines

were derived previously (Larriba et al., 2011)

Transient transfections

cells were seeded in plates at 50% confluence for plasmid transfection, using JetPei PolyPlus

reagent (Genycell Biotech), following the manufacturer's instructions. After 24 h cells were

treated as indicated for 24 h.

For Sirt1 siRNA, cells plated in six well plates at 50% confluence were transfected with

JetPRIME PolyPlus reagent (Genycell Biotech) following the manufacturer's instructions. After

2 days cells were treated 24 h with LiCl and then another 24 h with $1,25(OH)_2D_3$ before

harvesting to analyse by western blot.

Site directed mutagenesis

Directed mutagenesis was performed using the QuickChange Site-Directed Mutagenesis Kit

(Stratagene). Oligonucleotides containing the mutation (Table 1) were designed following the

manufacturer's recommendations. 2.5 U of PfuTurbo DNA polymerase was added to each

sample and they were amplified by performing the following cycles: 1 denaturation cycle at 95 °

C for 1 minute, 18 amplification cycles of 30 seconds at 95 ° C, 50 seconds at 60 ° C, 1 minute

/ kilobase of plasmid at 68 °C and the last extension cycle 7 minutes at 68ªC. Once amplified, 10

U of Dnp I enzyme were added and incubated at 37 ° C for one hour to digest the supercoiled

double-stranded DNA and eliminate the parental molecules that are not mutated.

The correct introduction of the mutation was evaluated by plasmid sequencing using the BigDye

Cycle Sequencing Kit (Applied Biosystems).

Preparation of cell extracts:

Whole cell extracts.

Cells were washed with iced PBS before extract preparation and scraped in RIPA buffer (10 mM

Tris HCl [pH 7.4], 5 mM EDTA, 5 mM EGTA, 1% Tryton X100, 10 mM $Na_4P_2O_7$ [pH 7.4], 10

mM NaF, 130 mM NaCl, 0.1% SDS, 0,5% Na-deoxycholate). After 5 min on ice, cells were

pelleted (12000 rpm for 5 min, 4ºC) and the supernatant was directly used as whole cell extract

or frozen at -80 ºC.

Fractionated cell extracts.

After washing as before, cells were scraped in hypotonic buffer (20 mM Hepes, [pH 8.0], 10 mM

KCl, 0,15 mM EDTA, 0,15 mM EGTA, 0,05% NP40 and protease inhibitors) and incubated on

ice for 10 min before adding 1:2 vol of sucrose buffer (50 mM Hepes [pH=8.0], 0.25 mM EDTA,

10 mM KCl, 70% sucrose). Lysates were fractionated (5000 rpm for 5 min at 4ºC) to obtain the

cytoplasmic fraction in the supernatant. Nuclear pellets were washed twice with washing buffer

(20 mM Hepes [pH 8.0], 50 mM NaCl, $MgCl_2$ 1.5 mM, 0,25 mM EDTA, 0,15 mM EGTA, 25%

glycerol and protease inhibitors), pelleted at 5000 rpm, 5 min at 4ºC and resuspended in nuclear

extraction buffer (20 mM Hepes[pH 8.0], 450 mM NaCl, $MgCl_2$ 1.5 mM, 0,25 mM EDTA, 0,15

mM EGTA, 0,05% NP40, 25% glycerol and protease inhibitors) before centrifugation at 12,000

rpm for 5 min at 4ºC to pellet and discard cell debris. The supernatants were used as nuclear

fractions.

SIRT1 Activity Assay

NAD^+-dependent deacetylase activity was measured on nuclear extracts of HT-29 cells treated

as indicated, using SIRT-Glo Assay kit (Promega), following the manufacturer's instructions.

For each reaction, 1 μg of protein was incubated with SIRT-Glo Reagent Mix at room

temperature for 3 min, and the luminescence was measured using a GloMax Microplate Reader

(Promega). Relative luciferase units (RLU) were calculated as fold induction relative to the

corresponding control.

Nuclear NAD^+ abundance

The NAD/NADH-Glo Assay Kit (Promega) was used to measure NAD^+ abundance in HCT 116

nuclear extracts. 20 μg of protein were diluted in 50 μl of lysis buffer (described for cell extracts)

for each reaction and 25 μl of 0.4 N HCl were added before incubation at 60 °C for 15 minutes.

This exploits the high stability of NAD^+ under acidic conditions, while NADH breaks down.

Samples were cooled down to room temperature for 10 min and neutralized with 25 µL 0.5 M

Trizma®. NAD^+ levels were then measured using a GloMax microplate reader (Promega) and

following the manufacturer's instructions.

Immunoprecipitation

For immunoprecipitation from fractionated extracts, the hypotonic buffer was modified by

adding 100mM NaCl and 0.1% NP40. For immunocomplex formation, protein A/G-coated

magnetic beads (Invitrogen) were washed 3 times with the extraction buffer before coating with

the primary antibody for 2 hr at 4 ºC in a rotating wheel, followed by 2 washes with the same

buffer to eliminates unbound antibody and then extracts were added O/N at 4ºC in the rotating

wheel. Immunocomplexes were washed twice and used for western blotting.

Western blotting

Proteins from lysed cells or immunoprecipitates were denatured and loaded on sodium dodecyl

sulfate polyacrylamide gels and then transferred to polyvinylidene difluoride membranes (Bio-

Rad). After blocking with 5% (w/v) BSA or milk the membrane was incubated with the

corresponding primary and secondary antibodies (Bio-Rad). The specific bands were analyzed

using Thyphoon or ChemiDoc Imaging Systems (Bio-Rad).

Immunoflorescence

Cells in cover slips were washed three times and fixed with 4% paraformaldehyde in PBS [pH

7.4] for 10min, washed again, permeabilized (PBS [pH7.4], 0.5% Triton X-100, 0.2% BSA) for

5min, blocked (PBS [pH7,4, 0,05% Triton X-100, 5% BSA) for 1h at room temperature and

incubated with primary antibody over night at 4ºC. After three washes (5 min) cells were

incubated with secondary antibody for 1h at room temperature. Slides were mounted and images

were acquired using a SP5 confocal microscope (Leica) with a 63x objective. Fluorescence

intensity was quantified using Image J software. For each experiment, 3 different fields were

evaluated per slide.

Cell-growth curves and cell cycle

Cell growth was determined by colorimetry. For that, cells were seeded at a density of 20,000

cells per well in a Corning 12-well plate and treated as indicated for 1 to 6 days. After that cells

were treated with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT)

(Sigma-.Aldrich) in a 1:10 ratio in a culture medium and incubated at 37 °C for 3 h. In living

cells, MTT is reduced to formazan, which has a purple color. The medium was removed, and the

formazan was resuspended in dimethylsulfoxide (DMSO), transferred to p96 plates and analyzed

by a Spectra FLUOR (Tecan) at 542 nm. Viability was measured in four independent

experiments and then duplicated.

For cell cycle analysis, cells were harvested by trypsinization, washed with PBS, fixed in 70%

ethanol, stained with 7-AAD (Santa Cruz Biotechnology) for 10 min at 37°C, and analyzed by

flow cytometry (FACSCalibur, Becton-Dickinson). The percentages of cells at each cell cycle

phase were analyzed using CXP software (Becton-Dickinson).

RT-qPCR

Total RNA was extracted from 3 replicates of colorectal cells using TRIzol reagent (Invitrogen).

Reverse transcription of 1µg of RNA was performed according to the manufacturer's instructions

Reagents and detection systems were from Applied biosystems. 18S ribosomal RNA primers

served as a nonregulated control. Relative expression was calculated using the Ct method,

expressed as $2^{-\Delta\Delta Ct}$ (Livak and Schmittgen, 2001). The PCR efficiency was approximately

100%.

TMA, immunohistochemistry, and quantification.

Tissue microarrays (TMA) containing 92 cores of stage 2 human CRC samples, were constructed

using the MTA-1 tissue arrayer (Beecher Instruments, Sun Prairie) for immunohistochemistry

analysis. Each core (diameter 0.6 mm) was punched from pre-selected tumour regions in

paraffin-embedded tissues. We chose central areas from the tumour, avoiding foci of necrosis.

Staining was conducted in 2-µm sections. Slides were deparaffinized by incubation at 60°C for

10 min and incubated with PT-Link (Dako, Agilent) for 20 min at 95°C in low pH to detect VDR,

SIRT1 and acetyl lysines. To block endogenous peroxidase, holders were incubated with

peroxidase blocking reagent (Dako, Agilent) and then with (1:50) dilutions of antibodies: anti-

VDR, anti-SIRT1 and (1:300) anti-acetyl lysine, overnight at 4°C. All previously described

antibodies presented high specificity. After that, slides were incubated for 20 min with the

appropriate anti-Ig horseradish peroxidase-conjugated polymer (EnVision, Dako, Agilent).

Sections were then visualized with 3,3'-diaminobenzidine (Dako, Agilent) as a chromogen for 5

min and counterstained with Harrys' Haematoxylin (Sigma Aldrich, Merck). Photographs were

taken with a stereo microscope (Leica DMi1). According to the human protein atlas (available at http://www.proteinatlas.org), a human colon tissue was used as a positive control for immunohistochemical staining to determine anti-VDR concentration and a human testis tissue for anti-SIRT1. Immunoreactivity was quantified blind with a Histoscore (H score) that considers both the intensity and percentage of cells stained for each intensity (low, medium, or high) following this algorithm (range 0–300): H score = (low%) × 1 + (medium%) × 2 + (high %) × 3. Quantification for each patient biopsy was calculated blindly by 2 investigators (MJFA and JMU). VDR and SIRT1 showed nuclear staining, whereas acetyl lysines were nuclear and extranuclear. Clinicopathological characteristics of patients are summarized in Table 2.

Bioinformatics analyses:

Gene expression (RNA-seq) data from the TCGA colon adenocarcinoma cohort (COAD) was downloaded from the cBioportal for Cancer Genomics (http://www.cbioportal.org) using CGDS-R (Cerami et al., 2012; Gao et al., 2013) following TCGA guidelines (http://cancergenome.nih.gov/publications/publicationguidelines). SIRT1 and VDR gene expression from samples of colorectal cancer and healthy tissue adjacent to the tumour were analysed with TNMplot.com (https://tnmplot.com/analysis/) tool (Bartha and Győrffy, 2021). An asymptotic Spearman correlation test using original log2 expression values was used to determine the significance of the Spearman rank correlation.

Statistical analysis

Results are presented as fold induction, mean ± SEM from three biological replicates. To determine whether calculated Hscores for protein levels were well-modelled by a normal distribution, Kolmogorov-Smirnov test was used. Association between SIRT1 and VDR protein levels was performed with Chi-square test. Comparisons between two independent groups (healthy versus tumors) were performed with Mann-Whitney U-test. For statistical association, Hscore of antigens were categorized as high- or low- expression levels using the median as a

cut-off point. Tests for significance between two sample groups were performed with Student's

t test and for multiple comparisons, ANOVA with Bonferroni's post-test.

ACKNOWLEDGMENTS: Funding for this work was provided by the Agencia Estatal de

Investigación (PID2019-104867RB-I00 / AEI/10.13039/501100011033, RTI2018-099343-B-

100 and PID2021-127645OA-I00); Instituto de Salud Carlos III (CIBERONC, CB16/12/00273

and CB16/12/00326); Comunidad de Madrid (Ayudas Atracción de Talento 2017-T1/BMD-

5334 and 2021-5A/BMD-20951; A385-DROPLET Young Reserchers R&D Project 2019 CAM-

URJC; PRECICOLON-CM, P2022/BMD7212); Universidad Rey Juan Carlos (ADIPOMELM,

Proyecto Puente de Investigación 2020).

We thank Lucille Banham for her valuable assistance in the preparation of the English

manuscript and María Gutiérrez Salmerón for help making figures.

Additional Information

FUNDER	GRANT REFERENCE NUMBER	AUTHOR
Agencia Estatal de Investigación	PID2019-110998RB-I00 AEI/10.13039/501100011033	Custodia Garcia-Jimenez
	PID2019-104867RB-I00	Alberto Muñoz & María Jesús Larriba
	RTI2018-099343-B-100	Antonio De la Vieja
	PID2021-127645OA-I00	Ana Chocarro-Calvo
Instituto de Salud Carlos III	CIBERONC CB16/12/00273	Alberto Muñoz
	CIBERONC CB16/12/00326	Antonio De la Vieja
Comunidad de Madrid	PRECICOLON-CM P2022/BMD7212	Custodia García-Jiménez Alberto Muñoz
	Ayudas Atracción de Talento 2017- T1/BMD-5334	Ana Chocarro-Calvo
	Ayudas Atracción de Talento 2021-5A/BMD-20951	
	A385-DROPLET Young Researchers R&D Project 2019	
Universidad Rey Juan Carlos	ADIPOMELM, Proyecto Puente de Investigación 2020	

Author contributions: CGJ, JMGM, AM, and MJL conceived the experiments, JMGM, ACC,

AB, JMU, MJFA, MCF, JCR performed the experiments. JMGM and ADLV performed the Figs.

All authors participated in the discussion. CGJ, AM and MJL wrote the manuscript and provided

supervision.

Ethics

The authors declare that they have no conflict of interest. All procedures followed were in

accordance with the ethical standards of the responsible committee on human experimentation

(institutional and national) as specified under the paragraph Human samples in the Materials and

Methods Section and with the Helsinki Declaration of 1975, as revised in 2000. Informed consent

was obtained from all patients for being included in the study.

REFERENCES

An B-S, Tavera-Mendoza LE, Dimitrov V, Wang X, Calderon MR, Wang H-J, White JH.
2010. Stimulation of Sirt1-Regulated FoxO Protein Function by the Ligand-Bound
Vitamin D Receptor. *Mol Cell Biol* **30**:4890–4900. doi:10.1128/MCB.00180-10

Bartha Á, Győrffy B. 2021. Tnmplot.Com: A web tool for the comparison of gene expression
in normal, tumor and metastatic tissues. *Int J Mol Sci* **22**:1–12.
doi:10.3390/ijms22052622

Bikle D, Christakos S. 2020. New aspects of vitamin D metabolism and action — addressing
the skin as source and target. *Nat Rev Endocrinol* **16**:234–252. doi:10.1038/s41574-
019-0312-5

Bonkowski MS, Sinclair DA. 2016. Slowing ageing by design: The rise of NAD+ and
sirtuin-activating compounds. *Nat Rev Mol Cell Biol* **17**:679–690.
doi:10.1038/NRM.2016.93

Cancer Genome Atlas Network. 2012. Comprehensive molecular characterization of human
colon and rectal cancer. *Nature* **487**:330–337. doi:10.1038/nature11252

Carafa V, Altucci L, Nebbioso A. 2019. Dual Tumor Suppressor and Tumor Promoter
Action of Sirtuins in Determining Malignant Phenotype. *Front Pharmacol* **10**:38.
doi:10.3389/fphar.2019.00038

Carlberg C, Muñoz A. 2022. An update on vitamin D signaling and cancer. *Semin Cancer Biol* **79**:217–230. doi:10.1016/j.semcancer.2020.05.018

Carlberg C, Velleuer E. 2022. Vitamin D and the risk for cancer: A molecular analysis. *Biochem Pharmacol* **196**. doi:10.1016/J.BCP.2021.114735

Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, Antipin Y, Reva B, Goldberg AP, Sander C, Schultz N. 2012. The cBio Cancer Genomics Portal: An open platform for exploring multidimensional cancer genomics data. *Cancer Discov* **2**:401–404. doi:10.1158/2159-8290.CD-12-0095

Chandler PD, Chen WY, Ajala ON, Hazra A, Cook N, Bubes V, Lee I-M, Giovannucci EL, Willett W, Buring JE, Manson JE. 2020. Effect of Vitamin D 3 Supplements on Development of Advanced Cancer. *JAMA Netw Open* **3**:e2025850. doi:10.1001/jamanetworkopen.2020.25850

Chen X, Sun K, Jiao S, Cai N, Zhao X, Zou H, Xie Y, Wang Z, Zhong M, Wei L. 2014. High levels of SIRT1 expression enhance tumorigenesis and associate with a poor prognosis of colorectal carcinoma patients. *Sci Rep* **4**. doi:10.1038/SREP07481

Chocarro-Calvo A, García-Martínez JM, Ardila-González S, de la Vieja A, García-Jiménez C. 2013. Glucose-Induced β-Catenin Acetylation Enhances Wnt Signaling in Cancer. *Mol Cell* **49**:474–486. doi:10.1016/j.molcel.2012.11.022

Evans SR, Nolla J, Hanfelt J, Shabahang M, Nauta RJ, Shchepotin IB. 1998. Vitamin D receptor expression as a predictive marker of biological behavior in human colorectal cancer. *Clinical Cancer Research* **4**:1591–1595.

Bray F., Ferlay J., Soerjomataram I, Siegel RL, Torre LA, Jemal A. 2018. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* **68**:394–424. doi:10.3322/caac.21492

Fang J, Ianni A, Smolka C, Vakhrusheva O, Nolte H, Krüger M, Wietelmann A, Simonet NG, Adrian-Segarra JM, Vaquero A, Braun T, Bober E. 2017. Sirt7 promotes adipogenesis in the mouse by inhibiting autocatalytic activation of Sirt1. *Proc Natl Acad Sci U S A* **114**:E8352–E8361. doi:10.1073/pnas.1706945114

Feldman D, Krishnan A V., Swami S, Giovannucci E, Feldman BJ. 2014. The role of vitamin D in reducing cancer risk and progression. *Nat Rev Cancer*. doi:10.1038/nrc3691

Fernández-Barral A, Bustamante-Madrid P, Ferrer-Mayorga G, Barbáchano A, Larriba MJ, Muñoz A. 2020. Vitamin D Effects on Cell Differentiation and Stemness in Cancer. *Cancers (Basel)* **12**:2413–2432. doi:10.3390/cancers12092413

Ferrer-Mayorga G, Gómez-López G, Barbáchano A, Fernández-Barral A, Peña C, Pisano DG, Cantero R, Rojo F, Muñoz A, Larriba MJ. 2017. Vitamin D receptor expression and associated gene signature in tumour stromal fibroblasts predict clinical outcome in colorectal cancer. *Gut* **66**:1449–1462. doi:10.1136/gutjnl-2015-310977

Ferrer-Mayorga G, Larriba MJ, Crespo P, Muñoz A. 2019. Mechanisms of action of vitamin D in colon cancer. *J Steroid Biochem Mol Biol* **185**:1–6. doi:10.1016/j.jsbmb.2018.07.002

Firestein R, Blander G, Michan S, Oberdoerffer P, Ogino S, Campbell J, Bhimavarapu A, Luikenhuis S, de Cabo R, Fuchs C, Hahn WC, Guarente LP, Sinclair DA. 2008. The SIRT1 Deacetylase Suppresses Intestinal Tumorigenesis and Colon Cancer Growth. *PLoS One* **3**:e2020. doi:10.1371/journal.pone.0002020

Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, Cerami E, Sander C, Schultz N. 2013. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* **6**:1–34. doi:10.1126/scisignal.2004088

Giardina C, Nakanishi M, Khan A, Kuratnik A, Xu W, Brenner B, Rosenberg DW. 2015. Regulation of VDR expression in Apc-mutant mice, human colon cancers and adenomas. *Cancer Prevention Research* **8**:387–399. doi:10.1158/1940-6207.CAPR-14-0371

Grant WB, Boucher BJ, al Anouti F, Pilz S. 2022. Comparing the Evidence from Observational Studies and Randomized Controlled Trials for Nonskeletal Health Effects of Vitamin D. *Nutrients* **14**:1–31.

Gutiérrez-Salmerón M, García-Martínez JM, Martínez-Useros J, Fernández-Aceñero MJ, Viollet B, Olivier S, Chauhan J, Lucena SR, de la Vieja A, Goding CR, Chocarro-Calvo A, García-Jiménez C. 2020. Paradoxical activation of AMPK by glucose drives selective EP300 activity in colorectal cancer. *PLoS Biol* **18**:e3000732. doi:10.1371/journal.pbio.3000732

Han Y, Jin YH, Kim YJ, Kang BY, Choi HJ, Kim DW, Yeo CY, Lee KY. 2008. Acetylation of Sirt2 by p300 attenuates its deacetylase activity. *Biochem Biophys Res Commun* **375**:576–580. doi:10.1016/j.bbrc.2008.08.042

Henn M, Martin-Gorgojo V, Martin-Moreno JM. 2022. Vitamin D in Cancer Prevention: Gaps in Current Knowledge and Room for Hope. *Nutrients* **14**:1–32.

Kim H, Lipsyc-Sharf M, Zong X, Wang X, Hur J, Song M, Wang M, Smith-Warner SA, Fuchs C, Ogino S, Wu K, Chan AT, Cao Y, Ng K, Giovannucci EL. 2021. Total Vitamin D Intake and Risks of Early-Onset Colorectal Cancer and Precursors. *Gastroenterology* **161**:1208-1217.e9. doi:10.1053/j.gastro.2021.07.002

Kim H, Yuan C, Nguyen LH, Ng K, Giovannucci EL. 2022. Prediagnostic Vitamin D Status and Colorectal Cancer Survival by Vitamin D Binding Protein Isoforms in US Cohorts. *J Clin Endocrinol Metab*. doi:10.1210/CLINEM/DGAC742

Larriba MJ, González-Sancho JM, Barbáchano A, Niell N, Ferrer-Mayorga G, Muñoz A. 2013. Vitamin D is a multilevel repressor of Wnt/β-catenin signaling in cancer cells. *Cancers (Basel)*. doi:10.3390/cancers5041242

Larriba MJ, Martín-Villar E, García JM, Pereira F, Peña C, García de Herreros A, Bonilla F, Muñoz A. 2009. Snail2 cooperates with Snail1 in the repression of vitamin D receptor in colon cancer. *Carcinogenesis* **30**. doi:10.1093/carcin/bgp140

Larriba MJ, Muñoz A. 2005. SNAIL vs vitamin D receptor expression in colon cancer: therapeutics implications. *Br J Cancer* **92**:985–989. doi:10.1038/sj.bjc.6602484

Larriba MJ, Ordóñez-Morán P, Chicote I, Martín-Fernández G, Puig I, Muñoz A, Pálmer
HG. 2011. Vitamin D receptor deficiency enhances Wnt/β-catenin signaling and tumor
burden in colon cancer. *PLoS One* **6**:e23524. doi:10.1371/journal.pone.0023524

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**:402–408.
doi:10.1006/meth.2001.1262

Manson JE, Bassuk SS, Buring JE, VITAL Research Group. 2020. Principal results of the
VITamin D and OmegA-3 TriaL (VITAL) and updated meta-analyses of relevant
vitamin D trials. *J Steroid Biochem Mol Biol* **198**:105522.
doi:10.1016/j.jsbmb.2019.105522

Mendes KL, Lelis D de F, Santos SHS. 2017. Nuclear sirtuins and inflammatory signaling
pathways. *Cytokine Growth Factor Rev* **38**:98–105. doi:10.1016/j.cytogfr.2017.11.001

Muñoz A, Grant WB. 2022. Vitamin D and Cancer: An Historical Overview of the
Epidemiology and Mechanisms. *Overview of the Epidemiology and Mechanisms*
Nutrients **14**:1448. doi:10.3390/nu14071448

Oppenheimer H, Gabay O, Meir H, Haze A, Kandel L, Liebergall M, Gagarina V, Lee EJ,
Dvir-Ginzberg M. 2012. 75-kd sirtuin 1 blocks tumor necrosis factor α-mediated
apoptosis in human osteoarthritic chondrocytes. *Arthritis Rheum* **64**:718–728.
doi:10.1002/ART.33407

Pálmer HG, Larriba MJ, García JM, Ordóñez-Morán P, Peña C, Peiró S, Puig I, Rodríguez
R, de La Fuente R, Bernad A, Pollán M, Bonilla F, Gamallo C, García De Herreros A,
Muñoz A. 2004. The transcription factor SNAIL represses vitamin D receptor
expression and responsiveness in human colon cancer. *Nat Med* **10**:917–919.
doi:10.1038/nm1095

Patel KR, Scott E, Brown VA, Gescher AJ, Steward WP, Brown K. 2011. Clinical trials of
resveratrol. *Ann N Y Acad Sci* **1215**:161–169. doi:10.1111/j.1749-6632.2010.05853.x

Peña C, Garciá JM, Silva J, García V, Rodríguez R, Alonso I, Millán I, Salas C, García de
Herreros A, Muñoz A, Bonilla F. 2005. E-cadherin and vitamin D receptor regulation
by SNAIL and ZEB1 in colon cancer: Clinicopathological correlations. *Hum Mol Genet*
14:3361–3370. doi:10.1093/hmg/ddi366

Polidoro L, Properzi G, Marampon F, Gravina GL, Festuccia C, di Cesare E, Scarsella L,
Ciccarelli C, Zani BM, Ferri C. 2013. Vitamin D protects human endothelial cells from
H2O2 oxidant injury through the Mek/Erk-sirt1 axis activation. *J Cardiovasc Transl*
Res **6**:221–231. doi:10.1007/s12265-012-9436-x

Ren NSX, Ji M, Tokar EJ, Busch EL, Xu X, Lewis DA, Li Xiangchun, Jin A, Zhang Y, Wu
WKK, Huang W, Li L, Fargo DC, Keku TO, Sandler RS, Li Xiaoling. 2017.
Haploinsufficiency of SIRT1 Enhances Glutamine Metabolism and Promotes Cancer
Development. *Current Biology* **27**:483–494. doi:10.1016/j.cub.2016.12.047

Sabir MS, Khan Z, Hu C, Galligan MA, Dussik CM, Mallick S, Stone AD, Batie SF, Jacobs
ET, Whitfield GK, Haussler MR, Heck MC, Jurutka PW. 2017. SIRT1 enzymatically

potentiates 1,25-dihydroxyvitamin D3 signaling via vitamin D receptor deacetylation.
J Steroid Biochem Mol Biol **172**:117–129. doi:10.1016/J.JSBMB.2017.06.010

Song M, Lee IM, Manson JAE, Buring JE, Dushkes R, Gordon D, Walter J, Wu K, Chan
AT, Ogino S, Fuchs CS, Meyerhardt JA, Giovannucci EL. 2021. No Association
Between Vitamin D Supplementation and Risk of Colorectal Adenomas or Serrated
Polyps in a Randomized Trial. *Clinical Gastroenterology and Hepatology* **19**:128-
135.e6. doi:10.1016/j.cgh.2020.02.013

Strycharz J, Rygielska Z, Swiderska E, Drzewoski J, Szemraj J, Szmigiero L, Sliwinska A.
2018. SIRT1 as a Therapeutic Target in Diabetic Complications. *Curr Med Chem*
25:1002–1035. doi:10.2174/0929867324666171107103114

Vaziri H, Dessain SK, Eaton EN, Imai SI, Frye RA, Pandita TK, Guarente L, Weinberg RA.
2001. hSIR2SIRT1 functions as an NAD-dependent p53 deacetylase. *Cell* **107**:149–
159. doi:10.1016/S0092-8674(01)00527-X

Wang L, Du Y, Lu M, Li T. 2012. ASEB: a web server for KAT-specific acetylation site
prediction. *Nucleic Acids Res* **40**:W376–W379. doi:10.1093/nar/gks437

Wu S, Jiang J, Liu J, Wang X, Gan Y, Tang Y. 2017. Meta-analysis of SIRT1 expression as
a prognostic marker for overall survival in gastrointestinal cancer. *Oncotarget* **8**:62589–
62599. doi:10.18632/oncotarget.19880

Yousafzai NA, Jin H, Ullah M, Wang X. 2021. Recent advances of SIRT1 and implications
in chemotherapeutics resistance in cancer. *Am J Cancer Res* **Nov 15**:5233-5248.

Yuan Q, Zhang R, Sun M, Guo X, Yang J, Bian W, Xie C, Miao D, Mao L. 2022. Sirt1
Mediates Vitamin D Deficiency-Driven Gluconeogenesis in the Liver via mTorc2/Akt
Signaling. doi:10.1155/2022/1755563

Zheng W, Wong KE, Zhang Z, Dougherty U, Mustafi R, Kong J, Deb DK, Zheng H,
Bissonnette M, Li YC. 2012. Inactivation of the vitamin D receptor in APC(min/+)
mice reveals a critical role for the vitamin D receptor in intestinal tumor growth. *Int J*
Cancer **130**:10–19. doi:10.1002/IJC.25992

FIGURE LEGENDS

Fig. 1. *1,25(OH)$_2$D$_3$ increases SIRT1 levels in CRC cells and VDR is required to ensure basal*

levels.

(A)-(D) HCT 116 or HT-29 CRC cells cultured under standard conditions and where indicated,

treated with 1α,25-dihydroxyvitamin D$_3$ (1,25(OH)$_2$D$_3$), 100 nM, added 24 hours before

harvesting. (E)-(H) ShVDR cells were derived from HCT 116 by stable and specific knock-down

of vitamin D receptor (VDR) using specific shRNA and are compared to ShControl cells that

contain normal VDR levels (Larriba et al., 2011). Cell extracts were fractionated in all cases.

(A) Confocal imaging of SIRT1 (green) in CRC (HCT 116 and HT-29) cells and fluorescence

intensity quantification of 3 independent experiments using ImageJ software. For each

experiment 3 different fields were evaluated per slide. Scale bars:25μm.

(B) Western-blot analysis of $1,25(OH)_2D_3$ effects on levels of SIRT1 in nuclear extracts (NE)

from indicated CRC cells. Representative blots and statistical analysis using TBP as loading

controls.

(C)-(E) RT-qPCR analysis of the effect of $1,25 (OH)_2D_3$ in indicated colon cancer cells. Values

normalized with endogenous control (18S RNA) are referred as fold induction over cells without

$1,25(OH)_2D_3$.

(C) Effect of $1,25 (OH)_2D_3$ on SIRT1 gene expression.

(D) validation of $1,25 (OH)_2D_3$ activity on the canonical target gene cytochrome P450 family 24

subfamily A member 1, CYP24A1, in HT-29 colon cancer cells.

(E) Requirement of vitamin D receptor (VDR) for $1,25 (OH)_2D_3$ induction of SIRT1 gene

expression.

(F) Effect of VDR knock-down on SIRT1 protein content evaluated by confocal imaging of

ShControl and ShVDR HCT 116 colon cancer cells to detect SIRT1 (green). Scale bars represent

25 μm. On the right, quantification of fluorescence intensity using ImageJ software; for each

experiment, 3 different fields were evaluated per slide.

(G) Western blot analysis of the effect of $1,25 (OH)_2D_3$ on nuclear accumulation of SIRT1 in

cells depleted (ShVDR) or not (ShControl) of VDR.

(H) Specificity of $1,25 (OH)_2D_3$ effects on SIRT1 by western blot analysis of the alternative

nuclear sirtuin, SIRT7.

Statistical analysis of 3 independent experiments in each panel was performed by Student t-test

(A) to (D) and (H) or by One-Way ANOVA (E)-(F). For (G), statistical analysis of the four

groups by ANOVA is represented by * and comparison of the two indicated groups by Students

t test, by #. In all panels, values represent mean ± SEM of triplicates corresponding to biological

replicates; * or # $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Fig. 2. *1,25(OH)$_2$D$_3$ induces specific SIRT1 deacetylase activity independently of levels, in*

CRC cells.

Cells were cultured as in Fig 1.

(A) and (G) NAD$^+$-dependent deacetylase activity, (B) Evaluation of nuclear NAD$^+$ levels. (C)

to (F) Western blot analysis using TBP or H3 as loading controls. Representative western blots

and statistical analysis of 3 independent experiments.

(A) NAD$^+$-dependent deacetylase activity measured on nuclear extracts of HT-29 cells. Relative

luciferase units (RLU) were calculated as fold induction relative to the corresponding control.

(B) Evaluation of nuclear NAD$^+$ levels in HT-29 CRC cells in response to 1,25(OH)$_2$D$_3$, with

data expressed as fold induction over the control, untreated cells.

(C) 1,25(OH)$_2$D$_3$ effects on the interaction between vitamin D receptor (VDR) and SIRT1. Input

(5%) and flow through (FT) are shown

(D) 1,25(OH)$_2$D$_3$ effects on the acetylation status of the SIRT1 substrate FoxO3a in nuclear

extracts immunoprecipitated with anti-acetyl lysine antibodies. Input (5%) and flow through

(FT) are shown.

(E) 1,25(OH)$_2$D$_3$ effects on the nuclear acetylation of the SIRT1 substrate H3K9 (Ace H3K9).

(F) Requirement of VDR for deacetylation of Ace H3K9 in response to 1,25(OH)$_2$D$_3$

(G) Quantification of NAD^+-dependent deacetylase activity on SIRT1 immunoprecipitates from

nuclear extracts (NE) of HCT 116 CRC cells treated or not with $1,25(OH)_2D_3$. Values represent

mean ± SEM of n=3 independent experiments.

Statistical analysis by Student t test, n ≥ 3 biological replicates and values represent mean ±

SEM; *p<0.05; **p<0.01; *** p<0.001.

Fig. 3. *CRC biopsies exhibit discrepancies between SIRT1 protein levels and deacetylase*

activity

(A)-(B) TNM plot analysis of VDR (A) or SIRT1 (B) gene expression in 160 paired samples of

non-cancer adjacent tissue (normal) versus adenocarcinoma tissue (tumour). Data was collected

from TCGA-COAD database. SIRT1 or VDR expression is shown as transcripts per million

(TPM) in log2. Tumour and control samples were compared with Mann-Whitney U test. The

statistical significance cut off was set at p < 0.01.

(C) Association analysis between VDR and SIRT1 protein levels in human colon cancer samples,

stage 2. Cut-off points to separate proteins between high- or low- expression levels were for

SIRT1 expression median of Histoscore, and negative *versus* positive staining for VDR

expression. Statistical analysis was performed with Chi-square test.

(D) Representative micrographs at 40X from immunostaining for vitamin D receptor (VDR) or SIRT1
on human samples of healthy colon or CRC patients. Scale bars: 20μm. Immunostainings were
revealed with DAB (diaminebenzidine) and thus, positiveness is highlighted by light or dark brown
according to their low or high protein expression. Counterstaining with hematoxilin stains nuclei
in dark blue and cytoplasm in light blue.

(E) Profile for SIRT1 content from healthy to colon cancer human samples. The profile was

obtained from Histoscore variations. Tumour and control samples were compared with Mann-

Whitney U test. The statistical significance cut off was set at p < 0.01.

(F) Western-blot analysis of the levels of SIRT1 and its substrate Ace H3K9 in total lysates from intestinal healthy (HIEC6) or CRC cells. Representative western blot with ERK as loading control.

(G) Western-blot analysis from human patient samples, with patient number indicated at the top. CRC samples (T) and adjacent nontumor tissue (NT) were probed for SIRT1 and its substrate Ace H3K9 (as in F). Frozen samples were obtained from HUFA patients.

(H) NAD^+-dependent deacetylase activity on immunoprecipitated SIRT1 from patient lysates. Values for T and NT samples were corrected according to immunoprecipitated SIRT1 levels shown on the western underneath the graph. Values for T samples were referred to values of NT and data (duplicates) are expressed as fold induction. Patient number is indicated at the bottom.

(I) Western-blot analysis for SIRT1 and its substrate Ace H3K9 levels in total lysates from alternative human CRC samples (as in G). Patient number is indicated at the top.

(J) NAD^+-dependent deacetylase activity on immunoprecipitated SIRT1 from patient lysates presented in (I). Levels of immunoprecipitated SIRT1 are presented in the bottom panel. SIRT1 deacetylase activity in patients is presented as fold induction, referred to the first patient sample (patient #10). Patient number is indicated at the bottom.

Fig. 4. *1,25(OH)₂D₃ induces SIRT1 activity through auto deacetylation*

HCT 116 CRC cells were cultured under standard conditions in DMEM containing LiCl (40 mM) to mimic Wnt signalling before addition of 1,25(OH)$_2$D$_3$, 100 nM or SRT1720 (20 μM) for the last 24 h; after fractionation nuclear extracts (NE) were used. (A)-(C) and (E)-(F) Immunoprecipitation (IP) using anti-acetyl lysine antibody from nuclear extracts (NE) of indicated cells and western-blot analysis to evaluate changes in acetylation of SIRT1; TBP was

used in the Flow Through (FT) as loading control. Representative western-blots and statistical

analysis are shown in all panels.

(A) Effect of $1,25(OH)_2D_3$ on acetylation of SIRT1.

(B) Effect of SRT1720 on acetylation of SIRT1. (C) Requirement of VDR for $1,25(OH)2D_3$ or

SRT1720 effects on acetylation of SIRT1 in Sh Control and ShVDR HCT 116 cells.

(D) SIRT1 scheme with functional domains and putative acetylation sites (top) and sequence

alignment. Conserved putative acetylation targets for activation/inactivation of SIRT1 shown at

the bottom.

(E) Acetylation status of exogenous SIRT1 wild type (WT) or mutants (H363Y or K610R). HCT

116 were transiently transfected with pcDNA expression vectors: empty (-), myc-tagged SIRT1

wild type (WT) or mutants: H363Y inactive or K610R, 48 h before harvesting.

Immunoprecipitation using anti-acetyl-lysine antibodies and western blot detection using myc

antibodies. Representative western-blot and statistical analysis; MYC and TBP in the flow

through (FT) serve as loading controls.

(F) Western-blot analysis of the interaction of SIRT1 wild type and mutants, with vitamin D

receptor (VDR). Expression and analysis as in (E).

(G) Western-blot analysis of the effect of SIRT1 mutant expression on Ace H3K9.

(H) *In vitro* NAD^+-dependent deacetylase activity assays on equivalent amounts of exogenous

SIRT1 wild type (WT), K610R and inactive H363Y mutants immunoprecipitated using anti-myc

antibodies.

Statistical analysis in all panels except (D) and (G) by One Way ANOVA of n>3 independent

experiments. Values represent mean ± SEM; *p<0.05; **p<0.01; *** p<0.001.

Fig. 5. *SIRT1 activation rescues antiproliferative effects of* $1,25(OH)_2D_3$ *in unresponsive CRC*

cells.

Cells were cultured as in previous figures and treatments with NAA (300 uM), SRT1720 (20μM)

and 1,25 (OH)$_2$D$_3$ (100 nM) also were for the last 24 h.

(A) Flow cytometry analysis of cell cycle from indicated cells treated (+) or not (-) with

1,25(OH)$_2$D$_3$. Numbers correspond to the percentage of cells in the indicated phases expressed

as mean ± SEM.

(B) Effect of NAA on the 1,25(OH)$_2$D$_3$-driven blockades of HCT 116 CRC cell proliferation.

(C) Effects of SRT1720 on cell cycle of HCT 116 CRC cells by flow cytometry analysis.

Numbers correspond to the percentage of cells in the indicated phases expressed as mean ± SEM.

(D) Flow cytometry analysis of cell cycle response to 1,25 (OH)$_2$D$_3$ in ShControl and ShVDR

HCT 116 cells. Numbers correspond to the percentage of cells in the indicated phases expressed

as mean ± SEM.

(E) Proliferation curves of ShControl or ShVDR HCT 116 colon cancer cells in response to 1,25

(OH)$_2$D$_3$ (100 nM) (grey), compared to control with vehicle (green) or SRT1720 (red).

Statistical analysis by One-Way ANOVA of 3 independent experiments; values represent

mean ± SEM of triplicates; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Table 1. Key Materials & Resources.

Antibodies		
Rabbit Polyclonal anti-Acetylated-lysine	Cell Signaling	Cat# 9441
Rabbit monoclonal anti-Histone H3K9-Ace (C5B11)		Cat # 9649
Rabbit Polyclonal anti-Histone H3		Cat # 9715
Rabbit Polyclonal anti-Vitamin D3 receptor (D2K6W)		Cat #12550
Rabbit monoclonal anti-FoxO3a (75D8)		Cat# 2497
Rabbit Polyclonal anti-SIRT1 (D1D7)		Cat# 9475
Rabbit Polyclonal anti-SIRT1	Santa Cruz Biotechnology	Cat#SC-15404
Rabbit Polyclonal anti-TFIID-TBP (N12)		Cat# SC-204
Rabbit Polyclonal anti-SIRT7		Cat#SC365344
Rabbit Polyclonal anti-ERK (C14)		Cat# SC-154
Mouse Monoclonal anti-MYC (9E10)		Cat# SC-40
Donkey Anti-Rabbit-AlexaFluor488	Invitrogen	Cat# A21206
Mouse monoclonal anti-TBP (51841)		Cat# MA514739
Goat Anti-Rabbit IgG (H+L) HRPO	BIO-RAD	Cat #170-6515
Goat Anti-Mouse IgG (H+L) HRPO		Cat #170-6516
Mouse Monoclonal anti-Tubulin Clone B-5-1-2	Sigma-Aldrich	Cat# T5168
EnVision + Dual Link System-HRP (DAB+)	Dako (Agilent)	Cat#K4065
Chemicals, Peptides, and Recombinant Proteins		
1α,25-DihydroxyVD$_3$	Sigma-Aldrich	17936
Lithium Chloride	Sigma-Aldrich	Cat# L9650
BSA	Sigma-Aldrich	A7906
Nicotinamide (NAA)	Sigma-Aldrich	N3376
SRT1720	Selleckchem	S1129
TRIzol reagent	Invitrogen	Cat#15596026
Protease inhibitor Cocktail	Roche	Cat#04693132001
7-AAD	Santa Cruz Biotechnology	SC-221210

3-(4,5-dimetiltiazol-2-il)-2,5-difeniltetrazol (MTT)	Sigma-Aldrich	Cat# M5655	
DMEM media	Lonza	Cat# 12-604F	
Bovine Fetal Serum	Sigma	Cat# F7524	
JetPEI PolyPlus reagent	Genycell Biotech	Cat# 101-10N	
JetPRIME PolyPlus reagent	Genycell Biotech	Cat # 114-01	
QuickChange Site-Directed Mutagenesis Kit	Stratagene	Cat #200523	
SIRT-Glo Assay kit	Promega	G6450	
NAD/NADH-Glo Assay Kit	Promega	G9071	
DYNAbeads Protein A	Invitrogen	Ref 10002D	
DYNAbeads Protein G	Invitrogen	Ref 10004D	
Bacterial Strains			
E. coli 5α	New England Biolabs	NEB5α	
Experimental Models: Cell Lines			
HCT 116 (Male) colorectal carcinoma	ATCC	Cat# CCL-247, RRID:CVCL_0291	
HT-29 (Female). Rectosigmoid adenocarcinoma	ATCC	Cat# HTB–38, RRID:CVCL_0320	
HCT 116 ShControl	In house	(Larriba et al. 2011)	
HCT 116 ShVDR	In house	(Larriba et al. 2011)	
Oligonucleotides			
Human SIRT1 K610R mutagenesis primers F 5′-3′: GGTTCTAGTACTGGGGAGAGGAATGAAAGAACTTCAGTGG R 5′-3′: CCAGCCACTGAAGTTCTTTCATTCCTCTCCCCAGTACTAG	Sigma	This study	
Human CYCLIN D1 (CCND1) qPCR primers F 5′-3′: AAGATCGTCGCCACCTGG R 5′-3′: GGAAGACCTCCTCCTCGCAC	Sigma	This study	
Human c-MYC qPCR primers F 5′-3′: CTTCTCTCCGTCCTCGGATTCT R 5′-3′: GAAGGTGATCCAGACTCTGACCTT	Sigma	This study	
Human 18s qPCR primers F 5′-3′: AGTCCCTGCCCTTTGTACACA R 5′-3′ GCCTCACTAAACCATCCAATCG	Sigma	This study	
CYP24A1 TaqMan® probe	Applied Biosystems	Hs00167999_m1	
Recombinant DNA			
pcDNA3.1-SIRT1-Myc-His	Gift from Prof. Colin R Goding	N/A	
pcDNA3.1-SIRT1H363Y- Myc-His	Gift from Prof. Colin R Goding	N/A	
pcDNA3.1-SIRT1 K601R- Myc-His	This study	N/A	
Software and Algorithms			
LAS AF software	Leica	SP5	
3730xl Analyzer	Applied Biosystem	ABI 3730XL	
GraphPad Prism software	GraphPad Software	https://www.graphpad.com	

ImageLab	Bio-Rad	ChemiDoc XRS+ System
CXP software	Becton-Dickinson	FACSCalibur
ImageJ software	https://imagej.nih.gov/ij/download.html	N/A
SPSS Statistics version 26	IBM	N/A

Table 2. Clinicopathological characteristics of CRC patients included in the study.

Patients Characteristic	N (%)	
	Stage II	Stage IV
Gender		
Male	57 (60%)	27 (48%)
Female	38 (40%)	23 (41%)
N/A	0	6 (11%)
Tumor Location		
Cecum	13 (14%)	3 (5%)
Right	25 (26%)	10 (18%)
Transverse	7 (8%)	3 (5%)
Left	5 (5%)	2 (4%)
Sigma	24 (25%)	21 (38%)
Rectum	21 (21%)	12 (21%)
N/A	0	5 (9%)
Grade Primary Tumor		
Well differentiated	18 (19%)	15 (27%)
Moderately differentiated	69 (73%)	27 (48%)
Poorly differentiated	8 (8%)	5 (9%)
N/A	0	9 (16%)
Paired Liver Metastasis		54 (96%)
N/A		2 (4%)
Grade Metastasis Tumor		
Well differentiated		12 (21%)
Moderately differentiated		29 (52%)
Poorly differentiated		4 (7%)
N/A		11 (20%)
Metastasis at diagnosis		
Synchronous		35 (62%)
Metachronous		21 (38%)
Pt		
T1	3 (3%)	2 (4%)
T2	30 (32%)	5 (9%)
T3	61 (64%)	42 (75%)
T4	1 (1%)	4 (7%)
N/A		3 (5%)
pN		
N0	95 (100%)	27 (48%)
N1		16 (29%)
N2		9 (16%)
N3		1 (2%)
N/A		3 (5%)