Neural responses to repeated noise structure in sounds
are invariant to temporal interruptions

Björn Herrmann

Rotman Research Institute,

Baycrest, M6A 2E1, North York, ON, Canada

Department of Psychology,

University of Toronto, M5S 1A1, Toronto, ON, Canada

*Correspondence concerning this article should be addressed to Björn Herrmann, Rotman Research

Institute, Baycrest, 3560 Bathurst St, North York, ON, M6A 2E1, Canada. E-mail:

bherrmann@research.baycrest.org

Keywords: Electroencephalography, frozen noise, temporal regularity, auditory perception, neural

synchronization

26   Abstract

The ability to extract meaning from acoustic environments requires sensitivity to repeating sound

structure. Yet, how events that repeat are encoded and maintained in the brain and how the brain

responds to events that reoccur at later points in time is not well understood. In two

electroencephalography experiments, participants listened to a longer, ongoing white-noise sound

which comprised shorter, frozen noise snippets that repeated at a regular 2-Hz rate. In several

conditions, the snippet repetition discontinued for a brief period after which the noise snippet

reoccurred. The experiments aimed to answer whether neural activity becomes entrained by the regular

repetition of noise snippets, whether entrained neural activity self-sustains during the discontinuation

period, and how the brain responds to a reoccurring noise snippet. Results show that neural activity is

entrained by the snippet repetition, but there was no evidence for self-sustained neural activity during

the discontinuation period. However, auditory cortex responded with similar magnitude to a noise

snippet reoccurring after a brief discontinuation as it responded to a noise snippet for which the snippet

repetition had not been discontinued. This response invariance was observed for different onset times

of the reoccurring noise snippet relative to the previously established regularity. The results thus

demonstrate that the auditory cortex sensitively responds to, and thus maintains a memory trace of,

previously learned acoustic noise independent of temporal interruptions.

44   Introduction

The sounds humans encounter in everyday life, such as speech and music, are rich of structured

amplitude and frequency components that repeat over time, and humans are sensitive to such

repetitions (McDermott et al., 2013; Teki et al., 2013; Sohoglu and Chait, 2016; Herrmann et al., 2020).

Sensitivity to the repetition of sound structure is thought to support generating predictions about future

sounds (Jones and Boltz, 1989; Barnes and Jones, 2000; Bendixen, 2014; Henry and Herrmann, 2014;

Nobre and van Ede, 2018), enabling acoustic-change detection (Schröger, 2007; Winkler et al., 2009),

and facilitating the perception of speech (Idemaru and Holt, 2011; Giraud and Poeppel, 2012; Peelle and

Davis, 2012; Baese-Berk et al., 2014). Yet, how events that repeat are encoded and maintained in the

brain and how the brain responds to events when they reoccur is not fully understood.

A variety of different approaches have been used to investigate the perception and neural

correlates of repeating structure in sounds (Guttman and Julesz, 1963; Warren et al., 2001; Schröger,

2005; Agus et al., 2010; Agus and Pressnitzer, 2013; Bendixen, 2014; Andrillon et al., 2015; Barascud et

al., 2016; Teki et al., 2016; Viswanathan et al., 2016; Andrillon et al., 2017; Kang et al., 2017; Bianco et

al., 2020; Herrmann et al., 2021; Hodapp and Grimm, 2021; Dauer et al., 2022; Ringer et al., 2022). One

fruitful line of research has focused on acoustic noise as stimuli to investigate how individuals perceive

and learn repeating structure in sounds (Warren et al., 2001; Kaernbach, 2004; Agus et al., 2010; Agus

and Pressnitzer, 2013; Luo et al., 2013; Andrillon et al., 2015; Rajendran et al., 2016; Andrillon et al.,

2017; Dauer et al., 2022). The use of acoustic noise to examine the sensitivity to repeated structure has

the advantage that a listener will not have previously heard the specific instantiation of a noise sound

and will not be able to use semantic category labels because each noise consists of randomly generated

numbers. Although the specific paradigms and stimuli have varied in previous studies, in a subset of

studies, a repeated, short snippet of noise (~0.2 s) was embedded in a longer, ongoing noise stimulus

(Andrillon et al., 2015; Andrillon et al., 2017; Dauer et al., 2022). Both the short noise snippet and the

longer noise are generated based on the same random-number distribution (e.g., Gaussian white noise).

The short noise snippet is sometimes referred to as "frozen noise" because the exact array of random

values that makes up the snippet are "frozen" in place and repeated within the ongoing noise stimulus

(Warren et al., 2001; Andrillon et al., 2015; Rajendran et al., 2016; Dauer et al., 2022). The ongoing noise,

in contrast, consists of nonrepeating random values drawn from the same distribution (Figure 1). The

repetition of the short noise snippet in the ongoing noise forms a regular auditory pattern that a listener

can perceive.

Individuals have remarkable memory for such frozen, repeated noise snippets that form a regular

auditory pattern in an ongoing noise. Listeners are more likely to detect the presence of a pattern when

the same, compared to a new, noise snippet is used on different trials (Andrillon et al., 2015; Andrillon

et al., 2017; Dauer et al., 2022). Individuals further appear to have memory for specific auditory patterns

even days and weeks after they have been exposed to them (Agus et al., 2010; Viswanathan et al., 2016;

Bianco et al., 2020; Bianco et al., 2023). However, how the brain maintains memory representations for

a noise snippet after it has been repeated is unclear. The current work focuses on short, within-second

timescales to investigate the neural responses to a noise snippet reoccurring after a brief discontinuation

of a snippet repetition.

Low-frequency neural activity is thought to reflect fluctuations in neuronal excitability (Bishop,

1933; Adrian and Matthews, 1934; Lakatos et al., 2005) that entrains to periodic repetitions of sounds

or sound features (Picton et al., 2003; Stefanics et al., 2010; Herrmann et al., 2013; Herrmann et al.,

2023) such that high excitability periods align with occurrences of the sounds or sound features to

facilitate their processing (Thut et al., 2011; Henry and Herrmann, 2014; Lakatos et al., 2019; Obleser

and Kayser, 2019). The term entrainment is used broadly here because responses to periodic repetitions

of sounds could either reflect neural oscillations or evoked responses, and distinguishing between the

two response types is empirically challenging (Klimesch et al., 2007; Sauseng et al., 2007; Capilla et al.,

2011; Fellinger et al., 2011; Notbohm et al., 2016; Keitel et al., 2021).

Entrainment of neural activity has been observed for noise snippets that are repeated in an

ongoing noise (Andrillon et al., 2015; Andrillon et al., 2017; Ringer et al., 2023b, a), providing a potential

mechanism that enables the encoding of the noise snippet. Neural activity that is entrained by

temporally regular stimulation – if oscillatory in nature – is hypothesized to self-sustain at the repetition

rate when the stimulation discontinues (Pikovsky et al., 2001; Lakatos et al., 2013b; Henry and

Herrmann, 2014; Keitel et al., 2022). Self-sustainability of entrained neural activity may facilitate the

neural representation of a snippet even when its repetition temporarily discontinues and may support

neural responses to a subsequent reoccurrence of the noise snippet. However, whether neural activity

self-sustains after a snippet repetition discontinues and possibly facilitates responses to a reoccurring

noise snippet is unclear. Moreover, if the entrained response reflects an evoked response, no self-

sustainability of neural activity would be expected. Whether or not neural activity self-sustains may thus

speak to the underlying response mechanism.

Perception of sounds and neural entrainment can depend on the timing at which events occur

(Large and Jones, 1999; Barnes and Jones, 2000; Jones et al., 2002; Jones et al., 2006; Lange, 2010;

Kayser et al., 2015; Herrmann et al., 2016; Morillon et al., 2016; Herrmann et al., 2023). For example,

listeners detect auditory patterns made of repeating noise snippets in an ongoing noise better when the

noise snippet is repeated at regular compared to irregular intervals (Rajendran et al., 2016; Dauer et al.,

2022). Making perceptual inferences about a sound is also better when the sound occurs in-sync with a

temporally regular stimulation – that is, at a time predicted by the periodicity of a regular stimulus

repetition – compared to out-of-sync (Barnes and Jones, 2000; Jones et al., 2002; McAuley and Jones,

2003; Jones et al., 2006; Herrmann et al., 2016). The perceptual benefit of presenting a sound in-sync

with a previous stimulus regularity is thought to arise from entrained, self-sustained neural activity

(Large and Jones, 1999; Lakatos et al., 2013b; Henry and Herrmann, 2014), such that high-excitability

periods occur at in-sync times and low-excitability periods at out-of-sync times. Perceptual sensitivity

and neural responses to a reoccurring noise snippet may thus be facilitated if the snippet reoccurs in-

sync compared to out-of-sync with a previously established temporal regularity, but this has not been

investigated.

In two electroencephalography (EEG) experiments, the current study investigates the neural

responses to frozen noise snippets that repeat at a regular rate. The experiments aim to answer whether

neural activity self-sustains during a discontinuation of the repetition and how the brain responds to a

reoccurring noise snippet. Specifically, Experiment 1 is concerned with the neural activity during the

discontinuation period and the neural response to a reoccurring noise snippet. Experiment 2 is

concerned with the degree of synchrony of a reoccurring noise snippet relative to a previously

established stimulus regularity. The two EEG experiments will help understand how the brain maintains

neural representations of recently learned structure in sounds.

General methods

Participants

Thirty-seven younger adults (18–34 years) participated in two EEG experiments of the current study.

Each person participated only in one of the two experiments. Participants gave written informed consent

prior to the experiment and were paid $7.5 CAD per half-hour for their participation. Participants

reported having normal hearing abilities. The study was conducted in accordance with the Declaration

of Helsinki, the Canadian Tri-Council Policy Statement on Ethical Conduct for Research Involving Humans

(TCPS2-2014), and was approved by the Research Ethics Board of the Rotman Research Institute at

Baycrest.

Sound environment and stimulus presentation

Data collection was carried out in a sound-attenuating booth. Sounds were presented via Sennheiser

(HD 25-SP II) headphones and a RME Fireface 400 external sound card. Stimulation was run using

Psychtoolbox in MATLAB (v3.0.14; MathWorks Inc.) on a Lenovo T480 laptop with Microsoft Windows

7. Visual stimulation was projected into the sound booth via screen mirroring. All sounds were presented

at a comfortable listening level that was fixed across participants (~70-75 dB SPL).

Acoustic stimulation and procedure

Stimulus generation was based on procedures described in previous work (Agus et al., 2010; Agus and

Pressnitzer, 2013; Andrillon et al., 2015; Andrillon et al., 2017; Dauer et al., 2022). Each sound was a

Gaussian white noise with a duration of either 4.8 s (Experiment 1) or 5 s (Experiment 2) sampled at 44.1

kHz. Participants listened either to white-noise sounds without additional characteristics – that is, they

did not contain a pattern – or to white-noise sounds that had embedded a short, repeating white-noise

snippet that created a regular pattern. In detail, a short noise snippet (mean duration across

experiments: 0.253 s; range of durations: 0.079–0.411 s), generated from the same Gaussian white-

noise distribution, was embedded in the 4.8-s or 5-s longer noise at six or more instances at a constant

onset-to-onset interval of 0.5 s (2 Hz). A unique, newly created noise snippet was generated for each

trial. The onset of the first embedded noise snippet was 0.5 s after the onset of the longer white noise.

The specific number of repetitions depended on the experiment and condition, and is described for each

experiment individually below. There were no silences or other acoustic cues separating embedded

snippet repetitions from the ongoing noise, because snippets and ongoing noise were generated from

the same Gaussian white-noise (i.e., random-number) distribution (Agus et al., 2010; Agus and

Pressnitzer, 2013; Andrillon et al., 2015; Andrillon et al., 2017; Dauer et al., 2022). That no acoustic cues

are present in the noise stimuli is also depicted in the time course, spectrogram, and cochleogram

(McDermott and Simoncelli, 2011) of a sample noise stimulus that included a snippet repetition shown

in Figure 1. Because of the absence of acoustic cues separating embedded snippet repetitions from the

ongoing noise, perception of a pattern in these sounds required the detection of the noise-snippet

repetition within the ongoing noise stimulus (Figure 1B).

Figure 1: Sample noise stimulus. A: Time course of a sample noise stimulus with seven embedded noise snippets. The seven short black lines above the x-axis indicate the onset and the 0.2-s duration of each noise snippet. **B:** 30-ms zoomed-in time courses time-locked to the onset of the first (onset at 0.5 s; top panel) and second (onset at 1 s; middle panel) embedded noise snippet. The bottom panel shows the same time courses of the snippet again, overlaid and time-locked to snippet onset. The overlaid time courses demonstrate that snippets are identical (purple line and green line overlap), but that the noise stimulus differs prior to snippet onset. **C:** Spectrogram of the sample stimulus in panel A. **D:** Cochleogram of sample stimulus in panel A (cochleogram was calculated using procedures described in previous work: McDermott and Simoncelli, 2011). The time course (A), spectrogram (C), and cochleogram (D) show that no silences or other acoustic cues separate embedded snippet repetitions from the ongoing noise.

During the experimental procedures, participants listened to the 4.8-s or 5-s white-noise sounds,

one at the time, and were asked to judge whether a pattern – that is a noise-snippet repetition – was

present or not. Participants were required to withhold their response until 0.1 s after sound offset where

a happy and a sad emoticon smiley were presented visually side by side (Herrmann et al., 2011a;

Herrmann et al., 2011b; Herrmann et al., 2021). If participants heard a pattern, they had to press the

button for the happy smiley. If they did not hear a pattern, they had to press the button for the sad

smiley. The position of the happy and sad smileys (left vs. right) was random across trials and participants

did therefore not know which button to press during listening, that is, prior to the visual presentation

(Herrmann et al., 2011a; Herrmann et al., 2011b; Herrmann et al., 2021). The next trial started 1.9 s after

they had made a response.

Detection of a noise-snippet repetition depends on the duration of the noise snippet (Rajendran

et al., 2016). In order to ensure that pattern detection was difficult but manageable, each participant

underwent a threshold-estimation procedure at the beginning of the experimental session during which

the duration of the noise snippet was estimated that would yield a detection rate for snippet repetitions

of 0.75 (with a few exceptions detailed for each experiment below). To this end, participants listened to

stimuli with embedded noise snippets varying at eight, equally-spaced durations: 0, 0.057, 0.114, 0.171,

0.229, 0.286, 0.343, 0.4 s (a duration of 0 means that no snippet repetition was embedded). Each snippet

duration was presented 8 times in random order. This block lasted about 8 min. For each duration level,

the proportion of 'pattern present' responses was calculated. A logistic function was fit to the proportion

data and the duration corresponding to a proportion of 0.75 (with a few exceptions, see below) was

calculated and used for the participant during the main experimental procedures.

Analysis of behavioral data during main experimental procedures

A response was considered a hit if a participant responded 'pattern present' to a sound that contained

a repetition of a noise snippet. A response was considered a false alarm if a person responded 'pattern

present' to a sound that did not contain a repetition of a noise snippet. The hit rate was calculated for

each of the experimental conditions (see below for each experiment), whereas one false alarm rate was

calculated per participant (i.e., non-condition-specific). Because pattern-absent trials were presented

among the conditions of pattern-present trials, no condition-specific false alarm rate could be calculated

(cf. Agus et al., 2010; Agus and Pressnitzer, 2013; Andrillon et al., 2015; Andrillon et al., 2017; Kang et

al., 2017; Hodapp and Grimm, 2021; Kang et al., 2021; Dauer et al., 2022; Ringer et al., 2022). False alarm

and hit rates were used to calculate perceptual sensitivity (d-prime) for each participant and condition

(Macmillan and Creelman, 2004) using the formula d-prime = z(hit rate) - z(false alarm rate), where z is

the inverse of the normal cumulative distribution function (norminv in Matlab).

Electroencephalography recordings and preprocessing

Electroencephalographical signals were recorded from 16 scalp electrodes (Ag/Ag–Cl-electrodes; 10-20

placement) and the left and right mastoids using a BioSemi system (Amsterdam, The Netherlands). The

sampling frequency was 1024 Hz with an online low-pass filter of 208 Hz. Electrodes were referenced

online to a monopolar reference feedback loop connecting a driven passive sensor and a common-

mode-sense (CMS) active sensor, both located posteriorly on the scalp.

Offline analysis was conducted using MATLAB software. An elliptic filter was used to suppress

power at the 60-Hz line frequency. Data were re-referenced by averaging the signal from the left and

right mastoids and subtracting the average separately from each of the 16 channels. Data were filtered

with a 0.7-Hz high-pass filter (length: 2449 samples, Hann window). The high-pass filter was specifically

designed to provide strong DC suppression that avoids requiring baseline correction and removes slow

drifts (Maess et al., 2006; Herrmann et al., 2011a; Herrmann et al., 2011b). Data were further low-pass

filtered at 22 Hz (length: 211 samples, Kaiser window), divided into epochs ranging from 1 s prior to

sound onset to 1 s post sound offset (-1 to 5.8 s for Experiment 1, and -1 to 6 s for Experiment 2; time-

locked to sound onset), and downsampled to 512 Hz. Independent components analysis (runica method,

Makeig et al., 1996; logistic infomax algorithm, Bell and Sejnowski, 1995; Fieldtrip implementation

Oostenveld et al., 2011) was used to identify and remove components related to blinks and horizontal

eye movements. Subsequently to the independent components analysis, epochs that exceeded a signal

change of more than 200 µV for any electrode were excluded from analyses.

Analysis of neural synchronization

In order to investigate whether the periodic repetition of a noise snippet elicited a synchronized

response, a fast Fourier transform (FFT) was calculated for each trial that contained a noise-snippet

repetition using the data in the 1-s to 3.5-s time window (Hann window tapering; zero-padding). The 1-

s to 3.5-s time window was selected because it covered the first 5 snippet repetitions that were present

on all trials (cycles 2 to 6; trials without noise repetition were not analyzed due to their low number;

N=36, see below). Inter-trial phase coherence (ITPC) was calculated by dividing the complex coefficients

that resulted from the FFT by its magnitude and averaging the result across trials (Lachaux et al., 1999).

ITPC values in the frequency window ranging from 1.9 to 2.1 Hz were averaged to obtain an ITPC

estimate at the 2-Hz stimulation frequency (snippet onset-to-onset interval of 0.5 s). ITPC values were

averaged across a fronto-central electrode cluster (F3, Fz, F4, C3, Cz, C4) that is known to be sensitive to

neural activity originating from auditory cortex, including neural synchronization (Näätänen and Picton,

1987; Picton et al., 2003; Irsik et al., 2021). To statistically assess whether neural activity was

synchronized with the 2-Hz regular rate in the stimulation, a paired samples t-test was used to compare

ITPC at 2-Hz to the averaged ITPC across neighboring frequencies (1.8–1.9 Hz and 2.1–2.2 Hz; for a

similar procedure see Nozaradan et al., 2011; Nozaradan et al., 2012).

232   Experiment 1

Experiment 1 investigates whether neural activity entrained by a noise-snippet repetition self-sustains

throughout a brief interruption and the degree to which a noise snippet that reoccurs after the

interruption elicits a neural response. Experiment 1 will help understanding how neural representations

of frozen noise snippets are maintained.

Methods

Participants

Seventeen participants took part in Experiment 1 (median age: 22 years; range: 18–34 years; 4 male, 13

female).

Stimuli and procedure

Participants listened to 4.8-s white-noise sounds in four different conditions. Each condition contained

a repetition of a noise snippet, but the number and onset times varied. Conditions are labeled according

to the number of snippets embedded in the ongoing noise. Sounds for condition S6 contained six

embeddings of a noise snippet, at 0.5, 1, 1.5, 2, 2.5, and 3 s after the onset of the ongoing white noise.

For sounds of condition S7, one additional noise snippet was embedded at 3.5 s, and, for sounds of

condition S8, two additional noise snippets were embedded at 3.5 and 4 s. Hence, conditions S6, S7, and

S8 contained an uninterrupted 2-Hz stimulus rhythm with 6, 7, and 8 cycles, respectively (Figure 2A).

Sounds for condition S7* also contained seven embeddings of a noise snippet, but the stimulus rhythm

was interrupted and contained one snippet omission in the 7th of the 8 cycles. Snippet onset times for

condition S7* were 0.5, 1, 1.5, 2, 2.5, 3, and 4 s (Figure 2A). Comparing activity in the 7th cycle between

S6/S7* and S7/S8 allows the investigation of self-sustainability of entrained neural activity. Comparing

activity in the 8th cycle between condition S7* and the other three conditions enables the examination

of neural sensitivity to a reoccurring noise snippet following a brief discontinuation in the regular, 2-Hz

snippet repetition.

Figure 2: Schematic of stimulus conditions and behavioral data for Experiment 1. **A:** Schematic of the stimulus conditions. The light gray part represents the longer noise stimulus and black boxes represent the embedded noise snippet. Dashed, vertical lines mark the onsets of the noise snippets for 8 cycles. **B:** Proportion of 'pattern-present' responses as a function of snippet duration in the threshold-estimation procedure. **C:** Behavioral data from the main experiment. Left: Mean d-prime values across participants. Right: Same data as displayed on the left-hand side, but here the between-participant variance was removed (Masson and Loftus, 2003). Large dots reflect the mean d-prime across participants, whereas the smaller dots reflect the d-prime for each participant. Error bars reflect the standard error of the mean. *$p_{Holm} < 0.05$

Each participant underwent the procedure to estimate the snippet duration that would yield a

rate of 0.75 for the detection of a snippet repetition (described above). For three participants, the

targeted detection rate was minimally lowered (but always ≥0.7) because of numerical issues resulting

from the logistic function fit used to estimate the threshold. The estimated duration was used

throughout the main part of the experiment. Participants listened to six blocks while EEG was recorded.

For two participants, EEG data from one block could not be used due to technical issues during recording.

Each block comprised 18 trials for each of the four stimulus conditions. A unique, newly created noise

snippet was generated for each trial. In addition, 6 trials without any embedded frozen noise snippets

were also presented in each block. These 'pattern-absent' trials were used to calculate the false alarm

rate for each participant. Hence, participants listened overall to 108 trials for each of the four conditions

containing a noise-snippet repetition ('pattern present') and 36 trials without an embedded noise

snippet ('pattern absent'; for similar procedures see Ng et al., 2012).

Analysis of behavioral data

D-prime values were analyzed using a one-way repeated measures analysis of variance (rmANOVA) with

the factor Condition (S6, S7, S7*, S8), followed up by planned t-tests and multiple-comparisons

correction using Holm's method. The Inclusion Bayes Factors (BF_{incl}) are also provided for the analysis

effects based on a Bayesian rmANOVA in JASP software (van den Bergh et al., 2019; JASP, 2022).

Neural responses to stimulus cycles

Analyses of neural data focused on stimulation cycle #7 and #8, for which the four conditions differed

(all four conditions contained an embedded noise snippet in the first six cycles). Neural activity was

averaged across the fronto-central electrode cluster (F3, Fz, F4, C3, Cz, C4) to focus on auditory-cortex

responses (Näätänen and Picton, 1987; Picton et al., 2003). Responses to the first five repetitions of a

noise snippet (cycles 2–6) show a positive deflection in the 0–0.2-s time window and a negative

deflection in the 0.2–0.4-s time window after snippet onset, jointly reflecting the entrained neural

activity (Figure 3). To obtain a response estimate separately for cycles #7 and #8, the amplitude in the

0.2–0.4-s time window after cycle onset was subtracted from the amplitude in the 0–0.2-s time window.

This resulted in one response-magnitude estimate for each of the four conditions (S6, S7, S7*, S8) and

each cycle (#7, #8).

To investigate whether entrained neural activity self-sustains during a brief interruption of a

snippet repetition, analyses focused on activity during the 7^{th} cycle (onset at 3.5 s). No noise snippet was

presented during the 7^{th} cycle for conditions S6 and S7*, whereas conditions S7 and S8 contained a noise

snippet in cycle 7. If entrained activity self-sustains during the interruption of the snippet repetition, a

larger than zero response magnitude for the S6/S7* conditions and no response difference between the

S6/S7* and the S7/S8 conditions would be expected. If entrained activity does not self-sustain during

discontinuation, the S7 and S8 conditions should elicited a larger response than the S6 and S7*

conditions, and responses for S6 and S7* may not differ from zero.

To investigate whether a noise snippet reoccurring after the interruption elicits a neural

response, analyses focused on neural activity during the 8^{th} cycle (onset at 4 s). Conditions S7* and S8,

but not conditions S6 and S7, contained a noise snippet in the 8^{th} cycle. Comparison of responses to S7*

relative to S6 and S7 will test whether the auditory system is sensitive to the reoccurrence of the noise

snippet, whereas comparison of responses to S7* relative to S8 will test whether the response to the

reoccurring noise snippet differs relative to the response to a snippet presented in an uninterrupted

sequence.

A one-way rmANOVA with the within-participants factor Condition (S6, S7, S7*, S8) was

calculated, separately for cycles #7 and #8, followed up by planned t-tests and multiple-comparisons

correction using Holm's method. The Inclusion Bayes Factors (BF_{incl}) are also provided for the rmANOVA

effects.

Results

The rmANOVA for perceptual sensitivity (d-prime) revealed a significant effect of Condition ($F_{3,48}$ =

17.205, p = $1 \cdot 10^{-7}$, ω^2 = 0.025, BF_{incl} = 101678). D-prime was larger for S8 compared to S6 (t_{16} = 5.756,

p_{Holm} = $1.4 \cdot 10^{-4}$, d = 1.396), S7 (t_{16} = 4.315, p_{Holm} = 0.002, d = 1.046), and S7* (t_{16} = 6.234, p_{Holm} = 7.2 ·

10^{-5}, d = 1.512). D-prime was also larger for S7 compared to S6, but only when Holm-correction was

omitted (t_{16} = 2.156, p_{Holm} = 0.140, p = 0.047, d = 0.523). S7* did not statistically differ from S6 or S7

(p_{Holm} > 0.24), indicating that there was no benefit from the reoccurring noise snippet for pattern

detection.

Inter-trial phase coherence was larger at the 2-Hz snippet-repetition frequency compared to

neighboring frequencies (t_{16} = 3.157, p = 0.006, d = 0.766; Figure 3), showing that neural activity

synchronized with the periodic presentation of frozen noise snippets. In fact, the amplitude of the neural

response to the snippet increased from the first to the second cycle (t_{16} = 3.037, p = 0.008, d = 0.737),

but did not further increase from the second to the third cycle (t_{16} = 0.074, p = 0.942, d = 0.018). That

only one repetition of the snippet was needed to elicit a full-magnitude response indicates that the brain

had some form of memory trace of the running noise prior to the first snippet repetition. The

topographical distribution shows largest ITPC values at fronto-central electrodes, suggesting that neural

responses originate from auditory cortex (Näätänen and Picton, 1987; Picton et al., 2003).

Figure 3: Neural responses to periodic repetitions of embedded noise snippets. A: Neural response time courses (averaged across channels from a fronto-central electrode cluster) for the four conditions time-locked to sound onset. Vertical, dashed lines mark the onsets of the embedded noise snippets. B: Inter-trial phase coherence (ITPC; averaged across channels from a fronto-central electrode cluster) and the topographical distribution of ITPC values for the 2-Hz stimulation frequency. ITPC was calculated across the four conditions containing a noise-snippet repetition (focusing on data from cycles 2-6). *p < 0.05

Analyses for neural activity during cycle #7 tested whether neural activity self-sustains during an

interruption of a snippet repetition. The rmANOVA revealed a significant effect of Condition ($F_{3,48}$ =

8.968, p = 8 · 10^{-5}, ω^2 = 0.259, BF_{incl} = 1451). Comparisons show that the response magnitude for S7 and

S8 was larger compared to S6 and S7* (S7 vs S6: t_{16} = 3.164, p_{Holm} = 0.024, d = 0.767; S7 vs S7*: t_{16} =

4.983, p_{Holm} = 8.1 · 10^{-4}, d = 1.208; S8 vs S6: t_{16} = 3.058, p_{Holm} = 0.024, d = 0.742; S8 vs S7*: t_{16} = 3.578,

p_{Holm} = 0.013, d = 0.868). Moreover, the response magnitude was significantly different from zero for S7

(t_{16} = 7.014, p = 2.9 × 10^{-6}, d = 1.7) and S8 (t_{16} = 4.038, p = 9.5 × 10^{-4}, d = 0.979), but not for S6 (t_{16} =

0.499, p = 0.624, d = 0.121) and S7* (t_{16} = -0.497, p = 0.626, d = 0.121), providing little evidence for self-

sustained periodic neural activity despite the hard task that required participants to focus attention on

the noise-snippet repetitions that created the stimulus regularity (Figure 4A).

Figure 4: Neural activity in the 7th and 8th stimulation cycle. **A:** Neural activity in cycle #7. S7 and S8 conditions, but not S6 and S7* conditions, contained a snippet in the 7th cycle. The mean neural response for each condition is depicted in the right plot – the response was calculated as the subtraction of the amplitude in the 0.2–0.4-s time window from the amplitude in the 0–0.2-s time window (left). The large dots reflect the mean amplitude across participants, whereas the smaller dots reflect the mean amplitude for each participant. **B:** Neural activity in cycle #8. S7* and S8 conditions, but not S6 and S7 conditions, contained a snippet in the 8th cycle. The plots are similar to panel A. Error bars reflect the standard error of the mean. *p_{Holm} < 0.05.

Analyses for neural activity during cycle #8 tested whether a reoccurring noise snippet following

an interruption in the regular repetition elicits a neural response. The rmANOVA revealed a significant

effect of Condition ($F_{3,48}$ = 8.433, p = 1.3 · 10^{-4}, ω^2 = 0.240, BF_{incl} = 738). Comparisons focused on the

differences in neural response magnitude for the S7* relative to the other conditions. The response

magnitude for S7* was greater than the response magnitude for S6 and S7 (S7* vs S6: t_{16} = 2.789, p_{Holm}

= 0.039, d = 0.676; S7* vs. 7: t_{16} = 3.260, p_{Holm} = 0.025, d = 0.791), whereas it did not differ from the

response magnitude for S8 (t_{16} = 0.686, p_{Holm} = 1, d = 0.166). Moreover, the response magnitude was

significantly different from zero for S7* (t_{16} = 3.573, p = 0.003, d = 0.867) and S8 (t_{16} = 5.252, p = 7.9 ×

10^{-5}, d = 1.274), but not for S6 (t_{16} = -0.257, p = 0.801, d = 0.062) and S7 (t_{16} = -0.75, p = 0.464, d = 0.182).

These results demonstrates that the auditory system maintains a memory trace of the specific spectro-

temporal structure of a noise snippet embedded in an ongoing noise despite an interruption of a

previously established stimulus regularity (Figure 4B).

Summary

The results of Experiment 1 show little evidence that neural activity entrained by a stimulus rhythm self-

sustains after the rhythm discontinues (Figure 4A). Critically, a frozen noise snippet reoccurring after a

brief interruption elicited a neural response that appeared similar in magnitude compared to the neural

response elicited by a noise snippet that was preceded by uninterrupted snippet repetitions (Figure 4B).

The results show that, once the auditory cortex establishes a memory trace for the noise snippet, it

faithfully responds to another repetition of a noise snippet despite an interruption in rhythmic

stimulation.

Stimulation in Experiment 1 was regular at a 2-Hz rate and the snippet reoccurred after the

interruption in-sync – that is, at an integer interval – with the preceding stimulus rhythm. A large body

of behavioral and electrophysiological work has shown that perceptual and neural sensitivity to a

stimulus can depend on its temporal relation to a preceding stimulus rhythm, that is, whether it occurs

in-sync or out-of-sync with it (Jones and Boltz, 1989; Large and Jones, 1999; Barnes and Jones, 2000;

Jones et al., 2002; Schroeder and Lakatos, 2009; Cravo et al., 2013; Henry and Herrmann, 2014; Henry

et al., 2014; Herrmann et al., 2016; Nobre and van Ede, 2018). In Experiment 1, the timing of the

reoccurring noise snippet was in-sync with the preceding stimulus rhythm, which perhaps explains the

strong response it elicited. Experiment 2 was conducted in order to investigate directly whether the

temporal relation to a preceding stimulus rhythm affects neural sensitivity to a reoccurring noise

snippet.

361   Experiment 2

Methods and Materials

Participants

Twenty individuals participated in Experiment 2 (median age: 21 years; range: 18–34 years; 6 male, 14

female). None of the participants who took part in Experiment 1 took part in Experiment 2.

Stimuli and procedure

Participants listened to 5-s white-noise sounds in four different conditions (Figure 5). Each condition

contained a repetition of a noise snippet, as in Experiment 1. Sounds for the 'absent' condition contained

six embeddings of a noise snippet, at 0.5, 1, 1.5, 2, 2.5, and 3 s after the onset of the ongoing white noise

(same as S6 in Experiment 1). The label 'absent' here refers to the absence of a reoccurring noise snippet

after the regular snippet repetition discontinued. The other three conditions contained one additional

snippet reoccurring after a brief discontinuation in the stimulus rhythm. In the 'early', 'on-time', and

'late' conditions, the reoccurring snippet was embedded at 3.75, 4, and 4.25 s, respectively. The snippet

onset for the 'early' and 'late' conditions was temporally placed in anti-phase (out-of-sync) relative to

the 2-Hz stimulus rhythm established by the preceding 6 snippet embeddings. The snippet onset for the

'on-time' condition was placed in-sync/in-phase with the preceding stimulus rhythm (same as 7* in

Experiment 1).

Figure 5: Schematic of stimulus conditions and behavioral data for Experiment 2. **A:** Schematic of the stimulus conditions. The light gray part represents the longer noise stimulus and black boxes represent the embedded noise snippet. Dashed, vertical lines mark the onsets of the noise snippets. **B:** Proportion of 'pattern-present' responses as a function of snippet duration in the threshold-estimation procedure. **C:** Behavioral data from the main experiment. Left: Mean d-prime values across participants. Right: Same data as displayed on the left-hand side, but here the between-participant variance was removed (Masson and Loftus, 2003). Large dots reflect the mean d-prime across participants, whereas the smaller dots reflect the d-prime for each participant. Error bars reflect the standard error of the mean. The were no significant differences between conditions.

Similar to Experiment 1, each participant underwent the procedure to estimate the snippet

duration that would yield a rate of 0.75 for the detection of a snippet repetition (see descriptions above).

For one participant, the targeted detection rate was minimally lowered (but ≥0.7) because of numerical

issues resulting from the logistic function fit. The estimated duration was used throughout the main

experimental procedures. Participants listened to six blocks while EEG was recorded. Each block

comprised 18 trials for each of the four stimulus conditions. As for Experiment 1, a unique, newly created

noise snippet was generated for each trial. Six trials without any embedded frozen noise snippets were

also presented in each block to calculate the false alarm rate for each participant (Ng et al., 2012).

Participants listened overall to 108 trials for each of the four conditions containing a noise-snippet

repetition ('pattern present') and 36 trials without an embedded noise snippet ('pattern absent').

Analysis of behavioral data

D-prime values were analyzed using a one-way rmANOVA with the within-participants factor Condition

(absent, early, on-time, late). A significant effect was resolved by post hoc comparisons and multiple-

comparisons correction using Holm's method. The Inclusion Bayes Factors (BF_{incl}) are also provided for

the rmANOVA effects based on a Bayesian rmANOVA in JASP software (van den Bergh et al., 2019; JASP,

2022).

Neural responses to sequence-final noise snippet

As for Experiment 1, neural activity was averaged across electrodes of the fronto-central electrode

cluster (F3, Fz, F4, C3, Cz, C4), focusing on auditory-cortex responses. An estimate of the response

magnitude to the embedded noise snippet was obtained by subtracting the amplitude in the 0.2–0.4-s

time window after snippet onset from the amplitude in the 0–0.2-s time window.

Responses were extracted for the 'early' (3.75–4.25 s), 'on-time' (4–4.5 s), and 'late' (4.25–4.75

s) conditions in order to investigate whether the onset-time of the reoccurring noise snippet relative to

the preceding stimulus rhythm affects the neural response to the reoccurring snippet. As a control,

neural responses were extracted from the same time windows for 'absent' trials, during which no

snippet reoccurred at 3.75, 4, nor 4.25 s.

A two-way rmANOVA was calculated using the within-participants factors Condition (early, on-

time, late) and Snippet Presence (snippet absent, snippet present). A significant Condition effect was

resolved by post hoc comparisons and multiple-comparisons correction using Holm's method. Inclusion

Bayes Factors (BF_{incl}) are provided for the rmANOVA effects.

Results

409   The rmANOVA for the behavioral data was not significant ($F_{3,57}$ = 0.912, p = 0.441, ω^2 < 0.001, BF_{incl} =

410   0.174; Figure 5C), that is, behavioral performance did not differ between conditions. The absence of a

411   behavioral effect is consistent with the results from Experiment 1, where a snippet that reoccurred after

412   a brief interruption of a stimulus rhythm also did not lead to a pattern-detection benefit relative to when

413   no snippet reoccurred (S7* vs. S6; Figure 2C).

Figure 6: Neural synchronization to periodic repetitions of noise snippets. A: Neural response time courses (averaged across electrodes from a fronto-central cluster) for the four conditions time-locked to sound onset. Vertical, dashed lines mark the onsets of the noise snippets. B: Inter-trial phase coherence (ITPC; averaged across electrodes from a fronto-central cluster) and the topographical distribution of ITPC values for the 2-Hz stimulation frequency. ITPC was calculated across the four conditions containing a noise-snippet repetition (focusing on data from cycles 2-6). *p < 0.05

As for Experiment 1, neural activity synchronized with the regular repetition of embedded, frozen

noise snippets. That is, ITPC was larger at the 2-Hz frequency at which snippets were repeated compared

to ITPC at neighboring frequencies (t_{19} = 3.325, p = 0.004, d = 0.744; Figure 6). Similar to Experiment 1,

the amplitude of the neural response increased from the first to the second snippet (t_{19} = 5.857, p = 1.2

× 10^{-5}, d = 1.310), but did not further increase from the second to the third snippet (t_{19} = 0.334, p =

0.742, d = 0.075). The topographical distribution shows largest ITPC values at fronto-central electrodes,

again suggesting that synchronized neural activity originates from the auditory cortex (Näätänen and

Picton, 1987; Picton et al., 2003). Pooling ITPC data across the two experiments revealed that ITPC at

the 2-Hz stimulation frequency (relative to ITPC at neighboring frequencies) was larger for participants

who required a longer snippet duration for achieving a similar detection threshold (r = 0.453, p = 0.005;

independent of conditions, because the snippet duration was identical across conditions for each

participant).

Figure 7: Neural responses to a reoccurring noise snippet at different times following a brief interruption in the regular snippet repetition. A: Response time courses for each condition (early, on-time, late) and the response time course for the 'absent' condition for each of the corresponding time windows (early: 3.75–4.25 s; on-time: 4–4.5 s; late: 4.25–4.75 s; x-axis is time-locked to snippet onset for each of these time windows). **B:** Mean response amplitude, calculated as the subtraction of the amplitude in the 0.2–0.4-s time window from the amplitude in the 0–0.2-s time window. **C:** Mean response amplitude for the three timing conditions (early, on-time, late) normalized by the 'absent' condition (left; the 'absent' condition was subtracted from the other three conditions) and non-normalized (right). The large dots reflect the mean amplitude across participants, whereas the smaller dots reflect the mean amplitude for each participant. Error bars reflect the standard error of the mean. Responses were significantly larger for the 'snippet present' compared to the 'snippet absent' conditions, but there were no significant differences between timing conditions.

The noise snippet reoccurring after a brief interruption in the stimulus rhythm elicited a larger

neural response than was elicited by the 'absent' condition, for which no snippet reoccurred (main effect

of Snippet Presence: $F_{1,19}$ = 32.076, p = 1.8 · 10^{-5}, ω^2 = 0.48, BF_{incl} = 606; Figure 7B). There was no effect

of Condition ($F_{2,38}$ = 0.488, p = 0.618, ω^2 < 0.001, BF_{incl} = 0.265) and no Condition × Snippet Presence

interaction ($F_{2,38}$ = 2.186, p = 0.126, ω^2 = 0.04, BF_{incl} = 0.789; Figure 7C, left). In other words, the three

conditions containing a reoccurring snippet elicited a response (relative to snippet absence),

demonstrating that the auditory system maintains sensitivity to reoccurring noise snippets embedded

in a longer noise sound regardless of whether the snippet reoccurs in-sync or out-of-sync with a previous

stimulus rhythm. There was also no difference in the response magnitude between the three timing

conditions when responses were not normalized relative to the 'absent' condition (effect of Condition:

$F_{2,38}$ = 1.843, p = 0.172, ω^2 = 0.019, BF_{incl} = 0.835; Figure 7C, right).

<div align="center">Discussion</div>

The current study investigated how the brain maintains a neural representation of repeating structure in sounds. In two EEG experiments, participants listened to a longer, ongoing white-noise sound which comprised shorter, frozen noise snippets that repeated at a regular rate (Andrillon et al., 2015; Andrillon et al., 2017; Dauer et al., 2022). In several conditions, the snippet repetition discontinued for a brief period after which a noise snippet reoccurred. Results show that neural activity entrained with the regular snippet repetition, but that this periodic neural activity does not sustain during the discontinuation period. Nevertheless, the auditory cortex responds to the reoccurring noise snippet similarly compared to a noise snippet for which the snippet repetition is not discontinued. This response invariance holds regardless of the degree of synchrony of the reoccurring noise snippet relative to the previously established stimulus regularity. Overall, the current study demonstrates that, for the short timescale investigated here, the auditory cortex sensitively responds to a previously learned noise structure in an ongoing sound independent of temporal interruptions.

Self-sustainability of neural activity

Neural activity becomes entrained by repeating sounds or structure in sounds (Picton et al., 2003; Lakatos et al., 2008; Stefanics et al., 2010; Nozaradan et al., 2011; Thut et al., 2011; Henry and Obleser, 2012; Nozaradan et al., 2012; Lakatos et al., 2013b; ten Oever et al., 2017; Lakatos et al., 2019; Obleser and Kayser, 2019; Herrmann et al., 2023). The peak at 2 Hz in the phase-coherence spectrum of the current study is consistent with entrained activity (Figures 3 and 6). Entrained neural activity – if oscillatory in nature – is hypothesized to self-sustain after periodic stimulation discontinues (Pikovsky et al., 2001; Lakatos et al., 2013b; Henry and Herrmann, 2014; Keitel et al., 2014; Lin et al., 2021; van Bree et al., 2021; Keitel et al., 2022), but there was no evidence for self-sustainability in the current study. That is, there was no clear neural responses in the time window during which the next snippet would have occurred if the repetition had continued (S6 and S7* in Experiment 1; Figure 4). Moreover, neural activity was smaller than activity elicited by a snippet presented in the same time window as part of the snippet repetition (Figures 4; see responses for cycle #7: S6/S7* vs S7/S8; and responses for cycle #8: S6/S7 vs S8).

The current stimulus paradigm may arguably be well suited for the investigation of self-sustained neural activity, assuming its existence, because there are no acoustic edges or changes associated with the occurrence of the noise snippet nor with the discontinuation of the snippet repetition (Figure 1;

Agus et al., 2010; Agus and Pressnitzer, 2013; Luo et al., 2013; Andrillon et al., 2015; Rajendran et al.,

2016; Andrillon et al., 2017; Dauer et al., 2022). The longer, ongoing noise continues without

interruption. Moreover, the detection of the regular repetition of frozen noise snippets is challenging

(as indicated by the off-ceiling d-prime values in Figures 2 and 5, and the titration of the detection rate

to 0.75) and requires participants to focus attention to the sounds. Attention is likely a facilitator for

entrained neural activity and self-sustainability (Lakatos et al., 2013b; Lakatos et al., 2013a; Henry and

Herrmann, 2014; Ringer et al., 2023a). Since the current study finds little evidence for self-sustainability,

this adds to the ongoing discussion about whether entrained neural activity self-sustains when a stimulus

periodicity discontinues (Lakatos et al., 2013b; Lin et al., 2021; van Bree et al., 2021). The absence of

self-sustained activity may further raise the question of whether neural activity entrained by the stimulus

regularity reflects neural oscillatory activity or event-evoked activity (Klimesch et al., 2007; Sauseng et

al., 2007; Capilla et al., 2011; Fellinger et al., 2011; Notbohm et al., 2016; Keitel et al., 2021). Self-

sustained activity would not be expected for event-evoked activity. The response to a snippet

reoccurring after the regular repetition had discontinued was very similar to the response to the snippet

when it was part of the regular repetition (S7* vs S8 in Experiment 1), which is in line with event-evoked

neural activity in the current study.

Invariant neural response to the reoccurring frozen noise snippet

The current study shows that the neural response to a frozen noise snippet, reoccurring after a regular

snippet repetition had briefly discontinued, is robust and invariant to a brief discontinuation (Figure 4).

That is, the response to the reoccurring noise snippet did not differ from the response to a noise snippet

for which the regular repetition was not discontinued previously. The response was further invariant to

changes in the temporal relation between the onset of the reoccurring snippet relative to the previous

stimulus regularity (Figure 7). Overall, it appears that once a neural representation for a specific noise

snippet is established, the brain responds to the reoccurrence of the snippet invariantly, at least over

the short timescale investigated here.

Several previous works have shown that individuals better detect a noise snippet repetition when

the snippet that is repeated recurs across trials compared to a snippet that does not recur (Agus et al.,

2010; Agus and Pressnitzer, 2013; Andrillon et al., 2015; Andrillon et al., 2017; Dauer et al., 2022).

Previous work further shows that individuals show behavioral correlates of memory for random noise

or tone structure in sounds even days and weeks after exposure (Agus et al., 2010; Viswanathan et al.,

2016; Bianco et al., 2020; Bianco et al., 2023). That the brain responds with similar magnitude to a

reoccurring noise snippet following an interruption compared to a snippet that is part of a regular, non-
interrupted sequence is consistent with a memory representation for a noise snippet that lasts long once
the representation has been established.

Relation between neural responses and behavioral detection performance

The results of the current study show that participants are more sensitive to the presence of a repetition
of a frozen-noise snippet in an ongoing noise when the number of regular, uninterrupted repetitions
increases (Figure 2C; Experiment 1). However, an additional reoccurring noise snippet following a brief
interruption in the repetition did not improve detection performance (Figures 2C and 5C). The absence
of a benefit for repetition-detection was observed for reoccurring snippets that were in- and out-of-sync
with the preceding stimulus regularity. While some previous works have shown perceptual benefits for
in-sync compared to out-of-sync stimuli (Jones et al., 2002; Jones et al., 2006; Hickok et al., 2015;
Herrmann et al., 2016), other recent research has failed to find such a perceptual benefit (Bauer et al.,
2015; de Graaf and Duecker, 2022; Lin et al., 2022; Sun et al., 2022). Stimulus designs often differed
substantially among previous studies, making it hard to pinpoint under which conditions perceptual
benefits for in-sync compared to out-of-sync stimuli can be observed.

Although the brain responded robustly to the reoccurring noise snippet, this apparently did not
translate to a behavioral performance increase (Figures 2C and 5C). Moreover, neural responses to the
frozen noise snippet were already present for the first snippet repetition (cycle #2) and did not further
increase with the number of repetitions, whereas behavioral detection performance increased with the
number of repetitions (Figure 2C). Other work also suggests that one repetition of a noise snippet is
sufficient for the brain to elicit a strong response, although this was not strictly analyzed previously
(Andrillon et al., 2015; Ringer et al., 2023a). Moreover, neural entrainment to periodic sound stimulation
has been observed even for sub-threshold sound stimulation (e.g., tone beeps in noise at a low signal-
to-noise ratio; ten Oever et al., 2017). This mismatch between the pattern of neural responses and the
behavioral patterns perhaps suggests that a memory trace of the noise snippet is established relatively
quickly and stably, whereas the detection of the repetition requires another, possibly higher cognitive
process involving decision making.

Conclusions

The current study investigated how the brain maintains representations of repeating structure in sounds. Participants listened to a longer, ongoing white noise sound which comprised shorter, frozen noise snippets that repeated at a regular rate. The current study aimed to answer whether neural activity self-sustains during a brief interruption of the repetition and how the brain responds to a noise snippet that reoccurs after the interruption. No evidence for self-sustained neural activity during the interruption was observed. However, the auditory cortex responded to a reoccurring noise snippet with similar magnitude compared to a noise snippet for which the snippet repetition had not been interrupted. This response invariance was observed for different onset times of the reoccurring noise snippet relative to the previously established regularity. The current study thus demonstrates that the auditory cortex sensitively responds to a previously learned noise structure in an ongoing sound independent of temporal interruptions.

Acknowledgements

We thank Christie Tsagopoulos for her help with data collection for both experiments. The research was supported by the Canada Research Chair program (CRC-2019-00156, 232733) and the Natural Sciences and Engineering Research Council of Canada (Discovery Grant: RGPIN-2021-02602).

Declaration of conflict of interest

None.

References

Adrian ED, Matthews BHC (1934) The Berger Rhythm: Potential Changes From The Occipital Lobes in Man. Brain 57:355-385.

Agus TR, Pressnitzer D (2013) The detection of repetitions in noise before and after perceptual learning. The Journal of the Acoustical Society of America 134:464-473.

Agus TR, Thorpe SJ, Pressnitzer D (2010) Rapid Formation of Robust Auditory Memories: Insights from Noise. Neuron 66:610-618.

Andrillon T, Kouider S, Agus TR, Pressnitzer D (2015) Perceptual Learning of Acoustic Noise Generates
Memory-Evoked Potentials. Current Biology 25:2823-2829.

Andrillon T, Pressnitzer D, Léger D, Kouider S (2017) Formation and suppression of acoustic memories
during human sleep. Nature Communications 8:179.

Baese-Berk MM, Heffner CC, Dilley LC, Pitt MA, Morrill TH, McAuley JD (2014) Long-term temporal
tracking of speech rate affects spoken-word recognition. Psychological Science 25:1546-1553.

Barascud N, Pearce MT, Griffiths TD, Friston KJ, Chait M (2016) Brain responses in humans reveal ideal
observer-like sensitivity to complex acoustic patterns. Proceedings of the National Academy of
Sciences 113:E616-E625.

Barnes R, Jones MR (2000) Expectancy, Attention, and Time. Cognitive Psychology 41:254-311.

Bauer A-KR, Jaeger M, Thorne JD, Bendixen A, Debener S (2015) The auditory dynamic attending theory
revisited: A closer look at the pitch comparison task. Brain Research 1626:198-210.

Bell AJ, Sejnowski TJ (1995) An information maximization approach to blind separation and blind
deconvolution. Neural Computation 7:1129-1159.

Bendixen A (2014) Predictability effects in auditory scene analysis: a review. Frontiers in Neuroscience
8:Article 60.

Bianco R, Hall ETR, Pearce MT, Chait M (2023) Healthy ageing effects on implicit auditory memory: from
encoding to 6-month retention. BioRxiv.

Bianco R, Harrison PMC, Hu M, Bolger C, Picken S, Pearce MT, Chait M (2020) Long-term implicit memory
for sequential auditory patterns in humans. eLife 9:e56073.

Bishop GH (1933) Cyclic changes in excitability of the optic pathway of the rabbit. American Journal of
Physiology 103:213-224.

Capilla A, Pazo-Álvarez P, Darriba A, Campo P, Gross J (2011) Steady-State Visual Evoked Potentials Can
Be Explained by Temporal Superposition of Transient Event-Related Responses. PLoS ONE
6:e14543.

Cravo AM, Rohenkohl G, Wyart V, Nobre AC (2013) Temporal Expectation Enhances Contrast Sensitivity
by Phase Entrainment of Low-Frequency Oscillations in Visual Cortex. The Journal of Neuroscience
33:4002-4010.

Dauer TD, Henry MJ, Herrmann B (2022) Auditory perceptual learning depends on temporal regularity
and certainty. Journal of Experimental Psychology: Human Perception and Performance 48:755-
770.

de Graaf TA, Duecker F (2022) No effects of rhythmic visual stimulation on target discrimination: An
online alpha entrainment experiment. European Journal of Neuroscience 55:3340-3351.

Fellinger R, Klimesch W, Gruber W, Freunberger R, Doppelmayr M (2011) Pre-stimulus alpha phase-
alignment predicts P1-amplitude. Brain Research Bulletin 85:417-423.

Giraud A-L, Poeppel D (2012) Cortical oscillations and speech processing: emerging computational
principles and operations. Nature Neuroscience 15.

Guttman N, Julesz B (1963) Lower Limits of Auditory Periodicity Analysis. The Journal of the Acoustical
Society of America 35:610-610.

Henry MJ, Obleser J (2012) Frequency modulation entrains slow neural oscillations and optimizes human
listening behavior. Proceedings of the National Academy of Sciences 109:20095-20100.

Henry MJ, Herrmann B (2014) Low-Frequency Neural Oscillations Support Dynamic Attending in
Temporal Context. Timing & Time Perception 2:62-86.

Henry MJ, Herrmann B, Obleser J (2014) Entrained neural oscillations in multiple frequency bands co-
modulate behavior. Proceedings of the National Academy of Sciences 111:14935-14940.

Herrmann B, Maess B, Friederici AD (2011a) Violation of syntax and prosody - Disentangling their
contributions to the early left anterior negativity (ELAN). Neuroscience Letters 490:116-120.

Herrmann B, Augereau T, Johnsrude IS (2020) Neural Responses and Perceptual Sensitivity to Sound
Depend on Sound-Level Statistics. Scientific Reports 10:9571.

Herrmann B, Araz K, Johnsrude IS (2021) Sustained neural activity correlates with rapid perceptual
learning of auditory patterns. NeuroImage 238:118238.

Herrmann B, Maess B, Johnsrude IS (2023) Age-related changes in neural synchronization and the
sustained response to amplitude and frequency modulation of sound. Hearing Research
428:108677.

Herrmann B, Henry MJ, Grigutsch M, Obleser J (2013) Oscillatory Phase Dynamics in Neural Entrainment
Underpin Illusory Percepts of Time. The Journal of Neuroscience 33:15799-15809.

Herrmann B, Henry MJ, Haegens S, Obleser J (2016) Temporal expectations and neural amplitude
fluctuations in auditory cortex interactively influence perception. NeuroImage 124:487-497.

Herrmann B, Maess B, Hahne A, Schröger E, Friederici AD (2011b) Syntactic and auditory spatial
processing in the human temporal cortex: An MEG study. NeuroImage 57:624-633.

Hickok G, Farahbod H, Saberi K (2015) The Rhythm of Perception: Entrainment to Acoustic Rhythms
Induces Subsequent Perceptual Oscillation. Psychological Science 26:1006-1013.

Hodapp A, Grimm S (2021) Neural signatures of temporal regularity and recurring patterns in random tonal sound sequences. European Journal of Neuroscience 53:2740-2754.

Idemaru K, Holt LL (2011) Word Recognition Reflects Dimension-based Statistical Learning. Journal of Experimental Psychology: Human Perception and Performance 37:1939-1956.

Irsik VC, Almanaseer A, Johnsrude IS, Herrmann B (2021) Cortical Responses to the Amplitude Envelopes of Sounds Change with Age. The Journal of Neuroscience 41:5045-5055.

JASP (2022) JASP [Computer software]. In: https://jasp-stats.org/.

Jones MR, Boltz MG (1989) Dynamic Attending and Responses to Time. Psychological Review 96:459-491.

Jones MR, Johnston HM, Puente J (2006) Effects of auditory pattern structure on anticipatory and reactive attending. Cognitive Psychology 53:59-96.

Jones MR, Moynihan H, MacKenzie N, Puente J (2002) Temporal Aspects of Stimulus-Driven Attending in Dynamic Arrays. Psychological Science 13:313-319.

Kaernbach C (2004) The Memory of Noise. Experimental Psychology 51:240-248.

Kang H, Agus TR, Pressnitzer D (2017) Auditory memory for random time patterns. The Journal of the Acoustical Society of America 142:2219-2232.

Kang H, Macherey O, Roman S, Pressnitzer D (2021) Auditory memory for random time patterns in cochlear implant listeners. The Journal of the Acoustical Society of America 1934:1934-1944.

Kayser SJI, Robin A. A., Gross J, Kayser C (2015) Irregular Speech Rate Dissociates Auditory Cortical Entrainment, Evoked Responses, and Frontal Alpha. The Journal of Neuroscience 35:14691-14701.

Keitel C, Quigley C, Ruhnau P (2014) Stimulus-Driven Brain Oscillations in the Alpha Range: Entrainment of Intrinsic Rhythms or Frequency-Following Response? The Journal of Neuroscience 34:10137-10140.

Keitel C, Obleser J, Jessen S, Henry MJ (2021) Frequency-Specific Effects in Infant Electroencephalograms Do Not Require Entrained Neural Oscillations: A Commentary on Köster et al. (2019). Psychological Science 32:966-971.

Keitel C, Ruzzoli M, Dugué L, Busch NA, Benwell CSY (2022) Rhythms in cognition: The evidence revisited. European Journal of Neuroscience 55:2991-3009.

Klimesch W, Sauseng P, Hanslmayr S, Gruber W, Freunberger R (2007) Event-related phase reorganization may explain evoked neural dynamics. Neuroscience and Biobehavioral Reviews 31:1003-1016.

Lachaux J-P, Rodriguez E, Martinerie J, Varela FJ (1999) Measuring Phase Synchrony in Brain Signals.
Human Brain Mapping 8:194-208.

Lakatos P, Gross J, Thut G (2019) A New Unifying Account of the Roles of Neuronal Entrainment. Current
Biology 29:R890-R905.

Lakatos P, Schroeder CE, Leitman DI, Javitt DC (2013a) Predictive Suppression of Cortical Excitability and
Its Deficit in Schizophrenia. The Journal of Neuroscience 33:11692-11702.

Lakatos P, Karmos G, Mehta AD, Ulbert I, Schroeder CE (2008) Entrainment of Neuronal Oscillations as
a Mechanism of Attentional Selection. Science 320:110-113.

Lakatos P, Shah AS, Knuth KH, Ulbert I, Karmos G, Schroeder CE (2005) An Oscillatory Hierarchy
Controlling Neuronal Excitability and Stimulus Processing in the Auditory Cortex. Journal of
Neurophysiology 94:1904-1911.

Lakatos P, Musacchia G, O'Connel MN, Falchier AY, Javitt DC, Schroeder CE (2013b) The Spectrotemporal
Filter Mechanism of Auditory Selective Attention. Neuron 77:750-761.

Lange K (2010) Can a regular context induce temporal orienting to a target sound? International Journal
of Psychophysiology 78:231-238.

Large EW, Jones MR (1999) The Dynamics of Attending: How People Track Time-Varying Events.
Psychological Review 106:119-159.

Lin WM, Oetringer DA, Bakker-Marshall I, Emmerzaal J, Wilsch A, ElShafei HA, Rassi E, Haegens S (2022)
No behavioural evidence for rhythmic facilitation of perceptual discrimination. European Journal of
Neuroscience 55:3352-3364.

Lin Y-J, Shukla L, Dugué L, Valero-Cabré A, Carrasco M (2021) Transcranial magnetic stimulation entrains
alpha oscillatory activity in occipital cortex. Scientific Reports 11:18562.

Luo H, Tian X, Song K, Zhou K, Poeppel D (2013) Neural Response Phase Tracks How Listeners Learn New
Acoustic Representations. Current Biology 23:968-974.

Macmillan NA, Creelman CD (2004) Detection Theory: A User's Guide. New Jersey, USA: Psychology
Press.

Maess B, Herrmann CS, Hahne A, Nakamura A, Friederici AD (2006) Localizing the distributed language
network responsible for the N400 measured by MEG during auditory sentence processing. Brain
Research 1096:163-172.

Makeig S, Bell AJ, Jung T-P, Sejnowski TJ (1996) Independent component analysis of
electroencephalographic data. In: Advances in Neural Information Processing Systems (Touretzky
D, Mozer M, Hasselmo M, eds). Cambridge, MA, USA: MIT Press.

Masson MEJ, Loftus GR (2003) Using confidence intervals for graphically based data interpretation.
Canadian Journal of Experimental Psychology and Aging 57:203-220.

McAuley JD, Jones MR (2003) Modeling Effects of Rhythmic Context on Perceived Duration: A
Comparison of Interval and Entrainment Approaches to Short-Interval Timing. Journal of
Experimental Psychology: Human Perception and Performance 29:1102-1125.

McDermott Josh H, Simoncelli Eero P (2011) Sound Texture Perception via Statistics of the Auditory
Periphery: Evidence from Sound Synthesis. Neuron 71:926-940.

McDermott JH, Schemitsch M, Simoncelli EP (2013) Summary statistics in auditory perception. Nature
Neuroscience 16:493-498.

Morillon B, Schroeder CE, Wyart V, Arnal LH (2016) Temporal Prediction in lieu of Periodic Stimulation.
The Journal of Neuroscience 36:2342-2347.

Näätänen R, Picton TW (1987) The N1 wave of the human electric and magnetic response to sound: a
review and an analysis of the component structure. Psychophysiology 24:375-425.

Ng BSW, Schroeder T, Kayser C (2012) A Precluding But Not Ensuring Role of Entrained Low-Frequency
Oscillations for Auditory Perception. The Journal of Neuroscience 32:12268-12276.

Nobre AC, van Ede F (2018) Anticipated moments: temporal structure in attention. Nature Reviews
Neuroscience 19:34-48.

Notbohm A, Kurths J, Herrmann CS (2016) Modification of Brain Oscillations via Rhythmic Light
Stimulation Provides Evidence for Entrainment but Not for Superposition of Event-Related
Responses. Frontiers in Human Neuroscience 10.

Nozaradan S, Peretz I, Mouraux A (2012) Selective Neuronal Entrainment to the Beat and Meter
Embedded in a Musical Rhythm. The Journal of Neuroscience 32:17572-17581.

Nozaradan S, Peretz I, Missal M, Mouraux A (2011) Tagging the Neuronal Entrainment to Beat and Meter.
The Journal of Neuroscience 31:10234-10240.

Obleser J, Kayser C (2019) Neural Entrainment and Attentional Selection in the Listening Brain. Trends
in Cognitive Sciences 23:913-926.

Oostenveld R, Fries P, Maris E, Schoffelen JM (2011) FieldTrip: Open source software for advanced
analysis of MEG, EEG, and invasive electrophysiological data. Computational Intelligence and
Neuroscience 2011:Article ID 156869.

Peelle JE, Davis MH (2012) Neural oscillations carry speech rhythm through to comprehension. Frontiers
in Psychology 3:Article 320.

Picton TW, John SM, Dimitrijevic A, Purcell DW (2003) Human auditory steady-state responses.
International Journal of Audiology 42:177-219.

Pikovsky A, Rosenblum M, Kurths J (2001) Synchronization: A Universal Concept in Nonlinear Sciences.
Cambridge: Cambridge University Press.

Rajendran VG, Harper NS, Abdel-Latif KHA, Schnupp JWH (2016) Rhythm Facilitates the Detection of
Repeating Sound Patterns. Frontiers in Neuroscience 10.

Ringer H, Schröger E, Grimm S (2022) Perceptual Learning and Recognition of Random Acoustic Patterns.
Auditory Perception & Cognition 5:259-281.

Ringer H, Schröger E, Grimm S (2023a) Perceptual learning of random acoustic patterns: Impact of
temporal regularity and attention. BioRxiv.

Ringer H, Schröger E, Grimm S (2023b) Within- and between-subject consistency of perceptual
segmentation in periodic noise: A combined behavioral tapping and EEG study. Psychophysiology
60:e14174.

Sauseng P, Klimesch W, Gruber WR, Hanslmayr S, Freunberger R, Doppelmayr M (2007) Are event-
related potential components generated by phase resetting of brain oscillations? A critical
discussion. Neuroscience 146:1435-1444.

Schroeder CE, Lakatos P (2009) Low-frequency neuronal oscillations as instruments of sensory selection.
Trends in Neurosciences 32:9-18.

Schröger E (2005) The Mismatch Negativity as a Tool to Study Auditory Processing. Acta Acustica united
with Acustica 91:490-501.

Schröger E (2007) Mismatch Negativity: A Microphone into Auditory Memory. Journal of
Psychophysiology 21:138-146.

Sohoglu E, Chait M (2016) Detecting and representing predictable structure during auditory scene
analysis. eLife 5:e19113.

Stefanics G, Hangya B, Hernádi I, Winkler I, Lakatos P, Ulbert I (2010) Phase Entrainment of Human Delta
Oscillations Can Mediate the Effects of Expectation on Reaction Speed. The Journal of Neuroscience
30:13578-13585.

Sun Y, Michalareas G, Poeppel D (2022) The impact of phase entrainment on auditory detection is highly
variable: Revisiting a key finding. European Journal of Neuroscience 55:3373-3390.

Teki S, Chait M, Kumar S, Shamma SA, Griffiths TD (2013) Segregation of complex acoustic scenes based
on temporal coherence. eLife 2:e00699.

Teki S, Barascud N, Picard S, Payne C, Griffiths TD, Chait M (2016) Neural Correlates of Auditory Figure-
Ground Segregation Based on Temporal Coherence. Cerebral Cortex 26:3669-3680.

ten Oever S, Schroeder CE, Poeppel D, van Atteveldt N, Metha AD, Megevand P, Groppe DM, Zion-
Golumbic E (2017) Low-Frequency Cortical Oscillations Entrain to Subthreshold Rhythmic Auditory
Stimuli. The Journal of Neuroscience 37:4903-4912.

Thut G, Schyns PG, Gross J (2011) Entrainment of perceptually relevant brain oscillations by non-invasive
rhythmic stimulation of the human brain. Frontiers in Psychology 2:Article 170.

van Bree S, Sohoglu E, Davis MH, Zoefel B (2021) Sustained neural rhythms reveal endogenous
oscillations supporting speech perception. PLOS Biology 19:e3001142.

van den Bergh D, van Doorn J, Marsman M, Draws T, van Kesteren E-J, Derks K, Dablander F, Gronau Q,
Kucharský Š, Gupta ARKN, Sarafoglou A, Voelkel JG, Stefan A, Ly A, Hinne M, Matzke D,
Wagenmakers E-J (2019) A Tutorial on Conducting and Interpreting a Bayesian ANOVA in JASP.
PsychArXiv.

Viswanathan J, Rémy F, Bacon-Macé N, Thorpe SJ (2016) Long Term Memory for Noise: Evidence of
Robust Encoding of Very Short Temporal Acoustic Patterns. Frontiers in Neuroscience 10:490.

Warren RM, Bashford Jr. JA, Cooley JM, Brubaker BS (2001) Detection of acoustic repetition for very long
stochastic patterns. Perception & Psychophysics 63:175-182.

Winkler I, Denham SL, Nelken I (2009) Modeling the auditory scene: predictive regularity representations
and perceptual objects. Trends in Cognitive Sciences 13:532-540.

756