Control of telomere length in yeast by SUMOylated PCNA and the Elg1 PCNA unloader

Pragyan Singh, Inbal Gazy and Martin Kupiec*

[1]The Shmunis School of Biomedicine and Cancer Research, The George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv 69978, Israel

***Corresponding author: martin@tauex.tau.ac.il**

ABSTRACT

Telomeres cap and protect the linear eukaryotic chromosomes. Telomere length is determined by an equilibrium between positive and negative regulators of telomerase activity. A systematic screen for yeast mutants that affect telomere length maintenance in the yeast *Saccharomyces cerevisiae* revealed that mutations in any of ~500 genes affects telomere length. One of the genes that, when mutated, causes telomere elongation is *ELG1*, which encodes an unloader of PCNA, the processivity factor for replicative DNA polymerases. PCNA can undergo SUMOylation on two conserved residues, K164 and K127, or ubiquitination at lysine 164. These modifications have already been implicated in genome stability processes. We report that SUMOylated PCNA acts as a signal that positively regulates telomerase activity. We also uncovered physical interactions between Elg1 and the CST (Cdc13-Stn1-Ten) complex, and dissected the mechanism by which Elg1 and Stn1 negatively regulates telomere elongation, coordinated by SUMO. We present a model that provides mechanistic insights on how chromosomal replication and telomere elongation are coordinated.

INTRODUCTION

Telomeres cap the ends of linear eukaryotic chromosomes and safeguard them against genome instability. Telomeric DNA consists of G/C-rich DNA repeats, with the G-rich strand extending to form a 3' single-stranded overhang. Telomeres are primarily involved in counteracting the gradual shortening of the telomeric DNA caused by conventional DNA replication, thereby solving the "end-replication problem" (Bonnell, Pasquier, & Wellinger, 2021; Soudet, Jolivet, & Teixeira, 2014). Another important function of telomeres is to protect chromosome ends from improper recognition as a DNA double-strand break (DSB) [reviewed in (de Lange, 2018)].

An important, conserved protein complex involved in regulation of telomere length is the CST, which in *Saccharomyces cerevisiae* is composed of Cdc13, Stn1, and Ten1 (L. Y. Chen & Lingner, 2013; Ge et al., 2020; Lim et al., 2020; Puglisi, Bianchi, Lemmens, Damay, & Shore, 2008; Rice & Skordalakes, 2016). The CST has similarity to the ssDNA binding complex RPA and localizes specifically to the single-stranded telomeric DNA, where it is involved in chromosome end capping and telomere length regulation (Hughes, Weilbaecher, Walterscheid, & Lundblad, 2000; Lin & Zakian, 1996; Mersaoui & Wellinger, 2019). The CST facilitates telomerase-mediated telomere elongation (Evans & Lundblad, 1999, 2000; Nugent, Hughes, Lue, & Lundblad, 1996), participates in telomere processing during DNA replication (Soudet et al., 2014), and helps "capping" the telomere (Mersaoui & Wellinger, 2019). In late S-phase Cdc13 interacts with the Est1 subunit of telomerase and promotes telomerase activity (Chandra, Hughes, Nugent, & Lundblad, 2001; H. Chen et al., 2018). Stn1 binds Cdc13 in a region that overlaps with that bound by Est1. Recruitment of Stn1 evicts Est1 and prevents further elongation of the G-strand (Chandra et al., 2001; Wang et al., 2000). The amino terminus of Stn1 binds Ten1, whereas its C-terminus interacts with both Cdc13 and Pol12, a subunit of Polymerase Alpha (Grossi, Puglisi, Dmitriev, Lopes, & Shore, 2004; Petreaca et al., 2006; Puglisi et al., 2008). Thus, the CST coordinates leading strand elongation by telomerase with lagging strand DNA synthesis by the cell's replisome (Grossi et al., 2004).

Very often the integrity of the genome is compromised by internal and external sources of DNA damage. This vulnerability increases during S-phase, when the DNA is unpacked and copied. Elg1, the major subunit of a replication factor C-like complex (RLC), plays a central role in DNA replication and is critical for genome maintenance. In *Saccharomyces cerevisiae*, loss of the *ELG1* gene causes gross chromosomal rearrangements, chromosome losses, defective sister chromatid cohesion and recombination, increased sensitivity to DNA damaging agents and abnormal telomere length maintenance [reviewed in (Arbel, Choudhary, Tfilin, & Kupiec, 2021)]. The function of the Elg1 RLC is to unload the processivity factor PCNA from chromatin, in particular after Okazaki fragment processing and ligation in the lagging strand, and during DNA repair (Kubota, Katou, Nakato, Shirahige, & Donaldson, 2015; Kubota, Nishimura, Kanemaki, & Donaldson, 2013; Parnas et al., 2010; Shemesh et al., 2017). The mammalian ortholog of *ELG1* (ATAD5) participates in the Fanconi Anemia pathway (Kim et al., 2020; Yang et al., 2011) and, when mutated, leads to genome instability and cancer in mice and humans (Bell et al., 2011; Kuchenbaecker et al., 2015; Maleva Kostovska et al., 2016).

Post-translational modifications of PCNA (in particular ubiquitination and SUMOylation) orchestrate the activity of a large number of interacting proteins during DNA replication and repair. Many of these proteins interact through a PIP (PCNA Interacting Peptide) motif. PCNA can undergo SUMOylation on two conserved residues, K164 and K127, or ubiquitination at lysine 164. Although the Elg1 RLC shows affinity to SUMOylated PCNA, the complex is able to unload modified and unmodified versions of the clamp (Kang et al., 2019; Kubota et al., 2015; Kubota et al., 2013; Parnas et al., 2010; Shemesh et al., 2017; Shiomi & Nishitani, 2013). Thus SUMOylation may assist but is not essential for PCNA unloading. Retention and accumulation of PCNA on DNA is the major cause of genome instability in *elg1Δ* (Itzkovich, Choudhary, Arbel, & Kupiec, 2023; Johnson, Gali, Takahashi, & Kubota, 2016; Shemesh et al., 2017).

Previous studies found that telomeric proteins, and in particular the CST, can be regulated by SUMO modification (Hang, Liu, Cheung, Yang, & Zhao, 2011). Here we show that SUMOylation of PCNA plays a role in the regulation of telomere length, and uncover an interaction between the Elg1 RLC and the CST. We report that PCNA SUMOylation

92 positively regulates telomerase activity, and its unloading by the Elg1-RLC is essential for
93 normal telomere size regulation. The N-terminus of Elg1 interacts with Stn1 during late S
94 phase, and mediates the interaction between Stn1 and Cdc13 after PCNA has been
95 unloaded from the telomeres.

RESULTS

The long telomeres phenotype of *elg1Δ* strains requires PCNA SUMOylation

Previous results suggested that the Elg1 protein preferentially interacts with the SUMO
machinery, and with SUMOylated PCNA (Parnas, Amishay, Liefshitz, Zipin-Roitman, &
Kupiec, 2011; Parnas et al., 2010; Shemesh et al., 2017). To test whether PCNA
modifications play any role in the elongated telomere phenotype of *elg1Δ*, we combined
the *ELG1* deletion with mutations in the *POL30* gene, encoding PCNA. We mutated either
lysine 127, lysine 164, or both (hereafter referred as *pol30-K127R, pol30-K164R,* and
pol30-RR, respectively) in the *elg1Δ* background. **Figure 1** shows that the mutations in
POL30 have little effect by themselves. Importantly, whereas the single mutants did not
affect the long telomeres of *elg1Δ*, mutating both lysine residues of each PCNA subunit
completely abrogated the long telomere phenotype. Lysine 164 can be modified by both
SUMO and ubiquitin; mutating only this residue prevents ubiquitination, but not
SUMOylation; figure 1demonstrates that this mutant shows no effect. To ensure that
SUMO, and not ubiquitin, is the modification responsible for the long telomeres of *elg1Δ*,
we deleted *RAD18*, encoding the E3 ubiquitin ligase required to ubiquitinate PCNA
(Hoege, Pfander, Moldovan, Pyrowolakis, & Jentsch, 2002; Stelter & Ulrich, 2003). Figure
1 shows that lack of Rad18 has no effect on the elongated telomeres of *elg1Δ*. [Deletion of
the SUMO E3 ligases *SIZ1* and *SIZ2* (to prevent SUMOylation) by itself alters telomere
length (Hang et al., 2011) and thus they could not be used here]. We conclude that
SUMOylation of PCNA is required for the long telomeres phenotype of *elg1Δ* mutants.

These results suggest that SUMOylated PCNA may play a central role in determining
telomere length also in cells with wild type Elg1 activity. We thus next asked whether
artificially overexpressing modified and unmodified PCNA may mimic the long telomere

phenotype of *elg1Δ*. Since it is hard to force ubiquitination or SUMOylation of PCNA *in vivo*, we used covalent fusions between PCNA and either ubiquitin or SUMO. These have been shown in the past to mimic naturally modified PCNA molecules (Parker, Bielen, Dikic, & Ulrich, 2007; Takahashi, Wollscheid, Lowther, & Ulrich, 2020). We overexpressed either wild type Pol30, Pol30-RR (which cannot be modified) or Pol30-RR fused either to ubiquitin or to SUMO, in *pol30-RR* cells, unable to carry out modifications of the PCNA encoded by the genome. **Figure 1B** shows that telomeres become elongated upon overexpression of the wild type protein (which can be modified), or the Pol30-RR version fused to SUMO, but do not elongate upon overexpression of the unmodifiable Pol30-RR version, or the one fused to ubiquitin. These results, together with those of Figure 1A, show that accumulation of SUMOylated PCNA, and not unmodified or ubiquitinated PCNA, is both **necessary** and **sufficient** for telomere elongation in the presence or in the absence of Elg1.

Genetic and Physical interactions between Elg1 and Stn1

The elongated telomere phenotype of Elg1 was identified in systematic screens of the yeast knockout collection (Askree et al., 2004; Gatbonton et al., 2006), together with ~200 additional mutants of similar phenotypes. To identify the pathway of telomere length maintenance in which Elg1 participates, we systematically combined the *elg1Δ* mutant with all the other mutants with long telomeres. Most double mutants exhibited telomeres longer than each of the single mutants, indicating that they affect different pathways. One of the mutants showing clear epistasis (the double mutant was not longer than the single mutants) carried a deletion of *RRP8*, a gene involved in modification of the ribosomal RNA. Further analysis demonstrated that the knockout of *RRP8* exhibited the "neighboring gene effect", and in fact the deletion of *RRP8* caused a decrease in the expression of its neighboring gene, *STN1* (Ben-Shitrit et al., 2012). We confirmed the genetic interactions between *ELG1* and *STN1* alleles by combining *elg1Δ* with the *stn1-13* and *stn1-164* alleles (Grandin, Reed, & Charbonneau, 1997), which exhibit long telomeres. The telomeres of the double mutants are not longer than those of the single mutants (**Figure 2A**), indicating that Elg1 works in the same TLM pathway as Stn1.

Given the epistatic relations between the *ELG1* and *STN1* genes, we addressed, by co-Immunoprecipitation (IP) and Yeast-Two-Hybrid (YTH) assay, whether the proteins in the CST complex interact with Elg1. IP of epitope-tagged Elg1 pulled down Stn1 but not the other two members of the CST complex, Cdc13 or Ten1 (**Figure 2B**). For our YTH experiments, the Elg1 protein was divided into three main functional domains: an N-terminal domain (amino acids 1-234), a central AAA domain (amino acids 235-514) and a C-terminal domain (amino acids 515-791) (Itzkovich et al., 2023), **Figure 3A**. The AAA and C-terminal domains contain the conserved RFC boxes and are required for the interactions with the small RFC subunits, shared by all RFC-like domains. The N-terminal domain (hereafter referred to as NTD) is unique to Elg1 (Arbel et al., 2021). We thus concentrated on this domain, and used the C-terminus as a negative control, as it does not interact with any of the proteins tested here.

The NTD contains SUMO-interacting motifs (SIMs), which were previously shown to interact with SUMO and the SUMO machinery (Parnas et al., 2011; Parnas et al., 2010) (**Figure 3A**). We used the YTH technique to test potential interactions between Elg1 and the CST components, and their dependency on SUMO. The N-terminus of Elg1 interacted strongly with Cdc13 and Stn1, and weakly with Ten1 (**Figure 3B,C, Figure S1**). In contrast, no interaction was detected with Elg1's CTD. We could also detect interactions of Cdc13, Stn1 and Elg1's NTD with SUMO (Figure 3). These were used as positive controls in all our experiments. Since the results with Ten1 were much weaker than those with Cdc13 and Stn1, we concentrated on these two last proteins. Below, we dissect, using YTH in various genetic backgrounds, the interactions between Elg1, Stn1, Cdc13 and PCNA. The results are also summarized in **Table 1**.

Table1: Summary of all YTH interactions presented.

Strain	Plasmid	Interaction w/Stn1	Interaction w/Cdc13
wild type	Elg1-NTD	**YES**	**YES**
wild type	Elg1-sim	**NO**	**YES**
siz1Δ siz2Δ	Elg1-NTD	**NO**	**NO**
pol30-RR	Elg1-NTD	**YES**	**NO**

elg1Δ	Elg1-NTD	**YES**	**NO**
elg1-sim	Elg1-NTD	**YES**	**YES**
elg1-DD	Elg1-NTD	**YES**	**NO**
elg1-sim+DD	Elg1-NTD	**YES**	**NO**
elg1-WalkerA	Elg1-NTD	**YES**	**NO**
cdc13-snm	Elg1-NTD	**YES**	**YES**
wild type	Elg1-NTD	**-**	*snm:* **NO**

elg1-sim: mutation in the SUMO-interacting motif of Elg1; *elg1-DD: TT386/7DD*; *elg1-sim+DD*: combination of mutations in the SIM and in TT386/7; *elg1-WalkerA*: mutation that eliminates ATPase and unloading activity of Elg1. *cdc13-snm:* allele of Cdc13 that cannot be SUMOylated.

Interestingly, the interaction of Elg1's NTD with Stn1, but not with Cdc13, was abolished when the SIM motifs (amino acids I27A, I93K, I121A, and I122A) were mutated (**Figure 3B,C**). We confirmed these results by co-Immunoprecipitation (co-IP) (**Figure 2B**). These results suggest that the interaction with Stn1 is mediated by SUMOylation of a protein, whereas the interaction of Cdc13 with Elg1-NTD is independent of the SIM in the plasmid-borne copy of Elg1. We confirmed this result by deleting the genes encoding the SUMO-specific E3 enzymes, *SIZ1* and *SIZ2* in the genome (a double deletion is necessary, as many times each protein can compensate for the lack of the other). **Figure 3D** shows that indeed, deletion of the genes encoding these enzymes abolishes the interaction of Stn1 and Cdc13 with Elg1's NTD. We conclude that SUMO plays a role in mediating the Elg1-CST interactions. Since Elg1 preferentially interacts with SUMOylated PCNA (Parnas et al., 2010), and Stn1 can bind SUMO non-covalently in YTH (**Figure 3C**), we reckoned that SUMOylated PCNA may mediate the interaction between Stn1 and Elg1. We therefore tested whether abolishing PCNA's SUMOylation sites in the genome (*pol30-RR*) had an effect on the Stn1-Elg1 interaction. We found, however, that Elg1 and Stn1 could still interact in the *pol30-RR* strain (**Figure 3D**), despite the fact that PCNA cannot become SUMOylated. Consequently, a different SUMOylated target seems to mediate the interaction between Elg1 and Stn1, or else either one of the proteins is SUMOylated.

Despite numerous attempts, we were unsuccessful in observing SUMOylation of either

Stn1 or Elg1.

We also noted that, despite the fact that mutation in the plasmid-borne NTD's SIM had no

effect on the interaction with Cdc13 (**Figure 3B**), deletion of *SIZ1* and *SIZ2*, and mutation

in the SUMOylation sites in PCNA completely abolished the interactions between Elg1's

NTD and Cdc13 (**Figure 3D**). This implies that PCNA SUMOylation is necessary for the

interaction between Elg1 and Cdc13. We confirmed by co-IP that Cdc13 indeed physically

interacts with PCNA (**Figure 3E**). In summary, the interaction between Elg1 and Cdc13,

but not that of Elg1 with Stn1, is dependent on SUMOylation of PCNA.

Elg1's functional activity is essential for its interaction with Cdc13

When we repeated the YTH assays in a strain in which the *ELG1* gene is deleted from the

genome, we were surprised to see that now Cdc13 failed to interact with the plasmid-borne

Elg1-NTD (**Figure 4A**, compare to Figure 3B). In contrast, the interaction between Stn1

and Elg1-NTD was not affected by the deletion of the genomic *ELG1*.

Taken together, these results imply that whereas Stn1 interacts directly with the plasmid-

borne Elg1-NTD, via its SIM-mediated SUMO-binding, the interaction with Cdc13 require

both PCNA SUMOylation and **the genomic** *ELG1* gene. We can envision two formal

possibilities for the latter requirement: either 1) the interaction with Cdc13 requires parts

of the Elg1 protein not present in the plasmid-borne NTD (e.g., the AAA or CTD domain),

or 2) Elg1's **activity** (e.g., PCNA unloading) may be required to facilitate the physical

interaction of the NTD with Cdc13.

To distinguish between these two possibilities, we carried out YTH experiments with

strains carrying different alleles of *ELG1* in the genome. Three mutations were tested first:

elg1-sim (mutated in amino acids I27A, I93K, I121A I122A), which does not interact with

SUMO; *elg1-TT386/387DD*, hereafter referred to as *elg1-DD*, which affects the interface

Elg1-PCNA (Shemesh et al., 2017); and *elg1-sim+DD*, an allele that combines both

mutations. Whereas the *SIM* mutation has a very mild effect on the ability of cells to unload

PCNA, and displays no phenotypes, the *elg1-DD* is more impaired and has a stronger

phenotype, and the *elg1-sim+DD* allele has essentially a null phenotype, similar to that of

elg1Δ, but expressing the protein at normal levels (Itzkovich et al., 2023; Shemesh et al.,
2017).

We found that in strains carrying genomic *elg1-DD* and *elg1-sim+DD* alleles, Cdc13 failed
to interact with plasmid borne Elg1-NTD, whereas in the *elg1-sim* strain the interaction
was weak but clearly seen (**Figure 4B-D**). This observation indicates that the PCNA
unloading activity of Elg1 is essential to allow the interaction between Cdc13 and the
plasmid borne Elg1-NTD: this activity is high in *elg1-sim* mutants, intermediate in the
elg1-DD strain and is abolished by the *elg1-sim+DD* mutations. Note that the presence of
a functional SIM at the plasmid-borne NTD is sufficient to ensure a mild interaction with
Cdc13, but **only** if the genomic copy retains PCNA unloading activity (as in the *elg1-sim*
mutant). On the other hand, the presence of a functional SIM in the inactive *elg1-DD* allele
does not warrant an interaction of Elg1's NTD (which has SIM motifs) with Cdc13. We
confirmed this idea by analyzing the interaction between Cdc13 and Elg1's NTD in another
SIM-containing mutant strain devoid of PCNA unloading activity, *elg1-WalkerA*
(KK343/4AA), which lacks ATPase and PCNA unloading activity (Itzkovich et al., 2023).
This mutant expresses wild type levels of protein, and the mutant has essentially the same
phenotype of an *elg1Δ* allele. **Figure 4E** shows that no interaction between the plasmid-
borne Elg1 NTD and Cdc13 could be detected, despite the fact that functional SIMs are
present both at the NTD and in the genome-encoded Elg1-WalkerA protein. All the
mutants expressed the Elg1 protein at wild type levels (**Figure S2**).

We conclude that whereas the interaction with Stn1 is direct and mediated by the
SUMOylation of a protein (different from PCNA), the interaction between Elg1's NTD and
Cdc13 is indirect, and can only take place after PCNA has been SUMOylated and
unloaded. This suggests a model for the coordination of telomerase activity and DNA
replication: the presence of SUMOylated PCNA at telomeres may serve as a positive signal
for telomerase activity. Once Elg1 unloads PCNA, however, Stn1-Elg1 can bind Cdc13,
ending telomerase activity. In this context, it is possible to understand the results seen in
Figure 3D, showing that mutations in the SUMO E3 enzymes or in the sites of
SUMOylation of PCNA, both of which prevent PCNA SUMOylation, show the same effect

as the *sim+DD* or *walkerA* mutations in the genomic *ELG1* copy, which prevent PCNA
unloading activity.

SUMOylation of Cdc13 is needed for interaction with Elg1

Cdc13 undergoes SUMOylation, and this modification plays an essential role in the
negative regulation of telomere length (Hang et al., 2011). Thus, Cdc13 is a good candidate
for the target of SUMOylation required for the interaction between Elg1 and Stn1.
Accordingly, mutations of the Cdc13 SUMOylation sites (*cdc13-snm*, "SUMO no more")
weaken its interaction with Stn1 and lead to elongated telomere phenotype (Hang et al.,
2011). We therefore checked whether the *cdc13-snm* mutation affects also the interaction
with Elg1. **Figure 4F** shows that indeed preventing SUMOylation of Cdc13 abolishes the
interaction between this protein and Elg1. Note that the strain used expresses a wild type
genomic copy of *CDC13*. Thus, SUMOylation **of the plasmid-borne version of Cdc13**
(i.e., in *cis*) is required for these interactions in the YTH assay. No effect was observed
when the YTH plasmid carried a wild type copy of *CDC13*, and the genomic copy was the
cdc13-snm allele (**Figure S3A**). In summary, the N-terminus of Elg1 interacts with Cdc13
only if 1) Cdc13 becomes SUMOylated (**Figure 4F**), 2) PCNA becomes SUMOylated
(**Figure 3**D), and 3) PCNA is unloaded (**Figure 4A, C, D,E**).

The interaction of Elg1 with Stn1 takes place at telomeres only at late S-phase

The fact that the interactions of the Elg1 NTD with Cdc13 are dependent on PCNA
modification and unloading, whereas those with Stn1 are not, suggests that the interaction
with Cdc13 may be mediated by, or dependent on, the interaction of Elg1 with Stn1. To
further dissect this point, we divided the Stn1 protein into an NTD (first 281 amino acids)
and a CTD (amino acids 282-495). The later has been shown to be the region that interacts
with Cdc13 (Petreaca, Chiu, & Nugent, 2007). **Figure S3B** shows that the NTD of Elg1
also interacts with Stn1 via its C-terminal domain, both in *WT* and *elg1Δ* strains.

Having established Stn1 as the direct interactor of Elg1 in the CST complex, we next
examined the interaction between these proteins throughout the cell cycle phases. We
arrested a strain with tagged Elg1, Stn1 and DNA Polymerase Delta in G1 with alpha
factor. Cells were released into the cell cycle and samples were taken at intervals during

two cell cycles. This allowed us to map the timing of interaction between Elg1 and Stn1

proteins, and whether this interaction is codependent on the movement of replication fork.

Figure S4 shows that the total level of these proteins does not change throughout the cell

cycle. We immunoprecipitated Elg1, and measured the level of the other two proteins by

Western blot (**Figure 5A-C**). Pol3 was co-IPed with Elg1 strongly only during the S phase

(40-70 minutes, 120-140 minutes) whereas Stn1 was only detected in late S phase (60-70

minutes, 140-160 minutes) for both cell cycles and was barely detected in G1 (**Figure 5A-**

C). This pattern coincides with telomerase activity at telomeres, which is low in early to

mid S phase and peaks in late S-phase (Bianchi et al., 2008, Giraud et al., 2008, Taggart et

al., 2002).

To monitor the arrival of the replication fork to telomeres, we performed chromatin

immunoprecipitation at telomeres (Telo-ChIP) in synchronized cells. **Figure 5D** shows

that the timing of arrival of PCNA to the telomeres (60-70 minutes into the cell cycle)

coincides with the timing of the co-IP between Elg1 and Stn1 (**Figure 5A-C**). Thus, our

results are consistent with the idea that Elg1 moves with the replication fork, and interacts

with Stn1 at the end of S-phase, when telomeres are replicated and telomerase is activated.

The interaction between Cdc13 and Stn1 is dependent on Elg1

Our YTH and IP data suggest that unlike the relationship between Cdc13 and Elg1, the

interaction between Stn1 and Elg1 is direct (**Figures 2 and 3**). Since both Elg1 and Cdc13

bind Stn1 at its C-terminus [**Figure S3** and (Petreaca et al., 2007)], it is possible that the

absence of Elg1 may affect the interaction between the two CST members. We thus

measured the co-IP of Cdc13 and Stn1 in a wild type or an *elg1Δ* strain. Cells were cell-

cycle synchronized in S-phase, as the interaction between these two proteins was barely

seen when immunoprecipitated in an asynchronous culture. When Elg1 was absent, Stn1

exhibited a strong reduction in its co-precipitation with Cdc13 (**Figure 5E,F**). This

suggests that the interaction between Stn1 and Cdc13 is, at least partly, dependent on Elg1.

**Model: Elg1 negatively regulates the telomere length by forming an interaction with
the CST complex**

Taken together, our results suggest the following model (**Figure 6**):

Following firing of the origin proximal to the telomeres at late S-phase, Elg1 moves with the fork, unloading PCNA from the lagging strand as the fork progresses (Kubota et al., 2015; Kubota et al., 2013; Shemesh et al., 2017) (**Figure 6A**). SUMOylation of PCNA acts as a signal for the activation of telomerase (**Figures 1A,B and 6B**). Once Elg1 reaches the telomeres, it interacts with Stn1 (**Figure 5**). This interaction requires SUMOylation of Stn1 or of some other protein, which is not PCNA (**Figure 3D**) nor Cdc13 (**Figure S3A**) and could be Stn1 itself or another telomeric protein (Hang et al., 2011). The Elg1-Stn1 complex then unloads the SUMOylated PCNA at the telomeres, and Cdc13 becomes SUMOylated (**Figure 6C**), now interacting with Elg1-Stn1. SUMOylation of Cdc13 allows its interaction with Stn1, and presumably with Ten1 (Hang et al., 2011), creating the complete CST complex, and preventing further telomerase activity (**Figure 6D**). The fact that co-IP experiments detect a strong interaction with Stn1 but fail to detect an interaction with Cdc13 (**Figure 2B**) suggests that indeed the interaction Elg1-Cdc13 is very transient, and Elg1 leaves chromatin immediately, leaving the C-terminus of Stn1 free to interact with Cdc13. It is possible that a single "wave" of SUMOylation at the telomeres coordinates PCNA unloading with the interaction between Elg1 and Cdc13. Alternatively, unloading of PCNA by Elg1 may allow the recruitment of SUMO ligases to modify Cdc13. In support for this idea, YTH interactions between Stn1 or Cdc13 with SUMO are abolished in *siz1Δ siz2Δ* strains (**Figure 3**D).

DISCUSSION

We have presented evidence for the fact that two negative regulators of telomerase activity, Elg1 and Stn1, work in the same pathway to coordinate chromosomal DNA replication and telomere elongation. We present a model (**Figure 6**) in which the unloading of SUMOylated PCNA by Elg1 at telomeres facilitates the interaction between Stn1 and Cdc13 that negatively regulate telomerase activity.

This model is supported by several observations. First, the double mutants of Elg1 and Stn1 do not show increased telomere length phenotype in comparison to the single mutants, indicating that both proteins work in the same pathway that negatively regulates telomere

length (**Figure 2A**). In addition, the interaction between Elg1 and Stn1 seems direct and occurs exclusively during the late S-phase (**Figures 2B** and **5A-C**). This interaction is neither dependent on SUMOylation of PCNA (**Figure 3D**) nor on Cdc13 (**Figure S3A**). In contrast, the interaction between Cdc13 and Elg1 is dependent on both the SUMOylation of PCNA (**Figure 3D**) and of Cdc13 itself (**Figure 4F**). We suggest that Elg1 participates in the negative regulation of telomere length by unloading SUMOylated PCNA from the telomeres. This eliminates a positive signal for telomerase activity, and causes the exchange of Est1 by Stn1 as interactor of Cdc13, effectively terminating telomerase activity (Chandra et al., 2001; Gao, Cervantes, Mandell, Otero, & Lundblad, 2007; Ge et al., 2020; Hang et al., 2011; Puglisi et al., 2008). As expected from the model, the interaction Elg1-Cdc13 is transient, and cannot be detected by co-IP.

We present evidence for the fact that SUMOylated PCNA serves as a positive signal for telomere elongation (**Figure 1**). Preventing its unloading (by mutations in the *ELG1* gene) leads to elongated telomeres, and so does PCNA overexpression, provided the protein is able to be SUMOylated, or is fused to SUMO (thus mimicking SUMOylated PCNA) (**Figure 1**). *ELG1* is the only PCNA unloader in eukaryotes; however, it is not an essential gene, implying that there are alternative ways in which PCNA can get unloaded (either spontaneously or with the help of RFC) (Arbel et al., 2021). Once this step does take place, Stn1 can replace Est1 as Cdc13's partner, and stop the activity of telomerase, but at a longer telomere steady state equilibrium point. This explains both the long telomere phenotype of *elg1* mutants, and the fact that it is less severe than that of *stn1* alleles (**Figure 2**). The fact that *pol30-RR* mutants have stable, normal-sized telomeres (**Figure 1**), imply that despite lacking PCNA SUMOylation, they are able to solve the "end replication problem". Our results support the notion (Maicher et al., 2017) that there are two alternative modes of telomerase: one carrying "basal activity", with minimal genetic requirements, able to maintain most telomeres at normal size, and another, "sustained activity", required for increased elongating activity (Maicher et al., 2017). Previous results have shown that the first mode does not require Rnr1, the large subunit of the ribonucleotide reductase enzyme, whereas the second depends on it (Maicher et al., 2017). Similarly, SUMOylated PCNA is only needed for the second mode of telomerase.

Many proteins in the cell become SUMOylated at one point or another. This modification is usually transient, due to the activity of de-SUMOylases (which eliminate the modification) and STUbLs (SUMO-dependent ubiquitin ligases, which usually send the protein to degradation). It has been proposed that SUMOylation often works by targeting many physically interacting proteins together. This may be executed by the local recruitment of SUMO ligases to preassembled, co-localized protein groups. SUMOylation thus may increase their interactions, acting to increase their local concentration or effective interaction (Jentsch & Psakhye, 2013). SUMOylation of telomeric proteins seems to be one of those cases (Garg et al., 2014; Hang et al., 2011; Matmati et al., 2018; Miyagawa et al., 2014). In our model, PCNA, Cdc13 and possibly Stn1 undergo SUMOylation to ensure the proper sequence of events in the coordination between telomerase activity and chromosomal replication. Cdc13 binds Pol12, a subunit of Polymerase Alpha (Grossi et al., 2004; Petreaca et al., 2006; Puglisi et al., 2008), allowing the synthesis of the lagging strand on the newly synthesized ssDNA. Thus, the arrival of the replisome to a short telomere at the end of S-phase brings PCNA, whose SUMOylation can elicit both telomerase activity, and its subsequent repression when PCNA is unloaded by Elg1. This in turn primes synthesis of the complementary strand, thus coordinating telomere elongation with chromosomal replication.

MATERIALS AND METHODS

Yeast strains, plasmids and Media: All yeast strains used are described in **Table 2**. All plasmids are described in **Table 3.** Yeast extract-peptone-dextrose (YPD) medium was prepared with a ready-to-use mixture (Formedium). Synthetic complete (SC) minimal medium was prepared with 2% dextrose (Formedium), yeast nitrogen base without amino acids (Difco), and all necessary amino acids. 2% agar (Difco) was added for solid medium. Standard yeast genetic procedures were used to create single and double mutants.

Southern Teloblots: For each genotype, at least 3 independently created strains were tested after 10 consecutive passages on YPD plates. Genomic DNA was isolated by pelleting 15 ml saturated overnight yeast culture grown in YPD, then beating the cells using

0.5 mm glass beads in 500 µl lysis buffer and 500 µl phenol chloroform. Cells were spun

down for 10 min at 14 000 rpm, and ∼500 µl of the top clear solution, containing the

genomic DNA, was carefully taken out. DNA was precipitated using 1000 µl of 95%

ethanol and then pelleted. The supernatant was discarded, and 500 µl of 70% ethanol was

added. Pellets were left to dry before resuspension in 300 µl TE with RNaseA for 10

minutes at 65°C. The DNA was precipitated by adding 130 µl of 7.5 M ammonium acetate

(pH 7) and 1 ml of 95% ethanol at −20°C, spun, and resuspended in 50 µl of TE. About 10

µl of genomic DNA was digested using XhoI overnight at 37°C. Digested gDNA and

ladder were loaded onto a 0.8% agarose gel and electrophoresed overnight at ∼67V in 1X

TBE. The gel was denatured for 30 min in a rocking shaker (1.5 M NaCl, 0.5 M NaOH)

and was neutralized for 15 min (1.5 M NaCl, 0.5M Tris, pH 7.0). The DNA on the gel was

transferred onto a hybond nylon membrane (GE Healthcare GERPN303B) with 10X SSC

(1.5 M NaCl 0.17 M NaCitrate, dihydrate) and UV cross-linked before blocking in Church

buffer (0.5 M Na_2HPO_4, pH7.2, 7% SDS, 1 mM EDTA, 1% BSA) for ∼2 h at 65°C. ^{32}P

radiolabeled PCR fragments were added onto the membrane and left to incubate overnight.

The membrane was washed 3–4 times for 15 min each, with 1X SSC 0.1%SDS buffer

before drying and exposing it to X-ray film. The teloblots blots were exposed for 1-8 days

at -70°C before the film was developed.

Yeast Two-Hybrid: The yeast two-hybrid assays were performed using PJ69-4A

(James, Halladay, & Craig, 1996) strains cotransformed with a *LEU2*-marked plasmid

containing genes fused to the *GAL4* activating domain (pACT or pGAD424) and a *URA3*-

marked plasmid containing genes fused to the *GAL4* DNA binding domain (pGBU9).

Strains containing the test plasmids were grown for 24 h at 30°C in SC-Ura-Leu liquid

medium, and were plated on the reporter maker SC-His medium.

Protein extraction and immunoprecipitation (IP) assays: Cells were grown

to mid-logarithmic phase, washed once with water, and resuspended in lysis buffer (20

mM Tris-HCl, pH 7.5, 0.5 mM EGTA, 0.5 mM EDTA, 1 mM DTT, 125 mM potassium

acetate, 12.5% glycerol, 0.1% Triton X-100, protease inhibitor mixture, and 1

mM phenylmethylsulfonyl fluoride). Cells were broken for 45 min with glass beads,

centrifuged for 10 min at 10,000 × g and the supernatant was collected. 20-30 µg of total

protein extract was resolved on SDS-PAGE using 10% acrylamide gels. For immunoprecipitation, 1,000 µg of proteins were prepared and pre-cleared with 20 µl of protein A-Sepharose and protein G-Sepharose beads mixture (GE Healthcare). 2 µl of (HA or MYC) antibodies were added to the cleared extract and incubated overnight at 4 ^{0}C. The beads were washed once with lysis buffer, once with lysis buffer containing 0.5 M NaCl, and twice with buffer A (50 mM Tris-HCl, pH 7.5, 0.1 mM EGTA, 0.1% β-mercaptoethanol). The resulting immunoprecipitates were used for *in vitro* kinase assays.

Western blotting: Cells were collected by centrifugation, re-suspended in 600 ul of phosphate-buffered saline with 1% Triton X-100 (PBST), supplemented with a protease inhibitor cocktail (Roche) and subjected to mechanical rupture using glass beads. The cell debris were removed by centrifugation, and the supernatants were applied onto 0.1 M dithiothreitol, and incubated at 80∘C for 10 min before sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) (Resolving gel: 30% Acrylamide, 1.5 M Tris-HCl pH8.8, 10% SDS (pH 7.2), 9.7 ml H2O, 100 ul 10% APS and 10 ul TEMED. Stacking gel: 30% Bis/Acrylamide, 1 M Tris-HCl pH 6.8, 10% SDS (pH 7.2), 5.5 ml H2O, 800 ul 10% APS and 8 ul TEMED). The samples were run with SDS-PAGE buffer at 100 V until the samples have passed the stacking gel and then at 160 V until the samples have been fully separated. Transfer to nitrocellulose was done in transfer buffer (200 ml Methanol, 3.03 gr Tris Base, 14.4 gr Glycine) at 500 mAmp and verified by staining with Ponceau-S dye. The blot was blocked with Milk for at least 60 min at room temperature. Primary antibody was added for 12 h at 4∘C. The blot was washed 3 × 5 min with TBST (Tris-Buffered Saline Tween-20) and secondary antibody was added for 1 h. The blot was washed 3 × 5 min with TBST and subjected to electro-chemi-luminiscence (ECL).

Cell cycle synchronization: To assay Elg1 and Stn1 interaction during the cell cycle, the triple tagged strain was grown in YPD to OD$_{600}$ 0.6-0.8, followed by the addition of 500ng/ml α factor and was grown until ∼90% of the cells appeared unbudded or exhibit Shmoo formation (for ∼2 hrs). The α-factor was removed by centrifugation and washing cells 2-3 times with warm YPD. Cells were released into YPD at OD$_{600}$ 0.6-0.8 with addition of pronase at a final concentration of 0.1mg/ml. Cells were collected for both FACS and Western blot analysis at different time points after release. Cells for FACS

analysis were fixed in 70% ethanol, digested with RNase overnight, washed again and stained with propidium iodide (15 µg/mL), and analyzed by flow cytometry.

Flow cytometry: 200 µl of a logarithmic cell culture (0.6 OD600) were harvested, resuspended in 60 µl of 50 mM Tris pH 7.5 and 140 µl of ethanol was added; cells were then kept overnight at 4 °C. Fixed cells were centrifuged and washed once in 200 µl of 50 mM Tris pH 7.5 bufer and resuspended in 100 µl RNAse (0.2 mg/ml in 50 mM Tris pH 7.5) for 2 h at 37 °C. In addition, proteinase-K (0.2 mg/ml in 50 mM Tris pH 7.5) was added to each tube and cells were incubated for 60 additional minutes at 50 °C. 20 µl of the sample was taken into a new tube and a 180 µl of 18 µg/ml propidium iodide 50 mM Tris pH 7.5 was added. The samples were kept in the dark at 4 °C overnight, sonicated twice at low setting (20% power) for 3–5 s and stored in the dark at 4 °C. The fow cytometry MACSQuant system was used for reading. Results were analyzed using either the Flowing Software or the FlowJo program.

ACKNOWLEDGEMENTS

We thank David Shore, Hele Ulrich, Connie Nugent, Michel Charboneau and Vicky Lundblad for strains and plasmids. We thank all present and past members of the Kupiec lab for support and ideas. This research was supported by grants from the Israel Science Foundation, the Israel Cancer Research Fund, and the Recanati Fund.

References

Arbel, M., Choudhary, K., Tfilin, O., & Kupiec, M. (2021). PCNA Loaders and Unloaders-One Ring That Rules Them All. *Genes (Basel), 12*(11). doi: 10.3390/genes12111812

Askree, S. H., Yehuda, T., Smolikov, S., Gurevich, R., Hawk, J., Coker, C., . . . McEachern, M. J. (2004). A genome-wide screen for Saccharomyces cerevisiae deletion mutants that affect telomere length. *Proc Natl Acad Sci U S A, 101*(23), 8658-8663. doi: 10.1073/pnas.0401263101

Bell, D. W., Sikdar, N., Lee, K. Y., Price, J. C., Chatterjee, R., Park, H. D., . . . Myung, K. (2011). Predisposition to cancer caused by genetic and functional defects of mammalian Atad5. *PLoS Genet, 7*(8), e1002245. doi: 10.1371/journal.pgen.1002245

Ben-Shitrit, T., Yosef, N., Shemesh, K., Sharan, R., Ruppin, E., & Kupiec, M. (2012). Systematic identification of gene annotation errors in the widely used yeast mutation collections. *Nat Methods, 9*(4), 373-378. doi: 10.1038/nmeth.1890

Bonnell, E., Pasquier, E., & Wellinger, R. J. (2021). Telomere Replication: Solving Multiple End
Replication Problems. *Front Cell Dev Biol, 9*, 668171. doi: 10.3389/fcell.2021.668171
Chandra, A., Hughes, T. R., Nugent, C. I., & Lundblad, V. (2001). Cdc13 both positively and
negatively regulates telomere replication. *Genes Dev, 15*(4), 404-414. doi:
10.1101/gad.861001
Chen, H., Xue, J., Churikov, D., Hass, E. P., Shi, S., Lemon, L. D., . . . Lei, M. (2018). Structural Insights
into Yeast Telomerase Recruitment to Telomeres. *Cell, 172*(1-2), 331-343 e313. doi:
10.1016/j.cell.2017.12.008
Chen, L. Y., & Lingner, J. (2013). CST for the grand finale of telomere replication. *Nucleus, 4*(4),
277-282. doi: 10.4161/nucl.25701
de Lange, T. (2018). Shelterin-Mediated Telomere Protection. *Annu Rev Genet, 52*, 223-247. doi:
10.1146/annurev-genet-032918-021921
Evans, S. K., & Lundblad, V. (1999). Est1 and Cdc13 as comediators of telomerase access. *Science,*
286(5437), 117-120. doi: 10.1126/science.286.5437.117
Evans, S. K., & Lundblad, V. (2000). Positive and negative regulation of telomerase access to the
telomere. *J Cell Sci, 113 Pt 19*, 3357-3364. doi: 10.1242/jcs.113.19.3357
Gao, H., Cervantes, R. B., Mandell, E. K., Otero, J. H., & Lundblad, V. (2007). RPA-like proteins
mediate yeast telomere function. *Nat Struct Mol Biol, 14*(3), 208-214. doi:
10.1038/nsmb1205
Garg, M., Gurung, R. L., Mansoubi, S., Ahmed, J. O., Dave, A., Watts, F. Z., & Bianchi, A. (2014).
Tpz1TPP1 SUMOylation reveals evolutionary conservation of SUMO-dependent Stn1
telomere association. *EMBO Rep, 15*(8), 871-877. doi: 10.15252/embr.201438919
Gatbonton, T., Imbesi, M., Nelson, M., Akey, J. M., Ruderfer, D. M., Kruglyak, L., . . . Bedalov, A.
(2006). Telomere length as a quantitative trait: genome-wide survey and genetic mapping
of telomere length-control genes in yeast. *PLoS Genet, 2*(3), e35. doi:
10.1371/journal.pgen.0020035
Ge, Y., Wu, Z., Chen, H., Zhong, Q., Shi, S., Li, G., . . . Lei, M. (2020). Structural insights into telomere
protection and homeostasis regulation by yeast CST complex. *Nat Struct Mol Biol, 27*(8),
752-762. doi: 10.1038/s41594-020-0459-8
Grandin, N., Reed, S. I., & Charbonneau, M. (1997). Stn1, a new Saccharomyces cerevisiae protein,
is implicated in telomere size regulation in association with Cdc13. *Genes Dev, 11*(4), 512-
527. doi: 10.1101/gad.11.4.512
Grossi, S., Puglisi, A., Dmitriev, P. V., Lopes, M., & Shore, D. (2004). Pol12, the B subunit of DNA
polymerase alpha, functions in both telomere capping and length regulation. *Genes Dev,*
18(9), 992-1006. doi: 10.1101/gad.300004
Hang, L. E., Liu, X., Cheung, I., Yang, Y., & Zhao, X. (2011). SUMOylation regulates telomere length
homeostasis by targeting Cdc13. *Nat Struct Mol Biol, 18*(8), 920-926. doi:
10.1038/nsmb.2100
Hoege, C., Pfander, B., Moldovan, G. L., Pyrowolakis, G., & Jentsch, S. (2002). RAD6-dependent
DNA repair is linked to modification of PCNA by ubiquitin and SUMO. *Nature, 419*(6903),
135-141.
Hughes, T. R., Weilbaecher, R. G., Walterscheid, M., & Lundblad, V. (2000). Identification of the
single-strand telomeric DNA binding domain of the Saccharomyces cerevisiae Cdc13
protein. *Proc Natl Acad Sci U S A, 97*(12), 6457-6462. doi: 10.1073/pnas.97.12.6457
Itzkovich, Z., Choudhary, K., Arbel, M., & Kupiec, M. (2023). Effects of Defective Unloading and
Recycling of PCNA Revealed by the Analysis of ELG1 Mutants. *Int J Mol Sci, 24*(2). doi:
10.3390/ijms24021568

James, P., Halladay, J., & Craig, E. A. (1996). Genomic libraries and a host strain designed for highly
efficient two-hybrid selection in yeast. *Genetics, 144*(4), 1425-1436.
Jentsch, S., & Psakhye, I. (2013). Control of nuclear activities by substrate-selective and protein-
group SUMOylation. *Annu Rev Genet, 47*, 167-186. doi: 10.1146/annurev-genet-111212-
133453
Johnson, C., Gali, V. K., Takahashi, T. S., & Kubota, T. (2016). PCNA Retention on DNA into G2/M
Phase Causes Genome Instability in Cells Lacking Elg1. *Cell Rep, 16*(3), 684-695. doi:
10.1016/j.celrep.2016.06.030
Kang, M. S., Ryu, E., Lee, S. W., Park, J., Ha, N. Y., Ra, J. S., . . . Myung, K. (2019). Regulation of
PCNA cycling on replicating DNA by RFC and RFC-like complexes. *Nat Commun, 10*(1),
2420. doi: 10.1038/s41467-019-10376-w
Kim, S. J., Wie, M., Park, S. H., Kim, T. M., Park, J. H., Kim, S., . . . Lee, K. Y. (2020). ATAD5 suppresses
centrosome over-duplication by regulating UAF1 and ID1. *Cell Cycle, 19*(15), 1952-1968.
doi: 10.1080/15384101.2020.1785724
Kubota, T., Katou, Y., Nakato, R., Shirahige, K., & Donaldson, A. D. (2015). Replication-Coupled
PCNA Unloading by the Elg1 Complex Occurs Genome-wide and Requires Okazaki
Fragment Ligation. *Cell Rep, 12*(5), 774-787. doi: 10.1016/j.celrep.2015.06.066
Kubota, T., Nishimura, K., Kanemaki, M. T., & Donaldson, A. D. (2013). The Elg1 replication factor
C-like complex functions in PCNA unloading during DNA replication. *Mol Cell, 50*(2), 273-
280. doi: 10.1016/j.molcel.2013.02.012
Kuchenbaecker, K. B., Ramus, S. J., Tyrer, J., Lee, A., Shen, H. C., Beesley, J., . . . Brca. (2015).
Identification of six new susceptibility loci for invasive epithelial ovarian cancer. *Nat
Genet, 47*(2), 164-171. doi: 10.1038/ng.3185
Lim, C. J., Barbour, A. T., Zaug, A. J., Goodrich, K. J., McKay, A. E., Wuttke, D. S., & Cech, T. R. (2020).
The structure of human CST reveals a decameric assembly bound to telomeric DNA.
Science, 368(6495), 1081-1085. doi: 10.1126/science.aaz9649
Lin, J. J., & Zakian, V. A. (1996). The Saccharomyces CDC13 protein is a single-strand TG1-3
telomeric DNA-binding protein in vitro that affects telomere behavior in vivo. *Proc Natl
Acad Sci U S A, 93*(24), 13760-13765. doi: 10.1073/pnas.93.24.13760
Maicher, A., Gazy, I., Sharma, S., Marjavaara, L., Grinberg, G., Shemesh, K., . . . Kupiec, M. (2017).
Rnr1, but not Rnr3, facilitates the sustained telomerase-dependent elongation of
telomeres. *PLoS Genet, 13*(10), e1007082. doi: 10.1371/journal.pgen.1007082
Maleva Kostovska, I., Wang, J., Bogdanova, N., Schurmann, P., Bhuju, S., Geffers, R., . . . Dork, T.
(2016). Rare ATAD5 missense variants in breast and ovarian cancer patients. *Cancer Lett,
376*(1), 173-177. doi: 10.1016/j.canlet.2016.03.048
Matmati, S., Vaurs, M., Escandell, J. M., Maestroni, L., Nakamura, T. M., Ferreira, M. G., . . . Coulon,
S. (2018). The fission yeast Stn1-Ten1 complex limits telomerase activity via its SUMO-
interacting motif and promotes telomeres replication. *Sci Adv, 4*(5), eaar2740. doi:
10.1126/sciadv.aar2740
Mersaoui, S. Y., & Wellinger, R. J. (2019). Fine tuning the level of the Cdc13 telomere-capping
protein for maximal chromosome stability performance. *Curr Genet, 65*(1), 109-118. doi:
10.1007/s00294-018-0871-3
Miyagawa, K., Low, R. S., Santosa, V., Tsuji, H., Moser, B. A., Fujisawa, S., . . . Tanaka, K. (2014).
SUMOylation regulates telomere length by targeting the shelterin subunit Tpz1(Tpp1) to
modulate shelterin-Stn1 interaction in fission yeast. *Proc Natl Acad Sci U S A, 111*(16),
5950-5955. doi: 10.1073/pnas.1401359111

Nugent, C. I., Hughes, T. R., Lue, N. F., & Lundblad, V. (1996). Cdc13p: a single-strand telomeric
586   DNA-binding protein with a dual role in yeast telomere maintenance. *Science, 274*(5285),
587   249-252. doi: 10.1126/science.274.5285.249

588   Parker, J. L., Bielen, A. B., Dikic, I., & Ulrich, H. D. (2007). Contributions of ubiquitin- and PCNA-
589   binding domains to the activity of Polymerase eta in Saccharomyces cerevisiae. *Nucleic*
590   *Acids Res, 35*(3), 881-889. doi: 10.1093/nar/gkl1102

591   Parnas, O., Amishay, R., Liefshitz, B., Zipin-Roitman, A., & Kupiec, M. (2011). Elg1, the major
592   subunit of an alternative RFC complex, interacts with SUMO-processing proteins. *Cell*
593   *Cycle, 10*(17), 2894-2903. doi: 10.4161/cc.10.17.16778

594   Parnas, O., Zipin-Roitman, A., Pfander, B., Liefshitz, B., Mazor, Y., Ben-Aroya, S., . . . Kupiec, M.
595   (2010). Elg1, an alternative subunit of the RFC clamp loader, preferentially interacts with
596   SUMOylated PCNA. *Embo J, 29*(15), 2611-2622. doi: 10.1038/emboj.2010.128

597   Petreaca, R. C., Chiu, H. C., Eckelhoefer, H. A., Chuang, C., Xu, L., & Nugent, C. I. (2006).
598   Chromosome end protection plasticity revealed by Stn1p and Ten1p bypass of Cdc13p.
599   *Nat Cell Biol, 8*(7), 748-755. doi: 10.1038/ncb1430

600   Petreaca, R. C., Chiu, H. C., & Nugent, C. I. (2007). The role of Stn1p in Saccharomyces cerevisiae
601   telomere capping can be separated from its interaction with Cdc13p. *Genetics, 177*(3),
602   1459-1474. doi: 10.1534/genetics.107.078840

603   Puglisi, A., Bianchi, A., Lemmens, L., Damay, P., & Shore, D. (2008). Distinct roles for yeast Stn1 in
604   telomere capping and telomerase inhibition. *Embo J, 27*(17), 2328-2339. doi:
605   10.1038/emboj.2008.158

606   Rice, C., & Skordalakes, E. (2016). Structure and function of the telomeric CST complex. *Comput*
607   *Struct Biotechnol J, 14*, 161-167. doi: 10.1016/j.csbj.2016.04.002

608   Shemesh, K., Sebesta, M., Pacesa, M., Sau, S., Bronstein, A., Parnas, O., . . . Kupiec, M. (2017). A
609   structure-function analysis of the yeast Elg1 protein reveals the importance of PCNA
610   unloading in genome stability maintenance. *Nucleic Acids Res, 45*(6), 3189-3203. doi:
611   10.1093/nar/gkw1348

612   Shiomi, Y., & Nishitani, H. (2013). Alternative replication factor C protein, Elg1, maintains
613   chromosome stability by regulating PCNA levels on chromatin. *Genes Cells, 18*(11), 946-
614   959. doi: 10.1111/gtc.12087

615   Soudet, J., Jolivet, P., & Teixeira, M. T. (2014). Elucidation of the DNA end-replication problem in
616   Saccharomyces cerevisiae. *Mol Cell, 53*(6), 954-964. doi: 10.1016/j.molcel.2014.02.030

617   Stelter, P., & Ulrich, H. D. (2003). Control of spontaneous and damage-induced mutagenesis by
618   SUMO and ubiquitin conjugation. *Nature, 425*(6954), 188-191. doi: 10.1038/nature01965

619   Takahashi, T. S., Wollscheid, H. P., Lowther, J., & Ulrich, H. D. (2020). Effects of chain length and
620   geometry on the activation of DNA damage bypass by polyubiquitylated PCNA. *Nucleic*
621   *Acids Res, 48*(6), 3042-3052. doi: 10.1093/nar/gkaa053

622   Wang, M. J., Lin, Y. C., Pang, T. L., Lee, J. M., Chou, C. C., & Lin, J. J. (2000). Telomere-binding and
623   Stn1p-interacting activities are required for the essential function of Saccharomyces
624   cerevisiae Cdc13p. *Nucleic Acids Res, 28*(23), 4733-4741. doi: 10.1093/nar/28.23.4733

625   Yang, K., Moldovan, G. L., Vinciguerra, P., Murai, J., Takeda, S., & D'Andrea, A. D. (2011).
626   Regulation of the Fanconi anemia pathway by a SUMO-like delivery network. *Genes Dev,*
627   *25*(17), 1847-1858. doi: 10.1101/gad.17020911

630 **Table 2: Yeast strains used in this study.**

Strain Number	Name	Genotype
13237	BY4741	MATa his3del, leu2del, met15del, ura3del
12480	BY elg1Δ::HygMX	MATa his3del, leu2del, met15del, ura3del, elg1::HYG
14421	BY pol30-K127R	MATa his3del, leu2del, met15del, ura3del, pol30-K127R::LEU2
14425	BY pol30-K164R	MATa his3del, leu2del, met15del, ura3del, pol30-K164R::LEU2
14423	BY pol30-KK127,164RR	MATa his3del, leu2del, met15del, ura3del, pol30-RR::LEU2
14426	BY pol30-K127R elg1Δ	MATa his3del, leu2del, met15del, ura3del, pol30-K127R::LEU2, elg1::HYG
14430	BY pol30-K164R elg1Δ	MATa his3del, leu2del, met15del, ura3del, pol30-K164R::LEU2, elg1::HYG
14428	BY pol30-KK127,164RR elg1Δ	MATa his3del, leu2del, met15del, ura3del,pol30-RR::LEU2,elg1::HYG
14398	BY rad18Δ:: KanMX	MATa his3del, leu2del, met15del, ura3del, rad18::KanMX.
14401	BY rad18D::KanMX elg1D::HygMX	MATa his3del, leu2del, met15del, ura3del, rad18::KanMX, elg1::HYG
19606	BY pol30-KK127,164RR Elg1-HA	MATa his3del, leu2del, met15del, ura3del ELG1-HA-NAT pol30-KK127,164RR:HIS3
20622	BY bar1 CDC13-HA ELG1-HA	MATa his3del, leu2del, met15del, ura3del, bar1::LEU2, CDC13:3HA:HISMX, ELG1-HA-KanMX
18418	BY STN1-Myc Elg1-HA	MATa his3del, leu2del, met15del, ura3del stn1::HYG CEN LEU2 STN1-(G)9-(myc)7 ELG1-HA-KanMx
18790	BY bar1 STN1-Myc ELG1-HA POL3-FLAG	MATa his3del, leu2del, met15del, ura3del bar1::NatMX stn1::HYG CEN LEU2 STN1-(G)9-(myc)7 ELG1-HA-KanMx POL3-FLAG-URA3
19552	BY bar1 CDC13-HA POL30-FLAG	MATa his3del, leu2del, met15del ura3del, bar1:: NatMX CDC13-3HA::HISMX, POL30-FLAG::KanMX,

20625	BY CDC13-HA STN1-Myc	MATa his3del, leu2del, met15del, ura3del, CDC13-3HA::HISMX, stn1::HYG CEN LEU2 STN1-(G)9-(myc)7,
20626	BY CDC13-HA STN1-Myc elg1D	MATa his3del, leu2del, met15del, ura3del,CDC13-3HA::HISMX, stn1::HYG ,CEN LEU2 STN1-(G)9-(myc)7,elg1::KanMx
20623	BY TEN1-FLAG	MATa his3del, leu2del, met15del, ura3del trp1del Lys2del can1:: STE2pr-Sp_HIS5 ten1::KanMX, Elg1-HA-NAT CEN URA3 TEN1-(G)8-(FLAG)3
17611	W303-1a	MATa leu2-3, 112 ura3-1 his3-11,15, trp1-1, ade2-1, can1- 100
9842	W303 elg1Δ::HygMX	MATa leu2-3, 112 ura3-1 his3-11,15, trp1-1, ade2-1, can1- 100, elg1::KanMx
9551	W303 stn1-13	MATa leu2-3, 112 ura3-1 his3-11,15, trp1-1, ade2-1, can1- 100, stn1-13
9848	W303 elg1Δ::HygMX stn1-13	MATa leu2-3, 112 ura3-1 his3-11,15, trp1-1, ade2-1, can1- 100 elg1::KanMx, stn1-13
12357	W303 stn1-164	MATa leu2-3, 112 ura3-1 his3-11,15, trp1-1, ade2-1, can1- 100, stn1Δ::Hyg + pRS-stn1-L164A (pVL3571)
12358	W303 stn1-164 elg1Δ: KanMX	MATa leu2-3, 112 ura3-1 his3-11,15, trp1-1, ade2-1, can1- 100, elg1::KanMx, stn1Δ::Hyg+pRS-stn1-L164A (pVL3571)
12062	PJ69-4	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ
15017	PJ elg1D	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ elg1::HYG
19774	PJ siz1 siz2	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ siz1::KanMX siz2::HYG
11069	PJ pol30-RR	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ POL30-RR :: leu2 :: KANMX
20624	PJ STN1-Myc	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3

		met2::GAL7-lacZ elg1-sim::KanMx, stn1::HYG,CENLEU2 STN1-(G)9-(myc)7
19916	PJ elg1-sim-Myc	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ elg1-SIM-MYC-KANMX
18798	PJ elg1-DDMyc	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ elg1-DD-MYC-KANMX
19917	PJ elg1-DD+sim-Myc	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ elg1-DDsim-MYC-KANMX
19915	PJ elg1-Walker A	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ, elg1 KK343/4DD:MYC:KanMX
19486	PJ cdc13-snm	MAT@ trp1-901 leu2-3,112 ura3-52 his3-200 gal4del gal80del GAL2-ADE2 LYS2:: GAL1-HIS3 met2::GAL7-lacZ cdc13-snm

633 **Table 3: Plasmids used in this study**

Strain Number	Genotype	Source or Reference
4239	pGad424 (LEU2)	Kupiec lab
4238	pGad424-POL30	Helle Ulrich lab
4237	pGad424-pol30-k127r,k164r	Helle Ulrich lab
4236	pGad424-pol30-k127r,k164r-SMT3	Helle Ulrich lab
4235	pGAD424-pol30-k127r,k164r-UBI	Helle Ulrich lab
4147	CEN LEU2 STN1-(G)9-(myc)7	Victoria Lundblad lab
4144	CEN URA3 TEN1-(G)8-(FLAG)3	Victoria Lundblad lab
2201	pCN181 pACT2-STN1 (LEU2)	Constance Nugent lab
2205	pVL855 pACT2-CDC13	Constance Nugent lab
2168	pACT2 - TEN1	Michel Charbonneau lab
1775	pGBU9 (URA3)	Kupiec lab
1973	pGBU9-ELG1-NTD(1-234)	This Study
3301	pGBU9-ELG1-CTD(541-791)	This Study
2260	pGBU9-ELG1-NTDsim(1-234)	This Study
2241	pGBU9-SMT3	This Study
2419	pGAD424-Stn1-Nt(1-282)	David Shore lab
2420	pGAD424-Stn1-Ct(282-494)	David Shore lab
2169	pGAD424-Stn1-13	Michel Charbonneau lab
2418	pGAD424-Stn1-63	David Shore lab
4287	pACT-cdc13-snm	This Study

A

wild type | pol30-127R | elg1Δ | pol30-127R elg1Δ | wild type | pol30-164R | elg1Δ | pol30-164R elg1Δ | wild type | pol30-RR | pol30-RR elg1Δ | elg1Δ | wild type | rad18Δ | rad18Δ elg1Δ

-2044bp

-779bp

B

Empty Vector | pPol30 | pPol30-RR | pPol30-RR-SUMO | pPol30-RR-Ubi

-2044bp

-779bp

Figure1

Figure 1: SUMOylated PCNA regulates telomere length. A. Southern blot (Teloblot) showing that lack of SUMOylation of PCNA prevents telomere elongation. Independently-created colonies were passaged ten times, its DNA extracted, digested with *Xho*I and run in an agarose gel. The DNA was then transferred to a nitrocellulose membrane, which was incubated with a radioactive probe that detects telomeres, and a size marker. **B.** Teloblot showing that overexpression of wild type PCNA or Pol30-RR-SUMO fusion, but not Pol30-ubiquitin fusion or Pol30-RR causes telomere elongation.

A

wild type | elg1Δ | stn1-13 | stn1-13 elg1Δ | stn1-164 | stn1-164 elg1Δ

-2044 bp

-779 bp

B

Cdc13-HA: − − + +
Elg1-MYC: − + − +

IP HA
Cdc13
Elg1

INPUT
Cdc13
Elg1
Act1

Strain: wild type | elg1-sim

Elg1-HA: − − + + | − − + +
Stn1-MYC: − + − + | − + − +

IP HA
Elg1
Stn1

INPUT
Elg1
Stn1
Act1

Elg1-HA: − − + +
Ten1-FLAG: − + − +

IP HA
Elg1
Ten1

INPUT
Elg1
Ten1
Act1

Figure 2

Figure 2: Genetic and physical interaction between *ELG1* and *STN1*. A. Teloblot showing epistasis between *elg1∆* and *stn1* mutants. **B.** Co-Immunoprecipitation experiment showing physical interaction between Elg1 and Stn1 and reduced physical interaction between Elg1-sim and Stn1. No interaction could be detected between Elg1 and Cdc13 or Ten1.

Figure 3

Figure 3: Determinants of the Elg1-Stn1 interaction. A. Schematic representation of the Elg1 protein. The 3 SIM motifs, the WalkerA motif and the 2 threonines at the interface with PCNA are shown. **B.** Yeast-Two-Hybrid (YTH) interaction of Cdc13 and Elg1 in a wild type strain. AD: protein fused to the activating domain of Gal4; BD: protein fused to the DNA binding domain of Gal4. e.v.: empty vector.**C.** YTH interaction of Stn1 and Elg1 in a wild type strain. **D.** YTH experiments in the *siz1Δ siz2Δ* and *pol30-RR* background. **E.** Co-Immunoprecipitation experiment showing physical interaction between Cdc13 and PCNA.

Figure 4

Figure 4: The unloading activity of Elg1 and Cdc13 SUMOylation are necessary for the Elg1-Cdc13 interaction. A. YTH experiment in a *elg1Δ* strain. **B.** YTH experiment in a *elg1-sim* strain. **C.** YTH experiment in a *elg1-TT386/7DD* strain. **D.** YTH experiment in a *elg1-sim+DD* strain. **E.** YTH experiment in a *elg1-Walker AB* strain. **F.** Lack of interaction between *cdc13-snm* and Elg1 (in a wild type strain).

Figure 5

Figure 5: Timing of Elg1-Stn1 interaction. A. Co-IP experiment with synchronized cells. Aliquots were taken at time intervals, Elg1 was immunoprecipitated, and the level of Stn1 and Pol3 (the large subunit of DNA polymerase Delta) were monitored by Western blot. Strains with single tags are shown as controls. Whole Cell Extract results are shown in **Figure S4**. **B.** Quantitation of the Western shown in A. **C.** DNA content of the cells used in A, by cell cytrometry. **D.** Chromatin immunoprecipitation at telomeres (Telo-ChIP) in synchronized cells showing PCNA occupancy. **E.** Interaction between Stn1 and Cdc13 in a wild type and a *elg1Δ* strain. **F.** Quantitation of the experiment shown in E. ****:p<0.001.

A

Last origin firing: Elg1 moves with the fork towards the telomeres and unloads S-PCNA.

B

Arrival of S-PCNA with the fork to telomeres leads to the activation of telomerase and to telomere elongation.

C

Elg1 interacts with S-Stn1. Elg1 unloads PCNA, Cdc13 becomes SUMOylated.

D

Stn1 (and presumably Ten1) binds SUMOylated Cdc13, terminating telomerase activity.

Figure 6

Figure 6: Model for the role of SUMOylated PCNA and Elg1 in telomere length regulation. A. Cdc13 binds ssDNA at the telomeres, Elg1 moves with the replisome at the lagging strand, unloading PCNA in each Okazaki fragment. **B.** Arrival of the SUMOylated PCNA at the fork to Cdc13 promotes telomerase activity. **C.** Elg1 interacts with Stn1, which could be SUMOylated. Cdc13 becomes SUMOylated. Elg1 unloads PCNA and leaves telomeres. **D.** Stn1 is now able to interact with Cdc13, evicting Est1 and terminating telomerase activity.

Strain	AD	BD	-URA -LEU	-HIS
Wild Type	E.V.	E.V.		
	TEN1	E.V.		
		ELG1-NTD		
		ELG1-CTD		
		ELG1-NTDsim		
		SUMO		
	SUMO	ELG1-NTD		

Figure S1: Weak interaction between Elg1 and Ten1. Results of YTH experiments.

Figure S2: All *elg1* alleles used are expressed at similar levels. Western blot results.

A

Strain	AD	BD	-URA -LEU	-HIS
cdc13-snm	Stn1	E.V.		
		ELG1-NTD		
		ELG1-NTDsim		
		SUMO		
	Cdc13	E.V.		
		ELG1-NTD		
		ELG1-NTDsim		
		SUMO		

B

Strain	AD	BD	-URA -LEU	-HIS	Strain	-URA -LEU	-HIS
Wild Type	EV	EV			*elg1Δ*		
	Stn1-Nt	EV					
	Stn1-Nt	Elg1-NTD					
	Stn1-Ct	EV					
	Stn1-Ct	Elg1-NTD					

Figure S3

Figure S3: A. The *sumo-no-more* allele of Cdc13 has no effect on the YTH interactions between wt Cdc13 or Stn1 and the N-terminus of Elg1. **B.** The C-terminus of Stn1 interacts with the N-terminus of Elg1 in wild type and *elg1Δ* strains.

Time (minutes)

| | No Tag | Elg1-HA | Stn1-MYC | Pol3-FLAG | Aynchr. | 0' | 10' | 20' | 30' | 40' | 50' | 60' | 70' | 80' | 90' | 100' | 120' | 140' | 160' | 180' | 200' |

ELG1-HA

STN1-MYC

POL3-FLAG

ACT1

Figure S4

Figure Legends

Figure 1: SUMOylated PCNA regulates telomere length. A. Southern blot (Teloblot) showing that lack of SUMOylation of PCNA prevents telomere elongation. Independently-created colonies were passaged ten times, its DNA extracted, digested with *Xho*I and run in an agarose gel. The DNA was then transferred to a nitrocellulose membrane, which was incubated with a radioactive probe that detects telomeres, and a size marker. **B.** Teloblot showing that overexpression of wild type PCNA or Pol30-RR-SUMO fusion, but not Pol30-ubiquitin fusion or Pol30-RR causes telomere elongation.

Figure 2: Genetic and physical interaction between *ELG1* and *STN1*. A. Teloblot showing epistasis between *elg1Δ* and *stn1* mutants. **B.** Co-Immunoprecipitation experiment showing physical interaction between Elg1 and Stn1 and reduced physical interaction between Elg1-sim and Stn1. No interaction could be detected between Elg1 and Cdc13 or Ten1.

Figure 3: Determinants of the Elg1-Stn1 interaction. A. Schematic representation of the Elg1 protein. The 3 SIM motifs, the WalkerA motif and the 2 threonines at the interface with PCNA are shown. **B.** Yeast-Two-Hybrid (YTH) interaction of Cdc13 and Elg1 in a wild type strain. AD: protein fused to the activating domain of Gal4; BD: protein fused to the DNA binding domain of Gal4. e.v.: empty vector.**C.** YTH interaction of Stn1 and Elg1 in a wild type strain. **D.** YTH experiments in the *siz1Δ siz2Δ* and *pol30-RR* background. **E.** Co-Immunoprecipitation experiment showing physical interaction between Cdc13 and PCNA.

Figure 4: The unloading activity of Elg1 and Cdc13 SUMOylation are necessary for the Elg1-Cdc13 interaction. A. YTH experiment in a *elg1Δ* strain. **B.** YTH experiment in a *elg1-sim* strain. **C.** YTH experiment in a *elg1-TT386/7DD* strain. **D.** YTH experiment in a *elg1-sim+DD* strain. **E.** YTH experiment in a *elg1-Walker AB* strain. **F.** Lack of interaction between *cdc13-snm* and Elg1 (in a wild type strain).

Figure 5: Timing of Elg1-Stn1 interaction. A. Co-IP experiment with synchronized cells. Aliquots were taken at time intervals, Elg1 was immunoprecipitated, and the level of Stn1 and Pol3 (the large subunit of DNA polymerase Delta) were monitored by Western blot. Strains with single tags are shown as controls. Whole Cell Extract results are shown in **Figure S4**. **B.** Quantitation of the Western shown in A. **C.** DNA content of the cells used in A, by cell cytometry. **D.** Chromatin immunoprecipitation at telomeres (Telo-ChIP) in synchronized cells showing PCNA occupancy. **E.** Interaction between Stn1 and Cdc13 in a wild type and a *elg1Δ* strain. **F.** Quantitation of the experiment shown in E. **:$p<0.001$.

Figure 6: Model for the role of SUMOylated PCNA and Elg1 in telomere length regulation. A. Cdc13 binds ssDNA at the telomeres, Elg1 moves with the replisome at the lagging strand, unloading PCNA in each Okazaki fragment. **B.** Arrival of the SUMOylated PCNA at the fork to Cdc13 promotes

telomerase activity. **C.** Elg1 interacts with Stn1, which could be SUMOylated. Cdc13 becomes SUMOylated. Elg1 unloads PCNA and leaves telomeres. **D.** Stn1 is now able to interact with Cdc13, evicting Est1 and terminating telomerase activity.

Figure S1: Weak interaction between Elg1 and Ten1. Results of YTH experiments.

Figure S2: All *elg1* alleles used are expressed at similar levels. Western blot results.

Figure S3: A. The *sumo-no-more* allele of Cdc13 has no effect on the YTH interactions between wt Cdc13 or Stn1 and the N-terminus of Elg1. **B.** The C-terminus of Stn1 interacts with the N-terminus of Elg1 in wild type and *elg1Δ* strains.

Figure S4: Whole cell extract showing the level of Elg1, Stn1 and Pol3 in the cell cycle experiment shown in **Figure 5**.