Title

Synthetic lipid rafts formed by cholesterol nano-patch induce T cell activation

Authors

Yunmin Jung [1], Young-Joo Kim [1,2], Sunmi Lee [1], Jungmin Nam [1,2], Seonyeong Ha [1,2], Jinwoo Cheon [1,2,3,*], Minsuk Kwak [1,2,*]

Affiliations

[1] Institute for Basic Science Center for Nanomedicine, Seoul 03722, South Korea

[2] Department of Nano-Biomedical Engineering, Advanced Science Institute, Yonsei University, Seoul 03722, South Korea

[3] Department of Chemistry, Yonsei University, Seoul 03722, South Korea

*e-mail: minsuk.kwak@yonsei.ac.kr, jcheon@yonsei.ac.kr

Abstract

Lipid rafts are small and dynamic membrane compartments that are enriched with cholesterols or sphingolipids. The coalescence of small lipid rafts into a large lipid raft forms a signaling platform at the immunological synapse that plays a crucial role in controlling T cell activation. Cholesterol is a major component for the formation and function of lipid rafts, but the relationship between the spatial configuration of cholesterols and the lipid raft mediated membrane segregation as well as TCR signaling remains poorly understood. Here we used cholesterol conjugated DNA origami nanostructures, cholesterol nano-patch (CNP) to construct lipid rafts on intact live T cells. CNPs are two-dimensional DNA scaffolds that allow controlling the spatial presentation of cholesterols in the plasma membrane. CNPs efficiently bind onto the T cell membrane and rapidly induce large, polarized membrane cap structures which colocalize with flotillin-1. These CNP mediated membrane segregations, which we define as synthetic lipid rafts, drive membrane segregation of key signaling molecules including TCR and CD45, thereby triggering downstream TCR signaling and subsequently functional T cell responses without antigenic challenges. Our work highlights the potential of CNP in uncovering the mechanisms that lipid rafts induced by cholesterols drive membrane protein reorganization and control T cell signaling, and offers valuable insights into the design of antigen-independent immunotherapies.

Keywords

1   Introduction

Lipid rafts are specialized membrane compartments containing high cholesterol or sphingolipids [1,2]. Due to

their more ordered, thicker, and less fluidic characteristics, lipid rafts contribute to the spatial organization

of membrane proteins and influence abroad arrays of cellular signaling events [3-6]. Lipid rafts are nanoscale

assemblies (10–200 nm) that can be stabilized to form a large platform for controlling cell signaling [1,3].

Cholesterol is an essential component of the lipid raft and is involved in a large variety of cellular functions

[7-11]. In synthetic model membranes or in molecular simulations, high contents of cholesterol can drive

spontaneous liquid-liquid phase separation (LLPS) of the lipids in the membrane, and the concentration of

cholesterols showed a high correlation with the formation of liquid-ordered (Lo) phase lipid domain [12-15].

It is also known that cholesterols form a thicker lipid bilayer therefore it influences the positioning of the

membrane proteins [16-18].

In T cells, a large coalescence of lipid rafts is formed at the immunological synapses between T cells and

APCs and serve as the signaling platform for T cell activation by driving spatial reorganization of key

signaling molecules [19-22]. T cell receptors (TCR) and other signaling molecules, such as Lck, LAT, or

ZAP70 are recruited into the large lipid raft at IS, while phosphatase CD45 is excluded from it [6,23,24]. The

role of cholesterol in T cells has been studied by enrichment using squalene administration or inhibiting

cholesterol esterification, or by extraction with methyl-β-cyclodextrin in the whole cell membrane showing

that modulation of cholesterol homeostasis profoundly influences T cell functions [25-28]. There are two

opposing effects of cholesterol in regulating TCR at the molecular level has been suggested [29] that binding

of cholesterol to TCR stabilizes its inactive conformation [30,31] or binding of cholesterol causes TCR

nanoclustering which enhances the avidity of the TCR-antigen interaction and positively regulates T cell

activation [9]. Another recent study suggested that cholesterol regulates CD45 exclusion at the tip of

microvilli in T cells at their resting state [17]. However, due to the technical limitation for controlling the

distribution or density of cholesterols in the live cell membrane, how the spatial organization of cholesterols

modulates lipid-raft-driven protein segregation and downstream TCR signaling is largely unknown.

To directly investigate the roles of cholesterols in the coalescence of lipid rafts formation and protein

partitioning in T cells, we designed a DNA origami nanostructure, cholesterol nano-patch (CNP) as a

nanosized lipid raft-like monomer when it attaches to the membrane. On a single CNP, we can organize

multiple cholesterols in a programmable manner to artificially induce lipid rafts by creating a high local

density of cholesterols on the cell membrane. The controllable architecture of DNA origami has been

applied to manipulate the spatial configuration of biomolecules with nanoscale precision [32,33] . Our result

showed that the CNP binds to the outer leaflet of the T cell membrane and forms a large and stably polarized

cluster. We defined this large coalescence of multiple CNPs segregated into a distinctive domain via

enrichment of cholesterols as a synthetic lipid raft. This CNP-mediated synthetic lipid raft co-polarized

with lipid raft protein flotillin-1 [34]. We found that CD45 molecules are largely excluded from the synthetic

lipid rafts, supporting the lipid raft mediated CD45 exclusion mechanism. Furthermore, the formation of

the synthetic lipid raft drives TCR accumulation to it and enhances CD69 expression and ZAP-70

phosphorylation. These results illustrate that the local concentration of cholesterols drives the formation of

large, polarized lipid rafts on T cell membranes that induce the spatial reorganization of signaling molecules,

activating T cells independent of antigenic challenges. Our results demonstrate the capability of DNA

origami in interrogating the mechanism of lipid raft-driven protein segregation on T cell membrane and

reveal the role of nanoscale arrangement and local concentration of cholesterols in regulating T cell

signaling.

Results

Design of DNA origami structure for cholesterol patterning.

We fabricated sheet-like DNA nanostructures, DNA nano-patch (NP), employing a DNA origami method

to construct a molecular template where the number or arrangement of cholesterols can be programmed.

(Fig. 1). DNA origami is a molecular fabrication technique enabling the construction of a customized

nanostructure with desired shape and size [32,33]. By folding a long single-stranded DNA (ssDNA) into a

target geometry through rationally designed hybridizations with multiple short ssDNAs, a customized DNA

nanostructure can be easily constructed with high shape accuracy. The DNA NP is a square sheet with a

side of 60 nm and a height of 6 nm (**Fig. 1a, b and Supplementary Table. 1**). To control the number of

cholesterols on it, an array of anchors was designed on one side, where additional ssDNA (DNA handles)

can be extended to accommodate cholesterol-conjugated DNAs (Chol-DNAs). Each DNA NP can display

up to 72 Chol-DNAs in an array of 12 x 6 with rows and columns separated by 3.5 nm and 7 nm, respectively,

which enables inter-cholesterol spacing from up to 3.5 nm to 38.5 nm (**Fig. 1a**). The DNA NP also contains

Cy5-fluorophores at each of four corners to allow for visualization. Through hybridization of DNA handles

with Chol-DNA strands and assist DNAs, which formed a stable duplex with Chol-DNA, a DNA NP is

converted to a cholesterol nano-patch (CNP), which facilitates the hydrophobic interactions with lipid

bilayers (**Fig. 1b**). The estimated length of hybridized Chol-DNAs is about 10.88 nm (32 base-pairs). The

transmission electron microscope (TEM) images showed that the DNA NP was well folded with a high

folding yield (**Fig.1c** and **Supplementary Fig. 1**) and the CNP binds to the surface of synthetic small lipid

vesicles through hydrophobic interactions (**Fig. 1d**). To control the local concentration of cholesterols, we

varied the spatial distribution of cholesterols to allow patterning 18- (0.005 cholesterols/nm^2), 36- (0.01

cholesterol/nm^2), or 72- (0.02 cholesterols/nm^2) cholesterols per a single NP, named as 18-CNP, 36-CNP,

and 72-CNP, respectively (**Fig. 1e**).

Formation of synthetic lipid rafts on T cell membranes

Next, we examined whether these CNPs can drive the formation of lipid rafts by increasing the local

concentration of cholesterols on T cell membranes. Two different membrane binding strategies were

employed: (1) Chol-DNA, assist-DNA, and DNA NP was sequentially applied to Jurkat T cells

(**Supplementary Fig. 2a**); (2) CNPs were pre-assembled in a tube and added to T cells (**Fig. 1b**) as a single-

step reaction. Using both strategies, the addition of 72-CNP spontaneously formed micron-sized (~14.3 ±

5 μm in perimeter) polarized clusters, synthetic lipid rafts, on the T cell membranes (**Fig 2a, and**

Supplementary Fig 2b). The mean fluorescent intensity (MFI) of single cell-bound CNP was 3.4 times

higher in the single step binding reaction than it of the sequential reaction, indicating that direct treatment

of pre-complexed CNP is more efficient in membrane binding (**Supplementary Fig. 2c**). Next, to

investigate the impact of the nanoscale distribution of cholesterols on forming the CNP clusters and

polarization, we applied the CNP with different numbers of cholesterols (**Fig. 1e**). The NP without

cholesterol (0-chol) did not bind on Jurkat T cell membranes. CNP with more cholesterols showed higher

efficiency in the formation of polarized clusters of CNPs on the membrane (**Fig. 2b, c**). In CNP72-treated

samples, 69% of cells formed single-polarized synthetic lipid rafts on their membrane, and the percentages

of this single-polarized population dropped to 50% and 33% in 36-CNP- and 18-CNP-treated cells,

respectively. (**Fig. 2c, Supplementary Fig. 2d**). We also found that a minority of cells (< 5%) have more

than one polarized cluster (**Fig. 2c, Supplementary Fig. 2e**). Next, we tested the stability of CNP-induced

synthetic lipid rafts in a physiological condition and the effects on T cell proliferation. We found that T

cells treated with 72-CNP maintained the polarized caps of CNP for 4 hours while most of the 18- and 36-

CNPs detached from the cells, revealing that a higher local density of cholesterols enhances the stability of

synthetic lipid rafts (**Fig. 2d** and **Supplementary Fig. 3**). Binding of CNPs did not affect the proliferation

for 4 days regardless of the number of cholesterols on CNP (**Supplementary Fig. 4)**. The 72-CNP similarly

induced the formation of the synthetic lipid rafts in resting or effector primary mouse and human T cells

(**Fig. 2e, Supplementary Fig. 5a**), and these synthetic lipid rafts remained almost intact for 4 hours

(**Supplementary Fig. 5b, c**). Together, these results suggest that cholesterol-mediated binding of CNP

drives membrane segregation, and tighter cholesterol distribution on CNP enhances membrane binding

efficiency, synthetic lipid raft formation, and stability.

Synthetic lipid rafts are colocalized with Flot1-cap on T cells

To characterize the nature of the CNP mediated synthetic lipid rafts on T cell membrane, we examined the

spatial organization and dynamics of the CNP and flotillin-1 (Flot1), a lipid raft associated protein [35,36].

Flot1 is known to segregate into lipid rafts by forming multimeric complexes at the inner leaflets [37,38]. In T

cells, Flot1 is weakly polarized on the membrane at the resting state and its polarization is enhanced by

stimulations [39-43]. Here, we formed synthetic lipid rafts using the 72-CNP on the Jurkat T cells stably

expressing Flot1 with Halo-tag at the C-terminus (Flot1-Halo). We monitored the dynamics of Flot1 for 30

minutes after adding the CNPs using fluorophore-conjugated Halo ligands. As reported previously, Flot1

formed cap-like membrane microdomains (**Fig 3a, b**), and it was more strongly polarized (50%) upon 72-

CNP addition than untreated controls (32.8%) (**Supplementary Fig. 6a**). CNP induced synthetic lipid rafts

showed high colocalization with Flot1-caps after 30 min incubation (the mean Pearson's correlation

coefficient (*R'*) and Manders overlap coefficient (*MOC*) were 0.63 ± 0.14 for both). The high colocalization

of the CNP clusters and Flot1-caps indicates the CNP mediated synthetic lipid raft can indeed mimic the

coalescence of lipid rafts (**Fig. 3c**). We observed three cases in live cell time-lapse imaging: (1) CNP

initially bound homogeneously on Flot1-unpolarized cells and it induced co-clustering of Flot1-caps and

CNP clusters (40 %) (**Supplementary Fig. 6b and Supplementary Movie 1**); (2) CNP initially bound

homogeneously on Flot1-polarized cells and moved to the Flot1 caps (33 %) (**Supplementary Fig. 6c and**

Supplementary Movie 2); (3) CNP was preferentially bound on the weakly polarized Flot1 cap and

enhanced the Flot1 cap polarization (27 %) (**Supplementary Fig. 6d and Supplementary Movie 3**).

Overall, CNP-mediated synthetic lipid rafts were highly colocalized Flot1 caps in T cells, and the binding

of CNP on the membrane further enhanced the polarization of Flot1, similar to when T cells activated by

anti-CD3/CD28, chemokines, or lipid raft crosslinking [41-43]. Interestingly the CNP is mostly bound on the

flat cell body but not on the microvilli of T cells (**Supplementary Fig. 7**).

It has been reported that the lateral mobility of Flot1-cap is significantly slower compared to the rest of

Flot1 in the non-cap area [39]. We monitored the lateral mobility of the 72-CNP on the Jurkat T cell membrane

using fluorescence recovery after photobleaching (FRAP). [44] Interestingly, there was almost no detectable

recovery of 72-CNP fluorescence in the synthetic lipid rafts for 5 minutes after rapid photobleaching.

(**Supplementary Fig. 8 and Supplementary Movie 4 and 5**). This result suggests that CNP formed highly

stable membrane domains exhibiting very slow diffusion. In contrast, there was a slow but gradual recovery

of Flot1 fluorescence reached a plateau after ~200 seconds in the synthetic lipid rafts (**Fig. 3d-f** and

Supplementary Movie 5). The diffusion coefficient (D) of Flot1 in the non-cap (non-raft) areas was ~2.1-

fold higher ($D = 0.21 \ \mu m^2 s^{-1}$) than D of Flot1 in the cap area (synthetic lipid raft) ($D = 0.10 \ \mu m^2 s^{-1}$) (**Fig.**

3g). There was a significant increase in D for Flot1 in the naturally formed polarized cap area ($D = 0.15$

$\mu m^2 s^{-1}$) compared to that for Flot1 in the synthetic lipid rafts induced by CNP binding (**Fig. 3g**). Together,

these results demonstrate that stable CNP clusters are formed at the outer leaflets, and these also stabilize

the lipid rafts on the inner leaflets of the membrane.

CD45 is excluded from the synthetic lipid raft

CD45 exclusion from the immunological synapse (IS) is one of the crucial factors in T cell activation by

limiting abundant phosphatase activities [45-48]. It has been reported that artificially induced CD45 exclusion

can induce T cell activation without antigen [46]. Two models have proposed the CD45 exclusion mechanism:

the kinetic segregation model [49-51] and the lipid raft driven segregation model [17,24,52-54]. The former is ligation

dependent, explaining that CD45 is excluded from the close contacts due to the long extracellular domain

of CD45 (~23-50 nm) which is larger than the intermembrane junction formed by the TCR-pMHC

interaction (~15 nm) [49-51]. The latter is a ligation independent mechanism [17]. We examined whether CD45

is segregated from the CNP mediated synthetic lipid rafts independent of ligand association. Upon 72-CNP

addition, CD45 molecules partitioned away from the regions of the membrane where 72-CNP formed a

large synthetic lipid raft in the live T cell (**Fig. 4a, b**). The mean Pearson's *R'* and *MOC* between the CD45

and CNP obtained from 51 polarized cells were low as 0.18 ± 0.09 and 0.13 ± 0.07, respectively (**Fig. 4c**).

To quantify the level of exclusion, we segmented sub-areas of the cell membrane for the synthetic lipid raft

(raft, marked in yellow) and the non-raft (marked in light blue) regions (**Fig. 4d**) and calculated the

ratiometric intensity score (RIS) [17] between the CD45 and the CNP. RIS becomes zero when both molecules

have the same normalized intensities, +1 when CD45 is only present, and -1 when 72-CNP is only present.

The significant difference in RIS values between the raft (RIS= -0.67 ± 0.11) and the non-raft (RIS= +0.56

± 0.16) areas indicate CD45 is excluded by CNP-mediated synthetic lipid rafts (**Fig. 4e, f**). Furthermore,

the average intensity of CD45 within the synthetic raft was 66.9 % lower than that of CD45 localized in

non-raft regions (**Fig. 4g**). These results demonstrate that about two-thirds of CD45 molecules in a cell can

be segregated merely by lipid rafts on the T cell membrane without forming close contacts, supporting the

lipid raft mediated CD45 segregation model.

Synthetic lipid raft induces T cell activation

We then investigated whether the CNP-mediated synthetic lipid raft can directly activate T cells without

any antigenic challenge. During early T cell activation, TCRs are accumulated at the core of IS [55], and

cholesterol can bind to TCRβ and induces TCR nanoclustering [56]. Thus, we tested whether the CNP can

recruit TCRs. In contrast to CD45, which was largely segregated away from the synthetic lipid rafts (**Fig.**
4), we could observe CD3 enrichment at the microdomains of CNP (**Fig. 5a, b**) in some cells. The average
Pearson's *R* and *MOC* were 0.44±0.15 for both, and 37% of the analyzed cells (34 out of 92 cells) showed
strong colocalization (*MOC* > 0.5) between TCR and CNP clusters (**Fig. 5c**). Wide variations in CD3 and
CNP colocalization coefficients indicate that the efficiency of CD3 recruitment to the synthetic lipid rafts
varies from cell to cell.

Next, we examined whether the CNP could transmit downstream TCR signals. Phosphorylation of cytosolic
kinase TCR-zeta chain-associated 70-kDa tyrosine phosphoprotein (ZAP-70) is one of the early
downstream TCR signaling [57,58]. We examined phosphorylation at tyrosine at 319-residue of ZAP-70 (ZAP-
70 ^{Y319}) in the lysate of Jurkat T cells that were untreated or treated with 36- or 72-CNPs by immunoblot
assay [59]. While 36-CNP treated cells showed only a modest and insignificant enhancement of ZAP-70 ^{Y319}
phosphorylation (2.0 ± 0.76 folds), 72-CNP treatment induced a significant increase in phosphorylation of
ZAP-70 ^{Y319} (2.7 ± 0.5 folds) compared to the basal level of phosphorylation in untreated cells (**Fig. 5d, e**).

We then monitored the surface expression of CD69 as an early cellular event indicative of T cell activation
[60]. Jurkat T cells were incubated with 72-CNP or stimulated with PMA/ionomycin and analyzed for CD69
expression by flow cytometry. Although 72-CNP was less effective than PMA/ionomycin stimulation
(37.9%), the percentages of CD69+ cells increased in the presence of 36-CNP (27.7%) and 72-CNP (33.7%)
compared to untreated control (13%) after 4 hours of incubation (**Fig. 5f, g**). This indicates that the shorter
inter-cholesterol spacing and higher cholesterol density on the CNP induced more efficient T cell activation
in terms of the CD69 expression. In summary, these results demonstrate that the synthetic lipid rafts can
activate T cells by recruiting TCRs while excluding CD45 and transmitting TCR signaling without specific
antigenic challenges, and cholesterol density and patterning in the CNP affect the efficiency of downstream
TCR signaling as well as CD69 expression in T cell activation.

185
Discussion

In this work, we have demonstrated a novel approach to create synthetic lipid rafts, large clusters of
nanosized lipid raft-like monomers, on the T cell membranes that effectively drives membrane protein
segregations and early T cell activation. Previous studies using model membranes composed of synthetic
or cell membrane-derived lipids have demonstrated that high levels of cholesterols can induce a large-scale
liquid-liquid phase separation (LLPS) of the lipids [12]. However, this micrometer scale membrane
segregation formed by cholesterol has not been observed in the live cell membrane. Here, we developed a
DNA origami-based cholesterol modification platform called CNP that permits facile control of the number,
distance, or configuration of cholesterols on one side of DNA origami. The array of multiple cholesterols

on the CNP increased local concentration of cholesterols upon binding on the membrane, resembling the

nano-sized lipid raft monomer. The binding of CNPs drove a large-scale LLPS of the membrane, forming

a large synthetic lipid raft. The number of cholesterols linearly correlated to the formation of the segregation

of the domain. Our results suggest that the changes in local distribution of cholesterols shift the

thermodynamic equilibrium to form a large phase separation of the live cell membrane.

Another important enigma of the cell plasma membrane is the asymmetry between the two leaflets [61].

Although the physiological quantity of the cholesterols in the inner or outer leaflet is controversial [62], the

lipid composition of the outer leaflet lipid is known to be sufficient to form a lipid raft domain on the

membrane [63]. The lipid raft formation in the outer leaflet driven by cholesterol enrichment alters the spatial

distribution of other lipid components in the opposing leaflet, known as lipid-mediated transbilayer

coupling [63,64]. This may cause lipid raft formation on both sides of the plasma membrane and regulate T

cell signaling. In our system, we assume that the cholesterols complexed with the CNP exclusively insert

into the outer leaflet of the membrane because the bulky DNA origami structure is unlikely to undergo flip-

flop. A significantly decreased diffusion of Flot1 upon the CNP binding indicates that cholesterols in the

outer membrane can induce transbilayer coupling with the adjacent inner leaflet of the membrane. Our

nanotechnology platform can promote research efforts towards dissecting the mechanism driving

transbilayer coupling in T cells and expand our understanding of the roles of cholesterol-driven lipid rafts

in regulating T cell signaling.

Our results showed that ~67% of CD45 molecules were excluded from the synthetic lipid raft without

forming close contact with an APC. It strengthens the mechanism of lipid raft-driven partitioning of the

CD45 and supports the hypothesis that cholesterol is the cause of the segregation of CD45 [17]. The

incomplete segregation of CD45 in our study and the previously reported reconstitution study using

modification on length of ligations [65,66] may indicate that the CD45 exclusion at the IS is driven by both

lipid raft- and kinetic segregation mechanisms. We also demonstrated that the synthetic lipid raft

constructed with cholesterols could induce T cell activation without specific antigens. Note that although

we could observe TCR accumulation at the synthetic lipid rafts, some cells did not display TCR polarization

nor high colocalization with the CNP clusters. It could be partially caused by the fast internalization of the

TCR antibodies bound on the membrane or by the different localization efficiency on the microvilli-

localization [67]. The efficient TCR polarization might require other signaling mechanisms, such as actin

cytoskeleton rearrangement.

We found that the CNP is highly colocalized to the preassembled Flot1 cap on T cells. The preassembling

of the Flot1 cap is a unique feature found in hematopoietic cells [68]. It has been reported that Flot1

preassembled cap is resistant to cholesterol-disrupting agents [40]. Our results show that the assembly of Flot1

could be initiated and enhanced by the high local concentration of cholesterols. Also, once formed, the Flot1 clusters maintain their stability. The mechanism driving the preassembly of the Flot1 cap and the physiological roles of this spatially distinctive membrane remain poorly understood. The CNPs might serve as an optimal tool for dissecting the underlying mechanism of the Flot1 preassembly.

Our study reveals the critical role of cholesterols in mediating T cell signaling and function and the possibility of direct control of T cell activation using patterned cholesterols. Lipid rafts have been one of the controversial topics in biology for decades. Our strategy for controlling the spatial distribution of cholesterol can be easily substituted or extended with other major components of lipid rafts, such as sphingolipids or proteins. We anticipate that our strategy for forming and controlling synthetic lipid rafts can be widely applied and will become a versatile and powerful tool for studying the direct function of lipid rafts in various cell types.

References

Pike, L. J. Rafts defined: a report on the Keystone Symposium on Lipid Rafts and Cell Function. *J Lipid Res* **47**, 1597-1598, doi:10.1194/jlr.E600002-JLR200 (2006).

Simons, K. & Ikonen, E. Functional rafts in cell membranes. *Nature* **387**, 569-572, doi:10.1038/42408 (1997).

Lingwood, D. & Simons, K. Lipid Rafts As a Membrane-Organizing Principle. *Science* **327**, 46-50, doi:10.1126/science.1174621 (2010).

Sezgin, E., Levental, I., Mayor, S. & Eggeling, C. The mystery of membrane organization: composition, regulation and roles of lipid rafts. *Nature Reviews Molecular Cell Biology* **18**, 361-374, doi:10.1038/nrm.2017.16 (2017).

Stone, M. B., Shelby, S. A., Núñez, M. F., Wisser, K. & Veatch, S. L. Protein sorting by lipid phase-like domains supports emergent signaling function in B lymphocyte plasma membranes. *eLife* **6**, e19891, doi:10.7554/eLife.19891 (2017).

Varshney, P., Yadav, V. & Saini, N. Lipid rafts in immune signalling: current progress and future perspective. *Immunology* **149**, 13-24, doi:10.1111/imm.12617 (2016).

Sheng, R. *et al.* Cholesterol modulates cell signaling and protein networking by specifically interacting with PDZ domain-containing scaffold proteins. *Nature Communications* **3**, 1249, doi:10.1038/ncomms2221 (2012).

Levitan, I., Singh, D. & Rosenhouse-Dantsker, A. Cholesterol binding to ion channels. *Frontiers in Physiology* **5**, doi:10.3389/fphys.2014.00065 (2014).

Molnár, E. *et al.* Cholesterol and Sphingomyelin Drive Ligand-independent T-cell Antigen Receptor Nanoclustering*. *Journal of Biological Chemistry* **287**, 42664-42674, doi:https://doi.org/10.1074/jbc.M112.386045 (2012).

Igarashi, M., Honda, A., Kawasaki, A. & Nozumi, M. Neuronal Signaling Involved in Neuronal Polarization and Growth: Lipid Rafts and Phosphorylation. *Frontiers in Molecular Neuroscience* **13**, doi:10.3389/fnmol.2020.00150 (2020).

Wang, Y. *et al.* Cholesterol-Rich Lipid Rafts in the Cellular Membrane Play an Essential Role in Avian Reovirus Replication. *Frontiers in Microbiology* **11**, doi:10.3389/fmicb.2020.597794 (2020).

12   Crane, J. M. & Tamm, L. K. Role of cholesterol in the formation and nature of lipid rafts in planar
and spherical model membranes. *Biophys J* **86**, 2965-2979, doi:10.1016/s0006-3495(04)74347-7
(2004).
13   McConnell, H. M. & Vrljic, M. Liquid-Liquid Immiscibility in Membranes. *Annual Review of*
Biophysics and Biomolecular Structure **32**, 469-492,
doi:10.1146/annurev.biophys.32.110601.141704 (2003).
14   Risselada, H. J. & Marrink, S. J. The molecular face of lipid rafts in model membranes.
Proceedings of the National Academy of Sciences **105**, 17367-17372,
doi:10.1073/pnas.0807527105 (2008).
15   Macdermaid, C. *et al.* Molecular dynamics simulations of cholesterol-rich membranes using a
coarse-grained force field for cyclic alkanes. *The Journal of Chemical Physics* **143**, 243144,
doi:10.1063/1.4937153 (2015).
16   Smondyrev, A. M. & Berkowitz, M. L. Structure of dipalmitoylphosphatidylcholine/cholesterol
bilayer at low and high cholesterol concentrations: molecular dynamics simulation. *Biophys J* **77**,
2075-2089, doi:10.1016/S0006-3495(99)77049-9 (1999).
17   Jung, Y., Wen, L., Altman, A. & Ley, K. CD45 pre-exclusion from the tips of T cell microvilli prior to
antigen recognition. *Nature Communications* **12**, 3872 (2021).
18   Bretscher, M. S. & Munro, S. Cholesterol and the Golgi Apparatus. *Science* **261**, 1280-1281,
doi:10.1126/science.8362242 (1993).
19   Chichili, G. R., Westmuckett, A. D. & Rodgers, W. T cell signal regulation by the actin
cytoskeleton. *J Biol Chem* **285**, 14737-14746, doi:10.1074/jbc.M109.097311 (2010).
20   Owen, D. M. *et al.* High plasma membrane lipid order imaged at the immunological synapse
periphery in live T cells. *Molecular membrane biology* **27**, 178-189,
doi:10.3109/09687688.2010.495353 (2010).
21   Gaus, K., Chklovskaia, E., Fazekas de St. Groth, B., Jessup, W. & Harder, T. Condensation of the
plasma membrane at the site of T lymphocyte activation. *Journal of Cell Biology* **171**, 121-131,
doi:10.1083/jcb.200505047 (2005).
22   Horejsí, V. Lipid rafts and their roles in T-cell activation. *Microbes Infect* **7**, 310-316,
doi:10.1016/j.micinf.2004.12.004 (2005).
23   Kabouridis, P. S. Lipid rafts in T cell receptor signalling. *Molecular membrane biology* **23**, 49-57,
doi:10.1080/09687860500453673 (2006).
24   Rodgers, W. & Rose, J. K. Exclusion of CD45 inhibits activity of p56lck associated with glycolipid-
enriched membrane domains. *Journal of Cell Biology* **135**, 1515-1523,
doi:10.1083/jcb.135.6.1515 (1996).
25   Bietz, A., Zhu, H., Xue, M. & Xu, C. Cholesterol Metabolism in T Cells. *Front Immunol* **8**, 1664,
doi:10.3389/fimmu.2017.01664 (2017).
26   Surls, J. *et al.* Increased Membrane Cholesterol in Lymphocytes Diverts T-Cells toward an
Inflammatory Response. *PLOS ONE* **7**, e38733, doi:10.1371/journal.pone.0038733 (2012).
27   Kabouridis, P. S., Janzen, J., Magee, A. L. & Ley, S. C. Cholesterol depletion disrupts lipid rafts and
modulates the activity of multiple signaling pathways in T lymphocytes. *European Journal of*
Immunology **30**, 954-963, doi:https://doi.org/10.1002/1521-4141(200003)30:3<954::AID-
IMMU954>3.0.CO;2-Y (2000).
28   Yang, W. *et al.* Potentiating the antitumour response of CD8+ T cells by modulating cholesterol
metabolism. *Nature* **531**, 651-655, doi:10.1038/nature17412 (2016).
29   Pathan-Chhatbar, S. *et al.* Direct Regulation of the T Cell Antigen Receptor's Activity by
Cholesterol. *Frontiers in Cell and Developmental Biology* **8**, doi:10.3389/fcell.2020.615996
(2021).

30 Swamy, M. *et al.* A Cholesterol-Based Allostery Model of T Cell Receptor Phosphorylation.
Immunity **44**, 1091-1101, doi:https://doi.org/10.1016/j.immuni.2016.04.011 (2016).
31 Chen, Y. *et al.* Cholesterol inhibits TCR signaling by directly restricting TCR-CD3 core tunnel
motility. *Mol Cell* **82**, 1278-1287.e1275, doi:10.1016/j.molcel.2022.02.017 (2022).
32 Douglas, S. M. *et al.* Self-assembly of DNA into nanoscale three-dimensional shapes. *Nature* **459**,
414-418 (2009).
33 Rothemund, P. W. K. Folding DNA to create nanoscale shapes and patterns. *Nature* **440**, 297-302
(2006).
34 Ficht, X. *et al.* In Vivo Function of the Lipid Raft Protein Flotillin-1 during CD8+ T Cell–Mediated
Host Surveillance. *The Journal of Immunology* **203**, 2377-2387, doi:10.4049/jimmunol.1900075
(2019).
35 Ficht, X. *et al.* In Vivo Function of the Lipid Raft Protein Flotillin-1 during CD8(+) T Cell-Mediated
Host Surveillance. *J Immunol* **203**, 2377-2387, doi:10.4049/jimmunol.1900075 (2019).
36 Yokoyama, H. & Matsui, I. The lipid raft markers stomatin, prohibitin, flotillin, and HflK/C (SPFH)-
domain proteins form an operon with NfeD proteins and function with apolar polyisoprenoid
lipids. *Crit Rev Microbiol* **46**, 38-48, doi:10.1080/1040841x.2020.1716682 (2020).
37 Otto, G. P. & Nichols, B. J. The roles of flotillin microdomains--endocytosis and beyond. *J Cell Sci*
124, 3933-3940, doi:10.1242/jcs.092015 (2011).
38 Kwiatkowska, K., Matveichuk, O. V., Fronk, J. & Ciesielska, A. Flotillins: At the Intersection of
Protein S-Palmitoylation and Lipid-Mediated Signaling. *International Journal of Molecular*
Sciences **21** (2020).
39 Langhorst, M. F. *et al.* Preformed reggie/flotillin caps: stable priming platforms for macrodomain
assembly in T cells. *The FASEB Journal* **20**, 711-713, doi:https://doi.org/10.1096/fj.05-4760fje
(2006).
40 Rajendran, L. *et al.* Asymmetric localization of flotillins/reggies in preassembled platforms
confers inherent polarity to hematopoietic cells. *Proceedings of the National Academy of*
Sciences **100**, 8241-8246, doi:10.1073/pnas.1331629100 (2003).
41 Affentranger, S., Martinelli, S., Hahn, J., Rossy, J. & Niggli, V. Dynamic reorganization of flotillins
in chemokine-stimulated human T-lymphocytes. *BMC Cell Biology* **12**, 28, doi:10.1186/1471-
2121-12-28 (2011).
42 Giri, B. *et al.* CXCL12-induced partitioning of flotillin-1 with lipid rafts plays a role in CXCR4
function. *Eur J Immunol* **37**, 2104-2116, doi:10.1002/eji.200636680 (2007).
43 Slaughter, N. *et al.* The flotillins are integral membrane proteins in lipid rafts that contain TCR-
associated signaling components: implications for T-cell activation. *Clin Immunol* **108**, 138-151,
doi:10.1016/s1521-6616(03)00097-4 (2003).
44 Axelrod, D., Koppel, D. E., Schlessinger, J., Elson, E. & Webb, W. W. Mobility measurement by
analysis of fluorescence photobleaching recovery kinetics. *Biophysical Journal* **16**, 1055-1069,
doi:https://doi.org/10.1016/S0006-3495(76)85755-4 (1976).
45 Courtney, A. H. *et al.* CD45 functions as a signaling gatekeeper in T cells. *Science Signaling* **12**,
eaaw8151, doi:10.1126/scisignal.aaw8151 (2019).
46 James, J. R. & Vale, R. D. Biophysical mechanism of T-cell receptor triggering in a reconstituted
system. *Nature* **487**, 64-69, doi:10.1038/nature11220 (2012).
47 Sibener, L. V. *et al.* Isolation of a Structural Mechanism for Uncoupling T Cell Receptor Signaling
from Peptide-MHC Binding. *Cell* **174**, 672-687.e627, doi:10.1016/j.cell.2018.06.017 (2018).
48 Varma, R., Campi, G., Yokosuka, T., Saito, T. & Dustin, M. L. T cell receptor-proximal signals are
sustained in peripheral microclusters and terminated in the central supramolecular activation
cluster. *Immunity* **25**, 117-127, doi:10.1016/j.immuni.2006.04.010 (2006).

49 Wild, M. K. *et al.* Dependence of T Cell Antigen Recognition on the Dimensions of an Accessory
Receptor–Ligand Complex. *Journal of Experimental Medicine* **190**, 31-42,
doi:10.1084/jem.190.1.31 (1999).
50 Davis, S. J. & van der Merwe, P. A. The structure and ligand interactions of CD2: implications for
T-cell function. *Immunology Today* **17**, 177-187, doi:https://doi.org/10.1016/0167-
5699(96)80617-7 (1996).
51 Shaw, A. S. & Dustin, M. L. Making the T cell receptor go the distance: a topological view of T cell
activation. *Immunity* **6**, 361-369, doi:10.1016/s1074-7613(00)80279-4 (1997).
52 Hořejší, V. Lipid rafts and their roles in T-cell activation. *Microbes and Infection* **7**, 310-316,
doi:https://doi.org/10.1016/j.micinf.2004.12.004 (2005).
53 Janes, P. W., Ley, S. C., Magee, A. I. & Kabouridis, P. S. The role of lipid rafts in T cell antigen
receptor (TCR) signalling. *Semin Immunol* **12**, 23-34, doi:10.1006/smim.2000.0204 (2000).
54 Zhang, M. *et al.* CD45 Signals outside of Lipid Rafts to Promote ERK Activation, Synaptic Raft
Clustering, and IL-2 Production1. *The Journal of Immunology* **174**, 1479-1490,
doi:10.4049/jimmunol.174.3.1479 (2005).
55 Grakoui, A. *et al.* The immunological synapse: a molecular machine controlling T cell activation.
Science **285**, 221-227, doi:10.1126/science.285.5425.221 (1999).
56 Molnár, E. *et al.* Cholesterol and sphingomyelin drive ligand-independent T-cell antigen receptor
nanoclustering. *J Biol Chem* **287**, 42664-42674, doi:10.1074/jbc.M112.386045 (2012).
57 Wang, H. *et al.* ZAP-70: an essential kinase in T-cell signaling. *Cold Spring Harb Perspect Biol* **2**,
a002279, doi:10.1101/cshperspect.a002279 (2010).
58 Chan, A. C. & Shaw, A. S. Regulation of antigen receptor signal transduction by protein tyrosine
kinases. *Current Opinion in Immunology* **8**, 394-401, doi:https://doi.org/10.1016/S0952-
7915(96)80130-0 (1996).
59 Di Bartolo, V. *et al.* Tyrosine 319, a newly identified phosphorylation site of ZAP-70, plays a
critical role in T cell antigen receptor signaling. *J Biol Chem* **274**, 6285-6294,
doi:10.1074/jbc.274.10.6285 (1999).
60 Ziegler, S. F., Ramsdell, F. & Alderson, M. R. The activation antigen CD69. *Stem Cells* **12**, 456-465,
doi:10.1002/stem.5530120502 (1994).
61 Bretscher, M. S. Asymmetrical Lipid Bilayer Structure for Biological Membranes. *Nature New
Biology* **236**, 11-12, doi:10.1038/newbio236011a0 (1972).
62 Steck, T. L. & Lange, Y. Transverse distribution of plasma membrane bilayer cholesterol: Picking
sides. *Traffic* **19**, 750-760, doi:https://doi.org/10.1111/tra.12586 (2018).
63 Kiessling, V., Crane, J. M. & Tamm, L. K. Transbilayer Effects of Raft-Like Lipid Domains in
Asymmetric Planar Bilayers Measured by Single Molecule Tracking. *Biophysical Journal* **91**, 3313-
3326, doi:https://doi.org/10.1529/biophysj.106.091421 (2006).
64 Raghupathy, R. *et al.* Transbilayer lipid interactions mediate nanoclustering of lipid-anchored
proteins. *Cell* **161**, 581-594, doi:10.1016/j.cell.2015.03.048 (2015).
65 James, J. R. & Vale, R. D. Biophysical mechanism of T-cell receptor triggering in a reconstituted
system. *Nature* **487**, 64-69, doi:10.1038/nature11220 (2012).
66 Carbone, C. B. *et al.* In vitro reconstitution of T cell receptor-mediated segregation of the CD45
phosphatase. *Proceedings of the National Academy of Sciences* **114**, E9338-E9345,
doi:10.1073/pnas.1710358114 (2017).
67 Jung, Y. *et al.* Three-dimensional localization of T-cell receptors in relation to microvilli using a
combination of superresolution microscopies. *Proceedings of the National Academy of Sciences*
113, E5916-E5924, doi:10.1073/pnas.1605399113 (2016).
68 Rajendran, L. *et al.* Flotillins Are Involved in the Polarization of Primitive and Mature
Hematopoietic Cells. *PLOS ONE* **4**, e8290, doi:10.1371/journal.pone.0008290 (2009).

69   Dong, R. *et al.* DNA origami patterning of synthetic T cell receptors reveals spatial control of the
sensitivity and kinetics of signal activation. *Proceedings of the National Academy of Sciences*
118, e2109057118 (2021).
70   Douglas, S. M. *et al.* Rapid prototyping of 3D DNA-origami shapes with caDNAno. *Nucleic acids*
research **37**, 5001-5006 (2009).
71   Stahl, E., Martin, T. G., Praetorius, F. & Dietz, H. Facile and scalable preparation of pure and
dense DNA origami solutions. *Angewandte Chemie* **126**, 12949-12954 (2014).
72   Ohmann, A. *et al.* Controlling aggregation of cholesterol-modified DNA nanostructures. *Nucleic*
acids research **47**, 11441-11451 (2019).
73   Ellenberg, J. *et al.* Nuclear membrane dynamics and reassembly in living cells: targeting of an
inner nuclear membrane protein in interphase and mitosis. *J Cell Biol* **138**, 1193-1206,
doi:10.1083/jcb.138.6.1193 (1997).
74   Rodgers, J. L. & Nicewander, W. A. Thirteen Ways to Look at the Correlation Coefficient. *The*
American Statistician **42**, 59-66, doi:10.2307/2685263 (1988).
75   Manders, E. M. M., Verbeek, F. J. & Aten, J. A. Measurement of co-localization of objects in dual-
colour confocal images. *Journal of Microscopy* **169**, 375-382, doi:10.1111/j.1365-
2818.1993.tb03313.x (1993).

Methods

Synthesizing the cholesterol nano-patch (CNP)

The DNA nano-patch (NP) was designed on the honeycomb lattice with a long sing-stranded scaffold, type p8064 (Tilibit), by modifying the structure of R. Dong et al. [69] Sequence of 183 staple strands of the DNA NP was determined using the open-source program, caDNAno,[70] (Supplementary Table S1) and the corresponding staple strands were synthesized from Bioneer. A folding mixture consists of 20 nM of scaffold DNA, 100 nM of each staple strands, 20 mM $MgCl_2$ and 5 mM NaCl in TE buffer (5 mM Tris and 1 mM EDTA, pH 8). The annealing was performed following a temperature profile: 70 °C for 15 min, 60 to 20 °C by 1 °C/ho in a thermocycler. Excessive staple strands were removing by polyethylene glycol (PEG) precipitation [71] followed by two rounds of ultrafiltration with a 100 kDa cut-off filter. A concentration of the DNA NP was measured using UV spectrometer and adjusted using the working buffer (1×TE, 10 mM of $MgCl_2$, and 5 mM of NaCl). Purified samples were stored at 4 °C. To form the Chol-patch, we mixed the DNA NP with four-fold excessive assist-DNAs and Chol-DNAs to the total number of DNA handle on the DNA NP and incubated them at 30 °C for 12 h. At the 3' end of assist-DNAs, a six-nucleotides-thymine (6T) is appended to minimize undesired hydrophobic interaction between Chol-DNAs by wrapping the neighboring cholesterol [72]. To remove undesired hydrophobic interactions, Chol-DNA was heated at 65 °C for 1 h before use. The mixed solution was purified by ultrafiltration with a 100 kDa cut-off filter using phenol red free RPMI (Gibco) (buffer and its concentration was adjusted using the phenol red free RPMI.

Agarose gel electrophoresis

The assembled CNP was electrophoresed using 1% agarose gel containing 0.5×TBE (45 mM Tris-borate and 1 mM EDTA), 10 mM of $MgCl_2$ and 0.1 ul·ml^{-1} of SYBR Safe (Thermofisher). Electrophoresis was

performed for 90 min at 70 V bias voltage in a cold room. Agarose gel was scanned by a GelDoc and Image

Lab v5.1 program (Bio-Rad)

TEM Imaging

The concentration of CNP was diluted to 3 nM with the working buffer. The sample solution (5 μL) was

deposited on a glow-discharged formvar-supported carbon-coated grid for 3 min and stained by a 2% uranyl

formate and 25 mM NaOH for 40 sec. TEM images were taken at 40k-fold magnification using JEM-

2100Plus (JEOL).

Tissue culture

Jurkat T cells (Clone E6-1, ATCC) or Flot1-Halo transduced Jurkat T cells were grown in complete RPMI

(Gibco) media containing 2 mM L-glutamine, 100 U penicillin-streptomycin (PS) (Gibco) and 10% FBS

(Gibco). Human pan T cells were purified from frozen peripheral blood mononuclear cells (PBMCs,

STEMCELL Technologies, Cat #70025.2. Lot 201671701C) using a negative selection human Pan T Cell

Isolation Kit (Miltenyi Biotec) and mouse T cells were purified from mouse splenocytes using a negative

selection EasySep™ Mouse T Cell Isolation Kit (STEMCELL Technologies). Mouse splenocytes were

prepared from spleens collected from 8-18 weeks old male BL6 mice purchased from Orient bio. For each

experiment, a spleen was processed through a 40 μm filter (STEMCELL Technologies) and RBCs were

removed from the single cell suspension using RBC lysis buffer (eBioscience). Freshly isolated resting cells

for both human and mouse were used after incubation in complete T cell media, containing 10% (v/v) FBS,

100 U PS, 1x GlutaMAX supplement (Gibco), 1mM sodium pyruvate (Sigma Aldrich) and 1x nonessential

amino acids solution (Sigma Aldrich) in RPMI for ~2-4 h in a CO_2 (5%) incubator at 37 °C. For human

effector T cells, purified pan T cells were cultured with human anti-CD3/anti-CD28 dynabeads (Gibco) at

a 2:1 ratio in the complete T cell media supplementing with 50μM 2-mercaptoethanol and 100 U·mL^{-1} of

IL-2 upto a week.

Sample preparation for imaging

Jurkat or primary T cells were washed with calcium and magnesium free PBS (Gibco) by centrifugation at

200 g for 5 min (spin wash) and then resuspended in phenol red free RPMI. For the sequential reaction,

cells were spin washed and then incubated with 10 μM of Chol-DNAs for 10min at 37 °C. After spin

washing, 10 μM of DNA assist-DNAs was added and incubated for 10 min and then 10 nM of DNA NP

(containing 72-anchors) was added to the sample. The unbound DNAs are removed by a spin wash. For

the single step reaction, Chol- DNAs were complexed with DNA NP prior to use and then directly added

the CNP to spin washed cells at a final concentration of 10 nM in total 50 μl of phenol red free RPMI and

incubated in a CO_2 (5%) incubator for 30 min at 37 °C, otherwise, the incubation time was indicated. Cells

were spin washed with PBS, ready for using live cell imaging. For nuclei staining, cells were fixed with 4 %

(w/v) paraformaldehyde (PFA) (Electron Microscopy Sciences) in PBS for 15 min at Room temperature

(RT) and washed in PBS. Cells were stained with 10 uM of Hoechst 33342 (Thermo Fisher Scientific) for

5 min and then washed with PBS. For Flot1-Halo labeling, after washing cells with PBS, cells were stained

with Janelia Fluor (JF) 549 conjugated Halotag ligands (Promega) at 200 nM concentration in 2% FBS/PBS

at 37 °C for 15 min. The unbound Halo tags were spin washed with PBS and then cells were incubated with

CNP for 30 min, as described above. For CD45 labeling, the CNP labeled cells were washed and blocked

with 2% BSA in PBS for 10 min in the placed in a poly L- lysine (PLL) coated glass bottom chamber and

then labeled with 1ug·ml^{-1} of Alexa Flour (AF) 488 conjugated anti-human CD45 antibodies (Biolegend,

Cat# 304017, clone HI30) in 2% BSA in PBS for 20 min at RT. Cells were gently rinsed with PBS 4 times.

For CD3 labeling, the CNP labeled cells were spin washed and incubated with 10μg·ml^{-1} of AF488

conjugated anti-CD3 antibodies (Biolegend, Cat # 300415, clone UCHT1) in 2% BSA/PBS on ice for

30min and then spin washed. For imaging, 70~100 μl of freshly prepared cells resuspended in PBS were

PLL (0,01 %, Sigma Aldrich) or collagen (50 mg·ml^{-1}, Corning) coated glass-bottom chamber (35 mm dish,

7 mm glass, MatTek). The chambers were freshly prepared prior to use. For live time-lapse imaging,

Halotag labeled cells resuspended in phenol red free RPMI were placed on the PLL coated glass bottom

chamber and then 3~5 ul of CNP in RPMI were added and mixed to a final concentration of 3~5nM. Image

acquisitions were started 2 min after adding the CNP.

Plasmid construction and lentiviral transduction

The lentiviral pLV-FLOT1-Halo plasmid (gifted from Prof. Young-Wook Jun at the University of

California San Francisco) and two packaging plasmids pSPAX2 (Addgene) and pMD2.G (Addgene) were

cotransfected into Lenti-X293T cells (Takara) using lipofectamine 2000 (Thermo Fisher Scientific)

following the manufacturer's instructions. The supernatant was collected and filtered after 48 h using a 0.45

μm pore sized Millex PVDF syringe filter (Millipore). The filtered virus was concentrated by

ultracentrifugation at 10,000 g for 2 h and then applied to Jurkat T cells for two days. The transduced Jurkat

T cells were spin washed and maintained in complete RPMI media for three days and harvested for sorting.

The sorted Flot1-Halo transduced cells were cultured in complete T cell media for two days and maintained

in complete RPMI media.

Flow cytometry

The bindings of DNA NP with or without cholesterol modification on a single cell were analyzed using

flow cytometry. After 30 min incubation with DNA NP or CNP, cells were either washed or added 1 ml of

complete RPMI media and incubated for 4 h at 37 °C in a CO_2 (5%) incubator. The harvested cells incubated

for 30 min or 4 h were spin washed with PBS and stained with LIVE/DEAD™ Fixable Violet Dead Cell

Stain dye (L/D violet) (Invitrogen) diluted in PBS (1:1000) for 10 min at RT in dark. Cells were spin washed

and loaded on a Fusion cell sorter (BD Biosciences). Single cells were gated using forward vs. side scatter

(FSC vs. SSC) and then the live cells gated for the negative population of the L/D violet were analyzed for

the Cy5 fluorescence on CNP using FlowJo (BD Biosciences, versions 10.8.1) software. For CD69

expression analysis, cells were incubated without or with CNP, or PMA (10 ng·ml^{-1}) (Sigma Aldrich) and

ionomycin (0.5 µg·ml^{-1}) (Sigma Aldrich) in RPMI for 30 min at 37 °C, then 1 ml of complete RPMI was

added and incubated for 4 h at 37 °C in a CO_2 (5 %) incubator. Cells were harvested and spin washed with

PBS and then labeled with L/D violet for 10 min at RT. Cells were spin washed with 2% FBS/PBS and

then resuspended in 100 µl of 2 % FBS/PBS, followed by staining with Brilliant Violet (BV) 785 conjugated

anti-human CD69 (10 µg·ml^{-1}) (Biolegend, Cat # 310932, clone FN50) for 30 min on ice. Cells were spin

washed and loaded on a FACSAria Fusion cytometer (BD Biosciences). For compensation, ArC amine

reactive compensation bead (Invitrogen) stained with L/D violet and UltraComp eBeads (Invitrogen)

stained with AF 647 conjugated anti-IgG antibody for Cy5 or the BV785 conjugated CD69 antibody in

PBS were used. The live single cell, CD69 positive population was analyzed using the FlowJo software.

For cell proliferation tracking analysis, Jurkat T cells were harvested and spin washed in PBS and stained

with Cell Tracker Green CMFDA (1 µM) (Invitrogen) diluted in 2% FBS/PBS for 20 min at 37 °C in a CO_2

(5 %) incubator. After spin washing with 2 % FBS/PBS, cells were resuspended in RPMI and incubated

without or with 36- or 72-CNP at 10 nM for 30 min at 37 °C in a CO_2 (5 %) incubator. Cells were spin

washed (day 0 sample) or were grown in complete RPMI media for 5 days. Cells were harvested and

analyzed the diluted intensities of the CMFDA every day (day 0–day 4) by flow cytometry. After spin wash

the harvest cells, cells were stained with L/D violet in PBS (1:1000) for 10 min at RT. Cells were rinsed

with PBS and then loaded on the cytometer. The distribution of the intensities of the CMFDA on the live

Jurkat T cells was analyzed using the FlowJo software.

Fluorescence-activated single cell sorting (FACS)

For sorting, the Flot1-Halo transduced cells were spin washed with PBS and stained with L/D violet (1:1000)

for 10 min at RT. After spin washing, cells were stained with JF549 conjugated Halotag ligands (Promega)

at 200nM concentration in 2% FBS/PBS for 20 min at RT, followed by spin washing with 2 % FBS/PBS.

The cells were resuspended in 2 % FBS/PBS and kept on ice in the dark. Cells were immediately loaded

for sorting on a Fusion cell sorter (BD Bioscience). Live single cells were gated as described above and

then the JF549 labeled Flot1-Halo positive cells were analyzed and sorted using FACSDiva software (BD

Biosciences, version 8.0.2).

Confocal imaging

Confocal images were taken with a laser scanning Ti Microscope (Nikon) using an Apo 60 x Oil λS DIC

N2 objective lens (N.A., 1.4) (Nikon) with a 46~52 μm sized pinhole. 4 laser lines at 405, 488, 561, and

633 nm, and a combination of filter sets (dichroic: BS 20/80, emission bandpass: 450/50, 525/50, 595/50

and 700/75) were used for taking 9-20 series of z-plane confocal images with a step of 0.5 or 1 μm for the

Hoechst, AF488-, JF549- and Cy5-labeled samples, respectively. Bright field images were taken with a

transmission detector.

FRAP analysis

For FRAP experiments, time lapse confocal images were acquired using a Zeiss LSM 900 (Zeiss)

microscope, equipped with a C-Apochromat 40x Water Korr (N.A. 1.2) (Zeiss) objective lens with a 50 μm

sized pinhole. 561-, or 633-nm laser lines were used to excite, and bleach and emission fluorescence

between 400-635 nm and 645-700 nm were collected for the JF549-Flot1-Halo and Cy5-labeled CNP,

respectively. Time lapse series images were taken for 5 min with 3 s or 10 s intervals. After taking 3-5

frame images, a squared area of 0.5 μm^2 was selected and bleached for 3 s with 80 % of power for each

laser line. Drifting and motility of cells during the measurement were corrected that each frame image was

corrected to the next frame image except for the bleached frame using the 'imregtform' function (optimizer:

'rigid') in Matlab. The trace of the mean fluorescence intensities of the bleached area in the time-lapse

movie was analyzed using a custom-written Matlab (MathWorks, version R2018a) script. The diffusion

coefficient (D) for Flot1 was determined by analyzing the fluorescence recovery trace for 2.5 min after

bleaching using the following equation [73]: $I(t) = I_t = I_{final} (1 - (\omega^2 (\omega^2 + 4\pi Dt)^{-1})^{1/2}$, where I_t is

intensity as a function of time, I_{final} is final intensity reached after recovery, ω is width of the bleached

area, and D is one dimensional diffusion coefficient, which was determined by fitting using a built-in

function in Matlab.

Image analysis

ImageJ (version 1.52p) and Matlab (MathWorks, version R2018a) were used for image processing. For

counting the polarization population, each population of cells was counted from the maximum intensity

projection of 9-11 z-stack images using an ImageJ built in function, cell counter. The perimeters of the

single polarized clusters were manually measured from a z plane of the middle of each cell in the confocal

images using image J. For the colocalization test between channel X and channel Y, Pearson's correlation

coefficient [74] $\left(R' = \dfrac{\sum_{i=1}^{n}(X_i-\bar{X})(Y_i-\bar{Y})}{\sqrt{\sum_{i=1}^{n}(X_i-\bar{X})^2}\sqrt{\sum_{i=1}^{n}(Y_i-\bar{Y})^2}}\right.$, where n is the total number of pixels, X_i and Y_i are the pixel

intensities indexed with i. $\bar{X}$ and $\bar{Y}$ is the mean intensity of X and Y images, respectively) and Manders'

overlap   coefficient   [75]   (

$MOC = \dfrac{\sum_{i=1}^{n}(x_i)(y_i)}{\sqrt{\sum_{i=1}^{n}(x_i)^2}\sqrt{\sum_{i=1}^{n}(y_i)^2}}$, where n is the total number of pixels, x_i and y_i are the intensities of above-

threshold pixels indexed i in the X and Y images) were calculated using a custom built Matlab script. For

MOC, the first threshold values obtained using the 'multithresh' function in Matlab were used. For

quantifying the CD45 segregation, ratiometric intensity score (RIS) was calculated by following equation

[17] : $RIS = (X_i - Y_i\,(\bar{X}/\bar{Y}))/(X_i + Y_i(\bar{X}/\bar{Y}))$, where X_i and Y_i are the intensity of each pixel indexed i for

channel X (CD45) and channel Y (72-CNP); $\bar{X}$ and $\bar{Y}$ represent the mean of total X and Y, respectively. The

RIS and the mean intensities of CD45 for the raft and non-raft area were analyzed with data within the

segmented area presented in Fig. 4d. The original intensity values were used without subtracting

background for R', RIS, and CD45 intensity calculations. For the CD45 intensity plot in Fig. 4g, the

intensity of the segmented area for CD45 was normalized to 0-1 value for each cell and then it was

normalized to the mean of the non-raft intensities of all analyzed cells. The region of interest (ROI) for

induvial cells was selected by ImageJ or Matlab script. The raft and non-raft area of the CD45 and 72-CNP

dual-labeled images were segmented using a custom-written Matlab script. Briefly, channel 1 (CD45) and

channel 2 (72-CNP) images and the summed images of these two were binarized as BW1, BW2, and BW3

using the first threshold obtained from the 'multithresh' function in Matlab. The noise elements (less than

10 connected components) were removed from the BW3 using the 'bwconncomp' function. Next,

morphological closing was performed using the 'imclose' function on the nonzero element of the BW2 and

BW3 for the raft area, and on the nonzero element of the BW1 and BW3 for the non-raft area. Any element

that is less than 10 connected components was removed from the raft or non-raft areas using the

'bwconncomp' function. The nonzero element of raft or non-raft became the total segmented area. To

eliminate the boundary regions, any nonzero values of the raft area overlapped with the dilated non-raft

area using the 'imdilate' function with a disk structuring element with a radius of 10 pixels were removed.

The display ranges of the intensities of individual color channels in **Fig. 2**, **Fig. 3f**, **Supplementary Figs.**

2, 3, and 5 were independently optimized using ImageJ or Matlab. However, for comparison, the display

ranges of Cy5 intensity in Supplementary **Fig. 2b** were adjusted to that of **Fig. 2a**; the 0-, 18- and 36- CNP

images in **Fig. 2b** and **Supplementary Fig. 3** were adjusted to the 72-CNP image in **Fig. 2b**; Supplementary

Fig. 5a was adjusted to the **Fig. 2e**. The display ranges of the all figures in Supplementary **Fig. 6** and

Supplementary Movies 1-3 were equal and all other images in this study were displayed between the 1%

~ 99% intensity of individual channel image using a built-in function in Matlab.

Immunoblotting assay

Jurkat T cells were incubated with 10 nM of 72-CNP in RPMI for 30min at 37 °C. Cells were spin washed

with ice-cold PBS and lysed in lysis buffer containing 20 mM HEPES (pH 7.0), 1 % Triton X-100, 150

mM NaCl, 10 % glycerol, 1 mM EDTA, 2 mM EGTA, 1x Protease Inhibitor cocktail (Cell Signaling

Technology) and 1x phosphatase inhibitor cocktail (Cell Signaling Technology). Cell lysates were

centrifuged at 12,000 g for 10 min, and the protein concentrations were determined by Bradford assay (Bio-

Rad). Protein samples were mixed with SDS sample buffer containing 312.5 mM Tris (pH6.8), 10 % SDS,

30 % glycine, 1mg·ml^{-1} bromophenol blue and 1 M 2-mercaptoethanol, and boiled for 5 min. The proteins

were separated by SDS-PAGE and transferred to nitrocellulose membranes by electroblotting for 1 h 30

min. The membranes were blocked with 5 % dry skimmed milk in Tris-buffered saline containing 0.05 %

(v/v) Tween-20 (TBST) for 1 h and incubated with rabbit anti-phospho Zap-70(Tyr319)/ Syk (Tyr352)

(Cell Signaling Technology, Cat # 2717T, Clone 65E4), anti-β actin antibody (Cell Signaling Technology,

Cat # 4970, clone 13E5), or anti-α-tubulin antibody (Sigma Aldrich, Cat #T5168, Clone B-5-1-2) diluted

in the blocking buffer (1:1000) for overnight at 4 ℃. After washing 3 times with TBST, the membranes

were incubated with HRP conjugated anti-rabbit IgG secondary antibody (Cell Signaling Technology, Cat

#7074) diluted in blocking buffer (1:2000) for 1 h at RT and the washed with TBST 3 times. The immune-

reactive bands were detected using an enhanced chemiluminescence kit (Sigma Aldrich) and ChemiDoc

imaging system (Bio-Rad) and quantified with ImageJ software (version 1.52p).

Statistical information

At least three independent experiments were performed except **Supplementary Fig. 2 b-c, Fig. 4**, (two

independent experiments) and the statistical details for each experiment are indicated in the figure legends.

The p-values, mean, standard deviation (SD), and standard error of the mean (SE) were determined using

the built-in functions in Matlab (MathWorks, R2018a) or Prism 8 (GraphPad, version 8.0.1).

Data availability

The sequences for the DNA structures are listed in the Supplementary Table1. Data that are generated and

used in this study are available from the corresponding author upon reasonable request.

Code availability

Custom Matlab codes used in this study are available upon request to Y.J.

Acknowledgments

We thank Gilad Haran (Weizmann Institute of Science), Ronen Alon (Weizmann Institute of Science), and

Amnon Altman (La Jolla Institute for Immunology) for their kind advice and discussion for the manuscript.

This research was supported by Basic Science Research Program through the National Research Foundation

of Korea (NRF) funded by the Ministry of Education (2022R1I1A1A01070462) to Y.J.

Author contributions

Y.J. conceived and designed the experiments, Y.K. designed and synthesized the CNPs, Y.J., Y.K, S.L.

performed experiments. Y.J. analyzed data, J.N. performed transduction, S.H. performed manual

quantification. Y.J. Y.K. and M.K. wrote manuscript, M.K and J. C. co-supervised the project.

Competing interest declaration

The authors declare no competing interests.

Figures

Figure 1. DNA nano-patch (NP) displaying multiple cholesterols. a, A schematic drawing for the DNA-NP structure. Each cylinder represents a double-stranded DNA (dsDNA). Gray circles represent potential locations (anchors) where the DNA handles can be added for binding of Chol-DNAs. Cy5-conjugated ssDNAs were attached at the four corners. The DNA-NP is a square shape of 60 x 60 nm^2 with a 6 nm height. Each anchor is separated by 3.5 and 7 nm in rows and columns, respectively. **b,** Schematic illustration of the construction of cholesterol nano-patch (CNP). Chol-DNAs are complexed to the DNA handles on the DNA NP *via* DNA hybridization. Assist-DNA strands are added together to form a stable duplex form. **c-d,** Representative TEM images of DNA-NP without cholesterols (c) and CNP bound on the surface of small unilamellar vesicles (SUVs) (d). (Scale bars:100 nm). **e,** Schematic illustrations of the spatial arrangement of cholesterol molecules in 18-CNP, 36-CNP or 72-CNP. The positions for the cholesterols are marked with orange dots.

Figure 2. Synthetic lipid rafts on T cell membranes. a, Representative confocal images of Jurkat T cells incubated with 72-CNP for 30 min. Bright-field (BF) (gray, left), 72-CNP (magenta, middle), and the merged images (right) are shown. **b,** Effects of increasing the number of cholesterols on the formation of the synthetic lipid rafts. Representative confocal images of Jurkat T cells incubated with DNA NP without

cholesterol or complexed with 18- (18-CNP), 36- (36-CNP), or 72- (72-CNP) cholesterols for 30 min. Images of nuclei stained with Hoechst (blue, left), CNP (Magenta, middle left), and merged images (middle right) for each sample are presented. The orange dots on the schematic drawings (right) indicate the positions of cholesterols. **c,** The percentage of weak/unbound (light gray), unpolarized (gray), multi-polarized (dark gray), or single-polarized (black) synthetic lipid raft positive cells were counted from Jurkat T cells incubated with 18-(n=120), 36-(n=78), or 72-CNP (n=89) for 30 min (see the **Supplementary Fig. 2d-g**). **d,** Intensity distributions of the different CNPs on single cells incubated for 30 min (left) or 4 h (right) were analyzed by flow cytometry. **e,** Representative confocal images of freshly isolated mouse T cells (upper) or human T cells incubated with 72-CNP for 30 min. (scale bars in a, b and e are 20 µm).

Figure 3. CNP-mediated synthetic lipid rafts colocalized with Flot1-caps. a, Representative confocal images of a Jurkat T cell stably expressing Flot1-Halo incubated with 72-CNP for 30 min. Flot1-Halo labeled with Janelia 549 halotag ligand (green, left), 72-CNP (magenta, middle), and the merged images (right) are presented. **b,** Line profiles of the Flot1 (green) and 72-CNP (magenta) intensities along the blue dashed line in the merged image in **a**. **c,** Quantification of colocalization (Pearson's *R'* and *MOC*) between Flot1 and CNP calculated from 42 cells. Data are mean ± SD. **d-g,** FRAP analysis of the lateral mobility of Flot1 and 72-CNP bound on the membrane. Representative time-lapse confocal images of Flot1 (green) and 72-CNP (magenta) in the area marked by a square in the left images taken before and after bleaching (d). The mean FRAP tracings of Flot1 (green) and 72-CNP (magenta) after photobleaching for 5 minutes from live Jurkat T cells treated with 72-CNP. Data are mean ± SEM (n = 18 cells) (e). The filmstrip showing fluorescence recovery of the bleached area of the cell in **d** marked in **Supplementary Movie 5** (f). The mean diffusion coefficients (D) of Flot1 in untreated (left bar, n= 21) at the cap and in 72-CNP treated cells either at the cap (middle, n=26) or non-cap (right, n=14) areas analyzed from the FRAP (g). Each dot in **c** and **g** represents data collected from a single cell. Error bars are SD. *p*-values (*, $p \leq 0.05$; ***, $p \leq 0.001$, ****, $p \leq 0.0001$) were calculated by two-tailed unpaired t-test. (Scale bars in a and b: 5μm; in e: 1μm).

Figure 4. CD45 exclusion mediated by synthetic lipid raft. **a,** Representative confocal images of a Jurkat T cell incubated with 72-CNP for 30 min showing CD45 exclusion from a 72-CNP enriched synthetic lipid raft. CD45 labeled with AF488-conjugated antibody (green, left), 72-CNP (magenta, middle), and the merged images (right) are shown (scale bars: 5μm). **b,** Line profiles of the CD45 (green) and 72-CNP (magenta) intensities along the blue dashed line in **a**. **c,** Quantification of colocalization (Pearson's *R'* and *MOC*) between CD45 and 72-CNP. Each dot represents the data from a single cell (n = 51 cells). Data are mean ± SD. **d,** The segmented map for synthetic lipid raft (raft, yellow) and non-raft (light blue) regions of the cell displayed in **a**. **e,** Ratiometric intensity score (RIS) in each pixel of the entire segmented area presented in d. **f-g,** Paired analysis of the mean RIS values (f) and the mean normalized CD45 intensities (g) of pixels within the segmented areas of the raft vs. non-raft regions. Each grey line represents the data collected from a single cell (n = 51 cells) Data are mean ± SD.

Figure 5. Synthetic lipid rafts activate T cells. a, Representative confocal images of 72-CNP treated Jurkat T cells, showing accumulation of CD3 at 72-CNP induced synthetic lipid rafts. CD3 labeled with AF488-conjugated antibody (cyan, left), 72-CNP (magenta, middle), and the merged images (right) are shown (scale bars: 5μm). **b,** Line profiles of the intensities of the CD3 (cyan) and 72-CNP (magenta) along the red dashed line in the merged image in a. **c,** Quantification of colocalization (Pearson's *R'* and *MOC*) between CD3 and 72-CNP. Each dot represents the data from a single cell (n = 92 cells). Data are mean ± SD. **d-e,** Immunoblot analysis of phosphorylated ZAP-70 (p-ZAP-70) in Jurkat T cells incubated without or with 36- or 72-CNPs for 30 min. Tubulin levels represent the loading control. A representative image of immunoblotting (d) and quantification of p-ZAP-70 levels (e). The intensity of each p-ZAP-70 band relative

to respective tubulin or β-actin band was normalized to that of untreated samples. Each dot represents the band intensity quantified from an individual immunoblotting experiment. Data are mean ± SD. **f-g,** Flow cytometry analysis of CD69 expression induced by CNP. The increments of CD69+ population of cells in 72-CNP treated cells compared to untreated cells obtained from 6 independent experiments (f). A representative set of CD69+ cells in untreated, 36- or 72-CNPs, or in PMA and Ionomycin treated cells gated in a square for each (g). In **e** and **f**, p-values (*, $p \leq 0.05$; ***, $p \leq 0.001$) were calculated by the two-tailed ratio paired t-test.

656