Energy Coupling and Stoichiometry of Zn^{2+}/H^+ Antiport by the Cation Diffusion Facilitator YiiP

Adel Hussein[1]*, Shujie Fan[2]*, Maria Lopez-Redondo[1], Ian Kenney[2], Xihui Zhang[1], Oliver
Beckstein[§2], David L. Stokes[§1]

[1] Dept. of Cell Biology, NYU School of Medicine, New York, NY 10016 USA

[2] Dept. of Physics, Arizona State University, Tempe AZ

* A. Hussein and S. Fan contributed equally to this paper

§ O. Beckstein and D.L. Stokes serve as corresponding authors

ABSTRACT
YiiP is a prokaryotic Zn^{2+}/H^+ antiporter that serves as a model for the Cation Diffusion Facilitator
(CDF) superfamily, members of which are generally responsible for homeostasis of transition
metal ions. Previous studies of YiiP as well as related CDF transporters have established a
homodimeric architecture and the presence of three distinct Zn^{2+} binding sites named A, B, and
C. In this study, we use cryo-EM, microscale thermophoresis and molecular dynamics
simulations to address the structural and functional roles of individual sites and the interplay
between Zn^{2+} binding and protonation. Structural studies indicate that site C in the cytoplasmic
domain is primarily responsible for stabilizing the dimer and that site B at the cytoplasmic
membrane surface controls the structural transition from an inward facing conformation to an
occluded conformation. Binding data show that intramembrane site A, which is directly
responsible for transport, has a dramatic pH dependence consistent with coupling to the proton
motive force. A comprehensive thermodynamic model encompassing Zn^{2+} binding and
protonation states of individual residues indicates a transport stoichiometry of 1 Zn^{2+} to 2-3 H^+
depending on the external pH. This stoichiometry would be favorable in a physiological context,
allowing the cell to use the proton gradient as well as the membrane potential to drive the export
of Zn^{2+}.

INTRODUCTION

YiiP is a bacterial Zn^{2+}/H^+ antiporter and a well characterized representative of the Cation Diffusion Facilitator (CDF) superfamily. Members of this family play important roles in homeostasis of transition metal ions such as Zn^{2+}, Mn^{2+}, Co^{2+}, and Fe^{2+} (Montanini et al., 2007). Mn^{2+} transporters from this superfamily are prevalent in plants, where this metal ion is essential for oxygen generation by photosystem II as well as for a variety of other enzymatic functions (Alejandro et al., 2020). Zn^{2+} CDF transporters are widespread in organisms from all kingdoms of life reflecting a large influence of this ion on cell biology. An estimated 10% of proteins employ Zn^{2+} either as a catalytic co-factor (e.g., carbonic anhydrase and cytochrome C oxidase) or as a structural element (e.g., Zn-finger transcription factors and steroid receptors) (Maret, 2013). In mammals, high concentrations are found in a variety of intracellular vesicles that are key players in the immune system, synaptic transmission, insulin trafficking, and fertilization (Kambe et al., 2015). Zn^{2+} figures in host-pathogen interactions, with the host attempting to either deprive or poison pathogens residing in endosomes (Lonergan and Skaar, 2019). Despite its prevalence, the bulk of intracellular Zn^{2+} is bound to protein, with vanishingly small ($<10^{-12}$ M) concentrations of free Zn^{2+} in the cytoplasm (Liang et al., 2016; Outten and O'Halloran, 2001). Homeostasis is maintained by CDF transporters, which generally export Zn^{2+} from the cytoplasm, as well as members of the Zinc-regulated or Iron-regulated transport Proteins (ZIP), P-type ATPase and ATP-binding cassette (ABC) superfamilies (Yin et al., 2022).

YiiP serves as a model for studying structural and mechanistic properties of CDF transporters. Structures of homologs from both *E. coli* and *S. oneidensis* in different conformational states provide a framework for describing the archetypal alternating access mechanism (Lopez-Redondo et al., 2021; Lopez-Redondo et al., 2018; Lu et al., 2009; Lu and Fu, 2007). Molecular Dynamics has been used to study dynamics (Sala et al., 2019), to define the transport pathway (Sharma and Merz, 2022) and, together with in vitro binding assays, to characterize the K_d and pKa of Zn binding sites. Our understanding of CDF transporters is amplified by recent MD simulations of Znt2 (Golan et al., 2018; Golan et al., 2019), cryo-EM structures of Znt8 (Daniels et al., 2020; Xue et al., 2020; Zhang et al., 2023) and Znt7 (Han et al., 2022), as well as X-ray structures of isolated cytoplasmic domains from a number of different species (Udagedara et al., 2020). Together, these studies establish characteristics of Zn^{2+} binding sites, a conserved homodimeric architecture, and the nature of inward-facing (IF) and outward-facing (OF) states that presumably characterize all CDF transporters.

Despite this extensive work, there is uncertainty about key mechanistic questions that we have sought to address in the current work. Our primary goal was to obtain experimental evidence for the individual roles of the three Zn^{2+} binding sites found on each YiiP protomer (Cotrim et al., 2019). Site A is within the transmembrane domain (TMD) and features three Asp and one His residue; it is alternately exposed to the cytoplasm or to the periplasm in IF and OF states, respectively, and is thus directly responsible for transport of ions across the membrane. Although each molecule composing the dimer appear to have an independent transport pathway, there is unresolved potential for cooperativity. Site C within the C-terminal domain (CTD) is a binuclear site featuring four His and two Asp residues; one of these sites (C1) is well conserved whereas the other (C2) is not (Parsons et al., 2018). Site B is on the loop between transmembrane helices 2 and 3 (TM2-TM3) featuring two His and one Asp residue. Despite a lack of sequence homology for sites B and C2, the structure of Znt8 showed Zn ions at similar locations of the tertiary fold, but bound at different locations in the linear sequence, suggesting that these non-conserved, auxiliary sites may have functional significance (Xue et al., 2020).

Another goal was to address the role of protons in the transport process. There is ample
evidence supporting Zn^{2+}/H^+ antiport. In vitro assays with Znt1, Znt2, ZitB and CzcD showed pH
dependence of transport (Chao and Fu, 2004a; Cotrim et al., 2021; Guffanti et al., 2002;
Shusterman et al., 2014). Binding studies with Znt8 (Zhang et al., 2023) as well as
computational studies with Znt2 (Golan et al., 2019) and YiiP (Sharma and Merz, 2022) showed
an interplay between Zn^{2+} and H^+ binding at site A. However, there are discrepancies in
stoichiometry, with work on ZitB (Chao and Fu, 2004a) and YiiP (Chao and Fu, 2004b)
supporting a 1:1 exchange of Zn^{2+} and H^+, whereas studies of CzcD (Guffanti et al., 2002) and
Znt2 suggest 1:2 (Golan et al., 2019). There is also recent evidence for Zn^{2+}/Ca^{2+} antiport by
Znt1 in neurons (Gottesman et al., 2022).

For the current study, we have generated mutants of YiiP and have used cryo-EM and
microscale thermophoresis (MST) to measure the pH dependence and structural effects of Zn^{2+}
binding at each of the three sites. In addition, we have used molecular dynamics (MD)
simulations together with the experimental MST data to deduce the pKa of residues at these
sites and to address the stoichiometry of transport. From these data, we conclude that Zn^{2+}
binding at site C is responsible for the integrity of the homodimer. Release of Zn^{2+} from site B
triggers a conformational change in which the transport site A becomes occluded, suggesting a
potential relay of Zn^{2+} between these two sites. Occlusion of only one protomer breaks the
symmetry of the dimer and suggests that the transport process is not cooperative. In addition,
binding affinity at site A displays a dramatic pH dependence, which can be explained by the
protonation of 2 or 3 of the residues comprising this site. A corresponding Zn^{2+}/H^+ antiport
stoichiometry of 1:2 or 1:3, depending on pH, is consistent with energetic coupling of Zn^{2+} export
to the proton-motive force in a physiological setting.

 RESULTS

 Structural effects of Zn^{2+} binding

        In order to study effects of Zn^{2+} binding at individual sites, we produced mutants of YiiP
 from *Shewanella oneidensis*. The D51A mutation was introduced to preclude binding at site A
 and D70A to preclude binding at site B. For site C we mutated Asp287, because it bridges the
 two Zn^{2+} ions at that site and we anticipated that D287A would therefore eliminate both ions.
 The wild-type (WT) protein studied in previous work (Coudray et al., 2013; Lopez-Redondo et
 al., 2021; Lopez-Redondo et al., 2018) served as a positive control. For each of these
 constructs, protein was expressed in *E. coli*, solubilized in decyl-β-D-maltoside and purified by
 affinity chromatography. To ensure binding of Zn^{2+} rather than other metal ions picked up during
 growth and initial purification (e.g., Ni^{2+}), we incubated the protein overnight with chelators and
 then added 250 μM Zn_2SO_4 prior to the final purification step. As in previous work (Lopez-
 Redondo et al., 2021), we used an Fab antibody fragment to produce a larger complex
 amenable to cryo-EM analysis: 165 kDa comprising the YiiP homodimer (65 kDa) bound to two
 Fab's (50 kDa each).

 Site A: D51A mutation

        The structure of the D51A complex was solved at 3.6 Å resolution (Fig. 1b, Suppl. Fig. 1,
 Table 1) revealing an architecture very similar to untreated, WT YiiP (PDB code 7KZZ) that
 represents the IF, holo state (Lopez-Redondo et al., 2021). Specifically, YiiP formed a two-fold
 symmetric homodimer with Fab molecules bound near the C-terminus of each CTD. Density for
 Zn^{2+} ions is clearly visible at sites B and C, with the latter large enough to accommodate two
 ions along with additional water molecules (Suppl. Fig. 2). At site A, however, this structure
 lacked density, confirming that the D51A mutation effectively eliminated binding at this site.

        For comparison, we solved the structure from the WT construct that was similarly loaded
 with Zn^{2+}. As expected, this 3.8 Å resolution structure (Fig. 1a, Suppl. Fig. 1) is indistinguishable
 from 7KZZ, which is also from the WT construct but not treated with chelators or explicitly
 loaded with Zn^{2+} (RMSD 1.07 Å for all 562 Cα atoms), Despite overall similarity, comparison
 with the D51A structure showed movements of TM1, TM4 and TM5 (Fig. 2a), which resulted in
 a notably elevated RMSD for this region: 4.7 Å based on the corresponding 200 Cα atoms
 compared to 0.69 Å for the 352 residues composing the rest of the YiiP dimer. The cytoplasmic
 end of TM5 had the largest differences. Specifically, a kink is introduced near His155 and the
 TM4-TM5 loop is disordered, suggesting a flexibility of this region in the absence of Zn^{2+} at site
 A.

 Site B: D70A mutation

        An initial structure of the D70A construct revealed a global conformational change
 reminiscent of the previously solved WT, apo state (PDB 7KZX) produced by removal of Zn^{2+} by
 chelators (Lopez-Redondo et al., 2021). The most salient feature of this conformation is a kink
 between TMD and CTD, which disrupts the overall two-fold symmetry of the complex. During
 image processing, it became evident that there were in fact multiple conformations in the
 dataset from D70A, which ultimately generated two distinct structures at 4.0 and 3.9 Å
 resolution (Fig. 1c, Suppl. Fig. 3). Unlike the WT, apo structure (7KZX), both of these new
 structures revealed densities at sites A and C (Suppl. Fig. 2), consistent with Zn^{2+} binding to
 intact sites. However, the two structures differed both in the conformation of the TMD and in the
 orientation of the CTD. One of these structures, termed D70A_sym, has a TMD displaying local

Figure 1. Overview of the cryo-EM structures.

Density maps and corresponding atomic models are shown for each mutant. Fab molecules are shown in orange (light chain) and red (heavy chain) with YiiP colored in cyan, blue, purple and green, depending on the mutant. The homo-dimers adopt C2 symmetry for WT (A), D51A (B) and D287A (D) mutants, but a bend between TMD and CTD break this symmetry for D70A (C). The D287A/H263A mutant (E) forms a dimer of dimers in which the Fab molecules are rather disordered. Two conformations were observed for D70A, both of which are shown: D70A_sym on the left and D70A_asym on the right. Location Zn^{2+} binding sites, membrane boundaries as well as the topology of the YiiP protomer (inset) are shown in panel A; rectangles and arrows represent α-helices and β-sheets, respectively. Insets in panels B-D show the coordination geometry at the individual sites as seen in the WT structure.

Table 1. Structure determination of YiiP mutants

Data set	WT	Site A D51A	Site B[1] D70A asymTMD	Site B[1] D70A symTMD	Site C D287A	Site C[2] D287A/ H263A
Deposition						
PDB	8F6E	8F6F	8F6H	8F6I	8F6J	8F6K
EMDB	EMD-28881	EMD-28882	EMD-28883	EMD-28884	EMD-28885	EMD-28886
Data collection and processing						
Magnification	81,000	81,000	81,000	81,000	81,000	81,000
Voltage (kV)	300	300	300	300	300	300
Electron exposure ($e^-/Å^2$)	50	50	50	50	50	50
Defocus range (µm)	1.0-3.0	0.75-2.75	0.7-3.0	0.7-3.0	0.7-3.0	0.7-2.5
Pixel size (Å)	1.068	1.079	1.079	1.079	1.079	1.079
Symmetry imposed	C2	C2	C1	C1	C2	C2
Initial particle images (no.)	3,058,414	3,672,562	1,702,119	1,702,119	1,664,097	2,982,749
Final particle images (no.)	536,206	196,484	188,414	182,413	252,599	300,844
Map resolution	3.78	3.63	3.93	4.03	3.68	3.46
FSC threshold (Å)	0.143	0.143	0.143	0.143	0.143	0.143
B factor ($Å^2$)	200.5	151.0	146.1	126.0	150.9	135.2
Resolution range (Å)	3.0-5.5	3.0-5.2	3.4-5.8	3.4-6.9	3.0-5.5	3.0-5.2
Model Refinement						
Model composition						
Non-hydrogen atoms	10478	10348	10421	10377	10472	8714
Protein residues	1366	1344	1358	1351	1366	1132
Ligands	8	6	6	6	8	4
RMS deviations						
Bond lengths (Å)	0.003	0.003	0.002	0.002	0.003	0.002
Bond angles (°)	0.557	0.535	0.509	0.517	0.536	0.499
Validation						
MolProbity score	1.70	1.68	1.72	1.75	1.77	1.96
Clashscore	5.44	6.31	5.78	7.11	6.37	10.28
Rotamer outliers (%)	0.00	0.09	0.00	0.00	0.00	0.00
CaBLAM outliers (%)	2.66	3.05	2.60	3.40	3.64	4.21
Rama-Z score	-0.77	-0.37	-0.05	0.20	-0.50	-0.45
Ramachandran plot						
Favored (%)	94.19	95.27	93.99	94.86	93.59	93.51
Allowed (%)	5.44	4.57	5.86	4.99	6.04	5.96
Disallowed (%)	0.37	0.15	0.15	0.15	0.37	0.53
Model vs. Data CC (mask)	0.75	0.83	0.79	0.80	0.81	0.73

[1] Both D70A structures arose from the same set of micrographs and initial particle picks.
[2] Fab molecules were not included for the refinement of D287A/H263A due to poor density in this region of the map

Figure 2. Comparisons of atomic structures.

(A) Overlay of the TMD for WT (cyan) and D51A (blue) structures with a view prioritizing protomer A. Core helices (M2, M3, M6) align well but that there are substantial changes to the peripheral helices (M1, M4, M5), presumably due to lack of Zn^{2+} binding at the site bridging M2 and M5. (B) Overlay of the TMD for WT (cyan) and D70A_sym (purple) with a view along the two-fold axis shows a very similar configuration of helices and an intact dimeric interface. (C) Structure of D70A_asym viewed along the membrane plane showing the asymmetry between the two TMD's. The protomer on the left (chain B) adopts an IF conformation, whereas the protomer on the right (chain A) adopts a novel occluded conformation that includes a reconfigured TM2/TM3 loop (blue) making a novel interaction with the CTD. (D) Overlay of the TMD for WT (cyan) and D70A_asym (plum) showing the occluded conformation adopted by protomer A (top) and the IF conformation adopted by protomer B (bottom). L199 and L154 make van der Waals interactions that appear to stabilize the occluded conformation. (E-F) Overlays of D70A_asym (light purple) and D70A_sym (dark purple) structures after alignment of core helices TM3 and TM6. Structures are viewed parallel to the membrane plane in E and along the two-fold axis in F. A significant shift in the position of the CTD is apparent in E and bottom panel in F. Despite this shift and substantial conformational changes in the occluded protomer A, the dimer interface in the TMD (TM3 and TM6) is well aligned (F, top panel). (G) Domain swapped dimer of dimers adopted by the D287A/H263A construct. Dimerization of TMD's involves interaction between one dark-green and one light-green molecule (e.g., TMDa and TMDb, where "a" and "b" refer to chain ID's), whereas dimerization of CTD's involves interaction between either two dark-green or two light-green molecules (e.g., CTDa and CTDd). The linker between M6 and the CTD adopts a long straight helix in chains b and d, but remains an unstructured loop in chains a and c. (H) Overlay of the TMD for WT (cyan) and D287A/H263A structures viewed along the dimer axis shows a good match, indicating that disruption of site C affects mainly the configuration of the CTD.

two-fold symmetry, in which both protomers adopt the IF state (Fig. 2b). Like the WT, apo
structure, the TM2/TM3 loops, which harbor Asp70 and site B, are disordered. In the other
structure, termed D70A_asym, one TMD adopts the IF state with a disordered TM2/TM3 loop,
whereas the other TMD adopts a novel conformation in which the TM2/TM3 loop extends ~17 Å
away from the membrane to interact with H1 and β1 elements of the CTD from the opposing
protomer (Fig. 2c). In addition to reconfiguring this loop, there are movements in TM1,2,4,5 that
close off site A from the cytoplasm and thus generate an occluded state (Fig. 2d).

Despite these changes, local dimeric elements from the D70A structures superimpose
closely onto the corresponding elements from the WT, holo structure. In particular, alignment of
CTD's produces low RMSD's of 0.82 Å and 1.08 Å relative to D70A_sym and D70A_asym,
respectively (168 Cα atoms). Also, the core helices mediating the TMD dimer interface (TM3/6)
align closely with RMSD's of 0.79 Å and 1.22 Å for D70A_sym and D70A_asym (106 Cα atoms,
Fig. 2b, d). The entire TMD from the D70A_sym is generally consistent with the WT, holo
structure (Fig. 2b), though slight displacement of TM1,4,5 and bending of the cytoplasmic end of
TM2 lead to a somewhat elevated RMSD of 2.1 Å for 366 Cα atoms from both protomers after
aligning TM3/6. For D70A_asym, the TMD in the IF conformation has a similar RMSD of 2.6 Å
relative to WT, holo (183 Cα atoms from protomer B, Fig. 2d), whereas the occluded TMD has
an elevated RMSD of 5.8 Å (197 Cα atoms from protomer A after aligning TM3/6) due to
differences in the cytoplasmic ends of TM1,2,4,5.

The observed structural effects of the D70A mutation are broadly consistent with MD
simulations based on the WT structure. In particular, when site B was empty, the TM2/TM3 loop
exhibited greatly increased mobility documented by >2-fold increase in root mean square
fluctuations (RMSF, Fig. 3a); this result is consistent with disorder of this loop in D70A cryo-EM
structures. The CTD also exhibited increased mobility as documented both by the broader
distribution of the angles relative to the TMD (Fig. 3b) and by per residue RMSF, in particular
when the TMD was used as reference for alignment (Fig. 3c), indicating that the CTD moved as
a fairly rigid body relative to the TMD. In structures with Zn^{2+} bound at site B, the TM2/TM3 loop
from one protomer is close to the linker between TM6 and the CTD of the opposing protomer
(Fig. 3e,f, Suppl. Fig. 2). We therefore used the distance between Asp72 in the TM2/TM3 loop
and Arg210 in the TM6/CTD linker from the opposing chain as a collective variable for tracking
CTD movement. The corresponding distance distributions (Fig. 3d) show a prevalent peak at 2
Å from simulations of the WT, holo protein, whereas a broad distribution was obtained when site
B is empty. Together with the D70A_sym and WT, apo structures, this result suggests a salt
bridge in the holo state is disrupted and the TM2/TM3 loops become disordered when Zn^{2+} is
released from site B. However, the D70A_asym structure suggests that this disordering may be
transient, as the loop adopts the extended conformation and associates with different elements
of the CTD (Fig. 2c). Interestingly, this novel change in the TM2/TM3 loop is coupled with
changes in TM1,2,4,5 that close site A off from the cytoplasm (Fig. 3i-j). Mapping of electrostatic
charge shows that whereas the access channel to site A in the IF protomer is negatively
charged, positive charge dominates the cytoplasmic surface of the protomer in the occluded
conformation (Fig. 3k,l).

Site C: D287A and D287A/H263A mutations
The D287A mutant was chosen to disrupt site C since it coordinates both Zn^{2+} ions at
this site. However, the corresponding structure (3.7 Å resolution, Fig 1d, Suppl. Fig. 4) shows no
structural changes relative to the WT structure (1.18 Ä RMSD for all 562 Cα atoms) and Zn^{2+}-

A
RMSF (Å) vs Residue
— empty site B
— holo
M2/M3 loop
TMD   CTD

B
empty site B
CTD/TMD angle (°) vs time t (ns)
holo
md0
md1
md2
empty site B
holo

C
empty site B
RMSF (Å) vs Residue
alignment
All
TMD
CTD
TMD   CTD

D
Probability vs Distance D72-R210 (Å)
apo
empty site B
holo

E WT
TMD
CTD

G D70A_sym

I D70A_asym
IF
OCC
(L)

K D70A_asym
IF
OCC

F
M6/CTD
M2/M3

H

J
IF
OCC
(L)

L
CTD
OCC
M2/M3
-5.0   +5.0

Figure 3. Zn^{2+} removal from site B in the TM2/TM3 loop.

(A) Per-residue RMSF of the WT, holo structure with Zn^{2+} present (cyan) and absent (purple) at site B demonstrates a notable increase in fluctuations in the TM2/TM3 loop. The dashed line demarcates the boundary between TMD and CTD. (B) Angle between the TMD and CTD in three individual simulations in the absence (purple) and presence (cyan) of Zn^{2+} at site B. The distribution of angles, on the right, highlight greater mobility when site B is empty. (C) Per-residue RMSF for simulations with empty site B using three different alignment schemes: the entire molecule ("All"), transmembrane domain ("TMD"), or C-terminal domain ("CTD"). Analogous data for the WT, holo structure have previously been published (Fig. 3i in Lopez-Redondo et al., 2021). (D) Distributions of distance between Asp72 in one chain and Arg210 in the opposite chain during simulations of the WT, holo structure, the apo structure with Zn^{2+} absent from all sites, and the structure with site B empty. The sharp peak at ~2 Å from the holo structure suggests a salt bridge that is disrupted in other structures. (E,F) Structure of the WT, holo YiiP dimer showing global C2 symmetry about a vertical axis and juxtaposition of the TM2/3 loop (blue) with the TM6/CTD linker from the opposing protomer (orange-to-red). (G,H) D70A_sym structure shows a kink between TMD and CTD and disordering of the TM2/3 loop. Both protomers are in the IF state. (I,J) D70A-asym structure showing further twisting of the CTD and asymmetry of the TMD's. The TM2/3 loop is disordered in the protomer on the left (chain B), but adopts a novel interaction with the CTD in the protomer on the right (chain A). (K,L). Electrostatic surface of D70A_asym showing a negatively charged cavity leading to site A on the left, but an occluded cavity with positive charge on the right. Rainbow colors progress from blue to red moving from the N-terminus of one protomer to the C-terminus of the other protomer. Note that L is at an oblique angle looking down on the M2-M3 loop.

related densities are still clearly visible at site C (Suppl. Fig. 2). We therefore introduced a
second mutation, D287A/H263A, and the corresponding particles had an unusual appearance
consistent with formation of higher order oligomers (Suppl. Fig. 4b). This observation suggested
that site C may be responsible for the integrity of the homodimer. Indeed, elution profiles from
SEC show that the main peak from D287A/H263A is shifted from 11.1 to 10.1 ml relative to the
other mutants (c.f., Suppl. Figs. 1a,b, 3a, 4a,b). Elution profiles from both D70A and D287A
have an earlier peak at ~10 ml and image processing in both cases revealed a relatively small
subpopulation of particles forming a dimer of dimers (Suppl. Fig. 3c & 4e). For D287A/H263A,
this dimer of dimers represents the main peak and the only class of particles that could be
refined to high resolution (3.5 Å) (Suppl. Fig. 4h). In this case, higher order oligomers (Suppl.
Fig. 4f) explain the earlier elution peak at 8-9 ml.

The dimer of dimers from D287A/H263A is stabilized by an unprecedented domain swap
in which CTD's and TMD's affiliate with different protomers (Fig. 2g). This domain swap is
enabled by a structural transition of the linker between the TMD and the CTD. In particular, the
normally unstructured loop between TM6 and the first helix of the CTD (H1) is reconfigured into
a very long, continuous α-helix extending from the N-terminus of M6 (Trp178) to the C-terminus
of H1 (Glu226). Site C itself is completely disordered and the CTD's are splayed apart (Suppl.
Figs. 2g & 5c): i.e., the distance between Cα atoms of Arg237 and Glu281 is 22.7 Å compared
to ~12 Å in structures with an intact site C (Lopez-Redondo et al., 2021). This domain swap was
not observed in the dimer-of-dimers formed by D70A and D287A, where the CTD and site C
were unperturbed and Fab molecules were uniquely responsible for inter-dimer interactions.
Although it was not possible to refine a structure from the higher-order oligomers from
D287A/H263A, due to heterogeneity and preferred orientation, they appeared to comprise a
linear chain of molecules propagated via this domain swap. These surprising changes in the
CTD indicate that site C is indeed crucial in maintaining the integrity of the native homo-dimer.
Despite the domain swap, however, TMD's from the D287A/H263A complex are quite congruent
with the WT structure (RMSD 1.35 Å for 346 Cα atoms excluding the TM2/TM3 loop) and
density for Zn^{2+} ions is clearly visible at site A (Suppl. Fig. 2). Weak density is visible for the
TM2/TM3 loops, indicating that they adopt an extended configuration that disrupts site B and
allows them to interact with the CTD; it does not appear that Zn^{2+} is bound at site B in this
domain-swapped complex (Suppl. Fig. 2).

Zn^{2+} binding affinity

To assess the binding affinity of individual Zn^{2+} binding sites, we used MST to analyze a
series of mutants designed to isolate the individual sites. Specifically, the triple mutant
D70A/D287A/H263A was used to study site A, D51A/D287A/H263A to study site B, and
D51A/D70A to study site C. All of these mutants eluted from SEC columns at the same volume
as WT (Suppl. Fig. 6) indicating that they all adopted the native dimeric structure. This
observation suggests that the domain-swapped dimer of dimers and higher order oligomers
seen by cryo-EM were induced by the Fab molecules. To maintain accurate and reproducible
Zn^{2+} concentrations ranging from nanomolar to micromolar we used nitrilotriacetic acid (NTA, K_d
= 14 nM) to buffer Zn^{2+} for sites with high affinity, and citrate (K_d = 12 μM) for sites with lower
affinity. At pH 7, sites A and C displayed relatively high affinity (K_d = 16 and 33 nM, respectively,
Table 2, Fig. 4), whereas site B had considerably lower affinity (K_d = 1.2 μM). Site C is
binuclear, and the apparent affinity was reduced when further mutations were introduced to
isolate individual C1 (D51A/D70A/H263A with K_d = 153 nM) and C2 (D51A/D70A/H234A with K_d

Figure 4. pH dependence of Zn²⁺ binding affinity.

MST was used to experimentally determine Zn^{2+} binding affinity at each site (panels A, B and C). These data were combined with a thermodynamic model, represented in panel H, with parameters derived either from CpHMD (panels D and E) or MST inference (panels F and G). (A) For MST studies of site A, the D70A/D287A/H263A construct was used with Zn^{2+} buffered either with NTA (pH 7.0 and 7.4) or with citrate (pH 5.6, 6.0 and 6.5). Curves represent the law of mass action with K_d values listed in Table 3. The relatively poor fit at pH 7.4 may reflect the fact that the K_d (1 nM) is lower than the minimum protein concentration (8 nM) supported by the assay. (B). For MST studies of site B, the D51A/D287A/H263A construct was used and Zn^{2+} was buffered with citrate. (C) For MST studies of site C, the D51A/D70A construct was used and Zn^{2+} was buffered with NTA at pH 7 or with citrate at the other pH's. (D,E) Curves represent predictions of a thermodynamic model with pKa values for either site A residues (Asp47, Asp51, His153, Asp157) or site B residues (D70, H73, H77) taken from CpHMD simulations. These pKa values are listed in Table 4 and the distributions of protonation states are shown in Suppl. Figs. 10 and 11. Symbols represent experimental MST data as in A and illustrate a poor fit using these parameters. Thermodynamic modeling was not possible for site C because of instability of CpHMD simulations in the absence of Zn^{2+}. (F,G). Curves represent predictions of the MST inference algorithm with corresponding pK_a values listed in Table 4. Again, symbols represent experimental MST data and show the excellent fit using these refined parameters. (H) Schematic representation of the microscopic thermodynamic model for site A. Titratable residues (D47, D51, D159, H155) are represented either as black squares for the protonated state, or white squares for the deprotonated state. Binding of one Zn^{2+} ion to the site is indicated by a filled magenta circle. Transitions are only possible between states connected by lines (protonation/deprotonation) or corresponding states connected by double-sided arrows (Zn^{2+} binding/release). An analogous model for site B comprised D70, H73 and H77.

= 223 nM) sites, suggesting cooperative binding of ions at C1 and C2 as implied by the
coordination geometry of this site (Suppl. Fig. 1d inset).

To explore the basis for coupling of Zn^{2+} transport to the proton-motive force, we
measured Zn^{2+} binding affinity at pH values from 5.6 - 7.4. We found that affinity at site A
changed by five orders of magnitude: K_d ranging from 1 nM at pH 7.4 - consistent with the
cytosol - to 302 μM at pH 5.6 (Fig. 4, Table 2). Site B had only modest pH dependence ranging
from 1-16.6 μM, whereas Site C changed by two orders of magnitude from 0.033 - 6.7 μM.
Cooperativity at site C is consistent with a high Hill coefficient at pH 7 (n=2.9, Table 3)), which
fell below 1 at lower pH's. However, Hill coefficients obtained for other sites were quite variable
(Table 3), including those for sites C1 and C2, making conclusions about cooperativity
inconclusive.

We also used MD simulations in conjunction with the experimental MST data to address
pH dependence of Zn^{2+} binding and to evaluate the contribution of individual residues. To start,
we estimated pKa values for each titratable residue at site A (Asp47, Asp51, His155 and
Asp159) and site B (Asp70, His73 and His 77) using constant pH MD (CpHMD) simulations with
pH replica exchange in the absence of Zn^{2+}. These simulations generated probabilities for each
protonation state as a function of pH; these probabilities were fit with the Hill-Langmuir equation
to derive per-residue pK_a's (Suppl. Figs. 7-9, Table 4). For site A residues, His155 had the
highest pK_a (8.0), followed by Asp159 (4.9), Asp47 (3.8) and Asp51 (2.9). Predicted populations
of the various microstates indicated that two residues are likely protonated at pH 5.5 (Suppl.
Figs. 10a & b) with His155 getting protonated first followed by Asp159. Although these
simulations were conducted with the IF conformation, we do not expect significant differences
for the OF state, given the similarity of site A seen in the X-ray structure (Lu et al., 2009) of the
OF state (RMSD of 0.49 Å for 17 Ca atoms on TM2 and TM5 and 1.01 Å for all atoms
composing the four residues that coordinate the Zn^{2+} ion). For site B, His73 and His77 have
almost identical pK_a's of 8.1, and the dominant state at neutral pH consists of both His73 and
His77 protonated (Suppl. Figs. 11a & b). This analysis of sites A and B applies to the absence
of Zn^{2+}; however, it was not possible obtain data about site C because the Zn^{2+}-free CTD proved
unstable in CpHMD simulations.

We went on to use a thermodynamic model incorporating both protonated and Zn^{2+}-
bound states to assess the stoichiometry of transport (Fig. 4h). We used this model to generate
Zn^{2+} binding curves based on pK_a's of individual residues and binding free energies for Zn^2
derived from the experimental MST data (Table 2). Initially, pK_a's of each residue in the absence
of Zn^{2+} were taken from the Hill-Langmuir fits to the CpHMD data (Table 4, Suppll Fig. 9). The
CpHMD-based model did not agree with the experimental data (Figs. 4b & e), likely due to
inaccuracy in microstate pK_a values. We therefore employed a novel modeling method based on
the inverse *Multibind* approach (see Methods) to iteratively refine the CpHMD pK_a's and the
MST Zn^{2+} binding free energies using the MST data points as a target for Monte Carlo (MC)
minimization. This analysis, referred to henceforth as MST inference, allowed us to estimate
probabilities of all the microstates as a function of pH and Zn^{2+} concentration (Suppl. Figs. 10 &
11). After refinement of parameters by MST inference, the predicted Zn^{2+} binding curves closely
reproduced the observed pH-dependence of Zn^{2+} binding at both sites A and B (Figs. 4c & f).
The dramatically improved agreement is a consequence of substantial changes in the pK_a's of
individual residues (Table 4).

Table 2. Binding affinity for individual Zn^{2+} sites measured by MST* or deduced by the MST inference algorithm[§]

	site A D70A/D287A/H263A	site B D51A/D287A/H263A	site C D51A/D70A	site C1 D51A/D70A/H263A	site C2 D51A/D70A/H234A
pH 5.6	302 ± 107 µM (149 µM)	27.1 ± 2.8 µM (19.5 µM)	8.10 ± 1.55 µM		
pH 6.0	6.11 ± 1.32 µM (10.8 µM)	5.54 ± 2.0 µM (5.38 µM)	0.692 ± 0.28 µM		
pH 6.5	0.654 ± 0.173 µM (0.503 µM)	5.73 ± 1.75 µM (1.36 µM)	0.088 ± 0.032 µM		
pH 7.0	0.0163 ± 0.0041 µM (0.0325 µM)	1.18 ± 0.6 µM (0.503 µM)	0.033 ± 0.0087 µM	0.153 ± 0.0479 µM	0.223 ± 0.039 µM
pH 7.4	0.001 ± 0.001 µM (0.0048 µM)				

* Values correspond to K_d as determined by applying the law of mass action to the data.
[§] Values deduced by MST inference are shown in parentheses

Table 3. Binding affinity and Hill coefficients derived from MST data*

	site A D70A/D287A/H263A	site B D51A/D287A/H263A	site C D51A/D70A	site C1 D51A/D70A/H263A	site C2 D51A/D70A/H234A
pH 5.6	1490 ± 2460 µM n=0.46	28.4 ± 3.54 µM n=0.94	7.50 ± 2.20 µM n=0.9		
pH 6.0	6.34 ± 1.67 µM n=0.81	5.68 ± 1.57 µM n=1.60	1.39 ± 0.382 µM n=0.78		
pH 6.5	0.622 ± 0.198 µM n=0.54	27.9 ± 34.3 µM n=0.38	0.133 ± 0.0359 µM n=0.66		
pH 7.0	0.0247 ± 0.0057 µM n=1.0	4.23 ± 13.0 µM n=0.45	0.0486 ± 0.0038 µM n=2.9	0.116 ± 0.0119 µM n=2.96	0.212 ± 0.0328 µM n=1.2
pH 7.4	0.0039 ± 0.0009 µM n=0.56				

* Values correspond to EC_{50} and the Hill coefficient (n) as determined by fitting the Hill equation to the data.

Table 4. pK_a values of residues determined by CpHMD simulations (Hill-Langmuir) and MST inference

		CpHMD (Hill-Langmuir)	MST inference
Site A	Asp47	3.84 ± 0.18	-0.20 ± 0.51
	Asp51	2.92 ± 0.12	6.51 ± 0.21
	His155	7.97 ± 0.15	7.82 ± 0.15
	Asp159	4.87 ± 0.18	7.83 ± 0.15
Site B	Asp70	2.06 ± 0.46	1.12 ± 0.57
	His73*	8.08 ± 0.46	12.37 ± 0.61*
	His77*	8.12 ± 0.13	5.26 ± 0.47*

* Titration curves for His73 and His77 from the MST inference model cannot be fit individually with a simple Hill equation (c.f. Suppl. Figs. 11e & f). We therefore consider these two residues as a coupled system with two effective pK_a values as shown.

274   A more detailed look at the population of microstates predicted by MST inference allows
275 us to explore the interplay of Zn^{2+} and H^+ binding under physiological conditions. For site A in
276 the absence of Zn^{2+}, the dominant state at cytosolic pH of 7.5 consists of 2 protons bound by
277 His155 and Asp159 (Suppl. Figs. 10c & d). Unexpectedly, the model predicts that these two
278 residues form a coupled system in which protonation is highly cooperative with an effective pK_a
279 of 7.8 for both residues (Suppl. Fig 10e) and with the single-proton state largely suppressed
280 (Suppl. Fig 10d). At the lower pH of the periplasm, a third proton is recruited by Asp51, whose
281 pK_a increased to 6.5 during refinement by MST inference (Table 4). Regardless of the pH, as
282 Zn^{2+} binds at site A, all the residues are deprotonated (Suppl. Fig. 10f), supporting the idea that
283 Zn^{2+} transport can be coupled with a proton gradient across the plasma membrane. For site B,
284 the dominant state at cytosolic pH in the absence of Zn^{2+} has one protonated residue: either
285 His73 or His77 (Suppl. Fig. 11c & d). Although the microstate pK_a's are equivalent, these site B
286 residues behave quite differently from Site A residues and generate an anti-cooperative binding
287 curve with effective pK_a's of 5.6 and 12.4 (Suppl. Figs. 11e & f). Zn^{2+} binding at cytosolic pH
288 displaces all protons (Suppl. Fig. 11g). Even though protons may co-exist with Zn^{2+} at lower
289 pH's, this is not a condition that would be physiologically relevant given that site B is uniquely
290 exposed to the cytoplasm.

DISCUSSION

In this study, we have focused on the properties of individual Zn^{2+} binding sites of YiiP from *S. oneidensis*. Previous studies have shown that this protein forms a homodimer in the IF state with Zn^{2+} ions constitutively bound at three sites (Lopez-Redondo et al., 2018). Treatment with EDTA to remove Zn^{2+} from all three sites induced a conformational change to produce an occluded state, a necessary precursor to the OF state (Lopez-Redondo et al., 2021). For the current work, we generated mutations at each site to measure their binding affinities and to assess their respective roles in generating conformational change. We found that site A, considered to be the transport site in the middle of the TMD, has nanomolar affinity at a cytoplasmic pH of 7.4, which is reduced 10^5-fold at the more acidic pH of 5.6. Release of Zn^{2+} from site A induces only modest movement of the peripheral transmembrane helices, but has no global effect on the architecture of the homodimer. Site B, on the short loop between TM2 and TM3, has much lower affinity in the micromolar range with relatively little pH dependence. When site B lacks Zn^{2+}, the loop becomes disordered and the protein undergoes a global conformational change leading to an occluded conformation in one of the protomers. Site C, a binuclear site in the CTD that engages residues from both protomers, has relatively high affinity and intermediate pH dependence. When site C was fully disrupted with a double mutation, the homodimer became destabilized such that the Fab molecules used for cryo-EM imaging induced a domain swap of the CTD leading to higher order oligomers.

Elements stabilizing the homodimer

Dimerization appears to be a universal feature of Cation Diffusion Facilitators. All structures so far show similar elements contributing to a common dimer interface (Suppl. Fig. 12). In particular, a conserved salt bridge exists at the cytoplasmic surface of the membrane (Lys79-Asp209 for soYiiP); the cytoplasmic end of the CTD generally includes one or more Zn^{2+} sites, and extensive hydrophobic interactions involving TM3 helices mediate TMD interactions (Lopez-Redondo et al., 2021). In early work, the salt bridge was proposed to act as a fulcrum for alternating, scissor-like movements in TMD and CTD driven by relay of Zn^{2+} between sites A and C (Lu et al., 2009). Furthermore, the original X-ray structure of ecYiiP featured a V-shaped architecture in which TMD's were completely disengaged (Lu et al., 2009; Lu and Fu, 2007). However, subsequent work showed that stabilization of TMD interactions by cysteine crosslinking did not inhibit transport activity (Lopez-Redondo et al., 2018) and compact dimeric TMD interfaces were observed in cryo-EM structures of soYiiP, Znt8 and Znt7 (Han et al., 2022; Lopez-Redondo et al., 2018; Xue et al., 2020). Indeed, the TMD interface is a remarkably conserved feature in our new structures of D70A and D287A/H263A mutants, despite large-scale conformational changes involving the TM2/TM3 loop and domain swapping of the CTD. We therefore conclude it to be unlikely that the TMD's undergo large-scale scissoring movements as part of the transport cycle.

For isolated CTD's, metal ion dependent scissor-like movements have been well documented using truncated constructs from a variety of species (Cherezov et al., 2008; Udagedara et al., 2020; Zeytuni et al., 2014). We observed similar movements in full-length soYiiP: i.e., the CTD's move apart when Zn^{2+} is removed from site C either by chelation or mutation. However, other elements must contribute CTD dimerization, given that isolated CTD's dimerize even in the absence of metal ions and Zn^{2+} ions were not observed at all in the CTD of Znt7 (Han et al., 2022). Nevertheless, the Fab-induced domain swap seen in the D287A/H263A double mutant indicates that, for soYiiP, CTD interactions are weakened in the absence of Zn^{2+}.

Conversely, it seems likely that dimer stabilization, via TMD and salt-bridge interactions, lead to
enhanced Zn^{2+} binding in the CTD, as evidenced by the considerably higher Zn^{2+} affinity at site
C of full-length soYiiP (K_d of 33 nM at pH 7) compared to those measured from isolated CTD's
(K_d in micromolar range (Udagedara et al., 2020; Zeytuni et al., 2014)). In any case, the
relatively high affinity of site C at pH 7 suggests that this site will remain occupied and will serve
to ensure dimer stability under physiological conditions.

Role of the site B

Structural effects of the D70A mutation support our previous hypothesis that site B is
responsible for inducing a global conformational change. The most conspicuous change is the
kink between TMD and CTD, which breaks the global two-fold symmetry of structures that retain
Zn^{2+} binding at site B, in which local two-fold axes of CTD and TMD are aligned (Suppl. Fig. 12).
Although stable, kinked conformations were not achieved over the time-scale of MD simulations,
increased movement of the CTD relative to the TMD is seen both in the previously reported apo
state simulations (Lopez-Redondo et al., 2021) and with an empty site B (Fig. 3), indicating a
role of the Zn^{2+}-bound TM2/TM3 loop in stabilizing the position of the CTD. Indeed, van der
Waals interactions are observed between the structured loop from one protomer and the TM6-
CTD linker of the opposing protomer on both sides of the dimer (Fig. 3e,f), and MD simulations
indicate that a salt bridge between Asp72 and Arg210 may reinforce these interactions (Fig. 3d,
c.f., site B in Suppl. Fig. 2).

Both cryo-EM structures and MD simulations show that the TM2/TM3 loop becomes
disordered when Zn^{2+} is released from site B and this is a likely explanation for increased
mobility of the CTD. Two structures were obtained from the D70A mutation in which the CTD
becomes progressively more tilted and twisted. In one structure, the TMD's are symmetrical and
retain the IF conformation (D70A_sym), whereas in the other structure with more extreme CTD
movement (D70A_asym), the TMD transitions to an occluded state. We speculate that these
structures represent a sequence of conformational change leading to the occluded state.
Although both TM2/TM3 loops are disordered in the D70A_sym structure, one of these loops
reforms into an extended conformation in the occluded protomer of the D70A_asym structure
(Suppl. Fig. 5). This change results in a novel interaction between the loop and the CTD, which
may drive the displacement and bending of membrane helices thus leading to occlusion of the
transport site A. On the other side of the dimer, the CTD has lost contact with the TMD, allowing
the TM2/TM3 loop to remain disordered and the TMD to remain in the IF state. Interestingly, the
TM2/TM3 loops are also extended in the D287A/H263A domain-swapped structure (Suppl. Fig.
5), but in this case, the interactions of this loop with the reconfigured CTD's are quite different
and the TMD's remain in the IF state, indicating that a specific structural constraint is required to
instigate occlusion.

Occlusion of the transport site A is a result of bending of TM5 and tilting of TM1 and 4,
thus narrowing the gap between these helices and closing off access to the cytosol. These
movements also generate a dramatic difference in the electrostatic surface at the cytoplasmic
side of the TMD. In the IF state, the open cavity leading toward site A is negatively charged, as
also shown for Znt7 (Han et al., 2022), thus serving to attract Zn^{2+} toward this site. After
transition to the occluded state, not only is the cavity closed, but the surface becomes positively
charged thus repelling Zn^{2+} (Fig. 3k,l). Previous work has identified a "hydrophobic gate"
consisting of two residues at the cytoplasmic ends of TM5 and TM6: Leu154 and Leu199 in
soYiiP (Gupta et al., 2014). Our previous comparison of holo and apo states showed that these

two residues do come closer together in the occluded state (Lopez-Redondo et al., 2021) and
homologous residues in Znt7 are close in the OF state and separated in the IF state (Han et al.,
2022). Interaction of Leu154 and Leu199 is also seen in the occluded protomer in the
D70A_asym structure (Fig. 2d), but these residues are at the periphery of the cavity leading to
site A. This observation suggests that interaction of these residues may play an important role in
stabilizing this state, but that bending at the cytoplasmic end of TM5 (Val148-Ala151) and tilting
of TM1 and TM4 may be more directly responsible for occlusion of the transport site.

For soYiiP, the IF state appears to be a low-energy, ground state when Zn^{2+} is present at
all three sites. Indeed, the IF state has been seen for WT protein both in lipid-based helical
crystals (Lopez-Redondo et al., 2018) and detergent micelles (Lopez-Redondo et al., 2021), as
well as in D287A and D287A/H263A mutant structures, all with RMSD's <1.5 Å. Removal of
Zn^{2+} from site B can be viewed as a source of energy for inducing conformational change and
likely leading to transport. Interestingly, the occlusion only occurs in one protomer, suggesting a
lack of synchrony across the dimer axis. A mixture of conformational states was also observed
for Znt8, in which one protomer adopted the OF state while the other was in the IF state. These
observations imply that a given protomer may be able to undergo transport while the other
remains inactive, as illustrated in Fig. 5.

The proposed sequence of conformational change is consistent with the notion that Zn^{2+}
is initially recruited to site B and that transfer to site A would act as a trigger for the first step of
the transport cycle. Such a two-step mechanism might be necessary if Zn^{2+} were delivered to
YiiP as a chelate – e.g., with metallothionein or glutathione – that would not have direct access
to site A. In the case of Cu^+, metallochaperones are thought to play a role in delivery, but so far
there is very limited evidence for Zn^{2+} metallochaperones (Chandrangsu et al., 2019).
Nevertheless, essentially all of the Zn^{2+} in the cytoplasm exists in chelated form with vanishingly
small concentrations of free Zn^{2+} available for direct binding (Choi and Koh, 1998; Outten and
O'Halloran, 2001). The accessibility of site B makes it a plausible acceptor as an initial
recruitment site. Once bound at site B, two features would facilitate transfer to site A: the drastic
difference in affinity between sites A and B would generate a large free-energy gradient and the
negative charge of the cavity leading to site A, which lies almost directly below site B in the WT,
holo structures, would direct the ion toward the transport site. Although the amino acid
sequence of site B is not conserved amongst CDF transporters, a Zn^{2+} ion was observed in an
analogous juxtamembrane position on Znt8 (Xue et al., 2020) and many CDF transporters have
histidine-rich loops, typically between TM4 and TM5, which might fulfill a similar role in recruiting
Zn^{2+} and orchestrating conformational change. Indeed, for Znt7 this His-rich loop was shown to
bind two Zn^{2+} ions leading these authors also to speculate that it plays a role in recruitment and
shuttling to the transport site (Han et al., 2022).

Stoichiometry of transport and energy coupling
The proton-motive force is generally postulated to drive antiport mechanism utilized by
CDF transporters. In the case of YiiP, the antiport involves exchange of Zn^{2+} from the cytoplasm
for H^+ in the periplasm and the stoichiometry of this process is key to energy coupling. In
previous work, the Zn^{2+}:stoichiometry has been measured to be 1:1 for ZitB (Chao and Fu,
2004a) and deduced by cellular or computational studies to be 1:2 for CzcD (Guffanti et al.,
2002) and Znt2 (Golan et al., 2019), respectively. In addition, isothermal calorimetry was used
to deduce a binding stoichiometry for Cd^{2+} of 1:1 for ecYiiP (Chao and Fu, 2004b). For the
current work, we refined CpHMD simulation data with experimental MST data using the inverse

Figure 5. Transport cycle for YiiP.

According to the alternating access paradigm, YiiP toggles between the OF and IF states via an intermediate occluded state. We assume that these changes occur independently in each protomer based on asymmetry seen in structures of soYiiP and Znt8 (Xue et al., 2020). Thus, our model depicts changes in the right-hand protomer while the left-hand protomer remains in a resting IF state. Although these intermediate states are informed by current and past structural work, they do not precisely conform to solved structures but represent hypothetical states that we believe to exist under physiological conditions. (A) Zn^{2+} is released to the periplasm. The TM2/TM3 loop is depicted interacting with the CTD in a Zn^{2+}-free, extended conformation, as seen in our D70A_asym structure. (B) The release of Zn^{2+} is facilitated by the low pH of the periplasm and results in protonation of two residues in site A, or potentially three residues at lower pH. (C-D) The protonated form transitions to the IF state via an occluded state. In the IF state, the Zn^{2+}-free TM2/TM3 loop is disordered. (E) Zn^{2+} is recruited to site B, inducing an ordered conformation of the TM2/TM3 loop that folds onto the CTD. (F) Zn^{2+} is transferred from a relatively low affinity site B to the much higher affinity site A via a negatively charged access channel, thus displacing two protons. (G) This transfer induces a Zn^{2+}-bound, occluded conformation in which the CTD tilts toward the occluded protomer and interacts with the TM2/TM3 loop in its Zn^{2+}-free, extended conformation. Features of the model are illustrated in the middle, boxed panel with desaturated colors. Zn^{2+} ions are depicted as magenta spheres and protons with a "+". The CTD is pink with two Zn^{2+} ions constitutively bound at site C. The scaffolding membrane helices (TM3 and TM6) are blue and the transport domain (TM1,2,4,5) is yellow. The TM2/TM3 loop is blue and depicted with dashed lines in the disordered state. Created with biorender.com.

Multibind approach to infer the prevalent protonation states for site A and thus to address
stoichiometry. Our analysis is consistent with Zn^{2+} binding to a fully unprotonated site A
regardless of the pH. After Zn^{2+} release, the transport site becomes either doubly or triply
protonated, depending on the pH. Due to strong coupling between His155 and Asp159, the
singly protonated state is essentially not seen. Thus, this model is consistent with a
stoichiometry of at least 1:2, possibly 1:3 in more acidic environments (pH < 6).

The energetics of transport are governed by the Nernst equation describing
electrochemical potential. The overall transport cycle can be described as

435
$$mH_o + nZn_i \rightleftharpoons mH_i + nZn_o$$

436   with the free energy calculated as

437
$$\Delta G = RT\ln\frac{[H_i]^m}{[H_o]^m} + RT\ln\frac{[Zn_o]^n}{[Zn_i]^n} + zFV_m$$

where *m* is the number of protons, *n* is the number of Zn^{2+} ions, *z* is the net charge for the
reaction and V_m is the membrane potential. Based on this standard equation, the higher
stoichiometries of 1:2 or 1:3 provide the cell with increased leverage over Zn^{2+} transport. In the
absence of membrane potential, for example, a 10-fold proton gradient (ΔpH of 1) can produce
Zn^{2+} gradients at equilibrium of up to 10^2 or 10^3, respectively, instead of simply 10 for a 1:1
stoichiometry (more generally, $10^{m/n}$). Membrane potential represents an important component
of the proton motive force that will influence electrogenic transport, where *z* is non-zero. In
particular, a stoichiometry of 1:1 would produce net positive charge transfer out of the cell and
thus would require working against this membrane potential. Given a potential of -80 mV,
relatively normal in *E. coli* (Felle et al., 1980), a 1:1 stoichiometry would produce an unfavorable
energy term that would overcome a chemical gradient of one pH unit. In contrast, a 1:2
stoichiometry would be electroneutral and would therefore be unaffected by membrane potential
thus generating a Zn^{2+} gradient of 10^2 as described above. Finally, a stoichiometry of 1:3 would
benefit both from the membrane potential as well as from the increased number of protons to
theoretically generate a gradient of $10^{4.4}$.

These calculations are based on equilibrium thermodynamics, but the pH dependence of
Zn^{2+} binding also has implications for the kinetics of transport. In particular, the dramatic
difference in K_d at cytoplasmic pH of 7.4 (1 nM) compared to more acidic pH's in the periplasm
(e.g., 6 μM at pH 6) implies that the on-rate of Zn^{2+} is dominant in the cytoplasm but that the off-
rate is enhanced in the periplasm, assuming that the binding site in the OF state has similar pH
dependence. This assumption seems plausible given the similarity of transport site geometry of
YiiP in the OF state (Lu et al., 2009). Thus, although the final equilibrium concentrations of Zn^{2+}
are not affected by the pH dependence of the transport site, the rate of transport and the rate of
equilibration will be greatly enhanced (Tanford, 1983).

YiiP has three aspartates and one histidine at site A, whereas many CDF transporters
including the mammalian Znt's substitute Asp47 for a second histidine. Previous work has
focused on changes in ion specificity associated with this change (Hoch et al., 2012), but it also
seems likely to affect stoichiometry. At a minimum, two histidines would make the 1:2
stoichiometry almost certain and a 1:3 stoichiometry highly plausible. The resulting
electrogenicity could be desirable for eukaryotic cells in which these transporters operate
predominantly in intracellular organelles such as insulin secreting granules, synaptosomes,

golgi or zinc-o-somes that are responsible for the dramatic Zn^{2+} sparks during oocyte fertilization
(Chu, 2018; Hara et al., 2017). Since the pH gradient of these organelles is modest, additional
driving force from the membrane potential might be key in producing the high internal
concentrations that are sometimes required for function.

MATERIALS and METHODS

Protein expression and purification

YiiP was expressed in *E. coli* (BL21(DE3)-CodonPlus-RIPL) from a pET vector that
included an N-terminal decahistidine tag. Cells were grown in LB media supplemented with 30
μg/ml kanamycin at 37°C until they reached an OD_{600} of 0.8. After cooling the media to 20°C,
expression was induced by addition of 0.5 mM isopropyl-β-D-thiogalactoside followed by
overnight incubation at 20°C. Cells were harvested by centrifugation at 4,000xg for 1 h,
resuspended in lysis buffer (20mM HEPES, pH 7.5, 100mM NaCl, 10% glycerol, and 500 μM
tris(2-carboxyethyl)phosphine) - 100 ml of buffer per 20 g of cells - and then lysed with a high-
pressure homogenizer (Emulsiflex-C3; Avestin, Inc. Ottawa Canada). Protein was extracted
from the membrane by adding 1.5 g dodecyl-β-D-maltoside per 100 ml of cell lysate followed by
2h of incubation at 4°C. Insoluble material was removed by centrifugation at 100,000xg for 30
min. The supernatant was loaded onto a Ni-NTA affinity column pre-equilibrated in buffer A (20
mM HEPES, pH 7.5, 100 mM NaCl, 10% glycerol, and 0.05% dodecyl-β-D-maltoside). The
column was washed by addition of buffer A supplemented with 20 mM imidazole and protein
was then eluted using a gradient of imidazole ranging from 20 to 500 mM. Peak fractions were
combined, supplemented with tobacco etch virus (TEV) protease (1:10 weight ratio of TEV:YiiP)
to cleave the decahistidine tag, and dialyzed overnight at 4°C against buffer A. TEV protease
was removed by loading the dialysate onto an Ni-NTA column and collecting the flow-through
fractions. After concentration, a final purification was done with a Superdex 200 size-exclusion
chromatography (SEC) column (GE Healthcare, Chicago, Illinois) equilibrated with SEC buffer
(20 mM HEPES, pH 7.5,150 mM NaCl, 0.2% n-decyl-β-D-maltoside, and 1 mM tris(2-
carboxyethyl)phosphine).

Fab selection, modification, expression, and purification has been described previously
(Lopez-Redondo et al., 2021). Briefly, the construct designated Fab2r was expressed in *E. coli*
strain 55244 from a freshly prepared transformation. Cells were cultured for ~24 h at 30°C with
constitutive expression behind an innately leaky T4 promoter. Cell pellets were harvested by
centrifugation, lysed with a high-pressure homogenizer and Fab was purified with a 5-ml HiTrap
Protein G HP column (GE Healthcare). Pooled fractions were dialyzed against sodium
carbonate buffer (pH 5.0) overnight at 4°C and further purified with Resource-S cation exchange
column (GE Healthcare). Finally, pure Fab protein was pooled and dialyzed against SEC buffer.

Cryo-EM sample preparation and data analysis

To ensure that YiiP was fully loaded with Zn^{2+}, the purified protein was initially incubated
with metal ion chelators (0.5 mM EDTA and 0.5 mM N,N,N′,N′-tetrakis(2-pyridinylmethyl)-1,2-
ethanediamine [TPEN]) for 24hrs at 4°C to remove ions that co-purified with the protein, as seen
in previous work (Lopez-Redondo et al., 2021). The sample was then dialyzed against YiiP SEC
buffer containing 0.5 mM EDTA for 10 h to eliminate chelator-metal complexes. Finally, YiiP was
loaded with Zn^{2+} by dialysis (10 h with four buffer exchanges) against SEC buffer supplemented
with 0.25 mM $ZnSO_4$. YiiP was then incubated with Fab2r at a 1:1 molar ratio for 1 h at 20°C to
form the YiiP-Fab2r complex. This complex was purified by SEC using a Superdex 200 column
equilibrated with SEC buffer supplemented with 0.25 mM $ZnSO_4$. Peak fractions at 3-5 mg/ml
were used immediately for preparation of cryo-EM samples. Specifically, 3–4 μl were added to
glow-discharged grids (C-Flat 1.2/1.3-4Cu-50; Protochips, Inc.) that were blotted under 100%
humidity at 4°C and plunged into liquid ethane using a Vitrobot (Thermo Fisher Scientific, Inc.
Bridgewater NJ).

The YiiP-Fab2r complex was imaged with a Titan Krios G3i electron microscope
(Thermo Fisher Scientific, Inc) equipped a Bioquantum energy filter and K2 or K3 direct electron
detector (Gatan, Inc. Pleasanton CA) with a pixel size of ~1 Å and a total dose of ~50
electrons/Å^2. Micrographs containing crystalline ice, excessive contamination, or imaging
artifacts were removed and the resulting micrographs were imported into cryoSPARC v2.15
(Punjani et al., 2017) for analysis. Particles were picked based on templates generated in
previous work and an initial set of particles were subjected to two rounds of 2-D classification to
remove false positives. The resulting particles were then subjected to successive rounds of ab
initio reconstruction with C1 symmetry and a resolution cutoff starting at 12 Å and declining to 8
Å; at each step, the best of two output classes was carried forward to the next round. The
resulting particles were used for heterogeneous refinement against two or three reference
structures derived from the ab initio jobs, still with C1 symmetry. A final selection of particles
was then used for non-uniform refinement using both C1 and C2 symmetry. Postprocessing
steps included calculation of local resolution and evaluation of 3-D variability (Punjani and Fleet,
2020). For the D70A construct, RELION was used to process a class of particles corresponding
to a dimer of dimers (Suppl. Fig. 3). After exporting these particles from cryoSPARC, steps of
2D classification and ab initio reconstruction were repeated in RELION, followed by 3D
refinement of ~117,000 dimer-of-dimer particles with C2 symmetry. A mask encapsulating one
dimer was created with Chimera (Pettersen et al., 2004) and symmetry expansion followed by
signal subtraction was used to generate a new set with ~234,000 dimeric particles. This
expanded particle set was imported back into cryoSPARC and combined with the class of
isolated dimer particles. This combined particle set was used for hetero-refinement to segregate
dimer complexes with symmetrical and asymmetrical TMD's. These segregated particle sets
were then used for final non-uniform refinement.

For model building of symmetrical structures from WT, D287A, D51A, we started with
the deposited coordinates from wild type (PDB, 7KZZ). This model was docked as a rigid body
to the map and adjusted manually in Coot (Emsley et al., 2010) as a starting structure for
refinement using PHENIX (Adams et al., 2010). For D70A and D287A/H263A mutants, the CTD
and the TMD from either 7KZZ or 7KZX were separately docked to the map. After adjustment
and crude rebuilding in Coot, these models were submitted to NAMDINATOR for Molecular
Dynamics Flexible Fitting (Kidmose et al., 2019). The resulting models were using as starting
points for PHENIX refinement. This refinement consisted of multiple rounds that alternated
between real-space refinement and manual adjustment using Coot. Atomic models were then
displayed using CHIMERA (Pettersen et al., 2004) and the PyMOL Molecular Graphics System
(Schrödinger, LLC, New York NY).

MST experiments
YiiP mutants were labelled with Alexa fluor 488 fluorescent dye (Invitrogen life
Technologies, Carlsbad CA) by adding 2.5 μL of dye from a 16 mM stock solution in DMSO to
200 μL of protein at 1-2 mg/ml in SEC buffer supplemented with 10% glycerol. Zn^{2+} was
removed by adding 0.5 mM EDTA and 0.5 mM TPEN followed by overnight incubation at 4°C.
Excess dye and chelated Zn^{2+} were removed using several cycles of dilution with SEC buffer
and concentration with a 50 kDa cutoff concentrator (AMICON, Millipore Sigma, Burlington MA).
For titration, Zn^{2+} was buffered either by 0.5 mM sodium citrate or 0.2 mM NTA; the total amount
of added $ZnSO_4$ was varied to achieve the desired concentration of free Zn^{2+}, according to the
program MAXCHELTOR (Bers et al., 1994). Protein concentration during the titration varied
from 8 to 100 nM. After 1:1 mixing of protein and Zn^{2+} solutions, samples were centrifuged for 5

min at 2000xg, then loaded into standard treated capillaries for measurement with a Monolith
NT.115 MST instrument (NanoTemper Technologies, South San Francisco, CA);
measurements were taken at 37 °C with LED power ranges from 20-60% and medium MST
power. Data from three independent titrations were analyzed with the MO.Affinity Analysis
software v2.3 using MST on-time of 15 s. For determination of K_d, data was fitted with a curve
based on the law of mass action,

572
$$F(C_{Zn}) = F_u + \frac{(F_b - F_u) * \left(C_{Zn} + C_P + K_d - \sqrt{(C_{Zn} + C_P + K_d)^2 - 4 * C_P * C_{Zn}} \right)}{2 * C_P}$$

573
574 where $F(C_{Zn})$ is the fraction bound, C_{Zn} is concentration of free zinc, C_P is the concentration of
575 YiiP protein, F_b and F_u refer to the normalized fluorescence in the bound and unbound state,
576 and K_d is the affinity constant. Alternatively, the data was fit with the Hill equation to assess
577 cooperativity,

578
$$F(C_{Zn}) = F_u + \frac{(F_b - F_u)}{1 + \left(\frac{EC_{50}}{C_{Zn}}\right)^n}$$

579 where EC_{50} is the half-maximal effective concentration (akin to K_d) and n is the Hill coefficient.

Overview of MD simulations

Two types of all-atom, explicit solvent MD simulations were conducted to assess the
effects of proton and zinc binding on the structure of the YiiP dimer in a lipid bilayer. We used
fixed charge equilibrium MD simulations (using a force field with unchanging parameters) to
investigate the interactions between Zn^{2+} ions and YiiP. We also carried out replica exchange
constant pH MD (CpHMD) simulations to calculate the pK_a values of all titratable residues and
assess the microscopic protonation states of the ion binding sites in the absence of Zn^{2+}. Unlike
CpHMD simulations, fixed charge simulations do not allow for dynamic protonation or
deprotonation of residues. Despite this limitation, they are our preferred approach to study
protein-zinc ion interactions and conformational changes because they can be run more
efficiently than CpHMD. Microsecond-length equilibrium MD simulations with fixed protonation
states were performed to quantify the effect of zinc ions on the structure of YiiP, using a
parametrization of zinc ions with the non-bonded dummy model as in our previous work (see
Table 5 for details of the simulations) (Lopez-Redondo et al., 2021).

Fixed charge equilibrium MD simulations

MD simulations in the *holo* (with Zn^{2+} in A, B, and C sites) and *apo* (no Zn^{2+}) state were
taken from our previous study (Lopez-Redondo et al., 2021). To investigate the influence of the
binding site B, we generated an "*empty site B*" structure by removing the Zn^{2+} ions from site B in
both protomers of PDB ID 5VRF, the same starting structure used previously.

The membrane-protein systems were modeled by embedding the YiiP dimer into a 4:1
palmitoyloleoylphosphatidylethanolamine:palmitoyloleoylphosphatidylglycerol (POPE:POPG)
lipid bilayer, which approximates the composition of the plasma membrane from *E. coli* (Raetz,
1986), and solvating the system with water as well as sodium and chloride ions corresponding
to a concentration of 100 mM using CHARMM-GUI v1.7 (Jo et al., 2008; Jo et al., 2009; Lee et
al., 2016). We used GROMACS 2021.1 (Abraham et al., 2015) with the CHARMM36 force field,
the CMAP correction for proteins (MacKerell et al., 1998; Mackerell et al., 2004), CHARMM36

Table 5. MD simulations

Simulation	length	N of replicas	Total simulation time	software
Fixed charge *apo**	1000 ns	3	3000 ns	GROMACS
Fixed charge *holo**	1000 ns	3	3000 ns	GROMACS
Fixed charge *empty site B*	1000 ns	3	3000 ns	GROMACS
CpHMD apo	12 ns	30 (pHREX)	360 ns	CHARMM

"Fixed charge" simulations were repeated three times. The CpHMD simulations were performed as coupled pH-replica exchange simulations.

*Simulations for apo and holo YiiP were taken from our previous work (Lopez-Redondo et al., 2021)

lipids (Klauda et al., 2010), and the CHARMM TIP3P water model. The *empty site B* system
contained 117512 atoms whereas the apo system contained 117394 atoms, both in a
hexagonal simulation cell (101 Å × 101 Å × 135 Å). Default protonation states of all ionizable
residues were used based on the experimental pH of 7. The neutral HSD tautomer (proton on
the N_δ) was selected to model all histidines except H73 and H155, which were modeled with
HSE (proton on the N_ε) based on their orientation relative to Zn^{2+} ions in the cryo-EM structure
PDB ID 5VRF.

The systems were first energy minimized and underwent a 3.75-ns six-stage
equilibration procedure with position restraints on protein and lipids, following the CHARMM-
GUI protocol (Jo et al., 2008). Three copies of 1-μs production simulations were carried out with
empty site B, starting from the same initial system conformation but with different initial
velocities. All simulations were performed under periodic boundary conditions at constant
temperature (T = 303.15 K) and pressure (P = 1 bar). The velocity rescaling thermostat (Bussi
et al., 2007) was used to maintain the temperature with a time constant of 1 ps and separate
temperature-coupling groups for protein, lipids, and solvent. A semi-isotropic pressure coupling
scheme was implemented using the Parrinello-Rahman barostat (Parrinello and Rahman, 1981)
with a time constant of 5 ps, a compressibility of 4.6×10^{-5} bar^{-1}. Long-range electrostatics were
calculated with the smooth particle mesh Ewald method (Essmann et al., 1995) under tinfoil
boundary conditions with an initial cutoff of 1.2 nm, which was optimized during the simulation,
and interactions beyond the cutoff were calculated in reciprocal space with a fast-Fourier
transform on a grid 0.12-nm and fourth-order spline interpolation. The van der Waals
interactions were switched smoothly to 0 between 1.0 nm and 1.2 nm, and the interactions were
shifted over the whole range and reduced to 0 at the cutoff. The Verlet neighbor list was
updated dynamically by GROMACS for optimized performance with a buffer tolerance of 0.005
kJ/mol/ps. Bonds to hydrogen atoms were treated as rigid holonomic constrains with the P-
LINCS algorithm (Hess, 2008) with an expansion order of four and two LINCS iterations;
alternatively, SETTLE (Miyamoto and Kollman, 1992) was used for water molecules. The
classical equations of motion were integrated with the leapfrog algorithm with a time step of 2 fs.

CpHMD simulations with pH replica exchange

In conventional MD simulations, the protonation states of titratable groups in the system
are fixed. To investigate the role of protons in the Zn^{2+} transport of YiiP, we performed
membrane-enabled hybrid-solvent continuous constant pH MD simulations (Huang et al., 2021).
The currently available implementation of the membrane-enabled CpHMD method does not
take into account the direct effect of ions on the titration of nearby residues so we only ran
CpHMD simulations for the apo system.

CpHMD simulations were initialized with the apo model (PDB ID 5VRF with Zn^{2+} ions
removed). An initial 250-ns production simulation was run for the CpHMD apo system to fully
relax the membrane, using the same GROMACS 2021.1 protocol described above for fixed
charge equilibrium simulations.

The membrane-enabled hybrid-solvent continuous CpHMD simulations were performed
using the CHARMM program version c42a2 (Brooks et al., 2009) with the PHMD module
(Khandogin and Brooks, 2005; Lee et al., 2004) and the pH replica-exchange (REPDSTR)
module (Wallace and Shen, 2011). The conformations of YiiP were sampled using conventional
all-atom simulations with the CHARMM22/CMAP all-atom force field (MacKerell et al., 1998;

Mackerell et al., 2004), the CHARMM36 lipid force field (Klauda et al., 2010), and the CHARMM
modified TIP3P water model. The titration coordinates were propagated using the membrane
GBSW implicit-solvent model (Im et al., 2003) with the GBSW input radii for the protein taken
from Chen et al. (Chen et al., 2006). Based on the average distance between the C2 atoms of
the lipids in the cytoplasmic- and periplasmic-facing leaflets, the thickness of the implicit bilayer
was set to 40 Å with a switching distance of 5 Å for the transition between the low dielectric slab
and bulk solvent. The implicit membrane was excluded by two cylinders with a radius of 14 Å
placed at the center of mass of each protomer of YiiP. The radius was selected to maximize the
coverage of the interior of the protein with minimal overlapping of the implicit membrane.

The final snapshot of the equilibrium simulation was used as the initial structure for the
CpHMD simulation. Dummy hydrogen atoms were added to the carboxylate groups of acidic
residues following the documentation of the PHMD module (Wallace and Shen, 2011) in
CHARMM (Brooks et al., 2009) using the HBUILD facility. The system was then equilibrated
with energy minimization using 50 steps of steepest descent followed by 50 steps of adopted
basis Newton-Raphson algorithms and CpHMD at pH 7 for 1 ns, whereby the harmonic
restraints on the protein heavy atoms were reduced from 1 kcal·mol^{-1}Å^{-1} to zero.

The production simulation was then performed using hybrid-solvent CpHMD with the pH
replica-exchange protocol (Huang et al., 2021; Wallace and Shen, 2011), using 30 replicas with
pH ranging from 1.5 to 11.5. The specific pH conditions were 1.5, 1.75, 2, 2.5, 2.75, 3, 3.25, 3.5,
3.75, 4, 4.25, 4.5, 4.75, 5, 5.25, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 8.75, 9, 9.5, 10, 10.25, 10.5, 11, 11.5,
chosen to ensure that the exchange rate between the nearby replicas was higher than 0.2. Each
replica was simulated under periodic boundary conditions at constant temperature (T = 303.15
K), pressure (P = 1 bar), and specified pH. A modified Hoover thermostat method (Hoover,
1985) was used to maintain the temperature, while pressure was maintained using the Langevin
piston coupling method (Feller et al., 1995) with a piston mass of 2500 amu. Long-range
electrostatics were evaluated with the particle mesh Ewald method (Darden et al., 1993) with a
real-space cutoff of 1.2 nm, and interactions beyond the cutoff were calculated in reciprocal
space with a fast-Fourier transform on a grid with 0.09-nm spacing and sixth-order spline
interpolation. The Lennard–Jones forces were switched smoothly to 0 between 1.0 and 1.2 nm,
and the potential was shifted over the whole range and reduced to 0 at the cutoff. Bonds to
hydrogen atoms were constrained with the SHAKE algorithm to allow a 2 fs time step. To avoid
a spike in potential energy due to a lack of solvent relaxation (Wallace and Shen, 2011), a
GBSW calculation was executed every 5 MD steps to update the titration coordinates. An
attempt to exchange adjacent pH replicas was made every 1000 MD steps (corresponding to 2
ps). Each replica simulation lasted 12 ns for a total aggregate sampling time of 360 ns.

Analysis of MD simulations
Analysis on the trajectories was carried out with Python scripts based on MDAnalysis
(Gowers et al., 2016). RMSDs of C$_\alpha$ atoms of the whole protein, TMD, and CTD were calculated
using the qcprot algorithm (Liu et al., 2010) after optimally superimposing the structure on the
same C$_\alpha$ atoms of the cryo-EM structure. Similarly, root mean square fluctuation (RMSF) of C$_\alpha$
atoms of the whole protein, TMD, and CTD were calculated. To assess the relative motion
between the two domains, a CTD-TMD rotation angle was calculated from the rotation matrix
which minimized the RMSD of the CTD after superimposing the protein on the TMD domain of
the reference structure (PDB ID 5VRF), as in our previous work (Lopez-Redondo et al., 2021).

The existence of the salt-bridge Asp72-Arg210 was quantified using the shortest distance
between O_δ atoms of Asp72 and hydrogen atoms on the side chain of Arg210.

The titration coordinates were extracted from CpHMD output files with the CpHMD-
Analysis scripts (https://github.com/Hendejac/CpHMD-Analysis).

The pK_a value of a titratable site was obtained in two different ways. For the first method,
we directly used the microstate probabilities from CphMD as part of a thermodynamic model of
the binding site using the *Multibind* method, as described below.

Individual residue pK_a's were obtained from a fit to the generalized Henderson-
Hasselbalch equation (Hill equation) (Henderson et al., 2020; Huang et al., 2016; Huang et al.,
2021) for the deprotonated fraction S as a function of pH,

704
$$S(\text{pH}) = \frac{1}{1 + 10^{n(\text{p}K_a - \text{pH})}}$$

705   where n is the Hill coefficient which represents the slope of the response curve. The
706   deprotonation fraction S of a titratable site was calculated from the number of trajectory frames
707   N of the residue in the deprotonated and protonated states as

708
$$S = \frac{N_{\text{deprot}}}{N_{\text{deprot}} + N_{\text{prot}}}$$

where the site is identified as deprotonated when the CpHMD titration coordinate λ is greater
than 0.8 and protonated when the titration coordinate is less than 0.2. The mean pK_a of a
residue was calculated for each pair of equivalent residues in protomer A and protomer B
because these residues were sampled independently in the CpHMD simulation. The statistical
error was estimated as the absolute difference between the pK_a's of protomer A and B residues
and their mean value.

The second method for determining pK_a values for individual residues relied on a
conventional approach employing the heuristic Hill equation to fit the average deprotonated
fraction as a function of pH. For this analysis, we treated titration coordinate data from protomer
A and B as independent; thus, by concatenating them we effectively doubled our sampling.
Using the same criterion for bound/unbound protons and the definition of microstates for site A
(Supplementary Figure 10) and site B (Supplementary Figure 11) we generated separate
microstate time series for sites A and B. These time series were then histogrammed to derive
the distributions of the microscopic protonation states of site A and B.

Calculation of microscopic, state-dependent pK_a values with the *Multibind* method
The *Multibind* method generates thermodynamically consistent models for systems of
coupled reactions by combining kinetic or thermodynamic measurements from different sources
using a maximum likelihood approach (Kenney et al., in preparation). Given free energy
differences between states (either from simulations or experiments), *Multibind* generates a set
of free energies for all states in the form of a potential graph that obeys path-independence of
free energies and detailed balance while being maximally consistent with the input data. From
the state free energies, all macroscopic thermodynamic observables can be calculated. Without
such an approach, thermodynamically inconsistent models arise due to random errors in the
input measurements. For Zn^{2+} and proton binding, the free energy differences in the potential

graph are calculated as functions of the external parameters, namely the free Zn^{2+} concentration
[X] and the pH, and with pK_a and Zn^{2+} standard state binding free energy values as input. The
binding free energy of a reaction $A + X \rightleftharpoons A{:}X$ is calculated as

736
$$\beta \Delta G_{\text{bind}}\,([X]) = \ln\left(\frac{K_D}{[X]}\right) = \ln\left(\frac{K_D}{c_0}\right) - \ln\left(\frac{[X]}{c_0}\right) = \beta \Delta G_{\text{bind}}^0 - \ln\left(\frac{[X]}{c_0}\right)$$

737   where $\beta = \frac{1}{k_B T}$ (k_B is Boltzmann's constant), K_D is the dissociation constant, and the binding free
738   energy is written as a sum of $\Delta G^0 bind$, the binding free energy at the standard state
739   concentration c_0=1 M, and a term depending on the ligand concentration [X]. For proton binding,
740   the free energy difference is expressed equivalently as

741
$$\beta \Delta G_{\text{prot}}\,(\text{pH}) = \ln(10)\,(\text{pH} - pK_a),$$

where pK_a is the acid dissociation constant.

We used *Multibind* with its Python implementation
(https://github.com/Becksteinlab/multibind) to construct thermodynamic models for binding sites
A and B, with the assumption that all binding sites across both protomers are independent.
Each model describes the transitions between all possible states as either binding of a proton or
binding of a Zn^{2+} ion. With four titratable residues, site A has $M=2^4=16$ Zn^{2+}-free protonation
microstates (Suppl. Fig. 10) and 16 Zn^{2+}-bound protonation microstates for a model with 32
states in total (Fig. 4h). Site B has three titratable residues and thus $M=8$ for a model with 16
states in total (Suppl. Fig. 11). Because only free energy differences between states are
measurable, we can arbitrarily specify one state as a reference with zero free energy. Here we
chose the 0-proton unbound state S0 as the reference state with G_0=0, but calculated
observables are independent of this particular choice. The probability of observing the system in
the microstate i is

755
$$P_i(\text{pH}, [Zn^{2+}]) = \frac{1}{Z} e^{-\beta G_i}$$

756   where G_i is the free energy of any state i and Z is the partition function

757
$$Z(\text{pH}, [Zn^{2+}]) = \sum_{k=1}^{M} e^{-\beta G_{k,\text{unbound}}} + \sum_{k=1}^{M} e^{-\beta G_{k,\text{Zn}}}$$

which contains the partition functions of the unbound and bound states. We note that the
microstate probabilities P_i can also be directly obtained from the CpHMD simulations, which
enables us to use *Multibind* as an inverse method to infer microscopic pK_a from CpHMD data
(without resorting to the Hill equation fit), as described below.

The fraction of YiiP protomers that have a Zn^{2+} ion bound to the binding site in question,
the *bound fraction*, is

764
$$\langle X \rangle = Z^{-1} \sum_{k=1}^{M} e^{-\beta G_{k,\text{Zn}}}$$

The bound fraction can be obtained experimentally from MST measurements.

With the ability to calculate microstate probability or bound fraction at specific pH or Zn^{2+}
concentration, we devised an inverse approach to *infer* microscopic, state-dependent pK_a's and
standard binding free energies (ΔG^0_{bind}) from either CpHMD data (to avoid using the Hill
equation with the deprotonated fractions, which are aggregated over multiple states and may
therefore mask coupling between residues) or from experimental MST measurements.

The inverse *Multibind* approach generates successive microscopic models with a Monte
Carlo scheme and compares the calculated observables to the target values. Observables
correspond to microstate probability distributions P_i(pH) for CpHMD or bound fraction <X> as a
function of pH and Zn^{2+} concentration for MST. Once convergence is reached, the microscopic
model contains the inferred microscopic state-dependent pK_a values and Zn^{2+} binding free
energies. We implemented a simple finite-temperature Monte Carlo algorithm that minimizes the
RMSD between the given and computed macroscopic features. The model was initialized with
pK_a values from CpHMD (obtained with the Hill equation) or ΔG^0_{bind} binding free energies
calculated from the experimental K_d obtained from the MST analysis. For each MC step, the
target observable was calculated for a range of pH values and Zn^{2+} concentrations. For as long
as the RMSD between the computed and the target observable remained above a cutoff (see
Table 6), a new set of pK_a or ΔG^0_{bind} was generated by adding a random value drawn from a
uniform distribution ranging from –0.2 to 0.2 pK_a units (for pK_a) or –0.2 to 0.2 kB_T (for binding
free energies) to all microscopic parameters. A new set of changes was accepted with
probability (Metropolis criterion)

786
$$P(pK_a, \Delta G^0_{bind} \rightarrow pK_a', \Delta G^0_{bind}') = \min(1, e^{-\frac{\Delta \text{RMSD}}{r}})$$

787   where $\Delta RMSD$ is the difference between the RMSDs between the target and the computed
788   observable for the old set of pK_a's or ΔG^0_{bind}'s and the new set

789
$$\Delta RMSD = RMSD(pK_a', \Delta G^0_{bind}') - RMSD(pK_a, \Delta G^0_{bind})$$

The fictional temperature in the Metropolis criterion, r, was set to 0.0001 to be on the
order of typical per-step changes. The RMSD cutoffs (Table 6) were chosen from initial runs
using a zero-temperature Monte Carlo algorithm, which accepted a new set of changes only
when $\Delta RMSD < 0$. These initial runs were stopped after 2000 steps and after no changes had
been accepted in the last 100 or more steps. The zero-temperature MC ensured that the final
set generated the smallest RMSD, while the finite-temperature MC used in the production runs
allowed larger parameter space to be sampled. Fifty runs were performed for each target.
Averaged pK_a values and corresponding standard deviations were calculated from the 50
independent MC runs. Although these standard variations of our parameter estimates weakly
depend on the somewhat arbitrary choice of r, we chose to report them instead of a rigorous
statistical error (which is unavailable with the current approach) to provide a sense of variability
of the estimates.

For site B, the CpHMD simulations suggested that H73 and H77 behaved identically and
had almost the same microscopic pK_a values. We calculated the "coupling energy" (Ullmann,
2003) between H73 and H77,

805
$$W = pK_a(S0{\rightarrow}S2) - pK_a(S1{\rightarrow}S3) = pK_a(S0{\rightarrow}S1) - pK_a(S2{\rightarrow}S3)$$

Table 6. MST convergence criteria

Target	Binding site	RMSD cutoff
CpHMD microstate probability distributions	Site A	0.0382
CpHMD microstate probability distributions	Site B	0.0447
MST bound fraction	Site A	0.0760
MST bound fraction	Site B	0.0995

(expressed in pK_a units, i.e. energy divided by $kT \ln 10$), from the microscopic CpHMD state
pK_a's (see Supplementary Data). The coupling energy measures how the protonation of a
specific residue (e.g. H73) depends on the protonation of another residue (here, H77). The
coupling energy is the difference in pK_a when H77 remains deprotonated while H73 binds a
proton (S0→S2, see Supplementary Figure 11) compared to the situation when H77 is already
protonated (S1→S3). The H73-H77 coupling energy for the CpHMD simulation was W=+1.0 for
D70 deprotonated and W=+0.2 for D70 protonated. Because W>0, binding of a proton to one of
the histidines decreases the pK_a of the other, thus decreasing proton binding via anti-
cooperative coupling. Preliminary calculations with the MST data, however, showed that small
initial differences in the starting values could lead to large pK_a shifts in H73 and H77, with the
directions determined by the initial ordering of pK_a values. Because we had no specific evidence
that the two histidines should behave differently, we considered the behavior of the initial
unconstrained MST inference calculations to be problematic. This problem indicated that the
inverse approach can be sensitive to initial values and that there may not always be sufficient
target data to constrain the microscopic model. We therefore treated the two residues as
symmetrical, i.e., both should be behaving in the same way, and imposed a constraint in the
Multibind approach so that both were assigned the same pK_a at each step, thus effectively
imposing the coupling observed in the CpHMD simulations. The interaction energies between
H73 and H77 for the MST-inference data are W=+6.2 (D70 deprotonated) and W=−0.70 (D70
protonated). The dominant state is the one with D70 deprotonated due to the low pK_a of D70
and hence the overall behavior of H73 and H77 remains strongly anti-cooperative in the MST-
inference model.

For Site A, the initial CpHMD simulations did not indicate any degeneracy in pK_a values
such as the one for H73/H77 and so no additional constraints were applied for any of the Site A
MST-inference calculations. We also calculated the coupling energy for H155 and D159 in site
A from the microscopic pK_a values (see Supplementary Data) for the CpHMD and the MST-
inference models for (1) no protons bound to D51 and D47 (W_{MST}=−5.05, W_{CpHMD}=+1.26), (2)
D51 protonated (W_{MST}=−7.39, W_{CpHMD}=+0.59), (3) D51 protonated (W_{MST}=−5.26, W_{CpHMD}=+1.02),
and (4) both D47 and D51 protonated (W_{MST}=−2.39, W_{CpHMD}=+1.79).Thus, refinement against
the experimental MST data changes the model based on the CpHMD results from anti-
cooperative binding (W>0) to strong cooperative binding (W<0).

The *inverse* Multibind approach has some limitations when applied to binding curves
such as the MST data. The binding isotherms, even when covering a range of pH values, do not
contain enough data to determine the microscopic model fully. Therefore, it was necessary to
use the CpHMD pK_a values to initialize the model instead of arbitrary starting values. Thus, in
the current implementation, the MST inference approach should be viewed as a refinement
process for the simulation-derived parameters, restricted or guided by the experimental data.
The resulting thermodynamic model allows the calculation of state probabilities (and any other
properties) at pH values outside the experimental range but it should be noted that these
calculated quantities are extrapolations that may not be accurate due to the lack of experimental
data to constrain the model; for instance, conformational changes may occur that could
drastically alter the interactions but these changes would not have been captured in a pure
binding model such as the one in Fig. 4h.

An effective pK_a under the Zn^{2+}-free condition was estimated for each residue by fitting
the deprotonation fractions (generated from the microscopic *Multibind* model) to the Hill-

Langmuir equation. The deprotonation fraction of a specific residue was calculated by summing
up the probabilities of microscopic states where the residue was deprotonated. The Zn^{2+}
concentration was set to 10^{-20} M to approximate the Zn^{2+}-free condition. For the symmetrized
residues H73 and H77, the Hill-Langmuir equation did not produce a satisfactory fit. In order to
compute macroscopic pK_a's for such a coupled system, we followed previous work (Henderson
et al., 2020; Ullmann, 2003) and fitted the total number of protons bound to the two residues to
the so-called "coupled titration model"

858
$$N_{prot} = 2 - [S_1(\text{pH}) + S_2(\text{pH})] = \frac{10^{(\text{p}K_2 - \text{pH})} + 2 \times 10^{(\text{p}K_1 + \text{p}K_2 - 2\text{pH})}}{1 + 10^{(\text{p}K_1 - \text{pH})} + 10^{(\text{p}K_1 + \text{p}K_2 - 2\text{pH})}}$$

859 where N_{prot} is the total number of protons, S_1 and S_2 are the deprotonation fractions of the two
860 residues, and pK_1 and pK_2 are the two coupled pK_a's describing the binding of the first proton
861 and the second proton to the coupled titrating sites. These two pK_a's can be interpreted as
862 effective pK_a values of two uncoupled residues.

Acknowledgements

We are grateful for help with initial MST experiments from Brian Kloss and Renato Bruni in the Center on Membrane Protein Production and Analysis at the New York Structural Biology Center, which receives funding from NIH grant P41GM116799. Electron microscopy was performed at the Cryo-EM core facility at NYU Langone Health. The authors acknowledge High Performance Computing core facility at NYU Langone Health as well as Research Computing at Arizona State University for providing computational and storage resources that have contributed to the research results reported within this paper. We also thank Bjørn Panella Pederson for critical reading of the manuscript.

Funding for this work was provided by NIH grants R01GM125081 (D.L. Stokes) and R35GM144109 (D.L. Stokes). A portion of this research was also supported by NIH grant U24GM129547 and performed at the PNCC at OHSU and accessed through EMSL (grid.436923.9), a DOE Office of Science User Facility sponsored by the Office of Biological and Environmental Research. MD simulations were performed using PSC Bridges at the Pittsburgh Supercomputing Center (allocation TG-MCB130177), a resource of the Extreme Science and Engineering Discovery Environment, which is supported by National Science Foundation grant ACI-1548562.

Competing Interests

None.

References

Abraham, M.J., Murtola, T., Schulz, R., Páll, S., Smith, J.C., Hess, B., and Lindahl, E. (2015).
GROMACS: High performance molecular simulations through multi-level parallelism from
laptops to supercomputers. SoftwareX *1--2*, 19 - 25, 10.1016/j.softx.2015.06.001.
Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J.,
Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W*., et al.* (2010). PHENIX: a comprehensive
Python-based system for macromolecular structure solution. Acta Crystallogr D Biol
Crystallogr *66*, 213-221, 10.1107/S0907444909052925.
Alejandro, S., Holler, S., Meier, B., and Peiter, E. (2020). Manganese in Plants: From
Acquisition to Subcellular Allocation. Front Plant Sci *11*, 300, 10.3389/fpls.2020.00300.
Bers, D.M., Patton, C.W., and Nuccitelli, R. (1994). A practical guide to the preparation of Ca2+
buffers. Methods Cell Biol *40*, 3-29, 10.1016/s0091-679x(08)61108-5.
Brooks, B.R., Brooks III, C.L., Mackerell Jr., A.D., Nilsson, L., Petrella, R.J., Roux, B., Won, Y.,
Archontis, G., Bartels, C., Boresch, S*., et al.* (2009). CHARMM: The biomolecular simulation
program. *30*, 1545-1614, https://doi.org/10.1002/jcc.21287.
Bussi, G., Donadio, D., and Parrinello, M. (2007). Canonical sampling through velocity
rescaling. J Chem Phys *126*, 10.1063/1.2408420.
Chandrangsu, P., Huang, X., Gaballa, A., and Helmann, J.D. (2019). Bacillus subtilis FolE is
sustained by the ZagA zinc metallochaperone and the alarmone ZTP under conditions of
zinc deficiency. Mol Microbiol *112*, 751-765, 10.1111/mmi.14314.
Chao, Y., and Fu, D. (2004a). Kinetic study of the antiport mechanism of an Escherichia coli
zinc transporter, ZitB. JBC *279*, 12043-12050, 10.1074/jbc.M313510200.
Chao, Y., and Fu, D. (2004b). Thermodynamic studies of the mechanism of metal binding to the
Escherichia coli zinc transporter YiiP. JBC *279*, 17173-17180, 10.1074/jbc.M400208200.
Chen, J.H., Im, W.P., and Brooks, C.L. (2006). Balancing solvation and intramolecular
interactions: Toward a consistent generalized born force field. Journal of the American
Chemical Society *128*, 3728-3736, 10.1021/ja057216r.
Cherezov, V., Hofer, N., Szebenyi, D.M., Kolaj, O., Wall, J.G., Gillilan, R., Srinivasan, V.,
Jaroniec, C.P., and Caffrey, M. (2008). Insights into the mode of action of a putative zinc
transporter CzrB in Thermus thermophilus. Structure *16*, 1378-1388,
10.1016/j.str.2008.05.014.
Choi, D.W., and Koh, J.Y. (1998). Zinc and brain injury. Annu Rev Neurosci *21*, 347-375,
10.1146/annurev.neuro.21.1.347.
Chu, D.S. (2018). Zinc: A small molecule with a big impact on sperm function. PLoS Biol *16*,
e2006204, 10.1371/journal.pbio.2006204.
Cotrim, C.A., Jarrott, R.J., Martin, J.L., and Drew, D. (2019). A structural overview of the zinc
transporters in the cation diffusion facilitator family. Acta Crystallogr D Struct Biol *75*, 357-
367, 10.1107/S2059798319003814.
Cotrim, C.A., Jarrott, R.J., Whitten, A.E., Choudhury, H.G., Drew, D., and Martin, J.L. (2021).
Heterologous Expression and Biochemical Characterization of the Human Zinc Transporter
1 (ZnT1) and Its Soluble C-Terminal Domain. Front Chem *9*, 667803,
10.3389/fchem.2021.667803.
Coudray, N., Valvo, S., Hu, M., Lasala, R., Kim, C., Vink, M., Zhou, M., Provasi, D., Filizola, M.,
Tao, J*., et al.* (2013). Inward-facing conformation of the zinc transporter YiiP revealed by
cryoelectron microscopy. PNAS *110*, 2140-2145, 10.1073/pnas.1215455110.
Daniels, M.J., Jagielnicki, M., and Yeager, M. (2020). Structure/Function Analysis of human
ZnT8 (SLC30A8): A Diabetes Risk Factor and Zinc Transporter. Curr Res Struct Biol *2*, 144-
155, 10.1016/j.crstbi.2020.06.001.

Darden, T., York, D., and Pedersen, L. (1993). Particle Mesh Ewald - an N.Log(N) Method for
Ewald Sums in Large Systems. J Chem Phys *98*, 10089-10092, Doi 10.1063/1.464397.
Emsley, P., Lohkamp, B., Scott, W., and Cowtan, K. (2010). Features and development of Coot.
Acta Crystallographica Section D - Biological Crystallography *66*, 486-501.
Essmann, U., Perera, L., Berkowitz, M.L., Darden, T., Lee, H., and Pedersen, L.G. (1995). A
Smooth Particle Mesh Ewald Method. J Chem Phys *103*, 8577-8593, Doi 10.1063/1.470117.
Felle, H., Porter, J.S., Slayman, C.L., and Kaback, H.R. (1980). Quantitative measurements of
membrane potential in Escherichia coli. Biochem *19*, 3585-3590, 10.1021/bi00556a026.
Feller, S.E., Zhang, Y.H., Pastor, R.W., and Brooks, B.R. (1995). Constant-Pressure Molecular-
Dynamics Simulation - the Langevin Piston Method. J Chem Phys *103*, 4613-4621, Doi
10.1063/1.470648.
Golan, Y., Alhadeff, R., Glaser, F., Ganoth, A., Warshel, A., and Assaraf, Y.G. (2018).
Demonstrating aspects of multiscale modeling by studying the permeation pathway of the
human ZnT2 zinc transporter. PLoS Comput Biol *14*, e1006503,
10.1371/journal.pcbi.1006503.
Golan, Y., Alhadeff, R., Warshel, A., and Assaraf, Y.G. (2019). ZnT2 is an electroneutral proton-
coupled vesicular antiporter displaying an apparent stoichiometry of two protons per zinc
ion. PLoS Comput Biol *15*, e1006882, 10.1371/journal.pcbi.1006882.
Gottesman, N., Asraf, H., Bogdanovic, M., Sekler, I., Tzounopoulos, T., Aizenman, E., and
Hershfinkel, M. (2022). ZnT1 is a neuronal $Zn(2+)/Ca(2+)$ exchanger. Cell Calcium *101*,
102505, 10.1016/j.ceca.2021.102505.
Gowers, R.J., Linke, M., Barnoud, J., Reddy, T.J.E., Melo, M.N., Seyler, S.L., Dotson, D.L.,
Domański, J., Buchoux, S., Kenney, I.M.*, et al.* (2016). MDAnalysis: A Python package for
the rapid analysis of molecular dynamics simulations. Paper presented at: Proceedings of
the 15th Python in Science Conference (Austin, TX: SciPy).
Guffanti, A.A., Wei, Y., Rood, S.V., and Krulwich, T.A. (2002). An antiport mechanism for a
member of the cation diffusion facilitator family: divalent cations efflux in exchange for K+
and H+. Mol Microbiol *45*, 145-153.
Gupta, S., Chai, J., Cheng, J., D'Mello, R., Chance, M.R., and Fu, D. (2014). Visualizing the
kinetic power stroke that drives proton-coupled zinc(II) transport. Nature *512*, 101-104,
10.1038/nature13382.
Han, B.B., Watanabe, S., Nomura, N., Liu, K., Uemura, T., Inoue, M., Tsutsumi, A., Fujita, H.,
Kinoshita, K., Iwata, S.*, et al.* (2022). Cryo-EM structures of human zinc transporter ZnT7
reveal the mechanism of Zn^{2+} uptake into the Golgi apparatus. bioRxiv.
Hara, T., Takeda, T.A., Takagishi, T., Fukue, K., Kambe, T., and Fukada, T. (2017).
Physiological roles of zinc transporters: molecular and genetic importance in zinc
homeostasis. J Physiol Sci *67*, 283-301, 10.1007/s12576-017-0521-4.
Henderson, J.A., Huang, Y., Beckstein, O., and Shen, J. (2020). Alternative proton-binding site
and long-distance coupling in Escherichia coli sodium-proton antiporter NhaA. Proc Natl
Acad Sci U S A *117*, 25517-25522, 10.1073/pnas.2005467117.
Hess, B. (2008). P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation. Journal
of chemical theory and computation *4*, 116-122, 10.1021/ct700200b.
Hoch, E., Lin, W., Chai, J., Hershfinkel, M., Fu, D., and Sekler, I. (2012). Histidine pairing at the
metal transport site of mammalian ZnT transporters controls Zn2+ over Cd2+ selectivity.
PNAS *109*, 7202-7207, 10.1073/pnas.1200362109.
Hoover, W.G. (1985). Canonical Dynamics - Equilibrium Phase-Space Distributions. Physical
Review A *31*, 1695-1697, DOI 10.1103/PhysRevA.31.1695.
Huang, Y., Chen, W., Dotson, D.L., Beckstein, O., and Shen, J. (2016). Mechanism of pH-
dependent activation of the sodium-proton antiporter NhaA. Nat Commun *7*, 12940,
10.1038/ncomms12940.

Huang, Y., Henderson, J.A., and Shen, J. (2021). Continuous Constant pH Molecular Dynamics
Simulations of Transmembrane Proteins. In Structure and Function of Membrane Proteins, I.
Schmidt-Krey, and J.C. Gumbart, eds. (New York, NY: Springer US), pp. 275-287.
Im, W.P., Lee, M.S., and Brooks, C.L. (2003). Generalized born model with a simple smoothing
function. J Comput Chem *24*, 1691-1702, 10.1002/jcc.10321.
Jo, S., Kim, T., Iyer, V.G., and Im, W. (2008). CHARMM-GUI: a web-based graphical user
interface for CHARMM. J Comput Chem *29*, 1859-1865, 10.1002/jcc.20945.
Jo, S., Lim, J.B., Klauda, J.B., and Im, W. (2009). CHARMM-GUI Membrane Builder for Mixed
Bilayers and its Application to Yeast Membranes. BJ *97*, 50--58, 10.1016/j.bpj.2009.04.013.
Kambe, T., Tsuji, T., Hashimoto, A., and Itsumura, N. (2015). The Physiological, Biochemical,
and Molecular Roles of Zinc Transporters in Zinc Homeostasis and Metabolism. Physiol Rev
95, 749-784, 10.1152/physrev.00035.2014.
Khandogin, J., and Brooks, C.L. (2005). Constant pH molecular dynamics with proton
tautomerism. BJ *89*, 141-157, 10.1529/biophysj.105.061341.
Kidmose, R.T., Juhl, J., Nissen, P., Boesen, T., Karlsen, J.L., and Pedersen, B.P. (2019).
Namdinator - automatic molecular dynamics flexible fitting of structural models into cryo-EM
and crystallography experimental maps. IUCrJ *6*, 526-531, 10.1107/S2052252519007619.
Klauda, J.B., Venable, R.M., Freites, J.A., O'Connor, J.W., Tobias, D.J., Mondragon-Ramirez,
C., Vorobyov, I., MacKerell, J., Alexander D, and Pastor, R.W. (2010). Update of the
CHARMM all-atom additive force field for lipids: validation on six lipid types. J Phys Chem B
114, 7830-7843, 10.1021/jp101759q.
Lee, J., Cheng, X., Swails, J.M., Yeom, M.S., Eastman, P.K., Lemkul, J.A., Wei, S., Buckner, J.,
Jeong, J.C., Qi, Y.*, et al.* (2016). CHARMM-GUI Input Generator for NAMD, GROMACS,
AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive
Force Field. Journal of chemical theory and computation *12*, 405-413,
10.1021/acs.jctc.5b00935.
Lee, M.S., Salsbury, F.R., Jr., and Brooks, C.L., 3rd (2004). Constant-pH molecular dynamics
using continuous titration coordinates. Proteins *56*, 738-752, 10.1002/prot.20128.
Liang, X., Dempski, R.E., and Burdette, S.C. (2016). Zn(2+) at a cellular crossroads. Curr Opin
Chem Biol *31*, 120-125, 10.1016/j.cbpa.2016.02.008.
Liu, P., Agrafiotis, D.K., and Theobald, D.L. (2010). Fast determination of the optimal rotational
matrix for macromolecular superpositions. J Comput Chem *31*, 1561-1563,
10.1002/jcc.21439.
Lonergan, Z.R., and Skaar, E.P. (2019). Nutrient Zinc at the Host-Pathogen Interface. Trends
Biochem Sci *44*, 1041-1056, 10.1016/j.tibs.2019.06.010.
Lopez-Redondo, M., Fan, S., Koide, A., Koide, S., Beckstein, O., and Stokes, D.L. (2021). Zinc
binding alters the conformational dynamics and drives the transport cycle of the cation
diffusion facilitator YiiP. J Gen Physiol *153*, 10.1085/jgp.202112873.
Lopez-Redondo, M.L., Coudray, N., Zhang, Z., Alexopoulos, J., and Stokes, D.L. (2018).
Structural basis for the alternating access mechanism of the cation diffusion facilitator YiiP.
Proc Natl Acad Sci U S A, 10.1073/pnas.1715051115.
Lu, M., Chai, J., and Fu, D. (2009). Structural basis for autoregulation of the zinc transporter
YiiP. Nature structural & molecular biology *16*, 1063-1067, 10.1038/nsmb.1662.
Lu, M., and Fu, D. (2007). Structure of the zinc transporter YiiP. Science *317*, 1746-1748,
10.1126/science.1143748.
MacKerell, A.D., Bashford, D., Bellott, M., Dunbrack, R.L., Evanseck, J.D., Field, M.J., Fischer,
S., Gao, J., Guo, H., Ha, S.*, et al.* (1998). All-atom empirical potential for molecular modeling
and dynamics studies of proteins. J Phys Chem B *102*, 3586-3616, 10.1021/jp973084f.
Mackerell, A.D., Jr., Feig, M., and Brooks, C.L., 3rd (2004). Extending the treatment of
backbone energetics in protein force fields: limitations of gas-phase quantum mechanics in

reproducing protein conformational distributions in molecular dynamics simulations. J
Comput Chem *25*, 1400-1415, 10.1002/jcc.20065.

Maret, W. (2013). Zinc biochemistry: from a single zinc enzyme to a key element of life. Adv
Nutr *4*, 82-91, 10.3945/an.112.003038.

Miyamoto, S., and Kollman, P.A. (1992). Settle - an Analytical Version of the Shake and Rattle
Algorithm for Rigid Water Models. J Comput Chem *13*, 952-962, DOI
10.1002/jcc.540130805.

Montanini, B., Blaudez, D., Jeandroz, S., Sanders, D., and Chalot, M. (2007). Phylogenetic and
functional analysis of the Cation Diffusion Facilitator (CDF) family: improved signature and
prediction of substrate specificity. BMC Genomics *8*, 107, 10.1186/1471-2164-8-107.

Outten, C.E., and O'Halloran, T.V. (2001). Femtomolar sensitivity of metalloregulatory proteins
controlling zinc homeostasis. Science *292*, 2488-2492, 10.1126/science.1060331.

Parrinello, M., and Rahman, A. (1981). Polymorphic transitions in single crystals: A new
molecular dynamics method. J~Appl~Phys *52*, 7182-7190, 10.1063/1.328693.

Parsons, D.S., Hogstrand, C., and Maret, W. (2018). The C-terminal cytosolic domain of the
human zinc transporter ZnT8 and its diabetes risk variant. FEBS J *285*, 1237-1250,
10.1111/febs.14402.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and
Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research and
analysis. J Comput Chem *25*, 1605-1612.

Punjani, A., and Fleet, D.J. (2020). 3D Variability Analysis: Resolving continuous flexibility and
discrete heterogeneity from single particle cryo-EM. 2020.2004.2008.032466,
10.1101/2020.04.08.032466 %J bioRxiv.

Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. Nat Methods *14*, 290-296,
10.1038/nmeth.4169.

Raetz, C.R. (1986). Molecular genetics of membrane phospholipid synthesis. Annu Rev Genet
20, 253-295, 10.1146/annurev.ge.20.120186.001345.

Sala, D., Giachetti, A., and Rosato, A. (2019). An atomistic view of the YiiP structural changes
upon zinc(II) binding. Biochim Biophys Acta Gen Subj *1863*, 1560-1567,
10.1016/j.bbagen.2019.06.001.

Sharma, G., and Merz, K.M. (2022). Mechanism of Zinc Transport through the Zinc Transporter
YiiP. Journal of chemical theory and computation *18*, 2556-2568, 10.1021/acs.jctc.1c00927.

Shusterman, E., Beharier, O., Shiri, L., Zarivach, R., Etzion, Y., Campbell, C.R., Lee, I.H.,
Okabayashi, K., Dinudom, A., Cook, D.I.*, et al.* (2014). ZnT-1 extrudes zinc from mammalian
cells functioning as a Zn(2+)/H(+) exchanger. Metallomics *6*, 1656-1663,
10.1039/c4mt00108g.

Tanford, C. (1983). Translocation pathway in the catalysis of active transport. Proc Natl Acad
Sci (USA) *80*, 3701-3705.

Udagedara, S.R., La Porta, D.M., Spehar, C., Purohit, G., Hein, M.J.A., Fatmous, M.E., Casas
Garcia, G.P., Ganio, K., McDevitt, C.A., and Maher, M.J. (2020). Structural and functional
characterizations of the C-terminal domains of CzcD proteins. J Inorg Biochem *208*, 111087,
10.1016/j.jinorgbio.2020.111087.

Ullmann, G.M. (2003). Relations between protonation constants and titration curves in polyprotic
acids: A critical view. Journal of Physical Chemistry B *107*, 1263-1271, 10.1021/jp026454v.

Wallace, J.A., and Shen, J.K. (2011). Continuous Constant pH Molecular Dynamics in Explicit
Solvent with pH-Based Replica Exchange. Journal of chemical theory and computation *7*,
2617-2629, 10.1021/ct200146j.

Xue, J., Xie, T., Zeng, W., Jiang, Y., and Bai, X.C. (2020). Cryo-EM structures of human ZnT8 in
both outward- and inward-facing conformations. eLife *9*, 10.7554/eLife.58823.

Wallace, J.A., and Shen, J.K. (2011). Continuous Constant pH Molecular Dynamics in Explicit
Solvent with pH-Based Replica Exchange. J Chem Theory Comput *7*, 2617-2629,
10.1021/ct200146j.
Xue, J., Xie, T., Zeng, W., Jiang, Y., and Bai, X.C. (2020). Cryo-EM structures of human ZnT8 in
both outward- and inward-facing conformations. Elife *9*, 10.7554/eLife.58823.
Yin, S., Duan, M., Fang, B., Zhao, G., Leng, X., and Zhang, T. (2022). Zinc homeostasis and
regulation: Zinc transmembrane transport through transporters. Crit Rev Food Sci Nutr, 1-
11, 10.1080/10408398.2022.2048292.
Zeytuni, N., Uebe, R., Maes, M., Davidov, G., Baram, M., Raschdorf, O., Nadav-Tsubery, M.,
Kolusheva, S., Bitton, R., Goobes, G.*, et al.* (2014). Cation diffusion facilitators transport
initiation and regulation is mediated by cation induced conformational changes of the
cytoplasmic domain. PLoS One *9*, e92141, 10.1371/journal.pone.0092141.
Zhang, S.F., Fu, C.T., Luo, Y.B., Xie, Q.R., Xu, T., Sun, Z.Y., Su, Z.M., and Zhou, X.M. (2023).
Cryo-EM structure of a eukaryotic zinc transporter at a low pH suggests its Zn2+-releasing
mechanism. J Struct Biol *215*, ARTN 107926 10.1016/j.jsb.2022.107926.

1097

1098

1099

Suppl. Fig. 1

Site A   Site B   Site C

WT

D51A

D70A asym

D70A sym

D287A

D287A/ H263A

Suppl. Fig. 2

A

D70A

B

3,420 mics → motioncorr patchCTF template pick

1,702,119 ptcls → 2D classification ab initio

685,046 ptcls

hetero refinement

C

164,345 ptcls
double complex
6.48 Å

342,514 ptcls
single complex
6.02 Å

178,187 ptcls
junk
9.26 Å

2x hetero refinement

hetero refinement

E   RELION

119,101 ptcls
double complex
6.12 Å

2D classification

117,368 ptcls

InitialModel
Refine3D

C2: 4.4 Å

symmetry expansion
signal subtraction

234,736 ptcls   342,514 ptcls

hetero refinement   reference vols

D

138,713 ptcls
sym TMD
6.11 Å

139,328 ptcls
asym TMD
6.03 Å

15,825 ptcls
junk
10.86 Å

F

182,413 ptcls
sym TMD: 6.04 Å

188,414 ptcls
asym TMD: 5.32 Å

NU refinement

NU refinement

G

−3.6Å
−4.0Å
−4.5Å
−5.4Å
−6.5Å

sym TMD

sym: 4.03 Å; B=126 Å²

No Mask (4.6Å)
Loose (4.1Å)
Tight (4Å)
Corrected (4Å)

DC   18Å 8.9Å 5.9Å 4.5Å 3.6Å 3Å 2.5Å 2.2Å

asym: 3.93 Å; B=146 Å²

No Mask (4.4Å)
Loose (4.1Å)
Tight (3.9Å)
Corrected (3.9Å)

DC   18Å 8.9Å 5.9Å 4.5Å 3.6Å 3Å 2.5Å 2.2Å

H

−3.6Å
−4.0Å
−4.5Å
−5.4Å
−6.5Å

asym TMD

Suppl. Fig. 3

Suppl. Fig. 4

Suppl. Fig. 5

Suppl. Fig. 6

Suppl. Fig. 7

Suppl. Fig. 8

Suppl. Fig. 9

Suppl. Fig. 10

Suppl. Fig. 11

Suppl. Fig. 12

Supplemental Figure Legends

Suppl. Figure 1. Determination of WT and D51A structures by cryo-EM.

(A,B) SDS-PAGE, left, and elution profile from SEC purification. Molecular weight standards are shown on the right at 116, 66, 45, 35, 25, 18.4, 14.4 kDa, with the YiiP monomer running as expected at 32.5 kDa. (C,D) The cryo-EM workflow starts with correction of beam-induced motion and contrast transfer function (CTF) of movies, followed by template-based particle picking. The particle set was subjected to two rounds of 2D classification and ab initio structure determinations to remove false positives. (E,F) Hetero-refinement jobs were used to look for multiple conformations and to maximize the homogeneity of particles for a final non-uniform refinement job, in this case with C2 symmetry. (G,H) The final structure was characterized by local resolution, represented by the colored surface shown from two orthogonal directions, and the Fourier Shell Correlation.

Suppl. Figure 2. Density at Zn^{2+} sites in the cryo-EM maps.

An overview of each cryo-EM structure is shown on the left, followed by densities at sites A, B and C, respectively. Contour levels used to display the various maps were as follows. WT: 7 σ, D51A: 7 σ, D287A, 6.5 σ for sites A and C, 5.5 σ for site B, D70A_asym: 9 σ, D70A_sym: 9 σ, D287A/H263A: 10 σ. Side-chains from selected residues only are shown, for clarity. For D70A_sym, site B is disordered in both protomers. In D70A_asym, the TM2/TM3 loop forms a Zn^{2+}-free association with the CTD (H1 and C-terminus of M6) in one protomer and is disordered in the other protomer. In D287A/H263A, one of the TM2/TM3 loops associates with CTD's from opposing protomers (CTDb, CTDc and TMDd refer to chains B, C and D, respectively). The other TM2/TM3 loop associates with the TM6/CTD linker as shown in Suppl. Fig. 5.

Suppl. Figure 3. Determination of D70A structures by cryo-EM

(A) SDS-PAGE and elution profile for SEC purification; molecular weight markers are at 116, 66, 45, 35, 25, 18.4, 14.4 kDa. The workflow for image processing was complex, involving multiple attempts to segregate distinct classes via ab initio and hetero-refinement jobs. This chart is a summary of the final, productive steps leading to structures presented. (B) Initial steps included correction of beam-induced motion and CTF followed by template-based particle picking, rounds of 2D classification and ab initio structure determination. (C) Hetero-refinement was used to segregate two major classes: class 1 comprising the expected dimer and class 2 comprising a dimer of dimers. The dimer of dimers involved interactions between Fab domains and thus did not appear to affect the conformations adopted by YiiP. (D) Particles from class 1 were subjected to further hetero-refinement to segregate into an additional two classes: one with symmetric TMD's (purple) and the other with asymmetric TMD's (blue) as described in the text. (E) Particles from class 2 (dimer of dimers) were exported to RELION where symmetry expansion and signal subtraction was used to generate a larger set comprising isolated dimers. (F) These symmetry-expanded particles were grouped together with class 1 particles for hetero-refinement using reference volumes with symmetric and asymmetric TMD's as well as a junk collector (not shown). (G, H) The resulting final classes were then used for non-uniform refinement to produce D70A_sym and D70A_asym structures, each colored according to local resolution and shown from orthogonal angles.

Suppl. Figure 4. Determination of D287A and D287A/H263A structures by cryo-EM
(A,B) SDS-PAGE and elution profiles from SEC purifications. Images come from a single gel
with the molecular weight markers (116, 66, 45, 35, 25, 18.4, 14.4 kDa). Position and presence
of the double peak is consistent with the higher order oligomerization seen during image
analysis. (C,D) Corrections for motion and CTF were followed by two rounds of 2D classification
and ab initio structure determination. (E,F) Hetero-refinement was used to select a
homogeneous set of particles for non-uniform refinement. (G,H) Final structures are
characterized by FSC and by local resolution, illustrated by the plots and by surface coloring,
respectively.

Suppl. Fig. 5. Interactions between the TM2/TM3 loop and the CTD.
(A) In the WT structure, the TM2/TM3 loop (blue) contacts the TM6/CTD linker from the
opposing protomer with proximity of D72 and R210. (B) In the D70A_asym structure, the Zn^{2+}-
free TM2/TM3 loop extends to interact with the H1 helix in the CTD of the opposing protomer.
(C) In the D287A/H263A structure, the TM2/TM3 loop extends towards CTD's of two different
protomers (e.g., TM2/TM3 loop from chain B inserts between CTD's of chains A and D). (D) In
the other monomer of D287A/H263A, the TM2/TM3 loop interacts with the TM6/CTD linker of an
opposing chain (e.g., the loop from chain A interacts with the linker from chain B). In these
chains, the TM6/CTD linker has refolded into one, long continuous helix.

Suppl. Fig. 6. Measuring Zn^{2+} affinity by MST
(A-D) SEC elution profiles for YiiP constructs. WT protein (peak at 12.4 ml) is included for
comparison but was not analyzed by MST. The site A mutant, D70A/H263A/D287A, site B
mutant, D51A/H263A/D287A, and site C mutant, D51A/D70A, produced peaks at at 12.6 ml,
12.3 ml and 12.6 ml respectively. (E-G) Raw data from MST for the titrations at pH 7 plotted in
Fig. 4. The blue rectangles correspond to the time window used for initial fluorescence levels
and the red rectangles correspond to the time window used for the thermophoresis rates. Each
plot includes all Zn^{2+} concentrations, which were performed in triplicate.

Suppl. Fig. 7. Walk of replica simulations in CpHMD pH-replica exchange (REX) through pH
space.
Each panel shows how one replica simulation (Rep:0 to Rep:29) changes over the course of the
simulation (Frame number) its current pH state, as indicated by the "pH replica" on the ordinate.
The pH replicas range from pH 1.5 (pH replica 0) to 11.5 (pH replica 29). Multiple replicas
sample most of the pH range and many move across the especially relevant range between pH
replica 9 (pH 4) and pH replica 22 (pH 9), indicating that the REX procedure samples pH space
well.

Suppl. Fig. 8. Convergence of the deprotonated fraction for titratable residues in CpHMD
simulations.
Residues in site A (D47, D51, H155, D159) and site B (D70, H73, H77) from both protomers (A
and B) are shown. The instantaneous deprotonated fraction S is plotted as a function of
simulation time and pH value, sampled across all replicas in the replica exchange simulation.
The deprotonated fractions generally converge to a stable value after about 8 ns with similar
behavior in both protomers A and B, thus indicating sufficient sampling during the CpHMD
simulations.

1296
Residues in site A (D47, D51, H155, D159) and site B (D70, H73, H77) are shown from both
protomers (A and B). The unprotonated fraction S from the end of the CpHMD simulations is
shown as a function of pH (black points). The Hill equation (generalized Henderson-Hasselbalch
equation) is fitted to the data (black line). The estimated pK_a is indicated as a red line at S=0.5.
As described in the text, H73/H77 were considered to be coupled and were analyzed in aggregate;
the two pK_as for the combined system (H73AH77A and H73BH77B) are represented by the pair of
red lines in the last two panels.

1304
(A). Populations of protonated states obtained directly from CpHMD simulations for site A, in the
absence of Zn^{2+}. Data for protomer A and protomer B were combined. Color coded definitions of
individual states are shown below with a "1" indicating the protonated state and absence of a
number the unprotonated state of each corresponding residue. (B) Populations of protonated
states from an inverse *Multibind* model based on inferring microscopic pK_a values directly from
the CpHMD populations in (A). (C) Population of protonated states resulting from MST
inference, which relies on the MC method to refine microscopic model parameters based on
experimental MST data. Populations represent the Zn^{2+}-free states although the model
incorporates the whole range of Zn^{2+} bound states [see (F)]. Only states S0, S3, S7, S11 have
appreciable (>0.0001 maximum probability) although all states contribute to aggregated state
probabilities in D and E. (D) MST inference results represented as populations of aggregated
states defined by the total number of protons bound in the absence of Zn^{2+}. (E) Deprotonated
fraction of each site A residue as a function of pH, averaged over all possible states of the MST
inference model in the absence of Zn^{2+}. Data points (symbols) were generated from the model;
then the Hill-Langmuir equation was fit to the generated data in order to obtain an effective per-
residue pK_a (listed in Table 4). (F) Population of states derived from the MST inference model
as a function of Zn^{2+} concentration at various pH values. S0Zn (dotted line) denotes the fully
deprotonated, Zn^{2+}-bound state, which is the only populated Zn^{2+}-bound state predicted by the
MST inference.

1324
(A). Populations of protonated states obtained directly from CpHMD simulations for site B, in the
absence of Zn^{2+}. Data for protomer A and protomer B were combined. Color coded definitions of
individual states are shown below with a "1" indicating the protonated state and absence of a
number the unprotonated state of each corresponding residue. (B) Populations of protonated
states from an inverse *Multibind* model based on inferring microscopic pK_a values directly from
the CpHMD populations in (A). (C) Population of protonated states resulting from MST
inference. For this analysis, a symmetry restraint was imposed on H73 and H77 as described in
the text. Populations are shown in the absence of Zn^{2+} although the model incorporates the
whole range of Zn^{2+} concentrations [see (G)]. (D) MST inference results represented as
populations of aggregated states defined by the total number of protons bound in the absence
of Zn^{2+}. (E) Deprotonated fraction of each site B residue as a function of pH, averaged over all
possible states of the MST inference model in the absence of Zn^{2+}. Data points were generated
from the model and the Hill-Langmuir equation was fit to D70 data (solid black line) to obtain an
effective per-residue pK_a; data for H73 and H77 are not correctly modelled by a Hill fit (not
plotted). (F) H73 and H77 were treated as a coupled system and the total deprotonated fraction
as a function of pH was fit with the "coupled titration model" (solid black line), resulting in two

effective pK_a values. (G) Population of states from the MST inference mode as a function of
Zn^{2+} concentration at various pH values. Unlike site A, multiple Zn^{2+}-bound states (S0Zn, S1Zn,
S2Zn, S3Zn) are populated at lower pH values.

Suppl. Fig. 12. Summary of structures from CDF transporters.
Cartoon representations of most structures so far determined. The structures have been
grouped according to conformational state: symmetric IF, symmetric OF and asymmetric states.
PDB accession codes are indicated for previously published structures together with
concentrations of Zn^{2+} present in the solution and the number of ions observed in the structures.
Accession codes for current work are listed in Table 1. Although the sites been named
differently for Znt7 and Znt8, the A,B,C nomenclature has been used to simplify the comparison.
The helices in the TMD are shown in rainbow colors starting with blue for TM1 and red for TM6.
CTD's are represented as a triangle. Zn^{2+} ions as magenta spheres. Hydrophobic residues
forming the so-called hydrophobic gate (Leu154 and Leu199 in SoYiiP) are shown as sticks.
Disordered in TM2/TM3 loops is indicated by a blurred line.

1355

1356