Evolution of quantitative traits under directional selection: Selective sweeps or small allele-frequency changes?

February 22, 2023

Hannah Götsch[a,b,*], Reinhard Bürger[a]

[a]*Faculty of Mathematics, University of Vienna, 1090 Vienna, Austria*
[b]*Vienna Graduate School of Population Genetics, Austria*

Abstract

We study the response of a quantitative trait to exponential directional selection in a finite haploid population, both on the genetic and the phenotypic level. We assume an infinite sites model, in which the number of new mutations per generation in the population follows a Poisson distribution (with mean Θ) and each mutation occurs at a new, previously monomorphic site. Mutation effects are beneficial and drawn from a distribution. Sites are unlinked and contribute additively to the trait. Assuming that selection is stronger than random genetic drift, we model the initial phase of the dynamics by a slightly supercritical Galton-Watson process. This enables us to obtain time-dependent results. We show that the copy-number distribution of the mutant in generation n, conditioned on survival until n, is described accurately by the deterministic increase from an initial distribution with mean 1. This distribution corresponds to the absolutely continuous part W^+ of the random variable, typically denoted W, that characterizes the stochasticity accumulating during the mutant's sweep. In bypassing, we prove the folklore result that W^+ is approximately exponentially distributed, but in general not exactly so by deriving its second and third moment. A proper transformation yields the approximate mutant frequency dynamics in a Wright-Fisher population of size N. Then we derive explicitly the (approximate) time dependence of the expected mean and variance of the trait and of the expected number of segregating sites. Unexpectedly, we obtain highly accurate expressions even for the quasi-stationary phase, when the expected per-generation response has equilibrated. These refine classical results, and they are derived from first principles. We find that Θ is the main determinant of the pattern of adaptation at the genetic level, i.e., whether the response is mainly due to sweeps or to small allele-frequency shifts, and the number of segregating sites is an appropriate indicator for it. The selection strength determines primarily the rate of adaptation. The accuracy of our results is tested for a Poisson offspring distribution by comprehensive simulations in a Wright-Fisher framework.

---

[*]Corresponding author
Email address: `hannah.goetsch@univie.ac.at` (Hannah Götsch)

Keywords: polygenic adaptation, mutation, infinite-sites model, Galton-Watson branching process, Wright-Fisher model

1. Introduction

Phenotypic adaptation is ubiquitous and can occur rapidly as a response to artificial selection or to a gradual or sudden change in the environment. The response of the mean phenotype can often be predicted or described by simple equations, such as the breeder's equation or its evolutionary counterpart, Lande's (1976) equation, thus on the basis of accessible phenotypic measurements (Falconer and Mackay, 1996, Walsh and Lynch, 2018). During this process genotype frequencies change, and new genotypes and new alleles can be generated by recombination and mutation. Even if the response of the mean phenotype is simple to describe, it is a highly complex, often insurmountable, task to infer the response on the genetic level.

There are multiple reasons for this problem. The response of a trait may be due to small frequency changes at a large number of loci of minor effect (Hill and Kirkpatrick, 2010, Barghi et al., 2019, Boyle et al., 2017), as originally proposed by Fisher (1918) and idealized and formalized in the infinitesimal model (Bulmer, 1980, Barton et al., 2017). The selection response may as well be due to selective sweeps at a few loci of large effect (Kaplan et al., 1989, Hermisson and Pennings, 2005, Messer and Petrov, 2013, Stephan, 2019). Finally, a mixture of both is a third realistic scenario (Lande, 1983, Chevin and Hospital, 2008, Chevin, 2019). In principle, different genetic architectures in combination with different initial conditions can lead to similar phenotypic responses, at least as the mean phenotype is concerned. Even if loci contribute additively to the trait and their number and effects are known, it is notoriously difficult to derive the allele-frequency dynamics at individual loci and the resulting dynamics of the trait's mean and variance from the selection pressure acting on the trait. Additional complications arise if linkage disequilibrium, epistasis, or pleiotropy have to be taken into account (Turelli and Barton, 1990, Bürger, 2000, Barton and Keightley, 2002, Zhang and Hill, 2005a, Walsh and Lynch, 2018).

For quite some time the focus in studying adaptation at the genetic and molecular level has been on selective sweeps, first hard sweeps and more recently soft sweeps which can occur from standing variation or immigration and are not restricted to mutation-limited evolutionary scenarios (Hermisson and Pennings, 2017). Although footprints of hard and soft selective sweeps are detectable in genetic data by various methods (Nielsen et al., 2005, Huber et al., 2016, Setter et al., 2020), they seem to be sparse in the data, especially hard sweeps in humans (Pritchard et al., 2010, Hernandez et al., 2011). The detection of polygenic responses by small allele-frequency changes at loci of minor effect is much more demanding. It became feasible at a larger scale only when sequencing became

cheaper (Berg and Coop, 2014). Genome-wide association studies (GWAS) became an important tool in identifying associations between genetic variants, typically single-nucleotide polymorphisms (SNPs), and phenotypic traits. Although GWAS are highly valuable in elucidating the genetic architecture underlying phenotypic traits, they provide no direct information on the contribution of the causal variants to adaptation of the trait, which can be deleterious, neutral or advantageous. More refined approaches, such as experimental evolution methods that yield time-series data, have been pursued successfully to demonstrate truly polygenic adaptation (reviewed in Sella and Barton, 2019, Barghi et al., 2020).

Theoreticians started only recently to explore when adaptation of polygenic traits is likely to be based on selective sweeps at a few loci vs. on small polygenic allele-frequency changes. Essential for this line of research is a good understanding of which genetic architectures and parameters (number and effects of loci, distribution of mutation effects, linkage, epistasis, pleiotropy) as well as population sizes and forms and strengths of selection can cause which adaptive patterns. By this we mean sweep patterns (hard or soft), polygenic patterns, or a mixture of both.

Some studies focused on polygenic traits under stabilizing selection, where the trait mean is close to the optimum or at most a few phenotypic standard deviations away and the selective response is due to standing genetic variation established previously by mutation-selection-(drift) balance. The findings are qualitatively similar, whether the population size is so large that random genetic drift can be ignored (De Vladar and Barton, 2014, Jain and Stephan, 2017a) or moderately large so that it plays a role but is weaker than selection (John and Stephan, 2020, Hayward and Sella, 2022). Roughly, if most standing variation is due to many polymorphic loci of small effect, then the response of the mean is mainly caused by small frequency changes. Only under exceptional circumstances do selective sweeps occur and drive the response. This result is not unexpected because (i) loci of small effect are more like to carry alleles of intermediate frequency and (ii) close to the phenotypic optimum fine tuning at the contributing loci is important.

Other studies focused on polygenic adaptation under directional selection (Höllinger et al., 2019), for instance caused by a sudden big environmental shift (Höllinger, 2018, Thornton, 2019). Although the assumptions on the genetic architecture and on the form of directional selection differ among these studies, they show that selective sweeps at several loci (parallel or successive) occur more often than in the above discussed scenario. The main determinant of the adaptive pattern is the so-called population-wide background mutation rate, but not the strength of selection or the population size (Höllinger et al., 2019).

In this study, we also explore how various evolutionary and genetic factors determine the rate and pattern of polygenic adaptation, both at the phenotypic and the genetic level. However, our assumptions depart in several respects from those imposed in previous studies. First, we

consider a trait subject to exponential directional selection, which is nonepistatic, does not induce linkage disequilibrium, and leads to a constant selection strength s. Second, in contrast to the investigations cited above, we focus on the response due to *de novo* mutations. Third, but similar to Thornton (2019), we assume an infinite sites model. The number of mutations per generation follows a Poisson distribution and each mutation occurs on a new, previously monomorphic locus. Loci are unlinked and contribute additively to the trait. Mutation effects are beneficial and follow a general distribution. This is justified by the assumption that in our population of size N selection is stronger than random genetic drift ($Ns > 1$), so that prolonged segregation or fixation of non-advantageous mutations can be ignored. These assumptions enable us to derive analytical expressions for the time dependence of the mutant frequency distribution at each locus, of the expected number of segregating sites, and of the expected mean and variance of the trait.

The paper is structured as follows. In Section 2, we introduce our basic model and give an overview of the analytical concepts regarding the one-locus model and its extension to infinitely many sites. Also our simulation setting is described. In Section 3, we derive an explicit, approximate expression for the mutant frequency distribution in a finite Wright-Fisher population as a function of time. This becomes feasible by using a slightly supercritical Galton-Watson process with arbitrary offspring distribution to describe the spreading of a beneficial mutant (see also Appendix A). Comparison with simulation results of a corresponding Wright-Fisher model demonstrates that our approximation is highly accurate for a wide range of parameters and a large time range. Explicit time-dependent expressions for the expected mean and variance of an additive quantitative trait under exponential selection are derived in Section 4 by assuming the infinite sites model. Interestingly, they even allow the derivation of expressions for the long-term, quasi-stationary response of the trait's mean and variance. These are not only derived from first principles but also recover and refine classical results. In Section 5 (and Appendix D), we derive explicit, approximate expressions for the evolution of expected number of segregating sites. Several of our results require a combination of branching process methods with approximations for the expected time to loss or to fixation of a new beneficial mutant. Some of them are new and derived from the diffusion approximations for sojourn times in Appendix B.

Because this paper is rather technical, we summarize, explain, and discuss our findings comprehensively in Section 6, where we also apply them to shed further light on the adaptive patterns underlying phenotpyic adaptation. We propose to use the expected number of segregating sites as an indicator for sweep vs. polygenic shift patterns. A central quantity in our theory is the expected number Θ of new beneficial mutations occurring per generation in the total population. We provide quantitative support to previous findings in related but different contexts that sweeps will occur if Θ is sufficiently much smaller than 1, whereas polygenic shifts will drive the selection response if

102 Θ is much greater than 1. Other model parameters have only a minor influence on the adaptive

103 pattern, but may have a major influence on the rate of response of the trait.

2. The model

We assume a haploid, panmictic population which has discrete generations and a constant

population size of N. Our main goal is to explore the dependence of pattern and rate of allele-

frequency change at the loci driving the adaptation of a quantitative trait on the model's parameters.

To this aim, we derive the dynamics from first principles. First we set up the basic model for the

quantitative trait, then the underlying one-locus model, the multilocus model, which we assume to

be an infinite-sites model, and the simulation model.

2.1. The quantitative-genetic model

We consider a quantitative trait subject to nonepistatic directional selection and investigate

the evolution of the distribution of allele frequencies at the loci contributing to this trait during

adaptation. The fitness of an individual with phenotypic value G is given by $W(G) = \exp(sG)$,

where the selection coefficient $s > 0$ is assumed to be constant in time. The trait is determined

additively by a (potentially infinite) number of diallelic loci at which mutation occurs. We ignore

environmental contributions to the phenotype which, in the absence of genotype-environment

interaction and of epistasis, would only weaken selection on the underlying loci. Therefore, genotypic

and phenotypic values are equal.

Each locus i can be in the ancestral state a_i or in the derived state A_i. The effect on the

phenotype of a locus in the ancestral state is zero, whereas a mutation at locus i contributes a value

$\alpha_i > 0$ to the phenotype G. Thus, the phenotype of an individual is given by $G = \sum_i \alpha_i \delta_{A_i}$, where

$\delta_{A_i} = 1$ if locus i is in the derived state, and $\delta_{A_i} = 0$ otherwise. The fitness of the ancestral allele

is normalized to 1. Therefore, also the fitness of the ancestral genotype, which has value $G = 0$,

is one. Because of our additivity assumption and the shape of the fitness function, a mutation

at locus i has a fitness of $\sigma_i = e^{s\alpha_i} > 1$, where $s > 0$. Hence, all mutations are advantageous.

Because we mostly assume $Ns > 1$, deleterious mutations are very unlikely to reach high frequency

or become fixed, and thus will not contribute to a permanent selection response of the mean

phenotype.

Unless stated otherwise, we assume that initially all loci are in the ancestral state and adaptation

occurs from *de novo* mutations. New mutations occur from the ancestral state to the derived state

and back-mutations are ignored. We define the population-scaled *advantageous* mutation rate

by $\Theta = NU$, where U is the total rate of mutations increasing the trait value, so that Θ is the

expected number of new mutations per generation that affect the trait. We assume an *infinite sites*

model, i.e., each mutation hits a new locus. This assumption is appropriate if per-locus mutation

136 rates are small enough such that the next mutation occurs only after the previous one became lost

or fixed.

In the infinite sites model we investigate two different scenarios. In the first, all mutation effects

are equal. In the second, mutation effects α_i at any locus i are drawn from a distribution defined

on $[0, \infty)$ that has a density f. Its mean is denoted by $\bar{\alpha}$ and its higher moments are assumed

to exist. Thus, the loci are exchangeable and all mutations are beneficial. For specification of

analytical results and their comparison with simulation results, we use an exponential distribution

(for empirical support, see Eyre-Walker and Keightley, 2007, Tataru et al., 2017) and a truncated

normal distribution (see Remark 4.10).

2.2. The one-locus model

We consider a single diallelic locus in isolation and assume that allele frequencies evolve

according to a haploid Wright-Fisher model with selection. The (relative) fitness of the ancestral

allele a and of the derived allele A (a new mutant) are set to 1 and $\sigma = e^s$, $s > 0$. Therefore,

the selective advantage of the mutant relative to the wildtype is $e^s - 1 \approx s$. We assume that in

generation $n = 0$ the new beneficial mutant A occurs as a single copy and no other mutation

occurs while this allele is segregating.

Because it seems unfeasible to derive analytic results for the time dependence of the allele

frequency for either the Wright-Fisher model or its diffusion approximation, we approximate the

stochastic dynamics in the initial phase, where stochasticity is most important, by a branching

process (e.g. Athreya and Ney, 1972, Allen, 2003, Haccou et al., 2005). During this initial phase,

interactions between mutant lineages can be ignored if N is large enough. The first step is to

approximate the evolution of the frequency distribution of the mutant by a Galton-Watson process.

In Section 3, we will approximate this discrete process by a continuous one and couple it with the

deterministic allele-frequency dynamics to obtain an analytically accessible model for the long-term

evolution of the mutant in a population of size N.

To be specific, we consider a Galton-Watson process $\{Z_n\}$, where $Z_n = \sum_{i=1}^{Z_{n-1}} \xi_i$, $Z_0 = 1$,

and ξ_i denotes the number of offspring of individual i in generation $n - 1$. Thus, Z_n counts the

number of descendants of the mutant, which emerged in generation 0. We assume that the ξ_i are

independent, identically distributed random variables satisfying

$$\mathrm{E}[\xi_i] = \sigma > 1 \ \text{ and } \ \mathrm{E}[\xi_i \ln(\xi_i + 1)] < \infty \,. \tag{2.1}$$

165 Therefore, the process $\{Z_n\}$ is supercritical. The ξ_i are also independent of n. We denote the

166 generating function of ξ_i by φ and observe that φ is also the generating function of Z_1. When we

167 compare our analytical results with simulation results from a Wright-Fisher model, we assume that

168 the offspring distribution is Poisson, because this provides a close approximation to the binomial

169 distribution if the population size N is sufficiently large, especially if $Ns \gg 1$ (cf. Lambert, 2006).

2.3. The infinite-sites model

We assume that the trait is determined by a potentially infinite number of loci at which new,
beneficial mutants can occur. We assume an infinite-sites model, so that every mutation occurs at
a new, previously monomorphic locus.

The initial population starts evolving in generation $\tau = 0$ in which the first mutations may
occur. Prior to generation 0, the ancestral allele is fixed at every site potentially contributing to the
trait. We assume that in each generation $\tau \geq 0$, the number of new mutants that emerge follows
a Poisson distribution with mean Θ. Each such mutant occurs at a new site (locus). Because
selection is multiplicative in our model and random genetic drift is a weak force relative to selection
(because N is assumed to be large), we ignore linkage disequilibrium among loci. Hence, we assume
that the allele-frequency distributions at the different loci evolve independently.

According to the mutation model, the ith mutation will occur at a random time τ_i. The
corresponding locus will be labeled locus i, and the mean offspring number of the mutant - when
the mutant is still rare in the population - is denoted by $\sigma_i > 1$. We approximate the distribution
of this waiting time τ_i by the Erlang distribution with density

$$m_{i,\Theta}(t) = \frac{\Theta^i t^{i-1}}{(i-1)!} \exp(-\Theta t), \quad \text{if } t > 0 \tag{2.2}$$

185 (and $m_{i,\Theta}(t) = 0$ if $t \leq 0$). Clearly, the mean waiting time until mutation i is i/Θ. We note that
186 if mutations occurred according to a Poisson point process with rate Θ, the Erlang distribution
187 would be the exact waiting-time distribution between i events.

188 The total number of mutation events k until generation τ is Poisson distributed with mean $\Theta\tau$

$$\text{Poi}_{\Theta\tau}(k) = \frac{(\Theta\tau)^k}{k!} \exp(-\Theta\tau). \tag{2.3}$$

189 In contrast, a non-recombining population could be considered. In this case, the model consists
190 of a trait determined by one locus with recurrent mutation, i.e., all mutations occur at the same
191 locus. This scenario and an extension to tightly linked, interfering sites will be studied elsewhere.

2.4. The simulation model

The simulation algorithm written in $C++$ is based on previous versions by Höllinger (2018)
and Wölfl (2019). With the above assumptions, we generate Wright-Fisher simulations forward in
discrete time to examine the adaptation of a trait under selection, mutation and random genetic
drift in the case of an originally monomorphic ancestral population. The first mutation events
can occur in generation 0. The algorithm acts on each locus because we assume that loci evolve
independently. The basic idea for the algorithm is that each generation is represented by a vector,

whose length is determined by the number of mutated loci. A mutation event occurs according to the mutation rate, adds a new locus and thus increases the length of the vector by one. The entries of the vector contain the number of mutants at each locus. Random genetic drift is included via binomial sampling with sampling weights due to selection (Ewens, 2004, p. 24). We use the *GNU Scientific Library* (Galassi et al., 2009) and the *MT19937* (Matsumoto and Nishimura, 1998) random number generator. Unless stated otherwise, a simulation for a set of parameters consists of $20,000$ replicate runs. Runs are stopped at a given number n (or τ) of generations. From these replicate runs, means (and variances) of the quantities of interest are computed, or histograms if distributions are the objects of interest.

3. Dynamics of allele-frequency distributions

It is well known that, conditioned on fixation, the frequency of a beneficial mutant grows faster than predicted by the corresponding deterministic model (e.g. Maynard Smith and Haigh, 1974, Otto and Barton, 1997). For models related to ours, it has been shown that conditioned on fixation and once a certain threshold has been exceeded, the evolution of the frequency of mutants can be approximated by a deterministic increase from a random variable, often labeled W, that describes the stochasticity that accumulates during the spread of the mutant (Desai and Fisher, 2007, Uecker and Hermisson, 2011). The latter authors called this the *effective initial mutant population size*. Martin and Lambert (2015) employed a variant of this approach and approximated the initial and the final phase of evolution by a Feller process conditioned on fixation. Based on this, they derived a semi-deterministic approximation for the distribution of the (beneficial) allele frequency at any given time. Below, we develop a refined variant that allows analytical treatment of our intended applications.

In Section 3.1, we prove the folklore result that in a slightly supercritical Galton-Watson process the absolutely continuous part of W is approximately exponentially distributed. However, we also show that in general it is neither exactly exponentially nor gamma distributed (Appendix A). In Section 3.2, we collect properties of the extinction probability until generation n for a Poisson offspring distribution. In Section 3.3, we use these results, condition on non-extinction until an arbitrary generation n in a slightly supercritical Galton-Watson process, and derive an approximation for the time-dependent distribution of the number of descendants of a single beneficial mutant. Theorem 3.6 is our main result and provides an explicit, analytically tractable and computationally efficient approximation for the density of a beneficial mutant in a diallelic haploid Wright-Fisher population of size N at any generation n. Finally, in Section 3.4 we apply Theorem 3.6 to the infinite-sites model.

 3.1. A general result for a single locus

Our starting point is the supercritical Galton-Watson process $\{Z_n\}$ defined in Section 2.2. The probability of ultimate extinction, denoted by $P_{\text{ext}}^\infty(\sigma)$, is the unique fixed point of the offspring generating function φ in $(0, 1)$, i.e., it satisfies $\varphi(P_{\text{ext}}^\infty) = P_{\text{ext}}^\infty$ and $0 < P_{\text{ext}}^\infty < 1$. When no confusion can occur, we suppress the dependence of P_{ext}^∞ on the mean offspring number $\sigma > 1$.

We define $W_n = Z_n/\sigma^n$. According to a classical result (Kesten and Stigum, 1966, Athreya and Ney, 1972, Haccou et al., 2005, p. 154) there exists a random variable W such that

$$\lim_{n \to \infty} W_n = W \tag{3.1}$$

with probability 1. The limiting random variable W has expectation $\mathrm{E}[W] = 1$, satisfies $W = 0$ precisely if the process dies out, and W is positive and continuous otherwise. We define the random variable W^+ corresponding to the absolutely continuous part of W by

$$\mathrm{Prob}[W \le x] = P_{\text{ext}}^\infty + (1 - P_{\text{ext}}^\infty)\,\mathrm{Prob}[W^+ \le x]\,, \quad \text{where } x \ge 0\,, \tag{3.2}$$

so that W^+ has a proper distribution with density w^+. We start by proving that W^+ is approximately exponentially distributed.

Lemma 3.1. *If the Galton-Watson process $\{Z_n\}$ defined above is slightly supercritical, then the positive, continuous part W^+ of the random variable W can be approximated by an exponential distribution with parameter $1 - P_{\text{ext}}^\infty$, where P_{ext}^∞ is the extinction probability of the process $\{Z_n\}$.*

Proof. We define the Laplace transform of W_n by $\Phi_n(u) = \mathrm{E}[e^{-uW_n}]$, where $u \ge 0$. Following Haccou et al. (2005, p. 155) we obtain

$$\Phi_n(u) = \mathrm{E}[e^{-uW_n}] = \mathrm{E}[\mathrm{E}[e^{-uW_n}|Z_1]] = \mathrm{E}[\mathrm{E}[e^{-\frac{u}{\sigma}W_{n-1}}]^{Z_1}] = \varphi(\Phi_{n-1}(u/\sigma))\,, \tag{3.3}$$

where φ is the generating function of Z_1 (see above). The third equality follows from the fact that, given Z_1, $Z_n = \sum_{i=1}^{Z_1} \chi_i$, where the χ_i are independent mutant population sizes with the distribution of Z_{n-1}. In the limit $n \to \infty$, this recursion yields the equation

$$\Phi(u) = \varphi(\Phi(u/\sigma))\,. \tag{3.4}$$

It is well known that this Poincaré functional equation is solved by a unique function $\Phi(u)$, which is the Laplace transform of the random variable W (e.g. Seneta, 1969, Haccou et al., 2005). Because Φ is monotone decreasing if $u > 0$ and we assume that $\{Z_n\}$ is supercritical, $1 > \Phi(u/\sigma) > \Phi(u) \ge \varphi(0)$ holds. In addition, because the process is slightly supercritical (i.e., $\sigma > 1$ but close to 1), $\Phi(u/\sigma)$ will be close to $\Phi(u)$, hence close to P_{ext}^∞, which satisfies $P_{\text{ext}}^\infty = \varphi(P_{\text{ext}}^\infty)$. Because P_{ext}^∞ is close to 1 in this case, $1 - \Phi(u/\sigma)$ is positive and close to zero.

Therefore, we obtain

$$\varphi(\Phi(u/\sigma)) = \mathrm{E}\big[(\Phi(u/\sigma))^{Z_1}\big] \approx (\Phi(u/\sigma))^{\mathrm{E}[Z_1]}$$
$$\approx \frac{1}{1 + \sigma(1 - \Phi(u/\sigma))} \,, \tag{3.5}$$

where we have used $\mathrm{E}[e^X] \approx e^{\mathrm{E}[X]}$, $\mathrm{E}[Z_1] = \sigma$, and $(1-x)^\sigma \approx \frac{1}{1+\sigma x}$ with $x = 1 - \Phi(u/\sigma)$. Together with (3.4), this yields

$$\Phi(u) \approx \frac{1}{1 + \sigma(1 - \Phi(u/\sigma))} \,. \tag{3.6}$$

It is known that for a Galton-Watson process with geometric offspring distribution (for which $\varphi(x) = (1 + \sigma(1 - x))^{-1}$), equality holds in (3.6) and, by inverting the Laplace transform, the resulting cumulative distribution has the form

$$\mathrm{Prob}[W \le x] = P_{\mathrm{ext}}^\infty + (1 - P_{\mathrm{ext}}^\infty)\,(1 - \exp[-(1 - P_{\mathrm{ext}}^\infty)x]) \,, \tag{3.7}$$

where for a geometric offspring distribution with mean σ the extinction probability is $P_{\mathrm{ext}}^\infty = 1/\sigma$ (Haccou et al., 2005, p. 155–156).

By comparing (3.7) with (3.2) and using continuity and uniqueness of the (inverse) Laplace transform, we conclude from (3.6) that the distribution of W^+ is approximately exponential with parameter $1 - P_{\mathrm{ext}}^\infty$. $\qquad\square$

Remark 3.2. Uecker and Hermisson (2011) derived an exponential distribution for W^+ for a birth-death branching processes with time-dependent mean offspring number (as they consider a model with temporal variation in its population size and the selection pressure). Martin and Lambert (2015) start the deterministic phase in their Wright-Fisher model with selfing from a gamma distribution because they admit one or more initially segregating mutants.

Remark 3.3. For a Poisson offspring distribution, the generating function of the offspring distribution and of Z_1 is given by $\varphi(x) = e^{-\sigma(1-x)}$. Therefore, the approximation (3.6) follows immediately from $e^{-\sigma(1-x)} \approx (1 + \sigma(1-x))^{-1}$.

Remark 3.4. By deriving the second and third moment, we show in Appendix A that in general the random variable W^+ defined in (3.2) is neither exactly exponentially nor gamma distributed. The expectation of W^+ is obviously $\mathrm{E}[W^+] = (1 - P_{\mathrm{ext}}^\infty)^{-1}$. Its variance is

$$\mathrm{Var}[W^+] = \frac{1}{1 - P_{\mathrm{ext}}^\infty}\left(\mathrm{Var}[W] + 1 - \frac{1}{1 - P_{\mathrm{ext}}^\infty}\right) \,, \tag{3.8}$$

where $\mathrm{Var}[W] = c_2/(1 + \sigma(\sigma - 1))$ and c_2 is the variance of the offspring distribution (cf. Haccou et al., 2005). If the offspring distribution is geometric with mean $\sigma > 1$, then $P_{\mathrm{ext}}^\infty = 1/\sigma$,

$\text{Var}[W_+] = \frac{\sigma^2}{(\sigma-1)^2} = \text{E}[W^+]^2$, and $\text{Skew}(W^+) = 2$ as is required for an exponential distribution. However, for a Poisson offspring distribution with mean $\sigma > 1$, we obtain

$$\text{Var}[W^+] = \frac{1 - \sigma P_{\text{ext}}^\infty}{(\sigma - 1)(1 - P_{\text{ext}}^\infty)^2}, \tag{3.9}$$

which is smaller than $\text{E}[W^+]^2$. With a Poisson offspring distribution, $\text{Skew}(W^+) = 2 - (m - 1) + \frac{1}{2}(m - 1)^2 + O((m - 1)^3)$. Appendix A contains derivations and more information on the moments of W and W^+. Series expansions of the density of W^+ and of the generating function are derived in Mieghem (2005), where also the densities of W^+ for Poisson and for geometric offspring distributions are compared graphically.

3.2. Extinction probability until generation n for a Poisson offspring distribution

Because we use a Poisson offspring distribution for comparisons of our analytic results with simulation results from the corresponding Wright-Fisher model, we summarize the properties of the probabilities of extinction of the mutant until generation n. We assume that the mutant starts as a single copy in generation $n = 0$, and the offspring distribution is Poisson with mean $\sigma = e^s$, $s > 0$. Then the probability $P_{\text{ext}}^\infty(\sigma)$ of ultimate loss, or extinction, of the mutant is the unique solution of $\varphi(x) = e^{-\sigma(1-x)} = x$ in $(0, 1)$. It can be expressed in terms of the (principal branch of the) product logarithm, or the Lambert function:

$$P_{\text{ext}}^\infty(\sigma) = -\frac{\text{ProductLog}[-\sigma \exp(-\sigma)]}{\sigma}. \tag{3.10}$$

With $\sigma = e^s$, this has the series expansion

$$P_{\text{ext}}^\infty(\sigma) = 1 - 2s + \frac{5s^2}{3} - \frac{7s^3}{9} + O(s^4). \tag{3.11}$$

The extinction probability $P_{\text{ext}}^\infty(e^s)$ decreases from 1 to 0 as s increases from 0 to ∞. We write $P_{\text{sur}} = 1 - P_{\text{ext}}^\infty$ for the (long-term) survival probability. It follows directly from (3.11) that Haldane's (1927) approximation, $2s$, serves as an upper bound for P_{sur} and approaches P_{sur} as $s \to 0$ (indeed, Haldane derived an equation equivalent to $e^{-\sigma(1-x)} = x$).

We define $P_{\text{ext}}(n, \sigma)$ as the probability that the mutation gets lost before generation n, i.e., $1 - P_{\text{ext}}(n, \sigma)$ is the probability that the mutation is still segregating in generation n. If no confusion can occur, we write $P_{\text{ext}}(n)$ instead of $P_{\text{ext}}(n, \sigma)$. Then $P_{\text{ext}}(n)$ satisfies the following recursion:

$$P_{\text{ext}}(0) = 0, \tag{3.12a}$$

$$P_{\text{ext}}(n) = \exp[-\sigma(1 - P_{\text{ext}}(n - 1))]. \tag{3.12b}$$

To demonstrate this, we write $p(k) = \frac{\sigma^k}{k!} \exp(-\sigma)$ for the probability that a mutant has k offspring. Because the mutation arises in generation zero, we have $P_{\text{ext}}(0) = 0$. The independence assumption

Figure 3.1: The solid curves show the extinction probability $P_{\text{ext}}(n, \sigma)$ in (3.12) until generation n as a function of n for Poisson offspring distributions with mean $\sigma = e^s$ as indicated. The dashed lines show the corresponding limit $P_{\text{ext}}^{\infty}(\sigma)$ from (3.10). The extinction probability of neutral mutations ($s = 0$) serves as an upper bound and is closely approached as the selection coefficient s gets small. Note that the branching process is no longer supercritical if $s = 0$.

among offspring lines in the branching process yields $P_{\text{ext}}(n + 1 | k \text{ mutants in generation } 1) = P_{\text{ext}}(n)^k$. Therefore, we obtain

$$P_{\text{ext}}(n + 1) = \sum_{k=0}^{\infty} p(k) P_{\text{ext}}(n)^k = \exp(-\sigma) \sum_{k=0}^{\infty} \frac{\sigma^k}{k!} P_{\text{ext}}(n)^k$$
$$= \exp(-\sigma) \exp[\sigma P_{\text{ext}}(n)] \tag{3.13}$$

and thus (3.12b). It follows that $P_{\text{ext}}(n, \sigma) \to P_{\text{ext}}^{\infty}(\sigma)$ for $n \to \infty$ and for every σ, as indeed holds for general offspring distributions with mean greater than 1.

The recursion is readily evaluated numerically, but an explicit formula for $P_{\text{ext}}(n)$ seems to be unavailable. However, the convergence of $P_{\text{ext}}(n, \sigma)$ to $P_{\text{ext}}^{\infty}(\sigma)$ is fast (Fig. 3.1). For instance, the number of generations needed for $P_{\text{ext}}(n, \sigma)$ to exceed $0.9 P_{\text{ext}}^{\infty}(\sigma)$ is $\leq \lceil 18 e^{-5s} \rceil$ if $s \geq 0$, where 18 is exact if $s = 0$, and $\lceil x \rceil$ denotes the smallest integer $\geq x$.

3.3. Evolution of the distribution of mutant frequencies at a single locus

Now we start the investigation of the evolution of the frequency distribution of a mutation that occurred as a single copy in generation $n = 0$. The following Lemma provides an approximation for the distribution of the number of beneficial mutants in an arbitrary generation n in an infinite population, provided the mutation is still present.

Figure 3.2: The probability density $\psi_n(y)$ of the number of descendants of a single mutation in generation $n = 5$ (blue) and $n = 20$ (red). Solid curves show the approximation (3.14). Dashed curves show the simplification of (3.14) obtained by approximating $P_{\text{ext}}(n)$ by P_{ext}^∞, which is equivalent to conditioning on long-term survival. The mean of the Poisson offspring distribution is $\sigma = e^s$ with $s = 0.1$. The histograms are from Wright-Fisher simulations with a population size of $N = 10^4$, conditioned on non-extinction until generation n.

Lemma 3.5. *Conditioned on non-extinction until generation n and for given σ, in an infinite population the (discrete) distribution of the number of descendants Z_n in generation n of a single mutation with fitness σ can be approximated by the distribution of the continuous random variable Y_n with density*

$$\psi_n(y) = (1 - P_{\text{ext}}(n))\sigma^{-n} \exp[-(1 - P_{\text{ext}}(n))\sigma^{-n}y]. \tag{3.14}$$

As with P_{ext}, we will usually suppress the dependence of ψ_n on σ.

Proof. We define the continuous random variable W_n^+ by its exponential density

$$w_n^+(x) = (1 - P_{\text{ext}}(n)) \exp[-(1 - P_{\text{ext}}(n))x], \; x \geq 0 \tag{3.15}$$

(and $w_n^+(x) = 0$ if $x < 0$). Because $\lim_{n \to \infty} P_{\text{ext}}(n) = P_{\text{ext}}^\infty$, Lemma 3.1 informs us that $\lim_{n \to \infty} W_n^+ = W^+$. Therefore, the continuous random variable

$$Y_n = \sigma^n W_n^+$$

can serve as an approximation of the distribution of Z_n conditioned on $Z_n > 0$, at least if n is not too small (because convergence to the exponential form will take some time). Hence, the distribution of the number of mutants Z_n in generation n can be approximated by the density

$$\psi_n(y) = w_n^+(y\sigma^{-n})\frac{d(y\sigma^{-n})}{dy}$$

of the continuous random variable Y_n. This yields (3.14). $\qquad\square$

Lemma 3.5 informs us that in an infinite population the distribution of the number of mutants after n generations, conditioned on non-extinction until generation n and originating from a single mutation, can be described by a deterministic exponential increase that starts with an exponentially distributed initial mutant population (according to ψ_0) of mean size 1. Then the mean of this population in generation n is $\sigma^n/(1 - P_{\text{ext}}(n))$, which approaches σ^n/P_{sur} as n increases, and the distribution remains exponential. Thus, according to this model a beneficial mutant destined to survival grows indeed faster than expected from the corresponding deterministic model (which has a growth rate of σ), in particular in the early generations; cf. (4.5).

As already noted, Desai and Fisher (2007), Uecker and Hermisson (2011), and Martin and Lambert (2015) used similar approaches, but conditioned on fixation, instead of non-extinction until the given generation. Therefore, they took the random variable W^+, or rather its exponentially distributed approximation with the fixation probability as parameter, as their *effective initial mutant population size* (which on average is larger than one). For large n, the distribution of $\sigma^n W^+$ has the same asymptotic behavior as ψ_n in (3.14), but our distribution ψ_n provides a much more accurate approximation for the early phase of the spread of the mutant (Figure 3.2) because in each generation it has the correct mean of $(1 - P_{\text{ext}}(n))^{-1}$. The distributions shown by dashed curves in Figure 3.2 are obtained by conditioning on long-term survival in the branching process.

Now we are able to formulate and prove a result for a finite population that will play a key role in the subsequent analyses.

Theorem 3.6. *Conditioned on non-extinction until generation n and for given N, the discrete distribution of the mutant frequency originating from a single mutation in generation 0 with fitness σ can be approximated by the distribution of the positive continuous random variable X_n with density*

$$g_a(x) = \frac{a}{(1-x)^2} \cdot \exp\left[-a\frac{x}{1-x}\right], \tag{3.16a}$$

where

$$a = a(n, \sigma, N) = N(1 - P_{\text{ext}}(n, \sigma))\sigma^{-n}. \tag{3.16b}$$

Proof. According to Lemma 3.5 the number of mutants Y_n in the branching process increases approximately exponentially in n. In a finite population this is impossible. Therefore, we follow the population genetics tradition and use relative frequencies (probabilities). Based on the well known relation between deterministic exponential and deterministic logistic growth (cf. eq. 4.5), the random variables X_n measuring the relative frequency of the mutants in generation n in a

population of size N can be defined by

$$X_n = \frac{Y_n}{N + Y_n},$$ (3.17)

whence $Y_n = NX_n/(1 - X_n)$. (For simplicity and because our analysis is intended to apply to large populations we use N instead of $N - 1$.) Thus, the density of X_n will be approximately

$$g_{a(n)}(x) = \psi_n\left(N\frac{x}{1-x}\right)\frac{d}{dx}\left(\frac{Nx}{1-x}\right),$$ (3.18)

where we recall that ψ_n in (3.14) depends on σ. This yields (3.16). $\qquad\square$

To interpret and understand the analytic results derived below, we first need to study the properties of the density $g_a(x)$ in (3.16) and of its constituent terms. Crucial quantities are $P_{\text{ext}}^\infty(\sigma)$ and $P_{\text{ext}}(n, \sigma)$. Figure 3.1 shows how fast $P_{\text{ext}}(n, \sigma)$ approaches $P_{\text{ext}}^\infty(\sigma)$ as n increases. The probability of extinction $P_{\text{ext}}(n, \sigma)$ until generation n converges quickly to the probability of extinction $P_{\text{ext}}^\infty(\sigma)$ if $s \gtrapprox 0.1$ and exceeds $0.9 P_{\text{ext}}^\infty(\sigma)$ in at most 18 generations (Section 3.2). Thus, simplifying the function g_a in (3.16) by substituting P_{ext}^∞ for $P_{\text{ext}}(n)$ will decrease the accuracy of the approximation considerably in the very early generations if the selective advantage is small.

The only parameter on which the mutant frequency distribution $g_a(x)$ in (3.16) depends is the compound parameter $a = a(n, \sigma, N)$. Obviously, $a(n, \sigma, N)$ is proportional to N. As a function of n, $a(n, \sigma, N)$ decreases approximately exponentially from its maximum value N, assumed at $n = 0$, to 0 as $n \to \infty$. As a function of σ (or s), $a(n, \sigma, N)$ decreases as well (see Fig. S.1).

We observe that $g_{a(n,\sigma,N)}(0) = a(n, \sigma, N)$. If $a > 2$, then $g_a(x)$ attains its maximum at $x = 0$ (Fig. 3.3). In this case $g_a(x)$ decays with increasing x (and approximately exponentially for small x). This will always be the case in the initial phase of the mutant's invasion. If $a < 2$, which will occur for sufficiently large n, then $g_a(x)$ has an unique maximum $g_a(\hat{x}) = \frac{4}{a}\exp(a - 2)$ at $\hat{x} = 1 - \frac{a}{2} \in (0, 1]$, which shifts from zero to one as a decreases (as caused by an increasing n).

Remark 3.7. Although the effective initial mutant population size $Y_0 = W_0^+$ in generation $n = 0$ is on average 1, the variable transformation (3.17) results in $\mathrm{E}[X_0] = \int x g_N(x)\,dx = 1 - Ne^N E_1(N) \approx \frac{1}{N} - \frac{2}{N^2}$, which deviates slightly from the initial condition in the Wright-Fisher and in the corresponding deterministic model. (Here, $E_1(a) = \int_a^\infty x^{-1}e^{-x}\,dx$ denotes the exponential integral; Abramowitz and Stegun, 1964, Chap. 5.1.) Nevertheless, growth is still considerably faster than in the deterministic model (Fig. 4.1).

Remark 3.8. Our expression (3.16) for the density $g_a(x)$ has the same structure as that given by Martin and Lambert (2015) in their equation (4). The difference is that their β_t decays exponentially with t with a constant parameter, whereas our $a = a(n, \sigma, N)$ decays exponentially with a parameter depending on n. This difference originates from the fact that we condition on

Figure 3.3: The probability density $g_a(x)$ in (3.16) is shown as a function of x and a.

non-extinction until n, whereas Martin and Lambert (2015) condition on fixation (see the discussion below Lemma 3.5).

Figures 3.4 and S.3 compare the analytic approximation (3.16) for the distribution of mutant frequencies at various generations with the corresponding histograms obtained from Wright-Fisher simulations. They show that the approximation is very accurate in the initial phase whenever $Ns \geq 1$. If $Ns \geq 100$, it remains accurate almost until fixation of the mutant. If Ns is of order 10, the approximation remains accurate for a surprisingly long period, essentially until the (mean) mutant frequency exceeds $\frac{1}{2}$, i.e., until it has become the dominating variant. That the branching-process approximation underestimates the effects of random genetic drift when the mutant reaches high frequency is an obvious consequence of its assumptions. In Section 4.1 we will see that the mean of g_a provides a highly accurate approximation to the true mean even if the mutation is close to fixation and Ns is much less than 100.

We note that Wright-Fisher simulations show that for $s > 0.5$ (i.e., very strong selection), the distribution of the mutant frequencies cannot be approximated by the density g_a. The reason is that for large σ and a Poisson offspring distribution, the density ψ_n of $Y_n = N\frac{X_n}{1-X_n}$ deviates too much from an exponential distribution. That for large σ, the variance becomes much smaller than the mean, can be inferred directly from (3.9). For a geometric offspring distribution, the limiting distribution is exponential, even for strong selection. In the following we exclude very strong selection and assume $s < 0.5$. In fact, we focus on parameters satisfying $s \leq 0.1$ and $Ns \geq 10$.

Figure 3.4: The curves show the probability density $g_{a(n,\sigma,N)}(x)$ in (3.16) of the mutant frequency in different generations n, as indicated in the legend. The histograms are obtained from simulations of the corresponding Wright-Fisher model. The population size is $N = 10^4$ and the mean of the Poisson offspring distribution is $\sigma = e^s$ with $s = 0.1$.

3.4. Evolution of the distribution of mutant frequencies in the infinite-sites model

We use the model outlined in Section 2.3. In particular, τ refers to the generation since the initial population started to evolve. Mutations $i = 1, 2, 3, \dots$ occur at generations τ_i, as outlined above. We note that τ differs from n, as used in the above subsections on a single locus, because for locus i the generation number in the Galton-Watson process is $n = 0$ at time τ_i. From (2.2) we recall the Erlang distribution and define

$$M_{i,\Theta}(\tau) = \int_0^\tau m_{i,\Theta}(t)dt = 1 - \frac{\Gamma(i, \Theta\tau)}{\Gamma(i)}, \qquad (3.19)$$

which approximates the probability that i mutations have occurred until generation τ. Here, $\Gamma(n, z) = \int_z^\infty x^{n-1}e^{-x}\,dx$ denotes the incomplete Gamma function (Abramowitz and Stegun, 1964, Chap. 6.5). We recall that $\lim_{z \to 0} \Gamma(n, z) = \Gamma(n) = (n-1)!$ (if n is a positive integer), and $\Gamma(0, z) = E_1(z)$.

In the following, we write $[t]$ for the nearest integer to t, and we define

$$a(t, \sigma_i, N) = N(1 - P_{\text{ext}}([t], \sigma_i))\sigma_i^{-t} \quad \text{for every (real) } t \geq 0\,, \qquad (3.20)$$

where σ_i is the fitness of the mutant at locus i. From Theorem 3.6 we infer that, conditioned on non-extinction until generation τ, the discrete distribution of the frequency of mutations at a locus at which the ith mutation event occurred, locus i for short, can be approximated by the

Figure 3.5: The curves in panel A show the absolutely continuous part $\tilde{h}_\tau^{(i)}(x)$ in (3.23) of the mutant frequency $\tilde{X}_\tau^{(i)}$ at the ith locus (indicated in the legend) in generation $\tau = 100$. The histograms are obtained from simulations of the corresponding Wright-Fisher model. The population-scaled mutation rate is $\Theta = 0.5$, the population size is $N = 10^4$ and the mean of the Poisson offspring distribution is $\sigma = e^s$ with $s = 0.1$. The red curve in panel B displays the probability $P_{\text{ext}}^{(i)}$ in (3.22) that the mutation at locus i gets lost until generation $\tau = 100$. The blue symbols show the simulation results from the corresponding Wright-Fisher model. For larger i, the simulated values would lie below the theoretical curve because in some replicate runs considerably less than 50 mutations occur until generation 100. We note that the shape of $h_\tau^{(i)}$ is essentially identical to that of $\tilde{h}_\tau^{(i)}$, except that the vertical range has to be rescaled. For smaller τ or smaller s slight differences become visible (results not shown).

distribution of the continuous random variable $X_\tau^{(i)}$ with density

$$h^{(i)}(x) = h_{\tau,\sigma_i,N,\Theta}^{(i)}(x) = \frac{1}{M_{i,\Theta}(\tau)} \int_0^\tau g_{a(\tau-t,\sigma_i,N)}(x)\, m_{i,\Theta}(t)\, dt \,. \qquad (3.21)$$

Here, we have used that given generation τ, the ith mutant has occurred approximately $[t]$ generations in the past, where t is Erlang distributed with parameters i and Θ.

The probability that the mutation at locus i gets lost until generation τ, i.e., that the ancestral type is again fixed at this locus in generation τ, can be approximated by

$$P_{\text{ext}}^{(i)}(\tau,\sigma_i,\Theta) = \frac{1}{M_{i,\Theta}(\tau)} \int_0^\tau P_{\text{ext}}(\tau - [t],\sigma_i)\, m_{i,\Theta}(t)\, dt \qquad (3.22)$$

(Fig. 3.5B). Below, we shall need unconditioned distributions of the mutant frequencies. Therefore, we define

$$\tilde{h}^{(i)}(x) = \tilde{h}_{\tau,\sigma_i,N,\Theta}^{(i)}(x) = \frac{1}{M_{i,\Theta}(\tau)} \int_0^\tau \left(1 - P_{\text{ext}}([\tau - t],\sigma_i)\right) g_{a(\tau-t,\sigma_i,N)}(x)\, m_{i,\Theta}(t)\, dt \qquad (3.23)$$

(see Fig. 3.5A). Then $\int_0^1 \tilde{h}_{\tau,\sigma_i,N,\Theta}^{(i)}(x)dx = 1 - P_{\text{ext}}^{(i)}(\tau,\sigma_i,\Theta)$ is the probability of non-extinction of the ith mutant until generation τ. Therefore, in the absence of conditioning on non-extinction, we obtain for the probability that in generation τ the frequency $\tilde{X}_\tau^{(i)}$ of the i-th mutant is less or equal $p \in [0,1]$:

$$\text{Prob}[\tilde{X}_\tau^{(i)} \le p] = P_{\text{ext}}^{(i)}(\tau,\sigma_i,\Theta) + \int_0^p \tilde{h}_\tau^{(i)}(x)\, dx \,. \qquad (3.24)$$

4. Evolution of the phenotypic mean and variance

Based on the results obtained above, we are now in the position to derive approximations for the expected response of the phenotype distribution from first principles. As usual in quantitative genetics, we concentrate on the evolution of the phenotypic mean and the phenotypic variance. For this purpose, we will need the first and second moments pertaining to the density $g_a(x)$. They are given by

$$\gamma_1(a) = \int_0^1 x g_a(x)\,dx = 1 - ae^a E_1(a) \tag{4.1a}$$

and

$$\gamma_2(a) = \int_0^1 x^2 g_a(x)\,dx = 1 + a - a(2+a)e^a E_1(a)\,, \tag{4.1b}$$

respectively, where $E_1(a) = \int_a^\infty x^{-1} e^{-x}\,dx$ is the exponential integral (cf. Remark 3.7). In addition, we will need

$$\gamma(a) = \int_0^1 x(1-x) g_a(x)\,dx = a(1+a)e^a E_1(a) - a\,. \tag{4.1c}$$

By the properties of the exponential integral, the mean $\gamma_1(a)$ decays from one to zero as a increases from 0 to ∞ (Abramowitz and Stegun, 1964, Chap. 5.1). Together with the properties of $a = a(t, \sigma, N)$, this implies that $\gamma_1(a(t))$ increases from approximately $1/N$ (Remark 3.7) to 1 as t increases from 0 to ∞ (see Fig. 4.1A). For the variance $\gamma(a)$, we find that $\gamma(a) \to 0$ as either $a \to 0$ or $a \to \infty$, and γ assumes a unique maximum at $a \approx 0.719$. As function of t, we have $\gamma(a(0)) \approx 1/N$ and $\gamma(a(t)) \to 0$ as $t \to \infty$ (see Fig. 4.1B). The value t at which the variance is maximized can be computed from (3.16b) with $a \approx 0.719$.

Remark 4.1. For large values of a (such as $a > 80$), we encountered numerical problems when using *Mathematica* to evaluate $\gamma_1(a)$ or $\gamma(a)$. The reason is that it multiplies the then huge value e^a with the tiny value $E_1(a)$, which is set to 0 if it falls below a prescribed value. However, the right-hand side of

$$e^a E_1(a) = \int_1^\infty \frac{1}{x} e^{a(1-x)}\,dx \tag{4.2}$$

can be integrated numerically without problems. An alternative efficient possibility is to use the approximation $e^a E_1(a) \approx \frac{1}{a}(1 - \frac{1}{a})$ (e.g., if $a > 50$), which is based on the asymptotic expansion $e^a E_1(a) = \sum_{k=0}^\infty (-1)^k \frac{k!}{a^{k+1}}$ (cf. Proof of Proposition 4.13).

The results derived below hold for offspring distributions having at least three finite moments (and mean greater than 1). A Poisson offspring distribution is used only for the numerical results (especially when compared to simulation results from a Wright-Fisher model) or if stated explicitly.

4.1. A single locus

We start by investigating the evolutionary dynamics of the mean and variance caused by the contribution of a single locus i. If α_i is the phenotypic effect of the mutant, $\sigma_i = e^{s\alpha_i}$ its fitness

effect, and $X_t^{(i)}$ the random variable describing its frequency t generations after its occurrence, the expected contribution to the phenotypic mean is $\alpha_i \, \mathrm{E}[X_t^{(i)}]$ and to the phenotypic variance $\alpha_i^2 \, \mathrm{E}[X_t^{(i)}(1 - X_t^{(i)})]$. By conditioning on non-extinction until t, we obtain from Theorem 3.6 and equations (4.1)

$$\bar{G}_i(t) = \alpha_i \int_0^1 x g_{a(t,\sigma_i)}(x) \, dx = \alpha_i \gamma_1(a(t,\sigma_i)) \tag{4.3}$$

and

$$V_{G_i}(t) = \alpha_i^2 \int_0^1 x(1-x) g_{a(t,\sigma_i)}(x) \, dx = \alpha_i^2 \gamma(a(t,\sigma_i)) \,. \tag{4.4}$$

In Figure 4.1 the branching process approximations $\bar{G}_i(t)$ and $V_{G_i}(t)$ (dashed curves) are compared with simulation results from the Wright-Fisher model (symbols) and with the deterministic dynamics (solid curve). In the latter, the mutant frequency $p(t)$ evolves according to

$$p(t) = \frac{\sigma^t p(0)}{1 + (\sigma^t - 1)p(0)} \,, \tag{4.5}$$

which is the solution of the difference equation $p' = \sigma p / \bar{w}$, where $\bar{w} = 1 + (\sigma - 1)p$ is the mean fitness (e.g., Bürger, 2000, p. 30). The figure shows excellent agreement between the branching process approximations and the Wright-Fisher simulations. It also demonstrates that in a finite population and conditioned on survival until generation t, the mutant spreads faster than in a comparable infinite population. The smaller the selective advantage, the faster is the (relative) initial increase of the mutant frequency, hence of the mean and the variance, on the time scale st. This is related to the fact that, to leading order in Ns (for $Ns \gg 1$), the scaled fixation time of a mutant of effect s in a Wright-Fisher model is $st_{\mathrm{fix}} \approx 2\ln(2Ns)$; cf. (B.2). For $N = 10^4$ and $s = 0.1$, $s = 0.001$, we obtain $st_{\mathrm{fix}} \approx 15.6$, 7.0, respectively, which provide quite good estimates for the time to completion of the sweep (Fig. 4.1).

The above described phenomenon is in fact well-known (e.g., Uecker and Hermisson, 2011). The intuitive explanation is that mutants that initially spread more rapidly than corresponds to their (small) selective advantage have a much higher probability of becoming fixed than mutants that remain at low frequency for an extended time period. The smaller the selection coefficient s, the more important is a quick early frequency increase.

4.2. Infinitely many sites

Now we proceed with the general case. Let α_i denote the effect of the ith mutation (at locus i) on the trait. Then its fitness effect is $\sigma_i = e^{s\alpha_i}$. The (unconditioned) distribution of the frequency $\tilde{X}_\tau^{(i)}$ of the ith mutant in generation τ was defined in (3.24) and its absolutely continuous part $\tilde{h}_\tau^{(i)}$ in (3.23). Below we shall need the k-th moment of $\tilde{X}_\tau^{(i)}$:

$$\mathrm{E}\left[(\tilde{X}_\tau^{(i)})^k\right] = \mathrm{E}\left[(X_\tau^{(i)})^k\right] = \int_0^1 x^k \tilde{h}_\tau^{(i)}(x) \, dx \,. \tag{4.6}$$

Figure 4.1: Contribution of a single locus to the expected phenotypic mean $\bar{G}_i(t)$ (panel A) and to the expected phenotypic variance $V_{G_i}(t)$ (panel B), t generations after the mutation has occurred. The dashed and the dotted curves show $\bar{G}_i(t)$ in (4.3) and $V_{G_i}(t)$ in (4.4) for $s = 0.1$ and $s = 0.001$, respectively. The symbols show the simulation results from the corresponding Wright-Fisher models. The solid curve shows mean and variance computed from the solution (4.5) of the deterministic model with $p(0) = 1/N$. In this case, the selection coefficient affects the evolution of the mutant frequency, hence of the mean and the variance, only through a proportional change in time scale. Rescaling time by the selection coefficient s makes the deterministic solution independent of s. In a finite population, weaker selection leads to a disproportionally faster spreading of the mutant, which is conditioned on non-extinction up to generation t. The population size is $N = 10^4$, and the mutation effect is normalized to $\alpha_i = 1$. Because we have $\alpha_i = 1$, $\bar{G}_i(t)$ and $V_{G_i}(t)$ coincide with the expected mean (4.1a) and variance (4.1c), respectively, of the mutant frequency distribution $g_a(x)$ defined in (3.16).

Now we are in the position to derive general expressions for the expectations of the mean phenotype $\bar{G}(\tau) = \bar{G}(\tau, f, s, N, \Theta)$ and of the phenotypic variance $V_G(\tau) = V_G(\tau, f, s, N, \Theta)$ in any generation τ, where we recall that the previously monomorphic population starts to evolve at $\tau = 0$ when the first mutation events may occur. The mutation effects α_i are drawn independently from the distribution f. Because we assume linkage equilibrium, the random variables $\tilde{X}_t^{(i)}$ ($i = 1, 2, 3, \ldots$) are independent in every generation t. For a given sequence of mutation events and given mutant frequency $x_\tau^{(i)}$ at locus i in generation τ, the phenotypic mean is $\sum_i \alpha_i x_\tau^{(i)}$ and the phenotypic variance is $\sum_i \alpha_i^2 x_\tau^{(i)}(1 - x_\tau^{(i)})$. Therefore, by taking expectations with respect to the Poisson distributed mutation events and the evolutionary trajectories of allele frequencies, we define the expected phenotypic mean and variance as follows:

$$\bar{G}(\tau) = \sum_{n=1}^{\infty} \text{Poi}_{\Theta\tau}(n) \left(\sum_{i=1}^{n} \int_0^\infty \alpha_i \, \mathrm{E}\left[\tilde{X}_{\tau, e^{s\alpha_i}, \Theta}^{(i)} \right] f(\alpha_i) \, d\alpha_i \right) \tag{4.7}$$

and

$$V_G(\tau) = \sum_{n=1}^{\infty} \text{Poi}_{\Theta\tau}(n) \left(\sum_{i=1}^{n} \int_0^\infty \alpha_i^2 \, \mathrm{E}\left[\tilde{X}_{\tau, e^{s\alpha_i}, \Theta}^{(i)} (1 - \tilde{X}_{\tau, e^{s\alpha_i}, \Theta}^{(i)}) \right] f(\alpha_i) \, d\alpha_i \right) . \tag{4.8}$$

The reader may note that the assumption of linkage equilibrium is not needed for the mean phenotype, and that assuming vanishing pairwise linkage disequilibria would be sufficient for the variance.

Proposition 4.2. *We assume the infinite sites model with mutation as introduced above. Then the expected mean and variance of the trait in any generation $\tau \geq 0$ are given by*

$$\bar{G}(\tau) = \Theta \int_0^\infty \alpha f(\alpha) \int_0^\tau \left(1 - P_{\text{ext}}([\tau - t], e^{s\alpha})\right) \gamma_1(a(\tau - t, e^{s\alpha})) \, dt \, d\alpha \qquad (4.9)$$

and

$$V_G(\tau) = \Theta \int_0^\infty \alpha^2 f(\alpha) \int_0^\tau \left(1 - P_{\text{ext}}([\tau - t], e^{s\alpha})\right) \gamma(a(\tau - t, e^{s\alpha})) \, dt \, d\alpha \, , \qquad (4.10)$$

respectively.

The proof is relegated to Sect. 4.5.

Remark 4.3. For a Wright-Fisher (or a related Markovian) model, these expressions hold only approximately because the density $g_a(x)$ derived in Theorem 3.6 is only an approximation of the true frequency distribution of the mutants. The proposition shows that the population-scaled mutation rate Θ affects the expected mean and the expected variance only as a multiplicative factor.

Figure 4.2 displays evolutionary trajectories of the expected mean and variance given in Proposition 4.2 and compares them with simulation results from corresponding Wright-Fisher models. On this scale of resolution the agreement is excellent, both for equal and exponentially distributed effects. After an initial phase the trajectories reach a stationary phase, where the equilibrium variance has stabilized and the response of the mean is approximately linear in time. The expected mean fixation times given in the figure caption provide accurate approximations for the average time required to enter the stationary phase. For refined comparisons with more explicit approximations for the stationary phase and for the initial phase we refer to Fig. 4.3 and to Fig. 4.4, respectively.

Notable in all panels of Fig. 4.2 is the faster increase of the mean and the variance for smaller selection coefficients on the time scale $s\tau$ (see the discussion in the connection with Fig. 4.1). In addition, for the same selection coefficient the evolutionary response is, especially initially, much faster for exponentially distributed mutation effects (panels C and D) than for equal mutation effects (panels A and B), as shown by different steepness (A vs. C) and shape (B vs. D) of the trajectories. The reason is that mutations of larger effect tend to fix earlier and more frequently than mutations of smaller effect. Therefore, large mutation effects speed up the response in the initial phase, whereas small mutation effects delay the entry into the stationary phase, with both types occurring in an exponential distribution. As shown by the numerical values of st_{fix} and $s\bar{t}_{\text{fix}}$ in the caption of Fig. 4.2, this results in similar mean fixation times for equal and for exponentially distributed mutation effects, and thus in similar average entry times into the stationary phase.

Figure 4.2: The scaled expected phenotypic mean $\bar{G}(s\tau)/\Theta$ (panels A and C) and the scaled expected phenotypic variance $V_G(s\tau)/\Theta$ (panels B and D) are shown for the selection coefficients s given in the legend. Generations τ are scaled by s to facilitate comparison of curves and data with differing s. In A and B all mutation effects are $\alpha_i = 1$, and in C and D the distribution f of mutation effects is exponential with mean $\bar{\alpha} = 1$. The solid curves show $\bar{G}(s\tau)/\Theta$ and $V_G(s\tau)/\Theta$ as computed for equal effects from (4.11) and (4.12), respectively, and for exponentially distributed effects from (4.9) and (4.10). The dashed lines show the equilibrium values given by (4.17) and (4.18). The values of the mean fixation times multiplied by s ($= 0.1, 0.01, 0.001$) are $st_{\text{fix}} \approx 15.6, 11.7, 7.0$ for equal effects $\alpha = 1$ in A and B, and $s\bar{t}_{\text{fix}} \approx 15.8, 10.6, 5.9$ for exponentially distributed effects in C and D, where t_{fix} is calculated from (B.4) and $\bar{t}_{\text{fix}}$ from (4.14) as described in Appendix B.2. The symbols are from Wright-Fisher simulations with $\Theta = \frac{1}{2}$. The population size is $N = 10^4$. Analogous results for a truncated normal distribution ($\alpha \geq 0$ and $\bar{\alpha} = 1$) are shown in Figure S.4.

Finally, with an exponential distribution the equilibrium variance is about twice as large as for equal effects (see also Corollary 4.8).

Whereas the simulation results for $2\Theta \geq 1$ are accurately described by the analytic approximations for the phenotypic mean and the phenotypic variance (Figure 4.2), for $2\Theta \ll 1$ much more fluctuation is observed in the simulations (Fig. S.2). The reason is that the time between successful mutation events is much larger for smaller Θ, and this results in more pronounced stochastic effects. For small Θ one sweep after another occurs, whereas for large Θ parallel sweeps are common. This will be quantified in Section 5 using the number of segregating sites in the population.

Remark 4.4. In general, the integrals in (4.9) and (4.10) cannot be simplified. Of course, for equal

effects α, $\int_0^\infty \alpha^k f(\alpha) \int_0^\tau \ldots dt\, d\alpha$ simplifies to $\alpha^k \int_0^\tau \ldots dt$. An alternative approach is to assume that the ith mutation occurs at its mean waiting time i/Θ and the number of mutation events until time τ is (approximately) $[\Theta\tau]$. For given mutation effects α_i, we obtain instead of eqs. (4.9) and (4.10) the simpler expressions

$$\bar{G}(\tau) \approx \sum_{i=0}^{[\Theta\tau]-1} \left(1 - P_{\text{ext}}([i/\Theta], \sigma_i)\right)\bar{G}_i\left(i/\Theta\right) \tag{4.11}$$

and

$$V_G(\tau) \approx \sum_{i=0}^{[\Theta\tau]-1} \left(1 - P_{\text{ext}}([i/\Theta], \sigma_i)\right)V_{G_i}\left(i/\Theta\right) , \tag{4.12}$$

respectively, where we have used the one-locus results (4.3) and (4.4). Their numerical evaluation is very efficient. The derivation is given in Appendix C.3. A meaningful application of these formulas requires $[\Theta\tau] > 1$. Then, for equal mutation effects, (4.11) and (4.12) are highly accurate approximations of (4.9) and (4.10), respectively (results not shown, but expected because the right-hand sides of (4.11) and (4.12) can be interpreted as approximating Riemann sums). For values of Θ much smaller than 1, few new mutations are expected until generation τ (unless τ is sufficiently large) and the rounded, expected number of mutation events, $[\Theta\tau]$, can be zero for many generations. Moreover, the Poisson distribution is highly asymmetric for small $\Theta\tau$ and gets more symmetric for larger $\Theta\tau$. This also implies that the phenotypic mean and variance may be only poorly described by their first moments when one sweep after another occurs (as with very small Θ). By Proposition 4.2, the mutation rate Θ enters the expected phenotypic mean and variance only as a multiplicative factor (see also Fig. S.2). Hence, we use $\bar{G}(\Theta) \approx \Theta \cdot \bar{G}(1)$ and $V_G(\Theta) \approx \Theta \cdot V_G(1)$ to obtain approximations from equations (4.11) and (4.12) for a small Θ. These are evaluated much faster than the integrals in Proposition 4.2 and are used in Fig. 4.2A,B and Fig. S.2.

In the following we derive simpler and more explicit expressions for the phenotypic mean and variance of the trait that describe their approximate dependence on the model parameters accurately. First, we focus on the long-term behavior of mean and variance, then on their initial increase.

4.3. Approximations for the stationary phase

Although our approach was designed to study the early phase of adaptation, we show here that it yields accurate and simple approximations for the stationary phase, when the equilibrium variance has stabilized and the response of the mean is approximately linear in time.

Because in diffusion theory quantities such as fixation probabilities or expected times to fixation are typically given as functions of the selection coefficient (and the population size), we denote the

selection coefficient of a mutant of effect α by

$$\tilde{s}(\alpha) = e^{s\alpha} - 1 \,. \tag{4.13}$$

For given N and s, we will need the expected mean fixation time of a single mutant (destined to fixation). The expectation is taken with respect to the distribution f of the mutation effects α which has mean $\bar{\alpha}$:

$$\bar{t}_{\text{fix}}(s, \bar{\alpha}, N) = \frac{\int_0^\infty t_{\text{fix}}(\tilde{s}(\alpha), N) P_{\text{fix}}(\tilde{s}(\alpha), N) f(\alpha) \, d\alpha}{\int_0^\infty P_{\text{fix}}(\tilde{s}(\alpha), N) f(\alpha) \, d\alpha} \,. \tag{4.14}$$

Here, $t_{\text{fix}}(\tilde{s}(\alpha), N)$ denotes the mean time to fixation of a single mutant with effect α and $P_{\text{fix}}(\tilde{s}(\alpha), N)$ its probability of fixation, both in a population of size N. For comparison with simulation results, we use the well-known diffusion approximation (B.1) for P_{fix} and the simplified version (B.4) of the diffusion approximation for t_{fix}. Details and an efficient numerical procedure to compute $\bar{t}_{\text{fix}}(s, \bar{\alpha}, N)$ are described in Appendix B.2. We note from (B.1) and (3.11) that $P_{\text{fix}}(\tilde{s}(\alpha), N) > 1 - e^{2\tilde{s}(\alpha)} > P_{\text{sur}}(e^{s\alpha})$ holds for every N. We emphasize that we distinguish between the fixation probability P_{fix} in the diffusion approximation of the Wright-Fisher model and the survival probability P_{sur} in the Galton-Watson process, although they are quantitatively very similar if Ns is sufficiently large and s small. If f is exponential with mean 1, a very accurate approximation for $\bar{t}_{\text{fix}}(\tilde{s}(\alpha), N)$ is $(2/s)\ln(2Ns)$ if $2/N \le s \le 0.2$, which is also a close approximation for $t_{\text{fix}}(s, N)$ in this range (Appendix B.2).

To proceed, we introduce

$$\tau_c = 1/\Theta + \bar{t}_{\text{fix}} \,, \tag{4.15}$$

the expected time by which the first mutation, which appeared at (about) $\tau = 1/\Theta$, becomes fixed. If $\tau \ge \tau_c$, then from the $\Theta\tau$ mutations, which are expected to arise until generation τ, $\Theta(\tau - \tau_c) + 1 = \Theta(\tau - \bar{t}_{\text{fix}})$ mutations are expected to have had enough time to go to fixation. Fixed mutations contribute α to $\bar{G}$ and 0 to V_G. Thus, the stationary phase, when the flux of incoming and fixing mutations balances is reached soon after about τ_c generations. The mutants from the remaining $\Theta\tau_c - 1 = \Theta\bar{t}_{\text{fix}}$ mutation events either segregate in the population or have been lost before generation τ. Segregating mutations contribute α to $\bar{G}$ and α^2 to V_G. We denote the values of $\bar{G}(\tau)$ and $V_G(\tau)$ in the stationary phase by $\bar{G}^*$ and V_G^*, respectively.

For notational simplicity, we set

$$\nu = \nu(s, \alpha, N) = N P_{\text{sur}}(e^{s\alpha}) \tag{4.16}$$

Here is our first main result.

Proposition 4.5. *At stationarity, when $\tau \gg \tau_c$, the phenotypic mean and the phenotypic variance*

can be approximated as follows:

$$\bar{G}^*(\tau, f, s, N, \Theta) \approx \Theta \int_0^\infty P_{\text{sur}}(e^{s\alpha})\frac{1}{s}\left(\exp(\nu e^{-s\alpha\bar{t}_{\text{fix}}})E_1(\nu e^{-s\alpha\bar{t}_{\text{fix}}}) - e^\nu E_1(\nu)\right)f(\alpha)\,d\alpha$$

$$+ \Theta(\tau - \bar{t}_{\text{fix}})\int_0^\infty \alpha P_{\text{sur}}(e^{s\alpha})f(\alpha)\,d\alpha \tag{4.17}$$

and

$$V_G^*(f, s, N, \Theta) \approx \Theta \int_0^\infty \frac{\alpha}{s}P_{\text{sur}}(e^{s\alpha})\nu e^\nu E_1(\nu)f(\alpha)\,d\alpha\,. \tag{4.18}$$

The proofs of this Proposition and of the Corollary below are relegated to Sect. 4.5.

Remark 4.6. For mutation effects equal to α, (4.18) simplifies to

$$V_G^*(\alpha, s, N, \Theta) \approx \Theta\frac{\alpha}{s}P_{\text{sur}}(e^{s\alpha})\nu e^\nu E_1(\nu)\,. \tag{4.19}$$

An analogous simplification applies to (4.17).

Remark 4.7. 1. The first term in (4.17) accounts for the contribution of the segregating mutations to the phenotypic mean. On average, these are the mutations that appeared less than $\bar{t}_{\text{fix}}$ generations in the past and had not yet enough time to become fixed. A permanent response of the mean is achieved by mutations that have gone to fixation. These are accounted for by the second term in (4.17), which is independent of the population size. Consequently, during the stationary phase, the phenotypic mean increases linearly with time as mutants reach fixation.

2. Because only segregating mutations contribute to the phenotypic variance, the number of segregating sites and the phenotypic variance remain constant over time once the stationary phase has been reached. The term $\frac{\alpha}{s}\nu e^\nu E_1(\nu)$ in (4.18) and (4.19) is precisely the total variance accumulated by a mutant with effect α during its sweep to fixation. This follows immediately from (4.4) and (C.3b).

The dashed straight lines in Figure 4.2 show that (4.17) and (4.18) provide excellent approximations for the Wright-Fisher simulations; see also Fig. 4.3. The following result provides much simpler approximations and not only recovers but refines well known results from evolutionary quantitative genetics. For $n \geq 2$ we write $\alpha_n(f) = \int_0^\infty \alpha^n f(\alpha)d\alpha$ for the nth moment about zero of the mutation distribution f. We remind the reader that we require $Ns \gg 1$, so that the branching process serves as an approximation for the Wright-Fisher model, and $s \ll 1$, so that also the diffusion approximation applies and the branching process is slightly supercritical.

Corollary 4.8. *In the stationary phase and for a Poisson offspring distribution, the following simple approximations hold.*

(1) The per-generation change of the mean phenotype at stationarity can be approximated as

follows:

$$\Delta \bar{G}^* \approx \Theta \int_0^\infty \alpha P_{\mathrm{sur}}(e^{s\alpha}) f(\alpha)\, d\alpha \tag{4.20a}$$

$$\approx 2\Theta s \left(\alpha_2(f) - \frac{5s}{6} \alpha_3(f) \right) . \tag{4.20b}$$

(2) For mutation effects equal to α, the change in the mean is

$$\Delta \bar{G}^* \approx 2\Theta s \alpha^2 (1 - \tfrac{5}{6} s\alpha) . \tag{4.21}$$

(3) For an exponential distribution f of mutation effects with expectation $\bar{\alpha}$, the change in the mean is

$$\Delta \bar{G}^* \approx 4\Theta s \bar{\alpha}^2 (1 - \tfrac{5}{2} s\bar{\alpha}) . \tag{4.22}$$

(4) The stationary variance has the approximation

$$V_G^*(f, s, N, \Theta) \approx \frac{1}{s} \Delta \bar{G}^* - \frac{\Theta \bar{\alpha}}{Ns} . \tag{4.23}$$

(5) If mutation effects are equal to α, then (4.23) and (4.21) yield

$$V_G^* \approx 2\Theta \alpha^2 \left(1 - \frac{5}{6} s\alpha - \frac{1}{2Ns\alpha} \right) . \tag{4.24}$$

(6) If f is exponential (with mean $\bar{\alpha}$), then (4.23) and (4.22) yield

$$V_G^* \approx 4\Theta \bar{\alpha}^2 \left(1 - \frac{5}{2} s\bar{\alpha} - \frac{1}{4Ns\bar{\alpha}} \right) . \tag{4.25}$$

Remark 4.9. (1) The approximations in (4.20a) and (4.23) hold for general offspring distributions. More explicit approximations can be derived from series expansions of the corresponding survival probabilities $P_{\mathrm{sur}}(e^{s\alpha})$. For weak selection, $P_{\mathrm{sur}}(e^{s\alpha})$ has typically an approximation of the form $cs\alpha$ for some $c > 0$. For instance, $c = 1$ for a geometric offspring distribution, and $c = 2$ for a Poisson distribution.

(2) The term $\frac{\Theta \bar{\alpha}}{Ns}$ in (4.23) term describes the reduction in variance caused by random drift. As is well known and also confirmed here, the latter does not affect $\Delta \bar{G}$ in the stationary phase. The derivation of (4.23) in (4.43) shows that, except for higher-order terms in $1/N$, this relation holds for general mutation distributions f.

In Figure 4.3 the approximations derived in Proposition 4.5 and Corollary 4.8 (shown as curves) are compared with results from Wright-Fisher simulations (shown as symbols). The figure shows that the approximation (4.18) for the stationary variance (with its simplification (4.19) for equal mutation effects) is highly accurate in a parameter range containing $5/N \le s\bar{\alpha} \le 0.5$ (dotted curves in A and B), although it has been derived under the assumption $Ns\bar{\alpha} \gg 1$. The corresponding

Figure 4.3: The scaled expected phenotypic variance V_G^*/Θ in the stationary phase for mutation effects equal 1 (panel A) and for exponentially distributed effects with mean 1 (panel B). The solid curves show the simple approximations (4.24) (in A) and (4.25) (in B) as functions of the selection coefficient s for three population sizes N (see legend). The most accurate available approximations are (4.19) and (4.18); they are shown as dotted curves in A and B, respectively. Simulation results from the Wright-Fisher model with $\Theta = \frac{1}{2}$ are shown as symbols (circles for $N = 10^4$, squares for $N = 10^3$, triangles for $N = 100$). In addition, the scaled expected per-generation change in the phenotypic mean, $\Delta \bar{G}^*/(\Theta s)$, is displayed (black dashed curves); it as calculated from (4.21) and (4.22) in (A) and (B), respectively. For the Wright-Fisher simulations, filled symbols are used for the scaled phenotypic variance and open symbols for the change in the scaled phenotypic mean. Note that in a Wright-Fisher model, $\Delta \bar{G}^*/(\Theta s)$ diverges to infinity if $s \to 0$ because neutral mutations can become fixed. Analogous results for a truncated normal distribution ($\alpha \geq 0$) with mean $\bar{\alpha} = 1$ are shown in Figure S.4.

simplified approximations (4.24) and (4.25) (solid curves), derived under the additional assumption

$s \ll 1$, are accurate in a range containing $5/N \leq s\bar{\alpha} \leq 0.1$. Whereas these approximations

correctly predict a decrease in V_G^*/Θ as N decreases, the analytically derived dependence of V_G^*/Θ

on s (proportional to $(Ns)^{-1}$) exaggerates the decrease of variance for small s observed in the

Wright-Fisher simulations. Indeed, in the limit of $s \to 0$ (neutrality), V_G^*/Θ should converge to

(approximately) α^2 if mutation effects are equal to α, and to $2\bar{\alpha}^2$ if they are exponentially distributed

with mean $\bar{\alpha}$. This follows from the classical result (e.g., Clayton and Robertson, 1955, Lynch and

Hill, 1986) that in the absence of selection, the stationary variance is to a close approximation

$2\Theta \int_{-\infty}^{\infty} \alpha^2 f(\alpha) d\alpha$, where f is the density of a mutation distribution with arbitrary mean and this

Θ includes mutations with positive and negative effect. The reason for the discrepancy is that for

weak selection ($Ns\bar{\alpha} \leq 1$), fixations are increasingly driven by random genetic drift which is not

reflected by the branching-process approximation. Finally, we note that the decrease in variance for

small s would not occur if we chose mutations from a symmetric distribution with positive and

negative effects. For large Ns such a choice would make almost no difference because strongly

deleterious mutations are lost very quickly. In the limit of $Ns \to 0$, however, the variance would be

approximately twice as high as with exclusively beneficial mutations because then both types of

mutations contribute nearly equally to the variance and even have similar chances of becoming

fixed.

The approximations (4.21) and (4.22) for $\Delta \bar{G}^*$ (black dashed curves in panels A and B of Fig. 4.3) accurately match the response calculated from Wright-Fisher simulations (open symbols) in a range containing $2/N \leq s\bar{\alpha} \leq 0.1$. If higher-order terms in s had been included, then these approximations would be accurate for $s\bar{\alpha}$ up to 0.5. (The dotted curves for the variance are accurate up to this value because they are computed from (4.19) and (4.18), which do not use series expansion in s.) Comparison of the terms of order s^2 in the approximations (4.21) and (4.22) with the term $\frac{\Theta\alpha}{Ns}$ in (4.23) shows that the latter can be neglected in (4.23) if $cs^2\bar{\alpha}^2 \gg 1/N$, where $c = \frac{5}{3}$ for equal mutation effects and $c = 10$ for exponentially distributed effect. The other way around, random genetic drift will distort the classical relation $\Delta\bar{G}^* \approx sV_G^*$ between the response of the mean and the genetic variance if (approximately) $s\bar{\alpha} < \sqrt{10/(cN)}$. For equal mutation effects ($c = \frac{5}{3}$), this becomes $s\bar{\alpha} < 2.4/\sqrt{N}$; for exponentially distribution effects ($c = 10$), it becomes $s\bar{\alpha} < 1/\sqrt{N}$. Both relations confirm the observations in Fig. 4.3, where $\bar{\alpha} = 1$.

Remark 4.10. In quantitative genetic models often a normal distribution with mean 0 is assumed for the mutation effects. To apply our analysis, which ignores mutations with a negative effect, to this setting, we have to adapt this distribution and use a truncated normal distribution instead. If the original normal distribution has variance v^2, then the truncated normal (restricted to positive effects) is defined by $f(\alpha) = \frac{2}{\sqrt{2\pi v^2}} \exp(-\alpha^2/(2v^2))$ if $\alpha \geq 0$ (and 0 otherwise). Also our mutation rate Θ will correspond to $\Theta/2$ in a quantitative genetic model in which mutations have (symmetric) positive and negative effects. This truncated normal distribution has mean $\bar{\alpha} = \sqrt{2v^2/\pi}$. Taking this as the parameter, we obtain $\alpha_2(f) = v^2 = \frac{1}{2}\bar{\alpha}^2\pi$ and $\alpha_3(f) = \bar{\alpha}^3\pi$. Therefore, instead of (4.22), (4.20b) yields

$$\Delta\bar{G}^* \approx \pi\Theta s\bar{\alpha}^2(1 - \tfrac{5}{3}s\bar{\alpha}), \tag{4.26}$$

and the stationary variance is approximately

$$V_G^* \approx \pi\Theta\bar{\alpha}^2 \left(1 - \frac{5}{3}s\bar{\alpha} - \frac{1}{\pi Ns\bar{\alpha}}\right). \tag{4.27}$$

After this excursion to the long-term evolution of the trait, we now investigate the initial phase of adaptation due to new mutations.

4.4. Approximations for the initial phase

In the initial phase, when $\tau < \tau_c$, on average no mutation had enough time to reach fixation. Therefore, the phenotypic variance (and the number of segregating sites) increases with the number of generations as more mutation events occur in the population. We define $\tau_1 = \lfloor 1/\Theta + 18e^{-5s} \rceil$, where $\lfloor x \rceil$ denotes the smallest integer $\geq x$. Then at least one mutation has occurred (on average) and $1 - P_{\text{ext}}(\tau, e^{s\alpha})$ can be approximated by $P_{\text{sur}}(e^{s\alpha})$ if $\tau \geq \tau_1$ (see Sect. 3.2). Better insight into the evolution of the phenotypic mean and the phenotypic variance than from Proposition 4.2 can be gained with the help of the following simplified approximations.

Proposition 4.11. *If $\tau_1 < \tau < \tau_c$, the phenotypic mean and the phenotypic variance can be approximated by*

$$\bar{G}(\tau, f, s, N, \Theta) \approx \Theta \int_0^\infty \frac{1}{s} P_{\text{sur}}(e^{s\alpha}) \Big(\exp(\nu e^{-s\alpha\tau}) E_1(\nu e^{-s\alpha\tau}) - e^\nu E_1(\nu) \Big) f(\alpha)\, d\alpha \qquad (4.28)$$

and

$$V_G(\tau, f, s, N, \Theta) \approx \Theta \int_0^\infty \frac{\alpha}{s} P_{\text{sur}}(e^{s\alpha}) \Big(\nu e^\nu E_1(\nu) - \nu e^{-s\alpha\tau} \exp(\nu e^{-s\alpha\tau}) E_1(\nu e^{-s\alpha\tau}) \Big) f(\alpha)\, d\alpha \,, \qquad (4.29)$$

where $\nu = N P_{\text{sur}}(e^{s\alpha})$.

The proofs of this and the following Proposition are relegated to Sect. 4.5. For a Poisson offspring distribution an insightful simplification is obtained by applying $P_{\text{sur}}(e^{s\alpha}) \approx 2s\alpha$. It demonstrates that mean and variance are not proportional to $1/s$, and the 'correct' exponents of α in the integrands become clearly visible.

Remark 4.12. For equal mutation effects α, the approximations (4.28) and (4.29) simplify to

$$\bar{G}(\tau, f, s, N, \Theta) \approx \frac{\Theta}{s} P_{\text{sur}}(e^{s\alpha}) \Big(\exp(\nu e^{-s\alpha\tau}) E_1(\nu e^{-s\alpha\tau}) - e^\nu E_1(\nu) \Big) \qquad (4.30)$$

and

$$V_G(\tau, f, s, N, \Theta) \approx \frac{\Theta\alpha}{s} P_{\text{sur}}(e^{s\alpha}) \Big(\nu e^\nu E_1(\nu) - \nu e^{-s\alpha\tau} \exp(\nu e^{-s\alpha\tau}) E_1(\nu e^{-s\alpha\tau}) \Big) \,, \qquad (4.31)$$

respectively. Again, it is insightful to keep in mind that for a Poisson offspring distribution $P_{\text{sur}}(e^{s\alpha}) \approx 2s\alpha$.

For the very early phase of adaptation much simpler approximations can be derived.

Proposition 4.13. *We assume that $\tau \ll (\ln N)/(s\bar{\alpha})$ and the offspring distribution is Poisson.*

(1) If f is exponential with mean $\bar{\alpha}$, the phenotypic mean and the phenotypic variance have the approximations

$$\bar{G}(\tau) \approx \frac{\Theta}{N} \frac{\bar{\alpha}\tau}{1 - s\bar{\alpha}\tau} \qquad (4.32a)$$

$$\approx \frac{\Theta\bar{\alpha}\tau}{N} \Big(1 + s\bar{\alpha}\tau + (s\bar{\alpha}\tau)^2 + (s\bar{\alpha}\tau)^3 \Big) \qquad (4.32b)$$

and

$$V_G(\tau) \approx \frac{\Theta\bar{\alpha}}{Ns} \left(\frac{1}{(1 - s\bar{\alpha}\tau)^2} - 1 \right) \qquad (4.33a)$$

$$\approx \frac{2\Theta\bar{\alpha}^2\tau}{N} \Big(1 + \tfrac{3}{2} s\bar{\alpha}\tau + 2(s\bar{\alpha}\tau)^2 + \tfrac{5}{2}(s\bar{\alpha}\tau)^3 \Big) \,. \qquad (4.33b)$$

Figure 4.4: The scaled phenotypic mean $\bar{G}(s\tau)/\Theta$ (panels A, C) and the scaled phenotypic variance $V_G(s\tau)/\Theta$ (panels B, D) in the initial phase for selection coefficients s as given in the legend. For equal mutation effects $\alpha = 1$, the solid curves in A and B show the approximations (4.30) and (4.31), respectively. The dashed curves show the simplified approximation in (4.34). For exponentially distributed effects with $\bar{\alpha} = 1$ the solid curves in C and D show the approximations (4.28) and (4.29), respectively. The dashed curves show the simplified approximations (4.32b) and (4.33b). In all cases, symbols present results from Wright-Fisher simulations with $\Theta = \frac{1}{2}$, the population size is $N = 10^4$, and the offspring distribution is Poisson.

(2) If the mutation effects are equal to α, then

$$\bar{G}(\tau) \approx \frac{1}{\alpha} V_G(\tau) \approx \frac{\Theta}{Ns} \left(e^{s\alpha\tau} - 1\right) . \tag{4.34}$$

In Figure 4.4 the approximations for the expected phenotypic mean and variance obtained in Propositions 4.11 and 4.13 are compared with Wright-Fisher simulations for various selection coefficients. For equal mutation effects (panels A and B) the approximations are accurate in a much wider range of values $s\tau$ than for exponentially distributed effects (panels C and D). In both cases, the more complicated approximations, (4.30) and (4.31) (solid curves in A and B) and (4.28) and (4.29) (solid curves in C and D), are accurate for a much larger time span than the corresponding simple approximations, (4.34) (dashed curves in A, B) and (4.32b), (4.33b) (dashed curves in C, D). For an exponential mutation distribution, the simple approximations are accurate if, approximately, $\tau < 1/(2s\bar{\alpha})$. For weak selection, this can still be quite a long time span.

 4.5. Proofs

726   Here, we give the proofs of the main results of Section 4.

727   *Proof of Proposition 4.2.* We will need the following integrals. Let $\phi(x)$ be a (continuous) function.

728   Then we obtain directly from the definition of $\tilde{h}^{(i)}$ by a change in the order of integration:

$$\int_0^1 \phi(x)\tilde{h}^{(i)}_{\tau,\sigma_i,\Theta}(x)\,dx = \int_0^\tau \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)}\left(1 - P_{\mathrm{ext}}([\tau-t],\sigma_i)\right)\int_0^1 \phi(x)g_{a(\tau-t,\sigma_i)}(x)\,dx\,dt\,. \quad (4.35)$$

729   With $\phi(x) = x$, we get from (4.1a) and (4.35)

$$\int_0^1 x\tilde{h}^{(i)}_{\tau,\sigma_i,\Theta}(x)\,dx = \int_0^\tau \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)}\left(1 - P_{\mathrm{ext}}([\tau-t],\sigma_i)\right)\gamma_1(a(\tau-t,\sigma_i))\,dt\,. \quad (4.36)$$

730   With $\phi(x) = x(1-x)$, we get from (4.1c) and (4.35)

$$\int_0^1 x(1-x)\tilde{h}^{(i)}_{\tau,\sigma_i,\Theta}(x)\,dx = \int_0^\tau \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)}\left(1 - P_{\mathrm{ext}}([\tau-t],\sigma_i)\right)\gamma(a(\tau-t,\sigma_i))\,dt\,. \quad (4.37)$$

731   We will also need

$$\sum_{n=1}^\infty \mathrm{Poi}_{\Theta\tau}(n)\left(\sum_{i=1}^n \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)}\right) = \Theta\,, \quad (4.38)$$

732   as is proved in Appendix C.1.

733   From the definition (4.7) of $\bar{G}(\tau)$ we deduce:

$$\bar{G}(\tau) = \sum_{n=1}^\infty \mathrm{Poi}_{\Theta\tau}(n)\left(\sum_{i=1}^n \int_0^\infty \alpha\,\mathrm{E}\left[\tilde{X}^{(i)}_{\tau,e^{s\alpha},\Theta}\right]f(\alpha)\,d\alpha\right) \quad (4.39a)$$

$$= \sum_{n=1}^\infty \mathrm{Poi}_{\Theta\tau}(n)\int_0^\infty \alpha f(\alpha)\left(\sum_{i=1}^n \int_0^1 x\tilde{h}^{(i)}_{\tau,e^{s\alpha},\Theta}(x)\,dx\right)d\alpha \quad (4.39b)$$

$$= \sum_{n=1}^\infty \mathrm{Poi}_{\Theta\tau}(n)\int_0^\infty \alpha f(\alpha)$$
$$\times \int_0^\tau \left(\sum_{i=1}^n \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)}\right)\left(1 - P_{\mathrm{ext}}([\tau-t],e^{s\alpha})\right)\gamma_1(a(\tau-t,e^{s\alpha}))\,dt\,d\alpha\,. \quad (4.39c)$$

734   Here, we obtained (4.39a) by using that the mutation effects α_i are drawn independently from

735   the same distribution f; (4.39b) by using (4.6); and (4.39c) by employing (4.36) and changing

736   the order of integration and summation. Finally, (4.39c) yields (4.9) by applying (4.38). Note also

737   that we suppressed the dependence on N in $\tilde{X}^{(i)}$, $\tilde{h}^{(i)}$, and a.

738   Analogously, by using (4.37) instead of (4.36), we obtain for the variance from its definition

 (4.8),

$$V_G(\tau) = \sum_{n=1}^{\infty} \text{Poi}_{\Theta\tau}(n) \left(\sum_{i=1}^{n} \int_0^{\infty} \alpha^2 \, \text{E}\left[\tilde{X}^{(i)}_{\tau,e^{s\alpha},\Theta}(1 - \tilde{X}^{(i)}_{\tau,e^{s\alpha},\Theta}) \right] f(\alpha) \, d\alpha \right)$$

$$= \sum_{n=1}^{\infty} \text{Poi}_{\Theta\tau}(n) \int_0^{\infty} \alpha^2 f(\alpha) \left(\sum_{i=1}^{n} \int_0^1 x(1-x) \tilde{h}^{(i)}_{\tau,e^{s\alpha},\Theta}(x) \, dx \right) d\alpha$$

$$= \sum_{n=1}^{\infty} \text{Poi}_{\Theta\tau}(n) \int_0^{\infty} \alpha^2 f(\alpha)$$

$$\times \int_0^{\tau} \left(\sum_{i=1}^{n} \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)} \right) (1 - P_{\text{ext}}([\tau - t], \sigma_i)) \gamma(a(\tau - t, \sigma_i)) \, dt \,, \tag{4.40}$$

740 which yields (4.10) with the help of (4.38). Note that for the derivation of the variance, the
741 independence of the random variables $\tilde{X}^{(i)}$ is crucial. $\qquad\square$

742 *Proof of Proposition 4.5.* We assume $\tau \gg \bar{t}_{\text{fix}}$ (in particular, $\tau > \tau_c$) and suppress the dependence
743 of a and ν on N. Then we obtain from (4.9)

$$\bar{G}(\tau) = \Theta \int_0^{\infty} \alpha f(\alpha) \int_0^{\tau} (1 - P_{\text{ext}}([\tau - t], e^{s\alpha})) \gamma_1(a(\tau - t, e^{s\alpha})) \, dt \, d\alpha \tag{4.41a}$$

$$\approx \Theta \int_0^{\infty} \alpha f(\alpha)(1 - P_{\text{ext}}^{\infty}(e^{s\alpha})) \int_0^{\tau} \gamma_1(a(\tau - t, e^{s\alpha})) \, dt \, d\alpha \tag{4.41b}$$

$$= \Theta \int_0^{\infty} \alpha f(\alpha) P_{\text{sur}}(e^{s\alpha}) \left(\int_0^{\tau - \bar{t}_{\text{fix}}} \gamma_1(a(\tau - t)) \, dt + \int_{\tau - \bar{t}_{\text{fix}}}^{\tau} \gamma_1(a(\tau - t)) \, dt \right) d\alpha \tag{4.41c}$$

$$\approx \Theta(\tau - \bar{t}_{\text{fix}}) \int_0^{\infty} \alpha f(\alpha) P_{\text{sur}}(e^{s\alpha}) \, d\alpha$$

$$+ \Theta \int_0^{\infty} f(\alpha) \frac{1}{s} P_{\text{sur}}(e^{s\alpha}) \left[\exp(\nu e^{-s\alpha \bar{t}_{\text{fix}}}) E_1(\nu e^{-s\alpha \bar{t}_{\text{fix}}}) - e^{\nu} E_1(\nu) \right] d\alpha \,. \tag{4.41d}$$

744 Here, we used $P_{\text{ext}}([\tau - t], e^{s\alpha}) \approx P_{\text{ext}}^{\infty}(e^{s\alpha})$ for sufficiently large τ to obtain (4.41b), a simple
745 integral splitting (and $P_{\text{sur}} = 1 - P_{\text{ext}}^{\infty}$) for (4.41c), and finally (C.6) and (C.5) to arrive at (4.41d),
746 which equals (4.17).

747   For the variance, we start from (4.10) and then apply (C.3b) to obtain the asymptotic variance:
748

$$V_G(\tau) \approx \Theta \int_0^{\infty} \alpha^2 f(\alpha) \int_0^{\tau} (1 - P_{\text{ext}}([\tau - t], e^{s\alpha})) \gamma(a(\tau - t, e^{s\alpha})) \, dt \, d\alpha$$

$$\approx \Theta \int_0^{\infty} \alpha^2 f(\alpha) P_{\text{sur}}(e^{s\alpha}) \int_0^{\tau} \gamma(a(\tau - t, e^{s\alpha})) \, dt \, d\alpha \tag{4.42a}$$

$$\xrightarrow[\tau \to \infty]{} \Theta \int_0^{\infty} \frac{\alpha}{s} f(\alpha) P_{\text{sur}}(e^{s\alpha}) \nu e^{\nu} E_1(\nu) \, d\alpha \,. \tag{4.42b}$$

749 This is (4.18). $\qquad\square$

750 *Proof of Corollary 4.8.* (1) From (4.17) we obtain immediately that the per-generation change of

the mean is $\Delta \bar{G}^* = \Theta \int_0^\infty \alpha P_{\text{sur}}(e^{s\alpha}) f(\alpha)\, d\alpha$, which is (4.20a). Therefore, (4.20b) follows from (3.11) by using $P_{\text{sur}}(e^{s\alpha}) \approx 2s\alpha - \frac{5}{3}(s\alpha)^2$.

(2) For equal effects, (4.20b) simplifies immediately to (4.21).

(3) follows from the properties of the exponential distribution.

(4) Using $\nu e^\nu E_1(\nu) \approx 1 - 1/\nu$ and $\nu = N P_{\text{sur}}(e^{s\alpha})$ in (4.18) we obtain

$$V_G^*(f, s, N, \Theta) \approx \Theta \int_0^\infty \frac{\alpha}{s} P_{\text{sur}}(e^{s\alpha})\Big(1 - \frac{1}{\nu}\Big) f(\alpha)\, d\alpha \tag{4.43a}$$

$$= \frac{\Theta}{s} \int_0^\infty \alpha P_{\text{sur}}(e^{s\alpha}) f(\alpha)\, d\alpha - \frac{\Theta}{sN} \int_0^\infty \alpha f(\alpha)\, d\alpha\,, \tag{4.43b}$$

which yields (4.23).

We prove the exponential case (6) before (5).

(6) In (4.18), we substitute $2Ns\alpha$ for ν and $2s\alpha - \frac{5}{3}(s\alpha)^2$ for $P_{\text{sur}}(e^{s\alpha})$ to obtain

$$V_G^*(f, s, N, \Theta) \approx 4\Theta Ns \int_0^\infty \alpha^3 \Big(1 - \frac{5}{6}s\alpha\Big) e^{2Ns\alpha} E_1(2Ns\alpha) f(\alpha)\, d\alpha \tag{4.44a}$$

$$= \frac{4\Theta Ns\bar{\alpha}^3}{(2Ns\bar{\alpha} - 1)^4}\Big[(2Ns\bar{\alpha} - 1)\big(8(Ns\bar{\alpha})^2 - 14Ns\bar{\alpha} + 11\big) - 6\ln(2Ns\bar{\alpha})\Big]$$

$$- \frac{20\Theta Ns^2\bar{\alpha}^4}{3(2Ns\bar{\alpha} - 1)^5}\Big[(2Ns\bar{\alpha} - 1)\big(24(Ns\bar{\alpha})^3 - 52(Ns\bar{\alpha})^2 + 46Ns\bar{\alpha} - 25\big) + 12\ln(2Ns\bar{\alpha})\Big]\,, \tag{4.44b}$$

where the integration has been performed with *Mathematica*. Now we fix s and let $Ns \to \infty$. Then, to first order in $1/(Ns\bar{\alpha})$, $s\bar{\alpha}$, and $1/N$, we obtain

$$V_G^*(f, s, N, \Theta) \approx 4\Theta\bar{\alpha}^2 \Big(1 - \frac{5}{2}s\bar{\alpha} - \frac{1}{4Ns\bar{\alpha}} + \frac{5}{12N}\Big)\,. \tag{4.45}$$

This yields (4.25) because $1/N \ll s\bar{\alpha}$ and $1/N \ll 1/(Ns\bar{\alpha})$.

(5) For equal effects, (4.44a) simplifies to $4\Theta Ns\alpha^3\big(1 - \frac{5}{6}s\alpha\big)e^{2Ns\alpha}E_1(2Ns\alpha)$. Using $e^x E_1(x) \approx x^{-1} - x^{-2}$ if $x \to \infty$, we obtain

$$V_G^*(\alpha, s, N, \Theta) \approx 2\Theta\alpha^2 \Big(1 - \frac{5}{6}s\alpha - \frac{1}{2Ns\alpha} + \frac{5}{12N}\Big)\,, \tag{4.46}$$

which yields (4.24), in which the term of smallest order, $\frac{5}{12N}$, has been omitted. $\qquad\square$

Proof of Proposition 4.11. We observe that we obtain (4.41b) from (4.9) if $\tau \geq \tau_1$. Then we apply (C.2a) with $t_1 = \tau$ to arrive at (4.28). The approximation (4.29) is obtained analogously from (4.42a), which follows from (4.10) if $\tau \geq \tau_1$, by employing (C.3a) instead of (C.2a). $\qquad\square$

Proof of Proposition 4.13. From the well-known asymptotic expansion $e^a E_1(a) = \sum_{k=0}^\infty (-1)^k \frac{k!}{a^{k+1}}$ the approximation $ae^a E_1(a) \approx 1 - 1/a$ follows if $a \gg 1$. Upon recalling the definition of $a(t, \sigma, N)$ in (3.16b), we find that $a \gg 1$ holds if $N \gg \sigma^\tau$, or $\tau \ll (\ln N)/(s\bar{\alpha})$ (see also Fig. S.1). Therefore,

we obtain from (4.1a),

$$\gamma_1(a(\tau - t, e^{s\alpha}, N)) \approx \frac{1}{a(\tau - t, e^{s\alpha}, N)} = \frac{e^{s\alpha(\tau - t)}}{N(1 - P_{\text{ext}}([\tau - t], e^{s\alpha}))} \tag{4.47}$$

for small τ.

Substituting this into (4.9), we obtain directly

$$\begin{aligned}
\bar{G}(\tau) &= \Theta \int_0^\infty \alpha f(\alpha) \int_0^\tau (1 - P_{\text{ext}}([\tau - t], e^{s\alpha}))\gamma_1(a(\tau - t, e^{s\alpha}))\, dt\, d\alpha \\
&\approx \frac{\Theta}{N} \int_0^\infty \alpha f(\alpha) \int_0^\tau e^{s\alpha[\tau - t]}\, dt\, d\alpha \\
&\approx \frac{\Theta}{Ns} \int_0^\infty f(\alpha)(e^{s\alpha\tau} - 1)\, d\alpha,
\end{aligned} \tag{4.48}$$

which yields (4.32a) if $f(\alpha) = \frac{1}{\bar{\alpha}}e^{-\alpha/\bar{\alpha}}$ provided $s\bar{\alpha}\tau < 1$ (otherwise, the integral diverges).

In a similar way, we get from (4.1c) that $\gamma(a) = 1/a + O(a^2)$ for large a. By substituting this into (4.10), we obtain

$$\begin{aligned}
V_G(\tau) &= \Theta \int_0^\infty \alpha^2 f(\alpha) \int_0^\tau (1 - P_{\text{ext}}([\tau - t], e^{s\alpha}))\gamma(a(\tau - t, e^{s\alpha}))\, dt\, d\alpha \\
&\approx \frac{\Theta}{N} \int_0^\infty \alpha^2 f(\alpha) \int_0^\tau e^{s\alpha[\tau - t]}\, dt\, d\alpha \\
&\approx \frac{\Theta}{Ns} \int_0^\infty \alpha f(\alpha)(e^{s\alpha\tau} - 1)\, d\alpha,
\end{aligned} \tag{4.49}$$

which yields (4.33a) if f is exponential with mean $\bar{\alpha}$.

The approximations (4.32b) and (4.33b) are then obtained by a series expansion. If the mutation effects are equal to α, (4.34) follows. $\qquad\square$

5. Segregating sites

An important component in examining whether sweeps or shifts determine the pattern of adaptation will be the number of segregating sites (see Section 6). If sweeps occur successively, at most one segregating site is expected for most of the time. As the number of segregating sites increases, parallel sweeps occur and the sweep will be transformed to a shift pattern. If adaptation is dominated by subtle allele frequency shifts, the number of segregating sites is expected to be large. Of course, there will be intermediate patterns, where no clear distinction between a sweep and a shift pattern can be made. To characterize the patterns of adaptation, an approximation for the expected number of segregating sites, $\mathrm{E}[S]$, is needed. This expectation is taken not only with respect to the distribution of the number of segregating sites, but also with respect to the mutation distribution f, and it assumes a Poisson offspring distribution. For a precise definition see Appendix D.

We assume the infinite sites model, as used in Sects. 4.2 – 4.5. However, we complement it by *diffusion approximations* for the fixation and the loss probabilities as well as for the corresponding expected times to fixation and loss (see Appendix B). To derive a simple approximation $\bar{S}$ for $E[S]$, we distinguish between mutations that eventually become fixed, which occurs with probability $P_{\text{fix}}(\tilde{s}(\alpha), N)$ (see (4.13) for $\tilde{s}(\alpha)$), and those that get lost, which occurs with probability $1 - P_{\text{fix}}(\tilde{s}(\alpha), N)$. In addition, we assume that mutations that are destined to fixation and occurred $t \geq t_{\text{fix}}(\tilde{s}(\alpha), N)$ generations in the past are already fixed. Thus, in a given generation τ, mutations destined to fixation contribute to $\bar{S}$ if and only if they occurred less or equal than $\min\{t_{\text{fix}}(\tilde{s}(\alpha), N) - 1, \tau\}$ generations in the past (including 0 generations in the past, i.e., now, in generation τ). Therefore, the number of generations in which mutants occur that contribute to the current variation is $\min\{t_{\text{fix}}(\tilde{s}(\alpha), N), \tau + 1\}$. This is consistent with the fact that mutations can occur already in generation $\tau = 0$. Analogously, mutations destined to loss contribute to $\bar{S}$ at time τ only if their time t of occurrence (prior to τ) satisfies $0 \leq t \leq \min\{t_{\text{loss}}(\tilde{s}(\alpha), N) - 1, \tau\}$, where $t_{\text{loss}}(s, N)$ denotes the expected time to loss of a single mutant with selective advantage s in population of size N; see (B.7). In summary, we define the approximation $\bar{S}$ as follows:

$$\bar{S}(\tau, f, s, N, \Theta) = \Theta \int_0^\infty f(\alpha) P_{\text{fix}}(\tilde{s}(\alpha), N) \min\{t_{\text{fix}}(\tilde{s}(\alpha), N), \tau + 1\}\, d\alpha$$
$$+ \Theta \int_0^\infty f(\alpha)(1 - P_{\text{fix}}(\tilde{s}(\alpha), N)) \min\{t_{\text{loss}}(\tilde{s}(\alpha), N), \tau + 1\}\, d\alpha. \qquad (5.1)$$

For the stationary as well as for the initial phase, we can specify relatively simple explicit approximations. We recall from (4.14) that $\bar{t}_{\text{fix}}(s, \bar{\alpha}, N)$ is the expected time to fixation of a single mutant with effect α on the trait drawn from the (exponential) distribution f; see also Appendix B.2. In the limit of large τ, i.e., $\tau \gg \bar{t}_{\text{fix}}$, we obtain from diffusion theory for $\bar{S}^*(f, s, N, \Theta) = \lim_{\tau \to \infty} \bar{S}(\tau, f, s, N, \Theta)$

$$\bar{S}^*(f, s, N, \Theta) \approx \Theta \int_0^\infty f(\alpha) P_{\text{fix}}(\tilde{s}(\alpha), N) t_{\text{fix}}(\tilde{s}(\alpha), N)\, d\alpha$$
$$+ \Theta \int_0^\infty f(\alpha)(1 - P_{\text{fix}}(\tilde{s}(\alpha), N)) t_{\text{loss}}(\tilde{s}(\alpha), N)\, d\alpha$$
$$\approx \begin{cases} 2\Theta\big[(1 - s\bar{\alpha})(\ln(N) + \ln(2Ns\bar{\alpha})) + 1 + \gamma - (\gamma + \frac{1}{2})s\bar{\alpha}\big] & \text{if } \alpha = \bar{\alpha}, \\ 2\Theta\big[(1 - s\bar{\alpha})(\ln(N) + \ln(2Ns\bar{\alpha})) + 1 - \frac{3}{2}s\bar{\alpha}\big] & \text{if } \alpha \sim \text{Exp}[1/\bar{\alpha}]. \end{cases}$$
$$(5.2)$$

To evaluate the integrals, we used $\tilde{s}(\alpha) \approx s\alpha$ from (4.13), $P_{\text{fix}}(s, N) \approx 1 - e^{-2s}$ from (B.1), $t_{\text{fix}}(s, N) \approx \frac{2}{s}(\ln(2Ns) + \gamma)$ from (B.2), and $t_{\text{loss}}(s, N) \approx -2(1 + s)\ln(2s) + 2(1 - \gamma) + s(3 - 2\gamma)$ from (B.7) and (B.5a). Finally, a series expansion for small s was performed. The approximate formulas in (5.2) are accurate if $Ns\bar{\alpha} > 2$ and $s\bar{\alpha} < 0.1$; see Figs. 5.1B and D.

We observe that in both cases in (5.2) the leading-order term is approximately $2\Theta \ln(N)$, which

Figure 5.1: The expected number of segregating sites (rescaled by Θ) as a function of $s\tau$ (panels A, C) and at stationarity as a function of s (panels B, D). In A and B the mutation effects are equal ($\alpha = 1$), in C and D they are drawn from an exponential distribution with mean $\bar{\alpha} = 1$. Analytic approximations are computed from (5.1) and are shown as curves. Wright-Fisher simulation results for $\mathrm{E}[S]$ (with $\Theta = \frac{1}{2}$) are shown as symbols. In panels A and C, $\bar{S}/\Theta$ is shown for $N = 10^4$ and $s = 0.1, 0.01, 0.001$ (solid curves and filled symbols in blue, red, black), and for $N = 10^3$ and $s = 0.1$ (dashed curve and open symbols in blue). In panels B and D, $\bar{S}^*/\Theta$ is displayed for $N = 10^4, 10^3, 10^2$ (solid, dashed, dotted curves, and corresponding open symbols). Black curves represent (5.1), red curves the approximation (5.2). Analogous results for a truncated normal distribution ($\alpha \geq 0$) with mean $\bar{\alpha} = 1$ are shown in Figure S.5.

is close to the classical neutral result $\mathrm{E}[S] = \Theta \sum_{i=1}^{N-1} \frac{1}{i} \approx \Theta(\ln(N) + \gamma)$, in which the factor 2 is contained in Θ (Ewens, 1974). We note that for equal mutation effects the stationary value $\bar{S}^*$ of $\mathrm{E}[S]$ is slightly higher than that for exponentially distributed effects.

In analogy to $\bar{t}_{\mathrm{fix}}(s, \bar{\alpha}, N)$, in Appendix B.3 we introduce the expected time to loss, $\bar{t}_{\mathrm{loss}}$, of a single mutant with effect α on the trait drawn from the (exponential) distribution f. If τ is small, but not tiny, i.e., $\bar{t}_{\mathrm{loss}} < \tau \ll \bar{t}_{\mathrm{fix}}$, then

$$\bar{S}(\tau, f, s, N, \Theta) \approx \Theta(\tau + 1) \int_0^\infty f(\alpha) P_{\mathrm{fix}}(\tilde{s}(\alpha), N) \, d\alpha$$
$$+ \Theta \int_0^\infty f(\alpha)(1 - P_{\mathrm{fix}}(\tilde{s}(\alpha), N)) t_{\mathrm{loss}}(\tilde{s}(\alpha), N) \, d\alpha$$

$$\approx \begin{cases} \Theta\tau(1 - e^{-2s\bar{\alpha}}) + 2\Theta\left[-(1 - s\bar{\alpha})(\ln(2s\bar{\alpha}) + \gamma) + 1 + \frac{1}{2}s\bar{\alpha}\right] & \text{if } \alpha = \bar{\alpha}\,, \\[3mm] \Theta\tau\dfrac{2s\bar{\alpha}}{1 + 2s\bar{\alpha}} + 2\Theta\left[-(1 - s\bar{\alpha})\ln(2s\bar{\alpha}) + 1 + \frac{3}{2}s\bar{\alpha}\right] & \text{if } \alpha \sim \mathrm{Exp}[1/\bar{\alpha}]\,, \end{cases} \tag{5.3}$$

where the same approximations as for (5.2) are used.

If $\tau < \bar{t}_{\mathrm{loss}}$, then (5.1) yields $\bar{S}(\tau, f, s, N, \Theta) \approx \Theta(\tau + 1)$, which is an overestimate except when $\tau = 0$ (cf. Fig. D.1).

As Figure 5.1 demonstrates, (5.1) yields a quite accurate approximation for the expected number of segregating sites, both as a function of time (panels A and C), and as function of s at stationarity (panels B and D). The piecewise linear shape of the curves in A results from the simplifying assumption of taking the minima of τ and t_{loss} or t_{fix}, i.e., the kinks occur at t_{loss} and at t_{fix}. In C, the curves are smooth because effects α are drawn from the distribution f. Panels A and C in Fig. 5.1 and, in more detail, Fig. D.1 show that as a function of $s\tau$ the rate of increase of the number of segregating sites is nearly independent of s, except for very early generations ($s\tau < 0.1$; cf. eq. 5.3). Therefore, if measured in generations, smaller values of s lead to a slower increase of the number of segregating sites, as is supported by the rough approximations in (5.3). The intuitive reason is that invasion becomes less likely. Panels B and D (but also A and C) show that the stationary value $\bar{S}^*$ of the number of segregating sites is maximized at an intermediate selection strength. For small values of s, $\bar{S}^*$ is increasing in s because larger s increases the probability of a sweep. For sufficiently large values of s, $\bar{S}^*$ drops again because selected mutants spend less time segregating.

Based on (5.2), it is straightforward to show that, as a function of s, the maximum value of $\bar{S}^*$ is achieved close to

$$s_{\mathsf{max}} = \frac{1}{\bar{\alpha}\,\mathrm{ProductLog}[2e^b N^2]} \approx \frac{1}{2\bar{\alpha}\ln(N)}\,, \tag{5.4}$$

where $b = \frac{3}{2} + 2\gamma \approx 2.65$ for equal effects and $b = \frac{5}{2}$ for exponentially distributed effects. In each case, the maximum values of $\bar{S}^*$ are very close to $2\Theta(2\ln(N) - \ln\ln(N))$. Thus, the maximum possible value of $\bar{S}^*$ under directional selection is about twice that under neutrality. This shows that once a stationary response to directional selection has been reached, the number of segregating sites depends only weakly on N, essentially logarithmically, and exceeds the neutral value by at most a factor of two, whatever the strength of selection. In summary, the by far most important factor determining the number of segregating sites at stationarity is Θ.

Our branching process model leads to an alternative approximation, $\tilde{S}$, for the expected number

of segregating sites. If all mutation effects are equal to α, the highly accurate approximation

$$\tilde{S}(\tau, \alpha, s, N, \Theta) \approx \Theta \sum_{j=0}^{[\tilde{\tau}]} (1 - P_{\text{ext}}(j, e^{s\alpha})) \tag{5.5}$$

is derived in Appendix D. Here, $\tilde{\tau} = \min\{t_{\text{fix}}(\tilde{s}(\alpha), N), \tau\}$, t_{fix} (eq. B.1) comes from the diffusion process, and P_{ext} (eq. 3.12b) from the branching process. For values $s\tau < 2$, this approximation is considerably more accurate than the approximation $\bar{S}$ given above. This is illustrated by Fig. D.1.A, which is based on a zoomed-in versions of panel A of Fig. 5.1. The disadvantage of the approximation (5.5) is that the terms $P_{\text{ext}}(j, e^{s\alpha})$ have to be computed recursively, whereas the above approximations $\bar{S}$ are based on analytically explicit expressions.

For mutation effects drawn from a distribution f, the following approximation of $\mathrm{E}[S]$ is derived in Appendix D under the assumption $\tau \ll \bar{t}_{\text{fix}}(\tilde{s}(\alpha), N)$:

$$\tilde{S}(\tau, f, s, \Theta) \approx \Theta \sum_{j=0}^{\tau} \left(1 - \int_0^\infty f(\alpha) P_{\text{ext}}(j, e^{s\alpha}) d\alpha\right). \tag{5.6}$$

As Figure D.1.B demonstrates, it is highly accurate if $s\tau < 2$. However, it is computationally more expensive than $\bar{S}$, which is based on the diffusion approximation and a considerable overestimate for such small values of τ.

Equations (5.5) and (5.6) show that the expected number of segregating sites is proportional to Θ. We already know that for large τ, i.e., in the stationary phase, the number of segregating sites depends at most logarithmically on N. The approximation (5.6) of $\tilde{S}$ is independent of N, and $\tilde{S}$ in (5.5) depends on N only through $\tilde{\tau}$, which can be assumed to equal τ if $\tau \ll \bar{t}_{\text{fix}}$. According to the approximation (B.15) of $\bar{t}_{\text{fix}}$, we have $\bar{t}_{\text{fix}} > 2.77/(s\bar{\alpha})$ if $Ns\bar{\alpha} \geq 2$, because $2 \ln 4 \approx 2.77$. Therefore, we can expect independence of $\tilde{S}(\tau)$ from N if $\tau \leq 2.7/(s\bar{\alpha})$. Comparison of the dashed blue curves with the solid blue curves in panels A and C of Fig. 5.1 shows that the expected number of segregating sites is essentially independent of N for a much longer time. Also the dependence of $\tilde{S}(\tau)$ on s is much weaker than that on Θ.

6. Discussion

The goal of our study has been to provide essential theory for answering the following questions: Which are the main evolutionary and genetic factors determining the rate and pattern of polygenic adaptation? How do these factors determine the response to selection, both on the phenotypic and on the genetic level? When is the selective response mainly due to sweeps at few loci and when is it due to small frequency shifts at many loci?

For this purpose, we developed and analyzed a simple population-genetic model, starting from a branching process approximation to describe the spreading of a beneficial mutant in a Wright-Fisher

population of size N. Our basic result is an accurate, analytically tractable, and computationally efficient approximation for the time dependence of the mutant-frequency distribution. Extension to an infinite sites model and the assumption of an additive quantitative trait under exponential directional selection enabled us to derive accurate approximations for the evolution of the expected mean phenotype, the expected phenotypic variance, and the expected number of segregating sites underlying the adaptive response. Our theory focuses on the response from new mutations and assumes that selection is stronger than random genetic drift ($Ns > 1$). In the sequel, and structured according to the main sections (3, 4, and 5) of the paper, we discuss the most relevant findings and their implications for answering the last question posed above.

6.1. Branching process approximation for the time dependence of the mutant frequency distribution

The spread of a new mutation in a population as a function of time can be described by the transition matrix of an appropriate Wright-Fisher model or the transition density of the approximating diffusion. Although diffusion theory leads to quite simple expressions for many important quantities, explicit and analytically tractable time-dependent results, tracing for instance the distribution of allele frequencies, seem to be out of reach (see Steinrücken et al., 2013, for a semi-explicit approach).

We combined two methods to derive an explicit and accurate time-dependent approximation for the mutant's density in any generation n: a branching process approach capturing the stochastic effects and the deterministic logistic growth model. We developed this approach quite generally with the help of a slightly supercritical Galton-Watson process that allows for quite general offspring distributions.

The approximation $g_a(x)$ in (3.16) for the mutant frequency density depends only on the compound parameter a, which in turn depends on the generation number n, the mean $\sigma = e^s$ of the offspring distribution, the extinction probability $P_{\text{ext}}(n, \sigma)$ until generation n in the branching process, and the population size N. Its shape is displayed in Figure 3.3 and compared with Wright-Fisher simulations in Figures 3.4 and S.3. It is highly accurate in the initial phase, already when $Ns \geq 1$. If $Ns \geq 10$ then it is accurate until about $\frac{1}{s}\ln(2Ns)$, which is approximately one half of the expected fixation time in the Wright-Fisher model. If $Ns \geq 100$, then it remains accurate for even longer and even if x is relatively close to 1. Because, in contrast to previous related investigations (Desai and Fisher, 2007, Uecker and Hermisson, 2011, Martin and Lambert, 2015), we condition on non-extinction until generation n, our approximation is more accurate than those obtained there, especially in the initial phase (cf. Figure 3.2, which shows the density ψ_n in the branching process, from which g_a is obtained by a proper transformation). However, by relying on a branching process to capture the inherent stochasticity, our model underestimates the effect of genetic drift near fixation, in particular if Ns is not much larger than 1. Remarkably, the

approximations for the mean and variance of this distribution are accurate in a much wider parameter range. In addition, they clearly display the well known fact that, conditioned on non-extinction (in our case, until the time of observation), the mutant frequency grows faster than predicted by the corresponding deterministic model, particularly when selection is weak (see Fig. 4.1 and the discussion in Sect. 3).

Finally, these results are extended to the infinite sites model, in which the number of new (beneficial) mutants emerging in the population per generation is Poisson distributed with mean $\Theta = NU$, and every mutation occurs at a new site. Here, U is the beneficial mutation rate affecting the trait. Sites are assumed to be unlinked (and in linkage equilibrium). The monomorphic initial population started to evolve at $\tau = 0$. Equation (3.21) provides the probability density $h^{(i)}(x) = h^{(i)}_{\tau, \sigma_i, N, \Theta}(x)$ of the ith mutation (which occurred at the random time τ_i at the ith site), conditioned on non-extinction until time τ. Mutations of unequal effects are admitted, as signified by σ_i. The integration is necessary because the waiting time distribution for the ith mutation event (an Erlang distribution) needs to be taken into account. Its cousin, the unconditioned distribution $\tilde{X}^{(i)}_\tau$ with absolutely continuous part $\tilde{h}^{(i)}_\tau(x)$ is defined in (3.23) and (3.24). It forms the basis for the applications to the quantitative genetic model.

6.2. Evolution of the mean phenotype and the genetic variance

The above discussed approximations (3.16) and (3.24) for the time dependence of density $g_{a(\tau)}$ and of the distribution of $\tilde{X}^{(i)}_\tau$ allow us to obtain highly accurate approximations for the evolution of the expected mean ($\bar{G}$) and variance (V_G) of an additive trait subject to exponential directional selection in a finite population of size N. We assume that potentially infinitely many, stochastically independent, loci (sites) can contribute additively to the trait. This assumption of global linkage equilibrium will be a good approximation if sites are physically unlinked because multiplicative fitnesses, as caused by exponential directional selection, do not generate linkage disequilibrium (Bürger, 2000, p. 79).

Equations (4.9) and (4.10) provide explicit approximations for the time dependence of the expected phenotypic mean and variance, which are astonishingly accurate for the whole evolutionary process, i.e., from the initial phase until the stationary response has been achieved (Figure 4.2). These expressions require integration with respect to two parameters: one over the time span until the time τ of interest, the other with respect to the distribution $f(\alpha)$ of mutation effects. The former is necessary because the extinction probability is time dependent, at least in the initial phase. The latter integration is unavoidable unless all mutation effects are equal, when accurate approximations are involving only the computation of a finite sum given in Remark 4.4. We note that these approximations hold for very general offspring distributions, so that they can cover cases where the effective population size N_e differs from the actual population size N. The offspring

distribution enters these equations through the extinction probabilities P_{ext}.

The general equations (4.9) and (4.10) do not provide much direct insight into the evolution of

$\bar{G}$ and V_G, except that the population-wide mutation rate Θ enters exclusively as a multiplicative

factor. However, their numerical evaluation is straightforward and efficient, and they yield excellent

and simple analytical approximations for the initial phase and for the stationary phase, when

emergence and fixation of new mutations balance and the genetic variance has reached a stationary

value.

6.2.1. The stationary phase

Typically, the stationary phase is reached once time exceeds the mean fixation time $\bar{t}_{\text{fix}}$ (4.14)

expected for the given mutation distribution, selection intensity, and population size (Figure 4.2).

The most accurate expressions for the stationary mean $\bar{G}^*(\tau)$, which responds linearly in time,

and the constant variance V_G^* are (4.17) and (4.18). They invoke the, in general, unavoidable

integration over the distribution of mutation effects. For equal effects, the simple approximation

(4.19) for the stationary variance is obtained. The equations (4.17) and (4.18) reveal the following

general and fundamental relation between the change of the mean and the variance:

$$\Delta \bar{G}^* \approx s V_G^* + U \bar{\alpha} \tag{6.1}$$

(from equation (4.23) by recalling $\Theta = NU$). Of course, with $U\bar{\alpha} \approx 0$, this is a special case

of Lande's (1976) equation, which was derived from the assumption of a normal distribution of

breeding values.

For the per-generation change of the mean phenotype at stationarity, we obtain in (4.20a) the

approximation $\Delta \bar{G}^* \approx \Theta \int_0^\infty \alpha P_{\text{sur}}(e^{s\alpha}) f(\alpha) \, d\alpha$, where P_{sur} is the survival probability of a single

mutant in the branching process. For a Poisson offspring distribution and an exponential distribution

f of mutation effects, this simplifies to $\Delta \bar{G}^* \approx 4\Theta s \bar{\alpha}^2 (1 - \frac{5}{2} s \bar{\alpha})$, which is approximately twice

the response of the mean when mutation effects are equal (see Corollary 4.8). Simple explicit

expressions for the variance V_G^* follow readily via (6.1). For large population sizes, $N \geq 10^4$,

the simple approximations in Corollary 4.8 are highly accurate for selection coefficients s ranging

over more than three orders of magnitude (Figure 4.3). The relation (6.1) is violated when the

population size is so small that $Ns < 2$, as can be observed in Fig. 4.3 by noting the deviation

of the open symbols from the black dashed curves. The reason is that for such small values of

Ns random genetic drift becomes an important force and the fixation probability of very slightly

beneficial mutations is higher than predicted by the branching process approximation.

Formally, the approximation (4.20a) for the per-generation change of the mean phenotype

at stationarity is equivalent to Hill's (1982) approximation for the asymptotic response of a

quantitative trait coming from fixation of new mutations. In our terminology and for a haploid

population, his general expression [3] becomes $\Delta \bar{G} = \Theta(N_e/N) \int_{-\infty}^{\infty} \alpha P_{\text{fix}}(\alpha) f(\alpha) \, d\alpha$, where his mutation distribution is not necessarily confined to positive values. Here, $P_{\text{fix}}(\alpha)$ is the diffusion approximation of the fixation probability in the Wright-Fisher model and thus nearly identical to P_{sur} because we assume $Ns > 1$. Thus, we recover classical results, but from a different perspective. It seems somewhat surprising that the branching process approach, which is best suited to study the initial fate of a mutation subject to selection, yields highly accurate approximations for the long-term response of a quantitative trait. For a more general discussion of the long-term response to various forms of directional selection, see Bürger (2000, pp. 331–338).

6.2.2. The initial phase

The general equations (4.9) and (4.10) for the evolution of the phenotypic mean and variance yield simpler and accurate approximations for the initial phase, during which the genetic variance is building up. However, we need to distinguish two phases.

During the first, very short phase when $\tau \ll (\ln N)/(s\bar{\alpha})$, the utterly simple approximations in Proposition 4.13 are obtained. Thus, for mutation effects equal to α, the mean phenotype grows approximately exponentially, i.e., $\bar{G}(\tau) \approx \frac{\Theta}{Ns}(e^{s\alpha\tau} - 1) \approx \frac{\Theta}{N}\alpha\tau\left(1 + \frac{s\alpha\tau}{2!} + \frac{(s\alpha\tau)^2}{3!}\right)$. This yields a very accurate approximation if $s\alpha\tau < \frac{1}{2}\ln N$, for small s even up to $\ln N$ (Figure 4.4). During this phase, variance and mean are related through $V_G \approx \alpha\bar{G}$. For an exponential distribution of mutation effects, the increase of the mean and the variance is considerably faster, indeed we have $\bar{G}(\tau) \approx \frac{\Theta}{N}\bar{\alpha}\tau\left(1 + s\bar{\alpha}\tau + (s\bar{\alpha}\tau)^2\right)$ (Proposition 4.13). This yields an increase of the mean that is slightly faster than that for equal effects with twice the selection intensity, $2s$. The reason is that with an exponential distribution of effects, mutations of large effect have a much higher probability of becoming fixed than mutations of small effect. In fact, the average effect of mutations drawn from an exponential distribution with mean $\bar{\alpha}$, conditioned on eventual fixation, is approximately $2\bar{\alpha}$. For exponentially distributed mutation effects, the approximations in Proposition 4.13 for the mean and variance are accurate only during the very short phase when (approximately) $s\bar{\alpha}\tau < \frac{1}{8}\ln N$.

In Proposition 4.11, we derived approximations for $\bar{G}(\tau)$ and $V_G(\tau)$ that are accurate for a much longer time span than above, but perform less well during the first generations. They perform well when $1/\Theta + 18e^{-5s} < \tau < \tau_c = 1/\Theta + \bar{t}_{\text{fix}}$. The lower bound results from the approximate waiting time for the first mutation plus the time span until at least 90% of mutants destined to loss are lost, i.e., such that $P_{\text{ext}}(\tau) \geq 0.9 P_{\text{ext}}^{\infty}$ (Section 3.2). This lower bound allows to avoid the iterative computation of the (time dependent) extinction probabilities $P_{\text{ext}}(t)$ in the general equations (4.9) and (4.10), hence the integration over the time. Thus, for equal mutation effects, no integration at all is required to evaluate $\bar{G}(\tau)$ and $V_G(\tau)$, which are then given by (4.30) and (4.31), respectively. The upper bound τ_c is chosen because then the contribution of fixed mutations to the response of the mean is small.

 6.2.3. Comparison of response from standing variation and from new mutations

1017   We developed our model for an initially monomorphic population and adaptation from *de novo*

1018 mutations. It is of interest to compare the response from new mutations with that from standing

1019 variation. From the classical breeder's equation it is well known that the per-generation response

1020 of the mean phenotype to directional selection is approximately

$$\Delta \bar{P} = h^2 \Delta_s \bar{P} = i h^2 \sigma_P \,, \tag{6.2}$$

1021 where $h^2 = \sigma_A^2/\sigma_P^2$ is the (narrow sense) heritability of the trait (with σ_A^2 and σ_P^2 the additive

1022 genetic and the phenotypic variance, respectively), $\Delta_s \bar{P}$ is the selection differential, i.e., the

1023 difference in mean phenotype before and after selection, and $i = \Delta_s \bar{P}/\sigma_P$ is the so-called selection

1024 intensity, or the standardized selection differential (e.g., Bürger, 2000, pp. 317-318). Under

1025 exponential directional selection one obtains $i = s\sigma_P$ (Bürger, 2000, p. 320), so that (6.2) becomes

1026

$$\Delta \bar{P} = h^2 s \sigma_P^2 \,. \tag{6.3}$$

Because classical quantitative genetic models assume that environmental effects have a dis-

tribution with mean zero, we have $\Delta \bar{G} = \Delta \bar{P}$. Thus, equation (6.1) predicts essentially the

same response as (6.3) because the per-generation contribution of new mutations will typically be

negligible. We use equation (6.3) to approximate the response of the mean from standing genetic

variation. Heritabilities are often in the range 0.1 to 0.7 (e.g., Table 10.1 in Falconer and Mackay,

1996; Chap. 7 in Lynch and Walsh, 1998), for morphological traits more often in the lower half of

this range.

Stationary phase. Our equation (4.20) informs us that, once stationarity is achieved, the response

from new mutations is approximately

$$\Delta \bar{G}^* \approx c\Theta s \alpha_2(f) = c\Theta s \bar{\alpha}^2 (1 + \mathrm{CV}_\alpha^2) \,, \tag{6.4}$$

where we have used $P_{\mathrm{sur}}(e^{s\alpha}) \approx cs\alpha$ (recall that $c = 2$ if the offspring distribution is Poisson,

whence $N_e \approx N$). Here, CV_α^2 is the squared coefficient of variation of the distribution of mutation

effects. For equal effects $\mathrm{CV}_\alpha^2 = 0$, for exponentially distributed effects $\mathrm{CV}_\alpha^2 = 1$, for Gamma-

distributed effects with density proportional to $\alpha^{\beta-1}e^{-\alpha}$, $\mathrm{CV}_\alpha^2 = 1/\beta$, and for effects drawn from

the truncated Gaussian in Remark 4.10, $\mathrm{CV}_\alpha^2 = \frac{1}{2}\pi - 1 \approx 0.507$. As already noted above, an

equation equivalent to (6.4) was derived and discussed by Hill (1982a,b). By comparison of (6.3)

with (6.4), we infer that for a Poisson offspring distribution and for exponentially distributed

mutation effects, the response due to new mutations exceeds that from standing variation if

$$4\Theta \bar{\alpha}^2 > h^2 \sigma_P^2 \,. \tag{6.5}$$

Inferences from various empirical estimates suggest that $\bar{\alpha}$ may often be not only of the same order of magnitude as σ_P, but actually quite similar to it (Walsh and Lynch, 2018, p. 1019). In fact, they suggest $\sigma_\alpha^2 \approx \sigma_E^2$, where σ_α^2 is the variance of a symmetric distribution of mutation effects with mean zero and σ_E^2 is the environmental variance. Under our assumption of exponentially distributed (positive) effects, we obtain $\sigma_\alpha^2 = 2\bar{\alpha}^2$. Thus, (6.5) becomes $2\Theta > h^2$ if we assume $\sigma_P^2 \approx \sigma_E^2 \approx \sigma_\alpha^2 = 2\bar{\alpha}^2$. Therefore, if $h^2 \approx 0.2$, a population-wide positive mutation rate of $\Theta \geq 0.1$ will be required for the response from new mutations to exceed that from standing variation. This reasoning for the stationary phase should apply after at most $\bar{t}_{\text{fix}} \approx \frac{2}{s\bar{\alpha}} \ln(2Ns\bar{\alpha})$ generations after the onset of directional selection (see eq. (B.15) and Fig. 4.2).

There is a caveat in the comparison in (6.5) because in the absence of new mutations the standing variation in the base population (before directional selection started) will slowly decrease and eventually be exhausted, i.e., h^2 will decrease to zero. The expected 'half-life', however, will be up to $1.4N_e$ generations (Robertson, 1960), which may considerably exceed the time needed to approach the stationary phase of response under new mutations (see above and recall that we assume $N \gg 1/(s\bar{\alpha})$). If the standing variation in the base population was due to mutation-selection-drift balance, and mutations continue to occur, then h^2 (and, of course, σ_P^2) will change very slowly, especially if the infinitesimal model applies (e.g., Barton et al., 2017, and references therein).

Equation (6.4) demonstrates that the response of the mean depends not only on the selection intensity, the mutation rate and the average effect of beneficial mutations, but also on the shape of the mutation distribution, in particular its coefficient of variation. Therefore, distributions with a large coefficient of variation of beneficial effects, such as Gamma-distributions with small β, lead to a faster response. This effect is already clearly visible by comparing panels A and C in Fig. 4.2, where in A we have $\text{CV}_\alpha^2 = 0$ and in B, $\text{CV}_\alpha^2 = 1$; see also panel A in Fig. S.4, where $\text{CV}_\alpha^2 = \frac{1}{2}\pi - 1$. Concerning the comparison of the response from standing variation to that from new mutations, (6.5) shows that the selection intensity plays no role in this regard, at least to this order of approximation.

Initial phase. A comparison of the response of the mean phenotype from standing variation and new mutations during the initial phase of adaptation is more delicate. Whereas for the response from standing variation (6.3) can still be used, the response from new mutations increases nonlinearly (see Fig. 4.4). In particular, equation (4.28) as well as its simplification (4.30) for equal effects, are too complicated to derive analytical comparisons between the two scenarios. Although numerical solution is simple and computationally fast, there are many scenarios to be compared with of how standing variation is maintained and how it is structured (e.g., De Vladar and Barton, 2014, Stephan and John, 2020, Hayward and Sella, 2022, and references therein). Thus, a detailed

 comparison has to be postponed to a future treatment.

 6.3. Number of segregating sites as an indicator for sweep vs. shift patterns

 Taking up the arguments in Sect. 5, we propose to use the number of segregating sites as a
 main indicator for the distinction between the scenarios of adaptation being mainly due to selective
 sweeps or mainly due to small frequency shifts at many loci. This is supported by our theory in
 Sect. 5 showing that once a stationary response to directional selection has been achieved, the
 expected number of segregating sites, $\bar{S}^*$, is proportional to Θ, depends logarithmically on N,
 exceeds the neutral expectation of $2\Theta \ln N$, but never by more than a factor of two if s is varied
 between 0 and 0.5 (see eq. (5.2) and Fig. 5.1). As a function of s, the maximum value of $\bar{S}^*$ is
 achieved if $\bar{\alpha} s \approx 1/(2 \ln N)$.

 For the initial phase of adaptation, the branching process approximations $\tilde{S}(\tau)$ in (5.5) and (5.6)
 are much more accurate than the simple approximations $\bar{S}(\tau)$ in (5.3) (see Fig. D.1). However,
 all these approximations show that the expected number of segregating sites is proportional to Θ,
 depends only very weakly on the product of s and the average mutation effect $\bar{\alpha}$, and is independent
 of the population size N.

 Further support for our proposition derives from Figure 6.1. Motivated by Höllinger et al.
 (2019), we explored the polygenic pattern of adaptation at the loci contributing to the trait at the
 generation T when the mean phenotype $\bar{G}$ in a population first reaches or exceeds a given value,
 which for this figure we chose to be 1 (in units of mutation effects α). For various parameter
 combinations and for equal and exponentially distributed mutation effects, Figure 6.1 displays the
 frequency distributions of the first four *successful* mutants at this generation T. By successful
 mutant we mean that at the stopping time T, we condition on the sites at which mutants are
 present (segregating or fixed), and we number them from 1 to 4, depending on the order in
 which these mutations occurred. Thus, the mutation at site 1 is the oldest in the population.
 This is a small subset of sites at which mutations have occurred because their majority has been
 lost. The histograms are obtained from Wright-Fisher simulations. We computed the average
 stopping time, $\bar{T}$, the average number of segregating sites at time T, $\bar{S}_T$, and the average of mean
 phenotypic values in generation T, $\bar{G}_T$. These values are given in each panel (with subscripts
 omitted for visibility reasons). Because time is discrete, $\bar{G}_T \geq 1$. For large selection coefficients
 (and large mutation effects) the per-generation response tends to be large once several mutations
 are segregating. Therefore, $\bar{G}_T$ can be noticeably larger than 1. Indeed, the distribution of $\bar{G}$
 (under the stopping condition $\bar{G} = 1$) has a sharp lower bound at 1 and may have substantial skew
 to the right, especially for exponentially distributed effects (results not shown).

 Figure 6.1 shows clear sweep patterns in the panels with $2\Theta = 0.1$ (where $\bar{S}_T < 2.5$) and
 distinctive small-shift patterns at many sites if $2\Theta = 10$ (where $\bar{S}_T > 35$). The patterns occurring

Figure 6.1: Histograms of the frequency x of the first four successful mutants at the first generation T in which the mean phenotype $\bar{G}$ in a single population reaches or exceeds the value 1. The averages of the stopping time T, of the number of segregating sites at T, and of the mean phenotypic values at T, denoted by $\bar{T}$, $\bar{S}(=\bar{S}_T)$, and $\bar{G}(=\bar{G}_T)$, are given in each panel. Histograms and averages are based on $20\,000$ replicate runs of the Wright-Fisher model. The dark blue color is for the first successful mutation, light blue for the second, red for the third, and orange for the fourth successful mutation. The selection coefficients s and the mutation effects α are indicated for each row. Columns show results for different values of Θ. The population size is $N = 10^4$ in all cases. One unit on the vertical axis corresponds to 200 runs. The expected times to fixation for rows 1, 2, and 3 are $t_{\text{fix}} \approx 7\,039$, $t_{\text{fix}} \approx 156$, and $\bar{t}_{\text{fix}} \approx 153$, respectively. The corresponding scaled stationary values of the number of segregating sites, $S^*/(2\Theta)$, are approximately 13.7, 16.5, and 16.0 (calculated from eq. (5.2)). Thus, in the first column S has reached its stationary value in rows 1 and 2, and 60% of it in row 3. The distributions of the stopping times T corresponding to the above panels are shown in Fig. S.6.

for $2\Theta = 1$ suggest a series of partially overlapping sweeps. The main effect of varying the selection strength s is on the time T to reach (or exceed) $\bar{G} = 1$. Successive sweeps are indeed expected if $2\Theta = 0.1$ because the average waiting time between successful mutation events is approximately $1/(2s\Theta)$. If $s = 0.001$, this waiting time is $10\,000$ and exceeds the expected fixation time of roughly $7\,000$ generations. If $s = 0.1$, the waiting time is 100 generations and falls short of the expected fixation time, which is about 156 generations (from (B.15)). In addition, we observe that for small Θ the first successful mutant corresponds mostly to the site with the highest mutant frequency at the time of stopping, whereas for large Θ the distributions of the first four successful mutations

are hardly distinguishable.

In the panels with $2\Theta = 0.1$, the expected stopping times $\bar{T}$ exceed their corresponding expected fixation times (given in the caption of Fig. 6.1) considerably. Hence, most of these populations have already reached the stationary phase. Somewhat unexpectedly, the observed average numbers of segregating sites (from top to bottom: 2.2, 2.1, 1.9) exceed their respective stationary values $\bar{S}^*$ (1.37, 1.65, 1.60) considerably. This observation seems to be at variance with the results in panels B and D of Figure 5.1, which show that $\bar{S}^*$ is a very accurate approximation of the numerically determined $\mathrm{E}[S]$. The reason for this discrepancy is the different stopping condition, more precisely the fact that the distribution of stopping times T is very wide and strongly skewed to the right if Θ is small (Fig. S.6). Slowly evolving populations may carry many more mutations once they reach $\bar{G} = 1$ than the average or fast evolving populations because they have to wait long for the first successful mutant(s). As a result, the stopping condition $\bar{G} = 1$ inflates the number of segregating sites compared to a process that is stopped at a given generation.

For $2\Theta \geq 1$ our branching process approximations (5.5) and (5.6) predict the observed average $\bar{S}_T$ very well if $\tilde{S}$ is evaluated at $\bar{T}$. Indeed, very accurate approximations of the observed $\bar{T}$ are obtained for equal effects by solving the equation $\bar{G}(t) = 1$ using (4.30). The reason is that the distribution of T becomes increasingly symmetric and less variable as Θ increases. It is nearly perfectly symmetric if $2\Theta = 10$ (Fig. S.6). For exponentially distributed effects, when the stopping time distribution is more skewed, an accurate approximation of $\bar{T}$ is obtained from (4.28) by solving $\bar{G}(t) = 1$ only if $2\Theta \geq 10$ (results not shown). Accordingly, for such large values of Θ, the Wright-Fisher simulations are stopped at the previously determined value $\bar{T}$ yield distributions very similar to those obtained by stopping when $\bar{G} = 1$ has been reached (results no shown).

Figure 6.1 confirms the above discussed finding that for an exponential distribution of mutation effects, the response of the mean is faster than for equal effects also under the current stopping condition. As a consequence, the number of segregating sites when $\bar{G} = 1$ is reached is smaller, and considerably so if $\bar{G} = 1$ is reached well before the stationary phase is entered, as is the case of $2\Theta \geq 1$. Unsurprisingly, the distribution of stopping times T is much wider than for equal effects (Fig. S.6) because mutations of larger (beneficial) effect have a higher fixation probability and lead to faster sweeps, especially for strong selection. Mutation distributions with a heavier tail than the exponential may intensify this effect.

If a small sudden shift in the optimum phenotype occurs, mutations of relatively small effect may be most conducive to selective sweeps because then large-effect mutations become deleterious when the population closely approaches the optimum. If a sudden shift in the optimum of appreciable magnitude occurs and the initial population has been in mutation-stabilizing selection balance before, then a kind of threshold behavior has been identified for deterministic models (De Vladar

and Barton, 2014, Jain and Stephan, 2017a). Under mutation-selection balance the initial variance will be high if allelic effects are small because such alleles are maintained at intermediate frequencies. By contrast, the initial variance will be low if allelic effects are very large because such alleles will be nearly fixed or absent. Alleles of very small effect will respond slowly after the onset of selection because they are nearly neutral. If their number is large, as assumed by the infinite sites model, then the total response of the mean phenotype can still be large (proportional to the variance, as in Lande's equation discussed above). Alleles of very large effect will initially also respond very slowly because they are so rare, and they will lead to a slow response of the mean phenotype because there is little variation on which selection can act. As a result, alleles with effects close to an intermediate threshold value will be the first to respond quickly and start sweeping. On a longer time scale and for a large shift in the optimum, the response from a few loci with large effects is comparable with that from many loci with small effects if the sum of effect sizes is the same. If most loci have large effects, then the response will clearly be much faster than the response from the same number of small-effect loci; the larger the effect the faster the sweeps (De Vladar and Barton, 2014, Jain and Stephan, 2017a,b).

In finite populations the conclusions are even more subtle and may deviate from those in the deterministic case. Mutations of small effect may contribute much less to the response than in large populations because their evolutionary fate is dominated by genetic drift and their initial distribution under mutation-stabilizing selection-drift balance will be U-shaped. Again, mutations of moderate effects may contribute most to the selective response after a sudden shift in the optimum (John and Stephan, 2020, Hayward and Sella, 2022). The latter authors also conclude that mutations of large effect are unlikely to sweep, but this is a consequence of their assumptions that the trait is highly polygenic (i.e., $\sqrt{\Theta} \gg 1$), the population is initially in mutation-stabilizing selection-drift balance (whence alleles of large effect are extremely rare), and there is only a small shift in the optimum of a few phenotypic standard deviations. The latter assumption implies that the equilibration phase is reached quickly, and then fine-tuning by alleles of small effect is essential for adaptation. If alleles of large effect contribute substantially to the initial variance and the shift is large, then they will sweep (Chevin and Hospital, 2008, Thornton, 2019, John and Stephan, 2020, Stephan and John, 2020).

For a gradually moving optimum, after a relatively short initial phase, a quasi-stationary phase is entered during which, except for stochastic fluctuations, a constant average variance is attained and the mean phenotype lags behind the optimum by an, on average, constant amount. Of course, a prolonged quasi-stationary response will depend crucially on a constant input of new mutations. During this phase, the trait experiences a mixture of directional and stabilizing selection, and the variance settles to a value that is lower than the variance under mutation-drift balance

(which is nearly the same as that achieved under exponential selection), but higher than that under mutation-stabilizing selection-drift balance (Lynch and Lande, 1993, Bürger and Lynch, 1995, Jones et al., 2004). Matuszewski et al. (2015) showed that compared to adaptation from *de novo* mutations, adaptation from standing variation proceeds by the fixation of more alleles of small effect. If the optimum is stopped after some time or new mutations are ignored, then the phenotypic dynamics becomes more complex and may depend on details of the underlying genetics (De Vladar and Barton, 2014).

In our finite-population model, in which initial mutant frequencies are $1/N$, the response of the mean is faster the larger the allelic effects. In addition, mutations that sweep, sweep faster than predicted by deterministic growth. Nevertheless, the response of the mean phenotype depends linearly on N through $\Theta = NU$ (Corollary 4.8 and Proposition 4.11), except in the very early phase when it depends mainly on $U\bar{\alpha}$ but is nearly independent of N (Proposition 4.13).

6.4. Conclusions, limitations, and outlook

We conclude from our analyses that Θ, the expected number of beneficial mutations occurring in the population per generation (and contributing to the trait) is the main determinant of the pattern of adaptation at the genetic level. Selective sweeps at a few loci are the dominant pattern if $\Theta \leq 0.1$, and small allele frequency shifts at many loci are observed if $\Theta \geq 10$. Given the same mean (beneficial) mutation effect, $\bar{\alpha}$, higher second and third moments of the mutation distribution favor sweeps. The expected number of segregating sites is an excellent indicator for the pattern of adaptation. However, it may be challenging to estimate the number of segregating sites contributing to a trait. The selection coefficient s and the average effect $\bar{\alpha}$ primarily determine the rate of adaptation via their product $s\bar{\alpha}$. Except for Θ, the population size has a weak effect on the rate of adaptation and on the population variance; often it enters expressions approximately logarithmically. We emphasize that our analysis is based on the assumption $Ns\bar{\alpha} \gg 1$, i.e., that selection is considerably stronger than random genetic drift. Most of our approximations, however, perform well when $Ns\bar{\alpha} > 2$.

Our conclusions concerning the influence of the parameters on the pattern of adaptation are similar to those of Höllinger et al. (2019), although their model and approach deviate from ours in important respects. Most significantly, their genetic architecture differs drastically from ours. They study a polygenic trait that assumes only two or very few values, such as resistance and non-resistance. The contributing loci thus are highly redundant, and a mutation at one or at very few loci is sufficient to complete adaptation. This is a very strong form of epistasis. In addition, mutation effects are the same at all loci. Selection on the discrete trait acts similarly as in our model: they assume a linear fitness function in continuous time, which corresponds to an exponential in discrete time. In contrast to our model, their initial populations are polymorphic

and (mostly) assumed to be in mutation-stabilizing selection-drift balance. Nevertheless, they find that their population-scaled background mutation rate Θ_{bg}, which corresponds closely to our 2Θ, explains the main differences in the patterns of adaptation, i.e., few sweeps versus many slight shifts. In the thesis by Höllinger (2018, Chap. 4), this main conclusion is confirmed for an additive trait subject to a sudden shift in the optimum phenotype, again for equal mutation effects.

The validity of our conclusions may be limited by a number of important simplifying assumptions, which are unlinked loci and linkage equilibrium, no pleiotropy or epistasis, and exponential directional selection. In addition, we assume that selection is stronger than random drift and we study only the selection response from new mutations. Our results are derived for haploid populations, but will apply as well to diploids without dominance by substituting $2N$ for N.

Our assumption of unlinked sites will be biologically reasonable unless the number of (simultaneously) segregating sites exceeds the number of chromosomes by far or many sites occur in regions of reduced recombination. Although exponential directional selection by itself does not induce linkage disequilibrium, deviations from this form of selection as well as random genetic drift will do so. Therefore, it would desirable to extend the above results to loosely linked loci and perform a kind of quasi-linkage equilibrium analysis.

Quite extensive simulation results for models of artificial directional selection (such as truncation selection) by Keightley and Hill (1983, 1987) suggest that linkage reduces the asymptotic variance, hence the response of the mean, compared to unlinked loci. The reduction will be small unless linkage is very tight. A similar observation was reported in Bürger (1993) for exponential directional selection. However, these results included only population sizes of a few hundred individuals. For a moving optimum model and population sizes up to about 10^4, a qualitatively similar finding was obtained in (Bürger, 2000, pp. 335–338). In all these investigations, mutation distributions with positive and negative effects were used. Then, mutations of both signs appear on the same chromosome, whence low recombination will reduce fixation probabilities, especially of weakly beneficial mutants, and hamper adaptation (Barton, 1995). In concordance with these results, a simulation study by Thornton (2019) found that weak or moderate linkage has almost no effect on the response of the mean and on the variance of the trait, and also little effect on the fixation probabilities of advantageous mutations, but the resulting interference increases fixation times and affects haplotype diversity. Analytical work on the effects of linkage is scarce, based on specific assumptions, and shows that linkage can increase or decrease the variance of a trait under directional selection, depending on the model assumptions and the magnitude of the parameters (e.g. Robertson, 1977, Hospital and Chevalet, 1996, Zhang and Hill, 2005b). Determining the effects of linkage on the likelihood of sweeps also poses a formidable challenge (e.g. Kim and Stephan, 2003). Thus, there is ample room for future research.

Moderate or strong linkage can lead to selective interference. An extreme scenario is provided by a single locus with recurrent mutation. For this case our theory remains applicable if $2N\mu \leq 1$, where μ is the per-locus mutation rate. Then it is sufficient to consider the first *successful* mutation event. If $2N\mu > 1$, competition of mutants from different mutation events can no longer be ignored and complicates the evolutionary dynamics (unpublished results by HG).

We have already pointed out that strong epistasis does not necessarily affect our main conclusions, but further research seems necessary. We expect that our assumption of exponential directional selection provides a good approximation for the earlier phases of adaptation under a wide variety of forms of directional selection, including truncation selection, a steadily moving optimum, and a sudden shift in the optimum phenotype. Some of the previously discussed results provide support for this expectation. Exponential directional selection will not be suitable as an approximation if selection is epistatic and strong, if it is stochastically or periodically fluctuating, or if there are pleiotropic fitness effects.

Our results concerning the time dependence of the distribution of sweeping *de novo* mutations will be applicable to very rare allelic variants already present in the population. This will be the case if their effect is large and the population has been under stabilizing selection before directional selection started. If more than one copy is initially present, the quasi-deterministic phase could be started from a gamma-distributed effective initial population size (Martin and Lambert, 2015) instead of an exponential. In each case, this will speed up the selection response.

Acknowledgments

We thank Joachim Hermisson, Emmanuel Schertzer, Ben Wölfl, and Ilse Höllinger for helpful discussions. Financial support by the Austrian Science Fund (FWF) through the Vienna Graduate School of Population Genetics (GrantDK W1225-B20) to HG is gratefully acknowledged.

Additional materials

We provide a comprehensive Wolfram *Mathematica* notebook, showing additional visualizations of the analytical predictions and efficient numerical evaluations of those. Upon publication, the notebook and the simulation code (C++) will be made available via github.

Appendices

A. Variance and skew of W^+

Here we show that, in general, the distribution of W^+ in Lemma 3.1 is not exponential. To this aim, we derive the variance and the skewness of W^+. Indeed, the method can be generalized

to derive higher-order cumulants of W^+. In the following we denote the nth cumulant of a random variable X by $\kappa_n[X]$. Hence, κ_2 is the variance and κ_3 the third central moment. For random variables X and Y that posses cumulants up to the given order, the following relations hold:

$$\mathrm{E}[X] = \mathrm{E}[\mathrm{E}[X|Y]]\,, \tag{A.1a}$$

$$\kappa_2[X] = \mathrm{E}[\kappa_2[X|Y]] + \kappa_2[\mathrm{E}[X|Y]]\,, \tag{A.1b}$$

$$\kappa_3[X] = \mathrm{E}[\kappa_3[X|Y]] + \kappa_3[\mathrm{E}[X|Y]] + 3\,\mathrm{Cov}[\mathrm{E}[X|Y], \kappa_2[X|Y]]\,. \tag{A.1c}$$

Here, the first two equations state the well known laws of total expectation and of total variance. The third is a special case of the so-called law of total cumulance, which holds for arbitrary multivariate cumulants (Brillinger, 1969).

Let $X = \sum_{i=1}^{Y} \xi_i$, where Y is a random variable with values in the positive integers and the $\{\xi_i\}$ are i.i.d. random variables with mean $\sigma = \mathrm{E}[\xi]$ and higher-order cumulants $c_n = \kappa_n[\xi]$. Then $\mathrm{E}[X|Y] = \sigma Y$ and $\kappa_n[X|Y] = c_n Y$, and we obtain

$$\mathrm{E}[X] = \sigma\,\mathrm{E}[Y]\,, \tag{A.2a}$$

$$\kappa_2[X] = \sigma^2 \kappa_2[Y] + c_2\,\mathrm{E}[Y]\,, \tag{A.2b}$$

$$\kappa_3[X] = \sigma^3 \kappa_3[Y] + 3\sigma c_2 \kappa_2[Y] + c_3\,\mathrm{E}[Y]\,. \tag{A.2c}$$

Now we apply these relations to the random variables Z_n in Lemma 3.1 (where $Z_0 = 1$) to obtain after brief calculations the following recursion equations for the second and third cumulant:

$$\kappa_2[Z_n] = \sigma^2 \kappa_2[Z_{n-1}] + c_2 \sigma^{n-1}\,, \tag{A.3a}$$

$$\kappa_3[Z_n] = \sigma^3 \kappa_3[Z_{n-1}] + 3\sigma c_2 \kappa_2[Z_{n-1}] + c_3 \sigma^{n-1}\,, \tag{A.3b}$$

where we have used $\mathrm{E}[Z_{n-1}] = \sigma^{n-1}$. These recursion relations can be solved successively and yield

$$\kappa_2[Z_n] = c_2 \frac{\sigma^{n-1}(\sigma^n - 1)}{\sigma - 1}\,, \tag{A.4a}$$

$$\kappa_3[Z_n] = \frac{\sigma^{n-1}(\sigma^n - 1)\left[3c_2^2(\sigma^{n-1} - 1) + c_3(\sigma - 1)(\sigma^n + 1)\right]}{(\sigma - 1)^2(\sigma + 1)}\,. \tag{A.4b}$$

A slightly more general expression for $\kappa_2[Z_n]$ (with arbitrary $\mathrm{E}[Z_0]$ and $\mathrm{Var}[Z_0]$) may be found in Haccou et al. (2005, p. 20). Now we apply them to $W_n = Z_n/\sigma^n$ and take the limit $n \to \infty$. This yields the second and third cumulant of the distribution W:

$$\kappa_2[W] = \frac{c_2}{\sigma(\sigma - 1)}\,, \tag{A.5a}$$

$$\kappa_3[W] = \frac{c_3}{\sigma(\sigma-1)(\sigma+1)} + \frac{3c_2^2}{\sigma^2(\sigma-1)^2(\sigma+1)}. \tag{A.5b}$$

We recall the relation (3.2) between W and W^+ which, in a less formal way, can be written as $W = \delta_0 P_{\text{ext}}^\infty + (1 - P_{\text{ext}}^\infty)W^+$. Somewhat tedious calculations lead to the following relations between the cumulants of W^+ and W:

$$\kappa_2[W^+] = \mathrm{E}[W^+](\kappa_2[W] + 1 - \mathrm{E}[W^+]), \tag{A.6a}$$

$$\kappa_3[W^+] = \mathrm{E}[W^+](\kappa_3[W] - 3\kappa_2[W](\mathrm{E}[W^+] - 1) + 2\,\mathrm{E}[W^+]^2 - 3\,\mathrm{E}[W^+] + 1). \tag{A.6b}$$

Explicit, but not very simple, expressions in terms of the properties of the offspring distribution are obtained by substituting $\mathrm{E}[W^+] = (1 - P_{\text{ext}}^\infty)^{-1}$ and (A.5) into these equations.

For a geometric offspring distribution with mean σ, so that $P_{\text{ext}}^\infty = 1/\sigma$, we obtain $\mathrm{E}[W^+] = \frac{\sigma}{\sigma-1}$ and

$$\kappa_2[W^+] = \frac{\sigma^2}{(\sigma-1)^2}, \tag{A.7a}$$

$$\kappa_3[W^+] = \frac{2\sigma^3}{(\sigma-1)^3}, \tag{A.7b}$$

as it should be for an exponential distribution of W^+.

For a Poisson offspring distribution with mean σ (and $c_2 = c_3 = \sigma$), we obtain

$$\kappa_2[W^+] = \frac{1 - \sigma P_{\text{ext}}^\infty}{(\sigma-1)(1 - P_{\text{ext}}^\infty)^2}, \tag{A.8a}$$

$$\kappa_3[W^+] = \frac{2 + \sigma[1 - (3 + 4\sigma - \sigma^2)P_{\text{ext}}^\infty + \sigma(\sigma+2)(P_{\text{ext}}^\infty)^2]}{(\sigma-1)^2(\sigma+1)(1 - P_{\text{ext}}^\infty)^3}, \tag{A.8b}$$

where $P_{\text{ext}}^\infty = P_{\text{ext}}^\infty(\sigma)$ is given by (3.10). It is not difficult to show that $\kappa_2[W^+] < \mathrm{E}[W^+]^2$ holds for all $\sigma > 1$ and $\kappa_2[W^+]/\mathrm{E}[W^+]^2 \to 1$ as $\sigma \to 1$ (from above). Series expansion yields

$$\kappa_2[W^+]/\mathrm{E}[W^+]^2 = 1 - \frac{2}{3}(\sigma-1) + \frac{4}{9}(\sigma-1)^2 + O((\sigma-1)^3). \tag{A.9}$$

Therefore, W^+ is not exponentially distributed if the offspring distribution is Poisson.

The skew of W^+ has the series expansion

$$\mathrm{Skew}[W^+] = \kappa_3[W^+]/\kappa_2[W^+]^{3/2} = 2 - (\sigma-1) + \frac{1}{2}(\sigma-1)^2 + O((\sigma-1)^3), \tag{A.10}$$

which is less than two if $\sigma > 1$. In fact, it can be shown that it is even less than the skew of a Gamma distribution with the same mean and variance. Such a Gamma distribution (proportional to $x^{\alpha-1}e^{-\beta x}$) has the shape parameter $\alpha = \frac{\sigma-1}{1 - \sigma P_{\text{ext}}^\infty}$ and the rate parameter $\beta = \frac{(\sigma-1)(1 - P_{\text{ext}}^\infty)}{1 - \sigma P_{\text{ext}}^\infty} > 1 - P_{\text{ext}}^\infty$. Therefore, the true density of W^+ decays (slightly) faster than the exponential derived

in Lemma 3.1.

Finally, we give a numerical example for a Poisson offspring distribution with mean $\sigma = 1.1$. Then $1 - P_{\text{ext}}^{\infty} \approx 0.176$, $\mathrm{E}[W^+] \approx 5.68$, $\kappa_2[W^+] \approx 30.22$, $\kappa_3[W^+] \approx 316.4$, $\kappa_2[W^+]/\mathrm{E}[W^+]^2 \approx 0.937$, $\mathrm{Skew}[W^+] \approx 1.905$. Thus, according to these moments, the distribution is still quite close to an exponential distribution. The Gamma distribution with mean $\mathrm{E}[W^+]$ and variance $\kappa_2[W^+]$ has $\beta \approx 0.188$ and a skewness of $2/\sqrt{\alpha} \approx 1.936$.

B. Mean times to fixation or loss of a mutant

Approximations for the expected times to absorption conditioned on either fixation or loss of a new mutation, for simplicity called mean fixation time or mean loss time, have been an important tool in theoretical population genetics ever since its inception in the 1920s. Even in large populations, the initial fate of a mutant, as well as its fate in the final stages before fixation, is subject to strong stochastic influences, and the derivation of simple and accurate approximations is still a topic of research (e.g., Charlesworth, 2020, and references therein). In this appendix, we add simple estimates to this literature that are particularly useful to compute the mean fixation or loss times of beneficial mutants drawn from a distribution of effects.

For the computation of mean times to fixation or loss in a finite population, we use diffusion approximations because the branching process model is not directly applicable. Although the approximation $g_a(p)$ in (3.16) for the distribution of the number of mutations is quite accurate for late stopping times, the derivation of useful approximations for the mean fixation and loss times requires more delicate transformations than that in the proof of Theorem 3.6 and is work in progress.

B.1. Diffusion approximations

Here, we collect results on diffusion approximations of the fixation probability and of the mean times to fixation or loss of favorable mutants with a fixed selection coefficient $s > 0$. In a haploid population, the well-known diffusion approximation for the fixation probability of a single beneficial mutant is

$$P_{\text{fix}}(s, N) = \frac{1 - e^{-2s}}{1 - e^{-2Ns}}. \tag{B.1}$$

In the applications to our model, we replace s by $\tilde{s}(\alpha)$ defined in (4.13). The diffusion approximation $P_{\text{fix}}(s, N)$ is always an upper bound to the true fixation probability in the Wright-Fisher model, and is highly accurate for every $N \geq 8$ if $0 < s \leq 0.1$ (Bürger and Ewens, 1995).

The mean times to fixation or loss can be expressed only in terms of integrals of the corresponding sojourn time densities (Ewens, 2004, Sects. 4.6 and 5.3). For the mean fixation time, the simple approximation

$$t_{\text{fix}}^{(\text{HP})}(s, N) = \frac{2}{s} \left(\ln(2Ns) + \gamma - \frac{1}{2Ns} \right) \tag{B.2}$$

was provided by Hermisson and Pennings (2005), where γ is the Euler gamma. It is very accurate if $2Ns \geq 3$.

If $2Ns < 3$, we use

$$t_{\text{fix}}^{(\text{small})}(s, N) = 2N \left(1 - 2Ns \frac{11 - 6\gamma - 6\ln 3}{27} \right) . \tag{B.3}$$

This is the linear function in s which connects the mean fixation time $2N$ at $s = 0$ with the value of $t_{\text{fix}}^{(\text{HP})}(s, N)$ at $s = 3/(2N)$. As a function of s, $t_{\text{fix}}^{(\text{HP})}$ assumes its maximum very close to this value and becomes negative for very small s. As illustrated by Fig. B.1, the function

$$t_{\text{fix}}(s, N) \approx \begin{cases} t_{\text{fix}}^{(\text{HP})}(s, N) & \text{if } s \geq 3/(2N) , \\ t_{\text{fix}}^{(\text{small})}(s, N) & \text{if } s < 3/(2N) , \end{cases} \tag{B.4}$$

obtained by concatenation of $t_{\text{fix}}^{(\text{HP})}$ and $t_{\text{fix}}^{(\text{small})}$ yields a highly accurate approximation of the true diffusion approximation.

For the mean loss time of a single advantageous mutant, we obtain (see below)

$$t_{\text{loss}}^{(\text{app})}(s, N) \approx -2(1 + s)\ln(2s) + 2(1 - \gamma) + s(3 - 2\gamma) + \frac{1}{Ns} \tag{B.5a}$$

$$\approx -2\ln(2s) + 2(1 - \gamma) + \frac{1}{Ns} , \tag{B.5b}$$

which is very accurate if $Ns \geq 2$ but breaks down if $Ns \leq 1$.

If $Ns < 2$, we use

$$t_{\text{loss}}^{(\text{small})}(s, N) = 2\ln(N) - sN \left(\ln 4 + \gamma - \tfrac{5}{4} \right) . \tag{B.6}$$

This is the linear function in s which connects the (approximate) mean loss time $2\ln(N)$ at $s = 0$ (Kimura and Ohta, 1969) with the value of $t_{\text{loss}}^{(\text{app})}(s, N)$ at $s = 2/N$. As illustrated by Fig. B.1, the function

$$t_{\text{loss}}(s, N) \approx \begin{cases} t_{\text{loss}}^{(\text{app})}(s, N) & \text{if } s \geq 2/N , \\ t_{\text{loss}}^{(\text{small})}(s, N) & \text{if } s < 2/N \end{cases} \tag{B.7}$$

yields a highly accurate approximation of the true diffusion approximation.

Proof of (B.5a). To derive this approximation, we start with the well-known diffusion approximations for the conditional sojourn time densities (eqs. (5.54) and (5.55) in Sect. 5.4 of Ewens, 2004). To account for haploidy, we set $\alpha = 2Ns$ and $p = 1/N$ in these expressions (p the initial mutant frequency). By integration with respect to x, we obtain with the help of *Mathematica* and after simple algebraic rearrangement

$$t_{\text{loss}}^{(1)}(s, N) = \frac{1}{(e^{2Ns} - 1)s} \left[e^{2Ns} \left(-\operatorname{Ei}(-2s) + \gamma + \ln(2Ns) - \ln(N - 1) \right) - \operatorname{Ei}(2Ns) + \operatorname{Ei}(2(N - 1)s) \right.$$

$$-\operatorname{Ei}(2s) + \gamma + \ln(2Ns) - \ln(N-1) + e^{2Ns}\big(\operatorname{Ei}(-2(N-1)s) - \operatorname{Ei}(-2Ns)\big)\bigg]$$

(B.8)

for the expected time the mutant frequency spends in $(0, 1/N]$, and

$$t_{\text{loss}}^{(2)}(s, N) = \frac{e^{2s} - 1}{(e^{2Ns} - 1)(e^{2Ns} - e^{2s})s}\bigg[-e^{4Ns}\operatorname{Ei}(-2s) + e^{2Ns}\operatorname{Ei}(2(N-1)s)$$
$$+ e^{4Ns}\operatorname{Ei}(-2Ns) + \operatorname{Ei}(2Ns) + 2e^{2Ns}\big(\gamma + \ln(2Ns) - \ln(N-1)\big)$$
$$- \operatorname{Ei}(2s) + e^{2Ns}\operatorname{Ei}(-2(N-1)s)\bigg]$$

(B.9)

for the time spent in $(1/N, 1)$. We note that the times $t_{\text{loss}}^{(1)}(s, N)$ and $t_{\text{loss}}^{(2)}(s, N)$ have been rescaled to generations, i.e., the diffusion time has been multiplied by N. Omitting terms of order e^{-2Ns} or smaller in (B.8) and using the approximation $\operatorname{Ei}(x) \approx e^x/(x-1)$ if $|x| > 4$, we obtain

$$t_{\text{loss}}^{(1)}(s, N) = \frac{1}{(e^{2Ns} - 1)s}\bigg[e^{2Ns}\big(-\operatorname{Ei}(-2s) + \gamma + \ln(2Ns) - \ln(N-1)\big) - \operatorname{Ei}(2Ns) + \operatorname{Ei}(2(N-1)s)\bigg]$$
$$\approx \frac{1}{s}\bigg[-\operatorname{Ei}(-2s) + \gamma + \ln(2s) + \ln(N) - \ln(N-1) - \frac{1}{e^{2Ns}}\big(\operatorname{Ei}(2Ns) - \operatorname{Ei}(2(N-1)s)\big)\bigg]$$
$$\approx 2 - s + 1/N.$$

The latter approximation is obtained by performing a series expansion up to order s^2 and by keeping Ns constant (as in the diffusion approximation). For $t_{\text{loss}}^{(2)}$ we obtain in an analogous way, by omitting terms of order e^{-2Ns} or smaller,

$$t_{\text{loss}}^{(2)}(s, N) \approx \frac{e^{2s} - 1}{(e^{2Ns} - 1)(e^{2Ns} - e^{2s})s}\bigg[-e^{4Ns}\operatorname{Ei}(-2s) + e^{2Ns}\operatorname{Ei}(2(N-1)s)\bigg]$$
$$\approx \frac{e^{2s} - 1}{s}\bigg[-\operatorname{Ei}(-2s) + e^{-2Ns}\operatorname{Ei}(2(N-1)s)\bigg]$$
$$\approx -2\gamma - 2(1 + s)\ln(2s) + 2(2 - \gamma)s + \frac{1 - s}{Ns}.$$

By summing up, i.e., $t_{\text{loss}} = t_{\text{loss}}^{(1)} + t_{\text{loss}}^{(2)}$, and omitting terms of order $1/N$, we arrive at (B.5a). $\quad\square$

B.2. Mean fixation time averaged over the mutation distribution f

Now we derive an approximation for the mean fixation time of a single mutant if its effect α on the selected trait is drawn from an exponential distribution f with mean $\bar{\alpha}$. For given N and s, we defined in (4.14) the corresponding expected mean fixation time by

$$\bar{t}_{\text{fix}}(s, \bar{\alpha}, N) = \frac{\int_0^\infty t_{\text{fix}}(\tilde{s}(\alpha), N) P_{\text{fix}}(\tilde{s}(\alpha), N) f(\alpha)\, d\alpha}{\int_0^\infty P_{\text{fix}}(\tilde{s}(\alpha), N) f(\alpha)\, d\alpha}.$$

(B.10)

We recall from (4.13) that $\tilde{s}(\alpha) = e^{s\alpha} - 1$ is the selection coefficient of a mutant of effect α. For $P_{\text{fix}}(\tilde{s}(\alpha), N)$ we use the approximation (B.1), and for $t_{\text{fix}}(\tilde{s}(\alpha), N)$ we use (B.4). With this

Figure B.1: Panels A and C demonstrate the accuracy of our method of approximating the true diffusion approximation for the mean time to fixation (t_{fix}) or loss (t_{loss}) by concatenation of a linear approximation for small s and a simplified approximation, i.e., of eqs. (B.4) and (B.7). In panel A, the mean fixation times $t_{\text{fix}}^{(\text{small})}(s, N)$ for $s < 3/(2N)$ (orange, (B.3)) and $t_{\text{fix}}^{(\text{HP})}(s, N)$ for $s \geq 3/(2N)$ (red, (B.2)) are compared with the exact diffusion approximation (black) for $N = 10^4$. Panel C is analogous to A and displays the mean loss times $t_{\text{loss}}^{(\text{small})}(s, N)$ for $s < 2/N$ (orange, (B.6)) and $t_{\text{loss}}^{(\text{app})}(s, N)$ for $s \geq 2/N$ (red, (B.5a)). In both panels, the exact diffusion approximations are computed by numerical integration of the corresponding sojourn time densities (Ewens, 2004, Sects. 4.6 and 5.3), and the dashed red curves show $t_{\text{fix}}^{(\text{HP})}$ or $t_{\text{loss}}^{(\text{app})}$, where they are inaccurate and substituted by $t_{\text{fix}}^{(\text{small})}$ or $t_{\text{loss}}^{(\text{small})}$, respectively. In addition, for mutation effects drawn from an exponential distribution with mean 1, the expected mean times to fixation ($\bar{t}_{\text{fix}}$ in (B.10)) or loss ($\bar{t}_{\text{loss}}$ in (B.16)) are displayed in blue. Panel B shows $t_{\text{fix}}(\tilde{s}(\alpha), N)$ from (B.4) (dashed curves) and $\bar{t}_{\text{fix}}(s, 1, N)$ from (B.10) (solid curves) as functions of $s \geq 10^{-4}$ for different population sizes (blue for $N = 10^4$, purple for $N = 10^3$, red for $N = 10^2$). Simulation results from the Wright-Fisher model for equal mutation effects $\alpha = 1$ and exponentially distributed effects with mean $\bar{\alpha} = 1$ are shown as open and filled symbols, respectively. Panel D is analogous and depicts $t_{\text{loss}}(\tilde{s}(\alpha), N)$ from (B.7) (dashed curves) and $\bar{t}_{\text{loss}}(s, 1, N)$ from (B.16) (solid curves). In accordance with (B.5a), panel D illustrates that t_{loss} and $\bar{t}_{\text{loss}}$ depend only very weakly on N if $s > 2/N$.

simplified diffusion approximation for $t_{\text{fix}}(\tilde{s}(\alpha), N)$, the integrals in (B.10) are readily computed numerically. Some values are given in the legend of Fig. 4.2.

An analytically explicit, still quite accurate, approximation of $\bar{t}_{\text{fix}}$ can be derived as follows. We use the approximations $P_{\text{fix}}(s, N) \approx 1 - e^{-2s}$ and $t_{\text{fix}}^{(\text{HP})}(s, N) \approx \frac{2}{s}\left(\ln(2Ns) + \gamma\right)$, which are valid for sufficiently large Ns. In addition, we assume that the selective advantage of the mutant is $s\alpha$

instead of $\tilde{s}(\alpha)$. If f is exponentially distributed with mean $\bar{\alpha}$, we obtain

$$\int_0^\infty P_{\text{fix}}(s\alpha, N)f(\alpha)\,d\alpha \approx \frac{2s\bar{\alpha}}{1 + 2s\bar{\alpha}} \tag{B.11}$$

and

$$\int_0^\infty t_{\text{fix}}^{(\text{HP})}(s\alpha, N)P_{\text{fix}}(s\alpha, N)f(\alpha)\,d\alpha \approx \frac{\ln(1 + 2s\bar{\alpha})}{s\bar{\alpha}}\left[2\ln(2Ns\bar{\alpha}) - \ln(1 + 2s\bar{\alpha})\right]. \tag{B.12}$$

Therefore,

$$\bar{t}_{\text{fix}}(s, \bar{\alpha}, N) \approx \frac{(1 + 2s\bar{\alpha})\ln(1 + 2s\bar{\alpha})}{2(s\bar{\alpha})^2}\left[2\ln(2Ns\bar{\alpha}) - \ln(1 + 2s\bar{\alpha})\right]. \tag{B.13}$$

The high accuracy of this approximation is demonstrated in the supplementary *Mathematica* notebook. Assuming $s\bar{\alpha} \ll 1$ and performing a series expansion in $s\bar{\alpha}$, we arrive at

$$\bar{t}_{\text{fix}}(s, \bar{\alpha}, N) \approx 2\ln(2Ns\bar{\alpha})\left(\frac{1}{s\bar{\alpha}} + 1 - \frac{2}{3}s\bar{\alpha}\right) - 2. \tag{B.14}$$

If the selective coefficient of the mutants is $\tilde{s}(\alpha)$, instead of $s\alpha$, then some of the above integrations cannot be performed analytically. However, numerics shows that the simple approximation

$$\bar{t}_{\text{fix}}(s, \bar{\alpha}, N) \approx \frac{2}{s\bar{\alpha}}\ln(2Ns\bar{\alpha}). \tag{B.15}$$

still works very well (see *Mathematica* notebook). This is the approximation given in Sect. 4.3.

B.3. Mean loss time averaged over the mutation distribution f

In analogy to (4.14) and (B.10), we define the expected mean loss time of a single mutant, with effect α on the selected trait drawn from an exponential distribution f with mean $\bar{\alpha}$, by

$$\bar{t}_{\text{loss}}(s, \bar{\alpha}, N) = \frac{\int_0^\infty t_{\text{loss}}(\tilde{s}(\alpha), N)(1 - P_{\text{fix}}(\tilde{s}(\alpha), N))f(\alpha)\,d\alpha}{\int_0^\infty (1 - P_{\text{fix}}(\tilde{s}(\alpha), N))f(\alpha)\,d\alpha}. \tag{B.16}$$

Here, $t_{\text{loss}}(\tilde{s}(\alpha), N)$ is taken from (B.7). With this simplified diffusion approximation for $t_{\text{loss}}(\tilde{s}(\alpha), N)$, the integrals in (B.16) are readily computed numerically (see *Mathematica* notebook). For graphs of t_{loss} and $\bar{t}_{\text{loss}}$, see Fig. B.1. It shows that, unless s is very small, $\bar{t}_{\text{loss}} > t_{\text{loss}}$. The reasons are that mutations of larger effect are lost with lower probability than those of smaller effect, and small effect mutations stay longer in the population. The following is a simple and accurate approximation if (approximately) $Ns\bar{\alpha} > 1$ and $s\bar{\alpha} < 0.2$ (see *Mathematica* notebook):

$$\bar{t}_{\text{loss}}(s, \bar{\alpha}, N) \approx 2(-\ln(2s\bar{\alpha}) + 1 + 2s\bar{\alpha}) - \frac{1.42 - \ln(Ns\bar{\alpha})}{Ns\bar{\alpha}}. \tag{B.17}$$

Usually, these times to loss are very small. For instance, if the mean of the exponential distribution is $\bar{\alpha} = 1$, $N = 10^4$, we obtain $\bar{t}_{\text{loss}} \approx 5.5$, 8.8, and 14.5 for $s = 0.1$, 0.01, and 0.001, respectively.

1418 For larger values of N almost the same numbers are obtained.

C. Technical results

1419

C.1. Poisson and Erlang distribution

1420

1421   Here, we prove (4.38).

$$
\begin{aligned}
\sum_{n=1}^{\infty} \mathrm{Poi}_{\Theta\tau}(n)\left(\sum_{i=1}^{n} \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)}\right) &= \sum_{n=1}^{\infty} e^{-\Theta\tau} \frac{(\Theta\tau)^n}{n!} \sum_{i=1}^{n} e^{-\Theta t} \frac{\Theta(\Theta t)^{i-1}}{\Gamma(i)M_{i,\Theta}(\tau)} \\
&= \Theta e^{-\Theta t} \sum_{i=1}^{\infty} \frac{(\Theta t)^{i-1}}{\Gamma(i)M_{i,\Theta}(\tau)} \sum_{n=i}^{\infty} e^{-\Theta\tau} \frac{(\Theta\tau)^n}{n!} \\
&= \Theta e^{-\Theta t} \sum_{i=1}^{\infty} \frac{(\Theta t)^{i-1}}{\Gamma(i)M_{i,\Theta}(\tau)} M_{i,\Theta}(\tau) = \Theta\,. \qquad \text{(C.1)}
\end{aligned}
$$

1422   The first equality is simply based on the definitions of $\mathrm{Poi}_{\Theta\tau}(n)$ in (2.3) and $m_{i,\Theta}(t)$ in (2.2).
1423   The second is obtained by reordering the sums, the third from a well-known relation between the
1424   Poisson distribution and the incomplete Gamma function (e.g. Abramowitz and Stegun, 1964,
1425   6.5.13) and the definition of $M_{i,\Theta}(\tau)$ in (3.19), and the last from the exponential series.

C.2. Some integral formulas

1426

1427   In the proof of Propositions 4.5 and 4.11, the following integrals are needed. They were evaluated
1428   with *Mathematica*, but can be obtained by partial integration using $\int E_1(x)\,dx = -e^{-x} + xE_1(x)$.

$$
\int_0^{t_1} \gamma_1(\tau - t, e^{s\alpha})\,dt = \frac{1}{s\alpha}\left[\exp(\nu e^{-s\alpha\tau})E_1(\nu e^{-s\alpha\tau}) - \exp(\nu e^{-s\alpha(\tau-t_1)})E_1(\nu e^{-s\alpha(\tau-t_1)})\right],
$$
$$\text{(C.2a)}$$

$$
\int_{t_1}^{\tau} \gamma_1(\tau - t, e^{s\alpha})\,dt = \frac{1}{s\alpha}\left[\exp(\nu e^{-s\alpha(\tau-t_1)})E_1(\nu e^{-s\alpha(\tau-t_1)}) - e^{\nu}E_1(\nu)\right], \qquad \text{(C.2b)}
$$

1429   as well as

$$
\int_0^{\tau} \gamma(\tau - t, e^{s\alpha})\,dt = \frac{1}{s\alpha}\left[e^{\nu}\nu E_1(\nu) - \exp[\nu e^{-s\alpha\tau}]\nu e^{-s\alpha\tau}E_1(\nu e^{-s\alpha\tau})\right] \qquad \text{(C.3a)}
$$

1430   and

$$
\lim_{\tau\to\infty}\int_0^{\tau}\gamma(\tau - t, e^{s\alpha})\,dt = \lim_{\tau\to\infty}\int_0^{\tau}\gamma(t, e^{s\alpha})\,dt = \frac{\nu}{s\alpha}e^{\nu}E_1(\nu)\,. \qquad \text{(C.3b)}
$$

1431   With $t_1 = \tau - \bar{t}_{\mathrm{fix}}$, these yield

$$
\int_0^{\tau-\bar{t}_{\mathrm{fix}}} \gamma_1(\tau - t, e^{s\alpha})\,dt = \frac{1}{s\alpha}\left[\exp(\nu e^{-s\alpha\tau})E_1(\nu e^{-s\alpha\tau}) - \exp(\nu e^{-s\alpha\bar{t}_{\mathrm{fix}}})E_1(\nu e^{-s\alpha\bar{t}_{\mathrm{fix}}})\right], \quad \text{(C.4)}
$$

1432   and

$$
\int_{\tau-\bar{t}_{\mathrm{fix}}}^{\tau} \gamma_1(\tau - t, e^{s\alpha})\,dt = \frac{1}{s\alpha}\left[\exp(\nu e^{-s\alpha\bar{t}_{\mathrm{fix}}})E_1(\nu e^{-s\alpha\bar{t}_{\mathrm{fix}}}) - e^{\nu}E_1(\nu)\right]. \qquad \text{(C.5)}
$$

From the asymptotic properties of the exponential integral (Abramowitz and Stegun, 1964, Chap. 5.1), we derive

$$\lim_{T \to \infty} \frac{1}{s\alpha T} \left[\exp(\nu e^{-s\alpha(T_1+T)}) E_1(\nu e^{-s\alpha(T_1+T)}) - \exp(\nu e^{-s\alpha T_1}) E_1(\nu e^{-s\alpha T_1}) \right] = 1 \, .$$

Therefore, for sufficiently large τ, we obtain

$$\int_0^{\tau - \bar{t}_{\text{fix}}} \gamma_1(\tau - t, e^{s\alpha}) \, dt \approx \tau - \bar{t}_{\text{fix}} \, . \tag{C.6}$$

C.3. Mutations occur at their expected time

An alternative, and potentially, simpler approach than using the Erlang distribution is to assume that the ith mutation occurs at its mean waiting time i/Θ and the number of mutation events until time τ is (approximately) $[\Theta\tau]$. Then (4.6) becomes

$$\mathrm{E}\left[(\tilde{X}_\tau^{(i)})^k\right] \approx (1 - P_{\text{ext}}([\tau - i/\Theta])) \int_0^1 x^k g_{a(\tau - i/\Theta)}(x) \, dx \, . \tag{C.7}$$

Consequently, (4.9) and (4.10) simplify to

$$\bar{G}(\tau) \approx \int_0^\infty \alpha f(\alpha) \left(\sum_{i=1}^{[\Theta\tau]} (1 - P_{\text{ext}}([\tau - i/\Theta], e^{s\alpha})) \, \gamma_1(a(\tau - i/\Theta, e^{s\alpha})) \right) d\alpha \tag{C.8a}$$

$$= \int_0^\infty \alpha f(\alpha) \left(\sum_{j=0}^{[\Theta\tau]-1} (1 - P_{\text{ext}}([j/\Theta], e^{s\alpha})) \, \gamma_1(a(j/\Theta, e^{s\alpha})) \right) d\alpha \, , \tag{C.8b}$$

and

$$V_G(\tau) \approx \int_0^\infty \alpha^2 f(\alpha) \left(\sum_{i=1}^{[\Theta\tau]} (1 - P_{\text{ext}}([\tau - i/\Theta], e^{s\alpha})) \, \gamma(a(\tau - i/\Theta, e^{s\alpha})) \right) d\alpha \tag{C.9a}$$

$$= \int_0^\infty \alpha^2 f(\alpha) \left(\sum_{j=0}^{[\Theta\tau]-1} (1 - P_{\text{ext}}([j/\Theta], e^{s\alpha})) \, \gamma(a(j/\Theta, e^{s\alpha})) \right) d\alpha \, , \tag{C.9b}$$

respectively. For the second equations we have reversed the summation order, i.e., $j = [\Theta\tau] - i$.

Instead of the approximations (4.17) and (4.18) in Proposition 4.5, this approach yields the following. We obtain from (C.8b) for sufficiently large τ by substituting $P_{\text{ext}}^\infty(e^{s\alpha})$ for $P_{\text{ext}}([j/\Theta], e^{s\alpha})$,

$$\bar{G}^*(\tau) \approx \int_0^\infty \alpha f(\alpha)(1 - P_{\text{ext}}^\infty(e^{s\alpha})) \sum_{j=0}^{[\Theta\tau]-1} \gamma_1\big(a([j/\Theta], e^{s\alpha})\big) \, d\alpha$$

$$\approx \int_0^\infty \alpha f(\alpha) P_{\text{sur}}(e^{s\alpha}) \int_0^{\Theta\tau} \gamma_1\big(a(t/\Theta, e^{s\alpha})\big) \, dt \, d\alpha \, ,$$

which becomes (4.41b) after a simple substitution. Following the derivation in (4.41d), we arrive at (4.17). Analogous arguments and results apply to the variance at stationarity.

For small τ we assume (4.47), as in the proof of Proposition 4.13. Substituting this into (C.8b) yields

$$
\begin{aligned}
\bar{G}(\tau) &\approx \frac{1}{N} \int_0^\infty \alpha \left(\sum_{j=0}^{[\Theta\tau]-1} e^{s\alpha[j/\Theta]} \right) f(\alpha)\, d\alpha \\
&\approx \frac{1}{N} \int_0^\infty \alpha \left(\sum_{j=0}^{\Theta\tau-1} e^{s\alpha j/\Theta} \right) f(\alpha)\, d\alpha \\
&= \frac{1}{N} \int_0^\infty \alpha \frac{e^{s\alpha\tau}-1}{e^{s\alpha/\Theta}-1} f(\alpha)\, d\alpha \\
&\approx \frac{1}{N} \int_0^\infty \frac{\alpha(e^{s\alpha\tau}-1)f(\alpha)}{(s\alpha/\Theta)+\frac{1}{2}(s\alpha/\Theta)^2+\frac{1}{6}(s\alpha/\Theta)^3}\, d\alpha \\
&\approx \frac{\Theta\tau}{N} \left[1 + s\left(\tau-\frac{1}{\Theta}\right) + s^2\left(\tau^2-\frac{3\tau}{2\Theta}+\frac{1}{2\Theta^2}\right) \right] + O(s^3)\,, \qquad \text{(C.10)}
\end{aligned}
$$

where in the last step we assumed $f(\alpha) = e^{-\alpha}$; cf. (4.32b). The leading-order term provides an accurate approximation if (approximately) $s\tau \leq \frac{1}{2}$. With the second-order term, the approximation is accurate if (approximately) $s\tau \leq 0.8$. It also requires that (approximately) $\Theta/s \geq 10$.

D. Segregating sites: A branching process approximation

Here we derive an alternative approximation for the number of segregating sites by relying on our branching process approach, similar as in Sect. 4.2, and amending it by a diffusion approximation for the mean fixation time. At a given generation τ, $\Theta\tau$ mutations have occurred on average. A mutant with fitness σ is still present in the population with probability $1 - P_{\text{ext}}(\tau - t, \sigma)$, where t is the number of generations since its occurrence and $P_{\text{ext}}(n, \sigma)$ is the probability of loss until generation n; see (3.12). On average, $t = i/\Theta$ for the ith mutation event. Some of those mutations may have already reached fixation and are no longer segregating in the population. Fixation occurs on average after $\bar{t}_{\text{fix}}$ generations (see eqs. 4.14 and B.15).

With this in mind and in the spirit of the derivations for the phenotypic mean and variance (Sect. 4.2), we define random variables $S^{(i)}_{\tau,e^{s\alpha},\Theta}$, which are 1 when locus i is segregating in the population and 0 otherwise. Using the expected fixation time t_{fix} in (B.1) from the diffusion approximation, we define

$$
\tilde{\tau} = \min\{t_{\text{fix}}(\tilde{s}(\alpha), N), \tau\}\,, \qquad \text{(D.1)}
$$

which is the expected time a mutant present in generation τ has already been segregating. Then,

$$
\text{E}[S^{(i)}_{\tau,e^{s\alpha},\Theta}] \approx \int_0^{\tilde{\tau}} \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)}(1 - P_{\text{ext}}([\tilde{\tau}-t], e^{s\alpha}))dt\,. \qquad \text{(D.2)}
$$

We define the expected number of segregating sites in our model, where mutation effects are drawn

Figure D.1: Scaled expected number of segregating sites, $E[S]/\Theta$ as a function of $s\tau$ in the early phase of adaptation ($s\tau \leq 2$) for the selection intensities $s = 0.1, 0.01, 0.001$ (blue, red, black). In A the mutation effects are equal ($\alpha = 1$), in B they are drawn from an exponential distribution with mean $\bar{\alpha} = 1$. The branching process approximations $\tilde{S}$ in (5.5) and (5.6) are shown by solid curves, the diffusion approximation $\bar{S}$ in (5.1) by dashed curves. The kinks occur at st_{loss}. Results from Wright-Fisher simulations (with $\Theta = \frac{1}{2}$) are shown as symbols. The mutation effects are $\alpha = 1$ and the population size is $N = 10^4$.

1467   from a distribution f and the offspring distribution is Poisson, by

$$\mathrm{E}[S](\tau, f, s, N, \Theta) = \sum_{n=1}^{\infty} \mathrm{Poi}_{\Theta\tau}(n) \left(\sum_{i=1}^{n} \int_{0}^{\infty} \mathrm{E}\left[\tilde{S}_{\tau, e^{s\alpha}, \Theta}^{(i)} \right] f(\alpha)\, d\alpha \right). \tag{D.3}$$

1468   By substituting the approximation (D.2) into (D.3), we obtain the following approximation for
1469 $\mathrm{E}[S]$:

$$\begin{aligned}
\tilde{S}(\tau, f, s, N, \Theta) &= \sum_{n=1}^{\infty} \mathrm{Poi}_{\Theta\tau}(n) \left(\sum_{i=1}^{n} \int_{0}^{\infty} \int_{0}^{\tilde{\tau}} \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)} \left(1 - P_{\mathrm{ext}}([\tilde{\tau} - t], e^{s\alpha})\right) dt\, f(\alpha)\, d\alpha \right) \\
&= \int_{0}^{\infty} \int_{0}^{\tilde{\tau}} \sum_{n=1}^{\infty} \mathrm{Poi}_{\Theta\tau}(n) \left(\sum_{i=1}^{n} \frac{m_{i,\Theta}(t)}{M_{i,\Theta}(\tau)} \right) \left(1 - P_{\mathrm{ext}}([\tilde{\tau} - t], e^{s\alpha})\right) dt\, f(\alpha)\, d\alpha \\
&= \Theta \int_{0}^{\infty} f(\alpha) \int_{0}^{\tilde{\tau}} \left(1 - P_{\mathrm{ext}}([\tilde{\tau} - t], e^{s\alpha})\right) dt\, d\alpha \\
&\approx \Theta \int_{0}^{\infty} f(\alpha) \sum_{i=0}^{[\tilde{\tau}]} \left(1 - P_{\mathrm{ext}}([\tilde{\tau}] - i, e^{s\alpha})\right) d\alpha \\
&= \Theta \int_{0}^{\infty} f(\alpha) \sum_{j=0}^{[\tilde{\tau}]} \left(1 - P_{\mathrm{ext}}(j, e^{s\alpha})\right) d\alpha. \tag{D.4}
\end{aligned}$$

1470   The above calculations are analogous to those in the derivation of (4.9) for $\bar{G}(\tau)$ in (4.39) upon
1471   substitution of $\alpha\, \mathrm{E}[\tilde{X}_{\tau, e^{s\alpha}, \Theta}^{(i)}]$ by $\mathrm{E}[S_{\tau, e^{s\alpha}, \Theta}^{(i)}]$.

1472   If all effects are equal to α, then $\tilde{\tau} = \tilde{\tau}(\alpha)$ is constant and the integration in (D.4) collapses to
1473   the computation of the sum, i.e., to (5.5) in the main text.

1474   Unless the distribution $f(\alpha)$ is a delta function, as for equal mutation effects, no further

simplification of (D.4) seems possible, except when $\tau \ll \bar{t}_{\text{fix}}(\tilde{s}(\alpha), N)$. Then, for given small τ, we have $\tilde{\tau} = \tau$ except for the (very rare) very large values of α in the tail of f for which $t_{\text{fix}}(\tilde{s}(\alpha), N) < \tau$. As a consequence, we obtain from (D.4) the approximation (5.6) in the main text.

 ## S. Supplementary figures

Figure S.1: Shape of $a(n, \sigma, N)$ in eq. (3.16b) as a function of the number of generations n for different selection coefficients s. The population size is $N = 10^4$ and the offspring distribution is Poisson with mean $\sigma = e^s$.

Figure S.2: The scaled expected phenotypic mean $\bar{G}(s\tau)/\Theta$ (panel A), the scaled expected phenotypic variance $V_G(s\tau)/\Theta$ (panel B), and the scaled expected number of segregating sites $\bar{S}(s\tau)/\Theta$ (panel C) are shown as functions of $s\tau$ for the selection coefficients $s = 0.1$ and $s = 0.001$ (blue and black, respectively). Equal mutation effects are assumed, i.e., $\alpha_i = 1$. Analytic approximations, with $\bar{G}(s\tau)$ computed from (4.11), $V_G(s\tau)$ from (4.12), and $\bar{S}(s\tau)$ from (5.1), are shown as solid curves. Wright-Fisher-simulation results with $2\Theta = 10$ and $2\Theta = 0.01$ (open and filled symbols, respectively) are shown as symbols. Whereas Θ is a multiplicative factor in the analytic approximations, these simulation results show fluctuations around the predicted curve if $2\Theta = 0.01$. Presumably, the reason is that with such small Θ, the waiting time for successful mutations is very long, about $1/(2s\Theta)$. On the time scale of the figure, this yields $1/(2\Theta) = 100$ units. Such rare successful mutations will entail substantial stochastic effects. We note that very slight fluctuations are also visible in Figures 4.2 and 5.1, where $2\Theta = 1$.

Figure S.3: The black curves show the probability density $g_a(x)$ in (3.16) of the mutant frequency in different generations n for various combinations of $a = a(n, \sigma, N)$ and $\sigma = e^s$ (conditioned on non-extinction until generation n). The population size is $N = 10^4$. The blue histograms are obtained from simulations of the corresponding Wright-Fisher model. The analytic approximation $g_a(x)$ underestimates the effect of random genetic drift near fixation (x large and $n > \frac{1}{2}t_{\text{fix}} \approx \frac{1}{s}\ln(2Ns)$), unless selection is very strong relative to random drift (top row, where $Ns = 10^3$). For $s = 0.1$, 0.01, 0.001, and 0.0001, we have $t_{\text{fix}} \approx 156$, 1060, 5991, and $13\,863$, respectively. In general, and according to intuition and assumptions, the accuracy of the approximation by $g_a(x)$ decreases with larger x and n, and smaller Ns. Nevertheless, for very early stopping times n relative to t_{fix}, the approximation remains useful even if $Ns = 1$ (left bottom). Its mean, given by (4.1a), stays accurate in a much wide parameter range than $g_a(x)$ (Fig. 4.1.A).

Figure S.4: The scaled expected phenotypic mean and variance, $\bar{G}(s\tau)/\Theta$ and $V_G(s\tau)/\Theta$ as function of scaled time (panels A and B), and the scaled expected phenotypic variance at stationarity, $V_G^*(s\tau)/\Theta$ (panel C) as a function of s for a truncated normal distribution with $\alpha \geq 0$ and $\bar{\alpha} = 1$ (cf. Remark 4.10). Panels A and B are analogous to Fig. 4.2 and panel C is analogous to Fig. 4.3. Analytic approximations are shown as curves, Wright-Fisher simulation results (for $\Theta = \frac{1}{2}$) as symbols.

In panels A and B, we have $N = 10^4$ and selection coefficients and color code are shown in the legends. The solid curves show $\bar{G}(s\tau)/\Theta$ and $V_G(s\tau)/\Theta$ as computed from (4.9) and (4.10). The dashed lines show the equilibrium values given by (4.17) and (4.18). The values of the mean fixation times multiplied by $s = 0.1, 0.01, 0.001$ are $s\bar{t}_{\text{fix}} \approx 16.1, 11.0, 6.2$, respectively (computed from (4.14)).

In panel C, different colors correspond to different population sizes, as shown in the legend. The most accurate, available approximation is (4.18); it is shown as dotted curves. In addition, the scaled variance V_G^*/Θ is compared with the scaled expected change in the phenotypic mean $\Delta\bar{G}^*/(\Theta s)$, as calculated from (4.26) (black dashed curve). For the Wright-Fisher simulation results (circles for $N = 10^4$, squares for $N = 10^3$, triangles for $N = 10^2$), filled symbols are used for the scaled phenotypic variance and open symbols for the change in the scaled phenotypic mean.

Figure S.5: The expected number of segregating sites (rescaled by Θ) as a function of $s\tau$ (panel A) and at stationarity as a function of s (panel B). This figure is analogous to the rows in Fig. 5.1 but mutation effects are drawn from a normal distribution truncated at 0 with mean $\bar{\alpha} = 1$ (cf. Remark 4.10). Analytic approximations are computed from (5.1) and are shown as curves. Wright-Fisher simulation results for $\mathrm{E}[S]$ (with $\Theta = \frac{1}{2}$) are shown as symbols. In A, $\bar{S}/\Theta$ is shown for $N = 10^4$ and $s = 0.1, 0.01, 0.001$ (solid curves and filled symbols in blue, red, black), and for $N = 10^3$ and $s = 0.1$ (dashed curve and open symbols in blue). In B, $\bar{S}^*/\Theta$ is displayed for $N = 10^4, 10^3, 10^2$ (solid, dashed, dotted curves, and corresponding open symbols).

Figure S.6: Histograms of the stopping times T at which the mean phenotype $\bar{G}$ in a single population first reaches or exceeds the value 1 (this figure complements Fig. 6.1). Histograms are based on $20\,000$ replicate runs of the Wright-Fisher model. The vertical black lines represent the corresponding means $\bar{T}$, which are given in Fig. 6.1. The corresponding minimum and maximum values of T are given in curly braces. The selection coefficients s and the mutation effects α are indicated for each row. Columns show results for different values of Θ. The population size is $N = 10^4$ in all cases.

References

Abramowitz, M., Stegun, I.A., 1964. Handbook of mathematical functions with formulas, graphs, and mathematical tables. volume 55. US Government Printing Office.

Allen, L.J.S., 2003. An introduction to stochastic processes with applications to biology. Pearson/Prentice Hall, Upper Saddle River, N.J.

Athreya, K.B., Ney, P.E., 1972. Branching processes. Springer, Berlin-Heidelberg.

Barghi, N., Hermisson, J., Schlötterer, C., 2020. Polygenic adaptation: A unifying framework to understand positive selection. Nature Reviews Genetics 21, 769—-781.

Barghi, N., Tobler, R., Nolte, V., Jakšić, A.M., Mallard, F., Otte, K.A., Dolezal, M., Taus, T., Kofler, R., Schlötterer, C., 2019. Genetic redundancy fuels polygenic adaptation in drosophila. PLoS Biology 17, e3000128.

Barton, N.H., 1995. Linkage and the limits to natural selection. Genetics 140, 821–841.

Barton, N.H., Etheridge, A.M., Véber, A., 2017. The infinitesimal model: Definition, derivation, and implications. Theoretical Population Biology 118, 50–73.

Barton, N.H., Keightley, P.D., 2002. Understanding quantitative genetic variation. Nature Reviews Genetics 3, 11–21.

Berg, J.J., Coop, G., 2014. A population genetic signal of polygenic adaptation. PLoS Genetics 10, e1004412–e1004412.

Boyle, E.A., Li, Y.I., Pritchard, J.K., 2017. An expanded view of complex traits: From polygenic to omnigenic. Cell 169, 1177–1186.

Brillinger, D.R., 1969. The calculation of cumulants via conditioning. Annals of the Institute of Statistical Mathematics 21, 215–218.

Bulmer, M.G., 1980. The mathematical theory of quantitative genetics. Clarendon Press.

Bürger, R., 1993. Predictions of the dynamics of a polygenic character under directional selection. Journal of Theoretical Biology 162, 487–513.

Bürger, R., 2000. The mathematical theory of selection, recombination, and mutation. Wiley, Chichester.

Bürger, R., Ewens, W.J., 1995. Fixation probabilities of additive alleles in diploid populations. Journal of Mathematical Biology 33, 557–575.

Bürger, R., Lynch, M., 1995. Evolution and extinction in a changing environment: A quantitative-genetic analysis. Evolution 49, 151–163.

Charlesworth, B., 2020. How long does it take to fix a favorable mutation, and why should we care? The American Naturalist 195, 753–771.

Chevin, L.M., 2019. Selective sweep at a QTL in a randomly fluctuating environment. Genetics 213, 987–1005.

Chevin, L.M., Hospital, F., 2008. Selective sweep at a quantitative trait locus in the presence of background genetic variation. Genetics 180, 1645–1660.

Clayton, G., Robertson, A., 1955. Mutation and quantitative variation. The American Naturalist 89, 151–158.

De Vladar, H.P., Barton, N.H., 2014. Stability and response of polygenic traits to stabilizing selection and mutation. Genetics 197, 749–767.

Desai, M.M., Fisher, D.S., 2007. Beneficial mutation selection balance and the effect of linkage on positive selection. Genetics 176, 1759–1798.

Ewens, W.J., 1974. A note on the sampling theory for infinite alleles and infinite sites models. Theoretical Population Biology 6, 143–148.

Ewens, W.J., 2004. Mathematical population genetics: Theoretical introduction. Springer, New York.

Eyre-Walker, A., Keightley, P.D., 2007. The distribution of fitness effects of new mutations. Nature Reviews Genetics 8, 610–618.

Falconer, D.S., Mackay, T.F.C., 1996. Introduction to quantitative genetics. Longman Group, UK.

Fisher, R.A., 1918. The correlation between relatives on the supposition of mendelian inheritance. Transactions of the Royal Society of Edinburgh 52, 399–433.

Galassi, M., Davies, J., Theiler, J., Gough, B., Jungman, G., 2009. GNU scientific library reference manual. 3 ed.

Haccou, P., Jagers, P., Vatutin, V.A., 2005. Branching processes: Variation, growth, and extinction of populations. volume 5 of *Cambridge studies in adaptive dynamics*. Cambridge University Press.

Haldane, J.B.S., 1927. A mathematical theory of natural and artificial selection, part V: Selection and mutation. Mathematical Proceedings of the Cambridge Philosophical Society 23, 838—-844.

Hayward, L.K., Sella, G., 2022. Polygenic adaptation after a sudden change in environment. eLife 11, e66697.

Hermisson, J., Pennings, P.S., 2005. Soft sweeps: Molecular population genetics of adaptation from standing genetic variation. Genetics 169, 2335–2352.

Hermisson, J., Pennings, P.S., 2017. Soft sweeps and beyond: Understanding the patterns and probabilities of selection footprints under rapid adaptation. Methods in Ecology and Evolution 8, 700–716.

Hernandez, R.D., Kelley, J.L., Elyashiv, E., Melton, S.C., Auton, A., McVean, G., Project, .G., Sella, G., Przeworski, M., 2011. Classic selective sweeps were rare in recent human evolution. Science 331, 920–924.

Hill, W.G., 1982a. Rates of change in quantitative traits from fixation of new mutations. Proceedings of the National Academy of Sciences - PNAS 79, 142–145.

Hill, W.G., 1982b. Predictions of response to artificial selection from new mutations. Genetics Research 40, 255–278.

Hill, W.G., Kirkpatrick, M., 2010. What animal breeding has taught us about evolution. Annual Review of Ecology, Evolution, and Systematics 41, 1–19.

Höllinger, I., 2018. Mathematical models of speciation and polygenic adaptation. Ph.D. thesis. University of Vienna.

Höllinger, I., Pennings, P.S., Hermisson, J., 2019. Polygenic adaptation: From sweeps to subtle frequency shifts. PLoS Genetics 15, 1–26.

Hospital, F., Chevalet, C., 1996. Interactions of selection, linkage and drift in the dynamics of polygenic characters. Genetics Research 67, 77–87.

Huber, C.D., Degiorgio, M., Hellmann, I., Nielsen, R., 2016. Detecting recent selective sweeps while controlling for mutation rate and background selection. Molecular Ecology 25, 142–156.

Jain, K., Stephan, W., 2017a. Rapid adaptation of a polygenic trait after a sudden environmental shift. Genetics 206, 389–406.

Jain, K., Stephan, W., 2017b. Modes of rapid polygenic adaptation. Molecular Biology and Evolution 34, 3169–3175.

John, S., Stephan, W., 2020. Important role of genetic drift in rapid polygenic adaptation. Ecology and Evolution 10, 1278–1287.

Jones, A.G., Arnold, S.J., Bürger, R., 2004. Evolution and stability of the g-matrix on a landscape with a moving optimum. Evolution 58, 1639–1654.

Kaplan, N.L., Hudson, R.R., Langley, C.H., 1989. The" hitchhiking effect" revisited. Genetics 123, 887–899.

Keightley, P.D., Hill, W.G., 1983. Effects of linkage on response to directional selection from new mutations. Genetics Research 42, 193–206.

Keightley, P.D., Hill, W.G., 1987. Directional selection and variation in finite populations. Genetics 117, 573–582.

Kesten, H., Stigum, B.P., 1966. A limit theorem for multidimensional galton-watson processes. The Annals of Mathematical Statistics 37, 1211–1223.

Kim, Y., Stephan, W., 2003. Selective sweeps in the presence of interference among partially linked loci. Genetics 164, 389–398.

Kimura, M., Ohta, T., 1969. The average number of generations until fixation of a mutant gene in a finite population. Genetics 61, 763–771.

Lambert, A., 2006. Probability of fixation under weak selection: A branching process unifying approach. Theoretical Population Biology 69, 419–441.

Lande, R., 1976. Natural selection and random genetic drift in phenotypic evolution. Evolution 30, 314–334.

Lande, R., 1983. The response to selection on major and minor mutations affecting a metrical trait. Heredity 50, 47–65.

Lynch, M., Hill, W.G., 1986. Phenotypic evolution by neutral mutation. Evolution 40, 915–935.

Lynch, M., Lande, R., 1993. Evolution and extinction in response to environmental change, in: Kareiva, P.M., Kingsolver, J.G., Huey, R.B. (Eds.), Biotic Interactions and Global Change, Sinauer Associates, Sunderland MA. pp. 234–250.

Lynch, M., Walsh, B., 1998. Genetics and analysis of quantitative traits. Sinauer Associates, Sunderland MA.

Martin, G., Lambert, A., 2015. A simple, semi-deterministic approximation to the distribution of selective sweeps in large populations. Theoretical Population Biology 101, 40–46.

Matsumoto, M., Nishimura, T., 1998. Mersenne twister: A 623-dimensionally equidistributed uniform pseudorandom number generator 8, 3–30.

Matuszewski, S., Hermisson, J., Kopp, M., 2015. Catch me if you can: Adaptation from standing genetic variation to a moving phenotypic optimum. Genetics 200, 1255–1274.

Maynard Smith, J., Haigh, J., 1974. The hitch-hiking effect of a favourable gene. Genetics Research 23, 23–35.

Messer, P.W., Petrov, D.A., 2013. Population genomics of rapid adaptation by soft selective sweeps. Trends in Ecology & Evolution 28, 659–669.

Mieghem, P.V., 2005. The limit random variable *W of a branching process. Technical Report. Delft University of Technology.*

Nielsen, R., Williamson, S., Kim, Y., Hubisz, M.J., Clark, A.G., Bustamante, C., 2005. Genomic scans for selective sweeps using snp data. Genome Research 15, 1566–1575.

Otto, S.P., Barton, N.H., 1997. The evolution of recombination: Removing the limits to natural selection. Genetics 147, 879–906.

Pritchard, J.K., Pickrell, J.K., Coop, G., 2010. The genetics of human adaptation: Hard sweeps, soft sweeps, and polygenic adaptation. Current Biology 20, R208–R215.

Robertson, A., 1960. A theory of limits in artificial selection. Proceedings of the Royal Society of London B 153, 234–249.

Robertson, A., 1977. Artificial selection with a large number of linked loci, in: Pollak, E.J., Kempthorne, O., Bailey, T.B., et al. (Eds.), Proceedings of the 1st International Conference on Quantitative Genetics, Iowa State University Press. pp. 307–322.

Sella, G., Barton, N.H., 2019. Thinking about the evolution of complex traits in the era of genome-wide association studies. Annual Review of Genomics and Human Genetics 20, 461–493.

Seneta, E., 1969. Functional equations and the galton-watson process. Advances in Applied Probability 1, 1–42.

Setter, D., Mousset, S., Cheng, X., Nielsen, R., DeGiorgio, M., Hermisson, J., 2020. Volcanofinder: Genomic scans for adaptive introgression. PLoS Genetics 16, e1008867.

Steinrücken, M., Wang, Y.R., Song, Y.S., 2013. An explicit transition density expansion for a multi-allelic wright–fisher diffusion with general diploid selection. Theoretical Population Biology 83, 1–14.

Stephan, W., 2019. Selective sweeps. Genetics 211, 5–13.

Stephan, W., John, S., 2020. Polygenic adaptation in a population of finite size. Entropy 22, 907.

Tataru, P., Mollion, M., Glémin, S., Bataillon, T., 2017. Inference of distribution of fitness effects and proportion of adaptive substitutions from polymorphism data. Genetics 207, 1103–1119.

Thornton, K.R., 2019. Polygenic adaptation to an environmental shift: Temporal dynamics of variation under gaussian stabilizing selection and additive effects on a single trait. Genetics 213, 1513–1530.

Turelli, M., Barton, N.H., 1990. Dynamics of polygenic characters under selection. Theoretical Population Biology 38, 1–57.

Uecker, H., Hermisson, J., 2011. On the fixation process of a beneficial mutation in a variable environment. Genetics 188, 915–930.

Walsh, B., Lynch, M., 2018. Evolution and selection of quantitative traits. Oxford University Press, Oxford.

Wölfl, B., 2019. Footprints of polygenic adaptation of a quantitative trait under stabilizing selection. Master's thesis. Vrije Universiteit Amsterdam and University of Amsterdam.

Wolfram Research, I., 2022. Mathematica, Version 13.2.

Zhang, X.S., Hill, W.G., 2005a. Genetic variability under mutation selection balance. Trends in Ecology & Evolution 20, 468–470.

Zhang, X.S., Hill, W.G., 2005b. Predictions of patterns of response to artificial selection in lines derived from natural populations. Genetics 169, 411–425.