CTCF supports melanoma cell migration by repressing cholesterol biosynthesis

Lukasz Stanislaw Kaczmarczyk[†1], Nehora Levi[†1], Mali Salmon-Divon[1,2,*] and Gabi Gerlitz[1,*]

[1]*Department of Molecular Biology, Faculty of Life Sciences and Ariel Center for Applied Cancer Research, Ariel University, Ariel 40700, Israel.*

[2]*Adelson School of Medicine, Ariel University, Ariel 40700, Israel*

[†]*Equal contribution*

Running title: CTCF supports cell migration.

Key words: SREBP / chromatin / transcriptional control / cell migration.

*Correspondence to: Department of Molecular Biology, Faculty of Natural Sciences, Ariel University, Kiryat Hamada, Ariel 40700, Israel. E-mail addresses: malisa@ariel.ac.il (M. Salmon-Divon), gabige@ariel.ac.il (G. Gerlitz).

Abstract

CCCTC-binding factor (CTCF) is a zinc finger protein that regulates correct three-dimensional chromatin folding in addition to activation and repression of transcription and splicing. CTCF was found to affect tumor cell migration, however the molecular mechanism is still obscure. To identify the molecular mechanism by which CTCF supports melanoma cell migration we used mouse melanoma cells with a partial loss of function (pLoF) of CTCF generated by the CRISPR-Cas9 system. Here we show that CTCF pLoF inhibited cell migration rate. Using RNA sequencing we identified an increase in the expression of multiple enzymes in the cholesterol biosynthesis pathway along with an increase in the cellular cholesterol levels. Inhibition of cholesterol synthesis in CTCF pLoF cells by Fatostatin restored the cellular migration rate, suggesting that CTCF supports cell migration by inhibition of cholesterol synthesis. Detailed analysis of the promoter of *Hmgcs1*, an upstream enzyme in the cholesterol synthesis pathway, revealed that CTCF pLoF led to reduced H3K27me3 levels in parallel to increased binding of the transcriptional activator SREBP2. Moreover, inhibition of H3K27 methylation in wildtype cells led to increased SREBP2 binding to this promoter. Taken together, our results suggest that CTCF represses transcription of *Hmgc1* by promoting H3K27 methylation to suppress SREBP2 binding. By that, CTCF fine tune the cellular cholesterol levels to support cell migration.

Introduction

Cell migration is a fundamental process in embryogenesis and in additional systems such as the immune system [1–3]. Mutations and deregulation of the cellular migration processes are linked to various human diseases including cancer [4]. During cell migration, the nucleus undergoes major changes in its position, structure and morphology [5–10]. One of these changes is global chromatin condensation, which was initially found in mouse melanoma cells migrating in the wound healing assay [11,12] and later on in additional cell types including human breast cancer cells [13], human and mouse CD4+ T-cells [14], rat tenocytes [15] and human fibrosarcoma cells [16]. Global chromatin condensation supports cell migration as concluded from dozens of experiments in which interfering with various chromatin condensation factors including histone modifying enzymes, DNA modifying enzymes and histone H1 altered cell migration rate [11,12,14,16,17,17–20]. Heterochromatin supports cell migration by altering the mechanical properties of the nucleus and by regulation of gene expression [19]. Chromatin condensation was shown to enhance nuclear deformability [20] and elasticity [15,21,22] that is thought to enable a bigger forward jump of the nucleus once it is extracted from a narrow pore, leading to an increase in cell migration rate [23]. The heterochromatin marker methylated H3K27 was found to support transcriptional changes in various genes as well as to prevent changes in transcription of hundreds of other genes [24].

To enhance the understanding of the importance of chromatin organization for cancer cell migration we focused on the chromatin architectural protein CCCTC-binding factor (CTCF). CTCF contains unstructured N' and C' termini with a central region that contains 11 zinc-finger domains, which are responsible for binding a long and complex DNA motif [25–27]. CTCF is involved in regulation of topologically associating domains (TADs) that are chromatin domains of strong self-contact comprising several hundred kb. TADs are thought to form by a loop-extrusion

mechanism in which cohesin complexes form chromatin loops by their ATP dependent molecular motor activity until halted by chromatin bound CTCF; thus, the resulted TAD is folded on itself while it is bound by cohesins and CTCF at its boundaries. Interactions between non-adjacent chromatin elements such as promoter and enhancer more commonly occur within a TAD rather than between separated TADs [27–29]. Loss of CTCF binding at TAD boundaries weakens the boundaries and their ability to insulate contacts between domains leading to transcriptional deregulation [30,31]. CTCF can affect transcription also by direct promoter binding to either repress transcription probably by recruitment of additional repressive factors such as the transcriptional repressor SIN3A [25,32] or to promote transcription by recruitment of transcription factors, repression of antisense transcription [25,32] and promotion of promoter-enhancer interactions [33,34].

CTCF involvement in cell migration is ambiguous; on one hand it was shown to repress breast cancer cell migration and invasion [35,36], while on the other hand it was shown to support migration and/or invasion of various primary and cancerous cells including human corneal epithelial cells [37,38], cortical neurons [39], skin-resident dendritic cells [40], prostate cancer cells [41,42], ovarian cancer cells [43], gastric cancer cells [44] and melanoma cells [45]. Although it is assumed that CTCF affects cell migration by its ability to regulate transcription, the exact mechanism by which CTCF controls transcription to control cell migration has not been determined yet.

Here we show that a partial loss of function (pLoF) of CTCF leads to slower migration of melanoma cells. Transcriptome analysis identified upregulation of cholesterol biosynthesis enzymes along with elevated cholesterol levels in CTCF pLoF cells. The elevated levels of cholesterol in CTCF pLoF cells interfered with their migration rate since inhibition of cholesterol biosynthesis restored their migration capabilities. Detailed analysis of the promoter of the cholesterol biosynthesis enzyme *Hmgcs1* showed that CTCF promotes recruitment of PRC2 to this

promoter to methylate H3K27. A methylation process that prevents binding of SREBP2 to the promoter and increased transcription of *Hmgcs1*. Thus, CTCF emerges as a new regulator of cholesterol biosynthesis.

Results and Discussion

To evaluate the effect of CTCF on cell migration we used the mouse melanoma B16-F1 cells in which we reduced CTCF expression levels by the CRISPR-Cas9 system [46]. Targeting of Cas9 to the beginning of the *CTCF* gene led to elimination of the full-length form of CTCF and the appearance of truncated forms of CTCF at low levels that are probably formed from downstream AUGs. The total amount of the CTCF shorter forms is 30-50% of the full-length protein in the parental cells. In addition, CTCF truncated forms have lower chromatin binding affinity than the full-length protein, thus these clones have a partial loss of function (pLoF) of CTCF. CTCF pLoF led to 20-40% reduction in the cellular migration rate in the wound healing assay and to 48-86% reduction in the cellular migration rate in the Transwell assay (Fig. 1a-c).

Fig. 1 **CTCF pLoF attenuates cell migration and elevates cellular cholesterol levels.** (**a**) CTCF pLoF attenuates the migration rate of B16-F1 cells in the wound healing assay. Confluent parental B16-F1 cells (WT) and two CTCF pLoF clones (C13 and C21) were induced to migrate for 8h. Representative pictures of the cells immediately after the scratch (0 hours) and eight hours after the scratch are shown. The wound edge is indicated with dashed white line. Scale bar: 50μm. (**b**) For quantification, the area covered by the cells after incubation of eight hours was measured using ImageJ and normalized to the parental B16-F1 cells (WT). The graph shows the mean relative scratch coverage in five independent experiments ± s.e. Statistical significance in comparison to WT cells was calculated by the Student's *t*-test, *$P<0.05$, **$P<0.01$. (**c**) CTCF

pLoF attenuates the migration rate of B16-F1 cells in the Transwell assay. Three hours after plating the WT and CTCF pLoF cells on top of 5 µm filters the cells were fixed, permeabilized, and stained with Hoechst reagent. In each experiment the fraction of the cells migrated to the lower side of the filter was calculated. The bar graph represents the mean percentage of migrating cells from the total number of cells on the filter in three independent experiments ± s.e. Statistical significance in comparison to WT cells was calculated by the Student's *t*-test, *$P<0.05$, **$P<0.01$. (**d**) Altered gene expression in CTCF pLoF clones. A volcano plot of altered genes comparing two combined results of two CTCF pLoF clones to WT; four repetitions for each cell type (WT, C13, C21). The graph represents unchanged genes (grey), genes with increased expression (red, 131 genes with fold change ≥2 and $P<0.05$) and genes with decreased expression (blue, 233 genes with fold change ≤-2 and $P<0.05$). (**e**) Top 5 pathways that are altered in CTCF pLoF clones identified by IPA analysis of 364 differentially expressed genes. The top 5 altered pathway were all activated. (**f**) Cellular cholesterol levels are increased in CTCF pLoF cells. The relative cholesterol level WT and CTCF pLoF clones normalized to the total protein level in relation to WT in three independent experiments ± s.e. Statistical significance in comparison to WT cells was calculated by the Student's *t*-test, *$P<0.05$, ****$P<0.0001$.

To look for the molecular mechanism for that phenotype we analysed the transcriptome of CTCF

pLoF cells and identified 364 significantly changed genes with at least 2-fold change in their

transcript level; 233 genes were downregulated, and 131 genes were upregulated (Fig. 1d, Sup.

Fig. 1). To identify affected pathways, we used the Ingenuity Pathway Analysis (IPA, Qiagen). As

shown in Fig. 1e, the most significantly up-regulated pathway in CTCF pLoF cells was cholesterol

biosynthesis. Cholesterol measurement confirmed that CTCF pLoF clones have an increase of 50-

70% in their cellular cholesterol levels (Fig. 1f). To validate the dependency of reduced cell

migration rate on elevated cholesterol levels in CTCF pLoF cells we repeated the migration assays

in cells treated with Fatostatin. Fatostatin is an inhibitor of activation of Sterol Regulatory-Element

Binding Proteins (SREBPs) that are the master regulators of transcriptional activation of

cholesterol biosynthesis enzymes [47,48]. As shown in Fig. 2, Fatostatin treatment elevated the

migration rate of CTCF pLoF cells to almost the migration rate of WT cells in both the wound

healing assay and the Transwell assay. Thus, it seems that elevated cholesterol levels in CTCF

pLoF cells are the main cause for their faulty migration capabilities.

Fig. 2 **Inhibition of cholesterol synthesis improves the migration capabilities of CTCF pLoF cells.** (**a**) Fatostatin improves the migration rate of CTCF pLoF cells in the wound healing assay. Confluent parental B16-F1 cells (WT) and two CTCF pLoF clones (C13 and C21) treated and non-treated with 5 μM Fatostatin for 24 hours prior to the experiment were induced to migrate for 8h. Representative pictures of the cells immediately after the scratch (0 hours) and eight hours after the scratch are shown. The wound edge is indicated with dashed white line. Scale bar: 50μm. (**b**) For quantification, the area covered by the cells after incubation of eight hours was measured using ImageJ and normalized to the non-treated parental B16-F1 cells (WT). The graph shows the mean relative

scratch coverage in three independent experiments ± s.e. Statistical significance was calculated by the Student's *t*-test, *P<0.05. (**c**) Fatostatin improves the migration rate of CTCF pLoF cells in the Transwell assay. Parental B16-F1 cells (WT) and two CTCF pLoF clones (C13 and C21) treated and non-treated with

5 µM Fatostatin for 24 hours prior to the experiment were plated on top of 5 µm filters. After three hours the cells were fixed, permeabilized, and stained with Hoechst reagent. In each experiment the fraction of the cells migrated to the lower side of the filter was calculated. The bar graph represents the mean percentage of migrating cells from the total number of cells on the filter in three independent experiments ± s.e. Statistical significance was calculated by the Student's *t*-test, **P<0.01.

IPA upstream regulator analysis suggested SREBPs as possible altered factors in CTCF pLoF cells (Sup. Fig. 2). We did not identify any increase at their RNA levels (Fig. 1) nor at their protein levels (Sup. Fig. 2). To analyze SREBPs function we focused on one of the first enzymes in the cholesterol biosynthesis pathway; 3-Hydroxy-3-Methylglutaryl-CoA Synthase 1 (Hmgcs1). ChIP analysis revealed an increased binding of SREBP2 to *Hmgcs1* promoter in CTCF pLoF cells in comparison to the parental cells (Fig. 3). We hypothesized that Srebp2 binding to *Hmgcs1* promoter is inhibited in the parental cells due to methylation at H3K27, since in a previous study we found that inhibition of H3K27 methylation in B16-F1 cells led to increased expression of cholesterol biosynthesis enzymes [24]. ChIP analysis of *Hmgcs1* promoter revealed reduced H3K27me3 levels in CTCF pLoF cells in comparison to the WT cells. To verified this, we also checked the accumulation of PRC2 that methylates H3K27 at this promoter and indeed binding of Suz12 (one of the subunits in PRC2) was reduced in CTCF pLoF cells (Fig. 3).

Fig. 3 **Altered chromatin state of *Hmgcs1* promoter in CTCF pLoF cells.** (**a**) A diagram of *Hmgcs1* promoter. SREBP motif is marked by red rectangle. Transcription start site is marked as TSS. Regions A-E were covered by five sets of primers used for ChIP-qPCR. (**b-d**) Mean Srebp2, H3K27me3 and Suz12 levels in the indicated parts of *Hmgcs1* promoter determined by ChIP-qPCR and presented as percentage of input in WT cells (white bars) and in CTCF pLoF

clones C13 (green bars) and C21 (orange bars) ± s.e. of six independent experiments. Normal rabbit IgG were used as a negative control to estimate non-specific binding of primary antibodies (grey bars). Statistical significance was calculated by the Student's *t*-test, *$P<0.05$, **$P<0.01$.

To prove that H3K27 methylation prevents SREBP2 binding to this promoter we repeated the ChIP analysis for SREBP2 in cells treated with a specific inhibitor for Ezh2, which is the catalytic subunit in PRC2 (GSK343). As shown in Fig. 4, GSK343 treatment led to increased SREBP2 binding to *Hmgcs1* promoter. Interestingly, GWAS query for the association of CTCF with human diseases revealed that the highest correlated disease with CTCF alterations was high density lipoprotein cholesterol (Sup. Fig. 3). Thus, suggesting CTCF as an unknown regulator of cholesterol homeostasis.

Fig. 4 **H3K27 methylation prevents Srebp2 binding to *Hmgcs1* promoter.** (**a**) Srebp2 binding to *Hmgcs1* promoter was determined in B16-F1 WT cells after 24 hours of GSK343 treatment (blue bar) and in control cells treated with DMSO (white bar) and expressed as a mean percentage of input ± s.e. of six independent experiments. Normal rabbit IgG were used as a control to estimate non-specific binding of target primary antibodies (grey bars). Statistical significance was calculated by the Student's *t*-test, *$P<0.05$. (**b**) A model for the transcriptional control of the cholesterol biosynthesis enzyme Hmgcs1 by CTCF that recruits the H3K27 methyltransferase PRC2 to elevate H3K27me3 at the promoter to prevent the binding of Srebp2.

In summary our results suggest that CTCF is crucial for correct cell migration due to its repressive effect on the cholesterol biosynthesis pathway. Reduced levels of CTCF led to impaired migration of the cells in parallel to increased RNA levels of cholesterol biosynthesis enzymes and elevated cellular cholesterol levels. Inhibition of cholesterol synthesis in CTCF pLoF cells alleviated their impaired migration rate. Cholesterol was shown to affect cell migration capabilities by changing membrane mechanical properties such as rigidity as well as by altering the function of membranal proteins and the activation of signal transduction pathways [49]. Detailed analysis of the promoter of *Hmgcs1* that encodes a cholesterol biosynthesis enzyme revealed that CTCF is not bound directly to it, but it is important for recruitment of PRC2 to this promoter to methylate H3K27. This methylation prevents the binding of SREBP2 to this promoter, and thus suppresses transcription of *Hmgcs1* and cholesterol biosynthesis (Fig. 4b). The exact mechanism by which CTCF recruits PRC2 to this promoter is still an open question; CTCF was shown to be able to bind PRC2 [50], thus CTCF may recruit PRC2 to a distant chromatin position from which it spreads towards the promoter. Another option is CTCF-dependent specific 3D contacts of the promoter that either generate a contact with a chromatin site to which PRC2 is already bound or/and prevent the generation of a contact with an active enhancer. Either way, CTCF appears as a new regulator

of cholesterol biosynthesis. Interestingly, recently CTCF was also found to regulate metabolic switch related to carbohydrates and protein metabolism in the morula-to-blastocyst transition during development [51]. Thus, CTCF emerges as a key player in metabolic regulation.

Acknowledgments

We thank Ariel University for funding.

Author contributions

Conceptualization, L.S.K., N.L., M.S.D. and G.G.; Methodology, L.S.K. and N.L.; Investigation, L.S.K. and N.L.; Formal Analysis - L.S.K., N.L. and M.S.D.; Writing - Original Draft, G.G.; Writing - Review & Editing, L.S.K., N.L., M.S.D. and G.G., Funding Acquisition, M.S.D. and G.G.

Declaration of interests

The authors declare no competing interests.

Methods

Cell culture

Mouse melanoma cells B16-F1 were purchased from the ATCC (CRL-6323, ATCC). Parental and CTCF pLoF clones C13 and C21 [46] were grown in DMEM (01-056-1A, Biological industries) supplemented with 10% Fetal Bovine Serum (FBS, 04-007-1A, Biological Industries, Israel), 0.5% penicillin-streptomycin antibiotic mix (03-031-1B, Biological Industries), and 1% L-glutamine

(03-020-1B, Biological Industries, Israel) at 37°C, 7% CO_2. Inhibitors used were and 3 µM of GSK343 (SML0766, Sigma-Aldrich, Rehovot, Israel) for 24 hours.

Motility assays

For the wound-healing assay, cells plated in fibronectin-coated 12-well plates were grown to confluence. Then the cells were scratched with the edge of a 10 µl tip, washed three times with DMEM and incubated in DMEM supplemented with 0.5% FCS. When indicated 5µM of Fatostatin (13562, Cayman chemical, Ann Arbor, Michigan, USA) was added 24 hours prior to the migration assay and was kept during the assay. Images of the same fields were collected with an Olympus SC100 camera mounted on an Olympus CKX41 microscope at two time points: immediately after the scratch and eight hours after the scratch. The area covered by the migrating cells was calculated using ImageJ/Fiji by comparison of the same fields between 0 hours and 8 hours. In each experiment four different fields were analyzed for each treatment (two in one well and another two in a second well).

The Transwell® assay was carried out in plate filters with pores diameter of 5 µm (3421, Corning Incorporated, Corning, NY, USA) as previously described [11]. Where indicated Fatostatin was added to the medium as in the wound healing assay. For counting purposes, the cells nuclei were stained with Hoechst reagent.

Transcriptome analysis

Total RNA was purified by the NucleoZOL kit (740404.200, MACHEREY-NAGEL, Duren, Germany) according to the manufacturer's instructions. For NGS sequencing by Cel-Seq2 [52] purified RNA from four repetitions for each cell type (WT, C13 and C21) was sent for poly A containing mRNA selection, library preparation and sequencing at the Technion Genome Center. For RT-qPCR 0.4µg of RNA was used to prepare cDNA with qPCRBIO cDNA Synthesis Kit

(PB30.11-10, PCR Biosystems, London, UK). qPCR was done with Fast SYBR Green Master Mix (4385612, Thermo Fisher Scientific, Vilnius, Lithuania). The various genes were normalized to the RNA levels of the ribosomal protein Rps29. Primers used: *Rps29*: forward: 5'-tcgttgggcgtctgaaggcaa-3', reverse: 5'- cggaagcactggcggcacat-3'; *Eva1a*: forward: 5'-ttgggaatggctctgctc-3', reverse: 5'- ggagatcctcatcaccaagg-3'; *Hs3st1*: forward: 5'-gtgtgaatttgctccaaagg-3', reverse: 5'- agcaaggctccctcttgatt-3'; *Pou3f1*: forward: 5'-aacaagtggctggaggagac-3', reverse: 5'- ggcttgggacacttgagaaa-3'; *Nov*: forward: 5'-tcgtctctgcatcgttcgg-3', reverse: 5'- catcactgcagaccccaca-3'; *Pgbd5*: forward: 5'-tacaaggtccagcccttcct-3', reverse: 5'- tttccgcttttcctcttcc-3'; *Hsd17b7*: forward: 5'-ggacgttgctcctacccata-3', reverse: 5'- cgcgtatttggtcagaggat-3'; *Hmgcs1*: Forward: 5'- cctggac-cgctgctattc-3', reverse: 5'- ttcaggaacatccgagctaga; *Lgals3*: forward: 5'- agacagtcagccttcccct-3', reverse: 5'- agttggctgatttcccggag-3'; *Egln3:* forward: 5'- caggttatgttcgccatgtg-3', reverse: 5'-ggctccacgtctgctacaa-3'; *Rgs16:* forward: 5'- cctgcgaggagttcaagaag-3', reverse: 5'- tggtagtgg-cagcttgtagg-3'; *Selenbp1:* forward: 5'- ctaacccacagaagccccg-3', reverse: 5'- gctggatcatctgagggcc-3'; *Col1a2*: forward: 5'- cccctggtcttactgggaac-3', reverse: 5'- caggtccttggaaaccttga-3'; *Mef2c*: for-ward: 5'- ggggactatggggagaaaaa-3', reverse: 5'- atctcacagtcgcacagcac-3'; *Dek:* forward: 5'- cag-tgacacaagggaagggt-3', reverse: 5'- ccactgaactgacccacgt-3'; *Ddit4:* forward: 5'- accag-ttcgctcacccttc-3', reverse: 5'- aaacgatcccagaggctagg-3'; *Slc2a1:* forward: 5'- ttgtt-gtagagcgagctgga-3', reverse: 5'- aaggccacaaagccaaagat-3'; *Egln1:* forward: 5'- cctggatcgaggg-caaaga-3', reverse: 5'- tagcctgttccgttgcctg-3'; *Frs3*: forward: 5'- gtgttcgagggcagaggac-3', reverse: 5'- cagagaccgtgtagccagc-3'; *Ctgf*: forward: 5'- cctgcgacccacacaagg-3', reverse: 5'- gctgcttt-ggaaggactcac-3'.

Cholesterol assay

The cholesterol assay was carried out according to the manufacturer protocol using the Cholesterol/Cholesterol Ester Assay Kit (ab65359, Abcam, Cambridge, UK) and normalized to total protein content as determined by Bradford Assay.

ChIP

Chromatin immunoprecipitation (ChIP) was performed as described previously [24] with some modifications: for each sample the lysate from 10^7 cells (total ~700µl) was divided into three fractions: sample (330µl), IgG control (330µl) and input (30µl). Samples were added to Magna ChIP™ Protein A+G Magnetic Beads (16-663, Sigma-Aldrich) precoated with specific antibody: 4µg of rabbit anti-Srebp2 (ab112046, Abcam, Cambridge, UK) or 4µg rabbit anti-H3K27me3 (07-449, Sigma-Aldrich) or 3µl of rabbit anti-Suz12 [3737, Cell Signaling, Danvers, MA, USA). IgG controls were added to magnetic beads precoated with 4µg of normal rabbit IgG (sc2027, Santa Cruz Biotechnology, Dallas, TX, USA). Quantitative PCR was carried with specific primers for the regions A-E in the promoter of *Hmgcs1*: (A) forward: 5'-tggaccaag tgccaaaataa-3', reverse: 5'-gaatgatgcagggcttgtct-3'; (B) forward: 5'-gccaaggtagtttaaattggaca-3', reverse: 5'-gcaaattgcatctgcatcac-3'; (C) forward: 5'- tacaggtgaagccgagctgt-3' or 5'- gaagccgagctgtggttg-3', reverse: 5'- actcctggggagagatggtc-3'; (D) forward: 5'-gaccttcaattggtcggaga-3', reverse: 5'-ggtaaggggtgggaacaaag-3'; (E) forward: 5'-agggaaaaccctagcgagtc-3', reverse: 5'- cggctac-ctctgccaatc-3'.

Western blot analysis

Whole cell lysate was prepared by incubation of cells with 2x Sample Buffer at 96°C for 10min and stored at -20°C. Antibodies used: mouse anti-Srebp1 (39939, Active Motif, Carlsbad, CA,

USA, dilution 1:1,000), rabbit anti-Srebp2 (112046, Abcam, Cambridge, UK, dilution 1:1,000), mouse anti-β-actin (A1978, Sigma-Aldrich, dilution 1:5,000).

Bioinformatic analysis

Sequencing reads from four replicates of RNA-seq samples for each pLoF clone and WT were aligned to the mouse genome (mm10) by Hisat2 program [53,54] after removal of adapter sequences and critical examination of quality controls using Trim Galore and FastQC respectively with default parameters. Next, the number of reads mapping each mouse gene (as annotated in Ensembl release GRCm38.100) was counted using the featureCounts program [55]. Differential expression analysis was performed using edgeR and Limma packages [56,57] from the Bioconductor framework [58]. Features with less than 1 read per million in four samples were removed. The remaining gene counts were normalized using the TMM method, followed by voom transformation [59]. Linear models as implemented by the Limma package were used to find differentially expressed genes between the two CTCF pLoF clones and the WT cells. Obtained lists with significantly changed genes ($p<0.05$, fold-change ≥2) were used in Ingenuity Pathway Analysis (IPA, Qiagen).

Supplementary data

Sup. Fig. 1 **Verification of RNA-seq based transcriptome profiling in CTCF pLoF cells by RT-qPCR.**
Relative RNA levels of up-regulated, down regulated and unchanged genes in the RNA-seq analysis were determined by RT-PCR. Graph bars represent means of ratios between RNA levels in CTCF pLoF cells

and WT cells ± s.e. of three independent experiments. Statistical significance was calculated by the Student's *t*-test, *P<0.05.

Sup. Fig. 2 **SREBPs in CTCF pLoF cells.** (**a**) Predicted upstream regulators by IPA analysis based on 364 differentially expressed genes in CTCF pLoF cells identified SREBPs (SREBFs) as possible activated transcription factors in these cells. (**b-c**) Protein levels of Srebp1 and Srebp2 in CTCF pLoF cells as determined by Western blot analyses.

Sup. Fig. 3 **Phenome-wide association studies for CTCF.** Association between CTCF variants and phenotypes across 4756 GWASs by GWAS-ATLAS (https://atlas.ctglab.nl/)

References

1. Blackley, D.G., Cooper, J.H., Pokorska, P., and Ratheesh, A. (2021). Mechanics of developmental migration. Seminars in Cell & Developmental Biology *120*, 66–74. 10.1016/j.semcdb.2021.07.002.

2. Sapir, T., Sela-Donenfeld, D., Karlinski, M., and Reiner, O. (2022). Brain Organization and Human Diseases. Cells *11*, 1642. 10.3390/cells11101642.

3. Delgado, M., and Lennon-Duménil, A.-M. (2022). How cell migration helps immune sentinels. Front. Cell Dev. Biol. *10*, 932472. 10.3389/fcell.2022.932472.

4. Novikov, N.M., Zolotaryova, S.Y., Gautreau, A.M., and Denisov, E.V. (2021). Mutational drivers of cancer cell migration and invasion. Br J Cancer *124*, 102–114. 10.1038/s41416-020-01149-0.

5. Wolf, K., Wu, Y.I., Liu, Y., Geiger, J., Tam, E., Overall, C., Stack, M.S., and Friedl, P. (2007). Multi-step pericellular proteolysis controls the transition from individual to collective cancer cell invasion. Nat Cell Biol *9*, 893–904. 10.1038/ncb1616.

6. Wolf, K., te Lindert, M., Krause, M., Alexander, S., te Riet, J., Willis, A.L., Hoffman, R.M., Figdor, C.G., Weiss, S.J., and Friedl, P. (2013). Physical limits of cell migration: Control by ECM space and nuclear deformation and tuning by proteolysis and traction force. The Journal of Cell Biology *201*, 1069–1084. 10.1083/jcb.201210152.

7. Friedl, P., and Wolf, K. (2009). Proteolytic interstitial cell migration: a five-step process. Cancer Metastasis Rev. *28*, 129–135. 10.1007/s10555-008-9174-3.

8. McGregor, A.L., Hsia, C.-R., and Lammerding, J. (2016). Squish and squeeze—the nucleus as a physical barrier during migration in confined environments. Current Opinion in Cell Biology *40*, 32–40. 10.1016/j.ceb.2016.01.011.

9. Lele, T.P., Dickinson, R.B., and Gundersen, G.G. (2018). Mechanical principles of nuclear shaping and positioning. J. Cell Biol. *217*, 3330–3342. 10.1083/jcb.201804052.

10. Yamada, K.M., and Sixt, M. (2019). Mechanisms of 3D cell migration. Nat Rev Mol Cell Biol *20*, 738–752. 10.1038/s41580-019-0172-9.

11. Gerlitz, G., Livnat, I., Ziv, C., Yarden, O., Bustin, M., and Reiner, O. (2007). Migration cues induce chromatin alterations. Traffic *8*, 1521–1529. 10.1111/j.1600-0854.2007.00638.x.

12. Gerlitz, G., and Bustin, M. (2010). Efficient cell migration requires global chromatin condensation. J. Cell. Sci. *123*, 2207–2217. 10.1242/jcs.058271.

13. Tanaka, H., Nishioka, Y., Yokoyama, Y., Higashiyama, S., Matsuura, N., Matsuura, S., and Hieda, M. (2012). Nuclear envelope-localized EGF family protein amphiregulin activates breast cancer cell migration in an EGF-like domain independent manner. Biochem Biophys Res Commun *420*, 721–726. 10.1016/j.bbrc.2012.03.045.

14. Zhang, X., Cook, P.C., Zindy, E., Williams, C.J., Jowitt, T.A., Streuli, C.H., MacDonald, A.S., and Redondo-Muñoz, J. (2016). Integrin α4β1 controls G9a activity that regulates epigenetic changes and nuclear

properties required for lymphocyte migration. Nucleic Acids Research *44*, 3031–3044. 10.1093/nar/gkv1348.

15. Zhang, B., Luo, Q., Chen, Z., Shi, Y., Ju, Y., Yang, L., and Song, G. (2016). Increased nuclear stiffness via FAK-ERK1/2 signaling is necessary for synthetic mechano-growth factor E peptide-induced tenocyte migration. Scientific Reports *6*, 18809. 10.1038/srep18809.

16. Hsia, C.-R., McAllister, J., Hasan, O., Judd, J., Lee, S., Agrawal, R., Chang, C.-Y., Soloway, P., and Lammerding, J. (2022). Confined migration induces heterochromatin formation and alters chromatin accessibility. iScience *25*, 104978. 10.1016/j.isci.2022.104978.

17. Maizels, Y., Elbaz, A., Hernandez-Vicens, R., Sandrusy, O., Rosenberg, A., and Gerlitz, G. (2017). Increased chromatin plasticity supports enhanced metastatic potential of mouse melanoma cells. Exp. Cell Res. *357*, 282–290. 10.1016/j.yexcr.2017.05.025.

18. Lu, Z., Zou, J., Li, S., Topper, M.J., Tao, Y., Zhang, H., Jiao, X., Xie, W., Kong, X., Vaz, M., et al. (2020). Epigenetic therapy inhibits metastases by disrupting premetastatic niches. Nature *579*, 284–290. 10.1038/s41586-020-2054-x.

19. Gerlitz, G. (2020). The Emerging Roles of Heterochromatin in Cell Migration. Front. Cell Dev. Biol. *8*, 394. 10.3389/fcell.2020.00394.

20. Madrazo, E., González-Novo, R., Ortiz-Placín, C., García de Lacoba, M., González-Murillo, Á., Ramírez, M., and Redondo-Muñoz, J. (2022). Fast H3K9 methylation promoted by CXCL12 contributes to nuclear changes and invasiveness of T-acute lymphoblastic leukemia cells. Oncogene *41*, 1324–1336. 10.1038/s41388-021-02168-8.

21. Furusawa, T., Rochman, M., Taher, L., Dimitriadis, E.K., Nagashima, K., Anderson, S., and Bustin, M. (2015). Chromatin decompaction by the nucleosomal binding protein HMGN5 impairs nuclear sturdiness. Nat Commun *6*, 6138. 10.1038/ncomms7138.

22. Strom, A.R., Biggs, R.J., Banigan, E.J., Wang, X., Chiu, K., Herman, C., Collado, J., Yue, F., Ritland Politz, J.C., Tait, L.J., et al. (2021). HP1α is a chromatin crosslinker that controls nuclear and mitotic chromosome mechanics. eLife *10*, e63972. 10.7554/eLife.63972.

23. Krause, M., Yang, F.W., te Lindert, M., Isermann, P., Schepens, J., Maas, R.J.A., Venkataraman, C., Lammerding, J., Madzvamuse, A., Hendriks, W., et al. (2019). Cell migration through three-dimensional confining pores: speed accelerations by deformation and recoil of the nucleus. Phil. Trans. R. Soc. B *374*, 20180225. 10.1098/rstb.2018.0225.

24. Segal, T., Salmon-Divon, M., and Gerlitz, G. (2018). The Heterochromatin Landscape in Migrating Cells and the Importance of H3K27me3 for Associated Transcriptome Alterations. Cells *7*, 205. 10.3390/cells7110205.

25. Braccioli, L., and de Wit, E. (2019). CTCF: a Swiss-army knife for genome organization and transcription regulation. Essays in Biochemistry *63*, 157–165. 10.1042/EBC20180069.

26. Wu, Q., Liu, P., and Wang, L. (2020). Many facades of CTCF unified by its coding for three-dimensional genome architecture. Journal of Genetics and Genomics *47*, 407–424. 10.1016/j.jgg.2020.06.008.

27. Dehingia, B., Milewska, M., Janowski, M., and Pękowska, A. (2022). CTCF shapes chromatin structure and gene expression in health and disease. EMBO Reports *23*. 10.15252/embr.202255146.

28. Lazniewski, M., Dawson, W.K., Rusek, A.M., and Plewczynski, D. (2019). One protein to rule them all: The role of CCCTC-binding factor in shaping human genome in health and disease. Semin Cell Dev Biol *90*, 114–127. 10.1016/j.semcdb.2018.08.003.

29. Misteli, T. (2020). The Self-Organizing Genome: Principles of Genome Architecture and Function. Cell *183*, 28–45. 10.1016/j.cell.2020.09.014.

30. Nora, E.P., Goloborodko, A., Valton, A.-L., Gibcus, J.H., Uebersohn, A., Abdennur, N., Dekker, J., Mirny, L.A., and Bruneau, B.G. (2017). Targeted Degradation of CTCF Decouples Local Insulation of Chromosome Domains from Genomic Compartmentalization. Cell *169*, 930-944.e22. 10.1016/j.cell.2017.05.004.

31. Wutz, G., Várnai, C., Nagasaka, K., Cisneros, D.A., Stocsits, R.R., Tang, W., Schoenfelder, S., Jessberger, G., Muhar, M., Hossain, M.J., et al. (2017). Topologically associating domains and chromatin loops depend on cohesin and are regulated by CTCF, WAPL, and PDS5 proteins. EMBO J *36*, 3573–3599. 10.15252/embj.201798004.

32. Sun, X., Zhang, J., and Cao, C. (2022). CTCF and Its Partners: Shaper of 3D Genome during Development. Genes *13*, 1383. 10.3390/genes13081383.

33. Ren, G., Jin, W., Cui, K., Rodriguez, J., Hu, G., Zhang, Z., Larson, D.R., and Zhao, K. (2017). CTCF-Mediated Enhancer-Promoter Interaction Is a Critical Regulator of Cell-to-Cell Variation of Gene Expression. Molecular Cell *67*, 1049-1058.e6. 10.1016/j.molcel.2017.08.026.

34. Kubo, N., Ishii, H., Xiong, X., Bianco, S., Meitinger, F., Hu, R., Hocker, J.D., Conte, M., Gorkin, D., Yu, M., et al. (2021). Promoter-proximal CTCF binding promotes distal enhancer-dependent gene activation. Nat Struct Mol Biol *28*, 152–161. 10.1038/s41594-020-00539-5.

35. Wu, J., Li, P.-C., Pang, J.-Y., Liu, G.-Y., Xie, X.-M., Li, J.-Y., Yin, Y.-C., Han, J.-H., Guo, X.-Z., and Qiu, L. (2017). CCCTC-binding factor inhibits breast cancer cell proliferation and metastasis via inactivation of the nuclear factor-kappaB pathway. Oncotarget *8*, 93516–93529. 10.18632/oncotarget.18977.

36. Lebeau, B., Jangal, M., Zhao, T., Wong, C.K., Wong, N., Cañedo, E.C., Hébert, S., Aguilar-Mahecha, A., Chabot, C., Buchanan, M., et al. (2022). 3D chromatin remodeling potentiates transcriptional programs driving cell invasion. Proc Natl Acad Sci U S A *119*, e2203452119. 10.1073/pnas.2203452119.

37. Wang, L., Deng, S.X., and Lu, L. (2012). Role of CTCF in EGF-Induced Migration of Immortalized Human Corneal Epithelial Cells. Invest. Ophthalmol. Vis. Sci. *53*, 946. 10.1167/iovs.11-8747.

38. Wang, L., Wu, X., Shi, T., and Lu, L. (2013). Epidermal Growth Factor (EGF)-induced Corneal Epithelial Wound Healing through Nuclear Factor κB Subtype-regulated CCCTC Binding Factor (CTCF) Activation. Journal of Biological Chemistry *288*, 24363–24371. 10.1074/jbc.M113.458141.

39. Elbert, A., Vogt, D., Watson, A., Levy, M., Jiang, Y., Brûlé, E., Rowland, M.E., Rubenstein, J., and Bérubé, N.G. (2019). CTCF Governs the Identity and Migration of MGE-Derived Cortical Interneurons. J. Neurosci. *39*, 177–192. 10.1523/JNEUROSCI.3496-17.2018.

40. Kim, T.-G., Kim, M., Lee, J.-J., Kim, S.H., Je, J.H., Lee, Y., Song, M.-J., Choi, Y., Chung, Y.W., Park, C.G., et al. (2015). CCCTC-binding factor controls the homeostatic maintenance and migration of Langerhans cells. J Allergy Clin Immunol *136*, 713–724. 10.1016/j.jaci.2015.03.033.

41. Shan, Z., Li, Y., Yu, S., Wu, J., Zhang, C., Ma, Y., Zhuang, G., Wang, J., Gao, Z., and Liu, D. (2019). CTCF regulates the FoxO signaling pathway to affect the progression of prostate cancer. J Cell Mol Med *23*, 3130–3139. 10.1111/jcmm.14138.

42. Xing, Z., Li, S., Liu, Z., Zhang, C., and Bai, Z. (2020). CTCF-induced upregulation of HOXA11-AS facilitates cell proliferation and migration by targeting miR-518b/ACTN4 axis in prostate cancer. Prostate *80*, 388–398. 10.1002/pros.23953.

43. Zhao, L., Yang, Y., Yin, S., Yang, T., Luo, J., Xie, R., Long, H., Jiang, L., and Zhu, B. (2017). CTCF promotes epithelial ovarian cancer metastasis by broadly controlling the expression of metastasis-associated genes. Oncotarget *8*, 62217–62230. 10.18632/oncotarget.19216.

44. Liu, C., Deng, L., Lin, J., Zhang, J., Huang, S., Zhao, J., Jin, P., Xu, P., Ni, P., Xu, D., et al. (2021). Zinc Finger Protein CTCF Regulates Extracellular Matrix (ECM)-Related Gene Expression Associated With the Wnt Signaling Pathway in Gastric Cancer. Front. Oncol. *10*, 625633. 10.3389/fonc.2020.625633.

45. Wen, X., Wang, H., Chai, P., Fan, J., Zhang, X., Ding, T., Jia, R., Ge, S., Zhang, H., and Fan, X. (2020). An Artificial CTCF Peptide Triggers Efficient Therapeutic Efficacy in Ocular Melanoma. Mol Ther Oncolytics *18*, 317–325. 10.1016/j.omto.2020.07.004.

46. Kaczmarczyk, L.S., Levi, N., Segal, T., Salmon-Divon, M., and Gerlitz, G. (2022). CTCF supports preferentially short lamina-associated domains. Chromosome Res *30*, 123–136. 10.1007/s10577-022-09686-5.

47. Kamisuki, S., Mao, Q., Abu-Elheiga, L., Gu, Z., Kugimiya, A., Kwon, Y., Shinohara, T., Kawazoe, Y., Sato, S., Asakura, K., et al. (2009). A Small Molecule That Blocks Fat Synthesis By Inhibiting the Activation of SREBP. Chemistry & Biology *16*, 882–892. 10.1016/j.chembiol.2009.07.007.

48. DeBose-Boyd, R.A., and Ye, J. (2018). SREBPs in Lipid Metabolism, Insulin Signaling, and Beyond. Trends in Biochemical Sciences *43*, 358–368. 10.1016/j.tibs.2018.01.005.

49. Maja, M., and Tyteca, D. (2022). Alteration of cholesterol distribution at the plasma membrane of cancer cells: From evidence to pathophysiological implication and promising therapy strategy. Front. Physiol. *13*, 999883. 10.3389/fphys.2022.999883.

50. Yan, H., Tang, G., Wang, H., Hao, L., He, T., Sun, X., Ting, A.H., Deng, A., and Sun, S. (2016). DNA methylation reactivates GAD1 expression in cancer by preventing CTCF-mediated polycomb repressive complex 2 recruitment. Oncogene *35*, 3995–4008. 10.1038/onc.2015.423.

51. Andreu, M.J., Alvarez-Franco, A., Portela, M., Gimenez-Llorente, D., Cuadrado, A., Badia-Careaga, C., Tiana, M., Losada, A., and Manzanares, M. (2022). Establishment of 3D chromatin structure after

fertilization and the metabolic switch at the morula-to-blastocyst transition require CTCF. Cell Reports *41*, 111501. 10.1016/j.celrep.2022.111501.

52. Hashimshony, T., Senderovich, N., Avital, G., Klochendler, A., de Leeuw, Y., Anavy, L., Gennert, D., Li, S., Livak, K.J., Rozenblatt-Rosen, O., et al. (2016). CEL-Seq2: sensitive highly-multiplexed single-cell RNA-Seq. Genome Biol *17*, 77. 10.1186/s13059-016-0938-8.

53. Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low memory requirements. Nat Methods *12*, 357–360. 10.1038/nmeth.3317.

54. Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907–915. 10.1038/s41587-019-0201-4.

55. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930. 10.1093/bioinformatics/btt656.

56. Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR : a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–140. 10.1093/bioinformatics/btp616.

57. Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research *43*, e47–e47. 10.1093/nar/gkv007.

58. Reimers, M., and Carey, V.J. (2006). [8] Bioconductor: An Open Source Framework for Bioinformatics and Computational Biology. In Methods in Enzymology (Elsevier), pp. 119–134. 10.1016/S0076-6879(06)11008-3.

59. Law, C.W., Chen, Y., Shi, W., and Smyth, G.K. (2014). voom: precision weights unlock linear model analysis tools for RNA-seq read counts. Genome Biol *15*, R29. 10.1186/gb-2014-15-2-r29.