Cortical interneurons: fit for function and fit to function? Evidence from development and evolution

Joram Keijser & Henning Sprekeler

April 7, 2023

Abstract

Cortical inhibitory interneurons form a broad spectrum of subtypes. This diversity suggests a division of labour, in which each cell type supports a distinct function. In the present era of optimisation-based algorithms, it is tempting to speculate that these functions were the evolutionary or developmental driving force for the spectrum of interneurons we see in the mature mammalian brain. In this study, we evaluated this hypothesis using the two most common interneuron types, parvalbumin (PV) and somatostatin (SST) expressing cells, as examples. PV and SST interneurons control the activity in the cell bodies and the apical dendrites of excitatory pyramidal cells, respectively, due to a combination of anatomical and synaptic properties. But was this compartment-specific inhibition indeed the function for which PV and SST cells originally evolved? Does the compartmental structure of pyramidal cells shape the diversification of PV and SST interneurons over development? To address these questions, we reviewed and reanalysed publicly available data on the development and evolution of PV and SST interneurons on one hand, and pyramidal cell morphology on the other. These data speak against the idea that the compartment structure of pyramidal cells drove the diversification into PV and SST interneurons. In particular, pyramidal cells mature late, while interneurons are likely committed to a particular fate (PV vs. SST) during early development. Moreover, comparative anatomy and single cell RNA-sequencing data indicate that PV and SST cells, but not the compartment structure of pyramidal cells, existed in the last common ancestor of mammals and reptiles. Specifically, turtle and songbird SST cells also express the *Elfn1* and *Cbln4* genes that are thought to play a role in compartment-specific inhibition in mammals. PV and SST cells therefore evolved and developed the properties that allow them to provide compartment-specific inhibition before there was selective pressure for this function. This suggest that interneuron diversity originally resulted from a different evolutionary driving force and was only later co-opted for the compartment-specific inhibition it seems to serve in mammals today. Future experiments could further test this idea using our computational reconstruction of ancestral Elfn1 protein sequences.

1 Introduction

Cortical inhibitory interneurons are a highly diverse group, differing in their morphology, connectivity, and electrophysiology [1]. Decades of experimental and theoretical work have suggested a role for interneurons in many functions [1, 2, 3], including the regulation of neural activity [4, 5], control of synaptic plasticity [6, 7], increasing temporal precision [8, 9], predictive coding [10, 11], and gain modulation [12, 13]. Many of these functions come down to the control of excitation.

Why would the control of excitation require a diversity of interneurons? A key reason could lie in the complexity of excitatory cells [14, 15]. Pyramidal cells (PCs) consist of several cellular compartments that have different physiological properties (e.g. sodium vs. calcium spikes [16]), receive different inputs (e.g., top-down vs. bottom up [17, 18], although see [19]) and follow distinct synaptic plasticity rules [20, 21, 22]. The control of different pyramidal cell compartments might therefore require inhibition from designated types of interneurons. Indeed, the two most common interneuron types—parvalbumin (PV)- and somatostatin (SST)-expressing cells— are classically distinguished by their connectivity with pyramidal cells: whereas

PV-expressing basket cells mainly target the somata of PCs, SST-expressing Martinotti cells mainly target their apical dendrites [1]. The cellular and synaptic properties of these interneurons also seem adapted to this purpose. SST interneurons receive facilitating synapses from PCs [23, 24], rendering them sensitive to bursts of action potentials [25, 26, 27] triggered by plateau potentials in the apical dendrite of PCs [16, 28]. Indeed, SST interneurons control dendritic excitability and bursting of PCs [26, 29, 30]. PV interneurons, on the other hand, receive depressing synapses [23, 31], rendering them less sensitive to these signals [32]. The presynaptic dynamics of PV and SST interneurons therefore seem particularly well-matched to the physiology of pyramidal cells, although both types also inhibit non-pyramidal cells and other interneurons (see e.g. [33, 34]). These and similar observations have led to the view that interneuron diversity can be understood from a functional perspective, in which the morphology and synaptic and cellular properties of different interneurons are fit to specific functions (Fig. 1a, [2, 14]). Consistent with this idea that interneurons are adapted to control different pyramidal cell compartments, we recently showed that properties (connectivity and short-term plasticity) of PV and SST interneurons emerge when optimising a network model for compartment-specific inhibition (Fig. 1b, [15]).

Figure 1: **Do development and evolution fit interneurons to function?** **(a)** The connectivity and short-term plasticity (STP) of PV and SST positive interneurons seem adapted to the morphology and electrophysiology of pyramidal cells, as highlighted by an optimization-based model **(b)**. In this model, optimising interneuron parameters to provide compartment- (soma/dendrite) specific inhibition causes interneurons to diversify into two groups that resembled PV and SST interneurons in their connectivity and short-term plasticity [15]. **(c)** The existence of PV and SST subtypes might therefore result from an developmental or evolutionary tuning of interneurons based on pyramidal properties. **(d)** This predicts that immature (or ancestral) circuits contain bursting pyramidal neurons and undiversified interneurons. Development (or evolution) then diversifies the interneurons into PV and SST subtypes. PC activity and short-term plasticity simulated with models from refs. [35, 36] and [37], respectively. Animal silhouettes from `https://beta.phylopic.org/`

The specialisation of PV and SST interneurons to pyramidal soma and dendrites, respectively, makes it tempting to speculate that the diversification of these interneuron subtypes was driven by pyramidal cell properties, either during evolution or during development (Fig. 1c). This hypothesis predicts a specific temporal order: During evolution or development, the compartmentalization of pyramidal cells should predate interneuron diversification (Fig. 1d).

Here, we evaluate this idea, with a focus on PC and interneuron properties that seem particularly well-adapted to each other: the active dendrites of pyramidal cells, and the connectivity and short-term plasticity

of interneurons. Reviewing and re-analysing recent evolutionary and developmental data, we reconstruct the developmental and evolutionary history of these three properties. We find no support for the idea that interneurons develop or evolved to control preexisting compartments of pyramidal cells. Instead, the central properties of PV and SST interneurons that led to this idea emerge before the PC properties they seem adapted to, in both development and evolution. Rather than pyramidal physiology driving interneuron diversification, this suggests a model in which existing interneuron properties enabled new pyramidal cell functions.

2 Developmental trajectory of compartment-specific inhibition

We first discuss the developmental trajectory of pyramidal cells and PV and SST interneurons in the mammalian cortex, to assess whether the diversification of PV and SST interneurons during development is driven by pyramidal cell properties. We mostly consider data from rodents, but many of the findings seem to be conserved among mammals [38, 39, 40, 41].

In contrast to pyramidal cells, interneurons are not born in the developing cortex, but subcortically ([42, 43], Box 1). It is only upon migrating to the cortex that interneurons acquire their mature morphology and physiology. The long period between interneuron birth and maturation has led to different models of interneuron development [2, 44]. One model attributes the late maturation of interneurons to a late specification of their cellular identity, possibly based on external cues within the circuit they embed themselves in [2, 44]. Alternatively, the late emergence of characteristic features could be due to the slow unfolding of a predetermined genetic program that happens independently of the surrounding circuit[45].

The malleability of interneuron properties during development is therefore currently an open question: Which properties are adapted to the surrounding circuit, and which are predetermined? Whatever properties are adapted, cellular identity (e.g., PV vs. SST) is probably not one of them [44, 45]. Interneuron types— at least on a coarse level—are determined by their time and place of birth. Future PV and SST interneurons, for example, are preferentially generated within different parts of the same embryonic structure ([46, 45],Box 1).

Recent data suggests that not just interneuron types (e.g., PV vs. SST), but also interneuron *sub*types (e.g., SST Martinotti vs. SST non-Martinotti) are specified early in development. Lim et al. [45] showed that Martinotti and non-Martinotti cells migrate to the developing cortex via different routes (Box 1). In addition, a developing interneuron's transcriptional profile can be used to predict its future fate [47, 48, 40, 49].

Box 1 Birth and migration of cortical interneurons

Cortical GABAergic interneurons are born in a transient region of the developing brain known as the ganglionic eminence [42, 45, 44], from where they tangentially migrate to the cortex [50]. The ganglionic eminence can be divided into multiple subregions patterned by unique combinations of transcription factors [43, 38, 39, 51], that activate distinct genetic programs. Since each genetic program corresponds to a different cell type, the majority of the cells born in the medial ganglionic eminence (MGE) will become PV and SST interneurons, whereas the caudal ganglionic eminence (CGE) generates, among others, vasoactive intestinal peptide (VIP)-expressing interneurons [52, 53, 54, 55, 56]). A key example for a patterning transcription factor that shapes interneuron identity is *Nkx2-1*, which is expressed within the MGE but not CGE [57, 55]. *Nkx2-1* knockout leads MGE-derived interneurons to adopt the fate of CGE-derived interneurons [58]. Molecular gradients have also been shown to contribute to interneuron diversity within the same eminence: the dorsal-caudal and rostral-ventral MGE preferentially generate SST and PV neurons, respectively [59, 60, 61, 62, 51, 63].

After birth, interneurons migrate to the developing cortex via two different routes: The superficial marginal zone (the MZ, which will develop into cortical layer 1) and the deeper subventricular zone (SVZ). These different migration routes are used by distinct layer 2-3 (L2-3) SST subtypes [45]. Whereas L2-3 SST Martinotti cells migrate via the marginal zone from where they descend to their future location , non-Martinotti cells migrate via the subventricular zone [64]. This indicates that SST subtypes (at least in L2-3) are predetermined before their arrival in cortex, possibly even before they "choose" one migratory route over the other. Future L2-3 Martinotti cells forced to migrate via the wrong route (the SVZ) still become Martinotti cells in terms of their transcriptional profile and electrophysiology, but they lack a fully developed layer 1 axon [64]. This suggests that developing L2/3 Martinotti cells cannot send their developing axon from deeper to upper layers, but have to leave it there while their cell body descends. Translaminar axons of other neurons such as a less studied PV subtype [64], and cerebellar granule cells [65] are established via a similar mechanism, suggesting it might be the only reliable way for neurons to develop translaminar projections.

Although interneurons are therefore likely hardwired to become a certain subtype, it is still possible that interneuron properties such as short-term plasticity or connectivity are subject to activity-dependent fine-tuning. For example, the development of short-term facilitation or a layer 1 axon of SST Martinotti cells might emerge in dependence on pyramidal neuron bursting. In this case, bursting should develop ahead of these SST features.

When do developing pyramidal cells first show dendrite-dependent bursting? Their electrophysiology matures relatively late: dendritic plateau potentials emerge only in the third postnatal week [66, 67]. This is consistent with the late maturation of their dendritic morphology. PCs develop their intricate apical arborization and tuft dendrites after the second postnatal week [66, 68]. For example, the tuft length increases almost twofold during the third postnatal week [68], and dendritic spikes fail to reach the soma on postnatal day 14 and 28 [66].

When does short-term facilitation (STF) of PC→SST synapses arise during development? Could its development be driven by bursting in pyramidal cells? Some of the early experiments showed such STF in rat cortex during the third postnatal week [69, 24, 23]. Evidence for an even earlier presence of STF in these synapses comes from molecular studies. The Gosh laboratory has shown that the short-term facilitation in hippocampal [70] and cortical [71] PC→SST synapses is due to the transmembrane protein Elfn1, which is expressed by SST mouse and human interneurons (Box 2, Fig. 2 In these experiments, STF was measured in the second postnatal week, and the expression of Elfn1 was detected already one week after birth [72, 73], providing an early molecular signature of short-term facilitation in SST cells. Short-term facilitation in PC→SST synapses is therefore present before dendrite-dependent bursting in PCs.

Figure 2: ***Elfn1* expression correlates with short-term facilitation in mammals.** **(a)** UMAP [77] plot of mouse interneurons coloured by subclass (left) and *Elfn1* expression (right). The two interneuron types — SST and VIP interneurons — known to receive facilitating synapses both express *Elfn1*. **(b)** Violin plot of *Elfn1* expression by subclass. CP10K: counts per 10 thousand. **(c,d)** As **(a,b)**, but for human interneurons. Data from from refs. [74] (a,b), and [75] (c,d).

What about the second difference between PV and SST neurons, their compartment-specific output synapses? SST and PV cells form compartment-specific synapses in visual cortical organotypic cultures that lack external inputs [78]. This strongly suggests a role for genetic encoding rather than experience-dependent activity. Indeed, recent work identified important molecular players in the establishment of compartment-specific synapses [73]. Several genes are involved in the formation of compartment-specific synapses. For example, suppressing *Cbln4* in SST interneurons decreased inhibition onto PC dendrites. An over-expression of the same gene in PV interneurons, on the other hand, increased inhibition onto PC dendrites [73]. Other genes contribute to somatic inhibition in a seemingly analogous way [73]. Both loss and gain of function were shown around P14. Similarly, somatic inhibition in CA1 abruptly emerges at the end of the second postnatal week [79]. It is therefore by the second postnatal week that PV and SST interneurons are committed as to where to direct their output synapses.

Intriguingly, *Cbln4* is only expressed in a subset of neurons (Fig. 3). Clustering revealed that these *Cbln4*+ neurons correspond to previously identified subtypes. The *Tac1* cluster labels non-Martinotti cells that target the dendrites of L4 cells [80, 81, 82], and the *Calb2* and *Etv1* clusters correspond to fanning-out Martinotti cells [80, 5], consistent with a role of *Cbln4* in establishing dendritic synapses. But only a subset of the *Myh8* cluster—corresponding to T-shaped Martinotti cells [5]— expressed *Cbln4*, suggesting diverse mechanisms for dendritic targeting.

An interneuron's cell type, the plasticity of their input synapses from PCs, and the PC compartments they target are therefore determined before interneurons are fully embedded within cortical circuits, and before pyramidal neurons develop their characteristic morphology and electrophysiology. This suggest that while PV and SST interneurons are fit for the function of compartment-specific inhibition of PCs, some of their characteristic properties are probably not developmentally driven by PC activity.

Figure 3: **Cbln4 is expressed in a subset of mammalian SST interneurons. (a)** UMAP plot of mouse and human interneurons, coloured by their expression of *Cbln4*, a gene that instructs synapse formation onto pyramidal dendrites in mice [73]. *Cbln4* is expressed in certain mouse and human interneuron subtypes, including SST cells. **(b)** UMAP of SST cells, clustered into subgroups. **(c)** *Cbln4* is expressed in clusters 0, 3, and 12, which also express marker genes *Tac1*, *Calb2* and *Etv1* **(d)**, respectively. **(e)** A subset of human Sst cells also express *Cbln4*. Data from from refs. [74] (a) and [75] (e).

Evolutionary trajectory of compartment-specific inhibition

On a much longer timescale than development, evolution also changes the properties of cell types. This raises the question whether the differentiation of PV and SST interneurons preceded the evolution of the compartmental complexity of pyramidal neurons.

If natural selection tuned PV and SST neurons to pyramidal cell properties, the brains of mammalian ancestors must have contained pyramidal cells with elaborate dendrites, while interneurons were still undifferentiated (Fig. 1c,d). This hypothesis cannot be tested directly since our mammalian ancestors are no longer alive, and their fossils provide no information regarding cell types. We therefore have to infer the evolutionary history of cell types by comparing data from modern-day species (Fig. 4a, [83, 84]). Although many cell type-specific properties such as short-term plasticity have not been measured in non-standard model organisms (see refs. [85, 86, 34] for recent exceptions), transcriptomic correlates can be studied using single cell RNA sequencing (scRNA-seq, [87]), offering a means for defining and comparing cell types across species [88, 84].

Interneuron conservation & principal neuron divergence

The first applications of scRNA-seq in neuroscience profiled cell types in mice [89, 90]. More recently, scRNA-seq was used to classify neuron types also in reptiles [91] and songbirds [92]. The evolutionary relationships of reptiles, birds and mammals suggest that a feature found in all three lineages predates their divergence, while a feature found exclusively in the mammalian lineage is, in fact, a mammalian invention. This idea enables inferring the evolutionary history of interneurons and pyramidal cells (Fig. 4a).

Let us first consider the general evolutionary trajectory of excitatory and inhibitory cell types. Tosches et al. [91] used scRNA-seq to analyse cells from the turtle and lizard forebrain and compare them with previously published mammalian data [90]. They found that reptilian inhibitory neurons cluster into groups that roughly correspond to mammalian interneuron types [91]. These results extend earlier findings that found similarities between turtle and mammalian interneurons based on marker genes and morphology [93, 94]. Colquitt et al. [92] recently made analogous observations regarding the similarity of songbird and mouse interneurons (Fig. 4b,c). The most parsimonious explanation of this sharing of interneuron types is that similar types already existed in a common ancestor of the three lineages, rather than convergent evolution in three lineages. This homology is likely due to shared developmental origins: the inhibitory interneurons of birds and reptiles are born within the conserved ganglionic eminences. The fact that interneurons of different lineages are homologous does not mean they are identical. For example, the correlation between mouse PV and SST cells and the best matching songbird clusters is 0.37 and 0.31, respectively (Fig. 4b). This is higher than the correlation between the best-matching glutamatergic types (0.19, see below), but lower than between some of the different cell types within the same species (mouse PV and SST cells: 0.58). Mammalian and non-mammalian interneurons, while homologous, therefore have likely undergone lineage-specific adaptations.

Figure 4: **Evolutionary conservation of GABAergic cell types** **(a)** Phylogenetic approach. **(b)** Pearson correlation between average RNA expression in clusters of songbird and mouse interneurons. Correlations between GABAergic neurons are typically larger. **(c)** UMAP plots of integrated gene expression data for GABAergic and glutamatergic neurons. GABAergic neurons first cluster by developmental origin (MGE vs. CGE, see Box 1) and then by species. Mouse data from ref. [74], songbird data and correlation analysis from ref. [92].

In contrast to inhibitory interneurons, excitatory neurons are probably not homologous between reptiles, songbirds, and mammals (Fig. 4b,c; [91, 92]). Excitatory cell types in different species are defined by different combinations of transcription factors. A clear example is given by the *Fezf2* and *Satb2* genes that specify subcortical [95] and callosal [96] projections, respectively, of mammalian pyramidal cells. Strikingly, these genes are mutually repressive in the mammalian neurons, but co-expressed in reptilian neurons [91, 97]. Comparing excitatory neurons in the songbird and the mammalian brain revealed an analogous pattern: Although excitatory neurons in the songbird forebrain express similar genes as their counterparts in mammalian neocortex, these genes are regulated by different transcription factors [92]. Instead, the transcription factors expressed by songbird glutamatergic neurons are similar to those in e.g. the mouse olfactory bulb and olfactory cortex. Since transcription factors specify cellular identity [98, 83] this suggests that excitatory neurons are not conserved across mammals, birds and reptiles [91, 92, 99].

Inhibitory cell types therefore seem more conserved than excitatory cell types, which appears broadly inconsistent with an evolutionary adaptation of interneurons to pyramidal cells. This is further confirmed when considering the evolutionary history of specific features of excitatory and inhibitory interneurons, in particular, elaborate dendrites and dendrite-dependent bursting and short-term plasticity.

Evolution of cell type-specific features

We are not aware of direct measurements of short-term facilitation in non-mammalian species and therefore aimed to infer its presence from the expression of *Elfn1* (Box 2). To this end, we re-analyzed publicly available gene expression data for reptilian and songbird interneuron types [91, 92]. We found that *Elfn1* is also expressed in the types corresponding to mammalian SST (and VIP) interneurons (Fig. 5), but not in the type corresponding to PV interneurons. This suggests that SST-like interneurons expressing *Elfn1* — and potentially faciliating glutamatergic input synapses — were already present in the last common ancestor of reptiles, songbirds and mammals. In terms of potential transcriptomic correlates of synaptic specificity, we find that *Cbln4* (Fig. 3) is expressed in certain subtypes of turtle SST neurons (Fig. 6a). Songbird SST neurons, on the other hand, do not express *Cbln4* (Fig. 6b). The expression of *Cbln4* by Sst interneurons therefore correlates with the presence of apical dendrites in pyramidal cells (Fig. 7, see next). The most parsimonious explanation is that *Cbln4* expression was lost in the songbird lineage. Alternatively, it could have evolved independently in the mammals and reptiles.

Figure 5: **Evolutionary conservation of *Elfn1* expression (a)** UMAP plot showing over-expression of *Elfn1* in SST-like and VIP-like interneurons in the turtle forebrain. Data from ref. [91].**(b)** Violin plots of *Elfn1* expression for each of the clusters. **(c,d)** As (a,b), but for zebra finch neurons. Data from ref. [92].

So not just interneuron subtypes, but also some of their specific properties seem evolutionarily conserved. In contrast, glutamatergic cell types in reptiles and birds show a very different dendritic morphology and physiology from their mammalian pyramidal counterparts. Turtle pyramidal cells have multiple apical dendrites, but no basal dendrites (Fig. 7, [100]). This clear morphological difference suggests that turtle pyramidal neurons are also electrophysiogically distinct. Larkum et al. [101] showed that, *in vitro*, turtle pyramidal neurons lack dendritic calcium spikes and dendrite-dependent bursting. Morphologically similar pyramidal cells in rodent piriform cortex also lack active dendrites ([102, 103], but see [104]). Interestingly, this is probably not due to an absence of calcium channels, but rather to the presence of A-type potassium channels [103]. Songbird excitatory cells have a stellate morphology, and differ therefore even more from mammalian pyramidal cells (Fig. 7, see e.g. [105, 106]).

Figure 6: ***Cbln4* expression in non-mammalian species**. *Cbln4* is expressed in certain subtypes of turtle SST neurons, but not in songbird SST neurons. Data from refs. [91, 92].

Figure 7: **Evolutionary divergence of projection neuron morphology.** Both turtle and mammalian projection neurons have a pyramidal morphology, but only mammalian pyramidal neurons have a single apical dendrite. Songbird projection neurons have a stellate, not pyramidal morphology. Turtle and mammalian neurons adapted from [101] (published under a Creative Commons License [107]). Songbird neuron adapted from [108] (published under a Creative Commons License [109])

The lack of dendrite-dependent bursting in reptiles and songbirds is consistent with comparative electro-physiology within the mammalian brain. Pyramidal neurons in the piriform cortex are homologous to certain types of glutamatergic turtle and songbird neurons [92], and also lack dendritic plateau potentials [102]. Pyramidal neurons with mammalian electrophysiological properties therefore evolved after interneurons differentiated into PV and SST cell types. (Fig. 8).

Ancestral Elfn1 reconstruction

The expression of *Elfn1* by zebra finch and turtle SST-like neurons suggests these cells—and therefore the ancestral SST-like cells— receive(d) facilitating inputs. But it is also possible that the ancestral Elfn1 protein had different functional properties. Previous work has used Elfn1 knockout [70, 110, 72] and targeted deletions [111] to discover functionally important domains of the mouse variant (Fig 9b). To determine the evolution of Elfn1 at the resolution of individual sites, we computationally reconstructed its ancestral state ([112, 113], see Methods), starting from the protein sequences of extant species (Fig. 9a). Alignment of the extant sequences revealed that on average across species 74.6% of the Elfn1 sites was identical to that of the mouse protein (Fig. 9c). Combining the sequence alignment with a probabilistic model of sequence evolution [114] and a phylogenetic species-tree allowed us to reconstruct the ancestral protein (Fig. 9d). The amount of conservation varied between protein domains and extant species: the zebra finch and turtle sequences were more similar to the ancestral sequence than the mammalian sequences (Fig. 9e). Future work could use the reconstructed sequences to determine the evolutionary history and the molecular mechanisms of short-term facilitation.

Figure 8: **Phylogenetic inference of interneuron and pyramidal evolution (a)** mice, humans, songbirds and turtles all have PV and SST interneurons. The most likely explanation for these similarities is that the interneuron types were already present in the last common ancestor of these lineages. **(b)** Only mammalian glutamateric neurons are known to exhibit dendritic plateau potentials that can elicit burst firing. Other lineages probably lack this trait. The most likely explanation is that dendritic bursting evolved only once, in the mammalian lineage.

Discussion

The suspicious match between the synaptic properties of PV and SST interneurons and the postsynaptic pyramidal cell compartments suggests that these interneuron properties could be the result of an adaptation to pyramidal cells. Here, we evaluated this idea of interneurons being 'fit to function' from an evolutionary and developmental perspective, and showed that the relevant interneuron properties predate those of pyramidal cells both during development and in evolutionary history.

Two lines of evidence indicate that the development of PV and SST interneurons is not induced by mature pyramidal cell activity. First, interneurons become committed to a particular cell type (e.g., PV or SST) before reaching the developing cortex. Interneuron fate therefore cannot be influenced by the activity of pyramidal cells. Second, at least some of the properties of PV and SST interneurons that strongly shape their control of pyramidal cells — short-term plasticity and output connectivity — emerge before the maturation of pyramidal cell morphology and dendritic activity (dendrite-dependent bursting). It should be noted that other interneuron properties clearly are influenced by pyramidal cell activity. Excitatory activity regulates both the survival of interneurons [116], and the formation of inhibitory synapses [117, 118]. Specific types of excitatory neurons determine the laminar allocation of interneurons [119, 120], and their activity can even change the intrinsic properties of mature interneurons [121]. Cell-extrinsic cues therefore play a role in the normal development of interneurons, but are unlikely to determine their identity and the properties we focused on here.

Analogous arguments suggest that the evolution of PV and SST interneurons also cannot be driven by the dendritic physiology of pyramidal cells. The lineages of birds, reptiles and mammals diverged over 300 million years ago, yet they all contain roughly similar interneuron types — evidence that these types were already present in a common ancestor of the three lineages. In contrast to interneurons, excitatory neurons are not conserved, and therefore probably evolved later. The second line of evolutionary evidence relates to two specific aspects of interneuron diversity: short-term plasticity and output connectivity. Recent scRNA-seq data [91, 92] show that reptilian and songbird SST interneurons express *Elfn1*, the gene that in mouse SST neurons is necessary and sufficient for short-term facilitation. Certain reptilian, but not songbird, SST subtypes also express *Cbln4* that plays a role in the synaptic specificity of mammalian SST cells [73].

These data suggest that ancestral interneurons already comprised PV- and SST-like cell types characterised by some of the genes for cell type-specific phenotypes in mammalian interneurons. It does not, however, imply that these phenotypes were actually present in ancestral cells. The expression of *Elfn1*, for example, is not sufficient for facilitating inputs, as shown in the case of VIP subtypes: Multipolar and bipo-

Figure 9: **Reconstruction of ancestral Elfn1 protein (a)** Species-tree showing the phylogenetic relationships of the species whose Elfn1 homologues were used to reconstruct the Elfn1 protein of the amniote ancestor. **(b)** Domain structure of mouse Elfn1 [115, 111]. LRR: leucine-rich repeat; CT: C-terminal domain; FN3: fibronectin type 3 domain; TM: transmembrane domain. **(c)** Per-site conservation across the tree shown in (a), computed as the fraction of extant species that share the mouse amino acid in a given site. Dashed lines correspond to gaps. Mean conservation: 0.746. **(d)** Posterior probability of ancestral protein. Gray: most likely (ML) sequence, red: 2nd most likely. Dashed line: cutoff for using the 2nd most likely base in "allAlt" sequence. Mean posterior: 0.986. **(e)** Multiple sequence alignment of protein domains shown in (a). Only the first 2 LRRs are shown for space reasons. Dots indicate identity to mouse site, dashes indicate gaps.

276 lar VIP neurons both express *Elfn1*, but only the multipolar subtype receives facilitating excitation [71]. It
277 will therefore be interesting to directly test the presence of PV- and SST-specific phenotypes in reptiles and
278 birds. If neither the reptile nor the songbird homologue of SST interneurons receives facilitating excitatory
279 inputs, *Elfn1* was likely reused for short-term facilitation in mammals. The emergence of short-term facilita-

tion in SST neurons would then be an adaptation to pyramidal bursting, co-opting pre-existing interneuron diversity for "pyramidal cell purposes". The anatomical connectivity of interneurons might similarly have been reused to control pyramidal cells. In the mammalian brain, PV and SST interneurons inhibit not just the somata and dendrites, respectively, of pyramidal cells but also of non-pyramidal cells. Ancestral PV and SST interneurons might therefore have specialised in compartment-specific inhibition, but not of pyramidal cells for which their presynaptic dynamics are so well-matched.

Although our results show that pyramidal cell bursting is unlikely the driver of the differentiation of PV and SST interneurons, this is not in conflict with the functional interpretation of these cell types. In fact, an evolution of active pyramidal cell dendrites before the presence of specialised interneurons would have resulted in aberrant excitation, as seen, e.g., in *Elfn1* mutants [110, 72]. This suggests an alternative picture, in which excitatory neurons can only evolve in a way that still allows the existing interneurons to regulate their activity. This still leaves open the question why interneuron diversity evolved in the first place, if it was not for compartment-specific inhibition. Although it is possible that the initial separation between PV and SST cell types was selectively neutral, this is unlikely given their evolutionary conservation. Instead, the existence of PV and SST cells presumably offers advantages to mammalian and non-mammalian brains alike. An important example of an conserved function could be the temporal coordination of inputs and outputs of pyramidal cells based on oscillations [122, 123].

Our findings have potential implications for the neuroscientific interpretation of optimisation-based models of neural networks, which have recently seen a renaissance [124, 125, 126, 127, 128]. Most of these models describe neural data at the relatively abstract level of dynamics and representations [129, 130]. Recently, such deep network models have also started to include circuit-level structure such as separate excitatory and inhibitory populations [131, 132], different neuronal timescales [133, 134], and short-term plasticity [135, 15]. Deep learning is therefore gradually making its way down from the level of dynamical systems to that of circuits, potentially revealing functional roles for circuit elements. Our findings highlight a challenge to achieving this goal: Multiple circuit-level features — such as the properties of interneuron and pyramidal cells — are interdependent. The function of one feature might depend on that of another and vice versa, raising the question which features should be optimized (e.g., interneurons), and which should be assumed as pre-exising constraints or opportunities (e.g., nonlinear PC dendrites). In other words, optimization-based models face the challenge of modelling processes such as co-evolution. Merging the functional and evolutionary/developmental perspectives will therefore be an important challenge for future work.

Acknowledgements

We thank Simon J.B. Butt, Loreen Hertäg and members of the Sprekeler Lab for comments on the manuscript.

3   Methods

Code was written in Python (version (v) 3.10.8 [136]) and R (v4.2.1 [137]), based on practices outlined in the Good Research Codebook [138]. Code for the transcriptomic analyses can be found at `https://github.com/JoramKeijser/interneuron_evolution` [139]. Code for the protein reconstruction can be found at `https://github.com/JoramKeijser/elfn1_reconstruction` [140]

Datasets

We analysed the following publicly available single cell RNA sequencing data sets: Mouse data from ref. [74] (downloaded from [141]), human data from ref. [75] (downloaded from [142]), zebra finch data (downloaded from [143]), and turtle data from ref. [91] (downloaded from [144]). The paper's code repository contains a script for automatically downloading the corresponding files.

For each data set, the starting point of our analysis was a matrix of gene counts per cell, together with the clustering of cells from the original publications. We converted each of the datasets to Seurat (v4 [145]) and AnnData (v0.8 [146]) objects for downstream analysis in Python and R, respectively. For visual comparison,

we labeled songbird and turtle cell clusters according to the most similar mammalian interneuron subclass, as determined in the original publications. This involved the merging of fine-level clusters that presumably capture within-subclass differences. For each dataset, we only visualized cells part of, or corresponding to, cortical interneurons. In particular, we did not visualize the correlation of the songbird GABAergic clusters 7, 8, and Pre, since these seem homologous to mouse olfactory bulb interneurons [92].

Dimensionality reduction and clustering

We used AnnData and Scanpy (v1.9.1 [147]) to visualize the expression of the *Elfn1* and *Cbln4* genes. This was done separately for each dataset. We first scaled the counts from each cell to counts per 10 thousand (CP10K) to account for differences in sequencing depth. We then used log plus one pseudo count (log1p) as variance-stabilizing transformation. Finally, we reduced the dimensionality of each dataset, by first finding highly variable genes, performed PCA followed by UMAP [77]. We used scanpy's default parameters for each of these steps. To investigate *Cbln4* expression within the SST population, we performed dimensionality reduction on all SST cells except long-range projecting Chodl cells. Clustering was done using the Leiden algorithm [148] with resolution 1.

Correlation analysis

We quantified the overall similarity of species-specific cell clusters by replicating the correlation analysis from refs. [91, 92]. We separately performed the following analysis on GABAergic and glutamatergic cells, and only compared zebra finch and mice. Specifically, we performed the following steps.

1. Select genes to compare across species. For each species, determine subclass-specific marker genes using Seurat's findAllMarkers (t-test, min.pct $= 0.2$, max.cells.per.ident $= 200$) and retain genes with Bonferroni adjusted p-value below 0.05.

2. Intersect the two species-specific lists to find genes that are differentially expressed in both species. This resulted in approximately 500 genes, depending on the cell type.

3. Average counts within each cluster and transform to log scale for variance-stabilization. Specifically, compute: $\log(1 + x) + 0.1$, with x the average count.

4. Divide each gene's value by its average across clusters to obtain a "specificity score" invariant to a genes' overall expression [91].

5. Compute the Pearson correlation between all pairs of mouse and songbird clusters.

We visualised the result using the R package pheatmap (v1.10.12 [149]).

Dataset integration

We used Seurat's anchor-based integration [150] to integrate the zebra finch and mouse data. We did this for GABAergic and glutamatergic neurons separately. First, we jointly performed normalization and variance stabilization for each dataset using Seurat's scTransform [151], with the percentage of mitochondrial counts as covariate. Next, we found the top 3000 most variable features across datasets, and used these to identify a set of anchors. These were then used to integrate the datasets. Finally, we jointly analysed the integrated datasets using Seurat's standard visualization pipeline: scaling and centering, PCA, and UMAP.

Ancestral Elfn1 reconstruction

We used the Topiary pipeline [113] to reconstruct the amino acid sequences of the ancestral Elfn1 protein based on sequences of extant species. To this end, we first constructed a source dataset consisting of the Elfn1 sequences from *Mus musculus* (mouse), *Homo sapiens* (human), *Taeniopygia guttata* (zebra finch), and *Pelodiscus sinensis* (Chinese softshell turtle). Next, we used Topiary's seed-to-alignment to find sequence homologs, perform reciprocal BLAST [152] to predict their orthology, reduce sequence redundancy, and align the remaining sequences using Muscle5 [153]. This resulted in 62 aligned sequences that were used as input to

Topiary's alignment-to-ancestors. This infers the maximum likelihood (ML) gene tree, the ML substitution model, and the ML ancestral sequences using RAxML-NG [154]. The posterior probability of an ancestral amino acid was computed using the amino acid's likelihood weighted by its prior probability, normalised by the sum over all amino acids. Topiary generates bootstrap replicates of the ML gene tree, and uses GeneRax [155] to reconcile the gene tree with the species tree. The number of bootstrap replicates was 700, as automatically determined by the software. This number—but not the reconstructed ML sequence—varied slightly between runs. Finally, we used topiary's bootstrap-reconcile that estimates the branch support for the reconciled tree. The ancestral sequence contained 16 ambiguous sites (based on a posterior probability cutoff of 0.25). Besides the ML sequence, we also report a worst case "altAll" sequence in which these ambiguous sites have been replaced by the next most-likely amino acid. Branch support for the amniote ancestor was 100/100, indicating very high confidence in the existence of this ancestor, as expected. We aligned extant and ancestral sequences using Muscle5, and visualised the resulting alignment using the R package Ggmsa [156].

References

[1] Robin Tremblay, Soohyun Lee, and Bernardo Rudy. Gabaergic interneurons in the neocortex: from cellular properties to circuits. *Neuron*, 91(2):260–292, 2016.

[2] Adam Kepecs and Gordon Fishell. Interneuron cell types are fit to function. *Nature*, 505(7483):318–326, 2014.

[3] Sadra Sadeh and Claudia Clopath. Inhibitory stabilization and cortical computation. *Nature Reviews Neuroscience*, 22(1):21–37, 2021.

[4] Tim P Vogels, Henning Sprekeler, Friedemann Zenke, Claudia Clopath, and Wulfram Gerstner. Inhibitory plasticity balances excitation and inhibition in sensory pathways and memory networks. *Science*, 334(6062):1569–1573, 2011.

[5] Yue Kris Wu, Christoph Miehl, and Julijana Gjorgjieva. Regulation of circuit organization and function through inhibitory synaptic plasticity. *Trends in Neurosciences*, 2022.

[6] Johannes J Letzkus, Steffen BE Wolff, and Andreas Lüthi. Disinhibition, a circuit mechanism for associative learning and memory. *Neuron*, 88(2):264–276, 2015.

[7] Leena E Williams and Anthony Holtmaat. Higher-order thalamocortical inputs gate synaptic long-term potentiation via disinhibition. *Neuron*, 101(1):91–102, 2019.

[8] Michael Wehr and Anthony M Zador. Balanced inhibition underlies tuning and sharpens spike timing in auditory cortex. *Nature*, 426(6965):442–446, 2003.

[9] Aanchal Bhatia, Sahil Moza, and Upinder Singh Bhalla. Precise excitation-inhibition balance controls gain and timing in the hippocampus. *Elife*, 8:e43415, 2019.

[10] Georg B Keller and Thomas D Mrsic-Flogel. Predictive processing: a canonical cortical computation. *Neuron*, 100(2):424–435, 2018.

[11] Loreen Hertäg and Claudia Clopath. Prediction-error neurons in circuits with multiple neuron types: Formation, refinement, and functional implications. *Proceedings of the National Academy of Sciences*, 119(13):e2115699119, 2022.

[12] Yu Fu, Jason M Tucciarone, J Sebastian Espinosa, Nengyin Sheng, Daniel P Darcy, Roger A Nicoll, Z Josh Huang, and Michael P Stryker. A cortical circuit for gain control by behavioral state. *Cell*, 156(6):1139–1152, 2014.

[13] Katie A Ferguson and Jessica A Cardin. Mechanisms underlying gain modulation in the cortex. *Nature Reviews Neuroscience*, 21(2):80–92, 2020.

[14] Gord Fishell and Adam Kepecs. Interneuron types as attractors and controllers. *Annual review of neuroscience*, 43:1–30, 2020.

[15] Joram Keijser and Henning Sprekeler. Optimizing interneuron circuits for compartment-specific feedback inhibition. *PLOS Computational Biology*, 18(4):e1009933, 2022.

[16] Matthew E Larkum, J Julius Zhu, and Bert Sakmann. A new cellular mechanism for coupling inputs arriving at different cortical layers. *Nature*, 398(6725):338–341, 1999.

[17] Leopoldo Petreanu, Daniel Huber, Aleksander Sobczyk, and Karel Svoboda. Channelrhodopsin-2–assisted circuit mapping of long-range callosal projections. *Nature neuroscience*, 10(5):663–668, 2007.

[18] Matthew Larkum. A cellular mechanism for cortical associations: an organizing principle for the cerebral cortex. *Trends in neurosciences*, 36(3):141–151, 2013.

[19] Julia MT Ledderose, Timothy A Zolnik, Maria Toumazou, Thorsten Trimbuch, Christian Rosenmund, Britta J Eickholt, Dieter Jaeger, Matthew E Larkum, and Robert NS Sachdev. Input to cortical layer 1 of somatosensory cortex. *bioRxiv*, pages 2021–11, 2022.

[20] Johannes J Letzkus, Björn M Kampa, and Greg J Stuart. Learning rules for spike timing-dependent plasticity depend on dendritic synapse location. *Journal of Neuroscience*, 26(41):10420–10429, 2006.

[21] P Jesper Sjostrom, Ede A Rancz, Arnd Roth, and Michael Hausser. Dendritic excitability and synaptic plasticity. *Physiological reviews*, 88(2):769–840, 2008.

[22] Matt Udakis, Victor Pedrosa, Sophie EL Chamberlain, Claudia Clopath, and Jack R Mellor. Interneuron-specific plasticity at parvalbumin and somatostatin inhibitory synapses onto ca1 pyramidal neurons shapes hippocampal output. *Nature communications*, 11(1):1–17, 2020.

[23] Alex Reyes, Rafael Lujan, Andrej Rozov, Nail Burnashev, Peter Somogyi, and Bert Sakmann. Target-cell-specific facilitation and depression in neocortical circuits. *Nature neuroscience*, 1(4):279–285, 1998.

[24] Gilad Silberberg and Henry Markram. Disynaptic inhibition between neocortical pyramidal cells mediated by martinotti cells. *Neuron*, 53(5):735–746, 2007.

[25] Thomas K Berger, Gilad Silberberg, Rodrigo Perin, and Henry Markram. Brief bursts self-inhibit and correlate the pyramidal network. *PLoS biology*, 8(9):e1000473, 2010.

[26] Masanori Murayama, Enrique Pérez-Garci, Thomas Nevian, Tobias Bock, Walter Senn, and Matthew E Larkum. Dendritic encoding of sensory stimuli controlled by deep cortical interneurons. *Nature*, 457(7233):1137–1141, 2009.

[27] Jesse H Goldberg, Clay O Lacefield, and Rafael Yuste. Global dendritic calcium spikes in mouse layer 5 low threshold spiking interneurones: implications for control of pyramidal cell bursting. *The Journal of physiology*, 558(2):465–478, 2004.

[28] Stephen R Williams and Greg J Stuart. Mechanisms and consequences of action potential burst firing in rat neocortical pyramidal neurons. *The Journal of physiology*, 521(Pt 2):467, 1999.

[29] Luc J Gentet, Yves Kremer, Hiroki Taniguchi, Z Josh Huang, Jochen F Staiger, and Carl CH Petersen. Unique functional properties of somatostatin-expressing gabaergic neurons in mouse barrel cortex. *Nature neuroscience*, 15(4):607–612, 2012.

[30] Matthew Lovett-Barron, Gergely F Turi, Patrick Kaifosh, Peter H Lee, Frédéric Bolze, Xiao-Hua Sun, Jean-François Nicoud, Boris V Zemelman, Scott M Sternson, and Attila Losonczy. Regulation of neuronal input transformations by tunable dendritic inhibition. *Nature neuroscience*, 15(3):423–430, 2012.

[31] Olivier Caillard, Herman Moreno, Beat Schwaller, Isabel Llano, Marco R Celio, and Alain Marty. Role of the calcium-binding protein parvalbumin in short-term synaptic plasticity. *Proceedings of the National Academy of Sciences*, 97(24):13372–13377, 2000.

[32] Frederic Pouille and Massimo Scanziani. Routing of spike series by dynamic circuits in the hippocampus. *Nature*, 429(6993):717–723, 2004.

[33] Xiaolong Jiang, Shan Shen, Cathryn R Cadwell, Philipp Berens, Fabian Sinz, Alexander S Ecker, Saumil Patel, and Andreas S Tolias. Principles of connectivity among morphologically defined cell types in adult neocortex. *Science*, 350(6264):aac9462, 2015.

[34] Luke Campagnola, Stephanie C Seeman, Thomas Chartrand, Lisa Kim, Alex Hoggarth, Clare Gamlin, Shinya Ito, Jessica Trinh, Pasha Davoudian, Cristina Radaelli, et al. Local connectivity and synaptic dynamics in mouse and human neocortex. *Science*, 375(6585):eabj5861, 2022.

[35] Richard Naud, Brice Bathellier, and Wulfram Gerstner. Spike timing prediction with active dendrites. *arXiv preprint arXiv:1311.3586*, 2013.

[36] Richard Naud and Henning Sprekeler. Sparse bursts optimize information transmission in a multiplexed neural code. *Proceedings of the National Academy of Sciences*, 115(27):E6329–E6338, 2018.

[37] Misha Tsodyks, Klaus Pawelzik, and Henry Markram. Neural networks with dynamic synapses. *Neural computation*, 10(4):821–835, 1998.

[38] David V Hansen, Jan H Lui, Pierre Flandin, Kazuaki Yoshikawa, John L Rubenstein, Arturo Alvarez-Buylla, and Arnold R Kriegstein. Non-epithelial stem cells and cortical interneuron production in the human ganglionic eminences. *Nature neuroscience*, 16(11):1576–1587, 2013.

[39] Tong Ma, Congmin Wang, Lei Wang, Xing Zhou, Miao Tian, Qiangqiang Zhang, Yue Zhang, Jiwen Li, Zhidong Liu, Yuqun Cai, et al. Subcortical origins of human and monkey neocortical interneurons. *Nature neuroscience*, 16(11):1588–1597, 2013.

[40] Yingchao Shi, Mengdi Wang, Da Mi, Tian Lu, Bosong Wang, Hao Dong, Suijuan Zhong, Youqiao Chen, Le Sun, Xin Zhou, et al. Mouse and human share conserved transcriptional programs for interneuron development. *Science*, 374(6573):eabj6641, 2021.

[41] Matthew T Schmitz, Kadellyn Sandoval, Christopher P Chen, Mohammed A Mostajo-Radji, William W Seeley, Tomasz J Nowakowski, Chun Jimmie Ye, Mercedes F Paredes, and Alex A Pollen. The development and evolution of inhibitory neurons in primate cerebrum. *Nature*, 603(7903):871–877, 2022.

[42] SA Anderson, DD Eisenstat, L Shi, and JLR Rubenstein. Interneuron migration from basal forebrain to neocortex: dependence on dlx genes. *Science*, 278(5337):474–476, 1997.

[43] Nuria Flames, Ramón Pla, Diego M Gelman, John LR Rubenstein, Luis Puelles, and Oscar Marín. Delineation of multiple subpallial progenitor domains by the combinatorial expression of transcriptional codes. *Journal of Neuroscience*, 27(36):9682–9695, 2007.

[44] Brie Wamsley and Gord Fishell. Genetic and activity-dependent mechanisms underlying interneuron diversity. *Nature Reviews Neuroscience*, 18(5):299–309, 2017.

[45] Lynette Lim, Da Mi, Alfredo Llorca, and Oscar Marín. Development and functional diversification of cortical interneurons. *Neuron*, 100(2):294–313, 2018.

[46] Carl P Wonders and Stewart A Anderson. The origin and specification of cortical interneurons. *Nature Reviews Neuroscience*, 7(9):687–696, 2006.

[47] Da Mi, Zhen Li, Lynette Lim, Mingfeng Li, Monika Moissidis, Yifei Yang, Tianliuyun Gao, Tim Xiaoming Hu, Thomas Pratt, David J Price, et al. Early emergence of cortical interneuron diversity in the mouse embryo. *Science*, 360(6384):81–85, 2018.

[48] Christian Mayer, Christoph Hafemeister, Rachel C Bandler, Robert Machold, Renata Batista Brito, Xavier Jaglin, Kathryn Allaway, Andrew Butler, Gord Fishell, and Rahul Satija. Developmental diversification of cortical inhibitory interneurons. *Nature*, 555(7697):457–462, 2018.

[49] Rachel C Bandler, Ilaria Vitali, Ryan N Delgado, May C Ho, Elena Dvoretskova, Josue S Ibarra Molinas, Paul W Frazel, Maesoumeh Mohammadkhani, Robert Machold, Sophia Maedler, et al. Single-cell delineation of lineage and genetic identity in the mouse brain. *Nature*, pages 1–6, 2021.

[50] Oscar Marín and John LR Rubenstein. A long, remarkable journey: tangential migration in the telencephalon. *Nature Reviews Neuroscience*, 2(11):780–790, 2001.

[51] Jia Sheng Hu, Daniel Vogt, Susan Lindtner, Magnus Sandberg, Shanni N Silberberg, and John LR Rubenstein. Coup-tf1 and coup-tf2 control subtype and laminar identity of mge-derived neocortical interneurons. *Development*, 144(15):2837–2851, 2017.

[52] Hynek Wichterle, Daniel H Turnbull, Susana Nery, Gord Fishell, and Arturo Alvarez-Buylla. In utero fate mapping reveals distinct migratory pathways and fates of neurons born in the mammalian basal forebrain. *Development*, 2001.

[53] Susana Nery, Gord Fishell, and Joshua G Corbin. The caudal ganglionic eminence is a source of distinct cortical and subcortical cell populations. *Nature neuroscience*, 5(12):1279–1287, 2002.

[54] Qing Xu, Inma Cobos, Estanislao De La Cruz, John L Rubenstein, and Stewart A Anderson. Origins of cortical interneuron subtypes. *Journal of Neuroscience*, 24(11):2612–2622, 2004.

[55] Simon JB Butt, Marc Fuccillo, Susana Nery, Steven Noctor, Arnold Kriegstein, Joshua G Corbin, and Gord Fishell. The temporal and spatial origins of cortical interneurons predict their physiological subtype. *Neuron*, 48(4):591–604, 2005.

[56] Goichi Miyoshi, Jens Hjerling-Leffler, Theofanis Karayannis, Vitor H Sousa, Simon JB Butt, James Battiste, Jane E Johnson, Robert P Machold, and Gord Fishell. Genetic fate mapping reveals that the caudal ganglionic eminence produces a large and diverse population of superficial cortical interneurons. *Journal of Neuroscience*, 30(5):1582–1594, 2010.

[57] Lori Sussel, Oscar Marin, Shioko Kimura, and JL Rubenstein. Loss of nkx2. 1 homeobox gene function results in a ventral to dorsal molecular respecification within the basal telencephalon: evidence for a transformation of the pallidum into the striatum. *Development*, 126(15):3359–3370, 1999.

[58] Simon JB Butt, Vitor H Sousa, Marc V Fuccillo, Jens Hjerling-Leffler, Goichi Miyoshi, Shioko Kimura, and Gord Fishell. The requirement of nkx2-1 in the temporal specification of cortical interneuron subtypes. *Neuron*, 59(5):722–732, 2008.

[59] Carl P Wonders, Lauren Taylor, Jelle Welagen, Ihunanya C Mbata, Jenny Z Xiang, and Stewart A Anderson. A spatial bias for the origins of interneuron subgroups within the medial ganglionic eminence. *Developmental biology*, 314(1):127–136, 2008.

[60] Matthew Fogarty, Matthew Grist, Diego Gelman, Oscar Marín, Vassilis Pachnis, and Nicoletta Kessaris. Spatial genetic patterning of the embryonic neuroepithelium generates gabaergic interneuron diversity in the adult cortex. *Journal of Neuroscience*, 27(41):10935–10946, 2007.

[61] Melis Inan, Jelle Welagen, and Stewart A Anderson. Spatial and temporal bias in the mitotic origins of somatostatin-and parvalbumin-expressing interneuron subgroups and the chandelier subtype in the medial ganglionic eminence. *Cerebral cortex*, 22(4):820–827, 2012.

[62] Miao He, Jason Tucciarone, SooHyun Lee, Maximiliano José Nigro, Yongsoo Kim, Jesse Maurica Levine, Sean Michael Kelly, Illya Krugikov, Priscilla Wu, Yang Chen, et al. Strategies and tools for combinatorial targeting of gabaergic neurons in mouse cerebral cortex. *Neuron*, 91(6):1228–1243, 2016.

[63] Melissa G McKenzie, Lucy V Cobbs, Patrick D Dummer, Timothy J Petros, Michael M Halford, Steven A Stacker, Yimin Zou, Gord J Fishell, and Edmund Au. Non-canonical wnt signaling through ryk regulates the generation of somatostatin-and parvalbumin-expressing cortical interneurons. *Neuron*, 103(5):853–864, 2019.

[64] Lynette Lim, Janelle MP Pakan, Martijn M Selten, André Marques-Smith, Alfredo Llorca, Sung Eun Bae, Nathalie L Rochefort, and Oscar Marín. Optimization of interneuron function by direct coupling of cell migration and axonal targeting. *Nature neuroscience*, 21(7):920–931, 2018.

[65] Pasko Rakic. Neuron-glia relationship during granule cell migration in developing cerebellar cortex. a golgi and electonmicroscopic study in macacus rhesus. *Journal of Comparative Neurology*, 141(3):283–312, 1971.

[66] J Julius Zhu. Maturation of layer 5 neocortical pyramidal neurons: amplifying salient layer 1 and layer 4 inputs by ca2+ action potentials in adult rat tuft dendrites. *The Journal of physiology*, 526(3):571–587, 2000.

[67] S Franceschetti, G Sancini, F Panzica, C Radici, and G Avanzini. Postnatal differentiation of firing properties and morphological characteristics in layer v pyramidal neurons of the sensorimotor cortex. *Neuroscience*, 83(4):1013–1024, 1998.

[68] Sandrine Romand, Yun Wang, Maria Toledo-Rodriguez, and Henry Markram. Morphological development of thick-tufted layer v pyramidal cells in the rat somatosensory cortex. *Frontiers in neuroanatomy*, 5:5, 2011.

[69] Michael Beierlein and Barry W Connors. Short-term dynamics of thalamocortical and intracortical synapses onto layer 6 neurons in neocortex. *Journal of neurophysiology*, 88(4):1924–1932, 2002.

[70] Emily L Sylwestrak and Anirvan Ghosh. Elfn1 regulates target-specific release probability at ca1-interneuron synapses. *Science*, 338(6106):536–540, 2012.

[71] Tevye Jason Stachniak, Emily Lauren Sylwestrak, Peter Scheiffele, Benjamin J Hall, and Anirvan Ghosh. Elfn1-induced constitutive activation of mglur7 determines frequency-dependent recruitment of somatostatin interneurons. *Journal of Neuroscience*, 39(23):4461–4474, 2019.

[72] Naoko H Tomioka, Hiroki Yasuda, Hiroyuki Miyamoto, Minoru Hatayama, Naoko Morimura, Yoshifumi Matsumoto, Toshimitsu Suzuki, Maya Odagawa, Yuri S Odaka, Yoshimi Iwayama, et al. Elfn1 recruits presynaptic mglur7 in trans and its loss results in seizures. *Nature communications*, 5(1):1–16, 2014.

[73] Emilia Favuzzi, Rubén Deogracias, André Marques-Smith, Patricia Maeso, Julie Jezequel, David Exposito-Alonso, Maddalena Balia, Tim Kroon, Antonio J Hinojosa, Elisa F Maraver, et al. Distinct molecular programs regulate synapse specificity in cortical inhibitory circuits. *Science*, 363(6425):413–417, 2019.

[74] Bosiljka Tasic, Zizhen Yao, Lucas T Graybuck, Kimberly A Smith, Thuc Nghi Nguyen, Darren Bertagnolli, Jeff Goldy, Emma Garren, Michael N Economo, Sarada Viswanathan, et al. Shared and distinct transcriptomic cell types across neocortical areas. *Nature*, 563(7729):72–78, 2018.

[75] Trygve E Bakken, Nikolas L Jorstad, Qiwen Hu, Blue B Lake, Wei Tian, Brian E Kalmbach, Megan Crow, Rebecca D Hodge, Fenna M Krienen, Staci A Sorensen, et al. Comparative cellular analysis of motor cortex in human, marmoset and mouse. *Nature*, 598(7879):111–119, 2021.

[76] Tevye Jason Stachniak, Rahel Kastli, Olivia Hanley, Ali Özgür Argunsah, Elianne Grietje Theodora van der Valk, George Kanatouris, and Theofanis Karayannis. Postmitotic prox1 expression controls the final specification of cortical vip interneuron subtypes. *Journal of Neuroscience*, 41(39):8150–8162, 2021.

[77] Leland McInnes, John Healy, and James Melville. Umap: Uniform manifold approximation and projection for dimension reduction. *arXiv preprint arXiv:1802.03426*, 2018.

[78] Graziella Di Cristo, Caizhi Wu, Bidisha Chattopadhyaya, Fabrice Ango, Graham Knott, Egbert Welker, Karel Svoboda, and Z Josh Huang. Subcellular domain-restricted gabaergic innervation in primary visual cortex in the absence of sensory and thalamic inputs. *Nature neuroscience*, 7(11):1184–1186, 2004.

[79] Robin F Dard, Erwan Leprince, Julien Denis, Shrisha Rao Balappa, Dmitrii Suchkov, Richard Boyce, Catherine Lopez, Marie Giorgi-Kurz, Tom Szwagier, Théo Dumont, et al. The rapid developmental rise of somatic inhibition disengages hippocampal dynamics from self-motion. *Elife*, 11:e78116, 2022.

[80] Nathan W Gouwens, Staci A Sorensen, Fahimeh Baftizadeh, Agata Budzillo, Brian R Lee, Tim Jarsky, Lauren Alfiler, Katherine Baker, Eliza Barkan, Kyla Berry, et al. Integrated morphoelectric and transcriptomic classification of cortical gabaergic cells. *Cell*, 183(4):935–953, 2020.

[81] Maximiliano José Nigro, Yoshiko Hashikawa-Yamasaki, and Bernardo Rudy. Diversity and connectivity of layer 5 somatostatin-expressing interneurons in the mouse barrel cortex. *Journal of Neuroscience*, 38(7):1622–1633, 2018.

[82] Federico Scala, Dmitry Kobak, Shen Shan, Yves Bernaerts, Sophie Laturnus, Cathryn Rene Cadwell, Leonard Hartmanis, Emmanouil Froudarakis, Jesus Ramon Castro, Zheng Huan Tan, et al. Layer 4 of mouse neocortex differs in cell types and circuit organization between sensory areas. *Nature communications*, 10(1):4174, 2019.

[83] Detlev Arendt, Jacob M Musser, Clare VH Baker, Aviv Bergman, Connie Cepko, Douglas H Erwin, Mihaela Pavlicev, Gerhard Schlosser, Stefanie Widder, Manfred D Laubichler, et al. The origin and evolution of cell types. *Nature Reviews Genetics*, 17(12):744–757, 2016.

[84] Maria Antonietta Tosches. From cell types to an integrated understanding of brain evolution: the case of the cerebral cortex. *Annual Review of Cell and Developmental Biology*, 37:495–517, 2021.

[85] Albert Gidon, Timothy Adam Zolnik, Pawel Fidzinski, Felix Bolduan, Athanasia Papoutsi, Panayiota Poirazi, Martin Holtkamp, Imre Vida, and Matthew Evan Larkum. Dendritic action potentials and computation in human layer 2/3 cortical neurons. *Science*, 367(6473):83–87, 2020.

[86] Lou Beaulieu-Laroche, Norma J Brown, Marissa Hansen, Enrique HS Toloza, Jitendra Sharma, Ziv M Williams, Matthew P Frosch, Garth Rees Cosgrove, Sydney S Cash, and Mark T Harnett. Allometric rules for mammalian cortical layer 5 neuron biophysics. *Nature*, 600(7888):274–278, 2021.

[87] Fuchou Tang, Catalin Barbacioru, Yangzhou Wang, Ellen Nordman, Clarence Lee, Nanlan Xu, Xiaohui Wang, John Bodeau, Brian B Tuch, Asim Siddiqui, et al. mrna-seq whole-transcriptome analysis of a single cell. *Nature methods*, 6(5):377–382, 2009.

[88] Amos Tanay and Arnau Sebé-Pedrós. Evolutionary cell type mapping with single-cell genomics. *Trends in genetics*, 37(10):919–932, 2021.

[89] Amit Zeisel, Ana B Muñoz-Manchado, Simone Codeluppi, Peter Lönnerberg, Gioele La Manno, Anna Juréus, Sueli Marques, Hermany Munguba, Liqun He, Christer Betsholtz, et al. Cell types in the mouse cortex and hippocampus revealed by single-cell rna-seq. *Science*, 347(6226):1138–1142, 2015.

[90] Bosiljka Tasic, Vilas Menon, Thuc Nghi Nguyen, Tae Kyung Kim, Tim Jarsky, Zizhen Yao, Boaz Levi, Lucas T Gray, Staci A Sorensen, Tim Dolbeare, et al. Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nature neuroscience*, 19(2):335–346, 2016.

[91] Maria Antonietta Tosches, Tracy M Yamawaki, Robert K Naumann, Ariel A Jacobi, Georgi Tushev, and Gilles Laurent. Evolution of pallium, hippocampus, and cortical cell types revealed by single-cell transcriptomics in reptiles. *Science*, 360(6391):881–888, 2018.

[92] Bradley M Colquitt, Devin P Merullo, Genevieve Konopka, Todd F Roberts, and Michael S Brainard. Cellular transcriptomics reveals evolutionary identities of songbird vocal circuits. *Science*, 371(6530), 2021.

[93] Mark G Blanton, Janet M Shen, and Arnold R Kriegstein. Evidence for the inhibitory neurotransmitter γ-aminobutyric acid in aspiny and sparsely spiny nonpyramidal neurons of the turtle dorsal cortex. *Journal of Comparative Neurology*, 259(2):277–297, 1987.

[94] Anton Reiner. Neurotransmitter organization and connections of turtle cortex: implications for the evolution of mammalian isocortex. *Comparative Biochemistry and Physiology Part A: Physiology*, 104(4):735–748, 1993.

[95] Simona Lodato, Bradley J Molyneaux, Emanuela Zuccaro, Loyal A Goff, Hsu-Hsin Chen, Wen Yuan, Alyssa Meleski, Emi Takahashi, Shaun Mahony, John L Rinn, et al. Gene co-regulation by fezf2 selects neurotransmitter identity and connectivity of corticospinal neurons. *Nature neuroscience*, 17(8):1046–1054, 2014.

[96] Elizabeth A Alcamo, Laura Chirivella, Marcel Dautzenberg, Gergana Dobreva, Isabel Fariñas, Rudolf Grosschedl, and Susan K McConnell. Satb2 regulates callosal projection neuron identity in the developing cerebral cortex. *Neuron*, 57(3):364–377, 2008.

[97] Tadashi Nomura, Wataru Yamashita, Hitoshi Gotoh, and Katsuhiko Ono. Species-specific mechanisms of neuron subtype specification reveal evolutionary plasticity of amniote brain development. *Cell reports*, 22(12):3142–3151, 2018.

[98] Oliver Hobert. Regulatory logic of neuronal diversity: terminal selector genes and selector motifs. *Proceedings of the National Academy of Sciences*, 105(51):20067–20071, 2008.

[99] Maria Antonietta Tosches. Different origins for similar brain circuits. *Science*, 371(6530):676–677, 2021.

[100] Barry W Connors and AR Kriegstein. Cellular physiology of the turtle visual cortex: distinctive properties of pyramidal and stellate neurons. *Journal of Neuroscience*, 6(1):164–177, 1986.

[101] Matthew E Larkum, Shigeo Watanabe, Nechama Lasser-Ross, Paul Rhodes, and William N Ross. Dendritic properties of turtle pyramidal neurons. *Journal of neurophysiology*, 99(2):683–694, 2008.

[102] Brice Bathellier, Troy W Margrie, and Matthew E Larkum. Properties of piriform cortex pyramidal cell dendrites: implications for olfactory circuit design. *Journal of Neuroscience*, 29(40):12641–12652, 2009.

[103] Friedrich W Johenning, Prateep S Beed, Thorsten Trimbuch, Michael HK Bendels, Jochen Winterer, and Dietmar Schmitz. Dendritic compartment and neuronal output mode determine pathway-specific long-term potentiation in the piriform cortex. *Journal of Neuroscience*, 29(43):13649–13661, 2009.

[104] Amit Kumar, Oded Schiff, Edi Barkai, Bartlett W Mel, Alon Poleg-Polsky, and Jackie Schiller. Nmda spikes mediate amplification of inputs in the rat piriform cortex. *Elife*, 7:e38446, 2018.

[105] Timothy J Devoogd and Fernando Nottebohm. Sex differences in dendritic morphology of a song control nucleus in the canary: a quantitative golgi study. *Journal of Comparative Neurology*, 196(2):309–316, 1981.

[106] Sam E Benezra, Rajeevan T Narayanan, Robert Egger, Marcel Oberlaender, and Michael A Long. Morphological characterization of hvc projection neurons in the zebra finch (taeniopygia guttata). *Journal of Comparative Neurology*, 526(10):1673–1689, 2018.

[107] https://creativecommons.org/licenses/by-nc-sa/3.0/.

[108] Jörgen Kornfeld, Sam E Benezra, Rajeevan T Narayanan, Fabian Svara, Robert Egger, Marcel Oberlaender, Winfried Denk, and Michael A Long. Em connectomics reveals axonal target variation in a sequence-generating network. *Elife*, 6:e24364, 2017.

[109] https://creativecommons.org/licenses/by/4.0/.

[110] Jackie Dolan and Kevin J Mitchell. Mutation of elfn1 in mice causes seizures and hyperactivity. *PloS one*, 8(11):e80491, 2013.

[111] Henry A Dunn, Dipak N Patil, Yan Cao, Cesare Orlandi, and Kirill A Martemyanov. Synaptic adhesion protein elfn1 is a selective allosteric modulator of group iii metabotropic glutamate receptors in trans. *Proceedings of the National Academy of Sciences*, 115(19):5022–5027, 2018.

[112] Georg KA Hochberg and Joseph W Thornton. Reconstructing ancient proteins to understand the causes of structure and function. *Annual review of biophysics*, 46:247–269, 2017.

[113] Kona N Orlandi, Sophia R Phillips, Zachary R Sailer, Joseph L Harman, and Michael J Harms. Topiary: Pruning the manual labor from ancestral sequence reconstruction. *Protein Science*, 32(2):e4551, 2023.

[114] David T Jones, William R Taylor, and Janet M Thornton. The rapid generation of mutation data matrices from protein sequences. *Bioinformatics*, 8(3):275–282, 1992.

[115] Jackie Dolan, Karen Walshe, Samantha Alsbury, Karsten Hokamp, Sean O'Keeffe, Tatsuya Okafuji, Suzanne FC Miller, Guy Tear, and Kevin J Mitchell. The extracellular leucine-rich repeat superfamily; a comparative survey and analysis of evolutionary relationships and expression patterns. *BMC genomics*, 8(1):1–24, 2007.

[116] Myrto Denaxa, Guilherme Neves, Adam Rabinowitz, Sarah Kemlo, Petros Liodis, Juan Burrone, and Vassilis Pachnis. Modulation of apoptosis controls inhibitory interneuron number in the cortex. *Cell reports*, 22(7):1710–1721, 2018.

[117] Natalia V De Marco García, Rashi Priya, Sebnem N Tuncdemir, Gord Fishell, and Theofanis Karayannis. Sensory inputs control the integration of neurogliaform interneurons into cortical circuits. *Nature neuroscience*, 18(3):393–401, 2015.

[118] Andre Marques-Smith, Daniel Lyngholm, Anna-Kristin Kaufmann, Jacqueline A Stacey, Anna Hoerder-Suabedissen, Esther BE Becker, Michael C Wilson, Zoltán Molnár, and Simon JB Butt. A transient translaminar gabaergic interneuron circuit connects thalamocortical recipient layers in neonatal somatosensory cortex. *Neuron*, 89(3):536–549, 2016.

[119] Simona Lodato, Caroline Rouaux, Kathleen B Quast, Chanati Jantrachotechatchawan, Michèle Studer, Takao K Hensch, and Paola Arlotta. Excitatory projection neuron subtypes control the distribution of local inhibitory interneurons in the cerebral cortex. *Neuron*, 69(4):763–779, 2011.

[120] Jason C Wester, Vivek Mahadevan, Christopher T Rhodes, Daniela Calvigioni, Sanan Venkatesh, Dragan Maric, Steven Hunt, Xiaoqing Yuan, Yajun Zhang, Timothy J Petros, et al. Neocortical projection neurons instruct inhibitory interneuron circuit development in a lineage-dependent manner. *Neuron*, 102(5):960–975, 2019.

[121] Nathalie Dehorter, Gabriele Ciceri, Giorgia Bartolini, Lynette Lim, Isabel del Pino, and Oscar Marín. Tuning of fast-spiking interneuron properties by an activity-dependent transcriptional switch. *Science*, 349(6253):1216–1220, 2015.

[122] Thomas Klausberger, Peter J Magill, László F Márton, J David B Roberts, Philip M Cobden, György Buzsáki, and Peter Somogyi. Brain-state-and cell-type-specific firing of hippocampal interneurons in vivo. *Nature*, 421(6925):844–848, 2003.

[123] Marlene Bartos, Imre Vida, Michael Frotscher, Axel Meyer, Hannah Monyer, Jörg RP Geiger, and Peter Jonas. Fast synaptic inhibition promotes synchronized gamma oscillations in hippocampal interneuron networks. *Proceedings of the National Academy of Sciences*, 99(20):13222–13227, 2002.

[124] Valerio Mante, David Sussillo, Krishna V Shenoy, and William T Newsome. Context-dependent computation by recurrent dynamics in prefrontal cortex. *nature*, 503(7474):78–84, 2013.

[125] Daniel LK Yamins, Ha Hong, Charles F Cadieu, Ethan A Solomon, Darren Seibert, and James J DiCarlo. Performance-optimized hierarchical models predict neural responses in higher visual cortex. *Proceedings of the national academy of sciences*, 111(23):8619–8624, 2014.

[126] Andrew M Saxe, James L McClelland, and Surya Ganguli. A mathematical theory of semantic development in deep neural networks. *Proceedings of the National Academy of Sciences*, 116(23):11537–11546, 2019.

[127] Blake A Richards, Timothy P Lillicrap, Philippe Beaudoin, Yoshua Bengio, Rafal Bogacz, Amelia Christensen, Claudia Clopath, Rui Ponte Costa, Archy de Berker, Surya Ganguli, et al. A deep learning framework for neuroscience. *Nature neuroscience*, 22(11):1761–1770, 2019.

[128] Laura Driscoll, Krishna Shenoy, and David Sussillo. Flexible multitask computation in recurrent networks utilizes shared dynamical motifs. *bioRxiv*, pages 2022–08, 2022.

[129] Nikolaus Kriegeskorte and Jörn Diedrichsen. Peeling the onion of brain representations. *Annual review of neuroscience*, 42(1):407–432, 2019.

[130] Saurabh Vyas, Matthew D Golub, David Sussillo, and Krishna V Shenoy. Computation through neural population dynamics. *Annual Review of Neuroscience*, 43:249, 2020.

[131] H Francis Song, Guangyu R Yang, and Xiao-Jing Wang. Training excitatory-inhibitory recurrent neural networks for cognitive tasks: a simple and flexible framework. *PLoS computational biology*, 12(2):e1004792, 2016.

[132] Laura B Naumann, Joram Keijser, and Henning Sprekeler. Invariant neural subspaces maintained by feedback modulation. *eLife*, 11:e76096, 2022.

[133] Robert Kim, Yinghao Li, and Terrence J Sejnowski. Simple framework for constructing functional spiking recurrent neural networks. *Proceedings of the national academy of sciences*, 116(45):22811–22820, 2019.

[134] Nicolas Perez-Nieves, Vincent CH Leung, Pier Luigi Dragotti, and Dan FM Goodman. Neural heterogeneity promotes robust learning. *Nature communications*, 12(1):1–9, 2021.

[135] Nicolas Y Masse, Guangyu R Yang, H Francis Song, Xiao-Jing Wang, and David J Freedman. Circuit mechanisms for the maintenance and manipulation of information in working memory. *Nature neuroscience*, 22(7):1159–1167, 2019.

[136] Guido vanRossum. Python reference manual. *Department of Computer Science [CS]*, (R 9525), 1995.

[137] R Core Team. R: A language and environment for statistical computing. r foundation for statistical computing, vienna, austria. 2012, 2021.

[138] Patrick Mineault and Kento Nozawa. patrickmineault/codebook: 1.0.0, 2021.

[139] JoramKeijser. JoramKeijser/interneuron_evolution: Code after revision, March 2023.

[140] JoramKeijser. JoramKeijser/interneuron_evolution: Code after revision, March 2023.

[141] Bosiljka Tasic, Zizhen Yao, Lucas T Graybuck, Kimberly A Smith, Thuc Nghi Nguyen, Darren Bertagnolli, Jeff Goldy, Emma Garren, Michael N Economo, Sarada Viswanathan, et al. `https://portal.brain-map.org/atlases-and-data/rnaseq/mouse-v1-and-alm-smart-seq`.

[142] Trygve E Bakken, Nikolas L Jorstad, Qiwen Hu, Blue B Lake, Wei Tian, Brian E Kalmbach, Megan Crow, Rebecca D Hodge, Fenna M Krienen, Staci A Sorensen, et al. `https://portal.brain-map.org/atlases-and-data/rnaseq/human-m1-10x`.

[143] Bradley M Colquitt, Devin P Merullo, Genevieve Konopka, Todd F Roberts, and Michael S Brainard. `https://cloud.biohpc.swmed.edu/index.php/s/nLicEtkmjGGmRF8?path=%2FHVC_RA`.

[144] Maria Antonietta Tosches, Tracy M Yamawaki, Robert K Naumann, Ariel A Jacobi, Georgi Tushev, and Gilles Laurent. `https://public.brain.mpg.de/Laurent/ReptilePallium2018/`.

[145] Yuhan Hao, Stephanie Hao, Erica Andersen-Nissen, William M Mauck III, Shiwei Zheng, Andrew Butler, Maddie J Lee, Aaron J Wilk, Charlotte Darby, Michael Zager, et al. Integrated analysis of multimodal single-cell data. *Cell*, 184(13):3573–3587, 2021.

[146] Isaac Virshup, Sergei Rybakov, Fabian J Theis, Philipp Angerer, and F Alexander Wolf. anndata: Annotated data. *bioRxiv*, 2021.

[147] F Alexander Wolf, Philipp Angerer, and Fabian J Theis. Scanpy: large-scale single-cell gene expression data analysis. *Genome biology*, 19(1):1–5, 2018.

[148] Vincent A Traag, Ludo Waltman, and Nees Jan Van Eck. From louvain to leiden: guaranteeing well-connected communities. *Scientific reports*, 9(1):5233, 2019.

[149] Raivo Kolde et al. Pheatmap: pretty heatmaps. *R package version*, 1(2):726, 2012.

[150] Tim Stuart, Andrew Butler, Paul Hoffman, Christoph Hafemeister, Efthymia Papalexi, William M Mauck III, Yuhan Hao, Marlon Stoeckius, Peter Smibert, and Rahul Satija. Comprehensive integration of single-cell data. *Cell*, 177(7):1888–1902, 2019.

[151] Christoph Hafemeister and Rahul Satija. Normalization and variance stabilization of single-cell rna-seq data using regularized negative binomial regression. *Genome biology*, 20(1):1–15, 2019.

[152] Stephen F Altschul, Warren Gish, Webb Miller, Eugene W Myers, and David J Lipman. Basic local alignment search tool. *Journal of molecular biology*, 215(3):403–410, 1990.

[153] Robert C Edgar. Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence homology and phylogeny. *Nature Communications*, 13(1):6968, 2022.

[154] Alexey M Kozlov, Diego Darriba, Tomáš Flouri, Benoit Morel, and Alexandros Stamatakis. Raxml-ng: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics*, 35(21):4453–4455, 2019.

[155] Benoit Morel, Alexey M Kozlov, Alexandros Stamatakis, and Gergely J Szöllősi. Generax: a tool for species-tree-aware maximum likelihood-based gene family tree inference under gene duplication, transfer, and loss. *Molecular biology and evolution*, 37(9):2763–2774, 2020.

[156] Lang Zhou, Tingze Feng, Shuangbin Xu, Fangluan Gao, Tommy T Lam, Qianwen Wang, Tianzhi Wu, Huina Huang, Li Zhan, Lin Li, et al. ggmsa: a visual exploration tool for multiple sequence alignment and associated data. *Briefings in Bioinformatics*, 23(4):bbac222, 2022.