Extensive re-modelling of the cell wall during the development of *Staphylococcus*

***aureus* bacteraemia.**

Edward J.A. Douglas[1,2]*, Nathanael Palk[1]*, Tarcisio Brignoli[1,3], Dina Altwiley[1], Marcia Boura[1],

Maisem Laabei[2], Mario Recker[4,5], Gordon Y.C. Cheung[6], Ryan Liu[6], Roger C. Hsieh[6],

Michael Otto[6], Eoin O'Brien[7], Rachel M. McLoughlin[7] and Ruth C. Massey[1,8,#].

1: School of Cellular and Molecular Medicine, University of Bristol, Bristol, BS8 1TD, UK.

2: Department of Life Sciences, University of Bath, BA2 7AY, UK.

3: Department of Biosciences, Università degli Studi di Milano, Milan, Italy.

4: Centre for Ecology and Conservation, University of Exeter, Penryn Campus, TR10 9FE,

UK.

5: Institute of Tropical Medicine, University of Tübingen, Tübingen, Germany.

6: Laboratory of Bacteriology, National Institute of Allergy and Infectious Diseases (NIAID),

US National Institutes of Health (NIH), Bethesda, Maryland, USA.

7: Host Pathogen Interactions Group, School of Biochemistry and Immunology, Trinity

College Dublin, Dublin 2, Ireland.

8: Schools of Microbiology and Medicine, University College Cork, and APC Microbiome

Ireland, Cork, Ireland.

* These authors contributed equally to this work.

For correspondence: ruth.massey@bristol.ac.uk; orcid id: https://orcid.org/0000-0002-

8154-4039

Introductory Paragraph / Abstract

The bloodstream represents a hostile environment that bacteria must overcome to cause bacteraemia. To understand how the major human pathogen *Staphylococcus aureus* manages this we have utilised a functional genomics approach to identify a number of new loci that affect the ability of the bacteria to survive exposure to serum, the critical first step in the development of bacteraemia. The expression of one of these genes, *tcaA,* was found to be induced upon exposure to serum, and we show that it is involved in the elaboration of a critical virulence factor, the wall teichoic acids (WTA), within the cell envelope. The activity of the TcaA protein alters the sensitivity of the bacteria to cell wall attacking agents, including antimicrobial peptides, human defence fatty acids, and several antibiotics. This protein also affects the autolytic activity and lysostaphin sensitivity of the bacteria, suggesting that in addition to changing WTA abundance in the cell envelope, it also plays a role in peptidoglycan crosslinking. With TcaA rendering the bacteria more susceptible to serum killing, while simultaneously increasing the abundance of WTA in the cell envelope, it was unclear what effect this protein may have during infection. To explore this, we examined human data and performed murine experimental infections. Collectively, our data suggests that whilst mutations in *tcaA* are selected for during bacteraemia, this protein positively contributes to the virulence of *S. aureus* through its involvement in altering the cell wall architecture of the bacteria, a process that appears to play a key role in the development of bacteraemia.

52
Main Text

Staphylococcus aureus is an important human pathogen and a significant global health

concern[1,2]. The most common interaction with its human host is as an asymptomatic coloniser;

however, it frequently transitions to a pathogenic state with the ability to cause a wide range

of diseases, ranging from relatively minor skin and soft tissue infections (SSTI) to more life-

threatening incidents of endocarditis or bacteraemia[1,2]. *S. aureus* is notorious for producing a

plethora of virulence factors ranging from pore forming toxins to various immune evasion

strategies[3]. The toxicity of *S. aureus* is generally accepted as playing an important role during

infection, with high toxicity isolates typically causing more severe symptoms and disease

progression[4,5]. However, for the more invasive diseases such as bacteraemia and pneumonia,

it has been shown that the causative isolates are often impaired in their toxin production and

instead rely on alternative virulence approaches to cause disease, such as being able to better

survive exposure to host immune defences[6,7].

As the most severe type of infection caused by *S. aureus*, bacteraemia has a mortality rate of

between 20 and 30%[8,9], and despite infection control measures that have reduced the

incidence of MRSA bacteraemia, the incidence of MSSA bacteraemia continues to increase

in several countries, suggesting we have much to learn about this disease process[10]. Entry

into the bloodstream is the first step in the development of bacteraemia and represents a major

bottle neck for the bacteria, whether they seed from an infection elsewhere in the body, or

take a more direct route through an intravenous device. As a heavily protected niche, the

bloodstream contains numerous humoral immune features with potent anti-staphylococcal

activity, including antimicrobial peptides (AMPs) and host defence fatty acids (HDFAs)[11-13].

However, the fact that cases of *S. aureus* bacteraemia occur demonstrates that these do not

represent an impregnable force and that the bacteria can adapt to resist these defensive

features.

In previous work we have pioneered the application of genome wide association studies

(GWAS) to characterise bacterial virulence, where it has proven to be a powerful approach to

define complex regulatory pathways[7, 15-19]. In related work we have also begun to characterise

a new category of *S. aureus* genes we refer to as MALs (mortality associated loci) and found

that many of these are involved in serum survival[20], with serum containing the bulk of the

humoral immune response present in blood, but without the added complexity of the cellular

components. Given the outstanding research questions surrounding the development of

bacteraemia we performed a GWAS on a collection of 300 clinical isolates with respect to

serum resistance. We found significant variability in how well individual isolates survive

exposure to serum and identified seven novel effectors of this activity. Of particular note was

the TcaA protein, which we have further characterised and demonstrate that it is a critical

protein for the bacteria involved in the remodelling of the bacterial cell wall during the

development of bacteraemia.

Results

Survival of S. aureus upon exposure to human serum is a variable trait. The human

bloodstream is a highly protected niche, and so to establish an infection in this environment

the bacteria must evade many aspects of host immunity, such as the bacterial membrane

damaging antimicrobial peptides (AMP) and host defence fatty acids (HDFAs) found in

serum[11-13,21-24]. To examine the level of variability that exists in serum-susceptibility in natural

populations of *S. aureus*, we exposed 300 clinical isolates to pooled human serum and

quantified the proportion of each culture that survived. These 300 isolates represent the two

major clones, as defined by multi-locus sequence typing: clonal complex 22 (CC22) and clonal

complex 30 (CC30), that are responsible for the majority of MRSA cases in the UK and

Ireland[16]. The isolates were collected from 300 individual cases of bacteraemia. This collection

is a unique resource as each strain has been sequenced, extensively phenotyped, and we

have clinical metadata for many of the patients the bacteria were isolated from[16]. We found

there to be similar levels of variability in the ability of the bacteria to survive exposure to human

serum across the two CCs, both with a >10^3 fold difference between the most and least

susceptible isolate with respect to the number of bacteria that survived (Fig. 1a &b).

Figure 1: Serum susceptibility is multifactorial and varies widely across closely
related *S. aureus* isolates. (**a & b**) The susceptibility of 300 bacteraemia isolates
from clonal complexes CC22 and CC30 were exposed to human serum and their
ability to survive this exposure quantified. The survival of each isolate was quantified
in triplicate and dot represents the mean of these data. (**c & d**) Manhattan plots
representing the statistical associations (on the y axis) between individual SNPs
across the genome (on the x axis) and serum survival. The dotted red line represents
the uncorrected significance threshold, and the dotted blue line represents the Sidak
corrected (for multiple comparisons) threshold.

Survival of S. aureus upon exposure to human serum is a polygenic trait. As the genome

sequence for each of the 300 clinical *S. aureus* isolates was available we performed a GWAS

(genome wide association study) to identify polymorphic loci or single nucleotide

polymorphisms (SNPs) that associated with the level of susceptibility of isolates to serum. For

this, the data from the two distinct clones were analysed independently, with population

structure within the clones being accounted for (Fig. 1c & d, Tables 1a & 1b). We applied both

uncorrected and corrected (for multiple comparisons) significance thresholds to this analysis,

as our previous work has demonstrated that the stringency of multiple correction approaches

increases the likelihood of type II errors (i.e. false negative results). For the CC22 collection

12 loci were associated with serum survival and for the CC30s there were 32 associated loci

(Fig. 1c & d, Tables 1a & 1b).

Tables 1: Loci associated with serum survival in the CC22 (Table 1a) and CC30

(Table 1b) collections. The SNP position is relative to the origin of replication in

the reference genomes: HO 5096 0412 (CC22) and MRSA252 (CC30). Locus

tags and where available gene names or putative protein functions have been

provided. NTML refers to the mutant available in the Nebraska transposon library.

Table 1a

SNP position	CC22 locus tag	Description	p Value	NTML
1945336	SAEMRSA15_RS09605	Intergenic	6.851546	N/A
375179	SAEMRSA15_03110	*metE*	5.606418	NE944
728154	SAEMRSA15_06310	*saeS*	5.553331	NE1296
1053036	N/A	Intergenic	3.60927	N/A
219764	SAEMRSA15_01760	putative oxidoreductase	3.575929	NE312
2280500	SAEMRSA15_20910	alcohol dehydrogenase	2.257841	NE1147
1196127	SAEMRSA15_10750	*yfhO*	2.212486	NE1597
408213	SAEMRSA15_03430	*guaA*	1.876101	N/A
2492381	SAEMRSA15_23120	*tcyB*	1.645759	NE228
1514441	SAEMRSA15_13420	hypothetical protein	1.600037	NE1349
2430758	SAEMRSA15_22540	*tcaA*	1.572702	NE285
134276	SAEMRSA15_01100	hypothetical protein	1.358386	NE599

141 **Table 1b**

SNP position	CC30 locus tag	Description	p Value	NTML
101174	SAR0097	putative DNA binding protein	18.466	N/A
1188605	SAR1143	*arcC*	12.786	NE1719
900409	SAR0856	phosphoglycerate mutase	11.226	NE1422
2667418	SAR2584	*gntR*	10.263	NE1124
2421610	SAR2347	putative membrane protein	10.160	N/A
1687821	SAR1615	*aroK*	9.793	N/A
532944	SAR0495	*veg*	9.751	NE1703
1252982	SAR1203	*recG*	8.502	NE1344
1816815	SAR1754	*clpX*	8.064	N/A
2206190	SAR2143	*ilvC*	8.050	NE1177
1443567	SAR1386	*trpA*	6.647	NE304
1167449	SAR1119	*uvrC*	6.439	NE1212
2442306	SAR2374	*ureC*	6.425	NE410
1784580	SAR1719	*tgt*	6.396	NE1885
880082	SAR0835	*est*	6.354	NE1122
1116463	SAR1070	*pdhD*	5.600	NE1610
289265	SAR0248	*tagB*	5.324	NE362
1442353	SAR1385	*trpB*	4.910	NE310
364861	SAR0319	NADH flavin oxidoreductase	4.373	N/A
1963781	SAR1877	*menE*	4.315	N/A
1946567	SAR1856	*arsB2*	4.122	NE1484
1416353	SAR1364	iron-sulfur cluster biosynthesis protein	2.829	NE1910
1887414	SAR1810	*fhs*	2.741	NE706
1133886	SAR0251	*tarF*	2.725	N/A
30969	SAR0023	*sasH*	2.507	NE295
330358	SAR0284	*essC*	2.486	NE1484
2220348	SAR2151	*tex*	1.704	NE1079
1468392	SAR1410	*hipO*	1.569	NE1688
1652613	SAR1582	*zwf*	1.556	N/A
2744774	SAR2656	*nrmA*	1.549	NE1223
2370812	SAR2287	*lacR*	1.392	NE346

Functional validation of the GWAS results identifies seven novel genes that affect serum

survival. Of the 44 loci associated with serum survival, there were transposon mutants

available for 32 in the Nebraska library[25]. To functionally verify our GWAS findings, each of

these mutants were tested for their ability to survive exposure to human serum. We found

seven mutants were significantly affected in this ability, and this effect was complemented by

expressing the inactivated gene *in trans* (Fig. 2). The pRMC2 plasmid used to complement the

mutated phenotypes requires the use of two antibiotics, one for selection of the plasmid and

one for induction of gene expression[26]. These antibiotics can often affect bacterial gene

expression. As such validation of complementation is based on the comparison between the

empty and target gene containing plasmid which can be grown under the same conditions.

The seven genes were *tcaA*, a gene associated with resistance to the antibiotic teicoplanin[27];

tarK, a gene involved in wall teichoic acid (WTA) biosynthesis[28]; *gntR*, which encodes

gluconate kinase[29]; *ilvC*, which encodes ketol-acid reductoisomerase[30]; *arsB*, which is an efflux

pump involved in arsenic resistance[31]; *yfhO*, which is involved in the glycosylation of

lipoteichoic acid[32]; and *pdhD*, which encodes a lipoamide dehydrogenase[33].

Figure 2: Functional verification of loci involved in serum sensitivity of *S. aureus*. The
effect of the inactivation of the genes associated serum sensitivity was examined
using transposon mutants of strain JE2. Of the 32 mutants tested seven were
significantly affected: (**a**) *tcaA*, (**b**) *tarK*, (**c**) *gntK*, (**d**) *ilvC*, (**e**) *arsB*, (**f**) *yhfO* and (**g**)
pdhD. The effect of the mutations on serum sensitivity was complemented by
expressing the gene from the expression plasmid pRMC2 (e.g. the *tcaA*
complementing plasmid is called p*tcaA*). The dots represent individual data points,

the bars the mean value, and the error bars the standard deviation. Significance was

determined as * <0.05, ** <0.01, *** <0.001.

The SNPs in the tcaA *gene of the clinical isolates are associated with decreased serum*

sensitivity. The *tcaA* gene has previously been reported to be involved in conferring increased

sensitivity to the antibiotic teicoplanin[27], although the molecular detail of how it achieves this

has yet to be determined, and our findings suggest it also confers increased sensitivity to

serum (Fig. 2a). To visualise the signal detected by the GWAS for this gene we have mapped

the position of the clinical isolates with the *tcaA* SNPs onto a figure displaying the range of

serum sensitivities of the collection (Fig. 3a). The majority of the isolates with the SNPs were

less sensitive to serum, suggesting the SNPs negatively affects the activity of the protein. The

most common SNP in the *tcaA* gene amongst the clinical isolates conferred a Phe (290) to

Ser change in the protein sequence. Using AlphaFold[34] we generated a structure for TcaA

where it is predicted to be membrane bound with a single membrane spanning domain; to

have a zinc finger presented intracellularly; with the majority of the protein including the

Phe290 displayed extracellularly (Fig. 3b). Given that phenylalanines are frequently involved

in protein-protein interactions[35] we examined the effect this change has on the activity of TcaA

by cloning the *tcaA* gene with this SNP into the complementing pRMC2 plasmid and

comparing its ability to confer serum resistance relative to that of the *tcaA* gene without the

SNP. The TcaA protein with Ser at position 290 instead of Phe were slightly more resistant to

serum killing, but not as affected as when the entire protein was inactivated, suggesting this

change has a subtle effect on the serum sensitising activity of the protein (Fig. 3c).

Figure 3: *The clinical* tcaA *mutations associate with increased serum resistance*. **(a)** The individual clinical isolates with polymorphism in the *tcaA* gene are indicated (in red) on a graph displaying the range of serum survival of the collection. **(b)** The structure of the TcaA protein as predicted by AlphaFold. **(c)** The most common *tcaA* SNP decreases the sensitivity of *S. aureus* to serum. The dots represent individual data points, the bars the mean value, and the error bars the standard deviation.

TcaA confers increased sensitivity to multiple antibacterial components of serum. To examine whether the effect of TcaA on serum resistance was specific to the JE2 background we transduced the *tcaA::tn* mutation from JE2 into *S. aureus* strains SH1000 and Newman. Attempts were made to transduce this mutation into a clinical CC22 isolate, but this lineage was refractory to any of the available transducing phage[36,37]. In all three backgrounds TcaA increased the sensitivity of the bacteria to serum (Fig. 4a) verifying that the effect was not limited to the JE2 background. To determine which aspects of human serum TcaA was

conferring increased sensitivity to, we measured the relative ability of the wild type and mutant

to survive exposure to some of the anti-bacterial factors found in serum: to the HDFA

arachidonic acid, and to two AMPs: HNP-1 and LL37. The strains producing TcaA (i.e. the WT

and complemented strain) were more sensitive to all three components of human serum (Fig.

4b-d). For the AMPs this sensitivity was significant after only 90 min exposure, whereas for

arachidonic acid sensitivity the bacteria were incubated overnight, and differences in

sensitivity determined by bacterial growth (OD_{600}).

Although none of the individual antibacterial components of serum tested here are involved in

S. aureus cell clumping, it is a possibility that some of the effects seen in serum is as a result

of this activity, artificially amplifying the effect. To examine this the wild type and *tcaA* mutant

were visually examined by microscopy in both PBS and serum where no differences in

clumping was observed (Supplementary Fig. 1). Of the other GWAS identified genes, the *ilvC*

mutant was less sensitive to arachidonic acid, the *yfhO* mutant was more resistant to HNP-1

and LL-37, while the *tarK, gntK, arsB* and *phdD* mutants were all more sensitive to HNP-1 and

LL37 when compared to the JE2 wild type strain (Supplementary Fig. 2).

Figure 4: TcaA production confers increased sensitivity to several antibacterial

components of serum. **(a)** The inactivation of the *tcaA* gene decreases the

sensitivity of three *S. aureus* strains (i.e., JE2, Newman and SH1000) to serum (90

min exposure). **(b)** TcaA production by the wild type and p*tcaA* complemented

strain confers increased sensitivity to arachidonic acid (100μM) following overnight

growth. **(c)** TcaA production by the JE2 wild type and p*tcaA* complemented strain

confers increased sensitivity to LL37 (5μg/ml, 90 min exposure) and **(d)** TcaA

production by the wild type and p*tcaA* complemented strain confers increased

sensitivity to HNP-1 (5μg/ml, 90 min exposure). The dots represent individual data

points, the bars the mean value, and the error bars the standard deviation.

Significance was determined as * <0.05, ** <0.01, *** <0.001.

Wall teichoic acids contribute to the arachidonic acid sensitivity of TcaA producing strains. The

antibacterial properties of HDFAs and AMPs share some common features in that they both

penetrate through the cell wall and attack the bacterial membrane[11-13,21-23]. AMPs are positively

charged molecules that rely on the relatively negative charge across the bacterial cell

envelope to penetrate, and thus resistance is frequently acquired by changing the charge

across the cell wall such that the AMPs are repelled[22,23]. Less is known about how HDFAs

penetrate the cell wall; however, resistance is associated with changes in the abundance of

wall teichoic acids (WTA) in the bacterial cell walls[24]. WTA are hydrophilic and when present

in the cell wall they are believed to be protective by interfering with the penetration of the

hydrophobic HDFAs through to the bacterial membrane[24]. It has also been shown that when

the ligation of WTA to the cell wall is affected, such that they are instead released into the

environment, this also decreases the sensitivity of the bacteria to HDFAs[13]. With distinct

modes of action and means of accessing their target, it is intriguing to consider how TcaA may

be contributing to increasing the sensitivity of *S. aureus* to both these types of anti-bacterial

molecules.

To examine whether the TcaA associated increase in sensitivity to arachidonic acid is a result

of an increase in the abundance of WTA in the bacterial cell we extracted and quantified WTA

from the wild type and *tcaA* mutant, where we found that the cell wall of the wild type had

significantly more WTA when compared to the *tcaA* mutant cell extract (Fig. 5a & b & c)). We

next quantified WTA in the bacterial supernatant, as recent work on a *S. aureus lcpA* mutant

demonstrated that another WTA related means of altering sensitivity to arachidonic acid is to

release the WTA the bacteria surface[11]. We found significantly less WTA in the extract of the

TcaA producing wild type strain compared to that of the *tcaA* mutant (Fig. 5a), which could

explain the observed difference in sensitivity to HDFAs. It was unclear from the previous work

on LcpA how the released WTA by the *lcpA* mutant affects arachidonic acid sensitivity, and

here we demonstrate that this mutant, like the *tcaA* mutant, is also less sensitive to human

serum (Fig. 5d). One hypothesis is that the WTA can either sequester or inactivate HDFAs in

the environment surrounding the bacteria, thereby neutralising them. To test this, we

harvested the supernatant of the wild type JE2, the JE2 *tcaA* mutant, and the JE2 *lcpA* mutant

from overnight growth. These supernatants were used to supplement fresh broth (at 10%) into

which the wild type JE2 strain was inoculated. In the absence of any arachidonic acid JE2

grew equally well regardless of the supernatant supplement (Fig. 5e). However, in the

presence of the supernatant of both the *tcaA* and *lcpA* mutant (both of which contain an

abundance of released WTA), JE2 was able to grow in the presence of arachidonic acid;

whereas when the JE2 supernatant was used as the supplement the bacteria were unable to

grow (Fig. 5e). To further verify that soluble WTA can neutralise arachidonic we performed

WTA extractions from an isogenic wild type and mutant of the *S. aureus* strain LAC where the

mutant has had the *tarO* gene deleted and consequently does not produce any WTA (Fig. 5e).

The incorporation of the wild type LAC purified WTA extract to broth at 2% created an

environment in which JE2 could grow in the presence of arachidonic acid, whereas the

equivalent WTA extract of the *tarO* mutant, which contains no WTA, did not (Fig. 5e). Together

these data demonstrate that WTA in the environment can neutralise arachidonic acid, and this

provides a likely explanation for the increased sensitivity to HDFAs associated with TcaA

production.

Figure 5: Wall teichoic acids are released from the cell wall in the *tcaA* mutant to affect resistance to HDFAs and AMPs. **(a)** WTA was extracted from both cells and supernatant of the wild type and *tcaA* mutant of *S. aureus* and visualised on and SDS-PAGE gel stained with 1 mg/ml Alcian blue. The *tcaA* mutant had significantly less WTA in the cell wall but more in the supernatant. **(b)** The WTA extractions were performed on the wild type and tcaA mutant in triplicate the relative density of the WTA quantified by densitometry **(c)** The phosphate content of the cell wall WTA extracts was quantified, which verified that the *tcaA* mutant has significantly less WTA. **(d)** The inactivation of the *lcpA* gene increases the sensitivity of *S. aureus* to killing by serum. **(e)** The wild type strain JE2 was grown in broth supplemented (at 10%) with supernatant of either itself (JE2), with that from a *tcaA* mutant, or with that from a *lcpA* mutant. These supernatants had no effect on the growth of JE2 in the absence of arachidonic acid. In the presence of arachidonic acid JE2 was unable to grow when supplemented with its own supernatant, however, when supplemented with the supernatant of the two mutants, which both contain soluble WTA, JE2 was able to grow. The addition of purified WTA extract (at 2%) from another *S. aureus* strain (LAC) also neutralised arachidonic acid, however an equivalent extract from an isogenic WTA mutant (LAC Δ*tarO*) did not. **(f)** The charge across the cell wall of the wild type and *tcaA* mutant was compared using cytochrome C, where the mutant

was found to be less negatively charged. An *mprF* mutant has been included as a

control. The dots represent individual data points, the bars the mean value, and the

error bars the standard deviation. Significance was determined as * <0.05, ** <0.01,

*** <0.001, **** <0.0001.

The reduced abundance of WTA in the bacterial cell wall contributes to the resistance of S.

aureus to AMPs. In additional to being hydrophilic, WTA is also predominantly negatively

charged[28]. Given that a change in charge across the bacterial cell wall is frequently associated

with resistance to AMPs[21-23], and that the wild type strain has more WTA in its cell wall, this

may explain the increased AMP sensitivity of the TcaA producing strain. To examine this, we

incubated the bacteria with cytochrome C, which is positively charged and through its

electrostatic-related ability binds to the bacterial cells, providing a measure of the charge

across bacterial cell walls. Using this we found that the wild type strain bound significantly

more cytochrome C and is therefore more negatively charged than the *tcaA* mutant, which

explains the increase in AMP sensitivity we have found associated with TcaA production (Fig.

5e). Together these results explain how TcaA production affects the sensitivity of *S. aureus* to

both HDFAs and AMPs. By retaining WTA within the cell wall, there is less released to

sequester or inactivate the fatty acids from the environment, and due to their negative charge,

increased amounts of WTA in the cell wall affects the charge across the cell wall and results

in increased electrostatic attraction of the positively charged AMPs.

Serum induced tcaA expression increases both the abundance of WTA in the cell wall and the

sensitivity S. aureus to teicoplanin. As mentioned previously, an additional interesting feature

of the TcaA protein is that is has also been associated with changing the sensitivity of the

bacteria to the antibiotic teicoplanin, although this has been reported to vary between

strains[27,38]. To examine this for our strains we determined the Minimum inhibitory

concentrations (MICs) for three independent sets of isogenic wild type and tcaA mutants (ie

in strains JE2, Newman and SH1000, all corresponding to CC8), where TcaA production

resulted in increased sensitivity to this antibiotic across all three strains (Fig. 6a). To examine

whether the clinical isolates with the SNPs in the *tcaA* gene associated with increased serum

resistance also had altered teicoplanin sensitivity, we compared the ability of 12 clinical

isolates containing the GWAS identified *tcaA* SNPs to grow in teicoplanin to that of 12 isolates

from the same collection with the wild type gene sequence, i.e., without the polymorphism.

The isolates with the polymorphism in the *tcaA* gene were on average less sensitive to

teicoplanin (Fig. 6b). Using the pRMC2 complementing plasmid expressing either the wild type

or SNP (conferring the Phe290-Ser substitution) containing *tcaA* gene, we demonstrated that

this mutation significantly reduces the ability of the protein to sensitize the bacteria to

teicoplanin (Fig. 6c), confirming that this mutation negatively affects the activity of the TcaA

protein.

As TcaA has been shown to be upregulated when exposed to teicoplanin[27,38], we examined

whether serum would also induce its expression. The wild type JE2 strain was exposed to

either sub-inhibitory (2.5% serum and 0.5μg/ml teicoplanin) or inhibitory (10% serum and

10μg/ml teicoplanin) concentrations of either serum or teicoplanin for 20 mins, total RNA was

extracted and the transcription of the *tcaA* gene quantified by qRT-PCR. For both serum and

teicoplanin, induction of expression of the *tcaA* gene was concentration dependent (Fig. 6d).

Given the association of WTA abundance in the cell wall with *tcaA* expression described above

(Fig. 5a-c) we next examined whether serum induced expression of *tcaA* would also result in

an increase in abundance of WTA in the bacteria cell wall. Both the wild type and *tcaA* mutant

were exposed to inhibitory concentrations of human serum for 90 mins and the WTA extracted

from the cell walls, where serum exposure resulted in an increase in WTA for the wild type

strain but not the *tcaA* mutant (Fig. 6e & f). To examine whether serum induced increased

expression of *tcaA* would affect sensitivity to teicoplanin, we exposed the wild type bacteria to

a sub-inhibitory concentrations of serum (2.5%) and then incubated these overnight in a

subinhibitory concentration of teicoplanin (0.5μg/ml). Pre-exposure to serum induced

increased sensitivity to teicoplanin suggesting that induction of *tcaA* expression by the initial

serum exposure was sufficient to induce cross-sensitivity to the antibiotic (Fig. 6g). To put this

sensitisation effect in the context of serum, the reverse experiment was also performed

whereby the wild type strain was exposed to a sub-inhibitory concentration of teicoplanin

(0.5μg/ml) overnight and a serum killing assay performed following this incubation (Fig. 6h).

Pre-exposure to teicoplanin also induced increased sensitivity to serum again highlighting this

cross-sensitivity effect.

To verify this induction effect was not specific to the JE2 lineage a *tcaA::gfp* reporter plasmid

was constructed and electroporated into JE2, SH1000, Newman and a clinical CC22 isolate

EMRSA15. In each background serum, arachidonic acid and LL37 reduced *tcaA* expression

with the exception of LL37 and the EMRSA15 strain (Supplementary. Fig. 3). These findings

suggest that upon entry into the bloodstream and exposure to many of the antibacterial

components of serum, expression of the *tcaA* gene is induced and this sensitizes *S. aureus*

to killing by these components of serum.

Figure 6: TcaA confers increased sensitivity to teicoplanin. **(a)** The inactivation of the *tcaA* gene increases the MICs in three *S. aureus* backgrounds. **(b)** The clinical isolates with polymorphisms in the *tcaA* gene were on average less sensitive to teicoplanin (0.5μg/ml) than those with the wild type gene. **(c)** The most common *tcaA* SNP decreases the sensitivity of *S. aureus* to teicoplanin (0.5μg/ml). **(d)** The expression of the *tcaA* gene was quantified qRT-PCR in both subinhibitory and inhibitory concentrations of serum and teicoplanin. There was a dose dependent effect on *tcaA* induction for all concentrations used. (**e**) Exposure of JE2 to inhibitory concentrations of human serum resulted in an increase in WTA abundance in the bacterial cell wall, an effect not seen when the *tcaA* mutant was exposed to serum. (**f**) The WTA extractions were performed on the wild type and tcaA mutant in triplicate the relative density of the WTA quantified by densitometry. (**g**) Growth of JE2 in the presence of sub-inhibitory concentrations of teicoplanin (0.5 μg/ml) over a 24h period was quantified following pre-exposure to subinhibitory concentrations of human serum (2.5%), where preexposure to serum increased sensitivity to teicoplanin. (**h**) The ability of *S. aureus* to survive exposure to subinhibitory concentrations of serum (2.5%) following overnight growth in a subinhibitory concentration of teicoplanin (0.5 μg/ml), where preexposure to teicoplanin

increases the sensitivity to serum. The dots represent individual data points, the bars the
mean value, and the error bars the standard deviation. Significance was determined as
* <0.05, ** <0.01, *** <0.001, **** <0.0001.

The inactivation of tcaA *alters the antibiotic resistance profile of S. aureus.* There is emerging

evidence that WTA play a role in cell division and peptidoglycan biosynthesis[39]. In support of

this, it has been reported that a *tarO* mutant, devoid of WTA, confers sensitization to β-lactam

antibiotics due to mislocalization of the PBPs[40]. As the target for teicoplanin is peptidoglycan,

more specifically the terminal D-Ala-D-Ala residue of lipid II, we hypothesised that the

increased ligation of WTA into the cell wall of the TcaA producing strain may affect

peptidoglycan biosynthesis and assembly, and that this may be responsible for the change in

sensitivity to teicoplanin. To test this further, we compared the sensitivity of the wild type and

mutant to a range of peptidoglycan attacking antibiotics, where we found the TcaA producing

wild type strain to be more sensitive to vancomycin (as shown previously[38]), oritavancin, but

not dalbavancin (all alternative members of the glycopeptide class of antibiotics), and

ramoplanin, a glycolipodepsipeptide antibiotic (Supp. Fig. 4). However, TcaA production

conferred decreased sensitivity to oxacillin, a β-lactam antibiotic that targets penicillin binding

protein 2 and moenomycin a phosphoglycolipid antibiotic that inhibits the transglycosylase

PBP enzymes and prevents the formation of peptidoglycan polymers (Supp. Fig. 4). This

suggests that TcaA production affects the overall composition and structure of peptidoglycan

in the wildtype strain.

To examine whether the crosslinking of peptidoglycan is affected by TcaA production, we

tested the susceptibility of the wild type and *tcaA* mutant to lysostaphin, which is

a glycylglycine endopeptidase capable of cleaving the pentaglycine crosslinks of

peptidoglycan[41]. We found that lysostaphin cleaved the cell wall of the wild type strain more

slowly than for the *tca*A mutant (Fig. 7a). We next performed a triton X100 induced autolysis

assay where the wild type exhibited a slower rate of autolysis compared to the mutant,

suggesting that the cell wall is more robust due to higher levels of crosslinking when TcaA is

produced (Fig. 7b). Together these data suggest that the integrity of the peptidoglycan, in

particular its crosslinking, is increased when TcaA is produced. This provides a likely

explanation for why the mutant is more resistant to teicoplanin, reduced crosslinking in the cell

wall increases the number of off target D-ala D-ala which teicoplanin will bind to instead of

reaching lipid II in the membrane and causing disruption to this. With changes to the quantity

and composition of WTA and peptidoglycan associated with TcaA production, we sought to

visualise the cell wall using transmission electron microscopy. We observed a clear difference

in the density and structure of the outer layer of the bacterial cell wall between the wild type

and *tcaA* mutant, confirming a role for TcaA in the structural integrity of the *S. aureus* cell wall

(Fig. 7c, additional images in Supp. Fig. 5)).

Figure 7: TcaA alters the structure of the *S. aureus* cell wall. **(a)** The rate of lysis

in the presence of lysostaphin was assayed, where the strain producing TcaA was

lysed at a slower rate than the *tcaA* mutant. A *pbp4* mutant was included as a

control. Turbidity (OD_{600}) and area under the curves (AUCs) of the OD_{600} data over

time are presented. **(b)** The rate of autolysis of the bacteria in the presence of

Triton x100 was assayed where the TcaA producing strain lysed at a slower rate

relative to the *tcaA* mutant. An autolysin (*atl*) mutant was included as a control.

Autolysis data and area under the curves (AUCs) of the autolysis data over time

are presented. The dots represent individual data points, the bars the mean value,

and the error bars the standard deviation. For both **(a)** and **(b)** significance was

determined as * <0.05, ** <0.01 following area under the curve (AUC) analysis of

three biological replicates **(c)** Transmission electron micrograph (TEM) of a wild

type JE2 and *tcaA* mutant cell at two magnifications showing the smooth and

consistent density of the cell wall when TcaA is produced, compared to the rough

and patchy density of the mutant cell wall.

Once established, TcaA contributes positively to the development of bacteraemia. Our data

shows that expression of *tcaA* is induced upon exposure to serum, that this can sensitize the

bacteria to serum killing but also increase the abundance of WTA in the cell envelope. From

a pathogenicity perspective these activities would appear to be in conflict, with one making it

less likely for a bacterium to cause bacteraemia (increased serum sensitivity) and the other

making it more likely for a bacterium to cause bacteraemia (increased WTA in the cell

envelope[42,43]). To understand this apparent dichotomy we first examined whether TcaA

production only affected survival in serum, whereas in whole blood where other components

of the human immune system may be present (e.g. complement, antibodies and phagocytes)

that this effect may be lost. Using human blood from three donors, we demonstrated that TcaA

production increases the sensitivity of the bacteria to whole blood killing (Fig. 8a). We next

interrogated our human clinical data, where in addition to the 300 bacteraemia strains that

form the basis of this study we have an additional 176 *S. aureus* isolates, which has allowed

us to include more genetic diversity to this analysis (i.e. from CC22 (n=138), CC30 (n=162),

ST8 (n=132) and ST93 (n=44)). The source of these isolates was also available to us (i.e.

bacteraemia (n=341), SSTI (n=80) or carriage (n=55)), as well as their genome sequences

(Supplementary Table 1). We examined the distribution of nonsynonymous mutations in the

tcaA gene across these, where it was present in 6% of the bacteraemia isolates compared

with only 1.5% of non-bacteraemia isolates (n= 21/341 and 2/135 respectively; p = 0.03, in a

Fisher's Exact test). However, when we examined the 30-day mortality rates for the 113

bacteraemia patients for whom we had this clinical data, the mortality rate of the proportion of

patients infected with a *S. aureus* strain with the wild type *tcaA* gene was significantly higher

compared to those infected with a mutated *tcaA* gene (Fig. 8b & c) (2-tailed chi squared test:

p = 0.03). So, although the incidence of mutations in *tcaA* appear to be relatively rare, that

isolates with variant *tcaA* genes are enriched amongst those causing bacteraemia suggests

that a functional TcaA may limit the propensity of *S. aureus* to cause a bloodstream infection,

such that it acts as a selective force for mutants of this gene. However, once the bacteraemia

has become established, TcaA positively contributes to the infection process, likely due to the

increased abundance of WTA in the cell envelope, and the greater structural integrity of the

cell wall.

To experimentally test these human findings we utilised a mouse model of bacteraemia where

the tail vein of C57/Bl6 mice are inoculated with 2×10^7 colony forming units (CFU) of bacteria

to establish a bloodstream infection, where they seed from there into organs such as the liver

spleen and kidneys. We performed mixed infections, where the wild type and mutant bacteria

had to compete during the development of the infection within the mouse. The mice were

infected with equal quantities of both bacteria and the change in ratio of the bacteria in the

mouse blood, liver, spleen and kidney recorded after 48hr, which allowed the relative virulence

ratio (RVR) of each bacterium to be calculated[44]. The RVR of the TcaA producing wild type

strain was significantly higher than that of the mutant at a systemic level (Fig. 8d). As such,

while the TcaA associated self-sensitization of the bacteria was not evident here, possibly due

to the size of the inoculum level needed, once an infection has become established, TcaA

contributes positively to disease severity as demonstrated by the wild type strain outcompeting

the *tcaA* mutant.

Figure 8: TcaA contributes to increased disease severity in mice and humans. **(a)** Whole human blood survival. The wild type JE2 and tcaA mutant were incubated in human blood from three donors (each in biological triplicate) for 90 minutes and their relative survival was quantified. TcaA production renders that bacteria more susceptible to killing. The dots represent individual data points, the bars the mean value, and the error bars the standard deviation. **(b & c)** A graph and contingency table showing data from 113 patient with *S. aureus* bacteraemia. The 30-day mortality rate was significantly higher for those infected with a *S. aureus* strain with a wild type *tcaA* gene compared to those with a mutated *tcaA* gene. **(d)** Mice were injected intravenously with 100 μl of PBS containing an equal mixture of wild type and *tcaA* mutant (2 x 10⁷ CFUs of each). After 48 hrs, the ratio of the wild type and mutant were quantified and the relative virulence ratio calculated. In the blood and all organs tested the wild type bacteria had a competitive advantage over the mutant demonstrating its increased relative virulence. The dots represent individual data points (n=5 per group), the bars the mean value, and the error bars the standard deviation. Significance was determined as * <0.05, ** <0.01, and *** <0.001.

Discussion

To develop novel therapeutic approaches for infectious diseases, we need to understand how microorganisms cause disease and evade the host's immune system. To address this, we have applied a population-based approach to analyse the pathogenicity of *S. aureus* and identified seven genes that affect the bacteria's ability to survive exposure to human serum, the first step in the development of bacteraemia. The dissection of the molecular detail of how these genes affect serum survival is underway to determine their potential for therapeutic intervention. However, our focus here has been on the *tcaA* gene where we show that its expression is induced upon exposure to serum, that it is involved in the ligation or retention of

WTA in the cell wall, and that this renders the bacteria more susceptible to killing by both

AMPs and HDFAs. We have also found that TcaA is associated with changes to the

crosslinking of peptidoglycan, such that the bacteria are more susceptible to some antibiotics

(e.g., teicoplanin) and less susceptible to others (e.g., oxacillin). It is interesting to note that

our GWAS approach identified *tcaA* due to the presence of function altering SNPs amongst

clinical isolates. Our human data suggests that bacteraemia and/or the antibiotics commonly

used to treat it such as teicoplanin and vancomycin may impose a selective pressure on the

bacteria to mutate this gene, due to the decreased sensitivity this confers to serum killing and

to clinically relevant antibiotics. However, both our human and mouse data have confirmed,

that once bacteraemia has become established, TcaA contributes positively to disease

progression, likely due to its cell wall re-modelling activity. A graphical summary of this has

been provided in Fig. 9.

Figure 9: Graphical summary of our hypothesis for TcaA activity**.** Upon exposure to

serum the expression of *tcaA* is induced in the wild-type strain, resulting in an increased

abundance of WTA in the cell wall and less in the extracellular milieu. This decreases

the negative charge of the bacterial surface, rendering the bacteria more susceptible

to AMPs. In addition, there is less WTA released into the extracellular milieu to

sequester HDFAs, rendering the bacteria more susceptible to killing by these

molecules. This self-sensitizing activity of TcaA limits the ability of *S. aureus* to

establish a bloodstream infection. However, once established a strain with wild-type

TcaA can cause more severe disease. In *tcaA* mutants, there is lower abundance of

WTA in the cell wall and a higher abundance in the extracellular milieu. This increases

bacterial resistance to AMPs as it increases the charge of the bacterial surface,

resulting in less electrostatic attraction. The release of WTA into the extracellular milieu

also sequesters HDFAs, preventing them from reaching the bacterial surface. As a

result, a *tcaA* mutant is more likely to establish a bloodstream infection. However, this

infection is less likely to cause severe disease due to the importance of TcaA in

remodelling the cell wall for the full virulence of *S. aureus*. (Image created using

BioRender.com).

The *tcaA* gene is part of a three gene locus alongside the *tcaR* gene, which is predicted to

encode a transcriptional regulator, and the *tcaB* gene which is predicted to encode an efflux

pump[27]. This locus is also reported to be part of the *S. aureus* cell wall stimulon, following a

number of studies that monitored the response of *S. aureus* to cell wall attack[45,46]. It is

interesting to note that the *tcaA* gene was not associated with serum resistance for the CC30

collection of isolates, however, there were very few isolates with variation in this gene amongst

that collection such that they would have fallen below the minimum allele threshold. While

work to understand the activity of each of the proteins encoded by the *tca* locus is underway,

our data suggests that TcaA play a role in the ligation or retention of WTA within the cell wall,

perhaps alongside the LcpA protein[13]. How this is achieved is yet to be determined. TcaA may

play a direct role in ligating WTA to the cell wall; it may play a role in cell wall turnover and the

subsequent release of WTA and other cell wall macromolecules into the external environment;

or it may interact and subsequently interfere with other cell wall biosynthetic processes that

are responsible for WTA ligation. It is interesting to note that *tarK*, which is one of the other

six genes associated here with serum survival is also involved in the biosynthesis of WTA.

Additionally, *yfhO* which encodes an enzyme that glycosylates lipoteichoic acid (LTA) is also

involved in serum resistance[32], providing further evidence for the importance of teichoic acid

associated molecules to this aspect of the biology and pathogenicity of *S. aureus*. For TcaR,

we hypothesise that it likely controls the transcription of the co-transcribed *tcaA* and *tcaB*

genes, and may directly respond to the external stimulus of teicoplanin and serum as reported

here. What is less clear is what role TcaB plays, as efflux pumps are more typically associated

with increasing the resistance of a bacterium to an antibiotic, or other toxic molecules.

In addition to *tcaA* this study has identified one other locus (*yfhO*) that negatively affects serum

survival, and added a further five loci (*tarK, gntR, ilvC, arsB*, and *pdhD*) to the list of genes

that contribute positively to serum survival by protecting the bacteria from HDFAs and AMPs.

This leads us to the questions of why *S. aureus* has loci in apparent conflict with regards to

the survival of the bacteria during bacteraemia. It is perhaps through the consideration of the

other ways in which *S. aureus* interact with humans that we may understand this. The vast

majority of interactions that occur between *S. aureus* and humans is as a commensal in the

human nose from which it can readily transmit. The five identified protective genes may

therefore also protect the bacteria during colonisation, which would confer a selective

advantage, and it is just an unfortunate coincidence that they also confer increased survival

in serum during the development of bacteraemia. While the focus of this work has been on

the role of TcaA in serum, it is also worth considering what role it may have during colonisation

of the nose, which involves the adherence of the bacteria to the nasal epithelium, to which

WTA has a well-established role. It is therefore possible that TcaA enhances the ability of *S.*

aureus cells to colonise the nose by retaining WTA within the cell wall. However, the role of

TcaA in responsiveness to cell wall attack is as yet unclear, but perhaps it contributes to the

competitiveness of *S. aureus* within the nasal microbiome in response to cell wall damage

inflicted by competitors.

The evolution of the virulence of a pathogen is heavily dependent upon their mode of

transmission, which for opportunistic pathogens raises interesting challenges. If within-host

invasiveness limits between-host fitness by preventing them from transmitting to a new host,

there are likely selective pressures acting on bacterial populations to evolve means of limiting

their ability to cause invasive disease. However, if increased invasiveness facilitates enhanced

within-host fitness, as the bacteria multiply and spread throughout the body, the 'short-sighted

evolution of virulence' hypothesis proposed by Levin and Bull[47] could apply and may explain

situations like *S. aureus* bacteraemia. We propose that TcaA may represent a system at the

cusp of both these scenarios, in that it responds to cell wall attack by serum by making itself

more susceptible to killing by serum, thereby potentially limiting its ability to cause

bacteraemia, which could have long-term between host benefits. However, once established

in the bloodstream and TcaA has remodelled the cell wall, the short-sighted view of virulence

hypothesis applies where the bacteria survive and thrive, allowing them to cause a successful

bacteraemia.

Entry into the bloodstream to establish an infection has long been considered a significant

bottle neck for the bacteria, but until now, the stringency of this bottleneck was believed to be

a result of the onslaught of all the immune mechanisms present in blood. Here we suggest

that *S. aureus* may contribute to this bottleneck by responding to serum induced cell wall

damage by further sensitizing itself to this attack as part of the pathway towards increasing

their pathogenic capabilities by remodelling the cell wall. Our findings presented here opens

up an entirely novel aspect to the biology of a major human pathogen and brings into question

the appropriateness of the term 'opportunistic' when we refer to such highly effective bacterial

pathogens.

Materials and Methods

Bacterial Strains and Growth Conditions

A list of the genetically amenable bacterial strains and mutants can be found in Table 2, and

the clinical strains are listed in Supplementary Table 1. All strains were cultured in Tryptic

soy broth (TSB) for 18 h at 37°C with shaking. Nebraska transposon mutant library (NTML)[23]

mutants were selected for using erythromycin (5 μg/ml). For the complemented pRMC2[45]

strains, chloramphenicol (10 μg/ml) and anhydroustetracycline (100 ng/ml) were added to

the media where indicated.

Table 2. Strains used in this study.

Strain	Description	Reference
JE2	USA300; CA -MRSA, type IV SCCmec; lacking plasmids p01 and p03; wild -type strain of the NTML	25
JE2 *tcaA:tn*	*tcaA* transposon mutant in JE2	25
JE2 *tcaA:tn* pRMC2	*tcaA* transposon mutant in JE2 transformed with empty pRMC2 vector	This study
JE2 *tcaA:tn* p*tcaA*	*tcaA* transposon mutant complemented with *tcaA* gene housed in pRMC2 expression plasmid	This study
JE2 *tarK:tn*	*tarK* transposon mutant in JE2	25
JE2 *tarK:tn* pRMC2	*tarK* transposon mutant in JE2 transformed with empty pRMC2 vector	This study

JE2 *tarK:tn* p*tarK*	*tarK* transposon mutant complemented with *tarK* gene housed in pRMC2 expression plasmid	This study
JE2 *gntR:tn*	*gntK* transposon mutant in JE2	25
JE2 *gntR:tn* pRMC2	*gntK* transposon mutant in JE2 transformed with empty pRMC2 vector	This study
JE2 p*gntR*	*gntK* transposon mutant complemented with *gntK* gene housed in pRMC2 expression plasmid	This study
JE2 *ilvC:tn*	*ilvC* transposon mutant in JE2	25
JE2 *ilvC:tn* pRMC2	*ilvC* transposon mutant in JE2 transformed with empty pRMC2 vector	This study
JE2 *ilvC:tn* p*ilvC*	*ilvC* transposon mutant complemented with *ilvC* gene housed in pRMC2 expression plasmid	This study
JE2 *arsB:tn*	*arsB* transposon mutant in JE2	25
JE2 *arsB:tn* pRMC2	*arsB* transposon mutant in JE2 transformed with empty pRMC2 vector	This study
JE2 *arsB:tn* p*arsB*	*arsB* transposon mutant complemented with *arsB* gene housed in pRMC2 expression plasmid	This study
JE2 *yfhO:tn*	*yfhO* transposon mutant in JE2	25
JE2 *yfhO:tn* pRMC2	*yfhO* transposon mutant in JE2 transformed with empty pRMC2 vector	This study
JE2 *yfhO:tn* p*yfhO*	*yfhO* transposon mutant complemented with *yfhO* gene housed in pRMC2 expression plasmid	This study
JE2 *pdhD:tn*	*pdhD* transposon mutant in JE2	25

JE2 *pdhD:tn* pRMC2	*pdhD* transposon mutant in JE2 transformed with empty pRMC2 vector	This study
JE2 *pdhD:tn* p*pdhD*	*pdhD* transposon mutant complemented with *pdhD* gene housed in pRMC2 expression plasmid	This study
RN4220	Restriction deficient mutant of 8325-4	48
DC10B	*Δdcm* mutant in *E. coli* DH10B (K12)	37
Mach1	One Shot™ Mach1™ T1 Phage-Resistant Chemically Competent *E. coli.*	Commercially available
SH1000	MSSA, laboratory strain, 8325-4 with a repaired rsbU gene; SigB positive.	49
SH1000 *tcaA::tn*	*tcaA* transposon mutant in SH1000	This study
Newman	MSSA, laboratory strain isolated from human infection	50
Newman *tcaA::tn*	*tcaA* transposon mutant in Newman	This study
tcaA::tn p*tcaA*SNP	*tcaA* transposon mutant complemented with *tcaA* gene harbouring F290S SNP housed in pRMC2 expression plasmid	This study
mprF::tn	*mprF* transposon mutant in Newman	25
Δ*lcpA*	In frame unmarked deletion of *lcpA* generated by allelic exchange	13
LAC	CA-MRSA USA300 type IV SCCmec	51
Δ*tarO*	*tarO* KO mutant in LAC	52
JE2 pSB2019:*tcaA*	JE2 transformed with the pSB2019:*tcaA* reporter fusion	This study
SH1000 pSB2019:*tcaA*	SH1000 transformed with the pSB2019:*tcaA* reporter fusion	This study

Newman pSB2019:*tcaA*	SH1000 transformed with the pSB2019:*tcaA* reporter fusion	This study
EMRSA15 pSB2019:*tcaA*	SH1000 transformed with the pSB2019:*tcaA* reporter fusion	This study

Serum & Blood Survival Assay

Normal human serum was prepared from blood obtained from 8 healthy volunteers using

Serum CAT tubes (BD). On the day of collection the blood was allowed to clot for 30 mins at

room temperature followed by 1h incubation on ice. Serum was extracted following two

rounds of centrifugation at 700 x g at 4°C for 10 min. Serum from individual donors were

pooled, aliquoted and immediately stored at -80°C, where each aliquot underwent only one

round of freeze and thaw. The bacteria were grown overnight, and their density normalised

to an OD_{600nm} of 0.1 and 20 µl used to incubate in 180 µl 10% pooled human serum (diluted

in PBS) for 90 min at 37°C with shaking. Serial dilutions were plated on tryptic soy agar

(TSA) to determine CFUs. The same number of bacterial cells inoculated into PBS, diluted,

and plated acted as a control. Survival was determined as the percentage of CFU in serum

relative to the PBS control. Relative survival was determined through normalisation to JE2,

which is presented ats 100% in the graphs. The survival of each bacterial isolate was

measures in triplicate and the mean of these datal presented here.

GWAS

Genome-wide association mapping was conducted using a generalized linear model, with

serum survival as the quantitative response variable. We accounted for bacterial population

substructure by adding to the regression model the first two component from a *principal*

component decomposition of SNP data for each set of clinical samples (CC22 and CC30).

The first two components accounted for 32% and 40% of the total variance for CC22 and

CC30, respectively. In both cases, three distinct clusters were identified. We further

considered a third model where we used cluster membership as covariates in our regression

model, where clusters were defined using K-means clustering analysis (setting K=3); this,

however, yielded identical results to the one based on PCA components. In total, 2066

(CC22) and 3189 (CC30) unique SNPs were analysed, the majority of which were

subsequently filtered out for exhibiting a minor allele frequency (maf) of <0.03, reducing the

data to 378 and 1124 SNPs, respectively. The P-values reported in Tables 1a and b are not

corrected for multiple comparison, however the Sidak method was used to correct for

multiple comparisons and both significance thresholds are indicated in the Manhattan plots.

Genetic manipulation of bacterial strains

Wildtype genes were amplified by PCR from JE2 genomic DNA using the primers shown in

Table 3 and KAPA HiFi polymerase (Roche). The PCR product was cloned into the

tetracycline inducible plasmid pRMC2[26] using *Kpn*I and *Sac*I restriction sites and T4 DNA

ligase (NEB). This was transformed into RN4220 and eventually into the respective NTML

mutants through electroporation.

Table 3. PCR primers used in this study.

Primer name	Primer sequence 5'-3'
tcaA FW	ataagcttgatggtaccaattgaagaggaacagaattgag
tcaA RV	tgaattcgagctcagatcatgacacatgcatcttatttaag
tarK FW	ataagcttgatggtacctttaataaaccgatgggggg
tarK RV	tgaattcgagctcagatgcaagtaatatatgccaacattag
gntK FW	ataagcttgatggtaccgtatgtcgtgtaatgaaggagtg
gntK RV	tgaattcgagctcagatttctttctttcattttaattcaacc
ilvC FW	ataagcttgatggtacccgaaacacaaaatatttaattatttgg
ilvC RV	tgaattcgagctcagattaacaactcctcattgtaggtctatc
arsB2 FW	ataagcttgatggtaccgtcataaccttaagactggtgaatgc

672	*arsB2* RV	tgaattcgagctcagatgagttgcctgtacatataaaataaattg
673	*yfhO* FW	ataagcttgatggtacctgaaagttgaggataatgttgtg
674	*yfhO* RV	tgaattcgagctcagatgtagcaacattatttttgtcttgc
675	*pdhD* FW	ataagcttgatggtaccgaattattattaatggaggggtaaaac
676	*pdhD* RV	tgaattcgagctcagatgcatgtcacatgttaacgatatctc

Site directed mutagenesis of the *tcaA* gene

Site directed mutagenesis was used to mutate the WT *tcaA* gene (Phe290) to the SNP *tcaA*

(Ser290). Primers (5'' ttttcaagttTcaaaacgtatggtc 3", 5" atacgttttgaAacttgaaaatgcc 3") were

designed to have a complementary overlap of 21 nucleotides at the 5' end, with the

nucleotide to be mutated at the centre of the overlap (the primers had non complementary

regions at the 3' end to facilitate annealing on the template). The p*tcaA* template was diluted

to 33 ng/µl, and 1ul was amplified using these primers and Phusion DNA polymerase. The

PCR product was subsequently checked by agarose gel electrophoresis and was treated

with 1 µl of DpnI directly in the PCR mix for 1 h at 37°C. Then 1 µl of the mixture was used to

transform One Shot™ Mach1™ T1 (ThermoFisher) competent cells by heat shock then into

RN4220 and finally *tcaA::tn* by electroporation.

Phage transduction of the *bursa aurealis* transposon

The *tcaA bursa aurealis* transposon was phage transduced using the *S. aureus* phage

Φ11[53]. The phage lysate of the donor strain (*tcaA::tn)* was prepared as follows: The donor

strain was grown overnight in TSB for 18 h at 37°C with shaking. Two hundred microlitres of

overnight culture was added to 25 ml of TSB containing 250 µl 1M $MgSO_4$, 250 µl 1M $CaCl_2,$

100 µl of Φ11 and left at 37°C with shaking until complete lysis was observed. The culture

was subsequently pelleted at 12000 x *g* for 3 min. The supernatant containing the lytic

phage was filtered sterilised and stored at 4°C. Next, recipient strains (Newman and

SH1000) were grown overnight in 20 ml of LK broth (1% Tryptone 0.5% Yeast Extract,

0.7% KCl) and pelleted at 2500 x *g* for 10 min. The supernatant was removed, and the

pellets resuspended in 1 ml of fresh LK broth. Two reactions were then set up, one

containing phage and one as a non-phage control. To the tube containing the phage add:

250 µl of the recipient strain (SH1000 or Newman), 250 µl of phage lysate and 750 µl of LK

broth containing 10 mM $CaCl_2$. To the tube containing the non-phage control add: 250 µl of

the recipient strain (SH1000 or Newman) and 1 ml od LK broth containing 10 mM $CaCl_2$.

Both reactions were incubated for 25 min at 37°C without shaking followed by 15 min at

37°C with shaking. Five hundred microlitres of ice cold 20 mM sodium citrate was added

followed by centrifugation at 10000 x *g* for 10 min. Pellets were resuspended in 500 µl of 20

mM sodium citrate and incubated on ice for 2 h. One hundred microlitres of culture was

plated out on TSA containing 5 µg ml^{-1} erythromycin to select for the transposon. For the

SH1000 phage transduction 100 µl was also plated out on oxacillin TSA plates as an added

control to ensure there was no contamination of the phage lysate with JE2 *tcaA::tn.*

Growth inhibition and MIC assays

To determine the relative inhibition of growth of the wild type and tcaA mutant by antibiotics

a broth microdilution method was used[54]. Briefly, overnight cultures were normalised to an

OD_{600nm} of 0.05 in cation adjusted Mueller Hinton broth (MHB++) and 20 µl of resultant

suspension used to inoculate 180 µl of fresh MHB++ containing a 1:2 dilution series of the

respective antimicrobial agent. The ability of the bacteria to survive these was determined by

quantifying bacterial growth (OD_{600nm}) using a CLARIOstar plate reader (BMG Labtech).

Antimicrobial peptide susceptibility

Antimicrobial peptide susceptibility Human neutrophil defensin-1 (hNP-1) (AnaSpec

Incorporated, California, USA) and LL-37 (Sigma) susceptibility assays were performed as

described previously[55]. Briefly, overnight cultures were normalised to an OD_{600nm} of 0.1 and

incubated with 5 µg/mL of hNP-1 or LL-37 for 2 h at 37 °C. Serial dilutions were plated on

tryptic soy agar (TSA) to determine CFUs. The same number of bacterial cells inoculated

into PBS, diluted, and plated acted as a control. Survival was determined as the percentage

of CFU on exposure to either LL-37 or HNP-1 relative to the PBS control. Relative survival

was determined through normalisation to JE2.

mRNA extraction

Overnight cultures were back diluted to an OD_{600nm} of 0.05 in 50 ml of fresh TSB and grown

to an OD_{600nm} of 2. Two hundred micro litres of either PBS, 25% serum (final concentration

2.5%, sub-inhibitory), 5 µg/ml teicoplanin (final concentration 0.5 µg/ml, sub-inhibitory),

100% serum (final concentration 10%, inhibitory), or 100 µg/ml teicoplanin (final

concentration 10 µg/ml, inhibitory) was added to 1.8 ml of bacterial culture and incubated for

20 min. RNA was extracted by Quick-RNA Fungal/Bacterial Miniprep Kit (Zymo Research)

according to the manufacturer's instructions. RNA integrity was checked by running 5 µl

aliquot of the RNA on a 1% agarose gel and observing the intensity of the ribosomal RNA

(rRNA). RNA samples were treated by TURBO™ DNase (Invitrogen) to eliminate any

genomic DNA contamination. To verify that the samples were free from any DNA

contamination, RNA samples were subjected to RT-qPCR alongside a no template control

and 2.5 ng of a known JE2 genomic DNA and threshold rates compared.

qRT-PCR

To quantify the expression of the *tcaA* RT-qPCR was performed using the housekeeping

gyrB gene as a control. Complementary DNA (cDNA) was generated from mRNA using

qScript® cDNA Synthesis Kit (Quantabio). Following the manufacturers protocol the cDNA

was used as a template for the RT-qPCR reaction. Primers used were as follows *gyrB* FW 5'

ggtgactgcattagatgtaaac 3', *gyrB* RV 5' ctgcttctaaaccttctaatacttgtatttg 3', *tcaA* FW 5'

tagtttgcgcttcaggtg 3', *tcaA* RV 5' tgtggacataaatttgatagtcgtc 3'. The RT-qPCR reaction was

performed as follows: 10 µl 2x KAPA SYBR Mix, 1 µl of 10 µM forward primer, 1 µl of 10 µM

reverse primer, 5 µl cDNA and RNase-free water up to a total of 20 µl volume. The RT-

qPCR was performed on a Mic qPCR cycler (bio molecular systems) and the cycling

conditions consisted of an initial denaturation step of 95°C for 2 min, followed by 40 cycles of

two step cycling: 95°C 15 s, 60°C 1 min. RT-qPCR was carried out in technical triplicate for

each sample and 3 biological repeats. The ratio of *tcaA* to *gyrB* transcript number was

calculated using the using the $2^{-(\Delta\Delta Ct)}$ method[56].

WTA preparations

Crude WTA from murein sacculi was extracted for analysis by PAGE using adaptations of a

previously described methodology[57,58]. Overnight cultures were washed once in buffer 1 (50

mM MES, pH 6.5) followed by centrifugation at 5,000 g. Cells were resuspended in buffer 2

(4% [wt/vol] SDS, 50 mM MES, pH 6.5) and boiled for 1 h. Sacculi were centrifuged at

5,000g and washed once in buffer 1, once in buffer 2, once in buffer 3 (2% NaCl, 50 mM

MES, pH 6.5), once more in buffer 1 and finally resuspended in digestion buffer (20 mM Tris-

HCl pH 8.0, 0.5% [w/v] SDS). To the digestion buffer suspension, 10 µl of proteinase K

solution (2 mg/ml) was added and incubated on a heat block 50°C for 4 h at 1,400 rpm.

Sacculi were centrifuged at 16,000 g and washed once in buffer 3, followed by 3 washes in

dH$_2$O to removed SDS. Sacculi were responded in 0.1M NaOH and incubated for 16 h at

room temperature at 1,400 rpm. Following the incubation, the sacculi were centrifuged at

16,000g, leaving the teichoic acids in the supernatant. 250 µl of 1 M Tris-HCL pH 6.8 was

added to neutralise the NaOH and stored at -20°C.

WTA was also precipitated from the supernatant of overnight culture by adding 3 volumes of

95% ethanol and incubation at 4°C for 2 h. Precipitated material was separated by

centrifugation at 16,000 g for 15 min, washed once in 70% ethanol and resuspended in 100

mM Tris-HCL (pH 7.5) containing 5 mM $CaCl_2$, 25 mM $MgCl_2$, DNase (10 µg/ml) and RNase

(50 µg/ml) and incubated for 3 h at 37°C. The enzymes were heat inactivated at 95°C for 3

min. The supernatant WTA preparations were similarly stored at -20°C before being

analysed by PAGE.

WTA PAGE

WTA preparations were separated on tricine polyacrylamide gels using a BioRad tetra cell

according to a previously described method[58]. The gels were separated at 4°C using a

constant amperage of 40 mA under constant stirring until the dye front reached the bottom.

Gels were washed 3 times in MilliQ H_2O followed by staining in 1 mg/ml Alcian blue

overnight. Gels were subsequently destained in in MilliQ H_2O, until the WTA became visible

and finally imaged.

Arachidonic acid supernatant conditioning

Overnight cultures of JE2, *tcaA*, and *lcpA* mutants were pelleted, and the supernatant saved.

Growth inhibition assays were subsequently performed by adding 20 µl of OD_{600nm} 0.1

bacterial suspension to 180 µl 10% overnight supernatant diluted in fresh MHB++ containing

400µM of arachidonic acid (Sigma). Purified WTA extracts were also added to the growth

media at a final concentration of 2%. The ability of the bacteria to survive the arachidonic

acid was determined by quantifying bacterial growth (OD_{600nm}) following 24 h at 37°C using a

CLARIOstar plate reader (BMG Labtech).

Cytochrome C binding assay

A cytochrome c binding assay was used to measure the relative surface charge of the

bacteria[59]. Briefly, overnight cultures were normalized to an OD_{600nm} of 8. The bacterial

suspensions were washed twice in MOPS buffer (20 mM pH 7.0) and finally resuspended in

200 µl of MOPS buffer. Samples were then combined with 50 µl of cytochrome C (equine

heart Sigma, 2.5 mg/ml in MOPS buffer) and incubated for 10 min at room temperature.

Finally, samples were pelleted (16,000 × g for 1 min) and 200 µl of supernatant read for

absorbance at Abs_{530nm} using a SUNRISE Tecan microplate reader.

Construction of a TcaA reporter fusion

A *tcaA* promoter-*gfp* fusion (pSB2019:*tcaA*) was constructed to determine expression of

tcaA in response to several antimicrobial components of serum. A region of 289 bp

upstream of the *tcaA* gene[60] was amplified by PCR from JE2 genomic DNA using the

primers *tcaA_promoter* FW 5'- atatgaattcagtattagaagtcatcaatca -3' and *tcaA_promoter* RV

5'-atatccccgggtttcacctcaattctgttcct-3'. The pSB2019 vector was prepared by digesting

pSB2031[61] from a previous study with EcoRI and SmaI to remove the RNAIII P3 promoter.

The *tcaA* promoter region was also digested with EcoRI and SmaI and ligated into pSB2019

using T4 DNA ligase (NEB). This was transformed into DC10B[37], a strain of *E. coli* which

mimics the adenine methylation profile of some *S. aureus* clonal complexes (including CC8

and CC22). The plasmid was then extracted and transformed into final strains via

electroporation.

TcaA induction assay

S. aureus strains carrying the pSB2019:*tcaA* plasmid were grown overnight in TSB with

10ug/ml chloramphenicol. Each strain was then normalized to an OD_{600nm} of 0.05 in fresh

TSB and subcultured to an OD_{600nm} of 0.5-0.6. Cultures were washed in PBS and

concentrated to an OD_{600nm} of 1 in PBS. 100 ul of bacteria was combined with 100 ul of the

appropriate antimicrobial compound (arachidonic acid or LL-37) in a black 96-well plate.

GFP Fluorescence (485_{nm} excitation/520_{nm} emission/1000 gain) was measured in a

PHERAstar FSX plate reader (BMG Labtech) over 2.5 h (readings every 30 min with

200 rpm shaking).

Lysostaphin turbidity assay

Overnight cultures were back diluted to an OD_{600nm} of 0.05 in 20 mL of fresh TSB and grown

to an OD_{600nm} 0.5-0.7. Cultures were normalised to give 1 ml of OD_{600nm} 0.6. Lysostaphin was

subsequently added to give a final concentration of 0.5 µg/mL. 200 µL of each strain was

then added to a 96-well plate and incubated for 3 h at 37°C in a CLARIOstar plate reader.

OD_{600nm} readings were taken every 10 min (500 rpm shaking before readings). Values were

blank corrected according to 200 µL of PBS. The PBP4*::tn* mutant was included as a control.

Triton-X100 induced autolysis assay

Overnight cultures were back diluted to an OD_{600nm} of 0.05 in 20 mL of fresh TSB and grown

to an OD_{600nm} 0.5-0.7. Cells were washed once in ice cold water (13,000 rpm 1 min) and

resuspended in autolysis buffer (water containing 0.1% Triton X-100) to give an OD_{600nm} of 1.

200 µL of each strain was then added to a 96-well plate alongside 200 µL of strains

resuspended in water as a negative control and 200 ul of water as a blank. The *atl::tn* strain

was used as a positive control. Strains were grown for 6 h at 37°C in a CLARIOstar plate

reader. OD_{600nm} readings were taken every 30 min (500 rpm shaking before readings). The

rate of autolysis was calculated as follows; (OD*t0*-OD*tn*/OD*t0*).

Transmission electron microscopy

S. aureus strains JE2 and the TcaA mutant were pelleted in microfuge tubes and fixed by

resuspending in 2.5% glutaraldehyde in cacodylate buffer (pH 7.3) kept at fridge

temperatures until fixation was complete and then stored in the fridge. Pellets were

resuspended in a BSA/glutaraldehyde gel at 10- 20°C which sets after the cells were re-

centrifuged into a pellet. The pellets were postfixed with osmium ferrocyanide/osmium

tetroxide mix in cacodylate buffer, *en bloc* stained with uranyl acetate (2%) and Walton's

lead aspartate solutions, prior to dehydration in ethanol and infiltration with propylene oxide

and Epon resin mix. Embedded blocks were polymerised for 24-48 hours at 60°C. Sections

(70nm) were cut on a Leica UC7 ultramicrotome and imaged using a FEI Tecnai T12

microscope.

In vivo **intra-venous challenge model**

For the mixed infection bacteraemia model, female, 7-8 week-old, C57BL/6NCrl mice

(Charles River Laboratories) were used. Bacteria were prepared as described previously[62].

Briefly, 1:100 dilutions of overnight bacterial cultures were grown to mid-log phase (~2 h) in

TSB with shaking at 180 rpm at 37 °C. Bacteria were then harvested, washed with PBS and

adjusted spectrophotometrically at OD_{600nm} to obtain the desired bacterial concentration of

$\sim 4.8 \times 10^8$ CFU/mL. A 50:50 mixture of JE2 and *tcaA::tn* were made and 100 μl was injected

intravenously via the tail vein. The final CFU for each strain per mouse was $\sim 2.4 \times 10^7$ CFU.

After 48 hours, terminal cardiac bleeds were performed on anesthetized mice and whole

blood was transferred into heparin tubes (Sarstedt) and mixed to prevent coagulation. Mice

were then euthanized by CO_2 inhalation. Organ homogenates of spleens, livers and kidneys

were performed as previously described[63] and serial dilutions of each sample were plated

onto TSA plates with and without erythromycin (5 μg/ml) for bacterial enumeration. The CFU

of JE2 was calculated by subtracting CFU on TSA with erythromycin (*tcaA::tn* only) from

CFU on TSA plates (JE2 and *tcaA::tn*).

Ethics statement

C57/Bl6 mice were bred in-house in Trinity College Dublin and C57BL/6NCrl mice were

purchased from Charles River Laboratories. Mice were housed under specific pathogen-free

conditions at the Trinity College Dublin Comparative Medicines unit and maintained under

pathogen-free conditions in an Association for Assessment and Accreditation of Laboratory

Animal Care (AAALAC)-accredited animal facility in NIAID, NIH, respectively. All mice were

female and used at 6–8 weeks. All animal experiments were conducted in accordance with

the recommendations and guidelines of the health product regulatory authority (HPRA), the

competent authority in Ireland and in accordance with protocols approved by Trinity College

Dublin Animal Research Ethics Committee and according to the regulations of NIAID's

Division of Intramural Research Animal Care and Use Committee (DIR ACUC), animal study

proposal LB1E. For the human serum collection, all healthy volunteers provided written

informed consent and all methods and experimental protocols were carried out in

accordance with the recommendations of the University of Bath, Research Ethics Approval

Committee for Health. The present study was approved by the University of Bath, Research

Ethics Approval Committee for Health [reference: EP 18/19 108].

Statistics

Paired two-tailed student t-test or One-way ANOVA (GraphPad Prism v9.0) were used to

analyse the observed differences between experimental results. A p-value <0.05 was

considered statistically significant and the Sidak method was used to correct for multiple

comparisons.

Acknowledgements: We would like to thank Prof. Mark Jepson, Chris Neal and Lorna

Hogdson at the Wolfson Bioimaging Suite. We would also like to thanks Eric Skaar and William

beavers for providing the *lcpA* mutant.

Funding information: EJAD was funded on a studentship provided by the School of Cellular

and Molecular medicine, University of Bristol. NP is funded on a BBSRC SWBio DTOP Phd

Studentship. DA is funded by a PhD studentship awarded by the Saudi Arabian Cultural

Bureau. RCMs and RMM are both Wellcome Trust Investigators (Grant reference numbers:

212258/Z/18/Z and 202846/Z/16/Z). MO is supported by the Intramural Research Program of

the National Institutes of Allergic and Infectious Diseases (NIAID), US National Institutes of

Health (NIH), project number ZIA AI000904. ML acknowledges funding from the Academy of

Medical Sciences (SBF006\1023).

Conflict of Interests: we the authors declare no conflict of interest associated with the work

described in this manuscript

Author Contributions: EJAD developed the methodology, conceptualized and performed the

experiments, analysed data, provided supervisory support and contributed to writing the

manuscript. NP developed methodology and performed experiments. TB developed

methodology, performed experiments and provided supervisory support. DA performed

experiments. ML provided materials, analysed data and provided supervisory support. MR

developed methodology, analysed data and contributed to writing the manuscript. GYCC,

RCH, RL performed serum resistance experiments with mouse serum and murine

bacteraemia experiments and GYCC analyzed data. MO provided supervisory support and

provided materials. EOB performed experiments and analysed data, RMM provided materials,

analysed data and contributed to writing the manuscript. RCM conceptualized the project,

provided materials, provided supervisory support, analysed data and contributed to writing the

manuscript.

REFERENCES

1. Gordon R.J. & Lowy F.D. Pathogenesis of methicillin-resistant Staphylococcus aureus

infection. *Clin Infect Dis* **5**, S350-9 (2008)

2. Lowy F.D. *Staphylococcus aureus* infections. *N Engl J Med* **339**, 520-32 (1998).

3. Powers M.E. &, Bubeck Wardenburg J. Igniting the fire: Staphylococcus aureus virulence

factors in the pathogenesis of sepsis. *PLoS Pathog.* **10**, e1003871 (2014).

4. Abdelnour A., Arvidson S., Bremell A., Ryden C. & Tarkowski A. The accessory gene

regulator (agr) controls Staphylococcus aureus virulence in a murine arthritis model. *Infect*

Immun **61**, 3879-85 (1993).

5. Cheung A.L. *et al*. Diminished virulence of a sar-/agr- mutant of Staphylococcus aureus in

the rabbit model of endocarditis. *J Clin Invest* **94**, 1815-22 (1994).

6. Rose H.R. *et al*. Cytotoxic Virulence Predicts Mortality in Nosocomial Pneumonia Due to

Methicillin-Resistant Staphylococcus aureus. *J Infect Dis* **211,** 1862-74 (2015).

7. Laabei M. *et al*. Evolutionary Trade-Offs Underlie the Multi-faceted Virulence of

Staphylococcus aureus. *PLoS Biol*. **13**, e1002229 (2015).

8. Kaash A.J. *et al.* Staphylococcus aureus bloodstream infection: a pooled analysis of five

prospective, observational studies. *J Infect* **68,** 242-51 (2014).

9. Bai A.D. *et al.* Staphylococcus aureus bacteraemia mortality: a systematic review and

meta-analysis. *Clin Microbiol Infect.* **28**,1076-1084 (2022).

10. https://www.gov.uk/government/statistics/mrsa-mssa-gram-negative-bacteraemia-

and-cdi-independent-sector-annual

11. Levy O. Antimicrobial proteins and peptides of blood: templates for novel antimicrobial

agents. *Blood* **96**, 2664-72 (2000).

12. Kenny J.G. *et al*. The Staphylococcus aureus Response to Unsaturated Long Chain Free

Fatty Acids: Survival Mechanisms and Virulence Implications. *PLoS One* **4**, e4344 (2009).

13. Beavers W.N. *et al*. Arachidonic Acid Kills Staphylococcus aureus through a Lipid

Peroxidation Mechanism. mBio **10**, 5 (2019).

14. Das U.N. Arachidonic acid and other unsaturated fatty acids and some of their metabolites

function as endogenous antimicrobial molecules: A review. *J Advanc Res* **11** 57-66 (2018).

15. Laabei M. *et al.* Predicting the virulence of MRSA from its genome sequence. Genome

Res. **24,** 839-49 (2014).

16. Recker M. *et al.* Clonal differences in Staphylococcus aureus bacteraemia-associated

mortality. *Nat Microbiol* **2**, 1381-1388 (2017).

17. Yokoyama M., *et al.* Epistasis analysis uncovers hidden antibiotic resistance-associated

fitness costs hampering the evolution of MRSA. *Genome Biol* **19**, 94 (2018).

18. Stevens E. *et al*. Targeted control of toxin production by a mobile genetic element in

Streptococcus pneumoniae. *Microb Genom* **8**, 000784 (2022).

19. Altwiley D. *et al.* A Functional Menadione Biosynthesis Pathway is Required for Capsule

Production by Staphylococcus aureus. *Microbiology* **167**, 001108 (2021).

20. Douglas E.J.A., Duggan S., Brignoli T. &, Massey RC. The MpsB protein contributes to

both the toxicity and immune evasion capacity of Staphylococcus aureus. *Microbiology* **167**,

001096 (2021).

21. Peschel A. *et al.* Inactivation of the dlt operon in Staphylococcus aureus confers sensitivity

to defensins, protegrins, and other antimicrobial peptides. *J Biol Chem* **274**, 8405-10 (1999).

22. Ernst C.M. & Peschel A. Broad-spectrum antimicrobial peptide resistance by MprF-

mediated aminoacylation and flipping of phospholipids. *Mol Microbiol* **80**, 290-9 (2011).

23. Hwang-Soo J. & Otto M. Mechanisms of resistance to antimicrobial peptides in

staphylococci. *Biochimica et Biophysica Acta (BBA) – Biomembranes* **1848**, 3055-3061

(2015).

24. Kohler T., Weidenmaier C. & Peschel A. Wall Teichoic Acid Protects Staphylococcus

aureus against Antimicrobial Fatty Acids from Human Skin. *J Bacteriol* **191**, 4482–4484

(2009).

25. Fey P.D. *et al.* A genetic resource for rapid and comprehensive phenotype screening of

nonessential Staphylococcus aureus genes. *mBio* **4**, e00537-12 (2013).

26. Corrigan R.M. & Foster T.J. An improved tetracycline-inducible expression vector for

Staphylococcus aureus. *Plasmid* **61**, 126-9 (2009).

27. Brandenberger M. *et al.* Inactivation of a novel three-cistronic operon tcaR-tcaA-tcaB

increases teicoplanin resistance in Staphylococcus aureus. *Biochim Biophys Acta.* **1523**,135-

9 (2000).

28. Xia G., Kohler T. & Peschel A. The wall teichoic acid and lipoteichoic acid polymers of

Staphylococcus aureus. *Int J Med Microbiol* **300**,148-54 (2010).

29. Fujita Y. & Fujita T. The gluconate operon gnt of Bacillus subtilis encodes its own

transcriptional negative regulator. *Proc Natl Acad Sci U S A* **84**, 4524-8 (1987).

30. Li K.H. *et al.* Biological Functions of ilvC in Branched-Chain Fatty Acid Synthesis and

Diffusible Signal Factor Family Production in Xanthomonas campestris. *Front Microbiol* **8**,

2486 (2017).

31.   Shen Z., Han J., Wang Y., Sahin O. & Zhang Q. The Contribution of ArsB to Arsenic

Resistance in Campylobacter jejuni. *PLoS One* **8** e58894 (2013).

32.   Rismondo J., Percy M.G. & Gründling A. Discovery of genes required for lipoteichoic acid

glycosylation predicts two distinct mechanisms for wall teichoic acid glycosylation. *J Biol Chem*

293, 3293–3306 (2018).

33. Hemilä H. Lipoamide dehydrogenase of Staphylococcus aureus: nucleotide sequence and

sequence analysis. *Biochim Biophys Acta* **1129**, 119-23 (1991).

34. Jumper J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583–9 (2021).

35. Ma B. & Nussinov R. Trp/Met/Phe hot spots in protein-protein interactions: potential targets

in drug design. *Curr Top Med Chem* **7**, 999-1005 (2007).

36. Waldron D.E. & Lindsay J.A. Sau1: a Novel Lineage-Specific Type I Restriction-

Modification System That Blocks Horizontal Gene Transfer into Staphylococcus aureus and

between S. aureus Isolates of Different Lineages. *J. Bacteriol* **188**, 5578-85 (2006).

37. Monk I.R. *et al.* Transforming the Untransformable: Application of Direct Transformation

To Manipulate Genetically Staphylococcus aureus and Staphylococcus epidermidis. *mBio* **3**,

e00277-11 (2012).

38. Maki H., McCallum N., Bischoff M., Wada A. & Berger-Bächi B. tcaA inactivation increases

glycopeptide resistance in Staphylococcus aureus. *Antimicrob Agents Chemother* **48**, 1953-9

(2004).

39. Schlag, M., et al. Role of staphylococcal wall teichoic acid in targeting the major autolysin

Atl. *Mol Microbiol.* **75**, 864-73 (2010).

40. Brown S, et al. Methicillin resistance in Staphylococcus aureus requires glycosylated wall

teichoic acids. *Proc Natl Acad Sci U S A.* **109**, 18909-14 (2012).

41. Kurokawa K., *et al.* Glycoepitopes of staphylococcal wall teichoic acid govern complement-

mediated opsonophagocytosis via human serum antibody and mannose-binding lectin. *J Biol*

Chem **288**, 30956-68 (2013).

42. Wanner S., *et al.* Wall teichoic acids mediate increased virulence in Staphylococcus

aureus. *Nature Microbiol* **2**, 16257 (2017).

43. Weidenmaier C., *et al.* Lack of Wall Teichoic Acids in Staphylococcus aureus Leads to

Reduced Interactions with Endothelial Cells and to Attenuated Virulence in a Rabbit Model of

Endocarditis. *Journal Infect Dis* **191**, 1771–7 (2005).

44. Monk IR., Casey P.G., Cronin M., Gahan C.G. & Hill C. Development of multiple strain

competitive index assays for Listeria monocytogenes using pIMC; a new site-specific

integrative vector. *BMC Microbiol.* **13**, 8:96 (2008).

45. Steidl R., *et al.* Staphylococcus aureus cell wall stress stimulon gene-lacZ fusion strains:

potential for use in screening for cell wall-active antimicrobials. *Antimicrob Agents Chemother*

52, 2923-5 (2008).

46. Utaida S., *et al.* Genome-wide transcriptional profiling of the response of Staphylococcus

aureus to cell-wall-active antibiotics reveals a cell-wall-stress stimulon. *Microbiology* **149**,

2719-32 (2003).

47. Levin BR & Bull JJ. Short-sighted evolution and the virulence of pathogenic

microorganisms. *Trends Microbiol.* **2**, 76-81 (1994).

48. Kreiswirth, B. N. et al. The toxic shock syndrome exotoxin structural gene is not detectably

transmitted by a prophage. *Nature* **305,** 709–712 (1983).

49. Horsburgh, M. J. *et al.* σ [B] Modulates Virulence Determinant Expression and Stress

Resistance: Characterization of a Functional *rsbU* Strain Derived from *Staphylococcus aureus*

8325-4. *J Bacteriol* **184**, 5457–5467 (2002).

50. Hawiger, J., Niewiarowski, S., Gurewich, V. & Thomas, D. P. Measurement of fibrinogen

and fibrin degradation products in serum by staphylococcal clumping test. *J Lab Clin Med* **75**,

93–108 (1970).

51. Diep, B. A. *et al.* Complete genome sequence of USA300, an epidemic clone of

community-acquired meticillin-resistant Staphylococcus aureus. *The Lancet* **367**, 731–739

(2006).

52. Schuster, C. F. *et al.* High-throughput transposon sequencing highlights the cell wall as

an important barrier for osmotic stress in methicillin resistant *Staphylococcus aureus* and

underlines a tailored response to different osmotic stressors. *Mol Microbiol* **113**, 699–717

(2020).

53. Krausz K.L. & Bose J.L**.** Bacteriophage Transduction in Staphylococcus aureus: Broth-

Based Method. *Methods Mol Biol.* **1373**, 63-8 (2016).

54. European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European

Society of Clinical Microbiology and Infectious Diseases (ESCMID). Determination of

minimum inhibitory concentrations (MICs) of antibacterial agents by broth dilution. *Clin*

Microbiol Infect **9**, ix-xv (2003).

55. Duggan S. *et al.* A Small Membrane Stabilizing Protein Critical to the Pathogenicity of

Staphylococcus aureus. *Infect Immun* **88**:e00162-20 (2020).

56. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-

Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402–408 (2001).

57. Kho, K. & Meredith, T. Extraction and Analysis of Bacterial Teichoic Acids. *Bio Protoc* **8**,

(2018).

58. Brignoli, T. *et al.* Wall Teichoic Acids Facilitate the Release of Toxins from the Surface of

Staphylococcus aureus. *Microbiol Spectr* (2022) doi:10.1128/spectrum.01011-22.

59. Douglas, E. J. A. *et al.* Antibacterial activity of novel linear polyamines against

Staphylococcus aureus. *Front Microbiol* **13**, (2022).

60. Dengler, V., Meier, P. S., Heusser, R., Berger-Bächi, B. & McCallum, N. Induction kinetics

of the Staphylococcus aureus cell wall stress stimulon in response to different cell wall active

antibiotics. *BMC Microbiol* **11**, 16 (2011).

61. Qazi, S. N. A. *et al. agr* Expression Precedes Escape of Internalized *Staphylococcus*

aureus from the Host Endosome. *Infect Immun* **69**, 7074–7082 (2001).

62. Nguyen, T. H. *et al.* Rapid pathogen-specific recruitment of immune effector cells in the

skin by secreted toxins. *Nat Microbiol* **7**, 62–72 (2022).

63. Bae, J. S. *et al.* Contribution of Staphylococcal Enterotoxin B to *Staphylococcus aureus*

Systemic Infection. *J Infect Dis* **223**, 1766–1775 (2021).