Title: A Gulf Stream frontal eddy harbors a distinct microbiome compared to adjacent

waters

Running Title: Gulf Stream frontal eddy microbiome

Jessica L. Gronniger[1], Patrick C. Gray[1], Alexandria K. Niebergall[2], Zackary I. Johnson[1,3],

Dana E. Hunt[1,3]*

[1]Marine Laboratory, Duke University, Beaufort NC USA

[2]Nicholas School of the Environment, Duke University, Durham NC USA

[3]Biology and Civil & Environmental Engineering, Duke University, Durham NC USA

*To whom correspondence should be directed:

Dana Hunt

135 Duke Marine Lab Rd

Beaufort NC 28516

(252) 504-7542

Abstract

Mesoscale oceanographic features, including eddies, have the potential to alter productivity and

other biogeochemical rates in the ocean. Here, we examine the microbiome of a cyclonic, Gulf

Stream frontal eddy, with a distinct origin and environmental parameters compared to

surrounding waters, in order to better understand the processes dominating microbial community

assembly in the dynamic coastal ocean. Our microbiome-based approach identified the eddy as

distinct from the surround Gulf Stream waters. The eddy-associated microbial community

occupied a larger area than identified by temperature and salinity alone, increasing the predicted

extent of eddy-associated biogeochemical processes. While the eddy formed on the continental

shelf, after two weeks both environmental parameters and microbiome composition of the eddy

were most similar to the Gulf Stream, suggesting the effect of environmental filtering on

community assembly or physical mixing with adjacent Gulf Stream waters. In spite of the

potential for eddy-driven upwelling to introduce nutrients and stimulate primary production,

eddy surface waters exhibit lower chlorophyll *a* along with a distinct and less even microbial

community, compared to the Gulf Stream. At the population level, the eddy microbiome

exhibited differences among the cyanobacteria (e.g. lower *Trichodesmium* and higher

Prochlorococcus) and in the heterotrophic alpha Proteobacteria (e.g. lower relative abundances

of specific SAR11 clades) versus the Gulf Stream. However, better delineation of the relative

roles of processes driving eddy community assembly will likely require following the eddy and

surrounding waters since inception; additionally, sampling throughout the water column could

better clarify the contribution of these mesoscale features to primary production and carbon

export in the oceans.

Introduction

Marine microorganisms are of the engines of biogeochemical cycles, supporting critical ecosystem functions on a global scale; yet these biogeochemical processes depend on the composition, function, and activity of microbial communities (1). Marine microbiology has emphasized environmental filtering (how environmental factors determine microbiome composition due to bottom up effects), in determining community assembly, including factors such as water temperature, salinity and nutrient availability (2, 3). However, processes affecting microbiome community assembly include biological interactions, dispersal limitation and stochasticity (4), as well as the microbiome's ecological history (5). These historical contingencies, the effects of prior conditions on current microbiome assemblages, have emerged as a significant force shaping communities (6, 7). For example, prior environment changes can result in microbiomes that are more resistant (composition remains unchanged after the disturbance) or resilient (composition is altered but eventually returns to pre-disturbance state) to future perturbations (8, 9). Similarly, microbiomes from stable environments may be more sensitive to environmental changes (9, 10). However, in aquatic ecosystems, establishing antecedent environmental conditions is particularly challenging due factors including water movement, unmeasured disturbances and environmental parameter seasonality (11-13). Further, although dispersal limitation is assumed be reduced in aquatic systems compared to other systems, such as soils, oceanographic features such as steep frontal gradients between water parcels or vertical stratification can limit dispersal (14), and thus constrain microbiome composition. Here, we examine a frontal eddy, which may be able to isolate and transport

microbial populations (15), to investigate the relative roles of environmental filtering, ocean

physics and historical contingencies in an aquatic system.

Frontal eddies form along western boundary currents when instabilities in the current, often

stemming from interactions with bathymetry, cause meanders to form and evolve into distinct

cyclonically rotating water parcels. While considered mesoscale eddies, frontal eddies are

smaller and less stable than typical mesoscale eddies such as Gulf Stream rings (16). Gulf Stream

rings form downstream from Cape Hatteras once the current is freely meandering and can "pinch

off", in contrast, frontal eddies do not detach from the Gulf Stream and stay trapped in the

meander of the current. The eddy core is shelf water that is entrained in this Gulf Stream

meander and a shallow warm streamer flows from the downstream meander crest around this

core, separating the eddy from the shelf (17). All frontal eddies are cyclonic, and thus rotation

typically uplifts colder, nutrient-rich waters into the euphotic zone, often stimulating

phytoplankton growth, though continued uplift is dependent on the eddy's age (16-18). Beyond

frontal eddies, there has been substantial oceanographic interest in eddies generally, particularly,

their roles in temporal and spatial variability in primary production, carbon export to the deep

sea, and rapid shifts in marine microbiomes (19-21). Ecologically, eddy formation partially

isolates waters, with the potential to develop microbiomes, environmental parameters, and

biogeochemical rates distinct from surrounding environments (22, 23). However, it is difficult to

predict eddy microbiomes; the specific microbial taxa stimulated vary among the phytoplankton

and include diatoms, prasinophytes, dinoflagellates, and *Synechoccocus* (24-26), with the

potential for concurrent increases in multiple phytoplankton (26), and with likely consequences

for heterotrophic composition and organic matter export (24, 25, 27). Differences among eddy

microbiomes have been attributed to eddy characteristics, such as the initial microbiome,

environmental factors (temperature, nutrient fluxes, mixed layer depth, etc.), and the eddy stage,

among others (24, 28). Thus, although mesoscale eddies are pervasive throughout the global

ocean, and frontal eddies are ubiquitous along western boundary currents, we have poor

predictive ability of their influence on microbiomes and associated biogeochemical processes.

This study takes place just northeast of Cape Hatteras, NC, USA (Figure 1A) which is a dynamic

point where the warm, salty, rapidly-flowing Gulf Stream, the cooler and fresher Slope Sea, and

the even colder and fresher mid-Atlantic Bight shelf water meet, and where the Gulf Stream

separates from the continental shelf. The study area is a region of high gradients, moving from

the productive continental shelf waters to the oligotrophic open ocean in a few dozen kilometers

across shore. Here, we compare a Gulf Stream frontal eddy's microbiome with those of the

adjacent continental shelf, continental slope, and Gulf Stream waters off the coast of Cape

Hatteras. Gulf Stream frontal eddies form ~every 3-7 days in the South Atlantic Bight (18), can

occupy nearly half of the front between the Gulf Stream and the continental shelf in this region

(29), and have been shown to play a significant role in nutrient transport into the euphotic zone,

enhancing primary and possibly secondary production both in the eddy and on the continental

shelf (17, 23, 30). Using a combination of field and satellite-based (temperature and chlorophyll

a) measurements, we take a microbiome-centric approach to identify water parcels and

population-level differences between water parcels in order to better understand the forces

shaping the microbiomes of these mesoscale ocean features.

Figure 1. Locations and microbiome of field samples (A) Map of sampling sites for five ~10-
15 km long ship-board transects off the coast of Cape Hatteras, North Carolina, USA September

5-7, 2021. Background color represents average sea surface temperatures from August 20th to
September 10th obtained from the Geostationary Operational Environmental Satellite 16 (GOES-
16). Each sampling point is shown along the transect and categorized by temperature and salinity
(shapes) and microbial-community based k-means clusters (colors), generated using the self-
organizing map (SOM) R package 'kohonen'. Isobaths for 500m (white), 1000m (grey) and
2000m (black) are included for context. (B) Non-metric multidimensional scaling (NMDS)
ordination computed based on Bray-Curtis dissimilarity for 16S rRNA gene libraries. Ellipses
show the multivariate t-distribution 95% confidence interval for the mean of each SOM cluster.
Environmental factors found to be statistically significant (Multiple regression permutation test,
$p < 0.05$) are indicated with an '*'.
126

Results & Discussion

Using satellite data, we followed a single Gulf Stream frontal eddy from formation on the

Charleston Bump until sampling ~2 weeks later (Figure 2). This sampled Gulf Stream frontal

eddy was visible via satellite as a cooler, chlorophyll *a* enriched region relative to the

surrounding Gulf Stream waters (31); however, there was no evidence of a surface bloom or

enhancement of chlorophyll *a* levels compared to the shelf waters. In fact, satellite- measured

surface chlorophyll *a* levels decline over time, likely either through nutrient depletion or mixing

with Gulf Stream waters (Figure 2)(31). Here, we *in situ* sampled the eddy (and surrounding

waters) as it moved past Cape Hatteras on September 5-7th 2021. At this point the eddy was

approximately 20 by 50 km (longer in the along stream dimension) and we collected five across-

stream transects of 10-15 km each (49 stations with 1-2 km between stations) to obtain high-

resolution measurements of surface microbial community composition and environmental

conditions (Figure 1A).

Figure 2. Satellite overview of the observed frontal eddy from formation to *in situ*
sampling. Black dotted lines are all 5 transects from this study. Black boxes outline the eddy
location across five days from August 27th to September 6th. SST is from GOES-16 and
chlorophyll *a* is from Sentinel-3's OLCI sensor. Boxes were determined manually by identifying
an anomaly in SST and ocean color measurements that is consistent with a cyclonic frontal eddy
and associated warm streamer.

Initially, surface temperature and salinity were used to identify water parcels (Figure S1) (31).

However, these physical parameter-based categories did not align with microbiome composition,

particularly among the Gulf Stream and eddy samples (Figure S2A). Thus, we used a

microbiome-based, self-organizing map (SOM) approach (32) to cluster the microbiome data

independently of environmental variables. We then compared these biologically-based clusters

(SOM) with the temperature and salinity-based water parcel classifications. Although these two

schemes did overlap, there were also a number of disagreements in the assignments (Figure 1B):

SOM cluster 1 largely corresponds to the continental shelf (87.5%), cluster 2 the continental

slope (67%), cluster 3 the Gulf Stream (77%) and cluster 4 the eddy (62.5%) stations. Satellite

data suggests that disagreements between these simple physical parameters and microbiome-

based classifications occurred at transitions between water parcels, particularly around the

periphery of the eddy (Figure S3), which could be due to complex mixing between the eddy and

surrounding Gulf Stream. We found that the microbiome-based (SOM) approach classified more

samples as eddy-associated. Therefore, identification of the eddy based on temperature and

salinity alone could underestimate the geographic extent of the eddy's biogeochemical influence

(Figure S3), or fail to identify mixing of the eddy microbiome into surrounding waters.

Henceforth, for clarity, these statistically-distinct SOM clusters (ANOSIM, $p<0.05$) will be

referred to by their most common location (1 = continental shelf, 2 = continental slope, 3 = Gulf

Stream, 4 = eddy), with the assumption that SOM clusters most accurately map potential

ecological differences between the samples. However, disagreements between clustering based

on surface properties (temperature, salinity) and microbiomes merits further investigation in both

the biological and physical oceanographic fields, particularly at eddy margins to better identify

microbiome origins and implications (e.g. changes in biogeochemical rates).

Next, we examined how environmental factors differed among these clusters; the shelf and slope

clusters are also clearly distinct on the NMDS, with relatively higher overlap for the Gulf Stream

and eddy clusters (Figure 1B). We examined environmental variables that might help explain the

observed microbiome patterns, the continental shelf and slope exhibit higher cell abundances

(total prokaryotes, *Synechococcus*) and higher PO_4 along with lower salinity and cooler water

temperatures compared to the eddy and Gulf Stream (Figures 3, S4; Pairwise Wilcoxon ranked

sum test $p < 0.05$). This is generally consistent with higher resource availability on the

continental shelf and slope compared to the Gulf Stream and eddy, indicating greater terrestrial

influence and resources in near-shore environments and oligotrophic conditions beyond the shelf

break (33). Ordination of the microbiome in relation to select environmental variables

[temperature, salinity, chlorophyll *a*, nutrients (SiO_4, NO_3, NH_4, PO_4), and flow-cytometric

abundances of bacteria and cyanobacteria] identified significant associations for variables other

than SiO_4, NO_3 and NH_4, (Figure 1B, Table S1), highlighting the factors which could explain the

microbiome composition. Here, we specifically focused on comparison of the Gulf Stream and

eddy samples as they are most similar in both environment and microbiome: the eddy is ~1 °C

cooler, has lower chlorophyll *a*, lower *Synechococcus* and higher non-phycoerythrin containing

picocyanobacteria (hereafter labeled *Prochlorococcus*) concentrations (Figures 3, S4; Pairwise

Wilcoxon ranked sum test $p < 0.05$). While PO_4 in the eddy is statistically higher than in the Gulf

Stream, eddy phosphate concentrations have a large range, suggesting potential variability in

nutrient fluxes/ availability within this water parcel (Figure 3D). Thus, in contrast to other

cyclonic eddies (23, 34), here, eddy surface waters generally exhibit oligotrophic characteristics

(low nutrients, high *Prochlorococcus* concentrations, lower chlorophyll *a,* but higher PO_4)

compared to the Gulf Stream, in spite of forming via entrainment of presumably higher nutrient

continental shelf water and the potential for upwelling-driven nutrient additions (Figures 3, S4).

Overall, these results are consistent with the satellite measurements that do not show enhanced

surface photosynthetic biomass of the eddy. Possible explanations for this result include that

photosynthetic biomass enhancement occurred primarily below the mixed layer and was not

visible at the surface, or that high stratification in the summer along with a deep euphotic depth

prevented the eddy-associated uplift from enhancing productivity. Examination of microbial

community characteristics (diversity etc.) as well as specific phylotypes associated with each

water parcel may offer additional clues to factors that shaped these communities in the coastal

ocean.

Figure 3. Key environmental factors for Self-Organizing Map (SOM) clusters. Numbered
clusters correspond to the most common environments: Continental shelf (Cluster 1), Continental
slope (Cluster 2), Gulf Stream (Cluster 3), Eddy (Cluster 4). For the box and whisker plots, the
center lines denote the median value while the box encompasses the 25^{th} to 75^{th} percentiles of the
dataset. Whiskers denote the 5^{th} and 95^{th} percentiles and values that fall outside these ranges are
shown as individual points. Brackets indicate p-values of Wilcoxon Rank Sum pairwise
comparisons. Additional environmental parameter comparisons are shown in Figure S4.
214

Microbial Assemblages

As eddies can alter microbial community parameters, potentially increasing diversity through

mixing of water parcels and influx of resources, or conversely reducing diversity and evenness

through competitive exclusion (20, 35-37), we compared diversity and evenness indexes in the

four water parcels. Notably, the eddy exhibits lower richness and evenness compared to the Gulf

Stream (Figure 4; Wilcoxon ranked sum test $p<0.05$), reflecting the dominance of specific taxa,

particularly *Prochlorococcus* (35) (Figure 5). While the eddy initially formed from continental

shelf waters (although off the coast of South Carolina) and retains comparable richness to shelf

waters in these transects (Figure 4), the eddy is statistically less even than the other water parcels

~two weeks after formation, suggesting continued development of the microbiome (Wilcoxon

ranked sum test $p<0.05$). While the eddy microbiome is most similar to the Gulf Stream's, if

dispersal limitation significantly shapes the eddy microbiome and if we assume the shelf

community where the eddy formed is similar to the shelf community off Cape Hatteras, we

might predict that presence-absence data (Sorenson index) would group the eddy with the

continental shelf and that relative abundances would shift to match the current environmental

conditions. However, the Sorenson index also groups the eddy and the Gulf Stream samples

(Figure S5), as was observed for Bray-Curtis dissimilarity (Figure 1B), suggesting that

environmental filtering or physical mixing shapes frontal eddy communities.

Figure 4. Alpha diversity metrics for 16S rRNA gene community composition for each Self-
Organizing Map (SOM) cluster. (A) Shannon's H index, (B) species richness, and (C) Pielou's
evenness were calculated based on the absolute abundance of ASVs using 'vegan' (v2.6.2) in R
(v4.0.0). SOM cluster numbers correspond to the following classifications: 1 –continental shelf,
2-continental slope, 3- Gulf Stream, 4- eddy. Brackets indicate p-values associated with
Wilcoxon Rank Sum test pairwise comparisons.
241
242
Figure 5. Average family-level composition for each Self-Organizing Map (SOM) cluster.
The average community was calculated by taking the mean relative abundance of each taxon
across all samples within that cluster, then normalizing relative abundances to 100%. ASVs with
a relative abundance of less than 1% are grouped as "Other." Amplicon Sequence Variants
(ASVs were generally grouped at the family level, but when family is undefined they are labeled

at the phylum level. Additionally, the family *Synechococcaceae* was resolved at the genus level
to illustrate the *Synechococcus-Prochlorococcus* inversion between clusters.
250

Across all samples, the microbiome composition reflects abundant coastal microbes including

Synechococcaceae (~25-40%), *Pelagibacteraceae* (~28-40%), Actinobacteria family OCS155

(~7%), the Archaeal Marine Group II clade (~2-4%) and the SAR406 clade A714017 (~2%;

Figure 5) (3, 33, 38, 39). Yet, we observed onshore-offshore spatial transitions (33), including

the switch within *Synechococcaceae* genera, with *Synechococcus* dominating the shelf and slope,

and *Prochlorococcus* the Gulf Stream and eddy (Figure 5). As specific taxa may offer additional

clues about the factors which shaped these microbiomes, we examined ASV relationships with

specific clusters, using 250 most abundant ASVs (representing ~73% of the microbial

community) with both linear discriminant effect size analysis (LEfSe) and a Bayesian

generalized joint attribute model (GJAM) (40, 41) (Tables S2, S3). Both LEfSe and GJAM

identify taxa associated with specific water parcels (Figure 6); moreover, these approaches

generally agree: identifying increased relative abundance of *Pelagibacter* and *Synechococcus* in

continental shelf and slope waters and enrichment of *Prochlorococcus* and *OCS155* in Gulf

Stream and eddy waters, which generally aligns with expectations of oligotrophic, warm water-

associated taxa offshore (33). GJAM identified a single difference between the eddy and Gulf

Stream associated taxa, a High Light clade I *Prochlorococcus* ASV (Tables S2,S3; Figure 6B)

that was enriched in both the eddy and slope samples, potentially suggesting either a continued

legacy of nearshore waters or mixing of slope waters into the eddy (Figure 6B). Although there

are similarities between the eddy and Gulf Stream samples, their differences help identify the

current (e.g. environmental filtering) and antecedent (e.g. historical contingencies) conditions

that shape the microbiome.

Figure 6. Amplicon sequence variant (ASV) associations with specific SOM groupings (A)
Heatmap of ln(absolute abundance+1) of the 22 ASVs identified with at least one significant
SOM cluster beta coefficient obtained from generalized joint attribute modeling (GJAM)
analysis of the 250 most abundant ASVs across all samples. Samples are organized by SOM
grouping (labeled at top) (B) Heatmap of statistically significant SOM cluster-associated beta
coefficients (95% confidence interval does not overlap 0), non-significant beta coefficients are
shown as white. Colored labels at the top of each column identifies the SOM cluster (as in panel
A). ASVs are organized maximum likelihood phylogenetic tree constructed using Smart Model
Selection in PhyML 3.0 with taxonomic assignments based on the RDP naïve Bayesian classifier
using the Greengenes version 13.5 database.
284

A comparison of just the eddy and Gulf Stream using both LEfSe and DESeq2 (42) more clearly

reveals distinctions between their microbiomes (Table S4; Figure 7). A number of SAR11

lineages were enriched in the Gulf Stream compared to the eddy (Figure 7). The SAR11 clade is

known to contain streamlined heterotrophs (43); however, we can only speculate that ASV

differences could be due to resource differences or stochastic effects. Several high-light clade

Prochlorococcus ASVs were enriched in the eddy over the Gulf Stream which is consistent with

flow-cytometric non-phycoerythrin containing picocyanobacteria enrichment in the eddy (Figure

3F) and their role as efficient nutrient recyclers (44). Interestingly, although inorganic nutrients

are low in both the eddy and Gulf Stream (Figures 3, S4), the Gulf Stream samples are enriched

in a putatively nitrogen-fixing cyanobacterium *Trichodesmium* ASV (Figure 7) (45). Thus,

within the cyanobacteria and despite similar inorganic nitrogen concentrations, the Gulf Stream

and eddy microbiomes may reveal differences in nutrient fluxes or the role of historical

contingencies in microbiome composition (46). Moreover, we acknowledge that standing stocks

(e.g. nutrient concentrations) are not equivalent to nutrient turnover rates, as resources can be

rapidly cycled in oligotrophic conditions (47). The development of the colonial, nitrogen-fixing

Trichodesmium may require additional time or lower nitrogen fluxes. As the eddy sustains higher

concentrations of *Prochlorococcus* and total bacterioplankton, and statistically higher levels of

PO_4, this data could indicate that the eddy has experienced higher nutrient fluxes, which could

support higher primary and secondary production, although this is not evident in standing

photosynthetic biomass (as measured by chlorophyll *a*, Figure 3C). Although there was no

satellite-detectable eddy surface bloom, and the eddy microbiome likely shifted toward that of

the Gulf Stream over time, microbial communities can offer clues about the processes occurring

in these dynamic mesoscale features.

308
Figure 7. Microbiome differences between the Gulf Stream and eddy. Left panel: Phylotypes
(ASVs) that exhibited significant differential abundance in the eddy relative to the Gulf Stream
as identified using DESeq2, left side indicates enrichment in the Gulf Stream, right side eddy.
Dots indicates the Log_2 fold changes for each ASV identified as significant (Benjamin-Hochberg
multiple hypothesis adjusted p<0.05), error bars indicate the standard error of the Log_2 fold
changes as calculated by DESeq2. Only ASVs with a minimum average relative abundance of
0.05% across the samples being compared were included in the analysis. Y-axis represents
maximum likelihood phylogenetic tree constructed using Smart Model Selection in PhyML 3.0.
Taxonomic assignments were obtained from the RDP database. Right hand column indicates
those ASVs that were also found to have a significant association with either the Gulf Stream
(dark blue) or the eddy (light blue) as identified using Linear discriminant analysis Effect Size
(LEfSE).
321

Conclusions

Here we show that a cyclonic, frontal eddy that formed along the western boundary of the Gulf

Stream harbored a microbial community distinct from those of surrounding waters. This eddy

community appears to be shaped by either environmental selection or mixing with the Gulf

Stream (16, 29), rather than dispersal limitation, as it was most similar to that of the Gulf Stream

rather than continental slope waters where it originated. While this study was limited to surface

samples and encompasses a brief period in the eddy's development, new technologies are

enabling Lagrangian sampling of eddy parameters during development and throughout the water

column to better quantify upwelling, eddy coherence and mixing with the Gulf Stream, across

depths (48-51). High-resolution, *in situ* sampling will enable the field to develop a more

quantitative assessment of how source populations and environmental drivers interact to shape

these dynamic mesoscale features over time. Although more research is needed to understand the

impact of eddies globally, this study deepens our understanding of the forces shaping Gulf

Stream frontal eddy microbiomes off the South Atlantic Bight and provides important directions

for future studies. While there was not support for this eddy contributing to the biodiversity

hotspot located off of Cape Hatteras (36, 52), other frontal eddies may enhance biodiversity by

delivering distinct microbial communities, enhanced zooplankton populations, and upwelling of

resources (31). Furthermore, the frequency of these eddies and their predicted nutrient injections

into the Gulf Stream may play a significant role in carbon sequestration and productivity along

the South Atlantic Bight. Although we found this eddy harbored a distinct microbial community,

further investigations will be needed to better understand the role of historical contingencies and

the temporal dynamics of eddy-associated communities and their potential role in global

biogeochemical cycles. Moreover, this approach of microbially-defined, environmental variable

agnostic sample clustering can be applied more broadly in oceanography to help disentangle

potentially ecologically-distinct microbiomes.

Methods

Sample collection and sea water parameter characterization

We tracked via satellite a cyclonic, frontal eddy that formed from shelf water off of Charleston,

South Carolina, USA on August 25-26[th] 2021, was isolated by warmer, Gulf Stream-derived

waters, and then traveled north along the Gulf Stream's western front (31). Seawater was

collected on board the *R/V Shearwater* on 5 transects from September 5-7[th], 2022; transects were

planned to cross the front between the shelf and eddy or shelf and Gulf Stream waters. Water

samples were collected during transit from the ship's flow-through seawater system (~1 m

depth). In general, the transects were 10-15 km long (over 2-3 hours) with 14-21 samples

collected per transect at approximately 10 min intervals for a total of 86 sampling points, 49 of

which included filtration for later nucleic acid analysis. Temperature and salinity were measured

with a SeaBird SBE38 Thermosalinograph. Chlorophyll *a* concentrations were measured by

filtering 100 mL of seawater onto a ~0.7 μm glass fiber filter (Whatmann APFF02500) using

gentle (<10mm Hg) vacuum. Chlorophyll *a* pigments were extracted in 100% methanol for 48h

at -20 °C and fluorescence was measured using a Turner Designs 10-AU fluorometer as

previously described (53). Duplicate whole seawater samples for flow-cytometric analyses were

collected, fixed with net 0.125% glutaraldehyde and stored at -80 °C until processing.

Prokaryotic phytoplankton populations were enumerated using a Becton Dickinson

FACSCalibur Flow Cytometer and categorized as previously described (53). Bacterioplankton

were quantified using SYBR Green-I on the Attune NxT acoustic flow cytometer (Life

Technologies) (54). Duplicate 0.22 μm Sterivex (Milipore) filtered seawater samples were

collected for dissolved inorganic nutrient quantification and stored at -80 °C until processing.

Nutrients (NO_2, NO_3, PO_4, SiO_4) were analyzed at the UC San Diego Scripps Institution of

Oceanography ODF chemical laboratory using a Seal Analytical continuous-flow AutoAnalyzer

3 (55, 56). Detection limits for the nutrients analyzed are as follows: $NO_2 = 20nM$, $NO_3 = 20nM$,

$PO_4 = 20nM$, $SiO_4 = 200nM$. No permits were required for environmental sampling within US

coastal waters.

DNA extraction and library preparation:

Microbial DNA was collected by filtering ~4 liters of whole seawater through a 0.22 micron

Sterivex filter (Millipore). Filters were stored at -20 °C while on the research vessel and then at -

80 °C upon return to the lab until extraction. Genomic DNA for 16S rRNA gene libraries was

extracted using the Gentra Puregene Yeast/Bacteria kit (QIAGEN) supplemented with 60

seconds of bead beating. Extracted DNA was quantified using a Nanodrop ND-100 and 16S

rRNA gene libraries were generated using gene primers targeting the V4-V5 region 515F- (5′-

GTGYCAGCMGCCGCGGTAA) and 926R (5′-CCGYCAATTYMTTTRAGTTT) (57, 58).

PCR reactions were performed in triplicate with 20 μL reactions containing 20 ng template

DNA, 1× Taq Buffer, 0.5 μM of each primer, 200 μM of dNTPs, and 0.4 U of Q5 DNA

polymerase (NEB). The thermal cycling conditions were 30 sec at 98 °C, followed by 25 cycles

of 10 sec at 98 °C, 30 sec at 50 °C, 30 sec at 72 °C, and a final extension at 72 °C for 2 min.

Triplicate PCR reactions were pooled and gel purified. Libraries were pooled and sequenced at

the Duke Center for Genomic and Computational Biology using V2 2 × 250 bp sequencing on

the Illumina MiSeq.

Sequence Processing:

Barcodes were removed and sequences were demultiplexed and assigned to samples using Sabre

(https://github.com/najoshi/sabre). Sequences were cleaned and clustered using VSEARCH

v2.5.1 (59). Low quality sequence ends were trimmed at the Phred quality score (Q) of 30 using

a 10 bp running window. Paired-end reads with 10 bp overlap and no mismatches were merged

and sequences with expected errors >1 and/or a length < 360 bp were removed. Minimum

entropy decomposition analysis (v2.1) was used to resolve amplicon sequences variants (ASVs)

(60). Sequencing-error associated noise was reduced by retaining only those ASVs with unique

sequences with a minimum abundance of 20 occurrences. Samples contained an average of

39,936 representative reads. Representative ASV sequences were assigned taxonomy using the

RDP classifier in MacQIIME v1.9.1. Mitochondrial sequences were removed and libraries were

subsampled to 14,187 reads per library. AmpliCopyrighter (v.0.46) was used to correct for

differential gene copy numbers across taxa (61). "Absolute abundance" was calculated by

multiplying relative abundances of ASVs by prokaryotic cell counts obtained from flow

cytometry analysis. NMDS plots were generated by calculating Bray-Curtis dissimilarity scores

based on ASV relative abundances using the *vegdist()* function in R.

Self-Organizing Map

The Kohonen package (32) in R (v. 4.0.0) was used to cluster samples based on microbial

community composition. A self-organizing map was constructed using ASV relative abundances

and each sample was assigned to a map unit based on community similarities. Our map was

composed of 4x4 circular nodes in a hexagonal, nontoroidal configuration. Clusters were

generated using k-means clustering and a reasonable value for k was chosen by estimating the

point of inflection in a scree plot of within-clusters sum of squares (Figure S6).

Discriminative taxa identification

Generalized joint attribute modelling (GJAM) was applied to model the 250 most abundant

ASVs and select environmental factors (temperature, salinity, chlorophyll *a*, PO_4), using the

GJAM v. 2.6.2 package in R. Iteration was set at 20,000 and burning at 5,000. Results were

visualized using the built-in function 'gjamPlot'. In order to identify discriminative taxa between

SOM clusters, the 250 most abundant ASVs were analyzed with LEfSe (41). The threshold for

significance on the logarithmic LDA score for discriminative features was two. P-values were

obtained from the Kruskall-Wallis test for differences in ASV abundances among SOM cluster

assignments. These two metrics differ in approach: LEfSe identifies phylotype enrichment in a

specific cluster while GJAM can identify the magnitude and directionality (positive or negative)

of associations across multiple clusters. ASVs exhibiting differential abundances between the

Gulf Stream and eddy were further identified using DESeq2 (42). Only ASVs with an average

relative abundance >0.05% averaged across samples being compared (Gulf Stream and eddy)

were included. Significant differences were considered those with a Benjamin-Hochberg

multiple hypothesis adjusted p<0.05. Quantification of the processes shaping microbiome

community assembly using Beta Mean Nearest Taxon Distance (βMNTD) (5) was not possible

because our data did not exhibit the required phylogenetic signal (non-significant Mantel

correlogram).

Satellite Data

Satellite data from two sources is used in this work. Chlorophyll *a* products were derived from

ocean color imagery acquired by the European Commission Copernicus programme's Sentinel-3

Ocean and Land Color Instrument (OLCI) using the CHL_OC4ME blue to green band ratio

algorithm (62). The OLCI data was used to track surface chlorophyll *a* within the eddy over its

lifetime. Sea surface temperature products were from the Geostationary Operational

Environmental Satellite 16 (GOES-16) which uses a retrieval from the long wave infrared bands

(8μm-12μm) (63). OLCI Level 2 and obtained from https://coda.eumetsat.int and GOES-16

hourly SST was acquired from

https://cwcgom.aoml.noaa.gov/erddap/griddap/goes16SSThourly.html.

Physical Water Parcel Delineations

Temperature and salinity delineations between water parcels are specific to this region and time.

They are based on satellite tracking of the eddy water mass and known Temperature-Salinity

properties of these waters. In general Gulf Stream waters are the hottest and saltiest, South

Atlantic Bight waters are slightly fresher and cooler, Mid Atlantic Bight slope waters are cooler

and fresher again, and Mid Atlantic Bight shelf water is the coolest and freshest (31, 64). To

determine the exact thresholds of temperature and salinity these general patterns were informed

and reinforced by observed sharp frontal gradients between water masses. This led to the

following thresholds: Gulf Stream > 28.2 °C, eddy <28.2 °C and > 35.75 PSU, continental slope

<35.75 PSU and > 34 PSU, and continental shelf <34 PSU (Figure S1) (31).

Data and Code Repository: Cruise field data is deposited: doi.org/10.5281/zenodo.7680135

and SSU rRNA library sequences are deposited as part of Bioproject PRJNA868376. Analysis

scripts are available at https://github.com/jgronniger/Gulf-Stream-Eddy-2021.

Acknowledgements:

We acknowledge the assistance of the captain and crew of the *R/V Shearwater.*

CRediT Contributions:

Jessica Gronniger- Conceptualization; Data Curation; Formal Analysis; Visualization; Writing-

original draft

Patrick C. Gray- Conceptualization; Data Curation; Formal Analysis; Visualization; Writing-

review and editing

Alexandria K. Niebergall- Formal Analysis; Writing-review and editing

Zackary I. Johnson- Resources; Writing-review and editing

Dana E. Hunt- Conceptualization; Resources; Writing-original draft;

References

1.   Falkowski PG, Fenchel T, Delong EF. The microbial engines that drive Earth's
biogeochemical cycles. Science. 2008;320(5879):1034-9.
2.   Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, et al. Structure
and function of the global ocean microbiome. Science. 2015;348(6237):1261359.
3.   Ward CS, Yung C-M, Davis KM, Blinebry SK, Williams TC, Johnson ZI, et al. Annual
community patterns are driven by seasonal switching between closely related marine bacteria.
The ISME Journal. 2017;11(6):1412-22.
4.   Zhou J, Deng Y, Zhang P, Xue K, Liang Y, Van Nostrand JD, et al. Stochasticity,
succession, and environmental perturbations in a fluidic ecosystem. Proceedings of the National
Academy of Sciences. 2014;111(9):E836-E45.
5.   Stegen JC, Lin X, Fredrickson JK, Chen X, Kennedy DW, Murray CJ, et al. Quantifying
community assembly processes and identifying features that impose them. The ISME Journal.
2013;7(11):2069.
6.   Hawkes CV, Keitt TH. Resilience vs. historical contingency in microbial responses to
environmental change. Ecology Letters. 2015;18(7):612-25.
7.   Stegen JC, Bottos EM, Jansson JK. A unified conceptual framework for prediction and
control of microbiomes. Current Opinion in Microbiology. 2018;44:20-7.

8.   Jurburg SD, Nunes I, Brejnrod A, Jacquiod S, Priemé A, Sørensen SJ, et al. Legacy
effects on the recovery of soil bacterial communities from extreme temperature perturbation.
Frontiers in Microbiology. 2017;8:1832.
9.   Waring BG, Hawkes CV. Short-term precipitation exclusion alters microbial responses to
soil moisture in a wet tropical forest. Microbial Ecology. 2015;69:843-54.
10.   Wang Z, Tsementzi D, Williams T, Juarez D, Garcia N, Johnson Z, et al. Environmental
stability impacts the differential sensitivity of marine microbiomes to increases in temperature
and acidity. The ISME Journal. 2021;15(1):19-28.
11.   Gronniger JL, Wang Z, Brandt GR, Ward CS, Tsementzi D, Mu H, et al. Rapid changes
in coastal ocean microbiomes uncoupled with shifts in environmental variables. Environmental
Microbiology. 2022.
12.   Evans S, Martiny JB, Allison SD. Effects of dispersal and selection on stochastic
assembly in microbial communities. The ISME Journal. 2017;11(1):176-85.
13.   Shen D, Langenheder S, Jürgens K. Dispersal modifies the diversity and composition of
active bacterial communities in response to a salinity disturbance. Frontiers in Microbiology.
2018;9:2188.
14.   Bracco A, Provenzale A, Scheuring I. Mesoscale vortices and the paradox of the
plankton. Proceedings of the Royal Society of London Series B: Biological Sciences.
2000;267(1454):1795-800.
15.   Villar E, Farrant GK, Follows M, Garczarek L, Speich S, Audic S, et al. Environmental
characteristics of Agulhas rings affect interocean plankton transport. Science.
2015;348(6237):1261447.
16.   Gula J, Molemaker MJ, McWilliams JC. Submesoscale dynamics of a Gulf Stream
frontal eddy in the South Atlantic Bight. Journal of Physical Oceanography. 2016;46(1):305-25.
17.   Lee TN, Yoder JA, Atkinson LP. Gulf Stream frontal eddy influence on productivity of
the southeast US continental shelf. Journal of Geophysical Research: Oceans.
1991;96(C12):22191-205.
18.   Glenn SM, Ebbesmeyer CC. Observations of Gulf Stream frontal eddies in the vicinity of
Cape Hatteras. Journal of Geophysical Research: Oceans. 1994;99(C3):5047-55.
19.   Falkowski PG, Ziemann D, Kolber Z, Bienfang PK. Role of eddy pumping in enhancing
primary production in the ocean. Nature. 1991;352(6330):55-8.
20.   Clayton S, Dutkiewicz S, Jahn O, Follows MJ. Dispersal, eddies, and the diversity of
marine phytoplankton. Limnology and Oceanography: Fluids and Environments. 2013;3(1):182-
97.
21.   Martin-Platero AM, Cleary B, Kauffman K, Preheim SP, McGillicuddy DJ, Alm EJ, et al.
High resolution time series reveals cohesive but short-lived communities in coastal plankton.
Nature Communications. 2018;9(1):266.
22.   Sweeney EN, McGillicuddy Jr DJ, Buesseler KO. Biogeochemical impacts due to
mesoscale eddy activity in the Sargasso Sea as measured at the Bermuda Atlantic Time-series
Study (BATS). Deep Sea Research Part II: Topical Studies in Oceanography. 2003;50(22-
26):3017-39.
23.   Baltar F, Arístegui J, Gasol JM, Lekunberri I, Herndl GJ. Mesoscale eddies: hotspots of
prokaryotic activity and differential community structure in the ocean. The ISME Journal.
2010;4(8):975-88.

24. Doblin MA, Petrou K, Sinutok S, Seymour JR, Messer LF, Brown MV, et al. Nutrient
uplift in a cyclonic eddy increases diversity, primary productivity and iron demand of microbial
communities relative to a western boundary current. PeerJ. 2016;4:e1973.
25. Lochte K, Pfannkuche O. Cyclonic cold-core eddy in the eastern North Atlantic. II.
Nutrients, phytoplankton and bacterioplankton. Marine Ecology Progress Series. 1987:153-64.
26. Wang L, Huang B, Chiang K-P, Liu X, Chen B, Xie Y, et al. Physical-biological
coupling in the western South China Sea: The response of phytoplankton community to a
mesoscale cyclonic eddy. PloS One. 2016;11(4):e0153735.
27. Ahern OM, Whittaker KA, Williams TC, Hunt DE, Rynearson TA. Host genotype
structures the microbiome of a globally dispersed marine phytoplankton. Proceedings of the
National Academy of Sciences. 2021;118(48).
28. Harke MJ, Frischkorn KR, Hennon GM, Haley ST, Barone B, Karl DM, et al. Microbial
community transcriptional patterns vary in response to mesoscale forcing in the North Pacific
Subtropical Gyre. Environmental Microbiology. 2021;23(8):4807-22.
29. Gula J, Molemaker MJ, McWilliams JC. Gulf Stream dynamics along the southeastern
US seaboard. Journal of Physical Oceanography. 2015;45(3):690-715.
30. Paffenhöfer G, Sherman B, Lee T. Summer upwelling on the southeastern continental
shelf of the USA during 1981. Zooplankton abundance and distribution. Prog Oceanog.
1987;19:403-36.
31. Gray PC, Gronniger J, Savelyev I, Dale J, Niebergall AK, Cassar N, et al. Investigating a
Gulf Stream Frontal Eddy North of Cape Hatteras: Questioning Overall Impacts on Ecology.
bioRxiv. 2023:2023.02. 22.529409.
32. Wehrens R, Buydens LM. Self-and super-organizing maps in R: the Kohonen package.
Journal of Statistical Software. 2007;21:1-19.
33. Wang Z, Juarez DL, Pan JF, Blinebry SK, Gronniger J, Clark JS, et al. Microbial
communities across nearshore to offshore coastal transects are primarily shaped by distance and
temperature. Environmental Microbiology. 2019;21(10):3862-72.
34. Benitez-Nelson CR, Bidigare RR, Dickey TD, Landry MR, Leonard CL, Brown SL, et al.
Mesoscale eddies drive increased silica export in the subtropical Pacific Ocean. Science.
2007;316(5827):1017-21.
35. Lévy M, Jahn O, Dutkiewicz S, Follows MJ, d'Ovidio F. The dynamical landscape of
marine phytoplankton diversity. Journal of The Royal Society Interface.
2015;12(111):20150481.
36. Barton AD, Dutkiewicz S, Flierl G, Bragg J, Follows MJ. Patterns of Diversity in Marine
Phytoplankton. Science. 2010;327(5972):1509-11.
37. Pajares S. Unraveling the distribution patterns of bacterioplankton in a mesoscale
cyclonic eddy confined to an oxygen-depleted basin. Aquatic Microbial Ecology. 2021;87:151-
66.
38. Chow C-ET, Sachdeva R, Cram JA, Steele JA, Needham DM, Patel A, et al. Temporal
variability and coherence of euphotic zone bacterial communities over a decade in the Southern
California Bight. The ISME Journal. 2013;7:2259–73.
39. Yilmaz P, Yarza P, Rapp JZ, Glöckner FO. Expanding the world of marine bacterial and
archaeal clades. Frontiers in Microbiology. 2016;6:1524.
40. Clark JS, Nemergut D, Seyednasrollah B, Turner PJ, Zhang S. Generalized joint attribute
modeling for biodiversity analysis: Median- zero, multivariate, multifarious data. Ecological
Monographs. 2017;87(1):34-56.

41.   Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic
biomarker discovery and explanation. Genome Biology. 2011;12(6):R60.
42.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology. 2014;15(12):1.
43.   Giovannoni SJ. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. Annual
Review of Marine Science. 2017;9(1):231-55.
44.   Polz MF, Hunt DE, Preheim SP, Weinreich DM. Patterns and mechanisms of genetic and
phenotypic differentiation in marine microbes. Philosophical Transactions of the Royal Society
B-Biological Sciences. 2006;361(1475):2009-21.
45.   Palter JB, Ames EJ, Benavides M, Goncalves Neto A, Granger J, Moisander PH, et al.
High N 2 fixation in and near the Gulf Stream consistent with a circulation control on
diazotrophy. Geophysical Research Letters. 2020;47(16):e2020GL089103.
46.   Conway TM, Palter JB, de Souza GF. Gulf Stream rings as a source of iron to the North
Atlantic subtropical gyre. Nature Geoscience. 2018;11(8):594-8.
47.   Benitez-Nelson CR, Buesseler KO. Variability of inorganic and organic phosphorus
turnover rates in the coastal ocean. Nature. 1999;398(6727):502-5.
48.   Ntaganou N, Kourafalou V, Beron-Vera FJ, Olascoaga MJ, Le Hénaff M, Androulidakis
Y. Influence of Caribbean eddies on the Loop current system evolution. Frontiers in Marine
Science. 2023;10:961058.
49.   Calil P, Richards K. Transient upwelling hot spots in the oligotrophic North Pacific.
Journal of Geophysical Research: Oceans. 2010;115(C2).
50.   Tarshish N, Abernathey R, Zhang C, Dufour CO, Frenger I, Griffies SM. Identifying
Lagrangian coherent vortices in a mesoscale ocean model. Ocean Modelling. 2018;130:15-28.
51.   Zhang Y, Ryan JP, Hobson BW, Kieft B, Romano A, Barone B, et al. A system of
coordinated autonomous robots for Lagrangian studies of microbes in the oceanic deep
chlorophyll maximum. Science Robotics. 2021;6(50):eabb9138.
52.   Byrd BL, Hohn AA, Lovewell GN, Altman KM, Barco SG, Friedlaender A, et al.
Strandings as indicators of marine mammal biodiversity and human interactions off the coast of
North Carolina. 2014.
53.   Johnson ZI, Shyam R, Ritchie AE, Mioni C, Lance VP, Murray JW, et al. The effect of
iron-and light-limitation on phytoplankton communities of deep chlorophyll maxima of the
western Pacific Ocean. Journal of Marine Research. 2010;68(2):283-308.
54.   Marie D, Partensky F, Jacquet S, Vaulot D. Enumeration and cell cycle analysis of
natural populations of marine picoplankton by flow cytometry using the nucleic acid stain SYBR
Green I. Applied and Environmental Microbiology. 1997;63(1):186-93.
55.   Atlas EL, Hager SW, Gordon LI, Park PK. A Practical Manual for Use of the Technicon
(Trade Name) Autoanalyzer (Trade Name) in Seawater Nutrient Analyses; Revised. OREGON
STATE UNIV CORVALLIS DEPT OF OCEANOGRAPHY; 1971.
56.   Gordon LI, Jennings Jr JC, Ross AA, Krest JM. A suggested protocol for continuous flow
automated analysis of seawater nutrients (phosphate, nitrate, nitrite and silicic acid) in the
WOCE Hydrographic Program and the Joint Global Ocean Fluxes Study. WOCE hydrographic
program office, methods manual WHPO. 1993(68/91):1-52.
57.   Needham D, Fichot E, Parada A, Yeh Y-C, Fuhrman J. Fuhrman Lab 515F-926R 16S
and 18S rRNA Gene Sequencing Protocol V.2. protocolsio. 2019.

58.   Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit
rRNA primers for marine microbiomes with mock communities, time series and global field
samples. Environmental Microbiology. 2016;18(5):1403-14.
59.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source
tool for metagenomics. PeerJ. 2016;4:e2584.
60.   Eren AM, Morrison HG, Lescault PJ, Reveillaud J, Vineis JH, Sogin ML. Minimum
entropy decomposition: Unsupervised oligotyping for sensitive partitioning of high-throughput
marker gene sequences. The ISME Journal. 2015;9(4):968-79.
61.   Angly FE, Dennis PG, Skarshewski A, Vanwonterghem I, Hugenholtz P, Tyson GW.
CopyRighter: a rapid tool for improving the accuracy of microbial community profiles through
lineage-specific gene copy number correction. Microbiome. 2014;2(1):11.
62.   Morel A, Huot Y, Gentili B, Werdell PJ, Hooker SB, Franz BA. Examining the
consistency of products derived from various ocean color sensors in open ocean (Case 1) waters
in the perspective of a multi-sensor approach. Remote Sensing of Environment. 2007;111(1):69-
88.
63.   Applications NCFS, Research. GHRSST NOAA/STAR GOES-16 ABI L2P Western
Atlantic SSTskin v2.70 dataset in GDS2. NASA Physical Oceanography Distributed Active
Archive Center; 2019.
64.   Seim HE, Savidge D, Andres M, Bane J, Edwards C, Gawarkiewicz G, et al. Overview of
the processes driving exchange at Cape Hatteras program. Oceanography. 2022;35(2).

Supplemental materials:

**Table S1. Multiple regression analysis of select environmental variables on Bray-Curtis
dissimilarity for 16S rRNA gene libraries using 'envfit' in R (v 4.0.0) with 9999
permutations.**
658

659   **Table S2. Linear discriminant analysis effect size (LEfSE) results for the 250 most
660   abundant ASVs across all samples.**
661

662   **Table S3. GJAM beta-coefficients of ASVs found to have a significant association with at
663   least one SOM cluster.**
664

665   **Table S4. Linear discriminant analysis effect size (LEfSE) results for the 250 most
666   abundant ASVs in Gulf Stream and eddy comparison.**
667

Figure S1. Environmental variables for physical parameter defined water parcels (A)
Temperature and (B) salinity for each of the temperature and salinity-defined water parcels

(Continental Shelf, Continental Slope, Gulf Stream, Eddy) as determined using the flow-through
seawater system at ~1m depth. Center lines denotes the median value while the box encompasses
the 25^{th} to 75^{th} percentiles of the dataset. Whiskers denote the 5^{th} and 95^{th} percentiles and values
that fall outside these ranges are shown as individual points. Brackets indicate p-values
associated with Wilcoxon Rank Sum test pairwise comparisons.
675

Figure S2. Non-metric multidimensional scaling (NMDS) ordination comparisons of
bacterial communities grouped by physical and biological parameters. NMDS computed
based on Bray-Curtis dissimilarity for 16S rRNA gene libraries and colored by (A)
oceanographic water parcels (temperature and salinity) and (B) Self-Organizing Map (SOM)
clusters based on the microbiome. Ellipses show the multivariate t-distribution 95% confidence
interval for the mean microbial community of each water parcel or SOM cluster.
682

Figure S3. Maps of sampling sites for five transects off the coast of Cape Hatteras, North
Carolina, USA from September 5-7, 2021. Background color represents sea surface
temperatures at time of sampling obtained from the Geostationary Operational Environmental
Satellite 16 (GOES-16). Each sampling point is shown along the transects and categorized by
oceanographic water parcel (shape), defined by temperature and salinity, and microbial-
community based k-means clusters (color), generated using the self-organizing map R package
'kohonen'. Black dotted lines indicate the approximate outline of the frontal eddy during the
time of sampling based on a qualitative analysis of the SST imagery.
691

Figure S4. Environmental conditions for Self-Organizing Map (SOM) clusters. Numbered
clusters correspond to the most common environments: Continental shelf (Cluster 1), Continental
slope (Cluster 2), Gulf Stream (Cluster 3), Eddy (Cluster 4). For the box and whisker plots, the
center lines denote the median value while the box encompasses the 25^{th} to 75^{th} percentiles of the
dataset. Whiskers denote the 5^{th} and 95^{th} percentiles and values that fall outside these ranges are
shown as individual points. Brackets indicate p-values of Wilcoxon Rank Sum pairwise
comparisons.
699

Figure S5. Non-metric multidimensional scaling (NMDS) ordination computed based on
Sørensen similarity for 16S rRNA gene libraries. Each sampling point is categorized by
oceanographic water parcel (shapes), defined by temperature and salinity, and microbial-
community based k-means clusters (colors), generated using the self-organizing map R package
'kohonen'. Ellipses show the multivariate t-distribution 95% confidence interval for the mean of
each Self-Organizing Map (SOM cluster).
706

707   **Figure S6**. **Self-Organizing Map within-clusters sum of squares plot for cluster number**
708   **selection**. Calculated using a 4x4 grid with hexagonal mapping units in the 'kohonen' (v.3.0.11)
709   in R (v4.0.0).
710

Figure 3

Figure 6

Click here to access/download

Supporting Information

Gronniger_Supplementary_Materials.docx