Balancing grain yield trade-offs in 'Miracle-Wheat'

Ragavendran Abbai[1*], Guy Golan[1], C. Friedrich H. Longin[2] and Thorsten Schnurbusch[1,3*]

[1]Research Group Plant Architecture, Leibniz Institute of Plant Genetics and Crop Plant
Research (IPK), OT Gatersleben, 06466 Seeland, Germany
[2]State Plant Breeding Institute, University of Hohenheim, Fruwirthstr. 21, 70599 Stuttgart,
Germany
[3]Martin Luther University Halle-Wittenberg, Faculty of Natural Sciences III, Institute of
Agricultural and Nutritional Sciences, 06120 Halle, Germany

[*]Correspondence:
Ragavendran Abbai, Email: abbai@ipk-gatersleben.de
Thorsten Schnurbusch, Email: schnurbusch@ipk-gatersleben.de

ORCID:
Ragavendran Abbai: 0000-0002-5712-910X
Guy Golan: 0000-0002-5255-393X
C. Friedrich H. Longin: 0000-0002-0737-1651
Thorsten Schnurbusch: 0000-0002-5267-0677

Running title:
Source-Sink strength in 'Miracle-Wheat'

Word count:
Introduction-1129; Results-1528; Discussion-1493; Conclusion-177

Main Figures:
Figs. 1-7

Supplementary data:
Figs. S1-S13 & Table S1.

HIGHLIGHT

Assimilate production and reallocation potential determines grain yield in the spike-branching

'Miracle-Wheat'.

ABSTRACT

Introducing variations in inflorescence architecture, such as the 'Miracle-Wheat' (*Triticum*

turgidum convar. *compositum* (L.f.) Filat.) with a branching spike, has relevance for

enhancing wheat grain yield. However, in the spike-branching genotypes, the increase in

spikelet number is generally not translated into grain yield advantage because of reduced

spikelet fertility and grain weight. Here, we investigated if such trade-offs might be a function

of source-sink strength by using 385 RILs developed by intercrossing the spike-branching

landrace TRI 984 and CIRNO C2008, an elite durum (*T. durum* L.) cultivar; they were

genotyped using the 25K array. Various plant and spike architectural traits, including flag

leaf, peduncle and spike senescence rate, were phenotyped under field conditions for two

consecutive years. On Chr 5AL, we found a new modifier QTL for spike-branching, *branched*

head^t 3 (*bh^t-A3*), which was epistatic to the previously known *bh^t-A1* locus. Besides, *bh^t-A3*

was associated with more grains per spikelet and a delay in flag leaf senescence rate.

Importantly, favourable alleles *viz., bh^t-A3* and *grain protein content* (*gpc-B1*) that delayed

senescence are required to improve spikelet fertility and grain weight in the spike-branching

RILs. In summary, achieving a balanced source-sink relationship might minimise grain yield

trade-offs in Miracle-Wheat.

Keywords:

Grain number, Grain weight, Grain yield, Inflorescence branching, QTLs, Senescence rate,

Source-Sink strength, Trade-offs

ABBREVIATIONS

Chr: Chromosome; RILs: Recombinant inbred lines; QTL: Quantitative trait locus

INTRODUCTION

Wheat (*Triticum* sp.) inflorescence – 'Spike' is a determinate structure harbouring the grain-bearing spikelets on its rachis in a distichous pattern. During immature spike development, the inflorescence meristem gives rise to multiple spikelet meristems in an acropetal manner. In turn, each spikelet meristem (indeterminate) produces florets, that potentially form grains (Kirby and Appleyard, 1984; Koppolu and Schnurbusch, 2019). However, some exceptions deviate from this standard developmental programme, such as the 'Miracle-Wheat' that produces a non-canonical spike with lateral branches instead of spikelets. Here, due to a single amino acid substitution in the *branched head^t* (*bh^t*) allele of *T. turgidum* convar. *compositum* (L.f.) Filat. accessions, encoding an APETALA2/ETHYLENE RESPONSIVE FACTOR (AP2/ERF) transcription factor, the spikelet meristems lose their identity and determinacy while partially behaving as inflorescence meristems, producing lateral branches or multiple spikelets per rachis node (Poursarebani *et al.*, 2015). Similarly, in hexaploid wheat, variations for supernumerary spikelet formation were also found for the wheat *FRIZZY PANICLE* (*WFZP*) (Dobrovolskaya *et al.*, 2015), *Photoperiod-1 (Ppd-1)* (Boden *et al.*, 2015), *TEOSINTE BRANCHED1 (TB1)* (Dixon *et al.*, 2018), and *HOMEOBOX DOMAIN-2 (HB-2)* (Dixon *et al.*, 2022). While branching spikes have considerably higher yield potential, i.e., more spikelet number, they often suffer from grain weight trade-offs, as observed in the tetraploid Miracle-Wheat (Poursarebani *et al.*, 2015). Moreover, despite the increase in overall grain number per spike, spikelet fertility (grains per spikelet) decreased in response to spike-branching (Wolde *et al.*, 2021).

A large body of evidence suggests that wheat grain yield is an outcome of multiple trait-trait interactions mediated by developmental, physiological and environmental factors across the entire lifespan, although some stages are more critical than others (Brinton and Uauy, 2019; Guo *et al.*, 2017; Guo *et al.*, 2018a; Guo *et al.*, 2016; Guo *et al.*, 2018b; Murchie *et al.*, 2023; Reynolds *et al.*, 2022; Slafer *et al.*, 2023). They can broadly be classified as source and sink strength related, which jointly determine a particular genotype's assimilate production and reallocation potential. Typically, green tissues of the plant – both foliar (leaves) and non-foliar (peduncle, spikes) are the photosynthesising organs that act as 'source' for resource generation (Chang *et al.*, 2022; Molero and Reynolds, 2020). In the pre-anthesis phase, assimilates are partitioned to both vegetative biomass establishment and developing spikes – that determine the overall yield potential (Fischer, 2011; Slafer, 2003). The inflorescence architecture, *viz.,* spikelet number per spike, floret number per spikelet, carpel size, rachis length etc., are

determined before anthesis (Brinton and Uauy, 2019; Kirby and Appleyard, 1984; Sakuma and Schnurbusch, 2020). For instance, the ovary size during flowering regulated floret and grain survival in a panel of 30 wheat genotypes (Guo *et al.*, 2016). Likewise, the duration of leaf initiation, spikelet initiation and stem elongation period influenced spike fertility in bread wheat (Roychowdhury *et al.*, 2023). The source strength is often characterised by radiation use efficiency (RUE), i.e., the ability for light interception and biomass production (Acreche and Slafer, 2009; Molero *et al.*, 2019). However, the balance between the resources allocated to the 'vegetative vs reproductive' tissues largely dictates the yield potential (Dreccer *et al.*, 2014; Ferrante *et al.*, 2013), a trait that has been under selection throughout the history of wheat breeding. The deployment of semi-dwarf *Rht-1* alleles ('green revolution' gene) significantly increased the harvest index and the grain number per unit area, possibly by enhancing the flow of assimilates (as the stem length is considerably reduced) to the juvenile spikes (Fischer and Stockman, 1986; Slafer *et al.*, 2023). However, other strategies might currently be required to further the resource allocation to early spike development as the semi-dwarf *Rht-1* allele is already a selection target (Peng *et al.*, 1999). Increasing the harvest index in the genotypes with high biomass (more robust source) might enhance grain yield (Sierra-Gonzalez *et al.*, 2021). Overall, the source strength from the terminal spikelet stage to anthesis majorly determines grain number and size in wheat.

After anthesis, the initiation of senescence in the foliar, but also non-foliar tissues drives extensive re-mobilization of resources into the developing grains; previous studies indicated that flag leaf and spike photosynthesis contribute to most of the assimilates during the grain filling phase (Distelfeld *et al.*, 2014; Molero and Reynolds, 2020). Hence, delayed flag leaf and spike senescence resulted in extended photosynthesis (functional stay-green), leading to higher grain yield (Chapman *et al.*, 2021b; Christopher *et al.*, 2016; Hassan *et al.*, 2021; Kichey *et al.*, 2007; Li *et al.*, 2022). However, the effect of delayed senescence was not consistent; for instance, prolonged photosynthesis influenced grain yield attributes only under low nitrogen conditions (Derkx *et al.*, 2012; Gaju *et al.*, 2011). The *GPC-B1* locus encoding *NO APICAL MERISTEM (NAM)*, a *NAC* transcription factor is the major regulator of senescence rate in wheat (Uauy *et al.*, 2006); but, despite a 40% increase in flag leaf photosynthesis, the *NAM* RNAi wheat lines had no advantage in grain weight compared to the control plants (Borrill *et al.*, 2015). In addition, the stay-green phenotype of *gpc-A1* and *gpc-D1* mutants did not influence grain yield determinants (Avni *et al.*, 2014). However, (Chapman *et al.*, 2021b) reported that novel *NAM-1* allele that delayed senescence was

associated with 14% increase in the final grain weight, possibly by enhancing resource re-mobilization. A plausible explanation for such discrepancies might be that grain yield in wheat is largely sink-limited (Lichthardt *et al.*, 2020; Reynolds *et al.*, 2005); the surplus water-soluble carbons that remain in the stem at physiological maturity supports this hypothesis (Serrago *et al.*, 2013). Thus, a higher sink capacity might be essential to capitalise on the extended photosynthetic period during the grain filling phase (Lichthardt *et al.*, 2020). In this context, a reductionist approach that focusses on characterising individual component traits might assist in the deeper understanding of source-sink dynamics but also be integrated to pin-point favourable combinations of alleles/haplotypes for improving wheat grain yield (Brinton and Uauy, 2019; Reynolds *et al.*, 2022).

As Miracle-Wheat has a stronger sink (more spikelet number), we hypothesized that delimited post-anthesis source strength might explain the spike-branching induced trade-offs on spikelet fertility and grain weight. To examine this, we developed a bi-parental wheat population comprising about 385 RILs by crossing the spike-branching TRI 984 with an elite durum CIRNO C2008. The idea was to evaluate this population under field conditions for various architectural traits, as well as the senescence rate of the flag leaf, the peduncles and the spike (details are in 'Materials and Methods' section). In summary, our current study aims to explain: i. The relationship between senescence rate and trade-offs regulating grain yield (spike-branching–grain number–grain weight); ii. The underlying genetics of such trade-offs; iii. Favourable trait and allele combinations of relevant QTLs to balance grain yield trade-offs; and iv. Finally, to verify if spike-branching might be a potential selection target to enhance grain yield in wheat.

MATERIALS AND METHODS

Population development

A bi-parental population comprising 385 RILs was developed by crossing the spike-branching Miracle-Wheat accession, 'TRI 984' and elite durum from CIMMYT, 'CIRNO C2008' (hereafter referred to as 'CIRNO'). A modified speed breeding method (Ghosh *et al.*, 2018; Watson *et al.*, 2018) was used for rapid generation advancement from F_3 to F_5. Initially, the grains were sown in the 96 well trays and grown in standard long day conditions *viz.*, 16h light (19°C) and 8h dark (16°C) for about two weeks. Later, the trays were transferred to speed breeding conditions *viz.*, 22h light (22°C) and 2h dark (17°C) to accelerate the growth. The spikes were harvested at maturity, and a similar method was used for the next cycle. Finally, the obtained F_5 plants were multiplied under field conditions during the spring of 2020, and the resulting F_6 grains (RILs) were genotyped and phenotyped (Fig. S1).

Genotyping and linkage map construction

The parental lines and three F_6 grains per RIL were sown in 96 well trays and were grown in standard greenhouse conditions for about two weeks. Leaves were sampled at the two-leaf stage from all the seedlings and stored at -80°C until further use. During the sampling, the leaves from the three replications of a particular RIL were pooled, and genomic DNA was extracted. The DNA integrity was evaluated on agarose gel, after which about 50 ng/μl aliquots were prepared for the genotyping. Eventually, the 25K wheat array from SGS-TraitGenetics GmbH (https://traitgenetics.com/index.php/disclaimer/2-uncategorised) was used for genotyping the 385 RILs along with the parental lines. However, only the 18K markers scored to the A & B sub-genome were considered for further analysis; we found that 5,089 makers were polymorphic (Fig. S2A). The linkage map was developed using the regression and maximum likelihood methods in JoinMap v4.1 (Stam, 1993). A subset of 2,128 markers was prepared after filtering, *viz.*, without segregation distortion (determined based on Chi-squared test), <10% heterozygosity and <10% missing (Fig. S2B&C). Haldane's mapping function was used in the regression method, while the maximum likelihood method involved the spatial sampling thresholds of 0.1, 0.05, 0.03, 0.02 and 0.01 with three optimisation rounds per sample. Outcomes from both these methods were used to determine the 14 linkage groups and final map order.

Experimental design

Greenhouse conditions

The genotypes were sown in 96 well trays with three replication each, and the two weeks old seedlings were vernalised at 4°C for one month. Then, the seedlings were transferred to 9 cm square pots, grown in standard long day conditions (16h light; 19°C & 8h dark; 16°C), and various traits were phenotyped. Standard fertilization was performed, and plants were treated with pesticides based on the requirement.

Field conditions

The genotypes were screened at IPK-Gatersleben (51°49□23□□N, 11°17□13□□E, 112m altitude) under field conditions for two growing seasons *viz.,* the F_6 derived RILs in the spring of 2021, and the F_7 derived RILs in the spring of 2022. They were grown in an α-lattice design with three replications, while each 1.5 m^2 plot had six 20 cm spaced rows comprising two genotypes (three rows each). Standard agronomic and management practices were in place throughout the growth cycle; however, the experimental trial was completely rainfed. Besides, a subset of genotypes (about 250 F_6 derived RILs) in one replication was evaluated at the University of Hohenheim (48°42'50"N, 9°12'58"E, 400 m altitude) in 2022.

Examining plant and spike architectural traits

Plants from the inner rows (at least five measurements per plot) were considered for all the phenotyping except for grain yield per meter row, where the mean of all three rows of a particular genotype was measured. Days to heading (DTH) was determined at 'Zadoks 55', i.e. when half of the spike has emerged (Zadoks *et al.*, 1974) in about 50% of the plants in a particular plot. Later, this was converted into growing degree days (GDD) to account for temperature gradients (Miller *et al.*, 2001). The distance from the tip of the flag leaf to its base was considered as the flag leaf length, while the flag leaf width was the end-to-end horizontal distance at the middle of the leaf. Flag leaf verdancy was measured at eight different locations along the leaf (Borrill *et al.*, 2019) at heading (both in the greenhouse and field) but also at 30 days after heading (only in the greenhouse) using the SPAD-502 chlorophyll meter (Konica Minolta). In the field, flag leaf senescence was screened at 30 days after heading using a four-point severity scale from '1' indicating the least senescence to '4' for the highest senescence (Fig. S3A). The number of senesced peduncles per 10 peduncles was counted from the inner rows to determine peduncle senescence (%) (Chapman *et al.*, 2021a). In this context, we

found a gradient of yellowness in the peduncle across the RILs; however, in the current study, this was not differentiated, i.e. we had only two classes – green and yellow (Fig. S3B). Days to maturity (DTM) was determined when most spikes turned yellow in a particular plot; later, this was converted to growing degree days similar to days to heading.

Spike weight, spike length (without awns) and straw biomass (dry weight of culm along with leaves) were measured after harvest. In addition, a scoring method was developed for estimating supernumerary spikelets (two spikelets per rachis node) and spike-branching (true branching with mini-spikes from the rachis nodes) (Fig. S4). '0' (standard spike), '1' (supernumerary spikelets only at the basal part of the spike), '2' (supernumerary spikelets until half of the spike), '3' (supernumerary spikelets throughout the spike) and '4' (proper branching). Floret number was measured from the non-branching genotypes from two spikelets at the centre of the spike at harvest. Besides, derived traits such as grains per spikelet, grain filling duration (Chapman *et al.*, 2021a), and harvest index was calculated as follows:

$$Grains\ per\ spikelet = \frac{\text{Grains per spike}}{\text{Spikelet number per spike}}$$

$$Grain\ filling\ duration\ = \text{Days to maturity} - \text{Days to heading}$$

$$Harvest\ index = \frac{\text{Grain weight per spike}}{\text{Straw biomass} + \text{Spike weight}}$$

'Marvin' digital grain analyser (GTA Sensorik GmBH, Neubrandenburg, Germany) was used to determine grains per spike, thousand-grain weight, grain length, and grain width. We also recorded the grain width and length of the parental lines manually using a Vernier calliper to reconfirm the observed trend from the 'Marvin' digital seed analyser (Fig. S5). All the above-mentioned traits were recorded at IPK-Gatersleben, while only the spike architectural traits were phenotyped from the experiment conducted at the University of Hohenheim.

Phenotypic and Genetic analyses

Genstat 19 (VSN International, Hemel Hempstead, UK) and GraphPad Prism 9.3.1 (GraphPad Software, San Diego, California, USA) were used for all the statistical analyses. Ordinary

one-way ANOVA followed by Dunnett's multiple comparisons test was employed for

multiple-range comparisons, whereas an unpaired *Student's t-test* was used to compare two

groups. Pearson correlation was used to study the relationship among the traits of interest;

besides, simple linear regression assisted in understanding the effect of a particular trait

(explanatory variable) on another (response variable). The corresponding figures contain all

the relevant details, such as P-value, R^2, and the number of samples compared.

QTL mapping was performed in Genstat 19 using the following criteria: i. step size of 10 cM,

ii. minimum cofactor proximity of 50 cM, iii. minimum QTL separation distance of 30 cM

and iv. genome wide significance 'α =0.05'. Simple interval mapping (SIM) was performed

as an initial scan to determine the positions of potential candidate QTL(s). These positions

were used as cofactors for multiple rounds of composite interval mapping (CIM); CIM was

repeated until similar results were obtained at least three consecutive times. Finally, QTL

backward-selection was carried out after CIM to estimate various QTL effects, including the

determination of QTL interval, high-value allele, additive effects, and phenotypic variance

explained. The QTLs were visualised using MapChart 2.32 (Voorrips, 2002).

RESULTS

Spike-branching affects spikelet fertility and thousand-grain weight

Consistent with previous findings using different germplasm (Wolde *et al.*, 2021), despite having more spikelets, the spike-branching landrace 'TRI 984' had fewer florets per spikelet compared to the elite durum 'CIRNO' (Fig. 1A-D). However, we found no difference in grain number per five spikes, while a considerably reduced thousand-grain weight in TRI 984, associated with shorter grains was observed (Fig. 1E-H). Although CIRNO flowered earlier (Figure 1I), it had greener flag leaves at heading (Fig. 1J) and also after 30 days of heading (Fig. 1K) along with greener peduncles (Fig. 1L). Besides, CIRNO had longer but narrower flag leaves (Fig. 1M&N), fewer tillers (Fig. 1O), shorter spikes (Fig. 1P), shorter plant stature (Fig. 1Q) and less straw biomass as opposed to TRI 984. Furthermore, there was no difference in the average spike weight (n=5) (Fig. 1R), while CIRNO had more grain yield per five spikes (Fig. 1S). These observations indicate a clear difference in terms of assimilate production and resource reallocation into sink organs between the two genotypes. As expected, CIRNO, a widely cultivated modern durum variety, had delayed flag leaf and peduncle senescence (more extended grain filling period) and higher thousand-grain weight. On the other hand, the spike-branching landrace TRI 984 exhibited a relatively poor resource production and reallocation potential, *viz.,* a less verdant/green flag leaf at heading, and quicker senescence rate (shorter grain filling period), poor spikelet fertility and thousand-grain weight. Besides, the resources required to maintain the vegetative parts might be higher in the case of TRI 984 because of the taller plant architecture and longer rachis internodes than CIRNO. Hence, we phenotyped the corresponding landrace-elite recombinants (TRI 984 x CIRNO) that vary in source-sink balance to obtain mechanistic insights into the negative effect of spike-branching on spikelet fertility and grain weight, two major components of the final grain yield.

Spikelet fertility and thousand-grain weight are associated with senescence rate

As expected, we witnessed a considerable diversity for all the plant and spike architectural traits (Fig. S6). Importantly, flag leaf and peduncle senescence rates were independent of the heading date; this implies that there is a possibility for the lines that flowered late to senesce early and vice-versa (Fig. 2A). The lines with delayed flag leaf senescence also had the tendency of retaining green/verdant peduncles for a longer duration (Fig. 2A). In addition, the

intensity of flag leaf greening (SPAD meter value) at heading had only a minor effect (R^2=0.031; p=0.047) on the progress of senescence (scored at 30 days after heading), indicating that these traits are largely independent (Fig. 2B). Flag leaf length and delay in senescence were positively related (R^2=0.046; p=0.0042), while flag leaf width did not influence the same (Fig. S7A&B). Moreover, we observed that the lines with more verdant/greener flag leaves at heading (higher SPAD value) also had a more significant number of florets per spikelet (R^2=0.085; p=0.0014), in line with the expected consequence of source strength on sink organ establishment before anthesis (Fig. S7C). Intriguingly, the number of florets and grains per spikelet, which is determined earlier, was associated with senescence rate, i.e., the lines with more florets and grains per spikelet displayed delayed flag leaf senescence (R^2=0.071; p=0.0009 & R^2=0.16; p<0.0001) (Figs. S7D & 2C). We mapped a QTL on Chr 5A (bh^t-A3) influencing grains per spikelet and flag leaf/peduncle senescence rate, which explains the underlying genetic basis of such an exciting relationship (Table S1). This trend implies a plausible pleiotropic regulation that requires further validation. Besides, the delayed senescence rate had a positive effect on thousand-grain weight (R^2=0.11; p<0.0001) (Fig. 2D). We realised that the observed increase in thousand-grain weight is primarily due to the change in grain width (R^2=0.097; p<0.0001) (Fig. 2E) and not grain length (Fig. S7E), implying that grain width is more plastic, influenced by resource reallocation compared to grain length. Nevertheless, it is clear that the longer duration of green flag leaf and peduncle is not simply 'cosmetic' – it influences grain yield determinants. This vital evidence supports our hypothesis that dissecting the source-sink relationship might have relevance in balancing the trade-offs that negatively regulate the final grain yield in 'Miracle-Wheat' like genotypes.

Genetic basis of source-sink dynamics in 'Miracle-Wheat'

The bh^t-A1 locus underlies sink and source capacity

Using a dosage-based scoring method (Fig. S4), we mapped a major effect QTL for spike-branching on Chr 2A (Fig. 3A, Table S1) that was tightly linked with the previously known locus bh^t-A1 (Poursarebani *et al*., 2015). Regardless of the increase in spikelet number per spike owing to the lateral branching (Fig. 3B), there was no difference in the total grain number per spike (Fig. 3C). Moreover, the bh^t-A1 locus while inducing spike-branching, was also associated with a reduction in grain length (Fig. 3D) and thousand-grain weight (Fig. 3E). Besides, we found that flag leaf verdancy at heading was negatively affected (Fig. 3F), as

with spike length (Fig. 3G). In principle, the TRI 984 allele at the bh^t-A1 locus induces spike-branching, but with a possible drawback on the source capacity. Overall, the spike-branching effect from bh^t-A1 locus could not be translated into any advantages for the final grain yield.

bh^t-A3, a novel spike-branching locus on Chr 5A reshapes source-sink dynamics

We named the newly identified spike-branching modifier locus as 'bh^t-A3' following the previously known bh^t-A1 (Poursarebani et al., 2015) and bh^t-A2 (Wolde et al., 2021) loci. Interestingly, the spike-branching effect of the bh^t-A3 locus (contributed by the CIRNO allele) manifests only in the presence of the mutated bh^t-A1 allele (Fig. 4A; Fig. S8A-C; Table S1). We divided the RILs into two sub-groups for QTL mapping viz., by fixing i. bh^t-A1, ii. BH^t-A1 and the outcome confirm the epistasis of the bh^t-A3 to bh^t-A1 locus (Fig. S8A). Possibly, this indicates that the plasticity for spike-branching is introduced by bh^t-A1, i.e., it might be first essential to have bh^t-A1 to disrupt the spikelet meristem identity and only then the bh^t-A3 locus might modify the branching intensity in the spikes. Moreover, in this region, we found co-localised QTLs for an array of traits influencing source-sink dynamics. The CIRNO allele contributed to spike-branching (Fig. 4B), delayed flag leaf senescence (Fig. 4C), more extended grain filling period (Fig. 4D), increased spikelet fertility (Fig. 4E) and grain yield per five spikes (Fig. 4F). Besides, we also found a subtle, yet positive effect on grain width (Fig. S9A), thousand-grain weight (Fig. 4G), florets per spikelet (Fig. S9B), straw biomass (Fig. S9C) and harvest index (Fig. S9D). Interestingly, we found that the observed variations in flag leaf senescence and thousand-grain weight were not dependent on the presence of bh^t-A1 (Fig. S8D&E). This pattern might imply that the phenotypic variation explained by the 5A QTL hotspot for spike-branching rate and senescence might be the outcome of at least two different genes. Nevertheless, the more extended photosynthetic period translated into grain number increases only in the spike-branching RILs – when bh^t-A1 is present (Fig. S8C). Taken together, this trend suggests that the favourable CIRNO allele (bh^t-A3) mediates enhanced assimilate production and reallocation of the resources to sink organs, including the lateral branches/supernumerary spikelets because of longer grain filling duration.

GPC-B1 is the major determinant of senescence rate and thousand-grain weight

A QTL on Chr 6B, which most likely is associated with GPC-B1 (Uauy et al., 2006), explained most of the observed phenotypic variance for the overall plant senescence rate (Fig. 5A; Table S1). Likewise, it was found that mutations in the NAC domain of NAM-A1 (GPC-A1) delayed peduncle and flag leaf senescence (Harrington et al., 2019). In the current study,

the CIRNO allele ensured delay in the flag leaf (Fig. 5B), peduncle (Fig. 5C) and spike senescence (days to maturity) (Fig. 5D). Therefore, there might be a possibility of more reallocation into the sink organs, leading to increase in grain width (Fig. 5E) and grain length (Fig. 5F). Accordingly, we observed a considerably higher thousand-grain weight in the RILs that senesce late (Fig. 5G). However, there was no meaningful difference in grain number per five spikes (Fig. S10A), straw biomass (Fig. S10B) and harvest index (Fig. S10C). Furthermore, our interaction analysis revealed that both bh^t-A1 (Fig. S11A-D) and bh^t-A3 (Fig. S12A-D) might function independent of gpc-B1.

Specific additive and epistatic interactions may increase yield potential in spike-branching genotypes

As the QTLs on Chr 2A, 5A and 6B explain variations in key source-sink attributes, we analysed their various allelic combinations to understand better the trade-off among spike-branching, spikelet fertility and thousand-grain weight (Figs. 6A-H & 7). Interestingly, the spike-branching lines carrying bh^t-A1 and bh^t-A3 loci along with gpc-B1 had higher grain number per five spikes (Fig. 6A, E) and were associated with a delay in post-anthesis flag leaf senescence. Eventually, they had higher thousand grain weight (Fig. 6B), higher grain yield per five spikes (Fig. 6C, G) and enhanced grain yield (per meter row) (Fig. 6D) as opposed to the early senescing branched spike RILs (bh^t-A1+BH^t-A3+GPC-B1) across all the three environments $viz.$, IPK-2021, IPK-2022 (Fig. 6A-D) and University of Hohenheim-2022 (Fig. 6E & G). However, the difference in thousand-grain weight was observed only at IPK (Figs. 6B & S13A, B), while this effect was absent in Hohenheim (Figs. 6F & S13C, D).

DISCUSSION

Over the course of domestication and breeding, grain yield determinants such as grain number and grain weight, but also grain quality traits under both favourable and stressful conditions, were the primary selection targets in all major cereal crops, including wheat (McSteen and Kellogg, 2022; Voss-Fels *et al.*, 2019). For instance, the selection of the semi-dwarf *Rht-1* allele was a vital driver of the 'green revolution' in wheat (Peng *et al.*, 1999); likewise, the prevalence of the less functional *GNI-A1* allele enabled higher floret fertility in the modern wheat cultivars (Golan *et al.*, 2019; Sakuma *et al.*, 2019). However, substantial genetic yield gaps [the difference between the genetic yield potential of a crop in a particular environment to that of the potential yield of the current local cultivar] suggest the presence of untapped genetic diversity for enhancing wheat grain yield (Senapati *et al.*, 2022). Grain yield can be optimised by fine-tuning various developmental processes (Mathan *et al.*, 2016) and introducing 'drastic variations' in crop breeding (Abbai *et al.*, 2020). The genetic pathways that coordinate inflorescence architecture are dissected in staple grasses (Kellogg, 2022; Koppolu *et al.*, 2022), which might have relevance for minimising the genetic yield gap.

Here, we considered the case of spike-branching Miracle-Wheat as a potential option for increasing sink strength (more spikelets and grains per spike). However, the genetic analysis of the TRI 984 x CIRNO recombinants revealed a couple of significant limitations. Firstly, we recorded inconsistencies in the expressivity (degree) of spike-branching (in the RILs that carried similar QTLs/alleles; Figs. 3B, 4B & S8B). Although final grain yield is the function of various events, it is conceivable that the relevant source-related component traits in the pre-anthesis (yield construction) phase (Murchie *et al.*, 2023; Slafer *et al.*, 2023) might play a significant role in determining the yield potential. Expectedly, we observed that the RILs that flowered late were the ones with more straw biomass and, in turn, showed increased spike-branching in the presence of relevant QTLs (Fig. S7F-H). Hence, a longer duration of the pre-anthesis phase might have enabled increased resource production and partitioning into the developing juvenile spikes, resulting in better expressivity of the spike-branching phenotype. This implies that spike-branching winter wheat might have higher yield potential given the longer duration of pre-anthesis phase, suggesting a possibility of extending the current findings to other populations. However, we also observed differences in the degree of spike-branching within the same genotype. Similarly, in an earlier study (Wolde *et al.*, 2021) reported that the expressivity of spike-branching in a particular genotype was higher in the outer rows as opposed to the inner rows of the plot. However, no new QTLs were mapped that

specifically explained such differences. In principle, field-grown plants experience

competition for various resources, including light (Huber *et al.*, 2021; Postma *et al.*, 2021),

especially in the inner rows (Rebetzke *et al.*, 2014). In this regard, future studies investigating

the response of various source and sink component traits in high-density plots or simulated

canopy shade (Golan *et al.*, 2022) are required to uncover the genetic framework of plant-

plant competition and its effect on spike-branching expressivity.

Next, (Poursarebani *et al.*, 2015) reported that the *bh^t-A1* locus increases grain number, but

with a grain weight trade-off. Likewise, we also observed considerably smaller grains in the

spike-branching genotypes (Figs. 1F&3E). However, in the current study, the *bh^t-A1* locus

showed no increase in the final grain number because of the spike-branching, suggesting poor

spikelet fertility (Figs. 1E&3C). Interestingly, there was no thousand-grain weight trade-off in

the spike-branching Bellaroi x TRI19165 semi-dwarf RILs (Wolde *et al.*, 2021) and also in

the Floradur NILs with supernumerary spikelets (Wolde *et al.*, 2019); thus, warranting the

analysis of source-sink dynamics in the non-canonical spike forms. Here, it is vital to

emphasise the relevance of the post-anthesis (yield realisation) events, chiefly related to the

transfer of assimilates to the previously established sink organs during grain filling (Murchie

et al., 2023; Slafer *et al.*, 2023). In this context, the senescence rate might have an impact on

grain filling duration (Chapman *et al.*, 2021b; Christopher *et al.*, 2016; Hassan *et al.*, 2021;

Kichey *et al.*, 2007; Li *et al.*, 2022), i.e., extended photosynthesis leading to more assimilate

production and allocation to the developing grains. But, final grain weight was not strongly

related with starch/sugar levels or the corresponding enzymatic capacity in 54 diverse wheat

genotypes, but it might be a function of early developmental events (Fahy *et al.*, 2018). In the

current study, we report that higher grain number per spikelet (Fig. 2C) and grain weight (Fig.

2D) is associated with delayed flag leaf, peduncle and spike senescence. As expected, the

observed effect of senescence rate might be because of the differences in various sink

strength-related traits such as rachis length, spikelet number per spike (spike-branching), and

floret number per spikelet in our RIL population (Fig. S6). As the sink strength increased,

perhaps the extended photosynthetic period was meaningful for influencing the final grain

yield. This trend further establishes the rationale for understanding the genetic and molecular

framework of source and sink-related component traits to enable grain yield gains (Brinton

and Uauy, 2019; Reynolds *et al.*, 2022). With this, the favourable alleles explaining the

source-sink dynamics might assist in balancing the trade-offs among spikelet fertility and

grain weight in the spike-branching genotypes. Here, we analysed the interactions among *bh^t-*

A1, *bhᵗ-A3,* and *gpc-B1*; the *bhᵗ-A1* and *bhᵗ-A3* loci regulated spike-branching, but also source strength, while *gpc-B1* delayed senescence rate and increased thousand-grain weight (Figs. 6&7). Transcriptional analysis of WT and *NAM* (*GPC*) RNAi lines revealed differential regulation of genes related to various processes, including photosynthesis and nitrogen metabolism, during flag leaf senescence (Andleeb *et al.*, 2022). Our preliminary genetic evidence indicates that *bhᵗ-A1* and *gpc-B1* function independently (Fig. S11); however, it might be interesting to verify the presence of any possible common downstream targets of *bhᵗ-A1* and *gpc-B1* to uncover subtler aspects of their regulation. In any case, as speculated, the spike-branching RILs with an extended photosynthetic period (delayed senescence) had considerably higher grain yield (per meter row) as opposed to branched spike genotypes that senesced early (Fig. 6D). In this case, the stay-green spike-branching RILs were associated with 15.82% (SEM: ±5.96%) more grains per spike (Fig. 6A). However, we believe that the grain number difference might be due to the interaction between floret number and flag leaf senescence, which is mediated by the *bhᵗ-A3* locus; the CIRNO allele increased florets per spikelet (Fig. S9B) and delayed flag leaf senescence (Fig. 4C). The pre-anthesis floret degeneration and post-anthesis flag leaf senescence might share a common genetic basis thereby primarily affecting the tip of the respective organs, i.e. spikelet meristem/rachilla and flag leaf, respectively. Therefore, it is conceivable that the underlying gene might have a pleiotropic effect on floret surivival and flag leaf senescence, thus explaining the grain number difference. Eventually at IPK-Gatersleben (2021 and 2022), we found a 9.35% (SEM: ±3.58%) increase in average grain weight (Fig. 6B) in the spike-branching genotypes that senesce late. The 2.61% (SEM: ±0.91%) rise in grain width (Fig. S13A) majorly contributed to the grain weight difference, as the grain length remained unaffected (Fig. S13B). Incidentally, it was found that grain width increased during wheat evolution under domestication (Gegas *et al.*, 2010). Besides, it might be interesting to evaluate the effect of expansin genes in the spike-branching lines as the ectopic expression of *TaExpA6* increased grain length (Calderini *et al.*, 2021).

However, we would like to emphasise certain limitations in our experimental setup: we used relatively small plots (only 1.5 m^2) with two genotypes in one plot; therefore, the influence of the border effect (Rebetzke *et al.*, 2014) cannot be excluded in grain yield per row calculations and besides, the evaluated population are landrace-elite recombinants, that might create another bias in the observed yield increase. Although there is a significant increase in grain number per five spikes in the stay-green spike-branching recombinants, the actual yield

advantage might be better understood by evaluating the effect in isogenic backgrounds (NILs) and larger plots in multiple environments. In this context, we are developing spike-branching CIRNO NILs for these follow-up experiments. Another trade-off associated with extending the grain filling duration that is not addressed here is its likely impact on grain nutrition profile; the functional *NAM-B1* allele improves grain protein, iron and zinc content by accelerating the senescence process (Uauy *et al.*, 2006). Then, the status of the stay-green spike-branching RILs under unfavourable conditions is also beyond the scope of the current study; however, previous reports indicate a positive effect of stay-green phenotypes on wheat grain yield under drought and heat (Lopes and Reynolds, 2012). Similarly, delay in senescence led to higher grain number and tiller number but lower thousand-grain weight under nitrogen-limiting conditions (Derkx *et al.*, 2012). In addition, a recent simulation study indicates the advantage of cultivating late-maturing wheat varieties in future climate scenarios (Minoli *et al.*, 2022), implying that a delay in senescence rate might eventually be beneficial.

CONCLUSION

The physiological and genetic analysis of TRI 984xCIRNO recombinants revealed that i. extended verdant flag leaf, peduncle and spike led to higher grain yield per spike as the traits influencing sink strength segregated, including spike-branching; ii. we identified three QTL regions–on Chr 2A (*bh^t-A1*), Chr 5A (*bh^t-A3*) and Chr 6B (*gpc-B1*) that regulated source-sink strength in the current bi-parental population; iii. upon analysing their various allele combinations, it was found that an increase in grain number and grain weight is predominantly possible among the stay-green, spike-branching genotypes. iv. Finally, as wheat grain yield is also sink-limited, we propose that introducing spike-branching as a breeding target might enable advancing genetic gains while minimising the gap between genetic yield potential and the actual realised yield. Although we provide insights into balancing spike-branching–spikelet fertility–grain weight trade-offs, it is still necessary to understand the basis of inconsistencies in the degree of spike-branching within the same genotype but also in diverse genetic backgrounds. To achieve this, tracking the source-strength dynamics during the early developmental stages might be necessary.

ACKNOWLEDGEMENT

We are grateful to Franziska Backhaus, Corinna Trautewig, Kerstin Wolf, Sonja Allner, Ellen Weiss, Ingrid Marscheider, and Angelika Püschel for their excellent technical assistance; Roop Kamal for helping with the field design; Prof. Nils Stein for his valuable comments on the project; Dr. Gemma Molero for sharing CIRNO grains; all members of Plant Architecture research group for the fruitful discussions and also for the support during harvest; Peter Schreiber and the team of gardeners for managing the field operations.

AUTHOR CONTRIBUTIONS

TS acquired funding and supervised the project. RA continued to develop the population further, generated the data, analysed and interpreted the results. TS and GG guided in analysis and interpretation of the results. FHL conducted the field experiment at the University of Hohenheim. RA wrote the manuscript with inputs from all the co-authors.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

FUNDING

TS thanks the European Fund for Regional Development (EFRE), the State of Saxony-Anhalt within the ALIVE project, grant no. ZS/2018/09/94616, the HEISENBERG Program of the German Research Foundation (DFG), grant no. SCHN 768/15-1 and the IPK core budget for supporting this study. GG was funded through the Alexander von Humboldt Foundation postdoctoral fellowship program.

REFERENCES

Abbai R, Singh VK, Snowdon RJ, Kumar A, Schnurbusch T. 2020. Seeking Crops with Balanced Parts for the Ideal Whole. Trends in Plant Science **25**, 1189-1193.

Acreche MM, Slafer GA. 2009. Grain weight, radiation interception and use efficiency as affected by sink-strength in Mediterranean wheats released from 1940 to 2005. Field Crops Research **110**, 98-105.

Andleeb T, Knight E, Borrill P. 2022. Wheat NAM genes regulate the majority of early monocarpic senescence transcriptional changes including nitrogen remobilization genes. G3 Genes|Genomes|Genetics.

Avni R, Zhao R, Pearce S, Jun Y, Uauy C, Tabbita F, Fahima T, Slade A, Dubcovsky J, Distelfeld A. 2014. Functional characterization of GPC-1 genes in hexaploid wheat. Planta **239**, 313-324.

Boden SA, Cavanagh C, Cullis BR, Ramm K, Greenwood J, Jean Finnegan E, Trevaskis B, Swain SM. 2015. Ppd-1 is a key regulator of inflorescence architecture and paired spikelet development in wheat. Nature Plants **1**, 14016.

Borrill P, Fahy B, Smith AM, Uauy C. 2015. Wheat grain filling is limited by grain filling capacity rather than the duration of flag leaf photosynthesis: a case study using NAM RNAi plants. PLoS One **10**, e0134947.

Borrill P, Harrington SA, Simmonds J, Uauy C. 2019. Identification of Transcription Factors Regulating Senescence in Wheat through Gene Regulatory Network Modelling. Plant Physiol **180**, 1740-1755.

Brinton J, Uauy C. 2019. A reductionist approach to dissecting grain weight and yield in wheat. J Integr Plant Biol **61**, 337-358.

Calderini DF, Castillo FM, Arenas□M A, Molero G, Reynolds MP, Craze M, Bowden S, Milner MJ, Wallington EJ, Dowle A. 2021. Overcoming the trade□off between grain weight and number in wheat by the ectopic expression of expansin in developing seeds leads to increased yield potential. New Phytologist **230**, 629-640.

Chang TG, Shi Z, Zhao H, Song Q, He Z, Van Rie J, Den Boer B, Galle A, Zhu XG. 2022. 3dCAP-Wheat: An Open-Source Comprehensive Computational Framework Precisely Quantifies Wheat Foliar, Nonfoliar, and Canopy Photosynthesis. Plant Phenomics **2022**, 9758148.

Chapman EA, Orford S, Lage J, Griffiths S. 2021a. Capturing and Selecting Senescence Variation in Wheat. Front Plant Sci **12**, 638738.

Chapman EA, Orford S, Lage J, Griffiths S. 2021b. Delaying or delivering: identification of novel NAM-1 alleles that delay senescence to extend wheat grain fill duration. Journal of experimental botany **72**, 7710-7728.

Christopher JT, Christopher MJ, Borrell AK, Fletcher S, Chenu K. 2016. Stay-green traits to improve wheat adaptation in well-watered and water-limited environments. Journal of experimental botany **67**, 5159-5172.

Derkx AP, Orford S, Griffiths S, Foulkes MJ, Hawkesford MJ. 2012. Identification of differentially Senescing mutants of wheat and impacts on yield, biomass and nitrogen Partitioning F. Journal of integrative plant biology **54**, 555-566.

Distelfeld A, Avni R, Fischer AM. 2014. Senescence, nutrient remobilization, and yield in wheat and barley. Journal of experimental botany **65**, 3783-3798.

Dixon LE, Greenwood JR, Bencivenga S, Zhang P, Cockram J, Mellers G, Ramm K, Cavanagh C, Swain SM, Boden SA. 2018. TEOSINTE BRANCHED1 Regulates Inflorescence Architecture and Development in Bread Wheat (Triticum aestivum). Plant Cell **30**, 563-581.

Dixon LE, Pasquariello M, Badgami R, Levin KA, Poschet G, Ng PQ, Orford S, Chayut N, Adamski NM, Brinton J, Simmonds J, Steuernagel B, Searle IR, Uauy C, Boden SA. 2022. MicroRNA-resistant alleles of HOMEOBOX DOMAIN-2 modify inflorescence branching and increase grain protein content of wheat. Sci Adv **8**, eabn5907.

Dobrovolskaya O, Pont C, Sibout R, Martinek P, Badaeva E, Murat F, Chosson A, Watanabe N, Prat E, Gautier N, Gautier V, Poncet C, Orlov YL, Krasnikov AA, Berges H, Salina E, Laikova L, Salse J. 2015. FRIZZY PANICLE drives supernumerary spikelets in bread wheat. Plant Physiol **167**, 189-199.

Dreccer MF, Wockner KB, Palta JA, McIntyre CL, Borgognone MG, Bourgault M, Reynolds M, Miralles DJ. 2014. More fertile florets and grains per spike can be achieved at higher temperature in wheat lines with high spike biomass and sugar content at booting. Functional Plant Biology **41**, 482-495.

Fahy B, Siddiqui H, David LC, Powers SJ, Borrill P, Uauy C, Smith AM. 2018. Final grain weight is not limited by the activity of key starch-synthesising enzymes during grain filling in wheat. Journal of experimental botany **69**, 5461-5475.

Ferrante A, Savin R, Slafer GA. 2013. Floret development and grain setting differences between modern durum wheats under contrasting nitrogen availability. Journal of experimental botany **64**, 169-184.

Fischer R. 2011. Wheat physiology: a review of recent developments. Crop and Pasture Science **62**, 95-114.

Fischer R, Stockman Y. 1986. Increased kernel number in Norin 10-derived dwarf wheat: evaluation of the cause. Functional Plant Biology **13**, 767-784.

Gaju O, Allard V, Martre P, Snape J, Heumez E, LeGouis J, Moreau D, Bogard M, Griffiths S, Orford S. 2011. Identification of traits to improve the nitrogen-use efficiency of wheat genotypes. Field Crops Research **123**, 139-152.

Gegas VC, Nazari A, Griffiths S, Simmonds J, Fish L, Orford S, Sayers L, Doonan JH, Snape JW. 2010. A Genetic Framework for Grain Size and Shape Variation in Wheat The Plant Cell **22**, 1046-1056.

Ghosh S, Watson A, Gonzalez-Navarro OE, Ramirez-Gonzalez RH, Yanes L, Mendoza-Suarez M, Simmonds J, Wells R, Rayner T, Green P, Hafeez A, Hayta S, Melton RE, Steed A, Sarkar A, Carter J, Perkins L, Lord J, Tester M, Osbourn A, Moscou MJ, Nicholson P, Harwood W, Martin C, Domoney C, Uauy C, Hazard B, Wulff BBH, Hickey LT. 2018. Speed breeding in growth chambers and glasshouses for crop breeding and model plant research. Nat Protoc **13**, 2944-2963.

Golan G, Abbai R, Schnurbusch T. 2022. Exploring the trade-off between individual fitness and community performance of wheat crops using simulated canopy shade. Plant, Cell & Environment.

Golan G, Ayalon I, Perry A, Zimran G, Ade-Ajayi T, Mosquna A, Distelfeld A, Peleg Z. 2019. GNI-A1 mediates trade-off between grain number and grain weight in tetraploid wheat. Theoretical and Applied Genetics **132**, 2353-2365.

Guo Z, Chen D, Alqudah AM, Röder MS, Ganal MW, Schnurbusch T. 2017. Genome☐wide association analyses of 54 traits identified multiple loci for the determination of floret fertility in wheat. New Phytologist **214**, 257-270.

Guo Z, Liu G, Röder MS, Reif JC, Ganal MW, Schnurbusch T. 2018a. Genome☐wide association analyses of plant growth traits during the stem elongation phase in wheat. Plant Biotechnology Journal **16**, 2042-2052.

Guo Z, Slafer GA, Schnurbusch T. 2016. Genotypic variation in spike fertility traits and ovary size as determinants of floret and grain survival rate in wheat. Journal of experimental botany **67**, 4221-4230.

Guo Z, Zhao Y, Röder MS, Reif JC, Ganal MW, Chen D, Schnurbusch T. 2018b. Manipulation and prediction of spike morphology traits for the improvement of grain yield in wheat. Scientific reports **8**, 14435.

Harrington SA, Overend LE, Cobo N, Borrill P, Uauy C. 2019. Conserved residues in the wheat (Triticum aestivum) NAM-A1 NAC domain are required for protein binding and when mutated lead to delayed peduncle and flag leaf senescence. BMC Plant Biology **19**, 407.

Hassan MA, Yang M, Rasheed A, Tian X, Reynolds M, Xia X, Xiao Y, He Z. 2021. Quantifying senescence in bread wheat using multispectral imaging from an unmanned aerial vehicle and QTL mapping. Plant physiology **187**, 2623-2636.

Huber M, Nieuwendijk NM, Pantazopoulou CK, Pierik R. 2021. Light signalling shapes plant–plant interactions in dense canopies. Plant, Cell & Environment **44**, 1014-1029.

Kellogg EA. 2022. Genetic control of branching patterns in grass inflorescences. Plant Cell **34**, 2518-2533.

Kichey T, Hirel B, Heumez E, Dubois F, Le Gouis J. 2007. In winter wheat (Triticum aestivum L.), post-anthesis nitrogen uptake and remobilisation to the grain correlates with agronomic traits and nitrogen physiological markers. Field Crops Research **102**, 22-32.

Kirby E, Appleyard M. 1984. Cereal development guide, 2nd edn. Stoneleigh: National Agricultural Centre Arable Unit.

Koppolu R, Chen S, Schnurbusch T. 2022. Evolution of inflorescence branch modifications in cereal crops. Curr Opin Plant Biol **65**, 102168.

Koppolu R, Schnurbusch T. 2019. Developmental pathways for shaping spike inflorescence architecture in barley and wheat. J Integr Plant Biol **61**, 278-295.

Li H, Liu H, Hao C, Li T, Liu Y, Wang X, Yang Y, Zheng J, Zhang X. 2022. The auxin response factor TaARF15-A1 negatively regulates senescence in common wheat (Triticum aestivum L.). Plant physiology.

Lichthardt C, Chen T-W, Stahl A, Stützel H. 2020. Co-evolution of sink and source in the recent breeding history of winter wheat in Germany. Frontiers in plant science **10**, 1771.

Lopes MS, Reynolds MP. 2012. Stay-green in spring wheat can be determined by spectral reflectance measurements (normalized difference vegetation index) independently from phenology. Journal of experimental botany **63**, 3789-3798.

Mathan J, Bhattacharya J, Ranjan A. 2016. Enhancing crop yield by optimizing plant developmental features. Development **143**, 3283-3294.

McSteen P, Kellogg EA. 2022. Molecular, cellular, and developmental foundations of grass diversity. Science **377**, 599-602.

Miller P, Lanier W, Brandt S. 2001. Using growing degree days to predict plant stages. Ag/Extension Communications Coordinator, Communications Services, Montana State University-Bozeman, Bozeman, MO.

Minoli S, Jägermeyr J, Asseng S, Urfels A, Müller C. 2022. Global crop yields can be lifted by timely adaptation of growing periods to climate change. Nature Communications **13**, 1-10.

Molero G, Joynson R, Pinera☐Chavez FJ, Gardiner LJ, Rivera☐Amado C, Hall A, Reynolds MP. 2019. Elucidating the genetic basis of biomass accumulation and radiation use efficiency in spring wheat and its role in yield potential. Plant Biotechnology Journal **17**, 1276-1288.

Molero G, Reynolds MP. 2020. Spike photosynthesis measured at high throughput indicates genetic variation independent of flag leaf photosynthesis. Field Crops Research **255**, 107866.

Murchie EH, Reynolds M, Slafer GA, Foulkes MJ, Acevedo-Siaca L, McAusland L, Sharwood R, Griffiths S, Flavell RB, Gwyn J, Sawkins M, Carmo-Silva E. 2023. A 'wiring diagram' for source strength traits impacting wheat yield potential. J Exp Bot **74**, 72-90.

Peng J, Richards DE, Hartley NM, Murphy GP, Devos KM, Flintham JE, Beales J, Fish LJ, Worland AJ, Pelica F. 1999. 'Green revolution'genes encode mutant gibberellin response modulators. Nature **400**, 256-261.

Postma JA, Hecht VL, Hikosaka K, Nord EA, Pons TL, Poorter H. 2021. Dividing the pie: A quantitative review on plant density responses. Plant, Cell & Environment **44**, 1072-1094.

Poursarebani N, Seidensticker T, Koppolu R, Trautewig C, Gawronski P, Bini F, Govind G, Rutten T, Sakuma S, Tagiri A, Wolde GM, Youssef HM, Battal A, Ciannamea S, Fusca T, Nussbaumer T, Pozzi C, Borner A, Lundqvist U, Komatsuda T, Salvi S, Tuberosa R, Uauy C, Sreenivasulu N, Rossini L, Schnurbusch T. 2015. The Genetic Basis of Composite Spike Form in Barley and 'Miracle-Wheat'. Genetics **201**, 155-165.

Rebetzke GJ, Fischer RTA, Van Herwaarden AF, Bonnett DG, Chenu K, Rattey AR, Fettell NA. 2014. Plot size matters: Interference from intergenotypic competition in plant phenotyping studies. Functional Plant Biology **41**, 107-118.

Reynolds M, Pellegrineschi A, Skovmand B. 2005. Sink☐limitation to yield and biomass: a summary of some investigations in spring wheat. Annals of Applied Biology **146**, 39-49.

Reynolds M, Slafer G, Foulkes J, Griffiths S, Murchie E, Carmo-Silva E, Asseng S, Chapman S, Sawkins M, Gwyn J, Flavell R. 2022. A wiring diagram to integrate physiological traits of wheat yield potential. Nat. Food **3**, 318-324.

Roychowdhury R, Zilberman O, Chandrasekhar K, Curzon AY, Nashef K, Abbo S, Slafer GA, Bonfil DJ, Ben-David R. 2023. Pre-anthesis spike growth dynamics and its association to yield components among elite bread wheat cultivars (Triticum aestivum L. spp.) under Mediterranean climate. bioRxiv.

Sakuma S, Golan G, Guo Z, Ogawa T, Tagiri A, Sugimoto K, Bernhardt N, Brassac J, Mascher M, Hensel G, Ohnishi S, Jinno H, Yamashita Y, Ayalon I, Peleg Z, Schnurbusch T, Komatsuda T. 2019. Unleashing floret fertility in wheat through the mutation of a homeobox gene. Proceedings of the National Academy of Sciences **116**, 5182-5187.

Sakuma S, Schnurbusch T. 2020. Of floral fortune: tinkering with the grain yield potential of cereal crops. New Phytol **225**, 1873-1882.

Senapati N, Semenov MA, Halford NG, Hawkesford MJ, Asseng S, Cooper M, Ewert F, van Ittersum MK, Martre P, Olesen JE. 2022. Global wheat production could benefit from closing the genetic yield gap. Nature Food **3**, 532-541.

Serrago RA, Alzueta I, Savin R, Slafer GA. 2013. Understanding grain yield responses to source–sink ratios during grain filling in wheat and barley under contrasting environments. Field Crops Research **150**, 42-51.

Sierra-Gonzalez A, Molero G, Rivera-Amado C, Babar MA, Reynolds MP, Foulkes MJ. 2021. Exploring genetic diversity for grain partitioning traits to enhance yield in a high biomass spring wheat panel. Field Crops Research **260**, 107979.

Slafer G. 2003. Genetic basis of yield as viewed from a crop physiologist's perspective. Annals of Applied Biology **142**, 117-128.

Slafer GA, Foulkes MJ, Reynolds MP, Murchie EH, Carmo-Silva E, Flavell R, Gwyn J, Sawkins M, Griffiths S. 2023. A 'wiring diagram' for sink strength traits impacting wheat yield potential. J Exp Bot **74**, 40-71.

Stam P. 1993. Construction of integrated genetic linkage maps by means of a new computer package: Join Map. Plant J **3**, 739-744.

Uauy C, Distelfeld A, Fahima T, Blechl A, Dubcovsky J. 2006. A NAC gene regulating senescence improves grain protein, zinc, and iron content in wheat. Science **314**, 1298-1301.

Voorrips RE. 2002. MapChart: software for the graphical presentation of linkage maps and QTLs. J Hered **93**, 77-78.

Voss-Fels KP, Stahl A, Wittkop B, Lichthardt C, Nagler S, Rose T, Chen T-W, Zetzsche H, Seddig S, Majid Baig M. 2019. Breeding improves wheat productivity under contrasting agrochemical input levels. Nature Plants **5**, 706-714.

Watson A, Ghosh S, Williams MJ, Cuddy WS, Simmonds J, Rey MD, Asyraf Md Hatta M, Hinchliffe A, Steed A, Reynolds D, Adamski NM, Breakspear A, Korolev A, Rayner T, Dixon LE, Riaz A, Martin W, Ryan M, Edwards D, Batley J, Raman H, Carter J, Rogers C, Domoney C, Moore G, Harwood W, Nicholson P, Dieters MJ, DeLacy IH, Zhou J, Uauy C, Boden SA, Park RF, Wulff BBH, Hickey LT. 2018. Speed breeding is a powerful tool to accelerate crop research and breeding. Nat Plants **4**, 23-29.

Wolde GM, Mascher M, Schnurbusch T. 2019. Genetic modification of spikelet arrangement in wheat increases grain number without significantly affecting grain weight. Molecular Genetics and Genomics **294**, 457-468.

Wolde GM, Schreiber M, Trautewig C, Himmelbach A, Sakuma S, Mascher M, Schnurbusch T. 2021. Genome-wide identification of loci modifying spike-branching in tetraploid wheat. Theor Appl Genet **134**, 1925-1943.

Zadoks J, Chang T, Konzak C. 1974. Decimal code for the growth-stages of cereals. Weed Res. **14**, 415-421.

Main figure legends

Fig. 1. TRI 984 has a poor source-sink balance as opposed to CIRNO. (A) Spike-branching phenotype of TRI 984 and (B) Standard spike of CIRNO. (C) Spikelet from the central part of the TRI 984 main rachis shows reduced florets relative to (D) a spikelet from a similar position in CIRNO. (E) Despite having more spikelets, there is no increase in grain number per 5 spikes in TRI 984. (F-H) However, the grains are smaller in TRI 984, leading to a reduction in thousand-grain weight. (I) TRI 984 exhibited delayed heading, (J) lower flag leaf verdancy at heading, (K, L) accelerated flag leaf and peduncle senescence. TRI 984 had (M, N) shorter but wider flag leaves, (O) more tillers at booting (under greenhouse conditions), (P) longer spikes, (Q) taller plant architecture. (R) There was no difference in spike weight. However, (S) grain yield per five spikes was reduced in TRI 984 compared to CIRNO.

Fig. 2. Functional 'Stay-green' phenotype was observed in the landrace-elite recombinants. (A) Flag leaf and peduncle senescence were strongly linked, while they were independent of days to heading. (B) Flag leaf verdancy at the heading had only a minor impact on the progression of senescence. The RILs that exhibited delayed senescence had (C) more grains per spikelet and (D) higher thousand-grain weight, majorly due to (E) wider grains. Note: (B-E) are linear regression plots with the explanatory variable on the x-axis, while the y-axis represents the response variable. R^2 is the phenotypic variance explained, and the corresponding P-values of the regression analysis are displayed.

Fig. 3. bh^t-A1 induces spike-branching but with a spikelet fertility and grain weight trade-off. (A) A major effect QTL hotspot for spike-branching, grain length and weight was mapped on the short arm of Chr 2A. (B) RILs with the TRI 984 allele showed spike-branching, (C) no difference in grains per 5 spikes, but a reduction in (D) grain length, (E) thousand-grain weight, (F) flag leaf verdancy at heading and (G) spike length. Note: In (B-G), 'n' represents the number of RILs that were compared for each allele class, *viz.,* TRI 984 allele data points are in 'black', while 'cyan' colored data points represent CIRNO allele. 'Unpaired t test' was used to determine the statistical significance, and the resulting P-values (Two-tailed analysis) are displayed for all the comparisons.

Fig. 4. bh^t-A3, a new modifier locus for spike-branching. (A) bh^t-A3 mediates spike-branching, flag leaf senescence rate, grain filling duration, spikelet fertility and grain yield per spike. The CIRNO allele (B) increases the expressivity of spike-branching (when bh^t-A1 is

present), (C) delays flag leaf senescence rate, (D) increases grain filling duration, (E) grains per spikelet, (F) grain yield per five spikes and (G) thousand-grain weight. Note: In (B-G), 'n' represents the number of RILs that were compared for each allele class *viz.,* TRI 984 allele data points are in 'black', while 'cyan' colored data points represent CIRNO allele. 'Unpaired t test' was used to determine the statistical significance, and the resulting P-values (Two-tailed analysis) are displayed for all the comparisons.

Fig. 5. *gpc-B1* regulates senescence rate and grain weight. (A) QTLs for overall senescence rate and thousand-grain weight co-localised on Chr 6B. (B) The modern (CIRNO) allele mediated delay in flag leaf senescence, (C) peduncle senescence and (D) days to maturity (spike senescence). The resulting increase in the post-anthesis phase is translated into (E) an increase in grain width, (F) grain length and eventually (G) thousand-grain weight. Note: In (B-G), 'n' represents the number of RILs that were compared for each allele class, *viz.,* TRI 984 allele data points are in 'black', while 'cyan' colored data points represent CIRNO allele. 'Unpaired t test' was used to determine the statistical significance, and the resulting P-values (Two-tailed analysis) are displayed for all the comparisons.

Fig. 6. *bh^t-A1*, *bh^t-A3*, and *gpc-B1* balance the trade-off between spikelet fertility and grain weight spike-branching recombinants. The RILs with various combinations of alleles were phenotyped at IPK-Gatersleben (2021 & 2022) and the University of Hohenheim (2022). At IPK, the spike-branching RILs that senesce late *(bh^t-A1+bh^t-A3+gpc-B1)* have (A) higher grain number per 5 spikes; (B) increased thousand-grain weight at IPK, (C) more grain yield per five spikes and (D) Finally, grain yield per meter row was also higher in the stay-green spike-branching RILs (calculated only at IPK). Likewise, at Hohenheim, (E) we observed more grains per five spikes, (F) but no change in thousand-grain weight; eventually, (G) there was an increase in grain yield per spike. (H) pictorial depiction of the various allelic combinations that are analysed in (A-G). Note: In (A-G), one-way ANOVA followed by Dunnett's test was used to determine the statistical significance. All the comparisons are made with respect to '*bh^t-A1+BH^t-A3+GPC-B1*' allelic combination, and the corresponding P-values are displayed in all the graphs (significant ones are in bold). The image (H) is partly created using biorender (https://biorender.com/). *bh^t-A1+BH^t-A3+GPC-B1*: Recombinants with one locus for spike-branching and two loci for accelerated senescence; *bh^t-A1+BH^t-A3+gpc-B1*: Recombinants with one locus each for spike-branching and delayed senescence; *bh^t-A1+bh^t-A3+GPC-B1*: Recombinants with two loci for spike-branching, while carrying one locus for delayed

senescence; bh^t-A1+bh^t-A3+gpc-B1: Recombinants with two loci each for spike-branching and delayed senescence.

Fig. 7. Summary of the physiological and genetic basis of interaction among bh^t-A1, bh^t-A3, and gpc-B1 in the current landrace-elite recombinant population. Note: The image is partly created using biorender (https://biorender.com/).

Fig. 1. TRI 984 has a poor source-sink balance as opposed to CIRNO. (A) Spike-branching phenotype of TRI 984 and (B) Standard spike of CIRNO. (C) Spikelet from the central part of the TRI 984 main rachis shows reduced florets relative to (D) a spikelet from a similar position in CIRNO. (E) Despite having more spikelets, there is no increase in grain number per 5 spikes in TRI 984. (F-H) However, the grains are smaller in TRI 984, leading to a reduction in thousand-grain weight. (I) TRI 984 exhibited delayed heading, (J) lower flag leaf verdancy at heading, (K, L) accelerated flag leaf and peduncle senescence. TRI 984 had (M, N) shorter but wider flag leaves, (O) more tillers at booting (under greenhouse conditions), (P) longer spikes, (Q) taller plant architecture. (R) There was no difference in spike weight. However, (S) grain yield per five spikes was reduced in TRI 984 compared to CIRNO.

Fig. 2. Functional 'Stay-green' phenotype was observed in the landrace-elite recombinants. (A) Flag leaf and peduncle senescence were strongly linked, while they were independent of days to heading. (B) Flag leaf verdancy at the heading had only a minor impact on the progression of senescence. The RILs that exhibited delayed senescence had (C) more grains per spikelet and (D) higher thousand-grain weight, majorly due to (E) wider grains. Note: (B-E) are linear regression plots with the explanatory variable on the x-axis, while the y-axis represents the response variable. R^2 is the phenotypic variance explained, and the corresponding P-values of the regression analysis are displayed.

Fig. 3. *bh^t-A1* induces spike-branching but with a spikelet fertility and grain weight trade-off. (A) A major effect QTL hotspot for spike-branching, grain length and weight was mapped on the short arm of Chr 2A. (B) RILs with the TRI 984 allele showed spike-branching, (C) no difference in grains per 5 spikes, but a reduction in (D) grain length, (E) thousand-grain weight, (F) flag leaf verdancy at heading and (G) spike length. Note: In (B-G), 'n' represents the number of RILs that were compared for each allele class, *viz.,* TRI 984 allele data points are in 'black', while 'cyan' colored data points represent CIRNO allele. 'Unpaired t test' was used to determine the statistical significance, and the resulting P-values (Two-tailed analysis) are displayed for all the comparisons.

Fig. 4. bh^t-A3, a new modifier locus for spike-branching. (A) bh^t-A3 mediates spike-branching, flag leaf senescence rate, grain filling duration, spikelet fertility and grain yield per spike. The CIRNO allele (B) increases the expressivity of spike-branching (when bh^t-A1 is present), (C) delays flag leaf senescence rate, (D) increases grain filling duration, (E) grains per spikelet, (F) grain yield per five spikes and (G) thousand-grain weight. Note: In (B-G), 'n' represents the number of RILs that were compared for each allele class *viz.,* TRI 984 allele data points are in 'black', while 'cyan' colored data points represent CIRNO allele. 'Unpaired t test' was used to determine the statistical significance, and the resulting P-values (Two-tailed analysis) are displayed for all the comparisons.

Fig. 5. *gpc-B1* regulates senescence rate and grain weight. (A) QTLs for overall senescence rate and thousand-grain weight co-localised on Chr 6B. (B) The modern (CIRNO) allele mediated delay in flag leaf senescence, (C) peduncle senescence and (D) days to maturity (spike senescence). The resulting increase in the post-anthesis phase is translated into (E) an increase in grain width, (F) grain length and eventually (G) thousand-grain weight. Note: In (B-G), 'n' represents the number of RILs that were compared for each allele class, *viz.,* TRI 984 allele data points are in 'black', while 'cyan' colored data points represent CIRNO allele. 'Unpaired t test' was used to determine the statistical significance, and the resulting P-values (Two-tailed analysis) are displayed for all the comparisons.

Fig. 6. *bh^t-A1*, *bh^t-A3*, and *gpc-B1* balance the trade-off between spikelet fertility and grain weight spike-branching recombinants. The RILs with various combinations of alleles were phenotyped at IPK-Gatersleben (2021 & 2022) and the University of Hohenheim (2022). At IPK, the spike-branching RILs that senesce late *(bh^t-A1+bh^t-A3+gpc-B1)* have (A) higher grain number per 5 spikes; (B) increased thousand-grain weight at IPK, (C) more grain yield per five spikes and (D) Finally, grain yield per meter row was also higher in the stay-green spike-branching RILs (calculated only at IPK). Likewise, at Hohenheim, (E) we observed more grains per five spikes, (F) but no change in thousand-grain weight; eventually, (G) there was an increase in grain yield per spike. (H) pictorial depiction of the various allelic combinations that are analysed in (A-G). Note: In (A-G), one-way ANOVA followed by Dunnett's test was used to determine the statistical significance. All the comparisons are made with respect to '*bh^t-A1+BH^t-A3+GPC-B1*' allelic combination, and the corresponding P-values are displayed in all the graphs (significant ones are in bold). The image (H) is partly created using biorender (https://biorender.com/).

Fig. 7. Summary of the physiological and genetic basis of interaction among *bh^t-A1*, *bh^t-A3*, and *gpc-B1* in the current landrace-elite recombinant population. Note: The image is partly created using biorender (https://biorender.com/).