1 **The effects of inversion polymorphisms on patterns of neutral genetic diversity**
2
3

Affiliation:

Institute of Ecology and Evolution, School of Biological Sciences, University of

Edinburgh, Edinburgh EH9 3FL, United Kingdom

Key words: Inversion polymorphisms, genetic diversity, sequence divergence,

recombination rate, population subdivision

Running title: Inversion polymorphisms and diversity

Corresponding author:

Name: Brian Charlesworth

Address: Institute of Ecology and Evolution, School of Biological Sciences, Ashworth

Laboratories, University of Edinburgh, Charlotte Auerbach Road, Edinburgh EH9

3FL, UK

Telephone: +44 131 650 5751/662 1725

Email: Brian.Charlesworth@ed.ac.uk

21
22

Abstract

The strong reduction in the frequency of recombination in heterozygotes for an inversion and a standard gene arrangement causes inversion and standard haplotypes to become partially isolated, resulting in sequence divergence between them and to changes in the levels of neutral variability at nucleotide sites within each arrangement class, as is also the case for other types of balanced polymorphisms. Previous theoretical studies have either assumed that the population is panmictic or that it is divided into two populations subject to divergent selection. Here, the theory is extended to a model of an arbitrary number of demes connected by migration, using a finite island. Recursion relations for mean pairwise coalescent times are used to obtain simple approximate expressions for diversity and divergence statistics relevant to an inversion polymorphism at equilibrium under recombination and drift, and for the approach to equilibrium following the sweep of an inversion to a stable intermediate frequency. The effects of an inversion polymorphism on patterns of linkage disequilibrium are also examined. The reduction in effective recombination rate caused by population subdivision can have significant effects on these statistics, and hence on estimates of the ages of inversions. The theoretical results are discussed in relation to population genomic data on inversion polymorphisms, with an emphasis on *Drosophila melanogaster*. Methods are proposed for testing whether or not inversions are close to recombination-drift equilibrium, and for estimating the rate of recombinational exchange in heterozygotes for inversions; difficulties involved in estimating the ages of inversions are also discussed.

Introduction

Naturally occurring genetic factors that massively reduce the rate of crossing over in *Drosophila melanogaster* when heterozygous were discovered by A.H. Sturtevant over 100 years ago (Sturtevant 1917). These were shown to be inversions of segments of chromosomes by Sturtevant (1926). A major mechanism by which crossing over is suppressed or reduced in frequency during *Drosophila* female meiosis in heterozygotes for a paracentric inversion and the standard arrangement was elucidated by Sturtevant and Beadle (1936), whose genetic data suggested that single crossovers produce dicentric and acentric chromosomes that fail to be included in the egg nucleus. While several factors contribute to crossover suppression in inversion heterozygotes, with the conversion of double strand breaks into non-crossover associated gene conversion events playing an important role (Gong et al. 2005; Crown et al. 2018; Li et al. 2022), there is no doubt that crossover suppression in inversion heterozygotes must cause different arrangements

to become substantially genetically isolated from each other. Recent studies of heterozygous

Drosophila inversions suggest a total suppression of crossing over within the regions covered by the

inversion and for a substantial distance outside it (Li et al. 2022; Koury 2023).

Inversion polymorphisms were long regarded as a curiosity of *Drosophila* and other

"higher" Diptera, where they can readily be detected cytologically using the polytene chromosomes

of the larval salivary glands. The largest body of information concerning the transmission and

population genetics of inversions has been accumulated in studies of numerous *Drosophila* species

(Krimbas and Powell 1992a; Kapun and Flatt 2019). However, the recent application of genome

sequencing technology to the natural populations of many organisms, including humans, has

revealed that inversions are much more abundant than was previously thought. They are sometimes

associated with striking phenotypic polymorphisms, such as the social chromosomes of ants and the

behavioral and color polymorphisms of the ruff and white-throated sparrow (Wellenreuther and

Bernatchez 2018; Villoutreix et al. 2021). This has led a surge in interest in the evolutionary

significance of inversion polymorphisms, although it should be borne in mind that the *Drosophila*

studies suggest that the vast majority of polymorphic inversions have little or no effects on visible

phenotypes, although effects on quantitative traits such as body size and some fitness components

have been detected (Krimbas and Powell 1992a; Kapun and Flatt 2019).

It is evident that the genetic isolation of different gene arrangements must play a major role

in the evolutionary processes that lead to the evolution of inversion polymorphisms, although the

nature of these processes is still an open research question, which may well have more than one

answer (Krimbas and Powell 1992b; Wellenreuther and Bernatchez 2018; Kapun and Flatt 2019;

Villoutreix et al. 2021). It is also clear that recombination suppression between arrangements is

often incomplete, with gene conversion and/or double crossing over causing exchanges of alleles

between arrangements in heterokaryotypes (Sturtevant and Beadle 1936; Chovnick 1973; Krimbas

and Powell 1992b; Navarro et al. 1997; Crown et al. 2018; Korunes and Noor 2019; Li et al. 2022).

Despite this potential for recombinational exchange between chromosome arrangements,

early studies using electrophoretic variants detected strong associations (linkage disequilibrium:

LD) between gene arrangements and molecular markers in *Drosophila pseudoobscura* (Prakash and

Lewontin 1968, 1971). These associations were initially interpreted as evidence for Dobzhansky's

"coadaptation" hypothesis for the maintenance of inversion polymorphisms by epistatic selection

(Dobzhansky 1947, 1950), with inversion heterozygosity suppressing recombination between

polymorphic loci with epistatic effects on fitness (Prakash and Lewontin 1968, 1971). It was soon

realized, however, that other processes, such as the genetic hitchhiking of alleles that happened to

be present in the chromosome in which an inversion originated, provided plausible alternative

explanations for strong LD between molecular markers and inversions (Mukai et al. 1974; Sheppard

1975, p.150; Nei and Li 1975). A model of hitchhiking that allowed for recombinational exchange

between inversion and standard arrangements in the presence of selection against recombinants was

developed by Ishii and Charlesworth (1977), who concluded that hitchhiking could probably

explain most, but not all, of the data available at the time. It is, of course, *a priori* unlikely that a

randomly chosen molecular marker will itself be involved in any selective process responsible for

maintaining an inversion polymorphism.

Several lines of evidence, including studies of changes in inversion frequencies in

experimental populations, temporal fluctuations in inversion frequencies and clinal patterns of

inversion frequencies, as well as direct measurements of fitness components, show that many

inversions are maintained at intermediate frequencies by natural selection (Krimbas and Powell

1992b; Mérot et al. 2018; Kapun and Flatt 2019). Once an inversion subject to balancing selection

has established itself at an intermediate frequency within a population, there will be a gradual

approach to mutation-drift-recombination equilibrium with respect to neutral or nearly neutral

variants, as recombinational exchange tends to homogenize the inverted and standard arrangements,

and new mutations arise and are either lost or spread to fixation by genetic drift (Navarro et al.

2000). The equilibrium properties of an inversion polymorphism with respect to neutral variability

are similar to those for neutral loci linked to a single locus subject to balancing selection, first

modeled by Ohta and Kimura (1970) using diffusion equations and by Strobeck (1983) using

identity probabilities. Over the following two decades, subsequent modeling work using the

structured coalescent process shed further light on patterns of neutral variability at sites linked to

loci under balancing selection or divergent local selection, e.g., Kaplan et al. (1988), Hudson and

Kaplan (1988), Hudson (1990), Takahata (1990), Charlesworth et al. (1997), Nordborg (1997) and

Navarro et al. (2000), reviewed by Charlesworth et al. (2003).

These studies showed that, when the balanced polymorphism has been maintained for a time

that is much longer than the mean coalescent time for a neutral locus, we expect to see LD between

the target of selection and neutral sites closely linked to the target of selection, LD among these

neutral sites, sequence divergence between haplotypes carrying the different alleles at the target of

selection, and enhanced variability in the population as a whole around the target of selection.

These patterns are all reflections of the same process of divergence by drift and mutation among the

partially isolated populations represented by the alleles at the target of selection, with a strong

analogy with the outcome of mutation and drift in a geographically structured population (Hudson

1990; Charlesworth et al. 2003). Inversions are only special because of the strong suppression of

recombination within the inversion, and in regions outside it but close to the inversion breakpoints

(Navarro et al. 1997; Crown et al. 2018; Korunes and Noor 2019; Li et al. 2022; Koury 2023), so

that these patterns are much more likely to be observed with inversions than with single locus

polymorphisms, due to the much larger region of the genome covered by inversions.

A basic finding of the early theoretical work was that significant associations between a

balanced polymorphism and variants at a neutral locus are only likely to be observed if the rate of

recombinational exchange in double heterozygotes for the loci involved is of the order of the

reciprocal of the effective population size, N_e, as is also the case for a pair of neutral loci (Hill and

Robertson 1968; Ohta and Kimura 1969, 1971). Andolfatto et al. (2001) applied the relevant

theoretical results on balanced polymorphisms to the available data on *D. melanogaster* inversions,

and concluded that large effects of inversions on patterns of variability are likely to be seen only for

loci close to inversion breakpoints, where exchange is likely to be most strongly inhibited, and that

the observed patterns suggested that the inversions in question were of relatively recent origin

compared with the age of the species.

Interest in the question of the effect of inversion polymorphisms on patterns of variability at

sites that are either inside inversions or closely linked to inversion breakpoints has increased with

the advent of whole genome sequencing, which has both greatly increased our ability to detect and

characterize inversions, and allowed much more fine-scaled analyses of patterns of variability in

genomic regions associated with inversions, e.g., Corbett-Detig and Hartl (2012), Cheng et al.

(2012), Mérot et al. (2021) and Kapun et al. (2023). Much more information of this kind is likely to

become available, and its interpretation will require a solid basis in population genetics theory. A

number of theoretical investigations on the effect of balanced polymorphisms on variability at

linked sites have extended the older work described above, without, however, greatly modifying the

basic conclusions, e.g., Innan and Nordborg (2003), Kirkpatrick and Guerrero (2014), Rousset et al.

(2014) and Zeng et al. (2021). A limitation of most of the theoretical work on the effects of

selectively maintained polymorphisms on neutral diversity is that it assumes a single, randomly

mating population, with the exception of Charlesworth et al. (1997), Nordborg (1997) and Guerrero

et al. (2012), who considered the case of divergent and/or balancing selection in a pair of

populations. Nordborg and Innan (2003) examined a more general model of population structure but

relied on coalescent process simulations to generate predictions.

While it is often considered that population subdivision in organisms like *Drosophila* is

likely to have only minor effects on genetic diversity, given the generally low levels of F_{ST} among

populations (Singh and Rhomberg 1987; Schaeffer 2002; Lack et al. 2015, 2016), there is evidence

from studies of allelism of recessive lethals that local deme sizes are in reality somewhat restricted

in size, with limited migration among them (Wright et al. 1942; Mukai and Yamaguchi 1974; Ives

and Band 1986); this conclusion has recently been confirmed by a resequencing study of a single

US population over time (Lange et al. 2022). It is known that population subdivision with a large

number of demes increases the amount of linkage disequilibrium among neutral loci when genomes

are sampled from the same deme (Wakeley and Lessard 2003), because population subdivision

increases local homozygosity, thereby reducing the effectiveness of recombination. It is therefore of

interest to examine the consequences of such subdivision for the effect of a biallelic balanced

polymorphism on variability at a linked neutral site, and this is the main topic of the present paper.

For brevity, the locus under selection is referred here to exhibiting an inversion

polymorphism, but the results apply to any Mendelian locus with the same properties. An island

model of a metapopulation of large size, divided into a finite number of demes of equal size, is

assumed, with the same migration rate between all pairs of demes. As shown by Wakeley and

Aliacar (2001), the properties of such a model are likely to provide a good approximation to more

realistic scenarios, such as a two-dimensional stepping stone model, provided that the number of

demes is large. In order to obtain simple results, it is assumed that the inversion is the derived state,

and that selection on the inversion is sufficiently strong that it has risen quickly to an equilibrium

frequency that is constant across demes. The properties of variability at the neutral locus, LD

between the neutral locus and the inversion, and the extent of divergence between karyotypes at

drift-mutation-recombination equilibrium, are studied first, followed by an examination of the

approach to equilibrium. Here, "karyotype" is used to denote the state of a haplotype with respect to

the arrangement which it carries, and "heterokaryotype" is used to denote a heterozygote for

inverted and standard arrangements. Approximations are used in order to generate simple analytical

results.

The model and its analysis
184

Assume that an autosomal inversion (In) is maintained at a frequency x and the standard

arrangement (St) has frequency $y = 1 - x$. Without loss of generality, we can assume $x \leq \frac{1}{2}$ when

considering equilibrium results; if this is not the case, then In and St can simply be interchanged.

Parameters for In and St are denoted by subscripts 1 and 2, respectively. The population is assumed

to be divided into local populations (demes) that are at equilibrium under mutation, genetic drift and

migration, and reproducing according to a Wright-Fisher model, so that the effective population

size of a deme is equal to its adult population size. An island model with a large number of demes,

$d,$ each with population size N is assumed, so that the migration effective population size (Nagylaki

1998) is $N_T = Nd$. Migration between populations occurs at rate m per generation.

The level of equilibrium neutral differentiation at autosomal loci that are independent of the

inversion is measured by F_{ST}, which is defined here as one minus the ratio of the mean coalescent

time for pairs of alleles sampled from within a population to the mean coalescent time for pairs of

alleles sampled randomly from the population as whole. Several other definitions of F_{ST} have been

used in the literature (Charlesworth 1998), and used in different estimation procedures

(Gammerdinger et al. 2020), but this one has the closest relationship to Wright's original definition

(Wright 1951), and is applicable to DNA sequence variability via the estimator γ_{ST} (Nei 1982) or its

equivalent the K_{ST} statistic of Hudson, Boos and Kaplan (1992). For large d, $F_{ST} \approx 1/(1 + 4Nm)$

(Charlesworth 1998). It should, however, be borne in mind that different authors use different

estimators; the widely used methods of Weir and Cockerham (1984) and Hudson, Slatkin and

Maddison (1992), which are mathematically equivalent, have much larger expected values than K_{ST}

when the number of populations being compared is small (Charlesworth 1998; Gammerdinger et al.

2020), so that caution needs to be used when comparing F_{ST} estimates from different studies. This is

especially important for comparisons between inverted and standard arrangements or between X

and Y chromosomes, where only two populations are involved (Gammerdinger et al. 2020).

Random mating within local populations and with respect to arrangement status is assumed.

The migration and recombination parameters, m and r, are assumed to be so small that second-order

terms can be neglected. At a neutral site within the region covered by the arrangement,

recombinational exchange between arrangements caused by gene conversion and/or double crossing

over occurs at rate r per generation. In the gametes produced by a heterokaryotype, In/St, there is a

probability r that a given neutral site associated with the In haplotype came from the St haplotype,

and that the homologous site associated with the St haplotype came from the In haplotype.

The dynamics of coalescent events are determined jointly by migration, gene conversion and

genetic drift. To determine pairwise coalescent times, we need to consider cases where two In

haplotypes are sampled (a sample of class 1/1), two St haplotypes are sampled (a class 2/2 sample),

and one of each type are sampled (a class 1/2 sample). The method of Nagylaki (1998) is used for

this purpose, which provides a simple way of determining expected coalescent times (see also

Charlesworth and Charlesworth 2010, p.316). For 1/1 samples, the probability that the focal neutral

site in one of the two haplotypes was associated with St in the previous generation is $2ry$ (ignoring

second-order terms in ry), since the chance that a given In haplotype was combined with an St

haplotype is y, and either of the two *In* haplotypes could have experienced a recombination event.

There is a probability $2m$ that one of the two haplotypes was present in a different deme from the

current one, and a probability of $1/(2Nx)$ that the two haplotypes coalesced in the previous

generation if they were both in the same deme; haplotypes from different demes cannot coalesce.

Similar considerations apply to 2/2 samples, except that x and y are interchanged. For 1/2 samples,

there is a probability y that the *In* haplotype was associated with an *St* haplotype in the previous

generation, so the probability that the focal neutral site in *In* derives from *St* is ry. Similarly, there is

a probability rx that the focal neutral site in the *St* haplotype derives from *In*; the net probability of

the focal site coming from a different karyotype from its current one is r. No coalescence of the two

haplotypes is possible in this case, but there is a probability $2m$ that one of them has moved from a

different deme.

We need to distinguish between a pair of haplotypes that are sampled from the same deme

(denoted by subscript w), and a pair of haplotypes sampled from two different demes (denoted by

subscript b). The expected coalescent time for a sample of class i/j for a within-deme sample is

denoted by t_{ijw} and the equivalent for a between-deme sample is t_{ijb}. For a between-deme sample, the

probability that a migrant haplotype came from the same deme as the non-migrant haplotype is $1/(d$

$- 1)$ and the probability that it came from a different deme is $(d - 2)/(d - 1)$. Following Nagylaki

(1998), we use the fact that, if no coalescent event occurs for a pair of haplotypes of a given class,

the relevant expected coalescent time is increased by one generation. Applying this principle to the

above definitions, simple recursions for the t's can be obtained, which are given by Equations (A1)

of the Appendix. Further simplification is provided by neglecting the products of $1/Nx$ and $1/Ny$

with m and r (Equations A2).

It is convenient to scale the migration rate and recombination rate by four times the deme

size, writing $M = 4Nm$ and $R = 4Nr$. In addition, coalescent times can be expressed relative to the

expected coalescent time for a pair of alleles sampled from the same deme at a locus that is

independent of the inversion, $T_{Sn} = 2N_T = 2dN$. Upper case T's are used to denote t's divided by T_{Sn}.

As shown in the Appendix, manipulation of the resulting equations leads to simple explicit

approximate expressions for the equilibrium values of the T_{ijb} and T_{ijw}, assuming a large number of

demes and $M \gg R$ (Equations A5 and A6).

It is useful to consider the mean scaled coalescent time for pairs of alleles sampled randomly

across karyotypes. For alleles from different demes, this is given by:

$$T_{Tb} = 2xyT_{12b} + x^2T_{11b} + y^2T_{22b} \qquad (1a)$$

258   Similarly, the mean scaled mean coalescent time for a pair of alleles sampled within

259 karyotypes, but from different demes, is:

261
$$T_{Sb} = xT_{11b} + yT_{22b} \qquad\qquad (1b)$$

263   A between-karyotype analogue of F_{ST} was defined by Charlesworth et al. (1997), analogous

264 to K_{ST} of Hudson, Boos and Kaplan (1992), is:

266
$$F_{ATb} = 1 - \frac{T_{Sb}}{T_{Tb}} \qquad\qquad (2a)$$

268 A quantity analogous to the $\langle F_{ST} \rangle$ statistic of Hudson, Slatkin and Maddison (1992) is:

270
$$\tilde{F}_{ATb} = 1 - \frac{T_{Sb}}{T_{12b}} \qquad\qquad (2b)$$

272 For equilibrium under recombination and drift, for which $T_{12} > 1$, $\tilde{F}_{ATb} > F_{ATb}$, since $T_{12b} > T_{Tb}$.

273   Corresponding expressions apply to alleles sampled within demes, replacing subscript b by

274 w. On general grounds, the Maruyama invariance principle that applies to structured coalescent

275 processes with conservative migration (Maruyama 1974; Nagylaki 1982) implies that at equilibrium

276 we have:

277
$$T_{Sw} = xT_{11w} + yT_{22w} = 1 \qquad\qquad (3)$$

If the terms in d^{-1} in Equations (A6f) and (A6g) are neglected, evaluation of the resulting expression

for T_{Sw} confirms that it is indeed equal to 1, to the accuracy of the approximations used to obtain

these equations. This result breaks down when $N_T r$ is close to zero; with $r = 0$, the two karyotypes

behave as separate populations, with migration effective population sizes of $N_T x$ and $N_T y$,

respectively, so that $T_{Sw} = x^2 + y^2$.

This result implies that the equilibrium expressions for F_{ATw} and $\tilde{F}_{ATw}$ simplify to:

286
$$F_{ATw} = 1 - \frac{1}{T_{Tw}} \qquad\qquad (4a)$$

287
$$\tilde{F}_{ATw} = 1 - \frac{1}{T_{12w}} \qquad\qquad (4b)$$

For equilibrium populations, therefore, the two F-statistics for within-population/between-

karyotype differentiation contain no more information than do T_{Tw} and T_{12w}, respectively; $T_{Tw} = 1/(1$

$- F_{ATw})$ and $T_{12w} = 1/(1 - \tilde{F}_{ATw})$. The latter two statistics have the advantage that they are not

constrained to be below one. For applications of these formulae to population genomic data, the

divergence statistics corresponding to the T's would need to be scaled by an estimate of mean

neutral nucleotide site diversity at sites independent of the inversion, or by the weighted mean of

the two within-karyotype diversities. The latter has the advantage that potential differences in

mutation rates among different genomic regions are factored out.

When the scaled migration rate M tends to infinity and F_{ST} tends to zero, the case of a

panmictic population with population size N_T is approached, and the subscripts w and b can be

dropped. The recursion relations for the within-deme statistics can be used for this case (Equations

A2), setting m to zero and the deme size to N_T. Application of the above method to this case, writing

$\rho = 4N_T r$ for the scaled recombination rate, and assuming that $\rho > 0$, gives the following expressions

for equilibrium:

303
$$T_{12} = 1 + 2\rho^{-1} \qquad (5a)$$

305
$$T_{11} = [x(1 + 2y) + \rho xy]/(1 + \rho xy) \qquad (5b)$$

307
$$T_{22} = [y(1 + 2x) + \rho xy]/(1 + \rho xy) \qquad (5c)$$

309   These are equivalent to Equations (A3) and (A4) of Nordborg (1997), Equations (6) of Navarro et

310   al. (2000), and Equation (8) of Zeng et al. (2021), which were derived by more complex methods. It

311   is easily seen that $T_S = xT_{11} + yT_{22} = 1$ at equilibrium, so that we have:

313
$$T_T = 1 + 4xy\rho^{-1} + xy(x - y)^2(1 + \rho xy)^{-1} \qquad (5d)$$

315
$$F_{AT} = 1 - \frac{1}{1 + 4xy\rho^{-1} + xy(x-y)^2(1 + \rho xy)^{-1}} \qquad (5e)$$

317
$$\tilde{F}_{AT} = \frac{2}{2 + \rho} \qquad (5f)$$

Theoretical results: equilibrium populations

General considerations

Navarro et al. (2000) have given a detailed description of results for the case of an equilibrium

panmictic population, which is the limiting case when $F_{ST} = 0$. The focus here is therefore on

subdivided populations. Numerical results from the approximations for the equilibrium T's given by

Equations (A5) and (A6), which assume large d and $m >> r$, are easily generated. When comparing

the theoretical results with observations, it is useful to note that the T's under the infinite sites

model of Kimura (1971) are proportional to the corresponding mean pairwise diversities or

divergences per basepair, taken over large numbers of neutral sites with the same mutation and

recombination rates (e.g., Hudson 1990). The mean levels of nucleotide site diversity and

divergence between arrangements within D. $melanogaster$ populations are such that the infinite

sites model fits the data reasonably well (Langley et al. 2012; Corbett-Detig and Hartl 2012; Kapun

and Flatt 2023), so that the T values and their ratios presented in the figures below can be taken as

indicating the corresponding diversity and divergence values when attempting to interpret data on

inversion polymorphisms in this species. Most multicellular organisms have lower diversity values

than this (Buffalo 2021), and so should present even less of a problem of interpretation.

Numerical results for subdivided populations

It is important to note that, under the assumptions made here, the equations for the T's involve only

the parameters $M = 4Nm$ and $\rho = 4N_T r = 4Ndr$; in turn, with large d we have $M \approx (1 - F_{ST})/F_{ST}$.

where F_{ST} is the measure of differentiation among local populations for neutral loci independent of

the inversion. Under these conditions, if F_{ST}, r and N_T are specified, the results are not affected by

the number of demes, the deme size (N) or the migration rate (m). Fig. 1 shows the results for the

case when the inversion is present at a frequency of 0.1 in all demes, and there are 200 demes each

with $N = 5000$, giving a total effective population size of $N_T = 10^6$. Fig. 2 shows similar results for

an inversion with a frequency of 0.5; in this case T_{11w}/ T_{22w} and T_{11b}/ T_{22b} are both equal to one, so

that these ratios are not shown.

These results are intended to give a somewhat crude representation of a set of populations in

a single geographical region that is isolated from other regions, with a within-deme neutral diversity

value like that found in population genomic surveys. The results from the $Drosophila$ Genome

Nexus Project, which has assembled genome sequence data for a large number of D. $melanogaster$

genomes from natural populations (Lack et al. 2015, 2016), show that $<F_{ST}>$ between pairs of

populations within a region is generally low, around 0.05 or even less, and is only about 0.2 even

between continents (Fig. 3 of Lack et al. 2016). F_{ST} values in animals (Roux et al. 2016) and

outcrossing flowering plants (Charlesworth 2003) rarely exceed 0.25, so that F_{ST} in the figures was

restricted to the range 0 – 0.25 for purposes of illustration, consistent with the approximations used

to generate the results. In what follows, special attention is paid to the results with $F_{ST} = 0.05$,

although the mean pairwise $<F_{ST}>$ for inversion-free genomes in Figure 3 of Lack et al. (2016)

between the Zambian *D. melanogaster* population (ZI) and two populations from South Africa (SP

and SI) is only 0.007. This suggests that panmixia is a good approximation for populations in this

region, which is thought be close to the center of origin of the species (Sprengelmeyer et al. 2020).

The ZI population is a focus of the intensive study of the *In(3R)P* inversion by Kapun et al. (2023),

which is discussed below in relation to the theoretical results.

Results are shown for four different values of the rate of recombination (*r*) between a neutral

site and the arrangement in *In/St* heterokaryotypes, which fall within the range reported for single

inversions in *Drosophila* (Navarro et al. 1997; Korunes and Noor 2019). The right-hand Y axes

correspond to the absolute values of the T_{ij}'s. The mean coalescent time between *In* and *St* relative

to $2N_T$, T_{12}, is almost the same for within-deme pairs of alleles as for alleles from separate demes

(see Equations A6d and A6e), so that only the between-deme values are shown. T_{12} provides a

measure of the expected sequence between a pair of alleles from the two different karyotypes,

relative to the within-deme neutral diversity for loci independent of the inversion, or to the mean

neutral diversity over *In* and *St* karyotypes sampled randomly from the same deme. Both of these

diversity measures are equal to $\theta = 4N_T u$, where *u* is the mutation rate per basepair (Nagylaki

1998)– see Equation (3). For $u = 5 \times 10^{-9}$ (Assaf et al. 2017) and $N_T = 10^6$, a value of $T_{12} = 1$

corresponds to a sequence divergence of $T_{12}\theta/2 = 0.01$ per basepair, which is approximately equal

to the within-population synonymous site diversity in *D. melanogaster* ancestral range populations

(Langley et al. 2012). As can be seen from Figs. 1 and 2, T_{12} is quite close to 1 for $r = 10^{-5}$ or 10^{-4}

with $F_{ST} \leq 0.05$, but takes much larger values for the two lower recombination rates. T_{12} increases

with F_{ST}, but only slowly at the two lower recombination rates; T_{12} is at most close to 1.7 for the two

higher *r* values.

Unless the recombination rate is very low, as may be the case near inversion breakpoints

(Navarro et al. 1997), the equilibrium level of sequence divergence between *In* and *St* is thus likely

to be less than 70% larger than the genome-wide mean neutral diversity, corresponding to an $\tilde{F}_{AT}$ of

0.41. Equation (A6d) shows that the effect of F_{ST} on T_{12} is caused by the increase in divergence

between alleles sampled within karyotypes and from different populations as the between-deme

coalescent times increase. The effect of recombination on T_{12} is controlled by the reciprocal of ρ, the

recombination rate scaled by $4N_T$, which is invariant with respect to the level of population

subdivision under the assumptions used here.

A related statistic is the net mean coalescent time for a pair of alleles sampled randomly

within demes without regard to karyotype (T_{Tw}), which determines the overall mean diversity at a

nucleotide site located within the region covered by the inversion. This is less sensitive to F_{ST} than

T_{12}, but increases somewhat with increasing F_{ST}, especially at high recombination rates. As would

be expected from the fact that it includes a contribution from within-karyotype coalescent events, it

is always smaller than T_{12}, even for an intermediate frequency inversion, so that the corresponding

diversity statistic provides a less useful index of between-karyotype differentiation than the

between-karyotype divergence. The corresponding statistic for alleles sampled from different demes

(T_{Tb}) has similar properties, but is always somewhat larger in magnitude, especially at high F_{ST}

values (see Table S1). Unless F_{ST} is rather high or r is very low, an inversion polymorphism at

recombination-drift equilibrium is not likely to have a large effect on the overall level of sequence

diversity in the population, consistent with the results for chromosome arm 3R of *D. melanogaster*

shown in Fig. 4 of Corbett-Detig and Hartl (2012).

The between-karyotype analogues of F_{ST}, F_{AT} and $\tilde{F}_{AT}$, (see Equations 4), are often used as a

measure of the extent of genetic differentiation among karyotypes, rather than the absolute

divergence measures just discussed. As noted by Charlesworth et al. (1997) in the context of a

single-locus balanced polymorphism (see also Zeng et al. [2021]), F_{AT} is equivalent to the

σ_d^2 measure of linkage disequilibrium of Ohta and Kimura (1971), treating the neutral site and

karyotype as a pair of loci, but it is considerably less tedious to estimate from population genomic

data. Its properties thus differ somewhat from those of T_{12} or T_{Tw}, especially as it is heavily

influenced by the within-karyotype coalescent time in the numerator of Equation (4a) (Charlesworth

1998). Figures 1 and 2 show plots of F_{AT} against F_{ST}, both for alleles from the same deme (F_{ATw}) and

for alleles from different demes (F_{ATb}); their values are displayed on the left-hand Y axes of each

panel.

F_{ATw} and F_{ATb} are again only weakly affected by the extent of population subdivision at the

lower two recombination rates, with F_{ATw} increasing slightly with F_{ST} whereas F_{ATb} decreases; F_{ATw} is

always larger than F_{ATb}, especially when F_{ST} and r are large, and is thus more useful as a measure of

between-karyotype differentiation. This difference reflects the fact that the inbreeding associated

with population subdivision reduces the effective rate of recombination among haplotypes from the

same deme (Wakeley and Lessard 2003). Both statistics are highly sensitive to r, with small values

at the two higher r values, especially for F_{ATb}. $\tilde{F}_{ATw}$ for equilibrium populations is given by $1 - 1/T_{12}$

(Equations 4b and 5f), and so behaves in a complementary fashion to T_{12} as a function of r and F_{ST}.

A moderately high level of population subdivision in an equilibrium system thus makes it

much easier to detect between-karyotype differentiation at low recombination rates, as measured by

either F_{ATw} or T_{12}, but makes F_{ATb} a less useful measure. Comparisons of the mean between-

karyotype divergence (either within or between demes) with the mean within-deme, within-

karyotype diversity at loci independent of the inversion (corresponding to a T_w of 1), or with the

mean within-deme, within-karyotype diversity (corresponding to a T_{Sn} of 1), are probably the most

useful measures of the extent of between-karyotype divergence, if equilibrium can be assumed.

It is of some interest to compare the values of F_{AT} for a polymorphism maintained at

constant frequencies with the value of σ_d^2 for a pair of neutral loci at statistical equilibrium under

recombination and drift in a subdivided population. Table 1 shows results for the within-deme

measure of F_{AT} compared with σ_d^2 for within-deme samples calculated from the equations in the

Appendix to Wakeley and Lessard (2003), for a range of F_{ST} values. For the intermediate and

highest recombination rates ($\rho = 4$ and $\rho = 40$), F_{AT} for $x = 0.5$ is very close to σ_d^2, whereas F_{AT} for x

$= 0.1$ is substantially smaller, except for the panmictic case ($F_{ST} = 0$); all three variables increase

considerably with increasing F_{ST}, reflecting the reduced effective recombination rate when there is

extensive population subdivision (Wakeley and Lessard 2003). In contrast, for the lowest

recombination rate ($\rho = 0.4$), F_{AT} for $x = 0.1$ is closer to σ_d^2, and there is only a small increase in the

three measures with F_{ST}. For a pair of neutral loci, σ_d^2 for a sample of haplotypes taken from

separate demes (which is equivalent to random sampling from the whole population with large d) is

independent of the extent of subdivision with the model used here (Wakeley and Lessard 2003), in

contrast to the behavior of F_{ATb} shown in Figs 1 and 2. The results for neutral loci thus shed only

limited light on what is to be expected for an inversion polymorphism maintained at a constant

frequency. The discrepancy between the statistics for the neutral case and the case with a constant

inversion frequency could in principle be used as a test for selection, although this would require

accurate knowledge of r.

When the inversion is present at a low frequency, as in Figure 1, the mean coalescent times

for pairs of alleles sampled within the inversion (T_{11}) are smaller than those for alleles sampled

within St (T_{22}), due to the lower effective population size of carriers of In; this of course also applies

to the corresponding diversity values (Nordborg 1997; Navarro et al. 2000).The converse is true if

In is more frequent than St. The ratios T_{11w}/T_{22w} and T_{11b}/T_{22b} are the black and red dashed curves,

respectively, in Figure 1, whose values are shown on the left-hand Y axes of each panel. T_{11w}/T_{22w}

decreases considerably as F_{ST} increases when r is relatively large, whereas T_{11b}/T_{22b} increases

slightly. Conversely, T_{11b}/T_{22b} increases considerably with F_{ST} when r is small, but T_{11w}/T_{22w} hardly

changes. T_{11w}/T_{22w} is, however, always greater than the value of 0.11 expected from the relative

frequencies of *In* and *St*, reflecting the effect of recombination; consistent with this, both T_{11w}/T_{22w}

and T_{11b}/T_{22b} increase with r.

All other things being equal, therefore, the departure of estimates of T_{11w}/T_{22w} from the ratio

of the frequencies of *In* and *St* provides an inverse measure of the extent of differentiation between

karyotypes, when the frequency ratio departs considerably from unity. The behavior of T_{11w}/T_{22w} as a

function of r reflects the fact that, for a low frequency inversion, T_{11w} increases considerably with

increasing r, whereas T_{22w} decreases slightly (see Table S1). For example, with $x = 0.1$ and $F_{ST} =$

0.05, $T_{11w} = 0.303, 0.430$ and 0.647 for $r = 10^{-7}, 10^{-6}$ and 10^{-5}, respectively. The corresponding

values of T_{22w} are 1.078, 1.063, and 1.040.

As expected from Equation (5a), a comparison of Figs. 1 and 2 shows that T_{12} is not greatly

affected by the inversion frequency; in contrast, F_{ATw} and F_{ATb} are larger with equal frequencies of

the two karyotypes than when the inversion is rare or very common. This reflects the smaller T_T

values with very low or high inversion frequencies, as can be seen from Equation (5d) for the

panmictic case, where T_T is approximately $1 + 2\rho^{-1}$ for $x = 0.5$, but approaches $1 + x(4\rho^{-1}+1)$ as x

tends to 0. This is another reason for using between-karyotype divergence relative to mean within-

deme diversity as a measure of differentiation between karyotypes.

Theoretical results: Approach to equilibrium

General considerations

The above results assume both that the inversion has reached its equilibrium frequency and that

there has been sufficient time for the effects of coalescence and recombination to equilibrate. In

reality, as was assumed in early hitchhiking models of associations between electrophoretic variants

and inversions (Ishii and Charlesworth 1977), a polymorphic inversion is likely to have arisen on a

single unique haplotype drawn at random from the initial population. Having survived early

stochastic loss, it will have approached its equilibrium frequency over a time that is inversely

related to the strength of selection acting on it. Molecular characterizations of inversion breakpoints

in *D. melanogaster* and *D. subobscura* strongly support the unique origin hypothesis (Corbett-Detig

and Hartl 2012; Orengo et al. 2019; Kapun et al. 2023). After this selective equilibrium has been

approached, there will be another extended period during which mutation-recombination-drift

equilibrium is approached. In the case of a panmictic population, both of these episodes can be

included in the same model, using phase-type theory (Zeng et al. 2021).

But the population genetics of the spread of a new mutation in a subdivided population, and

its hitchhiking effects on linked neutral sites, is much more complex (Barton et al. 2013), and has

not been applied to the case of balancing selection. In the present treatment of subdivided

populations, the initial approach to the equilibrium inversion frequency is assumed to occur

effectively instantaneously, so that only the second phase of the approach to equilibrium is studied.

Clearly, this is likely to cause the time taken to approach equilibrium and the effects of

recombination during this period to be underestimated, due to the additional time needed for a

mutation to spread through a subdivided population compared with panmixia, even if the habitat is

two-dimensional rather than one-dimensional (Barton et al. 2013).

By using this simplifying assumption, we can set $T_{22w} = T_{22b} = 0$, $T_{12w} = T_{11w} = 1$, and $T_{12b} =$

T_{11b}, with $T_{11b} \approx 1/(1 - F_{ST})$ when d is large. Using these as initial values, the change per generation

in the deviations of the T's from their equilibrium values can be calculated by means of the matrix

iteration $x_n = A\ x_{n-1}$, where x_n is the column vector of deviations of the T's in generation n from their

equilibrium values (see section S1 of the Supplementary Appendix for details). This approach

breaks down in the absence of recombination, since in this case the divergence between In and St

increases without limit as time increases; a separate treatment of this case is given in sections S6

and S7 of the Supplementary Appendix.

In order to speed up calculations, a relatively small total population size (10^5 for the case of

a subdivided population, and 10^4 for the equivalent panmictic population) was assumed, with

rescaling of parameters to keep their products with N_T identical with those used for the equilibrium

results. Insight into the asymptotic rate of approach to equilibrium when d is large can be obtained

from the eigenvalues of the A matrix. As shown in section S3 of the Supplementary Appendix, if

second-order terms in $1/N$, r and m are neglected, the structure of A is such that its 6 eigenvalues (3

in the panmictic case) are each approximately equal to one of its diagonal elements, i.e., to $1 - 2m -$

$2ry - 1/(2Nx)$, $1 - 2m/d - 2ry$, $1 - 2m - 2rx - 1/(2Ny)$, $1 - 2m/d - 2rx$, $1 - 2m - r$, and $1 - 2m/d - r$,

respectively. The asymptotic rate of approach to equilibrium is controlled by the largest of these

quantities; which of the six is the largest is determined by the relative values of Nx, Ny, rx, ry and

m. Since $m \gg r$ and $d \gg 1$ with the approximations used here, either $1 - 2m/d - 2ry$ or $1 - 2m/d -$

$2rx$ will be the largest eigenvalue, showing that the asymptotic rate of approach to equilibrium is

largely controlled by the product of $2r$ and the smaller of the two frequencies x and y when d is very

large.

In the case of a panmictic population, the system reduces to a three-dimensional one with

eigenvalues approximately equal to $1 - 2ry - 1/(2Nx)$, $1 - 2rx - 1/(2Ny)$, and $1 - r$, so that a similar

conclusion applies. The timescale of the approach of the coalescence times to equilibrium after the

inversion has approached its equilibrium frequency is thus of the order of the inverse of the

recombination frequency (unless this is very close to zero – see sections S6 and S7 of the

Supplementary Appendix for this case), and is approximately independent of population structure

with large d and $m \gg r$. The full solution for x_n in generation n in terms of the representation of A^n

by the eigenvalues and eigenvectors of A is given in section 3 of the Supplementary Appendix. In

practice, however, it simpler to iterate the basic matrix recursion, which takes only a few seconds to

iterate several million generations on a laptop computer.

It is also of interest to examine the initial rates of change of the Ts using the starting point of

an instantaneous sweep to equilibrium described above. The corresponding initial values of F_{ATw}

and F_{ATb} are then both equal to $1 - y/(2xy + y^2) = x/(1 + x)$; $\tilde{F}_{ATw} = \tilde{F}_{ATb} = x$. The initial F_{AT} statistics

can thus be substantially different from zero despite the absence of any divergence between In and

St, due to the assumed lack of variability within inverted chromosomes. This is seen in other

systems such as X-Y comparisons when a new Y chromosome lacks variability (Bergero et al.

2019); Toups et al. 2019); Gammerdinger et al. 2020). Normalizing Equations (A2) by dividing

each term by $2N_T$, neglecting second-order terms in the products of the deviations of the T's from

their equilibrium values with m, r and $1/N$, and exploiting the fact that $4Nm \approx (1 - F_{ST})/F_{ST}$ in order

to simplify Equations (A6c - A6e), the following simple relations hold for the initial changes per

generation in the T's:

537
$$\Delta T_{11w} \approx \frac{1}{2N_T} + 2ry \tag{6a}$$

538
$$\Delta T_{11b} \approx \frac{1}{2N_T} + \frac{2ry}{(1-F_{ST})} \tag{6b}$$

539
$$\Delta T_{22w} \approx \frac{1}{2N_T} + \frac{2mF_{ST}}{(1-F_{ST})} - \frac{1}{2Ny}$$

540
$$= \frac{1}{2N_T} + \frac{1}{2N} - \frac{1}{2Ny} \quad \text{(for } m > 0) \tag{6c}$$

541
$$\Delta T_{22b} \approx \frac{1}{2N_T} - \frac{2m}{d(1-F_{ST})} = \frac{1}{2N_T} - \frac{1}{2N_T} = 0 \tag{6d}$$

542
$$\Delta T_{12w} \approx \frac{1}{2N_T} + \frac{2mF_{ST}}{(1-F_{ST})} - rx$$

543
$$= \frac{1}{2N_T} + \frac{1}{2N} - rx \quad \text{(for } m > 0) \tag{6e}$$

544
$$\Delta T_{12b} \approx \frac{1}{2N_T} - \frac{rx}{(1-F_{ST})} - \frac{2mF_{ST}}{d(1-F_{ST})} \approx \frac{1}{2N_T} - \frac{rx}{(1-F_{ST})} \tag{6f}$$

546   Using the results in section S4 of the Supplementary Appendix, we also have:

$$\Delta F_{ATw} \approx -\frac{x}{(1+x)^2}\left[\frac{1}{2N_T y} - \frac{1}{Ny} + 2r(1+x)\right] \quad\quad (7a)$$

$$\Delta F_{ATb} \approx -\frac{x}{[1+x(1-F_{ST})]^2}\left[\frac{(1-F_{ST})}{2N_T} + 2r(1-2x)\right] \quad\quad (7b)$$

These results require rx and ry to both exceed $1/2N_T$, and hence are invalid for very low rates

of recombination, as in the case of $\rho = 0.4$ shown in Figures 3-5 below.

Equations (6a), (6c), (6e) and (7a) can be applied to the panmictic case, by setting the right-

hand terms involving m and $1/2N$ (but not $1/2Ny$) to zero, and equating N and N_T.

Results for a single randomly mating population

The results for a single, randomly mating population of size $N = N_T$ are considered first, initially

assuming an instantaneous sweep of a new inversion to its equilibrium frequency. This case was

previously studied by Navarro et al. (2000) using coalescent simulations. Some representative

results are shown in Fig. 3, for three different recombination rates (10^{-7}, 10^{-6} and 10^{-5}, i.e., scaled

recombination rates of $\rho = 4Nr = 0.4$, 4 and 400, respectively), and two different equilibrium

frequencies, $x = 0.1$ and $x = 0.5$. Time is measured relative to $2N = 2N_T$, and the results are thus

invariant with respect to N for constant ρ (unless N is sufficiently small that the assumptions of the

coalescent process are violated). The accuracy of the approximations used to generate these results

was checked by coalescent simulations and found to be excellent (see Table S2). Note, however,

that the standard deviations of the coalescent times are of similar size to the means, as is usual for

quantities generated by exponential distributions. Further details of results based on the recursion

relations are given in Table S3.

As shown above, the initial value of F_{AT} after an instantaneous sweep of the inversion to its

equilibrium frequency is simply $x/(1 + x)$. F_{AT} does not necessarily change monotonically over time,

as can be seen for the case of $x = 0.1$ with $r = 10^{-6}$, where it first decreases and then increases. With

the highest recombination rate, the final direction of change of F_{AT} can be opposite to that for T_{12}

because its initial value is greater than its equilibrium value, so that F_{AT} declines over time,

reflecting the negative term in r in Equation (7a). In contrast, T_{12} increases on the approach to

equilibrium, after an initial decline when r is large, as predicted by Equation (6e) (see the results for

$x = 0.5$ and $r = 10^{-5}$).

As expected intuitively, and as is consistent with Equation (6a), the mean coalescent time

for the inversion (T_{11}) always increases with time, reflecting the effect of recombination in causing

it to share ancestry with the standard sequence, and the fact that the carriers of the inversion

increase in numbers away from a complete bottleneck haploid population size of one. Even with the

lowest frequency of recombination ($\rho = 0.4$), there is a relatively fast approach of T_{11} towards its

equilibrium value compared with F_{AT} and T_{12}, over a timescale that is much less than the neutral

coalescent time with a low inversion frequency. Except for the lowest recombination rate, relatively

large values of F_{AT} and T_{12} (when compared with the values for the higher two recombination rates)

are approached over this timescale, reflecting their large equilibrium values. With the lowest

recombination rate, however, there is a long period of time when F_{AT} and T_{12} are both much lower

than their equilibrium values. Consistent with Equation (6c), T_{22} decreases initially for all three

recombination rates, although very slowly if Ny is close to N. In some cases (e.g., with $x = 0.5$ and r

$= 10^{-5}$), F_{AT} changes non-monotonically, with an initial decrease followed by an increase. This

reflects the fact that, for these parameter values, Equation (7a) predicts an initial decrease in F_{AT},

whereas its final value is greater than its initial value.

A simple analytical solution to the trajectories of the mean coalescent times for the case of

no recombination can also be derived (see section S6 of the Supplementary Appendix). In this case,

T_{12} increases linearly with time on the coalescent timescale (Equation S5a), whereas T_{11} and T_{22}

experience an exponential decay of their deviations from their respective equilibrium values of x

and y, with rate constants x and y. In the case of an instantaneous sweep of In to its equilibrium

frequency, this implies that T_{11} always increases over time, and T_{22} always decreases. For

sufficiently large T, these expressions imply that F_{AT} increases monotonically towards one, due to

the fact that T_{12} increases without bound in the absence of recombination. Approximations to these

expressions for small T show that F_{AT} always increases initially (see Equations (S5e) – (S5i) in the

Supplementary Appendix).

Results for a subdivided population

The numerical values of the population statistics were generated under the assumptions concerning

the total population size and local deme size used for Figs. 1 and 2. Figs. 4 and 5 show the results

for two different values of F_{ST} at neutral sites independent of the inversion, corresponding to scaled

migration rates of $4Nm = 19$ and 5.67, respectively. Further details are given in Tables S4 and S5.

The results are broadly similar to those for the panmictic case described above. The most

notable difference is that there is a short initial period with rapid increases in T_{12w} and F_{ATw}. For F_{ATw},

this period is followed by a monotonic decrease if the equilibrium value of F_{ATw} is lower than the

maximum values it achieves, or (in the case of $x = 0.1$ and $r = 10^{-6}$) a decrease followed by an

increase) when its equilibrium value exceeds its minimum value. In contrast, T_{12b} and F_{ATb} behave

initially much like T_{12} and F_{AT} in the panmictic case. For the case of $r = 0$ and a large number of

demes (d), when dMT is very small Equations (S9c) and (S9d) show that $T_{12w} \approx 1 + dT$, whereas T_{12b}

$\approx (1 + M)/M + T$. The numerical results used in Figs. 4 and 5 show, however, that T_{12w} and T_{12b}

quickly converge and have nearly identical trajectories after $T = 0.005$ (about 10,000 generations

with the parameters used here). It is easily shown from Equations (A2e) and (A2f) for the case of no

recombination and an instantaneous sweep of the inversion that:

$$T_{12b}(T) - T_{12w}(T) \approx [T_{12b}(0) - T_{12w}(0)]\exp{(-dMT)} = M^{-1}\exp{(-dMT)} \quad (8)$$

The two measures of T_{12} thus converge rapidly when $r = 0$; since it assumed here that migration is a

more powerful force than recombination, this is true more generally,

Similarly, Equation (S20e) shows that the coefficient of T in the numerator of the expression

for F_{ATw} with no recombination involves dx, so that F_{ATw} also increases very fast initially when the

number of demes is large, especially if x is close to one. In contrast, there is no contribution from d

to the expression for F_{ATb} (Equation S20f); F_{ATb} starts, however, from a higher initial value than F_{ATw}

because of population subdivision.

When the recombination rate is sufficiently high, Equations (6) and (7) for the initial rates of

change in the population statistics can be applied. ΔT_{12w} in Equation (6e) involves the sum of

$1/(2N_T)$ and $1/(2N)$, whereas the term in $1/(2N)$ is absent from the corresponding expressions for

ΔT_{12b} as well as ΔT_{12} in the panmictic case. Since $1/(2N)$ is larger than $1/(2N_T)$ by a factor of d, this

term has a large effect on the initial rate of change of T_{12w}. Similarly, the expression for ΔF_{ATw}

(Equation 7a) involves the positive term $(1 - 1/2d)/Ny$, whereas the corresponding expression for

ΔF_{ATb} involves $(1 - F_{ST})/(2Nd)$, which is negative. These results show how it is possible for the two

measures to change initially in opposite directions.

Discussion

General considerations

The results described here are broadly consistent with previous theory on the patterns of diversity

associated with balanced polymorphisms (Kaplan et al. 1988; Hudson and Kaplan 1988; Hudson

1990; Takahata 1990; Charlesworth et al. 1997; Nordborg 1997; Navarro et al. 2000; Charlesworth

et al. 2003; Innan and Nordborg 2003; Nordborg and Innan 2003; Guerrero et al. 2012; Kirkpatrick

and Guerrero 2014; Zeng et al. 2021), but extend it in several ways. From the purely technical point

of view, the recursion relations for mean pairwise coalescence times in a structured population

(Nagylaki 1998) provide a simple and computationally efficient method for calculating the expected

values of population statistics and their trajectories, on the assumption that the alleles at the target

of selection are maintained at constant frequencies. The only previous application of this method to

balanced polymorphisms appears to have been that by Zeng et al. (2021) for the case of a single

population.

The chief novelty is the application of this method to a subdivided population, for the simple

case of an autosomal locus and a finite island model of d demes of equal size N, under a Wright-

Fisher model of drift for which the effective population size of a single deme equals N. The

"migration effective population size" (Nagylaki 1998), which determines the mean coalescent time

for a pair of alleles drawn from the same deme at a locus independent of the balanced

polymorphism, is then given by $N_T = Nd$, provided that Nm is not very close to 0. This result applies

more widely to all conservative migration models, where the mean allele frequency across demes is

not changed by migration (Nagylaki 1982, 1998). For more general drift models, N can be replaced

by N_e, the effective population size of a deme, given certain restrictions (Charlesworth and

Charlesworth 2010, p.327). At recombination-drift equilibrium, the mean coalescent time for this

sampling scheme is also equal to the mean within-karyotype coalescent time, if coalescent times for

the two arrangements are weighted by the arrangement frequencies (Equation 3) and $N_T r$ is not

close to 0. For this reason, the various coalescent times used here have been expressed relative to

$2N_T$.

For simplicity, the rest of the discussion will refer only to two arrangements with respect to

an inversion polymorphism, but identical results apply to other types of biallelic polymorphisms. In

principle, it is possible to extend this approach to more general situations, such as polymorphisms

for multiple different arrangements, sex chromosomes, and non-random mating populations, as well

as changes in population size. Its main drawback is that it is limited to mean pairwise coalescent

times, and hence cannot provide information on sampling properties or on features such as the site

frequency spectrum at neutral sites linked to the target of balancing selection. For these properties,

either coalescent simulations or more complex analytical approaches, such as the phase-type theory

of Zeng et al. (2021), are needed.

Interpreting population genomic data on inversion polymorphisms

The results on equilibrium mean coalescent times for pairs of alleles, shown in Figures 1and 2,

suggest that the scaled mean coalescent time T_{12} for a pair of alleles sampled from the two different

arrangements (*In* and *St*) provides the most meaningful basis for interpreting their level of sequence

divergence. Somewhat surprisingly, for the model of population subdivision studied here, Equation

(A6e) shows that, when the number of demes is large, there is essentially no difference in

equilibrium T_{12} between alleles sampled from the same deme versus alleles from two different

demes, at least when F_{ST} at neutral sites independent of the inversion is moderate. This property

reflects the fact that, with large d, the probability that two alleles sampled from two distinct demes

were derived from the same deme in the recent past is very small compared with the probability that

alleles sampled from the same deme were derived from separate demes. This causes the properties

of pairs of alleles sampled from the same or different demes (other than their probabilities of

coalescing) to converge; there is, of course, no immediate chance of coalescence of two alleles

sampled from different arrangements.

The numerical results on the time courses of the T_{ij}'s with population subdivision (Tables S4

and S5) show that the lack of dependence of T_{12} on the nature of the sample (within or between

demes) holds very soon after the equilibrium frequency of the inversion has been reached. In

addition, except for the lowest scaled recombination rate considered here ($\rho = 0.4$), the ratio of T_{12}

to the weighted mean scaled coalescent time within karyotypes and within demes (T_{Sw}) approaches

its equilibrium value quite fast, despite the fact that both variables deviate considerably from their

equilibrium values For example, with panmixia, $x = 0.1$ and $\rho = 4$, the equilibrium value of T_{12}/T_S is

1.5 ($T_S = 1$), respectively; at time $T = 0.050$, $T_{12} = 1.039$, $T_S = 0.916$, and $T_{12}/T_S = 1.14$; at $T = 0.5$,

$T_{12} = 1.250$, $T_S = 0.926$, and $T_{12}/T_S = 1.37$. With $\rho = 40$, the equilibrium T_{12}/T_S is 1.05; at $T = 0.050$,

$T_{12}/T_S = 1.039$; by $T = 0.1$, $T_{12}/T_S = 1.04$. It is, however, not always the case that T_{12}/T_S is initially

lower than its equilibrium value; for $x = 0.5$, its initial value is 2 compared with 1.73 at $T = 0.1$ with

$\rho = 4$ and a final value of 1.55.

These findings are important for the interpretation of population genomic data for several

reasons. First, if there is evidence of significant population subdivision at neutral loci independent

of the inversion, the measure of sequence divergence between *In* and *St*, estimated by Nei's d_{xy} (Nei

1987), is best obtained from the mean of within-population samples, since its expectation is equal

to $2N_T T_{12} u$ and $4N_T u$ is equal to the expected within-population neutral diversity in the absence of an

inversion polymorphism. Second, T_{12}/T_{Sw} (which is equal to T_{12} for an equilibrium population) can

estimated by dividing an estimate of d_{xy} either by an estimate of the mean within-karyotype, within-

population diversity (π_{Sw}), or by an estimate of the corresponding mean diversity at independent

neutral sites (π_{nw}). This is because the expectations of both these statistics are equal to $4N_T u$ under

the infinite sites model of mutation (for the same mutation rate), unless there is a very low

frequency of recombination, or the inversion has arisen very recently. To control for possible

differences in mutation rate, it is preferable to use the ratio of an estimate of d_{xy} to an estimate of π_{Sw}

in order to obtain T_{12}. As discussed above in the section *The model and its analysis*, the theoretical

value of the commonly used statistic $\tilde{F}_{ATw}$, obtained by applying the $<F_{ST}>$ statistic of Hudson,

Slatkin and Maddison (1992) to *In* versus *St* haplotypes is equivalent to $1 - T_{Sw}/T_{12}$. However, some

published studies report estimates of $\tilde{F}_{ATw}$ or $\tilde{F}_{AT}$ statistics for samples pooled across different

populations, leading to some inaccuracies in using them for the purpose of estimation. In addition,

reciprocals have undesirable statistical properties if means are to be taken over sets of sequences; it

is preferable to estimate T_{12} directly using the formula described here.

This raises the question of whether recombinational exchange in heterokaryotypes occurs at

a sufficient rate that the condition $T_{Sw} = 1$ is likely to hold. At least for simple inversions (with only

a single pair of breakpoints) there is ample experimental evidence for recombination between *In* and

St at sites within and adjacent to the inversion in species of *Drosophila*, much of which appears to

be caused by gene conversion rather than double crossovers(Chovnick 1973; Korunes and Noor

2019; Li et al. 2022; Koury 2023), with $r = 10^{-5}$ per basepair per generation being a commonly

accepted typical rate (Chovnick 1973; Korunes and Noor 2019). Complex inversions, with three or

more breakpoints, might be expected to have much lower rates of exchange than simple inversions,

but heterozygotes for multiply inverted chromosomes in *D. melanogaster* have been found to

experience non-crossover associated gene conversion events at rates that are even higher than in the

absence of inversions (Crown et al. 2018), so that this expectation may not be well founded.

The occurrence of such "gene flux" (Navarro et al. 1997) is consistent with numerous

observations of substantial nucleotide site diversity within *In* karyotypes, but at a reduced level

compared with genome-wide diversity. In many *Drosophila* examples, F_{AT} and/or $\tilde{F}_{AT}$ increase

and/or within-inversion diversity levels decrease towards the inversion breakpoints (Andolfatto et al.

2001; Nobréga et al. 2008; Corbett-Detig and Hartl 2012; Kapun et al. 2016, 2023), suggesting that

flux is lowest near the breakpoints and highest towards the center of the inversion; an exception is

the polymorphism in *D. subobscura* for the overlapping pair of inversions O_{3+4} versus the standard

arrangement O_{st} (Munté et al. 2005; Papaceit et al. 2012).

Patterns of this kind do not, however, necessarily imply a complete absence of exchange at

the breakpoints. Rozas et al. (1999) obtained population genetic evidence for gene conversion events

near the breakpoints of arrangements in the O chromosome system of inversions in *D. subobscura*,

and Li et al. (2022) found no evidence for an effect of proximity to breakpoints for the *dl-49*

inversion of *D. melanogaster* on the rate of gene conversion. There are, however, some mechanistic

reasons for believing that exchange rates will be lowest at the breakpoints and highest in the center

of a simple inversion (Navarro et al. 1997). Population genomic data, which reveal the effects of

recombination over very long time spans compared with laboratory crosses, offer an excellent

opportunity for exploring the effects of breakpoints on recombination.

Estimating the ages of inversions: divergence between *In* and *St*

If exchange were indeed completely suppressed around inversion breakpoints, divergence between

In and *St* for sequences close to the breakpoints could be used to estimate the date of origin of an

inversion (Hasson and Eanes 1996; Andolfatto et al. 2001; Cáceres et 2001; Corbett-Detig and

Hartl 2012). In this case, it can no longer be assumed that $T_{Sw} = 1$ for the sequences concerned.

Given the reservations about whether such suppression of exchange is absolute, caution should

probably be used in making such inferences.

If, as is usually assumed, the new arrangement had a unique origin, the initial expectation of

T_{12} in the absence of exchange is equal to that for a pair of randomly chosen alleles in the ancestral

population. In the case of a panmictic Wright-Fisher population of constant size, T_{12} at time T since

the origin of the inversions is given by $1 + T$ (Equation S5a). With population subdivision, this

expression is no longer accurate; if $dMT \gg 1$, and d is large, $T_{12} \approx (1 - F_{ST})^{-1} + T$ for both within-

and between deme measures of T_{12} (see the approximation to Equation [S9b]). The first term here

corresponds to the mean coalescent time for pairs of alleles sampled from different demes. For

species like *D. melanogaster*, with N_T of approximately one million for populations in the ancestral

species range (Lack et al. 2015), a T value of 0.1 corresponds to 200,000 generations; with $F_{ST} =$

0.05, $(1 - F_{ST})^{-1}$ is approximately 50% of this, so that ignoring this term creates a substantial error in

the estimated time of origin. The situation would clearly be worse for a species with a higher level

of population subdivision.

Under these conditions, and assuming population size stability, if there is population

subdivision, the neutral diversity estimated from pairs of alleles sampled randomly across

populations (π_{nT}) can be treated as a proxy for the corresponding mean coalescence time, and

subtracted from d_{xy}. The ratio of this corrected value to the neutral diversity statistic then provides

an estimate of T as defined here. For a very recent origin of the inversion, the full Equation (S9b)

would usually have to be applied if there is significant population subdivision. The simple

approximations of Equations (S9c) and (S9d) could be used when $dMT \ll 1$, which corresponds to

an extremely recent origin when $M \geq 1$, as is likely to be the case for between-population

differences within species of most outcrossing species of animals and plants (Charlesworth 2003;

Roux et al. 2016).

In practice, different authors have used different methods for estimating T from divergence

between arrangements. Taking some of the pioneering studies as examples, Andolfatto et al. (1999)

used the number of fixed differences between In and St, which is strongly affected by sample size.

Cáceres et al. (2001), Hasson and Eanes (1996) and Corbett-Detig and Hartl (2012) used d_{xy}

corrected by subtraction of the within-St diversity. Somewhat different values will be generated by

each of these methods. Difficulties clearly arise if there is evidence for recent strong population

expansions or contractions, so that the current population statistics cannot be equated to their values

at the time of origin of In. There are also considerable estimation errors, especially as the d_{xy} values

are often very small for $D.$ $melanogaster$ inversions (see Table S2 in Corbett-Detig and Hartl

[2012]).

Estimating the ages of inversions: diversity within In

The nucleotide site diversities within arrangements also provide information on the age of an

inversion, assuming an absence of exchange. If the inversion frequency is low, however, the within-

inversion within-deme diversity ($\pi_{11w} = 2N_T T_{11w} u$) is only a small fraction of the diversity within the

standard sequence or at sites independent of the inversion (see Fig.1), so that a reduced level of

within-inversion diversity does not necessarily imply a recent origin of the inversion. In the case of

a subdivided population, provided that $dT \gg x(1 + Mx)$, Equation (S18a), which corresponds to the

exponential growth model used for simulation-based estimates of inversion age by Corbett-Detig

and Hartl (2012), implies that:

$$T \approx -x \ln(x - T_{11w}) \qquad\qquad (9)$$

This expression is conveniently independent of M. Clearly, it is only meaningful if $T_{11w} < x$.

Given values of x, mean diversity within the inversion (π_{11w}) and π_{nw} as defined above, it is possible

to estimate T from this formula, equating π_{11w}/π_{nw} to T_{11w}. For example, Table 1 of Andolfatto et al.

(1999) provides estimates of diversity values of 0.0009 and 0.0125 for a breakpoint of $In(2L)t$ and

for independent sites, respectively, in a $D.$ $melanogaster$ population with an inversion frequency of

0.23. T_{11w} is thus estimated as $0.0009/0.0125 = 0.072$, giving $T = 0.42$. Andolfatto et al. (1999)

estimated the time to the most common recent ancestor of the sequences around the breakpoint from

the standard neutral coalescent formula as 0.15 in the present notation; since this method ignores the

effect of the recovery from the sweep of the inversion on the gene genealogy, it overestimates the

opportunity for diversity to recover from its complete loss, and hence underestimates the time since

the origin of the inversion.

Similar calculations applied to the data in Figs. 2 and S1B of Kapun et al. (2023) on the

Zambian population of *D. melanogaster* for sites within 100kb of the distal and proximal

breakpoints of *In(3R)P* yield estimates of T_{11w} of 0.59 and 0.11, respectively. These are highly

discordant, reflecting the much lower diversity at the proximal breakpoint, which is also seen in the

European and North American samples. Given the estimated frequency of 0.11 for *In(3R)P* in this

population (Kapun and Flatt 2019, Table S1), these estimates of T_{11w} are inconsistent with a total

absence of exchange near the breakpoints.

The corresponding values of T_{22w} are 1.28 and 1.13, yielding estimates of T_{Sw} of 1.20 and

1.02, respectively, so that the data are reasonably consistent with recombination-drift equilibrium

for sites close to the breakpoints, although the flux rates must be substantially lower than the mean

of somewhat less than 10^{-5} estimated for the inversion as a whole (see the next section). The

numerical results in Supplementary Table S1 for the panmictic case suggest r values around 10^{-7}

and 10^{-6} for the distal and proximal breakpoints, respectively. Caution should, however, be used in

interpreting these conclusions, as they are highly sensitive to the frequency of the inversion in

question, and there is considerable continent-wide variation in the frequency of inversions such as

In(3R)P within Africa (Kapun and Flatt 2019, Table S1); Sprengelmeyer et al. (2020) estimated a

frequency of 0.23 for a small Zambian sample. However, the data for Zambian, Bechuanaland and

Swaziland samples in Table 1 of Aulard et al. (2002), yield an overall frequency of 0.11 with s.e. of

0.024, so that a frequency around 0.10 for this region of Africa seems reasonable.

An alternative has been to assume that the recent sweep of an inversion results in a star

phylogeny, so that the expected pairwise diversity within the inversion (in the absence of exchange)

is equal to $2Tu$ (Rogers 1995); Rozas et al. (1999) and Nobréga et al. (2008) used this method to

estimate the ages of several inversions of *D. subobscura*. Since a star phylogeny must have the

smallest mean divergence between a pair of alleles compared with a gene tree with one or more

coalescent events after the origin of the inversion, this method overestimates the age of the

inversion.

Estimating the exchange parameter

If equilibrium can be assumed, estimates of between-arrangement divergence and within-

arrangement diversities can be compared with the theoretical predictions for the corresponding

variables, in order to obtain estimates of the rate of exchange r (mutation rate estimates are also

needed to estimate r rather than ρ).

The figures show that T_{11w}/T_{22w}, as well as T_{12}, are strongly affected by the level of

population subdivision, unless recombination is rare or absent. Even a modest change from $F_{ST} = 0$

to $F_{ST} = 0.05$ causes this ratio to shift from close to 0.8 to approximately 0.6 when $r = 10^{-5}$ and the

inversion frequency is 0.1 (Figure 1C), reflecting the effect of population subdivision in multiplying

the effective rate of recombination within populations by $1 - F_{ST}$ (Wakeley and Lessard 2003).

Estimates of F_{ST} thus need to be included in any such attempts to estimate r. Another difficulty here

is that the current model assumes a long-term constancy of the inversion frequency over time, as

well as an absence of among-population variation in inversion frequencies between populations.

Many studies of *Drosophila* and other groups have revealed major differences among populations

in the frequencies of polymorphic inversions, often clinal in nature, which are likely to reflect

locally varying selection pressures on genes contained in the inversion, e.g., Krimbas and Powell

(1992b), Aulard et al. (2002), Cheng et al. (2012), Kapun and Flatt (2019) and Mérot et al. (2021).

Further work is needed to evaluate the consequences of relaxing the assumption of constancy.

This raises the question of whether equilibrium can safely be assumed in such analyses.

Studies of inversion polymorphisms in many *Drosophila* species suggest that they rarely persist

beyond species boundaries (Krimbas and Powell 1992a). There are examples of very close relatives

with totally different sets of inversion polymorphisms, and many fixed differences with respect to

gene arrangements, e.g., *D. miranda* versus *D. pseudoobscura* (Bartolomé and Charlesworth 2006).

It is thus quite likely *a priori* that the times of origin of many polymorphic inversions are only a

small fraction of the mean neutral coalescent time, $2N_T$, as was proposed by Andolfatto et al. (2001)

in their review of early data on the molecular population genetics of *Drosophila* inversions.

Figs. 3-5 show that for the two higher recombination rates 10^{-6} and 10^{-5} corresponding to

scaled recombination rates (ρ) of 4 and 40, the times for statistics such as T_{11}, T_{12} and F_{AT} to

approach their equilibrium values after the inversion has approached its equilibrium frequency are

either commensurate with $2N_T$ ($r = 10^{-6}$) or much smaller ($r = 10^{-5}$), consistent with the theoretical

prediction that the timescale of approach to equilibrium in terms of generations is of the order of

$1/r$, both for a panmictic population and a subdivided population with a large number of demes (see

section S2 of the Supplementary Appendix). In the absence of recombination, however, diversity

within the inversion recovers over a timescale of $2xN_T$ generations for a panmictic population and

$2(1 + Mx)N_T/M$ generations for a subdivided population (see Equations S5b and S18a), so that the

signature of the selective sweep of the inversion on within-karyotype diversity can then persist for a

long time, as has been noted previously (Navarro et al. 2000; Zeng et al. 2021). In addition, the fact

that the ratio T_{12}/T_{Sw} converges on its equilibrium value much faster than the absolute T_{ij}'s (see the

above section *Interpreting population genomic data on inversion* polymorphisms) means that it is

not even necessary to assume that the population is very close to equilibrium when using the

corresponding divergence to diversity ratio as a statistic. For a low frequency inversion, a similar

result applies to T_{11w}/T_{Sw}.

If recombination in heterokaryotypes is as frequent as is suggested by the data on

Drosophila (Korunes and Noor 2019), the assumption of recombination-drift equilibrium may thus

often be reasonably accurate as a predictor of observed patterns of population genomic statistics for

genomic regions covered by an inversion polymorphism, at least for sites that are located well away

from inversion breakpoints. The equilibrium assumption can be tested by comparing the within-*In*

neutral diversity level against the diversity at neutral sites independent of the inversion (or against

the within-*St* diversity). A very recent sweep of the inversion, leaving the system far from

equilibrium, would cause diversity across the whole length of the inversions to be much lower than

a fraction x of the diversity outside the inversion or $x/(1 - x)$ of diversity within the standard

arrangement – see the curves in Figures 3-5. It would also be expected to leave a signature of an

excess of low frequency variants at neutral sites compared with what is seen at comparable sites

independent of the inversions (Navarro et al. 2000; Andolfatto et al. 2001; Zeng et al. 2021),

although the effects of background selection and selective sweeps within the low recombination

environment created by a low-frequency inversion could also cause such a pattern (Charlesworth

and Jensen 2021). The inversion *In(1)Be* in African populations of *D. melanogaster* is an example

of a very recent spread of an inversion (Corbett-Detig and Hartl 2012).

The study by Kapun et al. of *In(3R)P* in the Zambian population (ZI) of *D. melanogaster*

provide an example of how to apply the theoretical results. As mentioned in the section *Theoretical*

results: equilibrium populations; Numerical results for subdivided populations, $<F_{ST}>$ for this

geographical region is so low that panmixia can be assumed, which implies that Equations (5) can

be used for the expected coalescence times. Let $X = T_{11}/T_{22}$. Using Equations (5b) and (5c), simple

algebra yields following formula for ρ, the scaled recombination rate:

$$\rho = \frac{X(1-2x)}{(1-X)xy} \qquad (10)$$

Fig. 2 of Kapun et al. (2023) show that the mean nucleotide site diversity (across all classes

of nucleotide sites) is significantly lower (0.00792) for the *In(3R)P* haplotypes than for the standard

haplotypes (0.00979), giving an estimate of $T_{11}/T_{22} = 0.00792/0.00979 = 0.81$ for the region

covered by the inversion; $x \approx 0.11$ for this population (Kapun and Flatt 2019, Table S1).

Substituting these numbers into Equation (10) yields an estimate of 34 for ρ, corresponding to $r =$

8.5×10^{-6} with $N^T = 10^6$, close to the estimate of 10^{-5} from crossing experiments (Korunes and Noor

2019). The estimate of diversity for regions outside the inversion on chromosome 3 is 0.00854,

giving an estimate of $T_{11}/T_S = (0.1 \times 0.00792 + 0.89 \times 0.00979)/0.00854$ of 1.12, slightly higher than

the theoretical value of 1 for equilibrium; this discrepancy probably reflects different levels of

selective constraints among the genome regions being compared and/or the fact that inversions

suppress exchange well outside their breakpoints (Li et al. 2022; Koury 2023).

Fig. 4 of Kapun et al. (2023) shows that $\tilde{F}_{AT}$ for regions of *In(3R)P* in the Zambian

population that are remote from the breakpoints is approximately 0.1. Equation 3a implies that:

921                    $$\rho = \frac{2(1-\tilde{F}_{AT})}{\tilde{F}_{AT}} \qquad (11)$$

This expression yields an estimate of $\rho = 18$, considerably less than the above value. This

may be due to the fact that the estimate of $\tilde{F}_{AT}$ involves taking the mean of a reciprocal over 100kb

windows, which biases it toward low values compared with using the mean of estimates of T_{12}.

The empirical results on *In(3R)P* are thus consistent with this inversion being close to

recombination-drift equilibrium, with a mean flux rate within the inversion of somewhat less than

10^{-5}. The noticeable increase in within-inversion diversity towards the middle of the inversion (Fig.

S1), and the increase in $\tilde{F}_{AT}$ near the breakpoints (Fig.4), strongly indicate that flux rates are

reduced close to the breakpoints, but the above analysis of the diversity data near the breakpoints

suggests that this reduction is not total.

The relevance of F_{AT} and $\widetilde{F}_{AT}$

Another point concerns the interpretation of F_{AT}, the analogue of F_{ST} for differentiation between *In*

and *St*), and the related statistic $\tilde{F}_{AT}$; see Equations (4). These statistics are sometimes loosely

referred to as measures of the extent of divergence between *In* and *St*, e.g., Guerrero, et al. (2012),

Kennington and Hoffmann (2013) and Kapun and Flatt (2019). As noted in the section *Theoretical*

results: Approach to equilibrium, General considerations, the fact that F_{AT} and $\tilde{F}_{AT}$ do not

necessarily measure the extent of divergence between karyotypes is brought out by their high initial

values immediately after an instantaneous sweep to an equilibrium frequency of x, for which $F_{AT} =$

$x/(1 + x)$, and $\tilde{F}_{AT} = x$. This reflects the fact that *In* and *St* are initially no more divergent on average

than a random pair of sequences, while there is no diversity within the inversion. Furthermore, in

the presence of recombination F_{AT} can reach an equilibrium level that is lower than its initial value

whereas T_{12} increases above 1 (e.g., Figure 4, lower middle panel), so that the two statistics can

change in opposite directions over time (see also Zeng et al. 2021, Fig. 8).

As described earlier, F_{AT} is equivalent to the measure of linkage disequilibrium σ_d^2 of Ohta

and Kimura (1971), which is approximately equal to R^2, the squared correlation coefficient between

two loci of Hill and Robertson (1968), so that the use of F_{AT} is effectively equivalent to estimating

linkage disequilibrium (LD) between SNPs and karyotype. This equivalence does not hold for $\tilde{F}_{AT}$,

however, which yields much larger values than F_{AT} for equilibrium situations (Charlesworth 1998;

Gammerdinger et al. 2020), and has frequently been used to characterize differentiation between

arrangements (e.g., Corbett-Detig and Hartl [2012], Kapun et al. 2023). This difference arises

because the theoretical value of $\tilde{F}_{AT}$ is equal to $1 - T_S/T_{12}$ whereas $F_{AT} = 1 - T_S/T_T$ with $T_T < T_{12}$ at

recombination-drift equilibrium (see Equations 4). The difference between them can be

considerable; Table 2 of Laayouni et al. (2003) shows both statistics for inversion polymorphisms

of *D. buzzatii*, with approximately four-fold larger values of $\tilde{F}_{AT}$ compared with F_{AT}. If data on

diversity within inverted and standard haplotypes are available, as well as an estimate of x, it is

possible to determine F_{AT} from an estimate of $\tilde{F}_{AT}$ and vice-versa. In the case of *In(3R)P* in the

Zambian population of *D. melanogaster*, the mean $\tilde{F}_{AT}$ value of 0.1 combined with the diversity

values in Figure 2 of Kapun et al. (2023) gives mean $F_{AT} \approx 0.033$, close to a direct estimate (Thomas

Flatt and Martin Kapun, personal communication). The magnitude of LD between single nucleotide

variants (SNPs) and an inversion polymorphism can therefore be rather small, even when there is

noticeable sequence differentiation between arrangements.

The larger expected equilibrium value of $\tilde{F}_{AT}$ compared with F_{AT}, as well as its closer

relationship with the divergence between arrangements, makes it a more powerful measure of the

extent of differentiation between arrangements relative to within-arrangement diversity. It is also

more convenient for use with population genomic data based on haploid genome sequences, as is

the case for much of the *Drosophila* Genome Nexus data (Lack et al. 2015, 2016) and for the data

in Kapun et al. (2023), because it does not require the reconstruction of the frequencies of diploid

genotypes used for calculating the mean diversity over a random set of individuals. As pointed out

above, however, direct estimates of T_{12} have better statistical properties.

Linkage disequilibrium among neutral variants associated with inversions

As just discussed, estimates of F_{AT} can be used to estimate LD between neutral variants such as

silent site SNPs and an inversion polymorphism. It is also of interest to examine LD between the

SNPs themselve, as this has been used to infer the existence of inversion polymorphisms in non-

model organisms from evidence for large blocks of LD in specific regions of the genome (e.g.,

Faria et al. 2019; Mérot et al. 2021). It is difficult to obtain exact analytical results on LD in

structured populations (Wakeley and Lessard 2023), but some approximations are easily obtained.

From Equation (A9b) of Charlesworth et al. (1997), if LD within arrangements is small compared

with LD between neutral sites and the inversion polymorphism, σ_d^2 for a pair of SNPS is

approximately equal to the product of σ_d^2 for each SNP and the inversion polymorphism. The

standard formula for a partial correlation coefficient implies that this is also true for the R^2 statistic

of Hill and Robertson (1968), if there is little or no correlation between SNPs within arrangements.

If there is LD within arrangements, these products somewhat underestimate the corresponding

statistics for the pairs of SNPs. If there is a major effect of divergence between arrangements on LD

between SNPs and karyotype, there should be little dependence of σ_d^2 or R^2 on the physical distance

between SNPs across the region where crossing over is suppressed by the inversion, in contrast to

what is expected for within-karyotype patterns of LD (e.g., Wakeley and Lessard 2003).

Figure S3 of Kapun et al. (2023) shows a pattern of elevated R^2 among SNPs across the

entire region of *In(3R)P* in the Zambian population, contrasted with the rapid decay of LD with

physical distance between SNPs within inverted and standard haplotypes. As expected from the

product formula, the magnitude of individual R^2 values is modest in this case – the estimate of

0.033 for F_{AT} given above gives an expected R^2 of approximately 0.001 for a pair of SNPs in

genomic regions affected by the inversion polymorphism. Much stronger patterns are, however,

seen in the non-African samples studied by Kapun et al. (2023), which may reflect the effects of

population bottlenecks or recent selection. This example illustrates the point that substantial

associations between SNPs and an inversion polymorphism may truly exist, but could be hard to

detect simply from LD patterns in samples without prior knowledge of the existence of the

inversion polymorphism. Conversely, false signals of an inversion polymorphism could be

generated from localized clusters of LD in bottlenecked populations (e.g., Haddrill et al. 2005). Of

course, if there are virtually complete associations between SNPs and arrangement, as in the

inversion polymorphisms of *Coelopa frigida* (Mérot et al. 2021), there will be a strong pattern of

localized LD blocks (see their Fig. 2).

1005

Effects of inversions on neutral divergence between populations

For the case of populations with a large number of demes, the extent of population differentiation at

a neutral locus associated with an inversion can be measured by F_{ST} as defined here (which in this

case is nearly the same as the $<F_{ST}>$ statistic of Hudson, Slatkin and Maddison 1992) for standard

and inverted haplotypes, denoted here by $F_{ST, St}$ and $F_{ST, In}$, respectively – see Equation (S23).

Provided that $m >> r$, which is likely to be true for sites within the inversion or close to the

breakpoints, the equilibrium values of $F_{ST, In}$ and $F_{ST, St}$ are equal to $1/(1 + Mx)$ and $1/(1 + My)$,

respectively, which are somewhat surprisingly independent of r (Equations [A6h] and [A6i]). These

expressions are similar in form to that for a subdivided population in the absence of a

polymorphism, with M replaced by Mx and My. For $r = 0$, it can be shown that these values are

reached almost instantaneously once the inversion has reached its equilibrium frequency (Equation

S23). Numerical examples show that this is also true for $r > 0$. The rapid equilibration of F_{ST} in

subdivided populations is well known (Crow and Aoki 1984; Pannell and Charlesworth 2000).

These results imply that within-arrangement haplotype F_{ST} or $<F_{ST}>$ statistics should be

larger for loci within the inversion than for loci that are independent of the inversion, independently

of location with respect to the inversion breakpoints. A low-frequency inversion should thus show a

considerable inflation in the between-populations F-statistics, even in the absence of differences in

karyotype frequencies between populations, whereas only a small effect should be seen for the

standard arrangement; such a pattern was reported by Kennington et al. (2013) for the *D.*

melanogaster inversion *In(2L)t*. Caution should therefore be used in interpreting such a pattern as

evidence for spatial differences in selection pressures.

In contrast, if inversion frequencies vary considerably between populations because of

divergent selection, the LD between neutral sites and karyotype will cause among-population

differentiation at neutral sites (Charlesworth et al. 1997; Guerrero et al. 2012), as has been found in

some studies of *Drosophila* (Pegueroles et al. 2013) and other taxa (Faria et al. 2019; Mérot et al.

2021). If exchange rates increase with distance from breakpoints, then higher values of $F_{ST, In}$ and

$F_{ST, St}$ (or the corresponding $<F_{ST}>$ statistics) are expected towards the centers of inversions, as has

been observed in some cases, such as Inv2La in *Anopheles gambiae* (Cheng et al. 2012).

1034

Conclusions

The theoretical results described here provide some pointers on how to interpret population

genomic data on inversion polymorphisms, especially caveats concerning methods for estimating

the ages of inversions. They are subject to several important limitations. In particular, they are

based purely on expectations for pairwise coalescence times. Estimates obtained by equating such

expectations to observed values are, of course, subject to considerable stochastic variation, and no

attempt has been made here to assess error bounds on estimates. In addition, many simplifying

assumptions have been made, including ignoring the consequences of selection on loci responsible

for maintaining the inversion polymorphisms, as well as the effects of Hill-Robertson interference

in the low recombination environment characteristic of low frequency inversions. There is,

therefore, plenty of scope for further work.

1046

1047 **Data Availability**

1048 No new data or reagents were generated by this research. The codes for the computer programs

1049 used to produce the results described below will be made available on Figshare on acceptance.

1050

1051 **Acknowledgments**

1052 I thank Deborah Charlesworth and Thomas Flatt for helpful discussions, and Thomas Flatt

1053 for his comments on a draft of this paper. Thomas Flatt and Martin Kapun generously shared

1054 their unpublished results on *In(3R)P*.

1055

1056

Literature cited

Andolfatto P, Wall JD, Kreitman M. 1999. Unusual haplotype structure at the proximal breakpoint of

In(2L)t in a natural population of *Drosophila melanogaster*. Genetics 153:1297-1311.

Andolfatto P, Depaulis F, Navarro A. 2001. Inversion polymorphisms and nucleotide

variability in Drosophila. Genet. Res. 77:1-8.

Assaf ZJ, Tilk S, Siegal ML, Petrov DA. 2017. Deep sequencing of natural and experimental

populations of *Drosophila melanogaster* reveals biases in the spectrum of new

mutations. Genome Res. 27:1988-2000.

Aulard S, David, J.R., Lemeunier F. 2002. Chromosomal inversion polymorphism in Afrotropical

populations of *Drosophila melanogaster*. Genet. Res. 79:49-63.

Bartolomé C, Charlesworth B. 2006. Rates and patterns of chromosomal evolution in

Drosophila pseudoobscura and *D. miranda*. Genetics 173:779–791.

Barton NH, Etheridge AM, Kelleher J, Véber A. 2013. Genetic hitchhiking in spatially

extended populations. Theor. Pop. Biol. 87:75-89.

Bergero R, Gardner J, Bader B, Yong L, Charlesworth d. 2019. Exaggerated heterochiasmy
in a fish with sex-linked male coloration polymorphisms. Proc. Natl. Acad. Sci. USA
116:6924-6931.

Buffalo V. 2021. Quantifying the relationship between genetic diversity and population size
suggests natural selection cannot explain Lewontin's paradox. Elife 10:e67509.

Cáceres M, Puig M, Ruiz A. 2001. Molecular characterization of two natural hotspots in the
Drosophila buzzatii genome induced by transposon insertions. Genome Res. 11:1353-
1364.

Charlesworth B. 2020. How good are predictions of the effects of selective sweeps on levels
of neutral diversity? Genetics 216:1217-1238.

Charlesworth B. 1998. Measures of divergence between populations and the effect of forces
that reduce variability. Mol. Biol. Evol. 15:538-543.

Charlesworth B, Charlesworth D. 2010. Elements of Evolutionary Genetics. Greenwood
Village, CO: Roberts and Company.

Charlesworth B, Jensen JD. 2021. The effects of selection at linked sites on patterns of genetic
variability. Ann. Rev. Ecol. Evol. Syst. 52:177-197.

Charlesworth B, Charlesworth D, Barton NH. 2003. The effects of genetic and geographic
structure on neutral variation. Ann. Rev. Ecol. Evol. Syst. 34:99-125.

Charlesworth B, Nordborg M, Charlesworth D. 1997. The effects of local selection, balanced
polymorphism and background selection on equilibrium patterns of genetic diversity in
subdivided populations. Genet. Res. 70:155-174.

Charlesworth D. 2003. Effects of inbreeding on the genetic diversity of plant populations.
Phil. Trans. Roy. Soc. B 358:1051-1070.

Chovnick A. 1973. Gene conversion and transfer of genetic information within the inverted
region of inversion heterozygotes. Genetics 74:123-131.

Corbett-Detig RB, Hartl DL. 2012. Population genomics of inversion polymorphisms in
Drosophila melanogaster. PloS Genet. 8:e1003056.

Crow JF, Aoki K. 1984. Group selection for a polygenic behavioral trait - estimating the
degree of population subdivision. Proc. Natl. Acad. Sci. USA 81:6073-6077.

Crown KN, Miller DE, Sekelsky J, Hawley RS. 2018. Local inversion heterozygosity alters
recombination throughout the genome. Curr. Biol. 28:2984-2990.

Dobzhansky T. 1947. Adaptive changes induced by natural selection in wild populations of
Drosophila. Evolution 1:1-16.

Dobzhansky T. 1950. Genetics of natural populations. XIX. Origin of heterosis through naturak selection in populations of *Drosophila pseudoobscura*. Proc. Natl. Acad. Sci. USA 35:288-302.

Faria R, Chaube P, Morales HE, Larsson T, Lemmon AR, Lemmon EM, Rafajlovic M, Panova M, Ravinet M, Johannesson K, et al. 2019. Multiple chromosomal rearrangements in a hybrid zone between *Littorina saxatilis* ecotypes. Mol. Ecol. 28:1375-1393.

Gammerdinger W, Toups M, Vicoso B. 2020. Disagreement in F_{ST} estimators: A case study from sex chromosomes. Mol. Ecol. Res. 20:1517-1525.

Gong WJ, McKim KS, Hawley RS. 2005. All paired up with no place to go: Pairing, synapsis, and DSB formation in a balancer heterozygote. PloS Genet. 1:e67.

Guerrero RF, Rousset F, Kirkpatrick M. 2012. Coalescent patterns for chromosomal inversions in divergent populations. Phil. Trans. R. Soc. B 367:430-438.

Haddrill PR, Thornton KR, Charlesworth B, Andolfatto P. 2005. Multilocus patterns of nucleotide variability and the demographic and selection history of *Drosophila melanogaster* populations. Genome Research 15:790-799.

Hasson E, Eanes WF. 1996. Contrasting histories of three gene regions associated with *In (3L)Payne* of *Drosophila melanogaster*. Genetics 144.

Hill WG, Robertson A. 1968. Linkage disequilibrium in finite populations. Theor. Appl. Genet. 38:226-231.

Hudson RR. 1990. Gene genealogies and the coalescent process. Oxf. Surv. Evol. Biol. 7:1-45.

Hudson RR, Boos DD, Kaplan NL. 1992. A statistical test for detecting geographic subdivision. Mol. Biol. Evol. 9:138-151.

Hudson RR, Kaplan NL. 1988. The coalescent process in models with selection and recombination. Genetics 120:831-840.

Hudson RR, Slatkin M, Maddison WP. 1992. Estimation of levels of gene flow from population data. Genetics 132:583-589.

Innan H, Nordborg M. 2003. The extent of linkage disequilibrium and haplotype sharing around a polymorphic site. Genetics 165:437-444.

Ishii K, Charlesworth B. 1977. Associations between allozyme loci and gene arrangements due to hitch-hiking effects of new inversions. Genet. Res. 30:93-106.

Ives PT, Band HT. 1986. Continuing studies on the South Amherst *Drosophila melanogaster* natural population during the 1970's and 1980's. Evolution 40:1289-1302.

Kaplan NL, Dardern T, Hudson RR. 1988. The coalescent process in models with selection and
recombination. Genetics 120:831-840.

Kapun M, Flatt T. 2019. The adaptive significance of chromosomal inversion
polymorphisms in *Drosophila melanogaster*. Mol. Ecol. 28:1263-1282.

Kapun M, Fabian DK, Goudet J, Flatt T. 2016. Genomic evidence for adaptive inversion clines in
Drosophila melanogaster. Mol. Biol. Evol. 33:1317–1336.

Kapun M, Durmaz Mitchell E, Kawecki T, Schmidt P, Flatt T. 2023. An ancestral balanced inversion
polymorphism confers global adaptation. BioRxiv 10.1101/2023.01.31.526462v1.

Kennington WJ, Hoffmann AA. 2013. Patterns of genetic variation across inversions,
geographic variation in the *In(2L)t* inversion in populations of *Drosophila*
melanogaster from eastern Australia. BMC Evol. Biol. 13:100.

Kimura M. 1971. Theoretical foundations of population genetics at the molecular level.
Theor. Pop. Biol. 2:174-208.

Kirkpatrick M, Guerrero RF. 2014. Signatures of sex-antagonistic selection on recombining
sex chromosomes. Genetics 197:531-541.

Korunes K, Noor MAF. 2019. Pervasive gene conversion in chromosomal inversion
heterozygotes. Mol. Ecol. 28:1302-1315.

Koury SA. 2023. Predicting recombination suppression outside chromosomal inversions in
Drosophila melanogaster using crossover interference theory. Heredity: in press.

Krimbas CB, Powell JR. 1992a. Drosophila Inversion Polymorphism. In. Boca Raton, FL:
CRC Press.

Krimbas CB, Powell JR. 1992b. Introduction. In: Krimbas CB, Powell JR, editors.
Drosophila Inversion Polymorphism. Boca Raton, FL: CRC Press. p. 1-52.

Laayouni H, Hasson E, Santos M, Fontdevila A. 2003. The evolutionary history of
Drosophila buzzatii. XXXV. Inversion polymorphism and nucleotide variability in
different regions of the second chromosome. Mol. Biol. Evol. 20:931–944.

Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Detig RB, Stevens KA, Langley
CH, Pool JE. 2015. The *Drosophila* Genome Nexus: A population genomic resource of
623 *Drosophila melanogaster* genomes, including 197 from a single ancestral range
population. Genetics 199:1229-1241.

Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. 2016. A thousand fly genomes: An
expanded *Drosophila* Genome Nexus. Mol. Biol. Evol. 33:3308-3313.

Langley CH, Stevens K, Cardeno C, Lee YCG, Schrider DR, Cardeno C., et al. 2012. Genomic
variation in natural populations of *Drosophila melanogaster*. Genetics 192:533-598.

Li L, Brent E, Hadjipanteli S, Galey M, Miller DE, Crown N. 2022. Heterozygous Inversion
breakpoints suppress meiotic crossovers by altering recombination repair
outcomes. BioRxiv 2022.11.09.515852.

Lange JD, Bastide H, Lack JB, Pool JE. 2022. A population genomic assessment of three
decades of evolution in a natural *Drosophila* population. Mol. Biol. Evol. 39:msab368.

Maruyama T. 1974. A simple proof that certain quantitities are independent of the geographic
structure of population. Theor. Pop. Biol. 5:148-154.

Mérot C, Llaurens V, Normandeau E, Bernatchez L, Wellenreuther M. 2018. Balancing selection via
life-history trade-offs maintains an inversion polymorphism in a seaweed fly. Nat. Commun.
11:1-11.

Mérot C, Berdan EL, Cayuela H, Djambazian H, Ferchaud A, Laporte M, Normandeau E,
Ragoussis J, Wellenreuther M, Bernatchez L. 2021. Locally adaptive inversions
modulate genetic variation at different geographic scales in a seaweed fly. Mol. Biol.
Evol. 38:3953-3971.

Mukai T, Watanabe TK, O. Y. 1974. The genetic structure of natural populations of
Drosophila melanogaster. XII. Linkage disequilibrium in a local population. Genetics
77:771-793.

Mukai T, Yamaguchi O. 1974. The genetic structure of natural populations of *Drosophila*
melanogaster. XI. Genetic variability in a local population. Genetics 76:339-366.

Munté A, Rozas J, Aguadé M, Segarra C. 2005. Chromosomal inversion oolymorphism leads
to extensive genetic structure: A multilocus survey in *Drosophila subobscura*. Genetics
169:1573-1581.

Nagylaki T. 1998. The expected number of heterozygous sites in a subdivided population.
Genetics 149:1599-1604.

Nagylaki T. 1982. Geographical invariance in population genetics. J. Theor. Biol. 99:159-
172.

Navarro A, Barbadilla A, Ruiz A. 2000. Effect of inversion polymorphism on the neutral
nucleotide variability of linked chromosomal regions in Drosophila. Genetics 155:685-
698.

Navarro A, Betrán E, Barbadilla A, Ruiz A. 1997. Recombination and gene flux caused by gene conversion and crossing over in inversion heterokaryotypes. Genetics 146:695-709.

Nei M. 1982. Evolution of human races at the gene level. In: Alan LR, editor. Human genetics, part A: The unfolding genome. New York: Liss. p. 167-181.

Nei M. 1987. Molecular Evolutionary Genetics. New York: Columbia University Press.

Nei M, Li W-H. 1975. Probability of identical monomorphism in related species. Genet. Res. 26:31-43.

Nobréga C, Khadem M, Aguadé M, Segarra C. 2008. Genetic exchange versus genetic differentiation in a medium-sized inversionof *Drosophila*: The A_2/Ast arrangements of *Drosophila subobscura*. Mol. Biol. Evol. 25:1534-1543.

Nordborg M. 1997. Structured coalescent processes on different time scales. Genetics 146:1501-1514.

Nordborg M, Innan H. 2003. The genealogy of sequences containing multiple sites subject to strong selection in a subdivided population. Genetics 163:1201-1213.

Ohta T, Kimura M. 1970. Development of associative overdominance through linkage disequilibrium in finite populations. Genet. Res. 18:277-286.

Ohta T, Kimura M. 1971. Linkage disequilibrium between two segregating nucleotide sites under steady flux of mutations in a finite population. Genetics 68:571-580.

Ohta T, Kimura M. 1969. Linkage disequilibrium due to random genetic drift. Genet. Res. 13:47-55.

Orengo DJ, Puerma E, Cereijo U, Aguadé M. 2019. The molecular genealogy of sequential overlapping inversions implies both homologous chromosomes of a heterokaryotype in an inversion origin. Scientific Reports 9:17009.

Pannell JR, Charlesworth B. 2000. Effects of metapopulation processes on measures of genetic diversity. Phil. Trans. R. Soc. B 355:1851-1864.

Pegueroles C, Aquadro CF, Mestres F, Pascual M. 2013. Gene flow and gene flux shape evolutionary patterns of variation in *Drosophila subobscura*. Heredity 110:520-529.

Prakash S, Lewontin RC. 1968. A molecular approach to the study of genic heterozygosity in natural populations. III. Direct evidence of coadaptation in gene arrangements of Drosophila. Proc. Natl. Acad. Sci. USA 59:398-405.

Prakash S, Lewontin RC. 1971. A molecular approach to the study of genic heterozygosity in
natural populations. V. Further direct evidence of coadaptation in inversions of
Drosophila. Genetics:405-408.

Rogers AR. 1995. Genetic evidence for a pleistocene explosion. Evolution 49:608-615.

Rousset F, Kirkpatrick M, Guerrero RF. 2014. Matrix inversions for chromosomal inversions: A
method to construct summary statistics in complex coalescent models. Theor. Pop. Biol. 97:1-10.

Roux C, Fraisse C, J. R, Anciaux Y, Galtier N, Bierne N. 2016. Shedding light on the grey
zone of speciation along a continuum of genomic divergence. PloS Biol. 14:e2000234.

Rozas J, Segarra C, Ribó G, Aguadé M. 1999. Molecular population genetics of the *rp49*
gene region in different chromosomal inversions of *Drosophila subobscura*. Genetics
151:189-202.

Schaeffer SW. 2002. Molecular population genetics of sequence length diversity in the Adh
region of *Drosophila pseudoobscura*. Genet. Res. 80:163-175.

Sheppard PM. 1975. Natural Selection and Heredity. 4th Edition. London: Hutchinson.

Singh RS, Rhomberg LR. 1987. A comprehensive study of genic variation in natural
populations of Drosophila melanogaster. II. Estimates of heterozygosity and patterns of
geographic differentiation. Genetics 117:255-271.

Sprengelmeyer QD, Mansourian S, Lange JD, Matute DR, Cooper BS, Jirle EV, Stensmyr MC, Pool
JE. 2020. Recurrent collection of *Drosophila melanogaster* from wild African environments
and genomic insights into species history. Mol. Biol. Evol. 37:627-638.

Strobeck C. 1983. Expected linkage disequilibrium for a neutral locus linked to a
chromosomal rearrangement. Genetics 103:545-555.

Sturtevant AH. 1926. A crossover reducer in *Drosophila melanogaster* due to inversion of a
section of chromosomes. Biol. Zentralbl. 46:697-702.

Sturtevant AH. 1917. Genetic factors affecting the strength of linkage in *Drosophila*. Proc.
Natl. Acad. Sci. USA 3:555-558.

Sturtevant AH, Beadle GW. 1936. The relations of inversions in the X chromosome of
Drosophila melanogaster to crossing over and disjunction. Genetics 21:554-604.

Takahata N. 1990. A simple genealogical structure of strongly balanced allelic lines and
trans-species evolution of polymorphism. Proc. Natl. Acad. Sci. USA 87:2419-2423.

Toups M, Rodrigues N, Perrin N, Kirkpatrick M. 2019. A reciprocal translocation radically
reshapes sex-linked inheritance in the common frog. Mol. Ecol. 28:1877-1889.

Villoutreix R, Ayala D, Joron M, Gompert Z, Feder JL, Nosil P. 2021. Inversion breakpoints
and the evolution of supergenes. Mol. Ecol. 30:2738–2755.
Wakeley J, Aliacar N. 2001. Gene genealogies in a metapopulation. Genetics 159:893-905.
Wakeley J, Lessard S. 2003. Theory of the effects of population structure and sampling on
patterns of linkage disequilibrium applied to genomic data from humans. Genetics
164:1043-1053.
Weir BS, Cockerham CC. 1984. Estimating *F*-statistics for the analysis of population
structure. Evolution 38:1358-1370.
Wellenreuther M, Bernatchez L. 2018. Eco-evolutionary genomics of chromosomal
inversions. Trnds. Ecol. Evol 33:6.
Wellenreuther M, Mérot C, Berdan EL, Bernatchez L. 2019. Going beyond SNPs: The role of
structural genomic variants in adaptive evolution and species diversification. Mol. Ecol. 28:1203-
1209.
White BJ, Hahn MW, Pombi M, Cassone BJ, Lobo NF, Simard F, Besansky NJ. 2007.
Localization of candidate regions maintaining a common polymorphic inversion (2La)
in *Anopheles gambiae*. PloS Genet. 3:e217.
Wright S. 1951. The genetical structure of populations. Ann. Eugen. 15:323-354.
Wright S, Dobzhansky T, Hovanitz W. 1942. Genetics of natural populations. VII. The
allelism of lethals in the third chromosome of *Drosophila pseudoobscura*. Genetics
27:363-394.
Zeng K, Charlesworth B, Hobolth A. 2021. Studying models of balancing selection using
phase-type theory. Genetics 218:iyab055.

1285

1286

1287 **APPENDIX**

1288 **Recursion relations for mean coalescence times**

1289 By ignoring second-order terms in *m* and *r*, the following recursion relations for the mean

1290 coalescence times defined in the main text can be obtained, with primes indicating their values in

1291 the next generation:

1292

1293 $$t'_{11w} = [1 - 2m - 2ry]\left[\frac{1}{2Nx} + \left(1 - \frac{1}{2Nx}\right)(1 + t_{11w})\right] + 2m(1 + t_{11b}) + 2ry(1 + t_{12w}) \quad \text{(A1a)}$$

1294 $$t'_{11b} = [1 - 2m - 2ry]t_{11b} + \frac{2m}{(d-1)}\left[\frac{1}{2Nx} + \left(1 - \frac{1}{2Nx}\right)(1 + t_{11w})\right]$$

1295
$$+ \frac{2m(d-2)}{(d-1)}(1 + t_{11b}) + 2ry(1 + t_{12b}) \tag{A1b}$$

1296
$$t'_{22w} = [1 - 2m - 2rx]\left[\frac{1}{2Ny} + \left(1 - \frac{1}{2Ny}\right)(1 + t_{22w})\right] + 2m(1 + t_{22b}) + 2rx(1 + t_{12w}) \tag{A1c}$$

1297
$$t'_{22b} = [1 - 2m - 2rx]t_{22b} + \frac{2m}{(d-1)}\left[\frac{1}{2Ny} + \left(1 - \frac{1}{2Ny}\right)(1 + t_{22w})\right]$$

1298
$$+ \frac{2m(d-2)}{(d-1)}(1 + t_{22b}) + 2rx(1 + t_{12b}) \tag{A1d}$$

1299
$$t'_{12w} = [1 - 2m - r][(1 + t_{12w})] + 2m(1 + t_{12b}) + r(1 + xt_{11w} + yt_{22w}) \tag{A1e}$$

1300

1301
$$t'_{12b} = [1 - 2m - r][(1 + t_{12b})] + \frac{2m}{(d-1)}(1 + t_{12w}) + \frac{2m(d-2)}{(d-1)}(1 + t_{12b})$$

1302
$$+ r(1 + xt_{11w} + yt_{22w}) \tag{A1f}$$

1303

1304 These equations can be simplified by neglecting products of $1/Nx$ and $1/Ny$ with m and r, and

1305 rearranging. This yields the simplified recursion relations:

1306

1307
$$t'_{11w} \approx 1 + \left[1 - \frac{1}{2Nx} - 2m - 2ry\right]t_{11w} + 2mt_{11b} + 2ryt_{12w} \tag{A2a}$$

1308
$$t'_{11b} \approx 1 + \frac{2m}{(d-1)}t_{11w} + \left[1 - \frac{2m}{(d-1)} - 2ry\right]t_{11b} + 2ryt_{12b} \tag{A2b}$$

1309
$$t'_{22w} \approx 1 + \left[1 - \frac{1}{2Ny} - 2m - 2rx\right]t_{22w} + 2mt_{22b} + 2rxt_{12w} \tag{A2c}$$

1310
$$t'_{22b} \approx 1 + \frac{2m}{(d-1)}t_{22w} + \left[1 - \frac{2m}{(d-1)} - 2rx\right]t_{22b} + 2rxt_{12b} \tag{A2d}$$

1311
$$t'_{12w} = 1 + rxt_{11w} + ryt_{22w} + [1 - 2m - r]t_{12w} + 2mt_{12b} \tag{A2e}$$

1312
$$t'_{12b} = 1 + rxt_{11b} + ryt_{22b} + \frac{2m}{(d-1)}t_{12w} + \left[1 - \frac{2m}{(d-1)} - r\right]t_{12b} \tag{A2f}$$

1313

1314 **Expressions for equilibrium mean coalescence times**

1315 Writing $M = 4Nm$ and $R = 4Nr$, we obtain the following expressions for equilibrium,

1316 equating the t's and ts:

1317

1318
$$t_{11w}(Mx + Ry + 1) = 2Nx + Mxt_{11b} + Rxyt_{12w} \tag{A3a}$$

1319
$$t_{11b}[2m + 2(d-1)ry] = d - 1 + 2mt_{11w} + 2(d-1)ryt_{12b} \tag{A3b}$$

1320
$$t_{22w}(My + Rx + 1) = 2Nx + Myt_{22b} + Rxyt_{12w} \tag{A3c}$$

1321
$$t_{22b}[2m + 2(d-1)rx] = d - 1 + 2mt_{22w} + 2(d-1)rxt_{12b} \tag{A3d}$$

1322
$$t_{12w}(2m + r) = 1 + 2mt_{12b} + r(xt_{11w} + yt_{22w}) \tag{A3e}$$

1323
$$t_{12b}[2m + (d-1)r] = d - 1 + 2mt_{12w} + (d-1)r(xt_{11b} + yt_{22b}) \tag{A3f}$$

1324

1325    This is a set of 6 linear equations that can be solved exactly (see the Supplementary

1326 Appendix, section S1). . However, it is possible to obtain informative approximate expressions by

1327 assuming that d is very large, with $2m \gg r$ but $dr \gg 2m$, which are plausible conditions for many

1328 real-life cases. This means that the terms in R or r on the left-hand sides of Equations (A3a), (A3c)

1329 and (A3e) can be neglected, as well as the terms in R or r on the right-hand sides of Equations

1330 (A3a), (A3c) and (A3e); in addition, $d-1$ can be replaced by d with neglible loss of accuracy.

1331    Using these approximations, we have:

1332

1333
$$t_{12w} \approx (2m)^{-1} + t_{12b} \qquad\qquad (A4a)$$

1334

1335 The intuitive interpretation of this expression is that, if recombination is infrequent compared with

1336 migration, a class 1/2 pair of alleles will wait an average of $1/(2m)$ generations before one of them

1337 migrates to a different deme, after which their expected coalescent time is t_{12b}. The various t values

1338 are of the order of $2N_T$ (see below), so that for large d and moderate F_{ST} values, $1/m$ is small

1339 compared with the t's, and the term in m in this equation can be neglected.

1340    Equations (A3a) and (A3c) can then be approximated as:

1341

1342
$$t_{11w}(Mx + 1) \approx 2Nx + Mxt_{11b} + 4Nxy[1 + r(xt_{11b} + yt_{22b})])$$

1343
$$\approx 2Nx(1 + 2y) + Mxt_{11b} \qquad\qquad (A4b)$$

1344
$$t_{22w}(My + 1) \approx 2Ny + Myt_{22b} + 4Nxy[1 + r(xt_{11b} + yt_{22b})])$$

1345
$$\approx 2Ny(1 + 2x) + Myt_{22b} \qquad\qquad (A4c)$$

1346

1347 From Equations (A3c) and (A4a) we obtain:

1348

1349
$$t_{11b}[2m + 2dry] \approx d + 2mt_{11w} + 2dry(r^{-1} + xt_{11b} + yt_{22b})$$

1350 so that:

1351
$$t_{11b}[2m + 2dry^2] \approx d(1 + 2y) + 2mt_{11w} + 2dry^2t_{22b}$$

1352

1353 Similarly,

1354
$$t_{22b}[2m + 2dryx^2] \approx d(1 + 2yx) + 2mt_{22w} + 2dryx^2t_{11b}$$

1355

1356 Using Equations (A4b) and (A4c), after some algebra we obtain:

1357

1358 $$t_{11b}[2m + 2dry^2(Mx + 1)] \approx d(1 + 2y)(Mx + 1) + 2dry^2(Mx + 1)t_{22b} \quad \text{(A4d)}$$

1359 $$t_{22b}[2m + 2drx^2(My + 1)] \approx d(1 + 2x)(My + 1) + 2drx^2(My + 1)t_{11b} \quad \text{(A4e)}$$

1360

For further work, it is convenient to scale the t's by division by the expected coalescent time

for a pair of neutral alleles sampled from the same deme, $T_{Sn} = 2N_T = 2dN$. Times on this scale are

denoted by upper-case T's. The corresponding scaled recombination rate is denoted by $\rho = 4N_T r$,

which replaces $2r$ in Equations (A4d) and (A4e) when using the rescaled t's; $2m$ is also replaced by

dM, and d can be cancelled from both sides of these equations, which then give a pair of linear

equations in T_{11b} and T_{22b}. After some further algebra, their solution can be written in the standard

form for a pair of linear equations as:

1368

1369 $$T_{11b} \approx (a_{22}b_1 - a_{12}b_2)/det, \quad T_{22b} \approx (a_{11}b_2 - a_{21}b_1)/det \quad \text{(A5a)}$$

1370 $$det = (a_{11}a_{22} - a_{12}a_{21}) \quad \text{(A5b)}$$

1371 $$a_{11} = M(Mx + 1)^{-1} + \rho y^2, a_{12} = -\rho y^2, a_{21} = -\rho x^2, a_{22} = M(My + 1)^{-1} + \rho x^2 \quad \text{(A5c)}$$

1372 $$b_1 = (1 + 2y), \quad b_2 = (1 + 2x) \quad \text{(A5d)}$$

1373

1374 Use of the expressions for the a_{ij} yields the following formulae:

1375

1376 $$det = M\{M(Mx + 1)^{-1} + M(My + 1)^{-1} + \rho[x^2(Mx + 1)^{-1} + y^2(My + 1)^{-1}]\} \quad \text{(A6a)}$$

1377 $$T_{11b} \approx [(1 + 2y)M(My + 1)^{-1} + \rho]/det \quad \text{(A6b)}$$

1378 $$T_{22b} \approx [(1 + 2x)M(Mx + 1)^{-1} + \rho]/det \quad \text{(A6c)}$$

1379

1380 These expressions can be substituted into the scaled versions of Equations (A4d)

1381 – (A4e) to obtain the complete approximate solution to the set of equations for the T's:

1382

1383 $$T_{12b} \approx 2\rho^{-1} + xT_{11b} + yT_{22b} \quad \text{(A6d)}$$

1384 $$T_{12w} \approx T_{12b} \quad \text{(A6e)}$$

1385 $$T_{11w}(Mx + 1) \approx d^{-1}x(1 + 2y) + MxT_{11b} \quad \text{(A6f)}$$

1386 $$T_{22w}(My + 1) \approx d^{-1}y(1 + 2x) + MyT_{22b} \quad \text{(A6g)}$$

1387

1388   If $d \gg 1$, as is often likely to be the case, the terms in d in Equations (A6f) and (A6g) can

1389 be neglected, so that:

1390
$$T_{11w} \approx Mx(1 + Mx)^{-1}T_{11b} \qquad \text{(A6h)}$$

1391

1392
$$T_{22w} \approx My(1 + My)^{-1}T_{22b} \qquad \text{(A6i)}$$

1393

1394   For calculating the approach to the equilibrium described by Equations (A6), it is convenient

1395 to eliminate the terms in 1 on the right-hand sides of Equations (A2) by using the vector of

1396 deviations of the T's from their equilibrium values, and applying a matrix A that contains the

1397 multipliers of the T's in Equations (A2). Details are given in the Supplementary Appendix .

1398

1399

1400

1401

1402 **Table 1 Comparisons of within-deme F_{AT} for a balanced polymorphism versus σ_d^2 for a pair**

1403   **of neutral loci**

1404 $\rho = 0.4$ $\rho = 4$ $\rho = 40$

F_{ST}	F_{AT} $x=0.1$	F_{AT} $x=0.5$	σ_d^2	F_{AT} $x=0.1$	F_{AT} $x=0.5$	σ_d^2	F_{AT} $x=0.1$	F_{AT} $x=0.5$	σ_d^2
0	0.489	0.714	0.380	0.117	0.200	0.156	0.021	0.024	0.023
0.05	0.492	0.718	0.419	0.130	0.232	0.200	0.051	0.072	0.070
0.10	0.496	0.723	0.454	0.143	0.265	0.241	0.073	0.120	0.114
0.15	0.500	0.728	0.488	0.157	0.299	0.281	0.092	0.167	0.156
0.20	0.504	0.733	0.521	0.171	0.333	0.320	0.110	0.216	0.197
0.25	0.509	0.739	0.552	0.186	0.368	0.358	0.128	0.263	0.238

1405

1406 The scaled recombination rates of $\rho = 0.4$, 4 and 40 correspond to r = 10^{-7}, 10^{-6} and 10^{-5},

1407 respectively, when $N_T = 10^6$, as assumed in the figures.

1408

1409

1410

1411

Figure 1. The equilibrium population statistics for an inversion polymorphism where the

inversion is maintained at a constant frequency of 0.1, for four different recombination

rates in heterokaryotypes (r), indicated at the top left of each panel. An island model of

population structure is assumed, with 200 demes and a total population size of $N_T = 10^6$, so

that individual demes have a population size of $N = 5000$. The X axis is the equilibrium F_{ST} for

neutral sites unlinked to the inversion; the product of $4N$ and the migration rate m is

approximately equal to $F_{ST}/(1 - F_{ST})$. The dashed curves are variables that fall between 0 and

1 (left-hand Y axis); black is the ratio T_{11w}/T_{22w}, red is the ratio T_{11b}/T_{22b}, brown is F_{ATw} and

blue is F_{ATb}. The solid curves are mean coalescent times, measured relative to $2N_T$ (right-

hand Y axis); red is T_{12}, black is T_{Tw} and blue is T_{22w}. Subscripts w and b indicates statistics for

alleles sampled within the same deme and between different demes, respectively. (For T_{12},

there is no significant difference between with- and between-deme values.) Subscripts 1

and 2 denote alleles sampled from the inversion and standard arrangement, respectively;

subscript T denotes pairs of alleles sampled without regard to karyotype. F_{AT} is the

standardised measure of divergence between the inverted and standard arrangements.

1427

1428

Figure 2. The equilibrium population statistics for an inversion polymorphism where the

inversion is maintained at a constant frequency of 0.5, under the same assumptions as in

Figure 1. In this case, $T_{11w} = T_{22w} = 1$ and $T_{11b}/T_{22b} = 1$ for all values of F_{ST} and are not

displayed. The dashed curves are variables that fall between 0 and 1 (left-hand Y axis); black

is F_{ATw} for $r = 10^{-7}$ and blue is F_{ATw} for $r = 10^{-6}$; brown is F_{ATb} for $r = 10^{-7}$ and red is F_{ATb} for $r =$

10^{-6}. The solid curves are mean coalescent times, measured relative to $2N_T$ (right-hand Y

axis); red is T_{12} for $r = 10^{-7}$ and black is T_{12} for $r = 10^{-6}$; blue is T_{Tw} for $r = 10^{-7}$ and brown is T_{Tw}

for $r = 10^{-6}$. The right-hand panel shows the results for $r = 10^{-5}$ and 10^{-4}, with colours and

line characteristics corresponding to those in the left-hand panel for $r = 10^{-7}$ and 10^{-6},

respectively.

1439

1440

1441

1442

1443

1444

1445

1446

Figure 3. The trajectories of change in the population statistics for the case of a panmictic

population of size $N = 10^6$, assuming that the time taken to approach the equilibrium

inversion frequency is negligible compared with the coalescent time of $2N$ generations. The

X axes display times in units of coalescent time following the sweep to equilibrium. Three

different recombination rates in heterokaryotypes are shown, as well as two different

frequencies of the inversion (0.1 in the upper panels and 0.5 in the lower panels). The

dashed curves are F_{AT}, whose values are displayed on the left-hand Y axes. The solid curves

are mean coalescent times, measured relative to $2N$; red is T_{12}, brown is T_T, black is T_{11} and

blue is T_{22} (right-hand Y axes). For the highest rate of recombination ($r = 10^{-5}$), only the first

N generations are shown, in order to capture the rapid changes at the start of the process.

The coloured bars inside the Y axes indicate the equilibrium values of the corresponding

statistics, for cases when these are substantially different from the final values of the

statistics.

1460

1461

1462

1463

1464

Figure 4. The trajectories of change in the population statistics for the case of a subdivided

(island model) population of total size $N_T = 10^6$, with 200 demes of size $N = 500$ and with an

F_{ST} for neutral sites independent of the inversion of 0.05 (i.e., $4Nm = 19$). It is assumed that

the time taken to approach the equilibrium inversion frequency is negligible compared with

the coalescent time of $2N_T$ generations. The X axes display times in units of coalescent time

following the sweep to equilibrium. Three different recombination rates in

heterokaryotypes are shown, as well as two different frequencies of the inversion (0.1 in the

upper panels and 0.5 in the lower panels). The solid curves represent within-population

statistics and the dashed curves are between-population statistics. The values of F_{ATw} and

F_{ATb} (brown curves) are given by the left-hand Y axes. The other curves are mean coalescent

times, measured relative to $2N_T$; red is T_{12w} (T_{12b} behaves almost identically, except for its

higher initial value and slower rate of increase when $T < 0.005$); black is T_{11} and blue is T_{22}

(right-hand Y axes). For the highest rate of recombination ($r = 10^{-5}$), only the first N_T

generations are shown. The coloured bars inside the Y axes indicate the equilibrium values

of the corresponding statistics, for cases when these are substantially different from the

final values of the statistics.

1481

1482

1483

1484   **Figure 5.** This is the same as Figure 4, except that F_{ST} = 0.15 (i.e., $4Nm$ = 5.67).

1485

Supplementary File S1- Appendix

S1 The exact equilibrium solution with population subdivision

The full solution for the equilibrium T's can be found from Equations (A3) as follows, considering each row in turn and after dividing both sides by $2N_T$.

$$T_{11w} = c_0 + c_1 T_{11b} + c_2 T_{12w}, c_0 = \frac{x}{dg_1}, \; c_1 = \frac{Mx}{g_1}, c_2 = \frac{Rxy}{g_1}, \; g_1 = Mx + Ry + 1 \quad \text{(S1a)}$$

$$T_{11b} = c_3 + c_4 T_{12w} + c_5 T_{12b}, c_3 = \frac{[(1-d^{-1})+c_0 M]}{g_2}, c_4 = \frac{c_2 M}{g_2}, c_2 = \frac{(\rho - R)y}{g_1},$$

$$g_2 = M(1 - c_1) + (\rho - R)y \quad \text{(S1b)}$$

$$T_{22w} = c_6 + c_7 T_{22b} + c_8 T_{12w}, c_6 = \frac{y}{dg_1}, c_7 = \frac{Mx}{g_1}, c_8 = \frac{Rxy}{g_3}, g_3 = My + Rx + 1 \quad \text{(S1c)}$$

$$T_{22b} = c_9 + c_{10} T_{12w} + c_{11} T_{12b}, c_9 = \frac{[(d-1)d^{-1}+c_6 M]}{g_4}, c_{10} = \frac{c_8 M}{g_4}, c_{11} = \frac{(\rho - R)x}{g_4},$$

$$g_4 = M(1 - c_7) + (\rho - R)x \quad \text{(S1d)}$$

These equations can be used to eliminate all the T's other than T_{12w} and T_{12b} from Equations (A1e) and (A1f), yielding a pair of linear equations:

$$b_{11} T_{12w} + b_{12} T_{12b} = e_1, \quad b_{21} T_{12w} + b_{22} T_{12b} = e_2 \quad \text{(S1e)}$$

where

$$e_1 = d^{-1} + \tfrac{1}{2} R[x(c_0 + c_1 c_3) + y(c_6 + c_2 c_9)], \; e_2 = 1 - d^{-1} + \tfrac{1}{2}\rho(xc_3 + yc_9)$$

$$b_{11} = M + \tfrac{1}{2} R[1 - x(c_2 + c_1 c_4) - y(c_8 + c_7 c_{11})], \; b_{12} = -\tfrac{1}{2} R[xc_1 c_5 + yc_7 c_{11}]$$

$$b_{21} = -[M + \tfrac{1}{2}\rho(xc_4 + yc_{10})], b_{22} = M + \tfrac{1}{2}\rho(1 - xc_5 - yc_{11})$$

This set of equations has the following standard form solution:

$$T_{12w} \approx (b_{22} e_1 - b_{12} e_2)/det, \quad T_{12b} \approx (b_{11} e_2 - b_{21} e_1)/det \quad \text{(S1f)}$$

$$det = (b_{11} b_{22} - b_{12} b_{21}) \quad \text{(S1g)}$$

S2 The approach to the approximate equilibrium solutions

Let the column vector $\boldsymbol{x} = (\delta T_{11w}, \delta T_{11b}, \delta T_{22w}, \delta T_{22b}, \delta T_{12w}, \delta T_{12b})^T$, where δ denotes the deviation from the equilibrium value of the element of the vector in question. Let $N_1 = 2Nx$, and $N_2 = 2Ny$.

1531 With large d, the 6 x 6 matrix A that describes the approach of x to a vector of zero elements, by

1532 iterating $x_n = Ax_{n-1}$, consists of the following elements:

$1-2m-2ry-1/N_1$	$2m$	0	0	$2ry$	0
$2m/d$	$1-2m/d-2ry$	0	0	0	$2ry$
0	0	$1-2m-2rx-1/N_2$	$2m$	$2rx$	0
0	0	$2m/d$	$1-2m/d-2rx$	0	$2rx$
rx	0	ry	0	$1-2m-r$	$2m$
0	rx	0	ry	$2m/d$	$1-2m/d-r$

S3 Approximations for the eigenvalues and eigenvectors of A

An approximation for the set of eigenvalues λ_i ($i = 1 - 6$) of A can be obtained as follows, by noting

that all its elements a_{ij} other than the diagonals a_{ii} are zero or of the order of m or r, so that second-

order terms in non-diagonal elements can be neglected. Ignoring the zero elements in the first row

of A, the characteristic equation of A, $\det(A - \lambda I)$, can be expanded as $(a_{11} - \lambda)A_{11} - a_{12}A_{12} + a_{15}A_{15}$,

where A_{ij} denotes the minor of element a_{ij}.

Similarly, $A_{11} = (a_{22} - \lambda)B_{22} - a_{34}B_{34} + a_{35}B_{35}$, where B_{ij} is the minor of a_{ij} in the submatrix

corresponding to the determinant A_{11}. Writing $O(\varepsilon)$ for terms of order m or r, and repeating this

procedure on the B_{ij}, it is easily seen that $B_{22} = (a_{22} - \lambda)(a_{33} - \lambda)(a_{44} - \lambda)(a_{55} - \lambda)(a_{66} - \lambda)$, $B_{34} =$

$O(\varepsilon)$, and $B_{35} = O(\varepsilon^2)$. Since $a_{34} = O(\varepsilon)$, the only significant term in $(a_{11} - \lambda)A_{11}$ is thus $(a_{11} - $

$\lambda)(a_{22} - \lambda)(a_{33} - \lambda)(a_{44} - \lambda)(a_{55} - \lambda)(a_{66} - \lambda)$.

We also have $A_{12} = a_{21}C_{21} - a_{26}C_{26}$, where C_{ij} is the minor of a_{ij} in the submatrix

corresponding to the determinant A_{34}. Since both a_{21} and a_{26} are $O(\varepsilon)$, their product with a_{12} can

be neglected. Similarly, A_{15} can be expanded as $a_{21}D_{21} - a_{22}D_{22} + a_{26}D_{26}$; the products of $a_{21}D_{21}$

and $a_{26}D_{22}$ with a_{15} can thus be neglected. D_{22} can in turn be expanded as $a_{33}E_{33} + a_{34}E_{34}$

where $E_{33} = 0$ and $E_{34} = O(\varepsilon)$.

The final result, therefore, is that $\det(A - \lambda I) = \Pi_i (a_{ii} - \lambda) + O(\varepsilon^2)$. This means that the six

eigenvalues of A can be equated to the corresponding diagonal elements, which are of order $1 - $

$O(\varepsilon)$.

This in turn allows approximations for the right and left eigenvectors of A, and hence its

spectral expansion, to be determined, giving a complete (but approximate) analytic solution to

the trajectory of deviations of the t's from the equilibrium values. The relevant components of

these vectors can be obtained from the equations $Az_i = \lambda_i z_i$ and $y_i A = \lambda_i y_i$, where z and y refer to

right and left eigenvectors, respectively, by successive elimination of elements of the vectors.

Using a similar procedure to that leading to Equation (S1), by considering only the non-

zero elements of A, examining each row A of in succession and writing $\lambda = 1 - \alpha$ for a given λ,

we have:

1574
$$z_1 = k_{12}z_2 + k_{15}z_2, \quad k_{12} = \frac{a_{12}}{1-\alpha-a_{11}}, \quad k_{15} = \frac{a_{15}}{1-\alpha-a_{11}} \tag{S2a}$$

1575

1576
$$z_2 = k_{25}z_5 + k_{26}z_6, \quad k_{25} = \frac{a_{21}k_{15}}{1-\alpha-a_{22}-a_{21}k_{12}}, \quad k_{26} = \frac{k_{26}}{1-\alpha-a_{22}-a_{21}k_{12}} \tag{S2b}$$

1577

1578
$$z_3 = k_{34}z_4 + k_{35}z_5, \quad k_{34} = \frac{a_{34}}{1-\alpha-a_{33}}, \quad k_{35} = \frac{a_{35}}{1-\alpha-a_{33}} \tag{S2c}$$

1579

1580
$$z_4 = k_{45}z_5 + k_{46}z_{56}, \quad k_{45} = \frac{a_{43}k_{35}}{1-\alpha-a_{44}-a_{43}k_{34}}, \quad k_{46} = \frac{a_{46}}{1-\alpha-a_{44}-a_{43}k_{34}} \tag{S2d}$$

1581

1582
$$z_5 = k_{56}z_6, \quad k_{56} = \frac{1-\alpha-a_{66}-a_{62}k_{26}-a_{64}k_{46}}{a_{62}k_{25}+a_{64}k_{45}+a_{65}} \tag{S2e}$$

1583

All elements of z can thus be evaluated explicitly as ratios with respect to z_6. The same

procedure can be carried out for the components of the left eigenvector, y; the coefficients

corresponding to the k_{ij} above are identical in form, except that the elements of the transpose of A

are used. Constructing the matrices Z and Y, whose columns and rows respectively are given by

the sets of column vectors z_i and row vectors y_i corresponding to each eigenvalue, we have the

standard result $A^n = Z\Lambda^n Y$, where Λ is the diagonal matrix whose non-zero elements are equal

to the eigenvalues of A, and the eigenvectors are normalized such that the product $y_i z_i = 1$.

Similar results can be obtained for the panmictic case, using rows and columns 1, 3 and 5

of A, with $m = 0$.

1593

1594   **S4 Approximations for the initial per generation changes in the F_{AT}'s**

1595   The first-order approximation for ΔF_{ATw} is as follows:

1596

1597
$$\Delta F_{ATw} \approx \Delta T_{11w}\partial_{T_{11w}}F_{ATw} + \Delta T_{22w}\partial_{T_{22w}}F_{ATw} + \Delta T_{12w}\partial_{T_{12w}}F_{ATw} \tag{S3a}$$

1598   Here, the derivatives are evaluated at the initial conditions: $T_{22w} = T_{22b} = 0$, $T_{12w} = T_{22w} = 1$,

1599   and $T_{22b} = T_{12b} \approx 1/(1 - F_{ST})$. Using the relation $F_{ATw} = 1 - T_{Sw}/T_{Tw}$, and the definitions of T_{Sw} and

1600 T_{Tw} given in the main text (Equations 1 - 4), we obtain the following expressions:

1601

1602 $\partial_{T_{11w}} T_{Sw} = x,\ \partial_{T_{22w}} T_{Sw} = y,\ \partial_{T_{12w}} T_{Sw} = 0;\ \partial_{T_{11w}} T_{Tw} = y^2;\ \partial_{T_{22w}} T_{Tw} = yx^2;\ \partial_{T_{12w}} T_{Tw} = 2xy$

1603   (S3b)

1604

Substituting these relations into Equations (6) and (S3), and using the fact that the derivative of

$F_{ATw} = 1 - T_{Sw}/T_{Tw}$ with respect to an arbitrary variable u is $\frac{T_S \partial_u T_{Tw} - T_{Tw} \partial_u T_{Sw}}{T_T^2}$, yields Equation

(7a), after some simplification.

A similar procedure can be used for F_{ATb} noting that terms involving the derivative of T_{22b}

can be ignored, because $\Delta T_{22b} \approx 0$ by Equation (6d). It is useful to note that the initial value of

T_{Tb} can be written as $(1 - F_{ST})^{-1} y[1 + x(1 - F_{ST})]$. We have:

1611

1612 $\partial_{T_{11b}} T_{Sb} = x,\ \partial_{T_{11b}} T_{Sb} = y;\ \partial_{T_{11b}} T_{Tb} = y^2;\ \partial_{T_{12b}} T_{Tb} = 2xy(1 - F_{ST})^{-1}$ (S3c)

1613

1614   This yields Equation (7b).

1615

1616   **S5 Inaccuracy in the Approximate Expressions for Equilibrium**

1617   Denote the true equilibrium vector of T's by $\widetilde{x} = x^* - \delta$, where x^* is the approximate value. Let

1618   the true value of x in generation n be x_n, where $n = 0$ for the initial generation under

1619   consideration. Let A be the true value of the recursion matrix discussed above. The true

1620   equilibrium satisfies $A\widetilde{x} + I = \widetilde{x}$, where I is the unit matrix. For generation 1, we can write:

1621

1622 $x_1 - x^* = x_1 - \widetilde{x} + \delta = Ax_0 + I - (A\widetilde{x} + I) - \delta = A[x_0 - (\widetilde{x} + \delta)] + \delta(A - I)$ (S4a)

1623

1624   In the next generation,

1625

1626 $x_2 - x^* = Ax_1 + I - (A\widetilde{x} + I) - \delta = A\{A[x_0 - (\widetilde{x} + \delta)] + \delta(A - I)\} + \delta(A - I) =$

1627 $= A^2[x_0 - (\widetilde{x} + \delta)] + (A + I)\delta(A - I)$ (S4b)

1628

1629   Continuing with the iteration, the general expression is:

$$x_n - x^* = A^n[x_0 - (\tilde{x} + \delta)] + (A^{n-1} + A^{n-2} + \cdots + I)\delta(A - I) \qquad \text{(S4c)}$$

From the analysis in section S3, the term in A^n must tend to zero as n increases, at least as far as first-order terms are concerned, so only the series in A need be considered. If we write P_i for the matrix formed by the vector product $z_i y_i$, such that $A^j = \sum_i \lambda_i^j P_i$, for large n this term approaches $\sum_i \frac{1}{(1-\lambda_i)} P_i$, provided that $|\lambda_i| < 1$, as must be the case from the results in section 3. This implies that the iteration of $x - x^*$ by the matrix A, as used in the numerical studies of the approach to equilibrium will yield a final value whose magnitude is of the same order as the error δ in the approximate equilibrium. If this value is small, the error involved in the approximation must also be small.

S6 Time courses of changes in coalescent times in the absence of recombination in a panmictic population

In this case, there is no distinction between within- and between-deme coalescence times, and the subscripts w and b can be dropped. The mean between-karyotype coalescent time after a time T has elapsed (where both quantities are scaled relative to the neutral coalescence time $2N_T$), is simply a linear function of T:

$$T_{12}(T) = T_{12}(0) + T \qquad \text{(S5a)}$$

The mean coalescence times within-karyotypes obey the rule for the approach to their equilibrium values described in section S2 above. In this case, the relevant expressions simplify to:

$$T_{11}(T) = x + [T_{11}(0) - x]\exp(-T/x) \qquad \text{(S5b)}$$

$$T_{22}(T) = y + [T_{22}(0) - y]\exp(-T/y) \qquad \text{(S5c)}$$

If the sweep of the inversion to its equilibrium frequency is assumed to be instantaneous, we have $T_{11}(0) = 0$ and $T_{12}(0) = T_{22}(0) = 1$; we also have $T_T(0) = 2xy + y^2 = y(1 + x)$, so that $F_{AT}(0) = x/(1 + x)$. For arbitrary T, $T_S(T) = x^2[1 - \exp(-T/x)] + y[y + x\exp(-T/y)]$ and $T_T(T) = 2xy(1 + T) + x^2[1 - \exp(-T/x)] + y^2[y + x\exp(-T/y)]$.

1661   These relations can be used to determine the time-course of F_{AT} following an

1662   instantaneous sweep, using the relation:

1663

1664
$$F_{AT}(T) \approx 1 - \frac{T_S(T)}{T_T(T)} \qquad \text{(S5d)}$$

1665

1666   For sufficiently large T, the exponential terms in the above equations can be neglected,

1667   and $T_S(T)/ T_T(T) \approx (1 - 2xy)/(2xyT)$. This implies that, as expected, F_{AT} always increases towards

1668   1 once a large T value has been reached. For small T, the first-order approximations $\exp(-z) \approx 1$

1669 $- z$ and $1/(1 + T) \approx 1 - T$ yield the following formulae:

1670

1671
$$T_{11}(T) \approx T \qquad \text{(S5e)}$$

1672
$$T_{22}(T) \approx 1 - (x/y)T \qquad \text{(S5f)}$$

1673
$$T_S(T) \approx y \qquad \text{(S5g)}$$

1674
$$T_T(T) \approx y(1 + x) + xT \qquad \text{(S5h)}$$

1675

1676   Substituting these approximations into Equation (S5d) and simplifying, we have:

1677

1678
$$F_{AT}(T) \approx \frac{x}{(1+x)}[1 + \frac{x}{y(1+x)}T] \qquad \text{(S5i)}$$

1679

1680   The multiplicand of T is always positive for $0 < x < 0$, implying that F_{AT} always increases

1681   initially with time.

1682

1683   **S7 Time courses of changes in coalescent times in the absence of recombination in a**

1684   **subdivided population**

1685   With no recombination, the recursion relations for T_{12w} and T_{12b}, which can be obtained from

1686   Equations (A3) reduce to:

1687
$$X(t) = c + A\,X(t - 1) \qquad \text{(S6a)}$$

1688

1689   where $X(t)$ is the column vector with elements $T_{12w}(t)$ and $T_{12b}(t)$; c is the column vector whose

1690   elements are both equal to $1/(2N_T)$; A is the 2 x 2 matrix with elements $a_{11} = 1 - 2m$, $a_{12} = 1 - 2m$,

1691 $a_{21} = 2m/(d - 1)$ and $a_{22} = 1 - 2m/(d - 1)$.

1692   Iteration of this equation yields the following expression:

1693

$$X(t) = A^t X(0) + \sum_{i=1}^{t} A^{t-i} c \qquad (S6b)$$

1695

1696   The powers of A can be evaluated using the spectral expansion of A in terms of its

1697   eigenvalues and eigenvectors. The eigenvalues of A satisfy the following characteristic equation:

1698

$$\lambda^2 - \lambda[2 - 2m(1 + d'^{-1})] + 1 - 2m(1 + d'^{-1}) \qquad (S7)$$

1700

where for convenience $d - 1$ is denoted by d'.

This equation has roots $\lambda_0 = 1$ and $\lambda_1 = 1 - 2m(1 + d'^{-1})$, with corresponding right and

left eigenvectors u_i and v_i, respectively ($i = 0, 1$). Use of the identities $Au_i = \lambda_i u_i$ and $v_i A = \lambda_i v_i$

give the result $u_{01} = u_{02}$ for the 1st and 2nd elements of u_0, respectively. Similarly, $u_{11} = (-d') u_{12}$,

$v_{01} = v_{02}/d$ and $v_{11} = -v_{12}$. In order to use these in the spectral expansion of A, we arbitrarily set

$u_{01} = u_{02} = 1$ and $u_{11} = 1$, with $u_{12} = -d'^{-1}$, and use the normalization requirement $u_i^{Tr} v_i = 1$,

where the superscript Tr denotes transposing a column vector into a row vector. This gives $v_{01} =$

$(1 + d')^{-1}$, $v_{02} = v_{11} = (1 + d'^{-1})^{-1}$, $v_{12} = -(1 + d'^{-1})^{-1}$. Using the formula for the spectral

decomposition of a matrix, $A = \sum_i \lambda_i u_i v_i$, we obtain:

1710

1711 $$A^t = \begin{pmatrix} (1+d')^{-1} & (1+d'^{-1})^{-1} \\ (1+d')^{-1} & (1+d'^{-1})^{-1} \end{pmatrix} + [1 - 2m(1+d'^{-1})]^t \begin{pmatrix} (1+d'^{-1})^{-1} & -(1+d'^{-1})^{-1} \\ -d'^{-1}(1+d'^{-1})^{-1} & d'^{-1}(1+d'^{-1})^{-1} \end{pmatrix} \quad (S8a)$$

1712

1713   For large d, this expression reduces to:

1714

1715 $$A^t \approx \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix} + [1 - 2m]^t \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} (1-2m)^t & 1 - (1-2m)^t \\ 0 & 1 \end{pmatrix} \qquad (S8b)$$

1716
1717

1718   This equation implies that:

1719 $$\sum_{i=1}^{t} A^{t-i} \approx \begin{pmatrix} \frac{1-(1-2m)^t}{2m} & t - \frac{1-(1-2m)^t}{2m} \\ 0 & t \end{pmatrix} \qquad (S8c)$$

1720

1721   If time is transformed to the coalescent timescale, $T = t/(2N_T)$, $(1 - 2m)^t$ can be

1722   approximated by $\exp(-MdT)$. On the assumption of an instantaneous approach of the inversion

1723   to its equilibrium frequency, so that the initial vector $X(0) = (1, (1 - F_{ST})^{-1})^{Tr} = (1, (1+M)/M)^{Tr}$

1724   Equations (S6b), (S8b) and (S8c) yield the following expression:

1725

$$X(T) \approx \begin{pmatrix} \exp(-dMT) & 1-\exp(-dMT) \\ 0 & 1 \end{pmatrix} (1, \ (1-F_{ST})^{-1})^{Tr} + \begin{pmatrix} \frac{[1-\exp(-dMT)]}{2m} & (2N_T T) - \frac{[1-\exp(-dMT)]}{2m} \\ 0 & 2N_T T \end{pmatrix} c \quad (S9a)$$

1726

1727

1728 which reduces to:

1729

$$X(T) \approx \exp^{-dMT} + (1-\exp^{-dMT})(1-F_{ST})^{-1} + T, \ (1-F_{ST})^{-1} + T \ )^{Tr} \quad (S9b)$$

1730

1731

1732 When $dMT \gg 1$, this expression implies that both T_{12w} and $T_{12b} \approx 1/(1-F_{ST}) + T \approx$

1733 $(1+M)/M + T$. For $dMT \ll 1$, T_{12w} increases faster with T than does T_{12b}, which always increases

1734 in direct proportion to T, especially when M is small. This can be seen by using the first-order

1735 approximation $\exp(-z) \approx 1 - z$ in Equation (S9b), which yields:

1736

$$T_{12w} \approx 1 - MT + dMT(1-F_{ST})^{-1} + T = 1 + (d+1)T \quad (S9c)$$

1737

$$T_{12b} \approx \frac{(1+M)}{M} + T \quad (S9d)$$

1738

1739

1740 The deviations of $T_{11w}(t)$ and $T_{11b}(t)$ from their equilibrium values, $T_{11w} = x$ and $T_{11b} =$

1741 $x/(1-F_{ST, In})$, where $F_{ST, In} =$ is the value of F_{ST} for the inversion subpopulation are described by

1742 the following matrix equation:

1743

$$Y(t) = B^t \ Y(0) \quad (S10a)$$

1744

1745 where $Y(t)$ is the column vector $(\delta T_{11w}(t), \ \delta T_{11b}(t))^{Tr}$. We have:

1746

$$B = \begin{pmatrix} 1 - \frac{(1+Mx)}{2Nx} & 2m \\ \frac{2m}{d'} & 1 - \frac{2m}{d'} \end{pmatrix} \quad (S10b)$$

1747

1748

1749 The characteristic equation of B can be written as:

1750

$$\lambda^2 - \lambda \left[2 - \frac{(1+Mx)}{2Nx} - 2md'^{-1} \right] + 1 + \frac{m}{Nd'x} - \frac{(1+Mx)}{2Nx} - 2md'^{-1} = 0 \quad (S11)$$

1751

1752

1753 This equation has the following roots:

1754
$$\lambda_0 = \frac{1}{2}\{2 - \frac{(1+Mx)}{2Nx} - 2md'^{-1} + \sqrt{\left[\frac{(1+Mx)}{2Nx} + \frac{2m}{d'}\right]^2 - \frac{4m}{Nd'x}}\}$$
(S12a)

1755
$$\lambda_1 = \frac{1}{2}\{2 - \frac{(1+Mx)}{2Nx} - 2md'^{-1} - \sqrt{\left[\frac{(1+Mx)}{2Nx} + \frac{2m}{d'}\right]^2 - \frac{4m}{Nd'x}}\}$$
(S12b)

1756

1757   The quantity inside the square term can be written as:

1758
$$\left[\frac{(1+Mx)}{2Nx} + \frac{2m}{d'}\right]^2 (1 - \varepsilon)$$
(S13a)

1759

1760   where
$$\varepsilon = \frac{4m}{Nd'x}\left[\frac{(1+Mx)}{2Nx} + \frac{2m}{d'}\right]^{-2}$$
(S13b)

1761

1762   Unless Nx is close to 1, $\varepsilon \ll 1$, and $\sqrt{1-\varepsilon} \approx 1 - \frac{1}{2}\varepsilon$, so that the eigenvalues can be

1763   approximated by:

1764
$$\lambda_0 \approx 1 - \frac{m}{Nd'x}\left[\frac{(1+Mx)}{2Nx} + \frac{2m}{d'}\right]^{-1} = 1 - \alpha$$
(S14a)

1765
$$\lambda_1 \approx 1 - \frac{(1+Mx)}{2Nx} - \frac{2m}{d'} + \frac{m}{Nd'x}\left[\frac{(1+Mx)}{2Nx} + \frac{2m}{d'}\right]^{-1} = 1 - \frac{(1+Mx)}{2Nx} - \frac{2m}{d'} + \alpha$$
(S14b)

1766   where

1767
$$\alpha = \frac{m}{Nd'x}\{\frac{[1+M(x+d'^{-1})]}{2Nx}\}^{-1} = \frac{M}{2Nd'}[1 + M(x + d'^{-1})]^{-1}$$
(S14c)

1768

1769   For large d, we have:

1770
$$\alpha \approx \frac{M}{2N_T(1+Mx)}$$
(S14d)

1771

1772   This expression implies that the asymptotic rate of approach to equilibrium on the

1773   coalescent timescale of $2N_T$ generations is approximately $M/(1 + Mx)$.

1774   The elements of the corresponding right and left eigenvectors, $\boldsymbol{u}'_i$ and $\boldsymbol{v}'_i$, are given by

1775   the following expressions:

1776
$$\frac{u'_{02}}{u'_{01}} = (2m)^{-1}[\frac{(1+Mx)}{2Nx} - \alpha] \approx \frac{(1+Mx)}{Mx}$$
(S15a)

1777

1778
$$\frac{u'_{12}}{u'_{11}} = -(2m)^{-1}(\frac{2m}{d'} - \alpha) \approx -d'^{-1}$$
(S15b)

1779

1780
$$\frac{v'_{02}}{v'_{01}} = d'(2m)^{-1}[\frac{(1+Mx)}{2Nx} - \alpha] \approx d'\frac{(1+Mx)}{Mx}$$
(S15c)

1781

1782
$$\frac{v'_{12}}{v'_{11}} = -d'(2m)^{-1}\left(\frac{2m}{d'} - \alpha\right) \approx -1 \qquad \text{(S15d)}$$

1783

1784   As before, we can arbitrarily set $u'_{01} = u'_{11} = 1$, and use the normalizations $\boldsymbol{u}'^{Tr}_i \boldsymbol{v}'_i = 1$

1785   to obtain the following relations:

1786

1787
$$u'_{02} = (2m)^{-1}\left[\frac{(1+Mx)}{2Nx} - \alpha\right] \approx \frac{(1+Mx)}{Mx} \qquad \text{(S16a)}$$

1788
$$v'_{01} = \{1 + d'[(1 + Mx)(Mx)^{-1}]^2\}^{-1} \qquad \text{(S16b)}$$

1789
$$v'_{02} = d'(2m)^{-1}\left[\frac{(1+Mx)}{2Nx} - \alpha\right]\{1 + d'[(1 + Mx)(Mx)^{-1}]^2\}^{-1}$$

1790
$$\approx d'\frac{(1+Mx)}{Mx}\{1 + d'[(1 + Mx)(Mx)^{-1}]^2\}^{-1} \qquad \text{(S16c)}$$

1791
$$u'_{12} = -(2m)^{-1}\left(\frac{2m}{d'} - \alpha\right) \approx -d'^{-1} \qquad \text{(S16d)}$$

1792
$$v'_{11} = \left[1 + d'(2m)^{-2}\left(\frac{2m}{d'} - \alpha\right)^2\right]^{-1} \approx (1 + d'^{-1})^{-1}$$

1793
$$v'_{12} = -v'_{02} \qquad \text{(S16e)}$$

1794

1795   The spectral decomposition of the power of a matrix can be used to write $\boldsymbol{B}^t =$

1796 $\sum_i \lambda_i^t \boldsymbol{u}'_i \boldsymbol{v}'_i$. This can be substituted into Equation (S10a) to obtain $T_{11w}(t)$ and $T_{11b}(t)$. When d is

1797   large, Equations (S16) yield:

1798

1799
$$\boldsymbol{B}^t \approx \begin{pmatrix} 0 & \frac{Mx}{(1+Mx)} \\ 0 & 1 \end{pmatrix}(1-\alpha)^t + \left[1 - \frac{(1+Mx)}{2Nx}\right]^t \begin{pmatrix} 1 & -\frac{Mx}{(1+Mx)} \\ 0 & 0 \end{pmatrix}$$

1800
$$= \begin{pmatrix} \left[1 - \frac{(1+Mx)}{2Nx}\right]^t & \frac{Mx}{(1+Mx)}\{(1-\alpha)^t - \left[1 - \frac{(1+Mx)}{2Nx}\right]^t\} \\ 0 & (1-\alpha)^t \end{pmatrix}$$

1801

1802
$$\approx \begin{pmatrix} \exp\left[-\frac{(1+Mx)t}{2Nx}\right] & \frac{Mx}{(1+Mx)}\{\exp(-\alpha t) - \exp\left[-\frac{(1+Mx)t}{2Nx}\right]\} \\ 0 & \exp(-\alpha t) \end{pmatrix} \qquad \text{(S17a)}$$

1803

1804   If time is measured in units of coalescent time (T), the last expression can be written as:

1805

1806
$$\boldsymbol{B}^T \approx \begin{pmatrix} \exp\left[-\frac{d(1+Mx)T}{x}\right] & \frac{Mx}{(1+Mx)}\{\exp\left[-\frac{MT}{(1+Mx)}\right] - \exp\left[-\frac{d(1+Mx)T}{x}\right]\} \\ 0 & \exp\left[-\frac{MT}{(1+Mx)}\right] \end{pmatrix} \qquad \text{(S17b)}$$

1807

1808   The terms involving $\exp[-d(1+Mx)T/x]$ decay much faster than those involving $\exp[-$

1809 $MT/(1+Mx)]$, unless d/x is of order l or less, and can therefore be neglected when d is large,

1810   except when T is of order $1/d$, i.e., for early stages in the process. For large d, the equilibrium

1811   values of T_{11w} and T_{11b} are x and approximately $(1+Mx)/M$, respectively; Equation (S17b)

1812   implies that, for $dT \gg x/(1+Mx)$, we have:

1813

1814 $$T_{11w}(T) \approx x + \left[T_{11b}(0) - \frac{(1+Mx)}{M}\right]\frac{Mx}{(1+Mx)}\exp\left[-\frac{MT}{(1+Mx)}\right] \qquad \text{(S18a)}$$

1815

1816 $$T_{11b}(T) \approx \frac{(1+Mx)}{M} + \left[T_{11b}(0) - \frac{(1+Mx)}{M}\right]\exp\left[-\frac{MT}{(1+Mx)}\right] \qquad \text{(S18b)}$$

1817

where $T_{11w}(0) = T_{11b}(0) = 0$ in the case of an instantaneous approach of the inversion to its

equilibrium frequency. In this case, Equation (S18a) has the same form as Equation (S5b) for the

panmictic case, except that $(1+Mx)/M$ replaces x in the exponential function. This corresponds

to the replacement of the migration effective population size, N_T, which determines the expected

coalescent time for pairs of allele sampled within a deme, by the "total effective population

size", $N_T/(1-F_{ST})$, which determines the expected coalescence time for pairs of alleles sample

randomly from the whole population {Charlesworth, 2010 #3373}, p.318. Since $x < (1+Mx)/M$,

this corresponds to a slower asymptotic rate of approach to equilibrium in the case of a

subdivided population, with the rate being an increasing function of M.

A similar treatment of changes in mean coalescent times can be applied to the St

subpopulation by replacing $T_{11w}(t)$ and $T_{11b}(T)$ with $T_{22w}(T)$ and $T_{22b}(T)$, and x with y, setting

$T_{22w}(0) = 1$ and $T_{11b}(0) = (1+My)/M$ in the case of an instantaneous approach of the inversion to

its equilibrium frequency.

The relevance of these asymptotic rates of change is not, however, clear, since the rates

of change caused by the smaller eigenvalue are substantial when d is large, as assumed here, so

that the equilibrium values of the T_{iiw} and T_{iib} ($i = 1$ or 2) may be approached quite early on. The

properties of the early stages of the process can be examined by approximating the exponentials

in Equation (S17), and its equivalent for T_{22w} and T_{22b}, by the first-order terms in the relevant

exponential functions, as was done to obtain Equations (S5) for the panmictic case in section S5.

But it is important to note that this approximation requires $dT(1+Mx)/x \ll 1$ in Equation (S17b),

so that it will be accurate only for $T \ll x/d(1+Mx)$, a brief period of time.

Applying the initial conditions $\delta T_{11w}(0) = -x$, $\delta T_{11b}(0) = -(1+Mx)/M$, $\delta T_{22w}(0) = x$ and

$\delta T_{22b}(0) = (1+M)M^{-1} - (1+My)M^{-1} = x$ to these equations, some simple algebra yields the

following expressions for the case of an instantaneous sweep:

1842

1843
$$T_{11w}(T) \approx \frac{Mx}{(1+Mx)}T \qquad\qquad \text{(S19a)}$$

1844
$$T_{11b}(T) \approx T \qquad\qquad \text{(S19b)}$$

1845
$$T_{22w}(T) \approx 1 - (dxy^{-1})T \qquad\qquad \text{(S19c)}$$

1846
$$T_{22b}(T) \approx \frac{(1+M)}{M} - \frac{Mx}{(1+My)}T \qquad\qquad \text{(S19d)}$$

1847

Equation (S19b) shows that T_{11b} is initially proportional to T, and Equation (S19a) shows

that T_{11w} increases with time at a rate that is a fraction $Mx/(1+Mx)$ of T. Equation (S19c) shows

that T_{22w} decreases rapidly with time when d is large, due to the dominance of the relevant term

in d in the relevant recursion equation; this means that the decline in T_{22w} dominates the

expressions for T_{Sw} and T_{Tw}. These expressions yield the following results for the mean within-

karyotype and overall coalescent times, assuming that d is large and $dMT << 1$:

1854

1855
$$T_{Sw}(T) \approx y - dxT \qquad\qquad \text{(S20a)}$$

1856
$$T_{Tw}(T) \approx y(1+x) - dxyT \qquad\qquad \text{(S20b)}$$

1857
$$T_{Sb}(T) \approx y(1+M)M^{-1} - x(1+My)^{-1}T \qquad\qquad \text{(S20c)}$$

1858
$$T_{Tb}(T) \approx y(1+x)(1+M)M^{-1} + x[1+y(1+M)](1+My)^{-1}T \qquad \text{(S20d)}$$

1859

1860   Combined with Equations (S9c) and (S9d), these expressions can be used to obtain

1861   approximations for F_{ATw} and F_{ATb} as functions of T when dT is small and d is large. After some

1862   algebra, the following expressions are obtained:

1863

1864
$$F_{ATw}(T) \approx \frac{x[y+d(2-x)T]}{y[(1+x)+dxT]} \qquad\qquad \text{(S20e)}$$

1865
$$F_{ATb}(T) \approx \frac{x}{(1+x)}\left\{1 + \frac{M(1-2x+My)}{y(1+x)(1+M)(1+My)}T\right\} \qquad \text{(S20f)}$$

1866

These equations imply that F_{ATw} always increases initially when there is no

recombination, for the case of an instantaneous sweep; F_{ATb} also increases initially, provided that

$y > (2x-1)/M$, a fairly light condition

It is also of interest to ask how the between-population measure of differentiation for the

In subpopulation behaves in the absence of recombination. This can be measured by considering

the within- and between-population mean coalescence times for pairs of randomly sampled

haploid genomes that both carry the inversion, giving $F_{ST,In} = 1 - T_{11w}/T_{11b}$, given that d is

assumed here to be very large. The general equations for coalescence times then imply that:

1875

1876
$$F_{ST,In}(T) \approx \frac{1}{(1+Mx)[1+g(T)]} \qquad (S21a)$$

1877   where:

1878
$$g(T) = [\frac{M}{(1+Mx)}T_{11b}(0) - 1]\exp[-\frac{MT}{(1+Mx)}]$$

1879

1880   Similarly,

1881
$$1 - F_{ST,In}(T) \approx \frac{Mx+(1+Mx)g(T)}{(1+Mx)[1+g(T)]} \qquad (S21b)$$

1882   and

1883
$$h(T) = \frac{1-F_{ST,In}(T)}{F_{ST,In}(T)} - Mx \approx -(1 + Mx)g(T) \qquad (S22c)$$

1884

1885   Equation (S22c) implies that:

1886
$$\frac{d\,h(T)}{dT} \approx -\frac{M\,h(T)}{(1+Mx)} \qquad (S22d)$$

1887

1888   i.e. the asymptotic proportional rate of change of the natural logarithm of the deviation of the

1889   ratio of $1 - F_{ST,In}$ to $F_{ST,In}$ from its equilibrium value of $1/(1 + Mx)$ is equal to $-M/(1 + Mx)$,

1890   which is the same as the asymptotic proportional rates of change of $\delta T_{11w}(T)$ and $\delta T_{11b}(T)$.

1891   As in the case of F_{AT}, the relevance of these asymptotic rates of change is unclear. For

1892   small T, following the same approach as used above for F_{AT}, we have:

1893

1894
$$F_{ST\,In}(T) \approx \frac{1}{(1+Mx)} \qquad (S23)$$

1895

with the somewhat surprising result that $F_{ST,In}$ is constant and equal to its equilibrium value, to

the order of the approximations used here, despite the fact that it is undefined when $T_{11b} = 0$.

This arises from the fact that Equation (S19a) shows that T_{11w} for small $T \approx MxT/(1+Mx)$

whereas Equation (S19b) shows that T_{12b} is proportional to T, so that $1 - T_{11w} / T_{11b} \approx 1/(1 + Mx)$.

Another way of obtaining this result is to note that, if α in Equation (S17a) is neglected (as

is reasonable for small t), the rate of change per generation in T_{11w}, given by **B**, is as follows:

1902

1903
$$\Delta T_{11w}(t) \approx -\frac{(1+Mx)}{2Nx}T_{11w}(t) + \frac{Mx}{(1+Mx)}T_{11b}(t) \qquad \text{(S24a)}$$

1904

1905   This expression is equal to zero when:

1906

1907
$$\frac{T_{11w}(t)}{T_{11b}(t)} = \frac{Mx}{(1+Mx)} \qquad\qquad\qquad \text{(S24b)}$$

1908

1909   which is equivalent to Equation (S23). The rate of change per generation in T_{11w} is initially very

1910   small, so that T_{11w} reaches a quasi-equilibrium such that its ratio with respect to T_{11b}, and hence

1911 $F_{ST,In}$, is approximately constant over time.

1912   A similar treatment of F_{ST} can be applied to the St subpopulation by replacing $T_{11w}(t)$ and

1913 $T_{11b}(T)$ with $T_{22w}(T)$ and $T_{22b}(T)$, and x with y.

1914

1915
1916
1917