# Coexistence in Periodic Environments

Alexa M Scott[1*], Carling Bieg[1,2], Bailey C McMeans[3], Kevin S McCann[1]

[1] Department of Integrative Biology, University of Guelph, Ontario, Canada

[2] Department of Ecology and Evolutionary Biology, Yale University, New Haven, Connecticut, USA

[3] Department of Biology, University of Toronto, Mississauga, Ontario, Canada

*corresponding author

ORCIDs and email addresses

*Alexa M Scott: 0000-0003-0776-0582 (ascott16@uoguelph.ca)

Carling Bieg: 0000-0003-1552-2007 (carling.bieg@gmail.com)

Bailey C McMeans: 0000-0002-9793-6811 (bailey.mcmeans@utoronto.ca)

Kevin S McCann: 0000-0001-6031-7913 (ksmccann@uoguelph.ca)

Keywords: Periodicities, species coexistence, competition, Lotka-Volterra, climate change

Word Count: 5340 (main body)

Submission Content: Main text, one table, six figures, supplementary material (4 sections)

**1  Abstract**

Climate change and other anthropogenic impacts are rapidly altering natural environmental

periodicities on a variety of time scales. Despite this, a general theoretical foundation describing

the role of periodic environmental variation in structuring species interactions and ecological

communities is still underdeveloped. Alarmingly, this leaves us unprepared to understand and

predict implications for the maintenance of biodiversity under global change. Here, we extend a

two-species Lotka-Volterra competition model that incorporates periodic forcing between

seasons of high and low production to investigate the effects of changing environmental patterns

on species coexistence. Towards this, we define coexistence criteria for periodic environments

by approximating isocline solutions akin to classical coexistence outcomes. This analytical

approach illustrates that periodic environments (i.e., seasonality) in and of themselves can

mediate different competitive outcomes, and these patterns are general across varying time

scales. Importantly, species coexistence may be incredibly sensitive to changes in these abiotic

periods, suggesting that climate change has the potential to drastically impact the maintenance of

biodiversity in the future.

**Introduction**

Nature is abundant with a diverse array of periodic climate signals (Jackson et al. 2021; Pokorný 2021). The complex variations in temperature over time (Jiang and Morin 2007; Klausmeier 2010), for example, can be decomposed into different lengths of underlying periodicities using spectral analysis, revealing a complex mosaic of short (e.g., seconds, minutes, hours, days), medium (e.g., months, years), long (e.g., multi-decadal), and very long natural periods (e.g., 100s to 1000s of years) (Forrest and Miller-Rushing 2010; Vasconcellos et al. 2011; Huntly et al. 2021; Joseph and Kumar 2021; Pokorný 2021). The regularity of these environmental periods allows for species to adapt and respond to them (Bernhardt et al. 2020; Fretwell 1972; Shuter et al. 2012; Tonkin et al. 2017), meaning that nature has evolved around, and within, these complex temporal abiotic signatures (Mathias and Chesson 2013; Varpe 2017; Rudolf 2019). Despite the long-known recognition of nature's complex abiotic palette (White and Hastings 2020), relatively little ecological research has considered the scope of nature's abiotic variability in maintaining species diversity (Abrams 2022).

Researchers have clearly argued that temporal variation (e.g., stochasticity) can promote species coexistence via fluctuation-dependent coexistence mechanisms (e.g., storage effect, relative nonlinearity) (Chesson 2000, 2018; Adler et al. 2006; Meyer et al. 2022). With these mechanisms, temporal niche differentiation enables coexistence between species with different competitive advantages (Angert et al. 2009; Mathias and Chesson 2013; Miller and Klausmeier 2017). Despite the broad focus on environmental variation, one specific class of variability that has been less-well explored, yet ought to provide a generalizable and analytically tractable

entryway into coexistence theory for variable environments, is that of periodic fluctuations – that

is, repeated and predictable fluctuations in environmental conditions. For examples, (Litchman

and Klausmeier 2001) use fast/slow approximations to elegantly show that night/day oscillations

in light can mediate competitive outcomes in phytoplankton – analytical solation that are tricky

to garner from environmentally stochastic models.

Studies are beginning to suggest that periodic environments may have significant implications

for species coexistence (Mathias and Chesson 2013; Miller and Klausmeier 2017; White and

Hastings 2020). For example, temporally changing resource conditions, which fluctuate between

seasons of high productivity to seasons of very little to no productivity (Fretwell 1972; Chesson

and Huntly 1997), may favour different species at different times of a period (Armstrong and

McGehee 1980; Litchman and Klausmeier 2001; Hastings 2012; Huntly et al. 2021). Similarly, it

has been suggested that certain life history trade-offs and temporal differentiation in competing

species' performance may alter coexistence outcomes in the face of periodic environments

(Litchman and Klausmeier 2001; McMeans et al. 2020). Recently, Mougi (2020) extended

competition results from single periodicities to polyrhythms (i.e., multiple interacting

periodicities) to show that the coupling of differently timed resource fluctuations may broaden

the range of coexistence between diverse species that rely on limited resources. While these

recent papers highlight the importance of periodic variation, the demand for a more general

theoretical understanding on the role of periodic conditions – either in isolation or as suites of

periodicities (i.e., polyrhythms) – remains (White and Hastings 2020; Abrams 2022). Notably,

we lack a general understanding of coexistence in periodic environments akin to our well-

established theoretical foundation framed around steady-state dynamics.

The development of such general theory is critical as climate change is currently altering the

nature of these environmental fluctuations (Dijkstra et al 2011; Shuter et al. 2012; Urban et al.

2012; Chesson 2018; Al-Habahbeh et al. 2020). Northern-hemisphere winters are becoming

shorter in length and more moderate (Caldwell et al. 2020; Edlund et al. 2017; Ficker et al. 2017;

Warne et al. 2020), and weather patterns across the globe are becoming more variable and

unpredictable (Fang and Stefan 1998; O'Reilly et al. 2015). In response, many communities have

experienced an increase in species extinction (Urban et al. 2012; Moor 2017; Fung et al. 2020)

and invasion rates (Stachowicz et al. 2002; Sharma et al. 2009; Dijkstra et al. 2011; Cerasoli et

al. 2019; Atkinson et al. 2020). Therefore, as climate change continues to alter the abiotic

conditions to which organisms have adapted to, the mechanisms regulating species coexistence

may be fundamentally altered (di Paola et al. 2012; Korpela et al. 2013; Tunney et al. 2014;

Anderson et al. 2015; Eloranta et al. 2016; Bartley et al. 2019; Caldwell et al. 2020). With all this

in mind, developing an understanding for the mechanisms behind the maintenance of

biodiversity in periodic environments becomes even more crucial.

Inspired by the fluctuating-light-driven coexistence results of Litchman and Klausmeier (2001),

we sought to develop a generalizable framework for coexistence in fluctuating environments.

Towards this, we extend upon the seasonal coexistence model first introduced by McMeans et al.

(2020) to more broadly explore the role of periodic forcing across time scales (e.g., days to

multi-decadal) via different biological parameter combinations. We also generalized our

approach by allowing high and low growth periods (not just high growth - no growth alone),

with temporally-differential competitive abilities. Specifically, we extend the classical Lotka-

Volterra coexistence criteria to include the role of environmental periodicities. This approach

allows our results to be phrased around classical coexistence conditions with temporally-scaled

inter- and intraspecific competition strengths. Here, we define a period as a unit of time that is

composed of two distinct seasons of variable length and seek to generally explore which

competitive outcomes may occur in these environments and under what biological conditions

(i.e., different growth rates). Towards this, we employ an analytical approach consistent with the

classical Lotka-Volterra phaseplane theory by developing a simple approximation that allows us

to solve for the isocline solutions of a time-separated periodic model. Specifically, this

approximation allows us to define coexistence criteria for periodic environments. We then

illustrate how periodic environments can, in and of themselves, drive bifurcations (i.e., changing

invasion criteria) such that competitive outcomes (i.e., stable coexistence, competitive exclusion,

and contingent coexistence) are mediated by the environment. We end by discussing our

competition results in light of how climate change is altering the nature of key underlying abiotic

periodicities.

**Methods**

We start by extending McMeans et al. (2020)'s annual seasonal model. Here, we define season

more generally as a discrete division in time that repeats itself, or is periodic, of any given length

within a period. As such, summer (more productive) and winter (less productive) seasons in

McMeans et al. (2020) repeat themselves with a periodicity of one-year, but we may also

similarly decompose other naturally shorter (e.g., seconds (Huntly et al. 2021)) and longer (e.g.,

El-Nino Southern Oscillations (Joseph and Kumar 2021)) periods of time into discrete seasons of

more or less productive conditions. Towards this general understanding of periodic

environments, we extend the Lotka-Volterra competition model (Chesson 2018) into a periodic

model that repeatedly alternates between two discrete seasons, a productive ($f_P$) and less

productive ($f_{LP}$) season. For each species, these functions are modelled with environmentally

specific parameter combinations to incorporate biological constraints within each season,

discussed below (Fig. 1a). The Lotka-Volterra model is defined as:

$$f_{S,j}(t) = \frac{dX_j}{dt} = r_{S,j}X_j(1 - \alpha_{S,jj}X_j - \alpha_{S,jk}X_k) \tag{1}$$

where $j$ and $k$ represent two competing species, and $S$ represents a season (either productive, $P$,

or less productive, $LP$). Here, $r_{S,j}$ is the intrinsic rate of population growth for species $j$ in a

season, $S$, $\alpha_{S,jj}$ is the intraspecific competitive coefficient for species $j$ in season $S$, and $\alpha_{S,jk}$ is

the interspecific competitive coefficient describing the effect of species $k$ on species $j$ in season

$S$.

When running simulations, and to maintain the spirit of the model assumptions (i.e., productive

and less productive season), we assumed the following simple biologically realistic assumptions:

(1) Maximal growth rates are larger in the productive season than the less productive season

121            for both species (i.e., $r_{P,j} > r_{LP,j}$), and;

(2) Since resources are more available in the productive season compared to the less

productive season, intraspecific competition will be lower in the productive season

compared to the less productive season (i.e., $\alpha_{P,jj} < \alpha_{LP,jj}$).

Further, to incorporate realistic biological trade-offs between competing species, we assumed

that species 1 is a better performer (in terms of growth and competition) in the productive season

compared to species 2, and the opposite is true in the less productive season. Keeping in mind

the previous seasonal constraints, this produces the following realistic parametric trade-offs for

the two species:

(1) Species 1 has a higher growth rate in the productive season (i.e., $r_{P,1} > r_{P,2}$) and a lower

growth rate in the less productive season (i.e., $r_{LP,2} > r_{LP,1}$) compared to species 2, and;

(2) Species 1 has a smaller intraspecific competitive coefficient in the productive season (i.e.,

$\alpha_{P,11} < \alpha_{P,22}$) and a larger intraspecific competitive coefficient in the less productive

season (i.e., $\alpha_{LP,11} > \alpha_{LP,22}$) compared to species 2.

These trade-offs are similar to an empirical case study by McMeans et al. (2020) where cold-

adapted fish (e.g., lake trout, *Salvelinus namaycush*) are seasonal generalists with moderate

growth rates year-round, and warm-adapted fish (e.g., smallmouth bass, *Micropterus dolomieu*),

the lake trout's competitor, are seasonal specialists with higher growth rates during the summer

and lower growth rates during the winter. Note that we have not set any seasonal constraints on

the interspecific competitive coefficients, therefore, $\alpha_{S,jk}$ can be any value.

With the above assumptions taken into consideration, our model is a periodic step function

(repeats every 1-time unit, or period) that goes through a productive season ($P$), and a less

productive season ($LP$), as follows:

$$\frac{dX_j}{dt} = \begin{cases} f_{P,j}(t) & i < t < i + \tau p \\ f_{LP,j}(t) & i + \tau p < t < i + \tau \end{cases} \tag{2}$$

Where $i$ is the period number (i.e. year) that runs on an integer step size of 1 from 0 to $t_{prd}$ (the

number of periods that the model runs for). $\tau$ governs the length of each period, and $p$ is the

proportion of each period that is considered productive (i.e., $\tau p$ is the length of the productive

season; leaving $\tau(1\text{-}p)$ as the length of the less productive season). The period length, defined by

$\tau$, allows us to examine how environmental periodicities, and associated biological trade-offs,

across time scales may influence coexistence. As above, $j$ represents one of the two competitive

species, either species 1 or 2. These periodic functions of $f_P$ and $f_{LP}$ are defined in equation (1)

as $f_S$.

We coded all numerical simulations in Mathematica 12.0. The models are integrated over

numerous periods until an asymptotic state, referred to hereafter as an equilibrium state, despite

within-period variation, has been reached (i.e., mean value from 900 – 1000 time-units) to

remove transient influences. However, within each time unit, as discussed above, they

sequentially follow first productive then less productive parameters corresponding to the given

productive seasonal fraction, $p$, within the period. The productive seasonal fraction, $p$, allows us

to change the proportion of the period that is under our productive conditions versus our less

productive conditions, ($1\text{-}p$) (e.g., increase $f_P$) (Fig. 1b).

Finally, all model parameterizations for our simulations can be found in our figure legends and in

the Supplementary Material. Below we first walk through our approximation and then our

general analytical results before highlighting the generality of our analytical solutions using

numerical simulations. For each outcome, we explore how changing the season lengths, via $p$,

influences competitive coexistence and exclusion (Fig. 1b; Table 1).

**Results**

Here, we present our approximated isocline solutions and investigate the resulting coexistence

criteria and behaviour under seasonality. Next, we reveal that changing season length, $p$, in

response to climate change, mediates coexistence, competitive exclusion, and contingent

coexistence (Fig. 3). Finally, we explore the robustness of our seasonally-mediated outcomes

(i.e., seasonally-driven bifurcations) across a range of period lengths (Figs. 4-6).

**Approximate Isocline Solutions for the Periodic Lotka-Volterra Model**

Although we use a periodically forced system, we can use equilibrium concepts to understand

the dynamics of our model. Specifically, our model reaches an attractor such that the densities

fluctuate modestly up and down on the attractor around a mean that does not change (i.e., an

asymptotic state or dynamic equilibrium; Fig.S2.2). Given this equilibrium-like dynamic, we are

interested in considering this asymptotic behaviour in a manner similar to the way we would for

a system that reaches a true equilibrium. Here, we provide an approximation for our model

isoclines and equilibria that mirror those of the original Lotka-Volterra competition model.

Note that the isoclines can be solved by recognizing that each period of $\tau$ time units would

necessarily have to result in zero net growth (i.e., no overall changes in density between the start

and end of each period), akin to classical zero-growth isoclines with a deterministic equilibrium.

That is, from System (2), the $X_j$ zero-net-growth condition occurs when $\frac{dx_j}{dt} = 0$. From this, the

isocline solution follows as:

$$\int_0^{\tau p} f_{P,j}(t) = -\int_{\tau p}^{\tau} f_{LP,j}(t) \tag{3}$$

This isocline solution (3) is not analytically tractable (note that each season has dynamics over a

time interval), making it seem as though an isocline approach appears infeasible.

Towards solving for an approximation for periodically forced isoclines, we take a slightly

different approach then the classical linearization of the full equations (see Litchman and

Klausmeier (2001) for an example). As our model is not tractable with this approach, we instead

proceed by assuming that we can find a linearized approximation to the isocline from the

periodic model by searching for points in phase space where the linearization of each seasons'

dynamics are exactly negated by each other (for both the $X_1$ and $X_2$ isoclines). We define such

points as **zero net growth** points in the phaseplane (i.e., a point on the forced model's isocline)

and we define the collection of these points as **the linearized isocline approximation of the**

**periodically-forced model**. Clearly, as the length of the period, $\tau$, goes to 0, the error in this

linear approximation will also go to 0. However, for longer periods or large growth rates ($r$),

non-linear dynamics may drive this simplification to work poorly (see Supplementary Material

S4). Because of this, we numerically check our analytical results with numerical calculations

throughout. We note that despite this, the results work surprisingly well across large parameter

values suggesting the linear approximation works even when some degree of nonlinear dynamics

are expressed.

By assuming that the instantaneous rates of change for each species at this $X_1$-$X_2$ co-ordinate are

linear, we approximate the dynamics over the time period, $\tau$, by solving the equations using the

Fundamental Theorem of Calculus over each period (see Supplementary Material S1). That is,

over the interval fraction, $\tau p$, the productive trajectory scales linearly on the co-ordinate $X_1$-$X_2$ as:

$$\frac{dX_1}{dt} = \tau p f_{P,1}(X_1, X_2) \tag{4}$$

213   and the less productive trajectory over the interval fraction $\tau(1-p)$ scales linearly as:

$$\frac{dX_1}{dt} = \tau(1-p) f_{LP,1}(X_1, X_2) \tag{5}$$

214   Given Equations (4) and (5), then the linearized dynamics at a $X_1$-$X_2$ co-ordinate negate each

215   other (i.e., resulting in the $X_1$-isocline) when:

$$\tau p f_{P,1}(X_1, X_2) = -\tau(1-p) f_{LP,1}(X_1, X_2) \tag{6}$$

This approximation (Equation (6)), if it works, can be solved symbolically as done elegantly for

the classical Lotka-Volterra model and thus allows an entry point into well-known coexistence

analyses and interpretations.

From Equation (6), we see that $\tau$ factors out. After substituting the productive and less

productive models of Equation (1) into Equation (6) and then some algebra, the resulting isocline

solutions, for both species, follow the following form:

$$X_1 = -\frac{pr_{P,1}\alpha_{P,12} + (1-p)r_{LP,1}\alpha_{LP,12}}{pr_{P,1}\alpha_{P,11} + (1-p)r_{LP,1}\alpha_{LP,11}}X_2 + \frac{pr_{P,1} + (1-p)r_{LP,1}}{pr_{P,1}\alpha_{P,11} + (1-p)r_{LP,1}\alpha_{LP,11}} \qquad (7)$$

The geometry of the approximated isoclines (Equation (7)) resembles that of the original Lotka-

Volterra isoclines, which therefore allows us to identify general rules for coexistence in periodic

environments akin to the original coexistence criteria (Fig. 2a). Here, the isocline approximation

solutions (and therefore coexistence criteria) are now based on temporal-growth-scaled inter-

versus intra-specific competition rates (whereas the original Lotka-Volterra conditions are not

dependent on growth rates; see Table 1 for Lotka-Volterra vs. Seasonal Coexistence Criteria).

Thus, the approximation enables us to generally determine and graphically explore how periodic

environments and biological rates may interact in regulating competitive outcomes (Fig. 2b).

Immediately, based on these conditions, we can see that the duration of each season, and the

corresponding biological conditions within them, play a critical role in regulating competitive

outcomes in periodic environments. That is, our parameter $p$ may drive bifurcations, and

therefore alter competitive outcomes under changing climatic conditions (Fig. 2b).

We found numerically that this approximation works quite well (see Supplementary Material S2

for examples) and was, for example, able to repeatedly and accurately calculate the transcritical

bifurcation points where an interior equilibrium intersects and exchanges stability, with an axial

solution. Other approaches similar to our approximation, such as Litchman and Klausmeier

(2001), have been used to accurately investigate species coexistence when organismal dynamics

are fast relative to their forcing speed. Given our assumption of linearity, the relative time scale

of forcing vs. dynamics ought to be important for our approximation's accuracy. Indeed, when

either the period length ($\tau$) or the growth rate ($r$) becomes larger, the apparent length of each

period increases, allowing for more nonlinearity within each season's dynamics, which results in

the approximation to fail at predicting the asymptotic behaviour (see Supplementary Material

S4). Nevertheless, the range at which this approximation holds is impressively robust.

**General outcomes of changing environments**

Given that changing season lengths may have powerful implications for species coexistence, we

now ask how changing season length, $p$, (i.e., altering seasonal asymmetry) affects competitive

outcomes. To do this we vary the season length from $p=0$ (i.e., the full period being entirely less

productive) to $p=1$ (i.e., the full period being entirely productive). We note that these endpoints

(i.e., $p=0$ or $p=1$) are constant (i.e., no seasonality) and thus, they reduce to the classic Lotka-

Volterra coexistence criteria based on the individual biological conditions for $P$ or $LP$ (Table 1).

These endpoints also give us reference conditions for exploring the effects of environmental

periodicity and seasonally-mediated coexistence outcomes.

Since McMeans et al. (2020) found the intriguing case of seasonally-mediated coexistence, we

were interested in using our analytical results to unpack other examples of how seasonal change

could fundamentally alter competitive outcomes. McMeans et al. (2020) noted that where the

boundary conditions yielded competitive exclusion (i.e., exclusion at $p=0$ and $p=1$), a seasonal

model could produce coexistence for intermediate $p$-values. This is intriguing as it immediately

suggests that alterations in $p$ (say from climate change) can drive exclusion. Importantly, our

analytical solutions (Table 1) suggest that seasonality can mediate all possible competitive

outcomes at intermediate $p$-values (Fig. 3). Indeed, we found seasonally-mediated coexistence,

competitive exclusion, and contingent coexistence (i.e., alternative states) (Fig. 3a-c

respectively).

For these seasonally-mediated outcomes, when the qualitative competitive outcome at

intermediate $p$ is fundamentally different from the extremes (i.e., $p = 0$ and 1), $p$ drives a series

of transcritical bifurcations that move the system between different conditions in Table 1 (shifts

between shaded and clear zones indicate transcritical bifurcations that alter the qualitative nature

of the attractor in Fig. 3). Specifically, under our given sets of parametric combinations,

changing $p$ always drives a series of two bifurcations for these seasonally-mediated outcomes

(Fig. 3). Other competitive outcomes of course are seasonally sensitive as they result in $p$-driven

bifurcations when coexistence outcomes transition between the two endpoint conditions.

Additionally, we note that these transcritical bifurcations can be considered as changes in the

invasion criteria – in this case, seasonally-altered invasion criterion – as they are commonly

referred to in competition theory. For example, in Fig. 3a, the gray shaded regions indicate that

one species cannot invade from small densities while in the white or unshaded region, it

suddenly can invade from small densities.

Finally, clearly not all parameter combinations are sensitive to $p$-driven bifurcations. In these

cases, $p$ may simply drive changes in species' densities rather than shifts in the equilibrium

structure (e.g., as seen in McMeans et al. (2020)). While here we broadly unpack $p$-driven

bifurcations to explore the generality of these outcomes, we note that all results below have

nearby solutions that do not undergo bifurcations yet and are qualitatively similar in terms of the

general effect of $p$ on the isocline and equilibrium geometry.

**Robustness of Seasonally-Mediated Competitive Outcomes**

To fully understand how robust the different seasonally-mediated outcomes are we looked at the

bifurcation structure in 2-dimensional parameter space, by using both season length, $p$, (as in Fig.

3) and the period length, $\tau$, as bifurcation parameters. We note that if seasonally-mediated

outcomes exist across a broad range of period lengths, $\tau$, then this suggests that seasonally-

mediated competitive outcomes could be found across a wide range of natural periodicities (from

diurnal to multi-decadal).

In Fig. 4a, we document cases where a lack of periodicity (i.e., at $p$=0 or 1) drives competitive

exclusion while for a broad range of $\tau$'s (ranging from 1-100 time units), we have species

coexistence at intermediate $p$-values. Our analytical solutions allow us to track the "mean

equilibrium" through each transcritical bifurcation from competitive exclusion of species 1 to

species coexistence and finally to competitive exclusion of species 2 as the length of the

productive period, $p$, increases (i.e., changing $p$ shifts between competitive exclusion and stable

coexistence conditions in Table 1; Figs. 4bi-iii, S3.1). The robustness of this pattern implies that

for any given $\tau$, seasonality alone can mediate coexistence at intermediate $p$-values. As the

period length ($\tau$) grows, the numerically-generated region of coexistence broadens, and the

second transcritical bifurcation point, the transition between species coexistence and competitive

exclusion of species 2, deviates away from the analytically-determined transcritical bifurcation

point (see more regarding approximation accuracy of seasonally mediated coexistence in S4.1).

This suggests that the analytical bifurcation structure is sensitive to different parametric values

(which is also suggested by the seasonal coexistence criteria shown in Table 1), but nonetheless,

the result of seasonally-mediated coexistence is qualitatively general. We note that the simple

linear prediction breaks down as r$\tau$ grows (i.e., larger $\tau r$ allows the nonlinear dynamics to

express themselves; see Supplementary Material S4.1). Thus, we see that seasonally-mediated

coexistence occurs very broadly suggesting that a large range of natural periodicities may be

powerful drivers of coexistence.

Similarly, in Fig. 5a, we document cases where species coexistence occurs in static environments

(i.e., at $p$=0 or 1), while competitive exclusion occurs at intermediate $p$-values over a broad range

of $\tau$'s (here, from 1-40 time units). Following McMeans et al. (2020)'s naming convention, we

refer to this as seasonally-mediated competitive exclusion. As in the above case, with increasing

$p$, our analytical approximation tracks the isoclines and the "mean equilibrium" from stable

coexistence to competitive exclusion of species 2 through a transcritical bifurcation on the $X_2$=0

axis, and finally back through the axis (via another bifurcation) to stable coexistence (Figs. 5b,

S3.2). The transcritical bifurcations points determined by our analytical solution accurately track

the numerical solutions over several period length values of $\tau$ before the seasonal bifurcations

cease at higher values of $\tau$. Again, we note that the simple linear prediction breaks down as $\tau r$

grows (i.e., larger $\tau r$ allows the nonlinear dynamics to express themselves; see Supplementary

Material S4.2). The 2-dimensional bifurcation diagram again shows a broad range in $\tau$-values

that yield competitive exclusion; however, it does not stretch across all $\tau$-values (i.e., very large

period lengths). While the exact range of parameter space that this seasonally-mediated

competitive exclusion occurs at is clearly dependent on other parameters, this result is still quite

general over a wide range of $\tau$-values, suggesting that periodic environments may also be

powerful drivers of competitive exclusion. Here, as $\tau$ grows, this tongue of competitive

exclusion slowly shrinks across intermediate values of $p$. Again, this is an interesting result,

which suggests that simple alterations in season length may fundamentally alter competitive

outcomes.

Finally, in Fig. 6a, similar to the above seasonally-mediated coexistence case with competitive

exclusion at the boundaries (i.e., $p=0$ or $1$), we now document cases where contingent

coexistence (i.e., alternative states) occurs at intermediate $p$ values across a broad range of $\tau$'s

(here, from 1-100 time units). Although this outcome is quite similar to seasonally-mediated

coexistence with competitive exclusion being located at each boundary (i.e., $p=0$ or $1$), its

isocline geometry, based off of parametric combinations, results in seasonally-mediated

contingent coexistence at intermediate $p$-values (Figs. 6bi-iii, S3.3). Our approximation tracks

the analytical non-trivial (interior) equilibrium, which in this case is unstable in positive ($X_1$, $X_2$)

state space (Fig. 6bii). Here, initial densities determine which species will dominate and which

species will become extinct with time (i.e., which of the two stable axial solutions, shown in Fig.

6bii, the system will end up at). The robustness of this pattern also implies that for any given $\tau$,

seasonality may be a powerful driver of contingent coexistence (i.e., changing $p$ moves from

competitive exclusion to contingent coexistence conditions in Table 1). As $\tau$ grows, the

numerically-generated contingent coexistence region expands across higher productive season

lengths before it narrows around lower lengths of the productive season. Again, our analytical

solution accurately determines at which $p$ values the transcritical bifurcations will occur for very

small $\tau$ values, but falls off with larger period lengths, $\tau$ (see Supplementary Material S4.3 for

approximation accuracy of seasonally mediated contingent coexistence), despite the general

phenomenon remaining across time scales.

These three seasonally-mediated outcomes suggest that seasonal alterations, due to climate

change, may drive precipitous changes in competitive systems. All of these seasonally-mediated

outcomes appear general and may be found across various ecosystems that experience abiotic

fluctuations of any period length. Similarly, for a given period length (e.g., annual variation),

these outcomes may be highly general for competing organisms with a range of life history

strategies (i.e., vital rates that dictate the speed of biotic dynamics relative to any environmental

variation). On the other hand, depending on parametric values, not all instances of periodic

variation will drive these series of bifurcations. Under some conditions, there are simply $p$-driven

changes in species densities between the boundary conditions where only one bifurcation (Figs.

S2.1e, S2.1f) or no bifurcations (Figs. S2.1d) are found under changing periodicities.

**Discussion**

Here, we present a linear approximation similar in approach to other periodic models (e.g., Han et al. (1999); Litchman and Klausmeier (2001)) to analytically solve for a periodic Lotka-Volterra competition model (see Table 1 for seasonal coexistence criteria). Our linear approximation quantitatively breaks down for large values in $\tau r$ but still largely operates to qualitatively predict the nature of periodic-forcing on competition (e.g., Fig. 4a,6a). This approximation, akin to the longstanding classical Lotka-Volterra coexistence conditions, importantly allows us to derive a parallel set of coexistence conditions for periodic environments. With the inclusion of periodic environmental forcing, we find that species coexistence depends on seasonal-growth-scaled inter- versus intra-specific competition strengths (Table 1). That is, where classical stable coexistence requires intraspecific competition to be greater than interspecific competition, for both species, we show that intraspecific competition, scaled by seasonal growth, must be greater than interspecific competition, scaled by seasonal growth, for both species.

These seasonal coexistence conditions importantly suggest that species' coexistence may be incredibly sensitive to changes in season length. While McMeans et al. (2020) found seasonally-mediated coexistence through empirically-motivated numerical simulations, our results show that seasonality can mediate coexistence, competitive exclusion, and contingent coexistence (Fig. 3). That is, seasonality, in and of itself, can drive all possible competitive outcomes (i.e., Table 1). As these seasonally-mediated outcomes appear across a vast range of period lengths (Fig. 4-6), this suggests that these seasonally-mediated outcomes are quite robust, and can be found across a

wide range of different temporal scales and life history strategies (i.e., fast vs. slow life history

strategies).

As the change in season length ($p$) may have an important role in mediating competitive

outcomes (Fig. 3), climate change is altering environmental periodicities across a range of

temporal scales (Dijkstra et al. 2011; Shuter et al. 2012; Urban et al. 2012; Chesson 2018; Al-

Habahbeh et al. 2020), and our results suggest that this could have sudden and drastic –

sometimes unpredictable – effects on coexistence. Often the responses to changing seasons are

nonlinear, and some of these outcomes are unexpected based on the classical Lotka-Volterra

conditions under non-seasonal environments (e.g., nonlinear paths between the boundary

conditions at $p = 0$ and 1; Figs. 3b, 5, S2.1b and d). As an example, we find cases where

competitive exclusion occurs under intermediate season lengths, even though coexistence is

expected under the same non-seasonal conditions (i.e., coexistence occurs for entirely low- or

entirely high-productive conditions; Fig. 3b). Alarmingly, these strong nonlinear effects of

changing season length, $p$, suggest that precipitous changes in species density and composition

may unexpectedly occur as climate change alters the nature of seasonality.

Although simple, the seasonal competition model we examined here is based on biologically

realistic assumptions about competitive species in temporally varying environments. First,

periodic environments are commonly reflected by times of high and low productivity, where

species tend to flourish during the productive seasons, and decline (e.g., mortality-dominated)

during the less productive seasons as resource availability becomes sparse (Fretwell 1972;

Litchman and Klausmeier 2001; Mutze 2009; Klausmeier 2010; Hastings 2012; Lyu et al. 2016;

Vihtakari et al. 2016). These periods therefore impact species' growth rates, but also offer the

potential for differential responses of competing species to seasonally-driven environmental

variation (Chesson and Huntly 1997; Forrest and Miller-Rushing 2010; Shuter et al. 2012; Gao et

al. 2016; Chesson 2018; Huang et al. 2019). It is well known that competing species can display

temporal trade-offs in stochastic environments that can promote coexistence (Angert et al. 2009;

Shuter et al. 2012; Mougi 2020). For instance, Litchman and Klausmeier (2001) discovered that

slow fluctuations in light promote stable coexistence between competing species of

phytoplankton who exhibit trade-offs in performance across temporally changing resource

availability. More recently, and consistent with our theoretical results here, empirical evidence is

beginning to suggest species may show trade-offs to regular seasonal fluctuations that might

promote coexistence. For example, a species may be a seasonal specialist (e.g., display a very

high growth rate during the summer and a low growth rate during the winter) while its

competitor may be more of a temporal generalist, who's able to maintain roughly the same

growth rate year-round (Niiyama 1990; Chan et al. 2009; Abrams et al. 2013; Korpela et al.

2013; Meyer et al. 2022).

While we concentrate on temporal forcing in our model, seasonal signals in competition may

often be related to spatial-temporal and behavioural patterns that govern species' competitive

outcomes (McMeans et al. 2020). As an example, in some cases, species may migrate (Holt and

Fryxell 2011; Teitelbaum et al. 2015; Tavecchia et al. 2016; le Corre et al. 2020; Moorter et al.

2021) or hibernate (Campbell et al. 2008; Giraldo-Perez et al. 2016) during less productive

seasons, opting out of competition when resources are scarce (Suski and Ridgway 2009). These

behavioural strategies may counteract potential negative effects of periodic variation to some

extent and thus promote coexistence by reducing competitive interactions during unfavourable

times. More empirical research is needed to further understand these spatial and behavioural

strategies, and seasonal trade-offs between competitive species, in order to understand their

mechanistic role in maintaining biodiversity with climate change.

As nature abounds with temporal variation, grasping a better understanding of coexistence

mechanisms in these fluctuating environments is now even more crucial with climate change.

We have provided an analytical solution to begin understanding these effects. Our research has

uncovered three competitive outcomes that could be found across a large range of natural

periodicities and life history strategies. As species coexistence appears to be incredibly sensitive

to periodic variation, climate change has the potential to drastically impact future competitive

outcomes in the natural world.

**Acknowledgements**

We would like to thank Cortland K. Griswold for his comments and suggestions that helped

improve this manuscript. We would also like to thank the University of Guelph's Elgin Card

Scholarship in Terrestrial Ecology (Grant number [I5023]) awarded to AMS, and the NSERC

Discovery Grant granted to KSM (Grant number [400353]) for funding this research.

**Code accessibility**

All code to reproduce the above analyses and figures have been archived on <u>Zenodo</u> (version

1.0).

**Literature Cited**

Abrams PA (2022) Competition in seasonal environments. In: Competition Theory in Ecology.
Oxford University Press, pp 171–208

Abrams PA, Tucker CM, Gilbert B (2013) Evolution of the storage effect. Evolution (N Y)
67:315–327. https://doi.org/10.1111/j.1558-5646.2012.01756.x

Adler PB, HilleRisLambers J, Kyriakidis PC, et al (2006) Climate variability has a stabilizing
effect on the coexistence of prairie grasses. Proc Natl Acad Sci U S A 103:12793–12798.
https://doi.org/10.1073/pnas.0600599103

Al-Habahbeh AK, Kortsch S, Bluhm BA, et al (2020) Arctic coastal benthos long-term responses
to perturbations under climate warming. Philos Trans A Math Phys Eng Sci 378:20190355.
https://doi.org/10.1098/rsta.2019.0355

Anderson KD, Heron SF, Pratchett MS (2015) Species-specific declines in the linear extension
of branching corals at a subtropical reef, Lord Howe Island. Coral Reefs 34:479–490.
https://doi.org/10.1007/s00338-014-1251-1

Angert AL, Huxman TE, Chesson P, Venable DL (2009) Functional tradeoffs determine species
coexistence via the storage effect. Proc Natl Acad Sci U S A 106:11641–11645.
https://doi.org/10.1073/pnas.0904512106

Armstrong RA, McGehee R (1980) Competitive Exclusion. American Society of Naturalist
115:151–170

Atkinson J, King NG, Wilmes SB, Moore PJ (2020) Summer and winter marine heatwaves favor
an invasive over native seaweeds. J Phycol 56:1591–1600.
https://doi.org/10.1111/jpy.13051

Bartley TJ, McCann KS, Bieg C, et al (2019) Food web rewiring in a changing world. Nat Ecol
Evol 3:345–354. https://doi.org/10.1038/s41559-018-0772-3

Bernhardt J, O'Connor M, Sunday J, Gonzalez A (2020) Life in fluctuating environments. Philos
Trans R Soc Lond B Biol Sci 375:1–22. https://doi.org/10.32942/osf.io/dgmr4

Caldwell TJ, Chandra S, Feher K, et al (2020) Ecosystem response to earlier ice break-up date:
Climate-driven changes to water temperature, lake-habitat-specific production, and trout
habitat and resource use. Glob Chang Biol 26:5475–5491.
https://doi.org/10.1111/gcb.15258

Campbell HA, Fraser KPP, Bishop CM, et al (2008) Hibernation in an Antarctic fish: On ice for
winter. PLoS One 3:. https://doi.org/10.1371/journal.pone.0001743

Cerasoli F, Iannella M, Biondi M (2019) Between the hammer and the anvil: How the combined
effect of global warming and the non-native common slider could threaten the european
pond turtle. Management of Biological Invasions 10:428–448.
https://doi.org/10.3391/mbi.2019.10.3.02

Chan B, Durrett R, Lanchier N (2009) Coexistence for a multitype contact process with seasons.
Annals of Applied Probability 19:1921–1943. https://doi.org/10.1214/09-AAP599

Chesson P (2018) Updates on mechanisms of maintenance of species diversity. Journal of
Ecology 106:1773–1794. https://doi.org/10.1111/1365-2745.13035

Chesson P (2000) Mechanisms of maintenance of speacies diversity. Annu Rev Ecol Syst
31:343–366. https://doi.org/10.1146/annurev.ecolsys.31.1.343

Chesson P, Huntly N (1997) The roles of harsh and fluctuating conditions in the dynamics of
ecological communities. Am Nat 150:519–553

di Paola A, Valentini R, Paparella F (2012) Climate change threatens coexistence within
communities of mediterranean forested wetlands. PLoS One 7:.
https://doi.org/10.1371/journal.pone.0044727

Dijkstra JA, Westerman EL, Harris LG (2011) The effects of climate change on species
composition, succession and phenology: A case study. Glob Chang Biol 17:2360–2369.
https://doi.org/10.1111/j.1365-2486.2010.02371.x

Edlund MB, Almendinger JE, Fang X, et al (2017) Effects of climate change on lake thermal
structure and biotic response in northern wilderness lakes. Water (Basel) 9:1–35.
https://doi.org/10.3390/w9090678

Eloranta AP, Helland IP, Sandlund OT, et al (2016) Community structure influences species'
abundance along environmental gradients. Journal of Animal Ecology 85:273–282.
https://doi.org/10.1111/1365-2656.12461

Fang X, Stefan HG (1998) Potential climate warming effects on ice covers of small lakes in the
contiguous U.S. Cold Reg Sci Technol 27:119–140. https://doi.org/10.1016/S0165-
232X(97)00027-X

Ficker H, Luger M, Gassner H (2017) From dimictic to monomictic: Empirical evidence of
thermal regime transitions in three deep alpine lakes in Austria induced by climate change.
Freshw Biol 62:1335–1345. https://doi.org/10.1111/fwb.12946

Forrest J, Miller-Rushing AJ (2010) Toward a synthetic understanding of the role of phenology
in ecology and evolution. Philosophical Transactions of the Royal Society B: Biological
Sciences 365:3101–3112. https://doi.org/10.1098/rstb.2010.0145

Fretwell SD (1972) Populations in a seasonal environment. Princeton University Press

Fung T, Chisholm RA, Anderson-Teixeira K, et al (2020) Temporal population variability in local forest communities has mixed effects on tree species richness across a latitudinal gradient. Ecol Lett 23:160–171. https://doi.org/10.1111/ele.13412

Gao G, Ponsard S, Lu Z, et al (2016) Different thermal responses in two coexisting aphids may account for differences in their seasonal abundance in cotton fields. Int J Pest Manag 62:288–294. https://doi.org/10.1080/09670874.2016.1185551

Giraldo-Perez P, Herrera P, Campbell A, et al (2016) Winter is coming: Hibernation reverses the outcome of sperm competition in a fly. J Evol Biol 29:371–379. https://doi.org/10.1111/jeb.12792

Han BP, Virtanen M, Koponen J, Straškraba M (1999) Predictors of light-limited growth and competition of phytoplankton in a well-mixed water column. J Theor Biol 197:439–450. https://doi.org/10.1006/jtbi.1998.0885

Hastings A (2012) Temporally varying resources amplify the importance of resource input in ecological populations. Biol Lett 8:1067–1069. https://doi.org/10.1098/rsbl.2012.0669

Holt RD, Fryxell JM (2011) Theoretical reflections on the evolution of migration. In: Milner-Gulland EJ, Fryxell JM, Sinclaire ARE (eds) Animal migration. Oxford University Press, New York, pp 17–31

Huang L, Xue W, Herben T (2019) Temporal niche differentiation among species changes with habitat productivity and light conditions. Journal of Vegetation Science 30:438–447. https://doi.org/10.1111/jvs.12741

Huntly N, Freischel AR, Miller AK, et al (2021) Coexistence of "cream skimmer" and "crumb picker" phenotypes in nature and in cancer. 9:1–18. https://doi.org/10.3389/fevo.2021.697618

Jackson MC, Pawar S, Woodward G (2021) The temporal dynamics of multiple stressor effects: From individuals to ecosystems. Trends Ecol Evol 36:402–410. https://doi.org/10.1016/j.tree.2021.01.005

Jiang L, Morin PJ (2007) Temperature fluctuation facilitates coexistence of competing species in experimental microbial communities. Journal of Animal Ecology 76:660–668. https://doi.org/10.1111/j.1365-2656.2007.01252.x

Joseph D, Kumar VS (2021) Response of ocean surface waves to the co-occurrence of Boreal Summer Intra-Seasonal Oscillation and El Niño Southern Oscillation. Clim Dyn 57:1155–1171. https://doi.org/10.1007/s00382-021-05763-3

Klausmeier CA (2010) Successional state dynamics: A novel approach to modeling nonequilibrium foodweb dynamics. J Theor Biol 262:584–595. https://doi.org/10.1016/j.jtbi.2009.10.018

Korpela K, Delgado M, Henttonen H, et al (2013) Nonlinear effects of climate on boreal rodent
dynamics: Mild winters do not negate high-amplitude cycles. Glob Chang Biol 19:697–710.
https://doi.org/10.1111/gcb.12099

le Corre M, Dussault C, Côté SD (2020) Where to spend the winter? The role of intraspecific
competition and climate in determining the selection of wintering areas by migratory
caribou. Oikos 129:512–525. https://doi.org/10.1111/oik.06668

Litchman E, Klausmeier CA (2001) Competition of phytoplankton under fluctuating light.
American Naturalist 157:170–187. https://doi.org/10.1086/318628

Lyu L, Deng X, Zhang Q bin (2016) Elevation pattern in growth coherency on the southeastern
Tibetan Plateau. PLoS One 11:1–15. https://doi.org/10.1371/journal.pone.0163201

Mathias A, Chesson P (2013) Coexistence and evolutionary dynamics mediated by seasonal
environmental variation in annual plant communities. Theor Popul Biol 84:56–71.
https://doi.org/10.1016/j.tpb.2012.11.009

McMeans BC, McCann KS, Guzzo MM, et al (2020) Winter in water: Differential responses and
the maintenance of biodiversity. Ecol Lett 23:922–938. https://doi.org/10.1111/ele.13504

Meyer I, Steinmetz B, Shnerb NM (2022) How the storage effect and the number of temporal
niches affect biodiversity in stochastic and seasonal environments. PLoS Comput Biol
18:1–21. https://doi.org/10.1371/journal.pcbi.1009971

Miller ET, Klausmeier CA (2017) Evolutionary stability of coexistence due to the storage effect
in a two-season model. Theor Ecol 10:91–103. https://doi.org/10.1007/s12080-016-0314-z

Moor H (2017) Life history trade-off moderates model predictions of diversity loss from climate
change. PLoS One 12:1–21. https://doi.org/10.1371/journal.pone.0177778

Moorter B van, Singh NJ, Rolandsen CM, et al (2021) Seasonal release from competition
explains partial migration in European moose. Oikos 130:1548–1561.
https://doi.org/10.1111/oik.07875

Mougi A (2020) Polyrhythmic foraging and competitive coexistence. Sci Rep 10:1–7.
https://doi.org/10.1038/s41598-020-77483-3

Mutze GJ (2009) Changes in body condition and body size affect breeding and recruitment in
fluctuating house mouse populations in south-eastern Australia. Austral Ecol 34:278–293.
https://doi.org/10.1111/j.1442-9993.2008.01929.x

Niiyama K (1990) The role of seed dispersal and seedling traits in colonization and coexistence
of Salix species in a seasonally flooded habitat. Ecol Res 5:317–331

O'Reilly CM, Sharma S, Gray DK, et al (2015) Rapid and highly variable warming of lake
surface waters around the globe. Geophys Res Lett 42:10773–10781.
https://doi.org/10.1002/2015GL066235

Pokorný V (2021) Polyrhythmic arrangements: Rhythm as a dynamic principle in the
constitution of environments. Open Philosophy 4:394–403. https://doi.org/10.1515/opphil-
2020-0192

Rudolf VHW (2019) The role of seasonal timing and phenological shifts for species coexistence.
Ecol Lett 22:1324–1338. https://doi.org/10.1111/ele.13277

Scranton K, Vasseur DA (2016) Coexistence and emergent neutrality generate synchrony among
competitors in fluctuating environments. Theor Ecol 9:353–363.
https://doi.org/10.1007/s12080-016-0294-z

Sharma S, Jackson DA, Minns CK (2009) Quantifying the potential effects of climate change
and the invasion of smallmouth bass on native lake trout populations across Canadian lakes.
Ecography 32:517–525. https://doi.org/10.1111/j.1600-0587.2008.05544.x

Shuter BJ, Finstad AG, Helland IP, et al (2012) The role of winter phenology in shaping the
ecology of freshwater fish and their sensitivities to climate change. Aquat Sci 74:637–657.
https://doi.org/10.1007/s00027-012-0274-3

Stachowicz JJ, Terwin JR, Whitlatch RB, Osman RW (2002) Linking climate change and
biological invasions: Ocean warming facilitates nonindigenous species invasions. Proc Natl
Acad Sci U S A 99:15497–15500. https://doi.org/10.1073/pnas.242437499

Suski CD, Ridgway MS (2009) Winter biology of centrarchid fishes. In: Cooke SJ, Philipp DP
(eds) Centrarchid Fishes. Blackwell Publishing Ltd., pp 264–292

Tavecchia G, Tenan S, Pradel R, et al (2016) Climate-driven vital rates do not always mean
climate-driven population. Glob Chang Biol 22:3960–3966.
https://doi.org/10.1111/gcb.13330

Teitelbaum CS, Fagan WF, Fleming CH, et al (2015) How far to go? Determinants of migration
distance in land mammals. Ecol Lett 18:545–552. https://doi.org/10.1111/ele.12435

Tonkin JD, Bogan MT, Bonada N, et al (2017) Seasonality and predictability shape temporal
species diversity. Ecology 98:1201–1216. https://doi.org/10.1002/ecy.1761

Tredennick AT, Adler PB, Adler FR (2017) The relationship between species richness and
ecosystem variability is shaped by the mechanism of coexistence. Ecol Lett 20:958–968.
https://doi.org/10.1111/ele.12793

Tunney TD, McCann KS, Lester NP, Shuter BJ (2014) Effects of differential habitat warming on
complex communities. Proc Natl Acad Sci U S A 111:8077–8082.
https://doi.org/10.1073/pnas.1319618111

Urban MC, de Meester L, Vellend M, et al (2012) A crucial step toward realism: Responses to
climate change from an evolving metacommunity perspective. Evol Appl 5:154–167.
https://doi.org/10.1111/j.1752-4571.2011.00208.x

Varpe Ø (2017) Life history adaptations to seasonality. Integr Comp Biol 57:943–960.
https://doi.org/10.1093/icb/icx123

Vasconcellos RM, Araújo FG, de Sousa Santos JN, de Araújo Silva M (2011) Diel seasonality in
fish biodiversity in a sandy beach in south-eastern Brazil. Marine Biological Association of
the United Kingdom 91:1337–1344. https://doi.org/10.1017/S0025315410000652

Vihtakari M, Renaud PE, Clarke LJ, et al (2016) Decoding the oxygen isotope signal for
seasonal growth patterns in Arctic bivalves. Palaeogeogr Palaeoclimatol Palaeoecol
446:263–283. https://doi.org/10.1016/j.palaeo.2016.01.008

Warne CPK, McCann KS, Rooney N, et al (2020) Geography and morphology affect the ice
duration dynamics of northern hemisphere lakes worldwide. Geophys Res Lett 47:1–10.
https://doi.org/10.1029/2020GL087953

White ER, Hastings A (2020) Seasonality in ecology: Progress and prospects in theory.
Ecological Complexity 44:1–6. https://doi.org/10.1016/j.ecocom.2020.100867

**Table 1:** Lotka-Volterra Coexistence Conditions vs. Seasonal Coexistence Conditions.

| | Lotka – Volterra Criteria | Seasonal Coexistence Criteria |
|---|---|---|
| Stable Coexistence | $\alpha_{11} > \alpha_{21}$ | $\dfrac{pr_{P,1}\alpha_{P,11} + (1-p)r_{LP,1}\alpha_{LP,11}}{pr_{P,1} + (1-p)r_{LP,1}} > \dfrac{pr_{P,2}\alpha_{P,21} + (1-p)r_{LP,2}\alpha_{LP,21}}{pr_{P,2} + (1-p)r_{LP,2}}$ |
| | $\alpha_{22} > \alpha_{12}$ | $\dfrac{pr_{P,2}\alpha_{P,22} + (1-p)r_{LP,2}\alpha_{LP,22}}{pr_{P,2} + (1-p)r_{LP,2}} > \dfrac{pr_{P,1}\alpha_{P,12} + (1-p)r_{LP,1}\alpha_{LP,12}}{pr_{P,1} + (1-p)r_{LP,1}}$ |
| Competitive Exclusion by $sp_1$ | $\alpha_{11} > \alpha_{21}$ | $\dfrac{pr_{P,1}\alpha_{P,11} + (1-p)r_{LP,1}\alpha_{LP,11}}{pr_{P,1} + (1-p)r_{LP,1}} > \dfrac{pr_{P,2}\alpha_{P,21} + (1-p)r_{LP,2}\alpha_{LP,21}}{pr_{P,2} + (1-p)r_{LP,2}}$ |
| | $\alpha_{22} < \alpha_{12}$ | $\dfrac{pr_{P,2}\alpha_{P,22} + (1-p)r_{LP,2}\alpha_{LP,22}}{pr_{P,2} + (1-p)r_{LP,2}} < \dfrac{pr_{P,1}\alpha_{P,12} + (1-p)r_{LP,1}\alpha_{LP,12}}{pr_{P,1} + (1-p)r_{LP,1}}$ |
| Competitive Exclusion by $sp_2$ | $\alpha_{11} > \alpha_{21}$ | $\dfrac{pr_{P,1}\alpha_{P,11} + (1-p)r_{LP,1}\alpha_{LP,11}}{pr_{P,1} + (1-p)r_{LP,1}} > \dfrac{pr_{P,2}\alpha_{P,21} + (1-p)r_{LP,2}\alpha_{LP,21}}{pr_{P,2} + (1-p)r_{LP,2}}$ |
| | $\alpha_{22} < \alpha_{12}$ | $\dfrac{pr_{P,2}\alpha_{P,22} + (1-p)r_{LP,2}\alpha_{LP,22}}{pr_{P,2} + (1-p)r_{LP,2}} < \dfrac{pr_{P,1}\alpha_{P,12} + (1-p)r_{LP,1}\alpha_{LP,12}}{pr_{P,1} + (1-p)r_{LP,1}}$ |
| Contingent Coexistence | $\alpha_{11} < \alpha_{21}$ | $\dfrac{pr_{P,1}\alpha_{P,11} + (1-p)r_{LP,1}\alpha_{LP,11}}{pr_{P,1} + (1-p)r_{LP,1}} < \dfrac{pr_{P,2}\alpha_{P,21} + (1-p)r_{LP,2}\alpha_{LP,21}}{pr_{P,2} + (1-p)r_{LP,2}}$ |
| | $\alpha_{22} < \alpha_{12}$ | $\dfrac{pr_{P,2}\alpha_{P,22} + (1-p)r_{LP,2}\alpha_{LP,22}}{pr_{P,2} + (1-p)r_{LP,2}} < \dfrac{pr_{P,1}\alpha_{P,12} + (1-p)r_{LP,1}\alpha_{LP,12}}{pr_{P,1} + (1-p)r_{LP,1}}$ |

646     **Figure 1**

**a)** Model Set Up

**b)** Model Experiment

**Fig. 1** Model schematic investigates the projected mean equilibrium density over the length of
the productive season when species compete under different periodic conditions. a) The model
set-up investigates two time-separated seasons with different environmental conditions
(productive (white box) and less productive (gray box)). In numerical simulations, competitive
species, $X_1$ (white fish) and $X_2$ (black fish), exhibit different trade-offs in response to these
differing environmental conditions, where each species may experience different levels of inter-
to intraspecific competitive interactions (thickness of arrows) depending on which species may
have an overall better performance (in terms of growth and competition) in one season compared
to the other. An experiment is projected over many periods (say 1000 time-units) of length $\tau$
until it reaches an asymptotic state (we refer to this as an equilibrium despite the within-year
variation). b) In the model experiment, the projected mean equilibrium density, which represents
a fluctuating' species density at its asymptote, is calculated as the duration of the productive
season (as a proportion of each time unit) varies from 0 – 1.

**Figure 2**

**Fig. 2** Isocline approximation of $X_1$ density on $X_2$ density. a) Stable coexistence isocline
approximation on phaseplane of $X_1$ (dashed line) and $X_2$ (solid line) densities. Approximate
isocline solutions for a given species occur when the instantaneous rate of change (represented
by vector arrows) in each season, scaled by seasonal duration, are equal and opposite of each
other (i.e., they "cancel" each other out such that the net change over a full period is null). Refer
to Table 1 for similarities to Lotka–Volterra coexistence conditions. b i-iii) illustrates a $p$-driven
bifurcation, showing the movement of the isoclines as the productivity season travels from i) $p =$
0.5; coexistence, to ii) $p = 0.6$; transcritical bifurcation, to iii) $p = 0.7$; competitive exclusion of
species 2. Parametric values: $\alpha_{P,11}$=0.44, $\alpha_{P,22}$=0.66, $\alpha_{P,12}$=0.25, $\alpha_{P,21}$=0.53, $\alpha_{LP,11}$=1.11,
$\alpha_{LP,22}$=0.83, $\alpha_{LP,12}$=1.83, $\alpha_{LP,21}$=0.48, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $r_{LP,1}$=0.3, $r_{LP,2}$=1.

**Figure 3**

**Fig. 3** Numerical simulations of mean density over length of productive season ($p$) for
seasonally-mediated outcomes. White zone represents stable coexistence, light gray zone
represents competitive exclusion, and dark gray zone represents contingent coexistence where
the equilibrium is unstable and multiple attractors exist between the two species. a) Seasonally-
mediated coexistence. Parametric values: $\alpha_{P,21}$=0.35, $\alpha_{P,12}$=0.165, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.436,
$r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.385, $\alpha_{LP,12}$=0,805, $\alpha_{LP,11}$=0.73, $\alpha_{LP,22}$=0.55, $r_{LP,1}$=0.3, $r_{LP,2}$=1. b)
Seasonally-mediated competitive exclusion. Parametric values: $\alpha_{P,21}$=0.29, $\alpha_{P,12}$=0.38,
$\alpha_{P,11}$=0.31, $\alpha_{P,22}$=0.44, $r_{P,1}$=4, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.27, $\alpha_{LP,12}$=1.02, $\alpha_{LP,11}$=1,67, $\alpha_{LP,22}$=1.18 $r_{LP,1}$=0.3,
$r_{LP,2}$=1. c) Seasonally-mediated contingent coexistence. Parametric values; $\alpha_{P,21}$=0.548,
$\alpha_{P,12}$=0.33, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.363, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.31, $\alpha_{LP,12}$=1.89, $\alpha_{LP,11}$=1.65,
$\alpha_{LP,22}$=0.66, $r_{LP,1}$=0.3, $r_{LP,2}$=1.

**Figure 4**

**Fig. 4** Two-dimensional bifurcation diagram of seasonally-mediated coexistence. a) seasonally-mediated coexistence expanding over a wide range of period lengths ($\tau$). The red dashed lines represent the isocline approximation's prediction of the two transcritical bifurcation points at $p$=0.115 and $p$=0.69 (see Fig. S3.1 for complete transition across $p$, and S4.1 for explanation of approximation accuracy of seasonally mediated coexistence). b) isocline approximation tracks the approximate equilibrium, which represents the mean asymptotic behavior, as the productive season ($p$) increases from i) $p$=0.1; competitive exclusion of species 1, to ii) $p$=0.4; stable coexistence, to iii) $p$=0.8; competitive exclusion of species 2. Filled in circles are stable equilibrium points, and open circles are unstable equilibrium points. Parametric values: $\alpha_{P,21}$=0.35, $\alpha_{P,12}$=0.165, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.436, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.385, $\alpha_{LP,12}$=0,805, $\alpha_{LP,11}$=0.73, $\alpha_{LP,22}$=0.55, $r_{LP,1}$=0.3, $r_{LP,2}$=1.

**Figure 5**

**Fig. 5** Two-dimensional bifurcation diagram of seasonally-mediated competitive exclusion. a)
seasonally-mediated competitive exclusion expanding over a wide range of period lengths ($\tau$).
The red dashed lines represent the isocline approximation's prediction of the two transcritical
bifurcation points at $p$=0.04 and $p$=0.97 (see Fig. S3.2 for complete transition across $p$, and S4.2
for explanation of approximation accuracy of seasonally mediated competitive exclusion). b)
isocline approximation tracks the approximate equilibrium, which represents the mean
asymptotic behaviour, as the productive season increases from i) $p$=0.0; stable coexistence, to ii)
$p$=0.3; competitive exclusion of species 2, and back to iii) $p$=1.0; stable coexistence. Filled in
circles are stable equilibrium points, and open circles are unstable equilibrium points. Parametric
values: $\alpha_{P,21}$=0.29, $\alpha_{P,12}$=0.38, $\alpha_{P,11}$=0.31, $\alpha_{P,22}$=0.44, $r_{P,1}$=4, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.27, $\alpha_{LP,12}$=1.02,
$\alpha_{LP,11}$=1,67, $\alpha_{LP,22}$=1.18 $r_{LP,1}$=0.3, $r_{LP,2}$=1.

**Figure 6**

**Fig. 6** Two-dimensional bifurcation diagram of seasonally-mediated contingent coexistence. a)
seasonally-mediated contingent coexistence expanding over a wide range of period length ($\tau$).
The red dashed lines represent the isocline approximation's prediction of the two transcritical
bifurcation points at $p$=0.07 and $p$=0.69 (see Fig. S3.3 for complete transition across $p$, and S4.3
for explanation of approximation accuracy of seasonally mediated contingent coexistence). b)
isocline approximation tracks the approximate equilibrium, which represents the mean
asymptotic behaviour, as the productive season increases from i) $p$=0.0; competitive exclusion of
species 1, to ii) $p$=0.4; contingent coexistence, to iii) $p$=0.8; competitive exclusion of species 2.
Filled in circles are stable equilibrium points, and open circles are unstable equilibrium points.
Parametric values; $\alpha_{P,21}$=0.548, $\alpha_{P,12}$=0.33, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.363, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.31,
$\alpha_{LP,12}$=1.89, $\alpha_{LP,11}$=1.65, $\alpha_{LP,22}$=0.66, $r_{LP,1}$=0.3, $r_{LP,2}$=1.

**Coexistence in Periodic Environments: Supplementary Material**

**S1: A Linearization of the Periodic Lotka-Volterra Dynamics**

If we consider any point on the phaseplane ($X_1$-$X_2$) due to the productive season that lasts from 0

to $\tau p$ where $\tau$ is the period length, and $p$ is the proportion of each period that is considered

productive, then we know:

$$X_{P,1}(\tau p) = X_{P,1}(0) + \int_0^{\tau p} \frac{dX_{P,1}}{dt} dt$$

(1)

$$X_{P,2}(\tau p) = X_{P,2}(0) + \int_0^{\tau p} \frac{dX_{P,2}}{dt} dt$$

(2)

731 These dynamics follow the differential equation over the trajectory from 0 to $\tau p$ starting at the

732 values $X_{P,j}(0)$. We linearize the trajectory over 0 to $\tau p$ by assuming the $\frac{dX_{P,j}}{dt}$ remains constant

733 (e.g., we calculate the instantaneous $\frac{dX_{P,j}}{dt}$ for a point in the phaseplane, say for time (0)). As this

734 is now a constant, we can use the Fundamental Theorem of Calculus to solve for Equations (1)

735 and (2) above yielding:

$$X_{P,1}(\tau p) - X_{P,1}(0) = \frac{dX_{P,1}}{dt} \, t \, \Big|_0^{\tau p}$$

(3)

$$X_{P,1}(\tau p) - X_{P,1}(0) = \tau p \frac{dX_{P,1}}{dt} - 0 \frac{dX_{P,1}}{dt}$$

(4)

$$X_{P,1}(\tau p) - X_{P,1}(0) = \tau p \frac{dX_{P,1}}{dt}$$

(5)

736 Similarly, we solve for the less productive season between $\tau p$ and $\tau$, giving us:

$$X_{LP,1}(\tau) - X_{LP,1}(\tau p) = \tau(1-p)\frac{dX_{LP,1}}{dt}$$

$$(6)$$

737     Thus, the linearization estimates the trajectory as a linear scale movement following the length of

738     the time interval (either $\tau p$, or $\tau(1-p)$). We can take this simple estimate to estimate the within

739     seasonal dynamics and use them to estimate the periodic 0-isoclines.

741     Given Equations (5) and (6), the $X_1$ isocline occurs when:

$$\tau p \frac{dX_{P,1}}{dt} = -\tau(1-p)\frac{dX_{LP,1}}{dt} \qquad (7)$$

743     With this approximation, we substitute the productive and less productive Lotka-Volterra models

744     into equation (7):

$$\tau p \, r_{P,1}X_1(1 - \alpha_{P,11}X_1 - \alpha_{P,12}X_2) = -\tau(1-p) \, r_{LP,1}X_1(1 - \alpha_{LP,11}X_1 - \alpha_{LP,12}X_2) \quad (8)$$

746     Noticing that $\tau$ cancels out, we can now solve for the isocline solution:

$$X_1 = -\frac{pr_{P,1}\alpha_{P,12} + (1-p)r_{LP,1}\alpha_{LP,12}}{pr_{P,1}\alpha_{P,11} + (1-p)r_{LP,1}\alpha_{LP,11}}X_2 + \frac{pr_{P,1} + (1-p)r_{LP,1}}{pr_{P,1}\alpha_{P,11} + (1-p)r_{LP,1}\alpha_{LP,11}} \qquad (9)$$

748     Similarly, we would perform the same steps to find the isocline solution for $X_2$. This isocline

749     approximation, for both species, allows us to determine coexistence criteria for seasonal

750     environments.

**S2: Isocline Approximation and Numerical Simulation Comparison**

**Fig. S2.1** Comparison of Isocline Approximation against Numerical Simulations. White zone represents stable coexistence, light gray zone represents competitive exclusion, and dark gray zone represents contingent coexistence. a) Seasonally mediated coexistence. Parametric values: $\alpha_{P,21}$=0.35, $\alpha_{P,12}$=0.165, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.436, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.385, $\alpha_{LP,12}$=0,805, $\alpha_{LP,11}$=0.73, $\alpha_{LP,22}$=0.55, $r_{LP,1}$=0.3, $r_{LP,2}$=1. b) Seasonally mediated competitive exclusion. Parametric values: $\alpha_{P,21}$=0.29, $\alpha_{P,12}$=0.38, $\alpha_{P,11}$=0.31, $\alpha_{P,22}$=0.44, $r_{P,1}$=4, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.27, $\alpha_{LP,12}$=1.02, $\alpha_{LP,11}$=1,67, $\alpha_{LP,22}$=1.18 $r_{LP,1}$=0.3, $r_{LP,2}$=1. c) Seasonally mediated contingent coexistence. The equilibrium is unstable in the dark gray zone and multiple attractors exist between the two species. Parametric values: $\alpha_{P,21}$=0.548, $\alpha_{P,12}$=0.33, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.363, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.31, $\alpha_{LP,12}$=1.89, $\alpha_{LP,11}$=1.65, $\alpha_{LP,22}$=0.66, $r_{LP,1}$=0.3, $r_{LP,2}$=1. d) Counterintuitive mean density change. Parametric values: $\alpha_{P,21}$=0.2, $\alpha_{P,12}$=0.155, $\alpha_{P,11}$=0.315, $\alpha_{P,22}$=0.335, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.57, $\alpha_{LP,12}$=0.4, $\alpha_{LP,11}$=1, $\alpha_{LP,22}$=0.715, $r_{LP,1}$=0.3, $r_{LP,2}$=1. e) Mean density changes with competitive exclusion in the less productive season. Parametric values: $\alpha_{P,21}$=0.288, $\alpha_{P,12}$=0.15, $\alpha_{P,11}$=0.336, $\alpha_{P,22}$=0.402, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.354, $\alpha_{LP,12}$=0.75, $\alpha_{LP,11}$=0.666, $\alpha_{LP,22}$=0.462, $r_{LP,1}$=0.3, $r_{LP,2}$=1. f) Mean density changes with competitive exclusion in the productive season. Parametric values: $\alpha_{P,21}$=0.35, $\alpha_{P,12}$=0.165, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.44, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.28, $\alpha_{LP,12}$=0.825, $\alpha_{LP,11}$=0.73, $\alpha_{LP,22}$=0.94, $r_{LP,1}$=0.3, $r_{LP,2}$=1.

**Fig. S2.2** Time Series at $p = 0.5$, and $\tau = 1$ for the comparison of isocline approximation against numerical simulations. a) Seasonally mediated coexistence. Parametric values: $\alpha_{P,21}$=0.35, $\alpha_{P,12}$=0.165, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.436, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.385, $\alpha_{LP,12}$=0,805, $\alpha_{LP,11}$=0.73,

$\alpha_{LP,22}$=0.55, $r_{LP,1}$=0.3, $r_{LP,2}$=1. b) Seasonally mediated competitive exclusion. Parametric values:

$\alpha_{P,21}$=0.29, $\alpha_{P,12}$=0.38, $\alpha_{P,11}$=0.31, $\alpha_{P,22}$=0.44, $r_{P,1}$=4, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.27, $\alpha_{LP,12}$=1.02,

$\alpha_{LP,11}$=1,67, $\alpha_{LP,22}$=1.18 $r_{LP,1}$=0.3, $r_{LP,2}$=1. ci) Seasonally mediated contingent coexistence. $X_1$

competitively excluded $X_2$. Parametric values: $\alpha_{P,21}$=0.548, $\alpha_{P,12}$=0.33, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.363,

$r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.31, $\alpha_{LP,12}$=1.89, $\alpha_{LP,11}$=1.65, $\alpha_{LP,22}$=0.66, $r_{LP,1}$=0.3, $r_{LP,2}$=1. cii)

Seasonally mediated contingent coexistence. $X_2$ competitively excluded $X_1$. Parametric values:

$\alpha_{P,21}$=0.548, $\alpha_{P,12}$=0.33, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.363, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.31, $\alpha_{LP,12}$=1.89,

$\alpha_{LP,11}$=1.65, $\alpha_{LP,22}$=0.66, $r_{LP,1}$=0.3, $r_{LP,2}$=1. d) Counterintuitive mean density change. Parametric

values: $\alpha_{P,21}$=0.2, $\alpha_{P,12}$=0.155, $\alpha_{P,11}$=0.315, $\alpha_{P,22}$=0.335, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.57, $\alpha_{LP,12}$=0.4,

$\alpha_{LP,11}$=1, $\alpha_{LP,22}$=0.715, $r_{LP,1}$=0.3, $r_{LP,2}$=1. e) Mean density changes with competitive exclusion in

the less productive season. Parametric values: $\alpha_{P,21}$=0.288, $\alpha_{P,12}$=0.15, $\alpha_{P,11}$=0.336, $\alpha_{P,22}$=0.402,

$r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.354, $\alpha_{LP,12}$=0.75, $\alpha_{LP,11}$=0.666, $\alpha_{LP,22}$=0.462, $r_{LP,1}$=0.3, $r_{LP,2}$=1. f) Mean

density changes with competitive exclusion in the productive season. Parametric values:

$\alpha_{P,21}$=0.35, $\alpha_{P,12}$=0.165, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.44, $r_{P,1}$=1.7, $r_{P,2}$=1.2, $\alpha_{LP,21}$=0.28, $\alpha_{LP,12}$=0.825,

$\alpha_{LP,11}$=0.73, $\alpha_{LP,22}$=0.94, $r_{LP,1}$=0.3, $r_{LP,2}$=1.

**S3: Tracking Isocline Approximation across Productivity Length for all Seasonally**

**Mediated Competitive Outcomes**

**Fig. S3.1** Seasonally mediated coexistence isocline approximation. Isocline approximation
tracking the interior equilibrium as the productive season increases in length from a) $p=0.1$;
competitive exclusion of species 1, to b) $p=0.115$; transcritical bifurcation, to c) $p=0.4$; stable
coexistence, to d) $p=0.69$; transcritical bifurcation, to e) $p=0.8$; competitive exclusion of species
2. Parametric values: $\alpha_{P,21}=0.35$, $\alpha_{P,12}=0.165$, $\alpha_{P,11}=0.33$, $\alpha_{P,22}=0.436$, $r_{P,1}=1.7$, $r_{P,2}=1.2$,
$\alpha_{LP,21}=0.385$, $\alpha_{LP,12}=0,805$, $\alpha_{LP,11}=0.73$, $\alpha_{LP,22}=0.55$, $r_{LP,1}=0.3$, $r_{LP,2}=1$.

**Fig. S3.2** Seasonally mediated competitive exclusion isocline approximation. Isocline
approximation tracking the interior equilibrium as the productive season increases in length from
a) $p$=0.0; stable coexistence, to b) $p$=0.04; transcritical bifurcation, to c) $p$=0.3; competitive
exclusion of species 2, to d) $p$=0.97; transcritical bifurcation, to e) $p$=1.0; stable coexistence.
Parametric values: $\alpha_{P,21}$=0.29, $\alpha_{P,12}$=0.38, $\alpha_{P,11}$=0.31, $\alpha_{P,22}$=0.44, $r_{P,1}$=4, $r_{P,2}$=1.2, $\alpha_{LP,21}$=1.27,
$\alpha_{LP,12}$=1.02, $\alpha_{LP,11}$=1,67, $\alpha_{LP,22}$=1.18 $r_{LP,1}$=0.3, $r_{LP,2}$=1.

**Fig. S3.3** Seasonally mediated contingent coexistence isocline approximation. Isocline
approximation tracking the interior equilibrium as the productive season increases in length from
a) $p$=0.0; competitive exclusion of species 1, to b) $p$=0.07; transcritical bifurcation, to c) $p$=0.4;
contingent coexistence, to d) $p$=0.69; transcritical bifurcation, to e) $p$=0.8; competitive exclusion
of species 2. Parametric values: $\alpha_{P,21}$=0.548, $\alpha_{P,12}$=0.33, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.363, $r_{P,1}$=1.7,
$r_{P,2}$=1.2, $\alpha_{LP,21}$=1.31, $\alpha_{LP,12}$=1.89, $\alpha_{LP,11}$=1.65, $\alpha_{LP,22}$=0.66, $r_{LP,1}$=0.3, $r_{LP,2}$=1.

## S4: Accuracy of the Linear Isocline Approximation

Our results hold for a broad range of period lengths but clearly depend on specific parameterizations that alter the timescale or the effects of timescale. Recall that our approximation operates by assuming a zero population growth rate exists in the phaseplane where linearization's of the Lotka-Volterra equations cancel out over the high growth and low growth periods (i.e., linearization of the high growth for species $X_1$ and species $X_2$ are both equal and opposite in sign to the low growth linearization). Given this assumption, we can immediately ask when we expect nonlinear dynamics to dominate and potentially threaten the validity of the linear assumption. First, the longer the period of the forced parameters ($\tau$), then the longer the time the dynamics have to fall off the linear assumption. Further, the larger the growth rates ($r$), the larger the potential for nonlinear dynamics even with smaller periods ($\tau$). As a result, we can say the larger the product $r\tau$, the more likely our assumption of linearity is threatened. Effectively, $\tau r$ sets the relative pace of the seasonal dynamics.

Recognizing that linear assumptions will lose accuracy when dynamics become nonlinear with longer periods and fast growth rates (i.e., there is more time for dynamics to become nonlinear), we first compare the dynamics (set by the original growth rates in the manuscript) with the approximation's accuracy when the period length ($\tau$) is increased. Next, we slow the dynamics by dividing all species' growth rates ($r$) by 10-units and compare the approximation's accuracy with these dynamics when the period length increases. Our results below show that indeed our approximation can fall off, but even here for large, combined values of $r\tau$, the approximation remains a reasonable qualitative predictor of steady state behaviour for seasonally-mediated

coexistence and contingent coexistent (S4.1 and S4.3 respectively). We note that the seasonally

mediated competitive exclusion is not as robust (S4.2).

As seen in Fig. 4a, the analytical approximation still accurately predicts the numerically

generated transcritical bifurcation at $p = 0.11$ (mean densities transition between competitive

exclusion of species 1 to coexistence of both species), even when the period length ($\tau$), reaches

100 time-units. This is due to the large difference between the productive and less-productive

growth rates of species 1. However, the approximation begins to inaccurately predict where the

second numerically generated transcritical bifurcation will occur ($p = 0.69$; mean density

transition between coexistence and competitive exclusion of species 2) when $\tau = 5$ time-units

(Fig. S4.11b). With the original growth rates used in the manuscript, when the period length is

small ($\tau = 1$), the seasonal dynamics are relatively linear, and the approximation accurately

predict the true mean densities from the numerical simulation (Figs. S4.11a and S4.12a).

However, as we increase the period length to 5 time-units, the dynamics become more nonlinear

as, with enough time, species' densities reach their seasonal equilibrium and spend more time at

these fixed points (Fig. S4.12b). Here, the numerically generated mean densities will be skewed

closer towards this equilibrium, while the linear approximation fails to capture this (Fig. S4.11b).

When all growth rates are slowed down (all $r$'s divided by 10 units), the seasonal dynamics

become more linear, and the approximation is more accurately able to track the mean densities

even when the period length ($\tau$) is increased from 1 (Figs S4.11c and S4.12c) to 5 time-units

(Figs S4.11d and S4.12d).

In Fig. 5a, the approximation still predicts seasonally mediated competitive exclusion, even

when species coexist at very large period lengths ($\tau > 33$ time units) across all $p$-values.

However, when $\tau$ is increased from 1 to 10 time-units with the original growth rates, the

dynamics become more nonlinear (Figs. S4.22a and S4.22b respectively) and the approximation

begins to fail at tracking the numerically-generated mean densities throughout the entire range of

$p$-values (Figs. S4.21a and S4.21b respectively), though it still captures the qualitative behavior

and presence of the bifurcations. When growth rates are smaller, the seasonal dynamics are now

more linear (Figs. S4.22c and FigsS4.22d), and the approximation more accurately tracks the

numerically generated mean densities as $\tau$ increases from 1 to 10 time-units (Figs. S4.22c and

FigsS4.22d respectively).

For seasonally mediated contingent coexistence, as the period length ($\tau$) increases, the region of

contingent coexistence from the numerically-generated results first broadens and then shrinks

(Fig. 6a). With the original growth rates, as the period length is increased from 1 to 20 time-

units, the seasonal dynamics become more nonlinear (Figs. S4.32a and S4.32b respectively), and

the approximation is unable to track the true bifurcations based on our numerical results (Figs.

S4.31a and S4.31b respectively). When the growth rates are a tenth of their original size (all $r$'s

divided by 10 units), the dynamics slow down and are now more linear as $\tau$ is increased (Figs.

S4.32c and S4.32d), allowing the approximation to more accurately track the numerically-

generated mean densities over the entire range of $p$-values (Figs. S4.31c and S4.31d).

**S4.1 Approximation Accuracy of Seasonally Mediated Coexistence**

**Fig. S4.11** Isocline approximation – mean density plots over the length of the productive season
($p$) for seasonally-mediated coexistence. White zones represent coexistence and light grey zones
represent competitive exclusion. a) and b) original growth rates (those used throughout
manuscript: $r_{P1}$ = 1.7, $r_{P2}$ = 1.2, $r_{LP1}$ = 1, $r_{LP2}$ = 0.3) when the period length $\tau$ = 1 and 5 time-
units respectively. c) and d) all growth rates have been divided by 10 units ($r_{P1}$ = 0.17, $r_{P2}$ = 0.12,
$r_{LP1}$ = 0.1, $r_{LP2}$ = 0.03) when the period length = 1 and 5 time-units respectively. Parametric
values: $\alpha_{P,21}$=0.35, $\alpha_{P,12}$=0.165, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.436, $\alpha_{LP,21}$=0.385, $\alpha_{LP,12}$=0,805, $\alpha_{LP,11}$=0.73,
$\alpha_{LP,22}$=0.55.

**Fig. S4.12** Isocline approximation – time series for seasonally-mediated coexistence. Length of productive season ($p$) = 0.5 represented by white zones, and light grey zones represent the less-productive season. a) and b) original growth rates (those used throughout manuscript: $r_{P1}$ = 1.7, $r_{P2}$ = 1.2, $r_{LP1}$ = 1, $r_{LP2}$ = 0.3) when the period length $\tau$ = 1 and 5 time-units respectively. c) and d) all growth rates have been divided by 10 units ($r_{P1}$ = 0.17, $r_{P2}$ = 0.12, $r_{LP1}$ = 0.1, $r_{LP2}$ = 0.03) when the period length = 1 and 5 time-units respectively. Parametric values: $\alpha_{P,21}$=0.35, $\alpha_{P,12}$=0.165, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.436, $\alpha_{LP,21}$=0.385, $\alpha_{LP,12}$=0,805, $\alpha_{LP,11}$=0.73, $\alpha_{LP,22}$=0.55.

## S4.2 Approximation Accuracy of Seasonally Mediated Competitive Exclusion

**Fig. S4.21** Isocline approximation – mean density plots over the length of the productive season (*p*) for seasonally-mediated competitive exclusion. White zones represent coexistence and light grey zones represent competitive exclusion. a) and b) original growth rates (those used throughout manuscript: $r_{P1} = 4$, $r_{P2} = 1.2$, $r_{LP1} = 1$, $r_{LP2} = 0.3$) when the period length $\tau = 1$ and 10 time-units respectively. c) and d) all growth rates have been divided by 10 units ($r_{P1} = 0.4$, $r_{P2} = 0.12$, $r_{LP1} = 0.1$, $r_{LP2} = 0.03$) when the period length = 1 and 10 time-units respectively. Parametric values: $\alpha_{P,21}=0.29$, $\alpha_{P,12}=0.38$, $\alpha_{P,11}=0.31$, $\alpha_{P,22}=0.44$, $\alpha_{LP,21}=1.27$, $\alpha_{LP,12}=1.02$, $\alpha_{LP,11}=1,67$, $\alpha_{LP,22}=1.18$.

**Fig. S4.22** Isocline approximation – time series for seasonally-mediated competitive exclusion.
Length of productive season ($p$) = 0.5 represented by white zones, and light grey zones represent
the less-productive season. a) and b) original growth rates (those used throughout manuscript: $r_{P1}$
= 4, $r_{P2}$ = 1.2, $r_{LP1}$ = 1, $r_{LP2}$ = 0.3) when the period length $\tau$ = 1 and 10 time-units respectively. c)
and d) all growth rates have been divided by 10 units ($r_{P1}$ = 0.4, $r_{P2}$ = 0.12, $r_{LP1}$ = 0.1, $r_{LP2}$ =
0.03) when the period length = 1 and 10 time-units respectively. Parametric values: $\alpha_{P,21}$=0.29,
$\alpha_{P,12}$=0.38, $\alpha_{P,11}$=0.31, $\alpha_{P,22}$=0.44, $\alpha_{LP,21}$=1.27, $\alpha_{LP,12}$=1.02, $\alpha_{LP,11}$=1,67, $\alpha_{LP,22}$=1.18.

 **S4.3 Approximation Accuracy of Seasonally Mediated Contingent Coexistence**

**Fig. S4.31** Isocline approximation – mean density plots over the length of the productive season
(*p*) for seasonally-mediated contingent coexistence. Light grey zones represent competitive
exclusion and grey zones represent contingent coexistence. a) and b) original growth rates (those
used throughout manuscript: $r_{P1}$ = 1.7, $r_{P2}$ = 1.2, $r_{LP1}$ = 1, $r_{LP2}$ = 0.3) when the period length $\tau$ =
1 and 20 time-units respectively. c) and d) all growth rates have been divided by 10 units ($r_{P1}$ =
0.17, $r_{P2}$ = 0.12, $r_{LP1}$ = 0.1, $r_{LP2}$ = 0.03) when the period length = 1 and 20 time-units
respectively. Parametric values: $\alpha_{P,21}$=0.548, $\alpha_{P,12}$=0.33, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.363, $\alpha_{LP,21}$=1.31,
$\alpha_{LP,12}$=1.89, $\alpha_{LP,11}$=1.65, $\alpha_{LP,22}$=0.66.

**Fig. S4.32** Isocline approximation – time series for seasonally-mediated contingent coexistence. Length of productive season ($p$) = 0.5 represented by white zones, and light grey zones represent the less-productive season. a) and b) original growth rates (those used throughout manuscript: $r_{P1}$ = 1.7, $r_{P2}$ = 1.2, $r_{LP1}$ = 1, $r_{LP2}$ = 0.3) when the period length $\tau$ = 1 and 20 time-units respectively. c) and d) all growth rates have been divided by 10 units ($r_{P1}$ = 0.17, $r_{P2}$ = 0.12, $r_{LP1}$ = 0.1, $r_{LP2}$ = 0.03) when the period length = 1 and 20 time-units respectively. Parametric values: $\alpha_{P,21}$=0.548, $\alpha_{P,12}$=0.33, $\alpha_{P,11}$=0.33, $\alpha_{P,22}$=0.363, $\alpha_{LP,21}$=1.31, $\alpha_{LP,12}$=1.89, $\alpha_{LP,11}$=1.65, $\alpha_{LP,22}$=0.66.