Cortical cell assemblies and their underlying connectivity: an *in silico* study

András Ecker[1*], Daniela Egas Santander[1*], Sirio Bolaños-Puchet[1], James B. Isbister[1], and Michael W. Reimann[1]

[1]Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Campus Biotech, Geneva, Switzerland

[*]Co-lead authors

Abstract

Recent developments in experimental techniques have enabled simultaneous recordings from thousands of neurons, enabling the study of functional cell assemblies. However, determining the patterns of synaptic connectivity giving rise to these assemblies remains challenging. To address this, we developed a complementary, simulation-based approach, using a detailed, large-scale cortical network model. Using a combination of established methods we detected functional cell assemblies from the stimulus-evoked spiking activity of 186,665 neurons. We studied how the structure of synaptic connectivity underlies assembly composition, quantifying the effects of thalamic innervation, recurrent connectivity, and the spatial arrangement of synapses on dendrites. We determined that these features reduce up to 30%, 22%, and 10% of the uncertainty of a neuron belonging to an assembly. The detected assemblies were activated in a stimulus-specific sequence and were grouped based on their position in the sequence. We found that the different groups were affected to different degrees by the structural features we considered. Additionally, connectivity was more predictive of assembly membership if its direction aligned with the temporal order of assembly activation, if it originated from strongly interconnected populations, and if synapses clustered on dendritic branches. In summary, reversing Hebb's postulate, we showed how cells that are wired together, fire together, quantifying how connectivity patterns interact to shape the emergence of assemblies. This includes a qualitative aspect of connectivity: not just the amount, but also the local structure matters; from the subcellular level in the form of dendritic clustering to the presence of specific network motifs. This connectivity-based characterization of cell assemblies creates an opportunity to study plasticity at the assembly level, and beyond strictly pairwise interactions.

1 Introduction

The formulation of the cell assemblies concept goes back to Hebb (1949), who defined them loosely as *"a diffuse structure comprising cells in the cortex"*. In the past 70 years, the sequential activation of groups of neurons, the Hebbian *"phase sequence"* was linked to several complex cognitive processes, reviewed in Harris (2005) and Buzsáki (2010). Hebb's idea was later paraphrased as *"cells that fire together, wire together"* (Shatz, 1992), giving it both a structural, and a functional side. In this article we will concentrate on quantifying how the cortical structure underlies its neurons' co-firing function, but linking these groups of co-active neurons to cognitive processes is outside of our scope.

Cell assembly research rejuvenated in the hippocampus field when spikes could be reliably sorted from recordings with tetrodes and therefore neurons could be grouped to co-firing ensembles (Harris et al., 2003; Dragoi and Buzsáki, 2006; Sasaki et al., 2006; Lopes-dos Santos et al., 2013). The introduction of modern *in vivo* two-photon calcium imaging into the field, with its improved scalability and stability over time, allowed Bathellier et al. (2012) and Carrillo-Reid et al. (2015) to detect cell assemblies in auditory and visual cortices, where they showed how even a small set of them can serve as a backbone for cortical coding. These, and studies that followed (Miller et al., 2014; Montijn et al., 2016; Pérez-Ortega et al., 2021) contributed greatly to our understanding of the functional role of the Hebbian cell assemblies, but they could not make claims about the patterns of synaptic connectivity they originate from, as they could only predict functional connectivity from correlations in neuronal activity, but did not have access to the underlying structural connectivity of the neurons recorded. Additionally, results based on calcium imaging are limited to the superficial layers of the cortex, missing potential assemblies in the deeper layers, which would be of great interest as they serve as the output of the cortex (Feldmeyer, 2012; Harris and Shepherd, 2015).

Early theoretical work in the field explored the potential link between memories and cells that fire and therefore wire together, concentrating on the storage and retrieval of memories in strongly recurrent networks, such as the CA3 area of the hippocampus (Hopfield, 1982). Theories evolved and improved, but modeling studies about cell assemblies still concentrate on plasticity rules underlying the learning, storage and recall of various patterns (Krotov and Hopfield, 2016; Fauth and Van Rossum, 2019; Kossio et al., 2021; Gastaldi et al., 2021). Thus, their focus lies on how function shapes structure, with little or no emphasis on the biologically accurate aspects of structural connectivity, such as low connection probabilities and an abundance of directed motifs (Song et al., 2005; Perin et al., 2011; Reimann et al., 2017b).

On the other hand, the perspective can be reversed: how does a more bio-realistic structural connectivity influences a neuron's membership in one or more assemblies, or on a more general level: how does structure determine function? Additionally, how does innervation from different sources, such as local connectivity and various thalamic afferents, interact to shape assembly membership? Finally, are the afferent synapses from fellow assembly neurons scattered across the dendritic tree, or clustered on single branches, employing the nonlinear computational capabilities of dendrites (Poirazi et al., 2003; Stuart and Spruston, 2015; Kastellakis and Poirazi, 2019).

In order to provide insights into these questions, we employed an *in silico* approach, using an improved version of the detailed, large-scale (somatosensory) cortical circuit model of Markram et al. (2015) (Figure 1A), simulating the activity of tens of thousands neurons in response to a stream of thalamic input patterns. In the model, we have access not only to the spiking activity of every neuron, but also to the entire connectome, including dendritic locations of synapses. We then considered the established, purely functional definition of cell assemblies as neurons that fire together more than expected. Therefore, functional assemblies across all cortical layers were detected using a combination of previously published methods (Pérez-Ortega et al., 2021; Herzog

Figure 1: Pipeline summary. A: Schematics and quick facts about the detailed, large-scale cortical microcircuit that was used several times before to study the relationship of cortical structure and function (Reimann et al., 2017b, 2022c; Nolte et al., 2020). **B:** Schematics of the assembly detection pipeline from the spiking activity of 186,665 excitatory neurons in the circuit. **C:** Analysis of the connectivity of cell assembly neurons. **D:** Derivation of assembly membership probability based on different features of stuctural connectivity.

et al., 2021) (Figure 1B). We then analyzed their underlying structural connectivity, searching for rules that could explain assembly membership (Figure 1C-D). This analysis of the structure-function relation could be readily applied to assemblies detected with other methods, such as Lopes-dos Santos et al. (2013); Isbister et al. (2021); van der Plas et al. (2023).

We found that the structural strengths of afferents from various sources explained significant portions of the uncertainty of a neuron's membership in an assembly (Figure 1D). Specifically, innervation from VPM (ventral posteriomedial nucleus of the thalamus) explained up to 30%, and up to 10% from POm (posterior medial nucleus of the thalamus). Strength of innervation through recurrent local connectivity explained up to 22%. The relative magnitudes of these effects differed between assemblies, with assemblies active immediately upon stimulus presentation being mostly determined by the structure of the thalamic innervation, and assemblies active 50 ms after stimulus onset being mostly determined by recurrent connectivity. Additionally, the effect of innervation strength on assembly membership can be much larger if the innervating population is highly interconnected within itself, and if the synapses are tightly clustered on the targeted dendrites (Figure 1D). The highly non-random structure of this connectome provides a more efficient wiring; implementing ensembles of reliably co-firing neurons with fewer synaptic connections than expected by chance.

2 Results

2.1 Presence of cell assemblies in network simulations

We simulated the electrical activity of a model of $2.4mm^3$ of cortical tissue, comprising 211,712 neurons in all cortical layers in an *in vivo*-like state. The model is a version of (Markram et al., 2015) with the anatomical improvements outlined in (Reimann et al., 2022b) (Figure 1A). We consider the activity to be *in vivo*-like, based on a comparison of the ratios of spontaneous firing rates of sub-populations, and responses to brief thalamic inputs to *in vivo* results from (Reyes-Puerta et al., 2015). A stream of thalamic input patterns was applied to the model (see

Methods), and the neuronal responses recorded (Figure 2A1). The circuit reliably responded to the brief stimuli with a transient increase in firing rate. This led to a slight shift to the right of the tail of the firing rate distribution from the spontaneous state (Figure 2A2), in line with (Wohrer et al., 2013). The stream consisted of repeated presentations of ten different input patterns in random order, similar to Reimann et al. (2022c). We designed the stimuli as 10 patterns with varying degrees of overlap (Figure 2B): 4 base patterns with no overlap (A, B, C, D), 3 patterns as combinations of two of the base ones (E, F, G), 2 patterns as combinations of three of the base ones (H, I), and 1 pattern as a combination of all four base ones (J). The overlap of these patterns can also be seen through the raster plots of their corresponding VPM fibers (Figure 2B bottom). For example the fibers corresponding to pattern A peak when stimulus A is presented, but also with 50% of the amplitude when E is presented.

Figure 2: *In vivo-like activity in silico.* **A1:** Raster plot of the microcircuit's activity with 628,620 spikes from 98,059 individual neurons and the population firing rates below. The y-axis shows cortical depth. (As cortical layers do not have the same cell density, the visually densest layer is not necessarily the most active - see A2 bottom.) **A2:** Single cell firing rates (in excitatory and 3 classes of inhibitory cells) and layer-wise inhibitory and excitatory population firing rates in evoked (showed in A1) and spontaneous (not shown) activity. **B:** Top: pyramid-like overlap setup of VPM patterns, then the centers of the VPM fibers in flat map space. Bottom: raster plots of VPM fibers forming each of the patterns for the stimulus stream in A1 (i.e., from pattern A at 2000 ms to pattern J at 6500 ms). On the right: same for non-specific (POm) input.

Using a combination of the algorithms of Pérez-Ortega et al. (2021) and Herzog et al. (2021), we detected functional assemblies in 125-second-long recordings of simulated neuron activity while receiving the random input stream (25 repetitions of all 10 patterns with 500 ms inter-stimulus interval, see Methods). Briefly, the assembly detection algorithm first groups neuronal activity into 20 ms time bins, and identifies those with significantly increased firing rates (Sasaki

et al., 2006; Carrillo-Reid et al., 2015) (see Methods, Figure 3A). Then, these time bins are hierarchically clustered based on the cosine similarity of their activation vector, i.e., the vector of the number of spikes fired in the time bin for each neuron (Montijn et al., 2016; Pérez-Ortega et al., 2021) (Supplementary Figure S1A, Figure 3B1). The threshold for cutting the clustering tree into clusters is determined by minimizing the resulting Davies-Bouldin index (Davies and Bouldin, 1979) (see Methods, Supplementary Figure S2 and Supplementary Figure S1B for lower dimensional representations). Finally, these clusters correspond to the functional assemblies, with a neuron being considered a member if its spiking activity correlates with the activity of an assembly significantly more strongly than chance level (Montijn et al., 2016; Herzog et al., 2021). This means that in each time bin only a single assembly is considered active, but neurons can be part of several assemblies (see Methods, Figure 1B, Figure 3B2).

We found that assemblies were activated in all stimulus repetitions and a series of two to three assemblies remained active for $110 \pm 30 ms$ (Figure 3B2). Activation probability and duration depended on stimulus identity; the stimulus associated with the strongest response elicited 1.6 times as many significant time bins than the stimulus associated with the weakest response. Not only were individual assemblies associated with only a subset of stimuli, but they also had a well-preserved temporal order, with some of them appearing early during a stimulus, and others later. Based on this, from now on we will refer to them as *early, middle*, and *late assemblies*, and will order them in the figures accordingly.

An assembly comprised on average $7\pm3.1\%$ of the simulated excitatory neurons, with late assemblies being about 3.7 times larger than early ones (Figure 3C). In terms of spatial distribution of assembly neurons, the early assemblies appear more *"patchy"* (small distinguishable clusters), and by visual inspection can be mapped back to the locations of VPM fibers corresponding to the stimuli that activated them (Figures 2B and 3C top). Moreover, their layer profile mimics that of VPM fiber innervation (Meyer et al., 2010; Reimann et al., 2022b) (Supplementary Figure S4A), indicating that these assemblies may be determined by direct thalamic innervation. On the other hand, the late assembly neurons are more evenly distributed, and cover the entire surface of the simulated circuit, even beyond the range of VPM fiber centers, and are found mostly in deeper layers of the cortex. Middle assembly neurons are somewhat in-between, both in spatial distribution and depth profile. Although we found late assemblies to be nonspecific (at the chosen clustering threshold), early and middle assemblies belonging to the same stimulus occupy similar regions in space and can be shown to have a relatively high (25%) overlap of neurons (Figure 3D left). This also means that single neurons belong to several assemblies (up to 7 out of 11, Figure 3D right). In conclusion, the time course of stimulus responses are well-preserved and reliable enough (although with some temporal jitter) to be simplified into a sequence of distinct sets of functional cell assemblies.

Although stimulus-specific, the time courses are not unique (see the responses to patterns H and I in Figure 3B2). Thus, we wondered to what degree is the overlap of assemblies associated with the overlap of the input patterns given as stimuli? When we compared the distances between the locations of the VPM fibers making up an input pattern, and the assembly sequences detected from the network activity, we found a significant linear trend, i.e., patterns that are close in the input space (e.g. H and I) are close in the *"output space"* defined as the counts of individual assemblies popping up for a given stimuli across repetitions (3E). Thus, the activation sequence of cell assemblies can be seen as a low-dimensional representation of the complex, high-dimensional activity of the circuit's response to different stimuli. The data points are highly variable for mid-sized input distances, and the linear trend gets weaker with increasing number of assemblies (Supplementary Figure S2C). In summary, increasing the number of assemblies by cutting the clustering tree differently improves the separation of inputs at the cost of reducing the correlation between input and output distance. As our aim was not to build an ideal decoder

Figure 3: Cell assembly detection. A: Population firing rate of excitatory neurons with the determined significance threshold. **B1:** Hierarchical clustering of the cosine similarity matrix of activation vectors of significant time bins (above threshold in A, see Methods). **B2:** Clustered significant time bins ordered by patterns presented. **C**: Number and location of neurons in each cell assembly: flat map view on top, depth-profile below. **D:** Jaccard similarity of cell assemblies and number of neurons participating in different number of assemblies. **E**: Input-output map: Input distance is calculated as the Earth mover's distance of the VPM fiber locations (see Figure 2B and Methods), while the output distance is the (normalized) Euclidean distance of pattern evoked time bin cluster counts (counts of different colors in the matrices above in B2, Methods).

of input patterns, in the following steps we analyze the assemblies resulting from a clustering that minimized the Davis-Bouldin index (Supplementary Figure S2A2).

2.1.1 Restricting assembly detection to superficial layers

Due to the limitations of traditional two-photon microscopy, *in vivo* experiments in the cortex can only detect cell assemblies in layer 2/3 (L2/3) (Bathellier et al., 2012; Miller et al., 2014; Carrillo-Reid et al., 2015; Montijn et al., 2016; Pérez-Ortega et al., 2021), whereas *in silico* we detect assemblies across all layers. This begs the question: are we detecting different assemblies,

or do they just cover more depth but contain the L2/3 ones? To answer this question, we used the same data and methodology as before, but restricted our analysis to L2/3 neurons only (Supplementary Figure S3). When detecting assemblies exclusively from the spiking activity of L2/3 pyramidal cells, we got the same results overall, but with some specific differences. First, we found significant time bins for a shorter range (up to 100 ms, which corresponds to our definition of early and middle assembly time windows). And second, stimuli could be distinguished better, e.g. stimuli H and I correspond to different early assemblies in L2/3 (compare Figure 3B2 and Supplementary Figure S3A2). We compared the L2/3 assemblies to the original full assemblies and found that the early ones can get mapped to the L2/3 ones relatively well, in terms of the Jaccard similarity of their respective intersections with L2/3 neurons, and by visual inspection of the spatial locations of assembly neurons (Supplementary Figure S3D). Thus, we predict that assemblies detected *in vivo* are the superficial subset of full assemblies, and may not include late assemblies with only a small fraction of neurons in superficial layers.

2.2 Determination of functional assemblies by structural features

It appears as if the spatial structure of the thalamic input stimuli strongly determines assembly membership. At the same time, neuronal assemblies are thought to be strongly recurrently connected (Song et al., 2005; Perin et al., 2011).

We generally observe that some features of structural connectivity can be predictive of the probability of assembly membership. Here, we consider features related to thalamic innervation, recurrent connectivity and synaptic clustering. We quantify the strength of this effect by means of a thresholded and signed version of the mutual information which we call their *normalized mutual information* and denote by nI; this allows us to compare the individual contributions (for details, see Methods). Its value has three basic properties: First, it is positive if the probability of assembly membership increases as the value of the structural feature increases, and is negative otherwise. Second, its absolute value is one if assembly membership can be completely predicted from the structural feature and it is only defined if the mutual information between the two processes is significantly larger than for randomly shuffled controls. Third, its absolute value does not require any assumptions about the shape of the dependency (e.g., linear, monotonic, etc.) between the structural feature and the probability of assembly membership.

2.2.1 Thalamic innervation

We began by considering the effect of direct thalamic innervation. Having confirmed that strong direct thalamic innervation facilitated assembly membership (Figure 4A, left), we formulated the following hypothesis: pairs of neurons are more likely to belong to the same assembly if they are innervated by overlapping sets of thalamic fibers. To test this, we first consider the common thalamic indegree of a pair of neurons, i.e., the number of thalamic fibers innervating both of them. Then, for each neuron, we use its *mean common thalamic indegree* over all cells in assembly A_n, as the structural feature to predict its membership in A_n. We performed this analysis separately for innervation from the VPM and POm nuclei.

In both cases, mean common thalamic indegree with an assembly increased the probability that a neuron is part of it (Figure 4A, second, olive curve for common thalamic indegree with the same assembly). The nI of the mean common thalamic indegree and membership in the same assembly was on average 0.165 for VPM and 0.054 for POm (Figure 4A, right, entries along the diagonals). More specifically, 0.157 and 0.034 for early assemblies, 0.198 and 0.087 for middle, and 0.082 and 0.037 for the late assembly. In addition, cross-assembly interactions were also observed, albeit at lower levels (Figure 4A, right, off-diagonal entries). The effect was strongest

for pairs of early and middle assemblies that responded to the same stimuli, e.g. assemblies 1, and 2, responding to pattern A.

The lower nI values for the late assembly are expected, as it contains many neurons in layers not directly innervated by thalamus (Figure 3C, Supplementary Figure S4A), and its activity is largely restricted to time bins in which the thalamic input is only weakly active (80 to 140 ms after onset; Figure 3B2 vs. 2A1). Interestingly, common innervation with POm has the highest nI for middle assemblies, which seems related to the prevalence of L5 neurons in them (2.3 times more L5 cells, than in the early ones). POm targets the upper part of L5, and more importantly L1 (Meyer et al., 2010; Reimann et al., 2022b) (Supplementary Figure S4A), where the apical tuft dendrites of thick-tufted L5 pyramidal cells reside (Harris and Shepherd, 2015; Ramaswamy and Markram, 2015). The delay caused by the long synapse to soma path distances (Supplementary Figure S4B) may explains the importance of common POm innervation 40-60 ms after stimulus onset.

Having confirmed that common innervation by thalamic fibers links pairs of neurons to the same assemblies, we then considered how much more assembly membership is determined by the identity of the specific patterns used. We hypothesized that direct innervation by fibers used in a pattern increases membership probability in assemblies associated with the same pattern. Specifically, we used as a structural feature the *pattern indegree* i.e., the total indegree of a neuron from VPM fibers used in each of the patterns.

As predicted from the previous results, probability of assembly membership grew rapidly with pattern indegree for early assemblies associated with the same pattern (Figure 4B, left). Every pattern had one early assembly strongly associated with it (except for pattern E, which only had middle and late assemblies Figure 3B2). On average, the nI of pattern indegree and assembly membership reached 0.26 (Figure 4B, right). A similar trend was observed for middle assemblies (mean: 0.215), while the late assembly was again an exception, with no value above 0.04, for reasons outlined above (Figure 4B third and fourth). In total, taking the stimulus patterns into account gives a 33% higher nI over the less specific mean common VPM innervation.

2.2.2 Recurrent connectivity

Even with perfect knowledge of thalamic innervation and pattern identity the nI did not exceed values of 0.3, leading to the question: What other factors determine the rest? The most commonly accepted structural correlate of cell assemblies is the overexpression of recurrent connectivity motifs between participating neurons (Harris, 2005; Song et al., 2005; Buzsáki, 2010; Perin et al., 2011). One particular class of motifs that has been linked to neuronal function are directed simplices of dimension k (k-simplices) (Reimann et al., 2017b). A k-simplex is a motif on $k+1$ neurons, which are all-to-all connected in a feed-forward fashion (Figure 4E left, inset); in particular 1-simplices are directed edges and 0-simplices are single cells. Indeed, we found a strong overexpression of directed simplices in the connectivity submatrices of cells within an assembly. In particular, the maximal simplex dimension found in assembly subgraphs is at least one higher than in the corresponding controls. Moreover, the peak of simplex counts in assembly graphs is in general one order of magnitude above the controls (Figure 4C).

Figure 4: Connectivity determines cell assembly membership. A: First: Effect of thalamic innervation (from VPM and POm nuclei) on participation in cell assemblies. Solid lines indicate the mean and the shaded areas indicate 95% confidence intervals. Second: probability of membership in an exemplary middle assembly against mean common POm indegrees with respect to all assemblies. Third and fourth: nI (normalized mutual information, see Methods) of mean common thalamic indegree and assembly membership. **B**: Probability of membership in exemplary early (first), middle (second), and late (third) assemblies against indegree with respect to all patterns. Fourth: nI of pattern indegree and assembly membership. **C:** Simplex counts within assemblies and random controls (same number of neurons with the same cell type distribution). (Caption continues on the next page.)

D: Probability of membership in exemplary early (first), middle (second) and late (third) assemblies against indegree with respect to all assemblies. Fourth: nI of indegree and assembly membership. **E:** First: Probability of membership in an examplary early assembly against k-indegree with respect to the same assembly. Inset: k-indegree is given by the number of k-simplices within an assembly (orange nodes) completely innervating a given neuron (black). nI of k-indegree and assembly membership for $k = 1$ (second), $k = 2$ (third) and $k = 3$ (fourth). (White: nI not defined.)

Based on this we define the *k-indegree with respect to an assembly* of a neuron i, as the number of k-simplices in the assembly such that all the cells in the simplex innervate i (see Figure 4E left, inset). For the case $k = 0$, i.e., the number of cells in the assembly innervating i, we found that it is a good predictor of membership in the same assembly, with an average nI value of 0.143 (Figure 4D). This time, late assembly membership could be predicted the best, with an nI of 0.224, compared to an average of 0.118 and 0.16 for early and middle assemblies, respectively (Figure 4D). Additionally, probability of late assembly membership also increased with 1-indegree with respect to all other assemblies. Conversely, 0-indegree with respect to the late assembly decreases the membership probability for early, and most of the middle assemblies (blue curves with negative slope on the first and second panels of Figure 4D). This reflects the temporal order of their activation and the fact that neurons in the deeper layers, that dominate the late assembly, mostly project outside the cortex and not back to superficial layers (Feldmeyer, 2012; Harris and Shepherd, 2015).

The nI values across the diagonals for $k = 1, 2$ are larger than for $k = 0$, for all assemblies except the late one. More precisely, we found nI averages of 0.152 and 0.161 for early, and 0.18 and 0.188 for middle assemblies for $k = 1$ and $k = 2$ respectively (Figure 4E second and third). On the other hand, off diagonal nI values drop for increasing k. This shows that not only the size of the presynaptic population has an effect on the activity of a neuron, but also the connectivity patterns between them. However, the effect of these non-local interactions are stronger within an assembly than across. For $k = 3$ the nI values drops, which can be explained by the narrower range of values the 3-indegree takes (Figure 4E first and fourth). As 0-indegree is the same as the general notion of indegree (the number of neurons in the afferent population) we will drop $k = 0$ and simply call it *indegree* in the following sections.

2.2.3 Synaptic clustering

So far, we have only considered features which can be extracted from the connectivity matrix of the system. However, our model also offers subcellular resolution, specifically, the dendritic locations of all synapses (Markram et al., 2015; Kanari et al., 2019), which we have demonstrated to be crucial for recreating accurate post-synaptic potentials (Ramaswamy et al., 2012; Ecker et al., 2020), and long-term-plasticity (Chindemi et al., 2022). This allowed us to explore the impact of co-firing neurons potentially sending synapses to the same dendritic branch i.e., forming synapse clusters (Wilson et al., 2016; Iacaruso et al., 2017; Kastellakis and Poirazi, 2019; Ujfalussy and Makara, 2020). We hypothesized that innervation from an assembly is more effective at facilitating membership in an assembly if it targets nearby dendritic locations. We therefore defined the synaptic clustering coefficient (SCC) with respect to an assembly A_n, based on the path distances between synapses from A_n on a given neuron (see Methods, Supplementary Figure S5). The SCC is a parameter-free feature, centered at zero. It is positive for intersynaptic distances that are lower than expected (indicating clustering) and negative otherwise (indicating avoidance).

Overall, we found similar trends as for the recurrent connectivity, although with a lower impact on assembly membership. Late assembly membership was explained the best by the SCC with 0.114 nI, while early and middle assemblies had an average nI of only 0.048 and

0.061 (Figure 5A). Although SCC was not that powerful by itself, we found that a given value of indegree led to a higher assembly membership probability if the innervation was significantly clustered (Figure 5B, first). This lead us to explore the correlation between indegree and the SCC (Figure 5B second), finding a weak but significant correlation between these measures. Note that the SCC controls for the decrease in distance between synapses expected from a higher indegree (see Methods), therefore we conclude that this is non-trival feature of the model. However, this means that the effects of indegree and SCC on assembly membership are partially redundant. To dissociate these effects, we compute the nI between SCC and assembly membership conditioned by indegree (see Methods), which is on average 0.025, reaching values up to 0.045 (Figure 5B third and fourth). This shows that indegree and SCC affect assembly membership both independently but more so in conjunction.

Figure 5: Synapse clustering coefficient determines cell assembly membership. A: Probability of membership in exemplary early (first), middle (second), and late (third) assemblies against synapse clustering coefficient (SCC, see Methods) with respect to all assemblies. Fourth: nI (normalized mutual information, see Methods) of SCC and assembly membership. **B:** Combined effect of SCC and indegree (as in Figure 4D). First: Probability of membership in an exemplary early assembly, against indegree with respect to the same assembly, grouped by SCC significance (see Methods). Second: Joint distribution of SCC and indegree. Third: nI of SCC and assembly membership conditioned by indegree. (White: nI not defined.) Fourth: Relation of nI and conditional nI grouped by postsynaptic early, middle, and late assemblies (i.e., rows of the matrices in A fourth, and B third). **C:** Simulation results for 10 selected neurons per assembly (with the highest indegree and significant clustering; red arrow in B left) with modified physiological conditions. Left: correlation of spike times with the assembly. Right: Single cell firing rates.

The observed effect of SCC can only be explained by nonlinear dendritic integration of synaptic inputs (Stuart and Spruston, 2015; Kastellakis and Poirazi, 2019; Goetz et al., 2021). Specifically, our model has two sources of nonlinearity: First, active ion channels on the dendrites (Stuart and Spruston, 2015; Goetz et al., 2021; Ujfalussy and Makara, 2020), causing Na^+ and Ca^{2+} spikes. Second, NMDA conductances, which open in a voltage dependent manner, leading to NMDA-spikes (Stuart and Spruston, 2015; Goetz et al., 2021). To show that these are the mechanisms by which the SCC acts, we studied how the removal of these non-linearities affects assembly membership of selected neurons. From each assembly we chose ten neurons whose probability of membership was most affected by their SCC i.e., those with highest indegree combined with significant clustering (red arrow in Figure 5B first). We subjected these neurons to the same input patterns they received in the simulations (see Methods), but with passive dendrites (Figure 5C green) or blocked NMDA receptors (Figure 5C orange). These modifications altered the neurons spiking activity, causing a non-negligible portion of them to drop out of their assembly, as their activity was no longer significantly correlated with it (Figure 5C first, see Methods). The manipulations resulted in a 45% drop in assembly membership for passive dendrites and 36% for blocking NMDA channels. Although, the manipulation resulted in an overall reduction of firing rate (Figure 5C second), this did not explain the drop in assembly membership (Figure 5C second: red dots with higher rate then black ones). We conclude that these nonlinearities contribute to the synchronization of activity within assemblies, underlying the observed effect of the SCC.

2.3 Robustness of assemblies across simulation instances

The results of our simulations are stochastic (Nolte et al., 2019), leading to different outcomes for repetitions of the same experiment, as in biology. To assess the robustness of our results, we repeated our *in silico* experiment 10 times with the same thalamic inputs but different random seeds. Changing the seed mostly affects the stochastic release of synaptic vesicles (Markram et al., 2015; Nolte et al., 2019) especially at the low, *in vivo*-like extracellular Ca^{2+} concentration used. The assemblies detected in the repetitions were similar to the ones described so far, in term of member neurons (Figure 6A), temporal structure (early and middle groups and a non-specific late one, not shown), and their determination by connectivity features (not shown).

We hypothesize that cell assemblies in cortical circuits are inherently stochastic objects, partially determined by the structural connectivity, input stimuli, and neuronal composition. Thus, each repetition yields a different (but overlapping) set of assemblies and neurons contained in them. In order to get an approximation of these stochastic objects we pooled the assemblies detected in all repetitions and determined which best corresponded to each other by clustering them based on the Jaccard distance of their constituent neurons (see Methods, Figure 6A). According to this distance, nearby assemblies have a large intersection relative to their size. We called the resulting clusters *consensus assemblies*, and the assemblies contained in each its *instances*. We assigned to neurons different degrees of membership in a consensus assembly, based on the fraction of instances they were part of, normalized by a random control, and called it its *coreness* (see Methods). We found, that as coreness increased so did the neurons' spike time reliability (defined across the repetitions of the experiment (Schreiber et al., 2003; Nolte et al., 2019), Methods); especially for the thalamus driven early assemblies (Figure 6B). We call the *union consensus assembly* the union of all its instances; and the *core consensus assembly* the set of cells whose coreness is significantly higher than expected (see Methods, see union vs. core distinction of Figure 6C).

When repeating our structural analysis on the core consensus assemblies, we found higher values of nI than before in all cases, except for the SCC (Figure 6D and Figure 7). This indicates that assembly membership in this model is not a binary property but exists on a spectrum from a

highly reliable and structurally determined core to a more loosely associated and less connected periphery. The notion of consensus assemblies is a way of accessing this spectrum.

Figure 6: Consensus assemblies. A: Jaccard similarity based hierarchical clustering of assemblies from different simulation instances. **B:** Spike time reliability (see Methods) of neurons belonging to given fraction of assembly instances. **C:** Locations of neurons in the union (all instances) and core (at least 9/10 instances, see Methods) of exemplary (pattern A responsive) early consensus assembly. **D:** nI (normalized mutual information, see Methods) of connectivity features and consensus assemblies membership.

Another way to take all ten repetitions into account would be to average the time-binned spike trains of the simulated neurons. Thus, we averaged the input instead of the output of the assembly detection pipeline and we call these the *average assemblies*. We first compared these to the assemblies obtained in a single repetition. The similarity of significant time bins was higher for average assemblies (Supplementary Figure S6B1), and their sizes were larger (Supplementary Figure S6D second panel), with neurons belonging to up to 10 assemblies out of the 13 detected (Supplementary Figure S6D third panel). On the other hand, the nI of the structural features and membership remained the same as for a single repetition (compare matrices in Figures 4 and 5 to Supplementary Figure S6C).

When contrasting the average and consensus assemblies, we found pairs with high Jaccard similarity (Supplementary Figure S6D and E1). A detailed comparison showed that all neurons that are part of at least 6 out of 10 assembly instances were all contained in their matching average assembly. Further lowering the cutoff began admitting neurons that were not part of the corresponding average assembly (Supplementary Figure S6E2). On the other hand, there are barely any cells in the average that are not contained in the union consensus assembly. This demonstrates how assembly membership becomes less determined towards the periphery, mirroring the reduced nI with the structural connectivity features. Furthermore, a neuron in a consensus assembly will most likely belong to all instances (Supplementary Figure S6E2), unlike for the binomial distribution expected by chance.

In summary, while average assemblies give similar results to the union consensus assemblies, the coreness values used in the consensus assembly framework assign different degrees of membership to its neurons that can be taken into account in downstream analyses, e.g., by con-

sidering only the functionally reliable core. Furthermore, for this core the determination by most structural features, measured by nI, is stronger (Figure 7).

Figure 7: Summary of nIs of all connectivity features and assembly membership. Assemblies from single simulation on top and consensus assemblies from 10 simulations on the bottom. Only within-assembly interactions (diagonals of nI matrices) shown, except for the patterns, where for each column (postsynaptic assembly) we used the maximum value (strongest innervating pattern). Not only the colors, but the radii of the pies code for features: Red (VPM) longer one depicts common-innervation (as yellow for POm) while the shorter one direct innervation from patterns. Green: Simplex dimension increases as length decreases (longest: indegree, shortest: 3-indegree). Blue: longer one is the SCC conditioned on indegree, while the shorter one is the same, but unconditioned (and that is why it always overlaps with indegree).

3 Discussion

Using a detailed, large-scale cortical network model we examined the link between cortical structure and function. Our principal findings from analyzing the connectivity of functional cell assemblies are as follows: (1) Different afferents dominated determination of membership in assemblies linked to different time windows: VPM innervation affected membership in early assemblies, POm innervation in middle assemblies, while recurrent innervation in the late assembly (Figure 7 red, yellow, green). (2) Recurrent innervation more strongly facilitated assembly membership the more the innervating neurons were wired among themselves, adding a non-local component to the structure-function relation. (3) Similarly, the innervation of a neuron by an assembly was significantly more powerful in facilitating membership when its synapses were clustered on the dendrites (Figure 7 blue). (4) In conjunction with the structure of cortical connectivity, features of subcellular physiology such as active dendritic channels and NMDA receptors, also influence assembly membership. (5) Interactions between assemblies emerged, where innervation by one assembly explained membership in another. Positive interactions (with increased membership probability) were found when the direction of innervation reflected the temporal order of assembly activation; otherwise, weaker or even negative interactions were found.

Point (1) above confirms our previous findings that, while the presence of an external stimulus makes the circuit much more reliable, this effect is not merely driven by direct innervation, but also requires recurrent connectivity (Nolte et al., 2019). Point (2) predicts a functional consequence for non-random features of neuronal connectivity, such as nodes with high centrality values and the presence of a rich club, that have been characterized in many species and regions,

at various levels of resolution (Bassett and Bullmore, 2017). Points (3) and (4) link the theory of neuronal assemblies to the literature on active dendritic computation (Poirazi et al., 2003; Stuart and Spruston, 2015; Kastellakis and Poirazi, 2019). Point (5) strengthens the case that the topology of connectivity is best studied in a directed way, since undirected networks (although more amenable to network science methods) miss an essential part of the picture. In particular, the relationship between the structural directionality of the connection and the temporal direction of the flow of activity (Reimann et al., 2017b).

Our analysis supports the idea that neuronal activity revolves around activation of assemblies. We have shown that assembly membership is determined by certain structural prerequisites, mostly amounting to increased membership probability when more afferent synapses from various sources are formed on a neuron. These may be costly to fulfill, both in terms of energy (Harris et al., 2012) and space taken up by wiring. Chklovskii et al. (2002) considered efficient layouts of wiring, i.e., axons and dendrites, given a certain connectivity matrix. They found that the layout is tightly constrained by the available space and close to a theoretical optimum. Here, we expanded on that idea, demonstrating that on top of it, the structure of the connectivity matrix is efficient. Indeed, on the sub-cellular level, we have shown, that synaptic clustering on dendrites increases efficiency, in that the same probability of assembly membership can be obtained with about 50% of the indegree when the synapses are highly clustered (Figure 5B, Supplementary Figure S7A). Furthermore, on the connectivity level, another potential mechanism lies in the non-local interaction of the presynaptic population measured by the k-indegree. We have shown that for higher values of k, a given membership probability is attained for lower k-indegree values, suggesting higher efficiency (Figure 4E, left). However, as k grows, k-indegree counts the innervation by larger and larger motifs, potentially requiring more incoming synaptic connections (Supplementary Figure S7B2). Here, we control for this by repeating our analysis not with respect to the number of motifs but the size of the presynaptic subpopulation forming them. When going up from $k = 0$ to $k = 1$, $k = 2$, and $k = 3$ the same membership probability can be obtained for a given neuron with about 80%, 34% and 6% of the incoming connections (Supplementary Figure S7B1), confirming that non-local interactions in the presynaptic population make afferent innervation more efficient. This can be explained by our earlier finding that the simplicial motifs we considered increase the correlations and reliability of the spiking activity of participating neurons (Reimann et al., 2017b).

It is possible that *wiring efficiency* can be optimized further, based on experience, through structural and functional plasticity. Recent modelling studies investigated how starting from a random initial connectivity, plasticity rules and network activity lead to assembly formation, maintenance and *competition for member neurons* (Fauth and Van Rossum, 2019; Kossio et al., 2021; Gastaldi et al., 2021). Conversely, our network model has strongly non-random connectivity, constrained by neuronal morphology (Reimann et al., 2015, 2017a), and can thus be viewed as a circuit in a non-random plastic state, but unshaped by experience. The presence of assemblies in such a naive circuit is in line with the belief that the brain is not a tabula rasa (Buzsáki, 2019). It is also in line with the recent *in vivo* experiments of Bathellier et al. (2012) and Trägenap et al. (2022), who found endogenous cell assemblies in mouse auditory and ferret visual cortices. Furthermore, Trägenap et al. (2022) also found that these endogenous assemblies *solidify* and become more reliable after eye opening. These points lead to the question: How does long-term plasticity affect the three aspects discussed above, i.e., assembly wiring efficiency, competition for member neurons and assembly solidification? To address this question, we are integrating the functional plasticity model of Chindemi et al. (2022) into our network simulations. We can then analyze these with the methods we have introduced here, which provide new, quantitative ways to characterize assemblies, their temporal evolution and the connectivity underlying them.

4 Methods

4.1 Network simulations

The most recent version of the detailed, large-scale cortical microcircuit model of Markram et al. (2015) was used for the *in silico* experiments in this study. Updates on its anatomy, e.g., atlas-based cell densities are described in Reimann et al. (2022b). Updates on the physiology of single cell models and multi-vesicular release of synapses are described in Reva et al. (2022) and Barros-Zulaica et al. (2019). The $2.4mm^3$ subvolume of the juvenile rat somatosensory cortex, containing 211'712 neurons, their structural connectivity and thalamic inputs is freely available at: https://doi.org/10.5281/zenodo.6906785 (Reimann et al., 2022a).

Although large-scale with bio-realistic counts of synapses originating from the local neurons, the neurons in the circuit still lacked most of their synapses (originating from other, non-modeled regions) (Markram et al., 2015). In order to compensate for this missing input, layer and cell-type specific somatic conductances following an Ornstein-Uhlenbeck process, similar to Destexhe et al. (2001) were injected to the cell bodies of all neurons. The algorithm used to determine the mean and variance of the conductance needed to put the cells into an *in vivo*-like high-conductance state, and the network as whole into an *in vivo*-like asynchronous firing regime with low rates and realistic responses to short whisker stimuli will be described elsewhere (see Data and code availability).

Simulations of selected cells with modified physiological conditions (active dendritic channels blocked or NMDA conductance blocked) used the activity replay paradigm of Nolte et al. (2019). In short, for each of these cells, spike times of its presynaptic population were recorded in the original network simulation. Then the selected cells were simulated in isolation by activating their afferent synapses according to the recorded spike times in the network simulation. Thus, the modified activity of the isolated cell did not affect the rest of the network.

Simulations were run using the `NEURON` simulator as a core engine with the Blue Brain Project's collection of hoc and NMODL templates for parallel execution on supercomputers (Hines and Carnevale, 1997; Awile et al., 2022). Simulating 2 minutes of biological time took 100'000 core hours, on our HPE based supercomputer, installed at CSCS, Lugano.

4.2 Distance metrics

This section gives a brief overview and justification of the various distance metrics used below.

Population activity in time bins were compared using their cosine similarity Carrillo-Reid et al. (2015). Two time bins with high cosine similarity have similar sets of firing neurons, thus detecting co-firing. Note that there is an increasing relationship between firing rate and cosine similarity (Cutts and Eglen, 2014) (Supplementary Figure S1A2).

Input patterns were defined using the Hamming distance between the sets of VPM fiber bundles involved to have specific values and thus specific sizes of intersections (Figure 2B). Input patterns were compared using Earth mover distances between the flat map locations of their contained fibers (Figure 3E).

Assemblies of neurons were compared using their Jaccard distances. Like Hamming, this also compares sizes of intersections but it is normalized with respect to the sizes of the sets involved.

Assembly sequences i.e., vectors of size the number of assemblies, with each entry counting the number of time bins the corresponding assembly was active in response to an input pattern, were compared using their normalized Euclidean distances. This is the Euclidean distance between vectors of normalized length.

Finally, afferent synapses were compared using their path distances. Specifically, the dendritic tree was represented as a graph with nodes being its branching points and edges between them

weighted according to the length of sections connecting them. Distances between synapses was computed as the path distance in this graph.

4.3 Thalamic input stimuli

The VPM input spike trains were similar to the ones used in Reimann et al. (2022c). In detail, the 5388 VPM fibers innervating the simulated volume were first restricted to be $\leq 500\mu m$ from the middle of the circuit in the horizontal plane to avoid boundary artefacts. To measure these distances we use a flat map, i.e., a two dimensional projection of the volume onto the horizontal plane, orthogonal to layer boundaries (Reimann et al., 2022b). Second, the flat map locations of the resulting 3017 fibers were clustered using k-means to form 100 bundles of fibers. To form the base patterns (A, B, C, and D), we randomly selected four non-overlapping groups of bundles, each containing 12% of them (corresponding to 366 fibers each). The remaining 6 patterns were derived from these base patterns with various degrees of overlap (see beginning of Results, Figure 2B). Third, the input stream was defined as a random presentation of these ten patterns with 500 ms inter-stimulus intervals, such that in every 30 second time intervals every pattern was presented exactly six times. Last, for each pattern presentation, unique spike times were generated for its corresponding fibers following an inhomogeneous adapting Markov process (Muller et al., 2007). When a pattern was presented, the rate of its fibers jumped to 30 Hz and decayed to 1 Hz over 100 ms. For the non-specific POm stimuli, a randomly selected 12% of the (unclustered) 3864 POm fibers were activated each time any pattern was presented (in every 500 ms). The spike trains were designed with the same temporal dynamics as described above for VPM, but with half the maximum rate (15 Hz). The implementation of spike time generation was based on `Elephant` (Denker et al., 2018).

4.4 Assembly detection

Our assembly detection pipeline was a mix of established techniques and consisted of five steps: binning of spike trains, selecting significant time bins, clustering of significant times bins via the cosine similarity of their activity, and determination of neurons corresponding to a time bin cluster and thus forming and assembly. Note, that time bins instead of neurons were clustered because this allows neurons to belong to several assemblies.

Spikes of excitatory cells were first binned using 20 ms time bins, based on Harris et al. (2003), who take a postsynaptic reader neuron specific point of view. They suggest 10-30 ms as an ideal integration time window of the presynaptic (assembly) spikes (Harris et al., 2003; Buzsáki, 2010). Next, time bins with a significantly high level of activity were detected. The significance threshold was determined as the mean activity level plus the 95th percentile of the standard deviation of shuffled controls. The 100 random controls were rather strict, i.e., all spikes were shifted only by one time bin forward or backward, based on Sasaki et al. (2006); Carrillo-Reid et al. (2015); Pérez-Ortega et al. (2021). Next, a similarity matrix of significant time bins was built, based on the cosine similarity of activation vectors, i.e., vectors of spike counts of all neurons in the given time bins (Carrillo-Reid et al., 2015; Pérez-Ortega et al., 2021). The similarity matrix of significant time bins was then hierarchically clustered using Ward's linkage (Montijn et al., 2016; Pérez-Ortega et al., 2021). Potential number of clusters were scanned between five and twenty, and the one with the lowest Davis-Bouldin index was chosen, which maximizes the similarity within elements of the cluster while maximizing the the between cluster similarity (Davies and Bouldin, 1979). These clusters corresponded to potential assemblies.

As the last step, neurons were associated to these clusters based on their spiking activity, and it was determined whether they formed a cell assembly or not. In detail, the correlations between the spike trains of all neurons and the activation sequences of all clusters were computed and

the ones with significant correlation selected. Significance was determined based on exceeding the 95th percentile of correlations of shuffle controls (1000 controls with spikes of individual cells shifted by any amount) (Montijn et al., 2016; Herzog et al., 2021). In relation to Figure 5C left, it is important to note, that these correlation thresholds were specific to a pair of a neuron and an assembly. Finally, it is possible to have a group of neurons that is highly correlated with one part of the significant time bins in a cluster, and another that is highly correlated with the rest, while the two groups of neurons have uncorrelated activity. To filter out this scenario, it was required that the mean pairwise correlation of the spikes of the neurons with significant correlations was higher than the mean pairwise correlation of neurons in the whole dataset (Herzog et al., 2021). Clusters passing this test were considered to be functional assemblies and the neurons with significant correlations their constituent cells.

For a test of the methods on synthetic data and comparison with other ways of detecting cell assemblies please consult Herzog et al. (2021). Assembly detection was implemented in Python, using common packages like `numpy` and `scipy` and parallelized to deal with the tens of millions of spikes recorded in the simulations.

4.5 Calculation of information theorectical measurements

To quantify the structural predictability of assemblies, *mutual information* of assembly membership (Y_n) and a structural feature of a neuron (X_m) was used, which is a measure of the mutual dependence between the two variables (Gray, 2011). More precisely, Y_n is a binary random variable that takes the value 1 if a neuron belongs to an assembly A_n, and 0 otherwise. On the other hand, X_m is a random variable determined by a structural property of the neuron with respect to the assembly A_m, e.g., the number of afferent connections into the neuron from all neurons in A_m.

To asses the dependence between these two variables, first the dependence of the probability of a neuron belonging to assembly A_n given a specific value of the structural feature measured by X_m was studied. More precisely, the function $f_{n,m}$ whose domain is the values of X_m and is given by:

$$f_{n,m}(x) = P(Y_n = 1 | X_m = x)$$

was considered. If this function has an increasing or decreasing trend, then the random variables Y_n and X_m can not be independent.

Their dependence was quantified by means of their mutual information. The value of mutual information is always non-negative and it is zero when the random variables are independent. In order to restrict this value to [0, 1] it was divided by the entropy of Y_n, which measures the level of inherent uncertainty of the possible outcome of the values of Y_n (Shannon, 1948). The calculation of mutual information is based on the probabilities of X_m and Y_n across all possible outcomes. If the number of possible values of X_m is large compared to the number of samples, there can be errors in determining these probabilities, possibly leading to inflated values of the mutual information. Therefore, the values of X_m were binned into 21 bins between the 1st and 99th percentile of all sampled values. The number of bins was determined such that the resulting value of mutual information in a shuffled control did not exceed 0.01. Shuffled controls were also used to threshold the mutual information values by considering only the pairs (n, m) whose mutual information was larger than the mean plus one standard deviation of all pairs in the shuffled controls. Finally, a negative sign was added to the significant mutual information value if the function $f_{n,m}$ was decreasing i.e., the probability of membership in A_n decreased as the values of X_m increased. This was assessed by the slope of a weighted (by the number of samples in each bins) linear fit of the function $f_{n,m}$. This thresholded and signed mutual information value was called *normalized mutual information* and denoted $nI(Y_n, X_m)$.

All the statements above can be made conditional with respect to a third random variable, yielding the *conditional normalized mutual information*, which was used when two structural features were inherently believed to be interacting as in the case of the *SCC* and indegree.

Calculations were done with the `pyitlib` package.

4.6 Synaptic clustering coefficient

To quantify the co-localization of synapses on the dendrites of a neuron i from its presynaptic population P_i with a single, parameter-free metric, synaptic clustering coefficient SCC was defined and calculated for all excitatory neurons in the circuit with respect to all assemblies. Based on these locations, D_i, the matrix of all pairwise path distances between synapses on i from P_i were calculated. Let $D_{i,p}$ be the submatrix for pairs of synapses originating from a subpopulation $p \subseteq P_i$. Then the nearest neighbour distance for p can be written as:

$$nnd(i, p) = \mathrm{mean}\left(\min_{rows}\left(D_{i,p}\right)\right)$$

In particular, for the subpopulation of P_i of neurons in the assembly A_n, denoted by p_n, $nnd(i, A_n) = nnd(i, p_n)$ was defined, where $p_n = P_i \cap A_n$. This value was normalized, using the nnd values of 20 random presynaptic populations from P_i of the same size as p_n. In summary, the SCC was defined as the negative z-score of $nnd(i, A_n)$ with respect to the distribution of control nnds (Supplementary Figure S5). Additionally, the significance of the clustering or avoidance of the synapse locations was determined with a two-tailed t-test of $nnd(i, A_n)$ against the 20 random samples with an alpha level of 0.05. SCC was implemented using `NeuroM` and other common Python packages.

4.7 Determination of consensus assemblies

Consensus assemblies were defined over multiple repetitions of the same input stream. These were groups of assemblies with similar sets of neurons. Additionally, all assemblies in a group were required to originate from a different repetition; noted as the *repetition separation criterion*. The Jaccard distance matrix between all pairs of assemblies from all repetitions were computed and modified by setting the distances between pairs of assemblies from the same repetition to twice the maximum of the whole matrix. The matrix was then hierarchically clustered using Ward's linkage, and the lowest number of clusters that satisfied the repetition separation criterion was chosen. The resulting clusters were the *consensus assemblies*, and the assemblies within them their *instances*.

The *union consensus assembly* was defined as the set of neurons given by the union of all instances. Its member neurons were assigned a membership degree based on number of instances they were part of in two ways. First, by simply using the fraction of the instances a neuron was part of. Second, in what was called the *coreness* value of a neuron, which is the number of instances a neuron is part of normalized by its expected value, given the number and sizes of its instances. This was calculated as a binomial distribution with n set to the number of instances and p to the mean size of the instances, divided by the size of the union consensus assembly. Based on this, the coreness of a neuron contained in r instances was defined as $-log_{10}(1 - B_{n,p}(r))$, where B is the cumulative binomial distribution. Neurons with a coreness value exceeding 4 were considered to be part of the corresponding *core consensus assembly*.

4.8 Calculation of spike time reliability

Spike time reliability was defined as the mean of the cosine similarities of a given neuron's mean centered, smoothed spike times across all pairs of repetitions (Schreiber et al., 2003; Cutts and

Eglen, 2014). To smooth the spike times, they were first binned to 1 ms time bins, and then convolved with a Gaussian kernel with a standard deviation of 10 ms.

Acknowledgements

The authors thank Nicolas Ninin for his involvement in the early stage of this project, Elvis Boci and Cyrille Favreau for their help with visualizations and Alberto Antonietti, Christoph Pokorny, Kathryn B. Hess, Ran Levi, and Henry Markram for discussions.

Funding

This study was supported by funding to the Blue Brain Project, a research center of the École polytechnique fédérale de Lausanne (EPFL), from the Swiss government's ETH Board of the Swiss Federal Institutes of Technology.

Author contributions

- Conceptualization: A.E., D.E.S., M.W.R.

- Data curation: A.E., J.B.I, S.B.-P.

- Formal analysis: A.E., D.E.S., M.W.R.

- Investigation: A.E.

- Methodology: A.E., D.E.S., M.W.R.

- Project administration: M.W.R.

- Software: A.E., D.E.S, M.W.R., S.B.-P., J.B.I.

- Supervision: M.W.R.

- Validation: A.E., D.E.S., M.W.R., J.B.I., S.B.-P.

- Visualization: A.E.

- Writing - original draft: A.E., D.E.S., M.W.R.

- Writing - review & editing: A.E., D.E.S., M.W.R., J.B.I., S.B.-P.

Data and code availability

Analysis code and processed data will be available on Zenodo by the time of the journal submission, which will happen after we finish the manuscript complementing Reimann et al. (2022b): *Part II: Modeling Physiology.*

References

Awile, O., Kumbhar, P., Cornu, N., Dura-Bernal, S., King, J. G., Lupton, O., Magkanaris, I., McDougal, R. A., Newton, A. J., Pereira, F., Săvulescu, A., Carnevale, N. T., Lytton, W. W., Hines, M. L., and Schürmann, F. (2022). Modernizing the NEURON Simulator for Sustainability, Portability, and Performance. *Frontiers in Neuroinformatics*, 16(884046).

Barros-Zulaica, N., Rahmon, J., Chindemi, G., Perin, R., Markram, H., Muller, E., and Ramaswamy, S. (2019). Estimating the Readily-Releasable Vesicle Pool Size at Synaptic Connections in the Neocortex. *Frontiers in Synaptic Neuroscience*, 11(29).

Bassett, D. S. and Bullmore, E. T. (2017). Small-World Brain Networks Revisited. *Neuroscientist*, 23(5):499–516.

Bathellier, B., Ushakova, L., and Rumpel, S. (2012). Discrete Neocortical Dynamics Predict Behavioral Categorization of Sounds. *Neuron*, 76(2):435–449.

Buzsáki, G. (2010). Neural Syntax: Cell Assemblies, Synapsembles, and Readers. *Neuron*, 68(3):362–385.

Buzsáki, G. (2019). *The brain from inside out.* Oxford University Press.

Carrillo-Reid, L., Miller, J. E. K., Hamm, J. P., Jackson, J., and Yuste, R. (2015). Endogenous sequential cortical activity evoked by visual stimuli. *Journal of Neuroscience*, 35(23):8813–8828.

Chindemi, G., Abdellah, M., Amsalem, O., Benavides-Piccione, R., Delattre, V., Doron, M., Ecker, A., Jaquier, A. T., King, J., Kumbhar, P., Monney, C., Perin, R., Rössert, C., Tuncel, M. A., van Geit, W., DeFelipe, J., Graupner, M., Segev, I., Markram, H., and Muller, E. (2022). A calcium-based plasticity model predicts long-term potentiation and depression in the neocortex. *Nature Communications*, 13(3038).

Chklovskii, D. B., Schikorski, T., and Stevens, C. F. (2002). Wiring optimization in cortical circuits. *Neuron*, 34(3):341–347.

Cutts, C. S. and Eglen, X. S. J. (2014). Detecting pairwise correlations in spike trains: An objective comparison of methods and application to the study of retinal waves. *Journal of Neuroscience*, 34(43):14288–14303.

Davies, D. L. and Bouldin, D. W. (1979). A Cluster Separation Measure. *IEEE Transactions on Pattern Analysis and Machine Learning*, PAMI-1(2):224–227.

Denker, M., Yegenoglu, A., and Grün, S. (2018). Collaborative HPC-enabled workflows on the HBP Collaboratory using the Elephant framework. In *Neuroinformatics 2018*.

Destexhe, A., Rudolph, M., Fellous, J. M., and Sejnowski, T. J. (2001). Fluctuating synaptic conductances recreate in vivo-like activity in neocortical neurons. *Neuroscience*, 107(1):13–24.

Dragoi, G. and Buzsáki, G. (2006). Temporal Encoding of Place Sequences by Hippocampal Cell Assemblies. *Neuron*, 50(1):145–157.

Ecker, A., Romani, A., Sáray, S., Káli, S., Migliore, M., Falck, J., Lange, S., Mercer, A., Thomson, A. M., Muller, E., Reimann, M. W., and Ramaswamy, S. (2020). Data-driven integration of hippocampal CA1 synaptic physiology in silico. *Hippocampus*, 30(11):1129–1145.

Fauth, M. J. and Van Rossum, M. C. (2019). Self-organized reactivation maintains and reinforces memories despite synaptic turnover. *eLife*, 8:e43717.

Feldmeyer, D. (2012). Excitatory neuronal connectivity in the barrel cortex. *Frontiers in Neuroanatomy*, 6(24).

Gastaldi, C., Schwalger, T., de Falco, E., Quiroga, R. Q., and Gerstner, W. (2021). When shared concept cells support associations: Theory of overlapping memory engrams. *PLoS Computational Biology*, 17(12):e1009691.

Goetz, L., Roth, A., and Häusser, M. (2021). Active dendrites enable strong but sparse inputs to determine orientation selectivity. *PNAS*, 118(30):e2017339118.

Gray, R. M. (2011). *Entropy and information theory.* Springer Science & Business Media.

Harris, J. J., Jolivet, R., and Attwell, D. (2012). Synaptic Energy Use and Supply. *Neuron*, 75(5):762–777.

Harris, K. D. (2005). Neural signatures of cell assembly organization. *Nature Reviews Neuroscience*, 6(5):399–407.

Harris, K. D., Csicsvari, J., Hirase, H., Dragoi, G., and Buzsáki, G. (2003). Organization of cell assemblies in the hippocampus. *Nature*, 424(6948):552–556.

Harris, K. D. and Shepherd, G. M. (2015). The neocortical circuit: Themes and variations. *Nature Neuroscience*, 18(2):170–181.

Hebb, D. O. (1949). *The Organization of Behavior; A Neuropsychological Theory.* John Wiley & Sons, Inc., New York.

Herzog, R., Morales, A., Mora, S., Araya, J., Escobar, M. J., Palacios, A. G., and Cofré, R. (2021). Scalable and accurate method for neuronal ensemble detection in spiking neural networks. *PLoS ONE*, 16(7):e0251647.

Hines, M. L. and Carnevale, N. T. (1997). The NEURON simulation environment. *Neural computation*, 9(6):1179–1209.

Hopfield, J. J. (1982). Neural networks and physical systems with emergent collective computational abilities. *PNAS*, 79(8):2554–2558.

Iacaruso, M. F., Gasler, I. T., and Hofer, S. B. (2017). Synaptic organization of visual space in primary visual cortex. *Nature*, 547(7664):449–452.

Isbister, J. B., Reyes-Puerta, V., Sun, J.-J., Horenko, I., and Luhmann, H. J. (2021). Clustering and control for adaptation uncovers time-warped spike time patterns in cortical networks in vivo. *Scientific Reports*, 11:15066.

Kanari, L., Ramaswamy, S., Shi, Y., Morand, S., Meystre, J., Perin, R., Abdellah, M., Wang, Y., Hess, K., and Markram, H. (2019). Objective Morphological Classification of Neocortical Pyramidal Cells. *Cerebral Cortex*, 29(4):1719–1735.

Kastellakis, G. and Poirazi, P. (2019). Synaptic Clustering and Memory Formation. *Frontiers in Molecular Neuroscience*, 12(300).

Kossio, Y. F. K., Goedeke, S., Klos, C., and Memmesheimer, R. M. (2021). Drifting assemblies for persistent memory: Neuron transitions and unsupervised compensation. *PNAS*, 118(46):e2023832118.

Krotov, D. and Hopfield, J. J. (2016). Dense associative memory for pattern recognition. *Advances in Neural Information Processing Systems*, 29:1172–1180.

Lopes-dos Santos, V., Ribeiro, S., and Tort, A. B. (2013). Detecting cell assemblies in large neuronal populations. *Journal of Neuroscience Methods*, 220(2):149–166.

Markram, M., Muller, E., Ramaswamy, S., Reimann, M. W., et al. (2015). Reconstruction and Simulation of Neocortical Microcircuitry. *Cell*, 163:456–492.

Meyer, H. S., Wimmer, V. C., Hemberger, M., Bruno, R. M., De Kock, C. P., Frick, A., Sakmann, B., and Helmstaedter, M. (2010). Cell type-specific thalamic innervation in a column of rat vibrissal cortex. *Cerebral Cortex*, 20(10):2287–2303.

Miller, J. E. K., Ayzenshtat, I., Carrillo-Reid, L., and Yuste, R. (2014). Visual stimuli recruit intrinsically generated cortical ensembles. *PNAS*, 111(38):E4053–E4061.

Montijn, X. J. S., Olcese, U., and Pennartz, X. C. M. A. (2016). Visual Stimulus Detection Correlates with the Consistency of Temporal Sequences within Stereotyped Events of V1 Neuronal Population Activity. *The Journal of Neuroscience*, 36(33):8624–8640.

Muller, E., Buesing, L., Schemmel, J., and Meier, K. (2007). Spike-Frequency Adapting Neural Ensembles: Beyond Mean Adaptation and Renewal Theories. *Neural Computation*, 19:2958–3010.

Nolte, M., Gal, E., Markram, H., and Reimann, M. W. (2020). Impact of higher order network structure on emergent cortical activity. *Network Neuroscience*, 4(1):292–314.

Nolte, M., Reimann, M. W., King, J. G., Markram, H., and Muller, E. B. (2019). Cortical reliability amid noise and chaos. *Nature Communications*, 10(3792).

Pérez-Ortega, J., Alejandre-García, T., and Yuste, R. (2021). Long-term stability of cortical ensembles. *eLife*, 10:e64449.

Perin, R., Berger, T. K., and Markram, H. (2011). A synaptic organizing principle for cortical neuronal groups. *PNAS*, 108(13):5419–5424.

Poirazi, P., Brannon, T., and Mel, B. W. (2003). Pyramidal Neuron as Two-Layer Neural Network. *Neuron*, 37:989–999.

Ramaswamy, S., Hill, S. L., King, J. G., Schürmann, F., Wang, Y., and Markram, H. (2012). Intrinsic morphological diversity of thick-tufted layer 5 pyramidal neurons ensures robust and invariant properties of in silico synaptic connections. *Journal of Physiology*, 590(4):737–752.

Ramaswamy, S. and Markram, H. (2015). Anatomy and physiology of the thick-tufted layer 5 pyramidal neuron. *Frontiers in Cellular Neuroscience*, 9(233).

Reimann, M. W., Bolaños-Puchet, S., Courcol, J.-D., Egas Santandre, D., et al. (2022a). A Model of Rat Non-barrel Somatosensory Cortex Anatomy.

Reimann, M. W., Bolaños-Puchet, S., Courcol, J.-D., Egas Santandre, D., et al. (2022b). Modeling and Simulation of Rat Non-Barrel Somatosensory Cortex. Part I: Modeling Anatomy. *bioRxiv*.

Reimann, M. W., Horlemann, A. L., Ramaswamy, S., Muller, E. B., and Markram, H. (2017a). Morphological diversity strongly constrains synaptic connectivity and plasticity. *Cerebral Cortex*, 27(9):4570–4585.

Reimann, M. W., King, J. G., Muller, E. B., Ramaswamy, S., and Markram, H. (2015). An algorithm to predict the connectome of neural microcircuits. *Frontiers in computational neuroscience*, 9(120).

Reimann, M. W., Nolte, M., Scolamiero, M., Turner, K., Perin, R., Chindemi, G., Dłotko, P., Levi, R., Hess, K., and Markram, H. (2017b). Cliques of neurons bound into cavities provide a missing link between structure and function. *Frontiers in Computational Neuroscience*, 11(48).

Reimann, M. W., Riihimäki, H., Smith, J. P., Lazovskis, J., Pokorny, C., and Levi, R. (2022c). Topology of synaptic connectivity constrains neuronal stimulus representation, predicting two complementary coding strategies. *PLoS ONE*, 17(1):e0261702.

Reva, M., Roessert, C., Arnaudon, A., Damart, T., Mandge, D., Tuncel, A., Ramaswamy, S., Markram, H., and Werner, V. G. (2022). A universal workflow for creation, validation and generalization of detailed neuronal models. *bioRxiv*.

Reyes-Puerta, V., Sun, J. J., Kim, S., Kilb, W., and Luhmann, H. J. (2015). Laminar and Columnar Structure of Sensory-Evoked Multineuronal Spike Sequences in Adult Rat Barrel Cortex in Vivo. *Cerebral Cortex*, 25(8):2001–2021.

Sasaki, T., Kimura, R., Tsukamoto, M., Matsuki, N., and Ikegaya, Y. (2006). Integrative spike dynamics of rat CA1 neurons: A multineuronal imaging study. *Journal of Physiology*, 574(1):195–208.

Schreiber, S., Fellous, J. M., Whitmer, D., Tiesinga, P., and Sejnowski, T. J. (2003). A new correlation-based measure of spike timing reliability. *Neurocomputing*, 52-54:925–931.

Shannon, C. E. (1948). A Mathematical Theory of Communication. *Bell System Technical Journal*, 27:623–656.

Shatz, C. J. (1992). The Developing Brain. *Scientific American*, 267(3):60–67.

Song, S., Sjöström, P. J., Reigl, M., Nelson, S., and Chklovskii, D. B. (2005). Highly nonrandom features of synaptic connectivity in local cortical circuits. *PLoS Biology*, 3(3):e68.

Stuart, G. J. and Spruston, N. (2015). Dendritic integration: 60 years of progress. *Nature Neuroscience*, 18(12):1713–1721.

Trägenap, S., Whitney, D. E., Fitzpatrick, D., and Kaschube, M. (2022). Experience drives the development of novel, reliable cortical sensory representations from endogenously structured networks. *bioRxiv*.

Ujfalussy, B. B. and Makara, J. K. (2020). Impact of functional synapse clusters on neuronal response selectivity. *Nature Communications*, 11(1413).

van der Plas, T. L., Tubiana, J., Le Goc, G., Migault, G., Kunst, M., Baier, H., Bormuth, V., Englitz, B., and Debrégeas, G. (2023). Neural assemblies uncovered by generative modeling explain whole-brain activity statistics and reflect structural connectivity. *eLife*, 11:e83139.

Wilson, D. E., Whitney, D. E., Scholl, B., and Fitzpatrick, D. (2016). Orientation selectivity and the functional clustering of synaptic inputs in primary visual cortex. *Nature Neuroscience*, 19(8):1003–1009.

Wohrer, A., Humphries, M. D., and Machens, C. K. (2013). Population-wide distributions of neural activity during perceptual decision-making. *Progress in Neurobiology*, 103:156–193.

Supplementary Material

Supplementary Figures

Figure S1: Spikes of (significant) time bins. A1 Cosine similarity matrix (same as in Figure 2B1, but unsorted). **A2**: Joint distribution of pair-wise mean firing rate (of time bins) and cosine similarity. **B1**: 2D linear projection of mean centered and normalized spike matrix. **B2**: 2D nonlinear projection of spike matrix (using cosine distance).

Figure S2: Different number of clusters (of significant time bins) tested. A1, B and **C**: as in Figure 3B, E. **A2:** Davis-Bouldin index (Davies and Bouldin, 1979) for the different number of clusters.

Figure S3: Cell assembly detection in L2/3. A-C: as in Figure 3B, D-E. **D:** Jaccard similarity of assemblies (detected across layers, but restricted to L2/3 here) and the ones detected in L2/3 on the left, and number and location: flatmap view on top, depth-profile below of exemplary pairs (pattern A and B responsive ones) of assemblies with high similarity to its right.

Figure S4: Anatomy of thalamocortical synapses. A: Density profile of VPM and POm synapses, digitized from Meyer et al. (2010). **B**: Synapse-to-soma path distances of different thalamocortical synapses on L5 neurons in an exemplary middle assembly (A_8). Ratios of box widths' represent the ratios of the number of synapses' from the inputs (e.g. most synapses on A_8 L5 pyramidal cells are coming from pattern G, in line with the indegree based A_8 membership probability on Figure 4B second).

Figure S5: Synaptic clustering coefficient. A1: Exemplary L5 pyramidal cell and all its afferent synapses from A_{11} (in blue) and from a control group (one out of the twenty) with the same number of presynaptic neurons (in gray). **A2**: Zoom in on A1. Soma, basal dendrites, and proximal apical dendrites are visible. Axon is not shown. The rendering was done with the `BioExplorer` package. **B1**: Distance matrix between all pairs of A_{11} synapses and distribution of nearest neighbour distances (minimum over the rows of the matrix) on its right (see Methods). **B2**: same as B1 but for the control group. The equations on the righmost part of the figure are motivated and explained in the Methods.

Figure S6: Assemblies detected from averaged spike matrices. A-B: as in Figure 3A-B. **C:** as in Figure 6C. **D**: Left: Jaccard similarity of consensus assemblies and average assemblies. Middle: Number of neurons in conesensus assemblies' union and core and average assemblies. Right: As middle, but number of neurons participating in given number of assemblies. **E1**: Jaccard similarity of consensus assemblies at different fraction thresholds (consensus assembly size grows to the right) and average assemblies. **E2**: Detailed comparison of the pair with the highest similarity in E1 at given number of assembly instances contained. (10 means that the consensus assembly neuron is part of 10/10 assembly instances, thus consensus assembly sizes grows to the right again.) Average \ consensus is negligible (419 neurons) and is not shown.

Figure S7: Efficiency of innervation. Probabilities of within assembly memberships for all assemblies. **A**: As in Figure 5B left. **B1**: Similar to Figure 4E left, but with different x-axis (presynaptic population size instead of simplex counts, see B2). **B2**: Illustration of the difference between k-indegree and presynaptic population size.