Diet influences resource allocation in chemical defence but not melanin synthesis in an aposematic moth

Cristina Ottocento[1,2]*, Bibiana Rojas[2,3]*, Emily Burdfield-Steel[2,4], Miriam Furlanetto[2],

Ossi Nokelainen[1,2], Sandra Winters[1], Johanna Mappes[1,2]

*Joint first-authorship

[1]*University of Helsinki, Organismal and Evolutionary Biology Research Programme, Faculty of*

Biological and Environmental Sciences, P.O. Box 65, Viikki Biocenter 3, Finland

[2]*University of Jyväskylä, Department of Biology and Environmental Science, P.O. Box 35,*

40014, Jyväskylä, Finland

[3]*Department of Interdisciplinary Life Sciences, Konrad Lorenz Institute of Ethology, University*

of Veterinary Medicine Vienna, Savoyenstraße 1, 1160, Vienna, Austria

[4]*University of Amsterdam, Institute for Biodiversity and Ecosystem Dynamics, Science Park 904,*

1098 XH Amsterdam, The Netherlands

Abstract

1. Aposematic organisms exhibit warning signals (primary defences) coupled with
   secondary defences, and competing for the same resources may lead to trade-offs in
   resource allocation. For animals that synthesise their chemical compounds *de novo*,
   resources, particularly proteins, can be expected to influence the investment in chemical
   defences and nitrogen-based warning colours such as melanin.

2. We manipulated protein availability in the larval diet of the wood tiger moth, *Arctia plantaginis*, to test how early life resource availability influences relevant life history traits, pigment production, and chemical defences. According to the resource allocation hypothesis, we expected individuals with less melanin (i.e., larger warning signals) to have more resources to allocate to chemical defences.

3. Then, we tested the efficacy of chemical defences of moths raised on diets with different protein content by presenting oat baits soaked with moth defensive fluids to wild-caught avian predators and observed their responses.

4. Early-resource environment affected chemical defences and developmental time of male moths but did not affect melanin expression in the wings. The amount of melanin in the wings, however, correlated with chemical defences: irrespective of the diet they were raised on, more-melanised individuals (i.e., smaller warning signal) produced more deterrent defensive fluids. Thus, the negative correlation between primary and secondary defences arises from more melanised individuals having better chemical defences, and not from resource allocation.

5. We conclude that the resources available in early life have an important role in the efficacy of chemical defences, but melanin-based warning colours are less sensitive to the variability of resources than other fitness-related traits.

1. Introduction

Organisms need to invest simultaneously in various life-history traits such as growth, energy maintenance, reproduction, predator avoidance and protection from pathogens (Stearns, 1992; Roff & Fairbairn, 2007). According to life-history theory, limited resources need to be

distributed among various competing traits, inevitably leading to trade-offs in resource allocation

(i.e., resource-allocation hypothesis; Bazzaz et al., 1987; Stearns, 1989; Glazier, 2002).

Examples include trade-offs between larval and adult traits (Stevens et al., 1999), or between

predator-avoidance and immune function (Rigby & Jokela, 2000). In general, allocation costs

lead to trade-offs between traits that shape the demography of populations (McCauley et

al.,1990; Boggs, 1992) and can significantly impact eco-evolutionary dynamics in nature

(Schroderus et al. 2010; Farahpour et al., 2018).

In aposematic organisms (Poulton 1890), those in which a primary defence (e.g., colour,

odour, sound) is coupled with secondary defences (chemical, morphological, behavioural), the

production of secondary defences may imply trade-offs in the allocation of resources needed for

the primary defence, or for other vital functions (Summer et al., 2015). However, experimental

studies testing trade-offs in resource allocation where both primary and secondary defences are

tested in the same system are rare. A study on the Asian lady beetle, *Harmonia axyridis*, found

no phenotypic correlation between the reflex bleeding response (frequency of secretion of the

defensive chemicals) and the beetle colouration, even though reflex bleeding is costly and affects

life-history traits (Grill and Moore, 1998). Recent work on monarch butterflies *Danaus plexippus*

showed a link between toxin sequestration and warning signals, with male conspicuousness

inversely correlated with oxidative damage (due to an increase in concentrations of sequestered

cardenolides) (Blount et al., 2023). Previous work by Lindstedt et al., (2010) showed that the

aposematic, generalist herbivore *Arctia plantaginis* (hereafter referred to as the wood tiger moth)

develops a paler, orange warning signal when reared on *Plantago lanceolata* with high

concentrations of iridoid glycosides (IG) compared to the more conspicuous, dark red warning

signal of individuals fed on a strain of the same plant with low concentrations of IG. This

suggests that the cost of conspicuousness arose via higher excretion costs rather than via

resource-allocation costs (Lindstedt et al. 2010).

Melanin plays an important role in the warning signals of many aposematic organisms

(Fabricant et al., 2013; Lindstedt, Suisto, Mappes, 2020). Melanin is a phenolic biopolymer

widely present across the animal kingdom. The melanogenesis process in mammals and insects

is regulated by different enzymes and substrates (Sugumaran, 2002), but melanin functions

similarly across species. In vertebrates, including humans, melanin has a protective role against

UV radiation (Jablonski & Chaplin 2017; Nicolaï et al. 2020), and in sexual and protective

colouration (Cuthill et al. 2017). In insects, melanin is also crucial in physiological and

ecological processes such as thermoregulation (True, 2003; Trullas et al., 2007; Lindstedt et al.

2009), immune defence (McNamara et al., 2021), wound healing (Bilandžija et al. 2017) and

predation (Hegna et al. 2013; Majerus, 1998). The expression of melanin is often genetically

based (Ellers and Boggs, 2002; Lindstedt et al. 2009; van't Hof et al. 2019) and is also tightly

related to protein resources (Lee, Simpson & Wilson, 2008). The proteins necessary for the

synthesis of melanin pigments can thus constrain the expression of life-history traits, colouration

or chemical defences (Lindstedt et al., 2020; Galarza 2021), making aposematic signals that

require melanin good candidates for testing the resource-allocation hypothesis.

Here, we use the aposematic wood tiger moth, which has a black melanin pattern that covers

approximately 20 - 70 % of its hindwings and 50 - 80% of its forewings (Hegna et al., 2013), to

investigate how early-life resource availability influences melanin expression in the wings, the

efficacy of chemical defences, and life history traits. In this species, melanin improves

thermoregulation at the expense of warning signal efficacy (Hegna et al. 2013). Thus, if the

synthesis of melanin is costly and uses the same "resource pool" as chemical defences, one

would expect a positive correlation between primary and secondary defences. We manipulated

the protein content of the diet of male wood tiger moths, to investigate how wild-caught, natural

predators (blue tits, *Cyanistes caeruleus*) respond to their chemical defences, and whether the

production of such defences leads to trade-offs with the production of other traits, such as wing

melanisation. We predicted that, compared to males raised on a high-protein diet, males raised

on a low-protein diet would have (1) higher melanisation, resulting from a trade-off between

melanin production and the effectiveness of secondary defences (i.e. more melanised individuals

(less efficient warning signal) have less deterrent chemical defences, and vice versa); (2) less

effective chemical defences, rendering them more palatable to bird predators; and (3) higher life-

history costs in terms of size or developmental time.

2. Materials & Methods

Study species

The aposematic wood tiger moth *Arctia plantaginis*, formerly *Parasemia plantaginis* (Rönkä et

al., 2016), displays a conspicuous colour polymorphism (Watson and Goodger, 1986; Chinery,

1993; Nokelainen, 2013) throughout the Holarctic region (Hegna, Galarza, Mappes, 2015). Male

hindwings across the species distribution range can have a red, yellow or white background with

black patterning that covers variable proportions (ca 20-70%) of the wing. White and yellow

colourations are partly produced by pheomelanin, whereas black is a dopamine-derived

eumelanin (Brien et al. 2022). Previous research has scored melanisation (the amount of

eumelanin) of the hindwings as "plus" (with stripes, high melanin) or "minus" (without stripes,

low melanin) (see Fig. 1), but this arbitrary categorisation does not necessarily represent the true

variation, in which subtle differences are not detectable by the human eye. Long-term natural

frequencies of male wood tiger moth melanin morphs in Estonia (classified by human eye) are

about 47% high-melanin and 53% low-melanin (O. Nokelainen, unpublished data, see Fig. 1).

As capital breeders, wood tiger moth adults do not feed; thus, resource acquisition only occurs

during the larval stage. To avoid predation by avian and terrestrial predators, these moths

produce two different types of defensive fluids: abdominal secretions deter predators such as

ants, and thorax secretions deter birds with no adverse effects on ants (Rojas et al., 2017). The

thoracic fluid contains two methoxypyrazines: 2-sec-butyl-3-methoxypyrazine (SBMP) and 2-

isobutyl-3-methoxypyrazine (IBMP), which are produced *de novo* (Burdfield-Steel et al., 2018).

Insects rearing, larval diet and thoracic fluid collection

Male wood tiger moths were obtained from a laboratory stock founded in 2013 with wild-caught

individuals from Estonia and kept at the greenhouse of the University of Jyväskylä. The stock

was supplemented annually with Estonian wild-caught individuals to preserve genetic diversity.

The greenhouse conditions approximately followed the outdoor temperatures and natural

daylight from May to August in Central Finland: between 20°C and 30°C during the day and

decreased to 15°C-20°C during the night. We picked 10 families that contained both low- and

high-melanin male morphs in the parental and grandparental generations, as classified by eye

(see Fig. 1). Laboratory crosses show that wing melanisation (both pattern and amount) is

strongly heritable (personal observation). After hatching, larvae of each of the 10 families were

kept together for 14 days and fed with lettuce and dandelion (*Taraxacum spp.)*. Then, using a

split family design, they were divided into two artificial diet treatments that differed in protein

content (high protein, low protein; see recipe in supplementary material). For each diet treatment,

we used 240 individuals which were kept in boxes of 10 individuals until reaching the pupal

stage. Each box was checked and watered daily and cleaned when needed. Larvae were fed daily,

ad libitum, with their corresponding diet treatment. Life-history traits (time to pupation, pupal

weight) and the degree of melanin (high +, low -) were recorded for each individual. In total, 480

larvae were followed from eggs to adulthood or death, 177 of which emerged as females and 184

as males. All larvae were grown by Furlanetto (2017) from May to August 2017. When the male

adults emerged from pupation, they were given water and stored at 4°C to slow their metabolic

rate and to maintain their condition; thoracic chemical fluid was then collected in June and July

2017. Prior to fluid collection, moths were kept at 20-25°C for thirty minutes. Then fluid was

extracted by squeezing just below the prothoracic section with tweezers and collecting expelled

fluid in a 10 µl glass capillary. Fluid samples were stored in glass vials at -18°C (Furlanetto,

2017).

Effect of diet manipulation on melanin

To measure variation in wing melanisation, we photographed the 53 male moths whose

defensive fluids were used in the predator response assay. The degree of melanisation varies

continuously and thus we measured the proportion of the hindwing that was melanised on a

continuous scale using image analysis.

Figure 1. Hindwing and forewing melanin variation in male wood tiger moths from Estonia. Left, (+) high-

melanin morph; right, (-) low-melanin morph. See text for details. Image from Nokelainen et al. 2013

(reprinted with permission).

Spread male moths were photographed using an established protocol (Nokelainen et al. 2017)

and scaled to the same resolution (pixels per mm). We used the MICA toolbox (Troscianko &

Stevens 2015) in ImageJ (v. 1.50f) for image analysis. From every image, a set of regions of

interest (ROIs) were selected from the dorsal side of the wings: (a) whole forewing area, (b)

forewing melanised region, (c) whole hindwing area, and (d) hindwing melanised region. The

relative proportion of the wings that were melanised was calculated by dividing the area of the

melanised ROIs by the whole wing ROIs, for the forewings:[(forewing melanised region area) /

(whole forewing area)]; and for the hindwings: [(hindwings melanised region area)/(whole

hindwings area)].

Effect of diet manipulation on the predator response to male's defensive fluid

Blue tits (*Cyanistes caeruleus*) are generalist feeders that have a similar distribution as *Arctia*

plantaginis in Europe and are known to attack this species. Blue tits are common in Finland, and

can be kept briefly in captivity for experiments (e.g., Rojas et al., 2017).

For the predator response assay, we used fluids from 53 male moths fed on high- and low-protein

diets; baits soaked with water were offered to 9 birds as a positive control. The volume of wood

tiger moth thoracic fluid has no effect on blue tit responses (Burdfield-Steel et al. 2019), so we

diluted each fluid sample with water to reach a total volume of 15 µl. The 15 µl was then divided

into two samples of 7 µl each, which were used as bait for the same bird. The same amount of

water was offered to control birds.

The birds used for the experiment were caught at Konnevesi Research Station (Central Finland)

from February to April 2018. The birds were trapped (Ham et al. 2006) then housed individually

in the aviary (see Ottocento et al. 2022).

Birds were first familiarised with the experimental boxes and trained to eat a bait (oat flakes)

(see Ottocento et al. 2022). Each bird then experienced four sequential trials, at 5-minute

intervals, in which they were presented with a plate containing one bait (see Fig. 2 A). The first

and last trials were done with baits soaked in tap water to ensure that the bird was motivated to

eat (first) and that the bird was still hungry (last). In the second and third trials, the bird was

presented with a bait soaked in 7µl of the defensive fluids of the same moth (30 raised under a

high protein diet, 23 raised under a low protein diet). The trial ended two minutes after the bird

had eaten the whole bait, or after a maximum duration of 5 minutes if the bird did not eat the

whole bait. In each trial, we recorded the proportion of the bait eaten (see Fig. 2 B), the beak

cleaning frequency (number of times the bird wiped its beak against a surface, e.g., the perch),

the latency to approach (how long it took for the bird to get close to the plate on which the bait

was offered), the latency to eat after approaching (the hesitation time between approach and

attacking) and the latency to eat.

A   B

Figure 2 A). Experimental setup of the predator assay illustrating the perch, water, camera, light source,

mesh opening for observation and hatch for inserting the plate with the bait (oat flake) into the enclosure.

2 B) Proportion of bait (oat soaked with defensive fluid or water) eaten by the predators.

The palatability was classified into four classes according to the proportion of the bait consumed by the

birds: 0%, 10% (0.10), 50% (0.50), 90% (0.90), and 100% (1).

Statistical analyses

All statistical analyses were done with the software R v. 4.1.2 (R Core Team, 2022) using the

RStudio v. 1.2.1335 interface (RStudio Team, 2019). The level of significance in all analyses

was set at $p < 0.05$.

Effect of diet manipulation on melanin

Our results show that the amount of melanin in the hindwings of wood tiger moths is best

measured as a continuous variable (see supplementary material), so we analysed how protein

content in the diet influences the proportion of the wing that is melanised and the size of the

wings using a linear mixed models with treatment as a fixed factor and family as a random

factor, using the lme4 package (Bates et al. 2015). The total area of the forewings, the total area

of the hindwings, the amount of melanin in the forewings (melanised area forewing/total area

forewing), and the amount of melanin in the hindwings (melanised area hindwing/total area

hindwing) were set as response variables (in separate models); the different types of diet

(high/low protein content) were set as predictors.

Effect of diet manipulation on the predator response to male's defensive fluid

To test the proportion of bait eaten, we used beta regression models using package glmmTMB

(Brooks et al., 2017) with family = beta_family(link= "logit); diet (low protein, high protein), the

amount of melanin, and trial set as fixed effects; bird ID as a random effect; and an offset

included to account for differences in observation time. To test for differences in beak wiping

events per minute, we used generalised linear mixed-effects models (GLMM; Knudson et al.,

2022) with a log link and Poisson distribution, fit by maximum likelihood (Laplace

approximation). The interaction between diet (low protein, high protein), the amount of melanin,

and trial were set as fixed effects, beak wiping frequency was set as the response variable, and

bird ID was set as a random effect. To test the latency to approach, the latency to eat after

approaching and the latency to eat, we used a Cox mixed-effects model using package *coxme*

(Therneau, 2020), fit by maximum likelihood. These three behaviours were set as response

variables; the interaction between diet (low protein, high protein), the amount of melanin and

trial were set as fixed effects; and bird ID was set as a random effect. The behaviours of the

predators were first compared to a water-only control to determine whether moth chemical

defences elicited adverse predator reactions. We used Spearman's rank correlations (Ojala et al.,

2005) to test the correlation between the amount of melanin and predator responses to chemical

defences (proportion of bait eaten, beak wiping, latency to approach, latency to eat after

approaching, latency to eat).

Effect of diet manipulation and hindwing melanin on life-history traits

To test whether male moths' development time is affected by dietary protein and the amount of

melanin, we used a mixed-effects Cox model (package coxme, Therneau et al., 2020), with

development time included as the response variable, diet treatment and melanisation included as

the predictor variables, and family as a random factor. We assessed the effect of diet treatment

on pupae weight with a generalised mixed-effects model (package lme4, Bates et al. 2015) with a

Gamma (link="log") distribution, with diet treatment and melanisation as the predictor variables

and family as a random factor. We used a linear mixed-effects model (package lme4, Bates et al.

2015) with diet and melanisation as predictor variables and family as a random factor to test the

effect of diet on the volume of the thoracic fluid. We tested the correlation between the pupal

weight and the volume of the thoracic fluid using a Pearson correlation. Spearman's rank

correlations (Ojala et al., 2005) were used to test the relationships between the amount of

melanin and the developmental time, pupal weight, and volume of thoracic fluid.

Results

Effect of diet manipulation on melanin

Neither the wing melanin content nor the size of the hind- or forewings ($p > 0.05$; see Table 1S,

2S, 3S, 4S; supplementary material; Fig. 3A, 3B and Fig. 4A, 4B) differed between moths raised

on a high- and low-protein diet.

Figure 3. Hindwing (A) and forewing (B) melanin areas do not differ between individuals raised on low-

and high-protein diets. Boxes show the median and the 25th and 75th percentiles of data distribution.

Vertical lines show the data range.

Figure 4. The size of both hindwings (A) and forewings (B) (in pixels) does not vary as a function of the

protein content in the diet individuals were raised on. Boxes show the median and the 25th and 75th

percentiles of data distribution. Vertical lines show the data range.

Effect of diet manipulation and hindwing melanin on the predator response to male's

defensive fluid

Baits soaked in the chemical defences of moths raised on a high-protein diet were eaten by the

predators in lower proportions than baits soaked in water (control) (coef $\pm$ s.e -2.54 $\pm$ 0.87, z = -

2.91, *p = 0.004*; Fig. 5), while baits soaked in fluid from moths fed with low protein diet did not

differ to the predators' response to baits soaked in water (coef $\pm$ s.e -1.56 $\pm$ 0.87, z = -1.79, *p =*

0.074; Fig. 5).

Irrespective of diet type, the chemical defences of more melanised individuals were also more

deterrent against predators (Spearman's rank correlation, r_s = 0.27; *p = 0.005*, Fig. 6A), as baits

soaked in these defences were eaten in lower proportions than those from less melanised males

(coef $\pm$ s.e -5.20 $\pm$ 1.96, z = -2.66, *p = 0.007*, Fig. 6A). Similarly, the predators' latency to eat

the bait increased with higher amounts of melanin (Spearman's rank correlation, r_s = 0.26; *p =*

0.012, Fig. 6B), indicating an adverse reaction of the predators to the chemical defence of highly

melanised individuals, regardless of the diet they were raised on (coef $\pm$ s.e =-4.92 $\pm$ 2.17, z = -

2.26, *p = 0.024,* Fig. 6B). There was no correlation between the amount of melanin and other

behavioural variables (latency to approach; latency to eat after approach; frequency of beak

wiping; *p > 0.05* for all comparisons; supplementary material).

Figure 5. Baits with defensive fluids from moths raised on a high-protein diet are eaten at a slower rate

than those with water. Boxes show the median and the 25th and 75th percentiles of data distribution.

Vertical lines show the data range.

Figure 6 A) Palatability (proportion of defence fluid-soaked bait eaten by birds) and **B)** latency to eat the

defence fluid-soaked bait compared to hindwing melanisation for male moths raised on high- and low-

protein diets.

We found no significant differences in the other predator behaviours (i.e., latency to approach;

latency to eat after approach; frequency of beak wiping) between birds exposed to baits soaked

in either fluid from moths raised on high-protein or low-protein diets, and those exposed to

water-soaked baits ($p > 0.05$ for all comparisons, see Table 7S, 9S, 11S; supplementary

material).

Effects of diet manipulation and hindwing melanin on the life-history traits

The developmental time from egg to pupa varied among diet treatments: individuals raised on a

low-protein diet took significantly longer to pupate than those raised on a high-protein diet (coef

$\pm$ s.e =-0.92$\pm$ 0.32, z = -2.87, $p = 0.004$). There were no differences in body mass (coef $\pm$ s.e=

0.03 $\pm$ 0.04, t = 0.75, $p = 0.47$, see Table S6) or in the volume of the thoracic defensive fluid

(coef $\pm$ s.e =1.05$\pm$ 0.87, t = 1.2, $p = 0.23$) between individuals raised on high- and low-protein

diets. The volume of the chemical defence did not correlate with the pupae weight (t = 0.96, df =

188, $p = 0.34$). There was no correlation between the melanin amount and the larval

developmental time ($r_s = $ 0.11; $p = 0.435$), the body mass ($r_s = 0.003$; $p = 0.98$), or the volume of

defensive fluid ($r_s = -0.03$; $p = 0.79$).

Discussion

Investigating how resource allocation affects life-history traits and defences is vital for

understanding which factors drive variation in aposematic signals. Phenotypic variation in

aposematic signals is puzzling because predators avoid defended prey that are common but

attack rarer ones (Müller 1878). Trade-offs in resource allocation have been suggested to account

for some of the variation found in natural aposematic populations (Wang, 2011; Burdfield-Steel

et al., 2019; Blount et al., 2023). However, the phenotypic correlations between primary and

secondary defences do not necessarily prove this connection, as interactions between such

complex traits can be shaped by genetic correlations, pleiotropic effects or simply variable

environmental conditions (e.g., food availability, environmental stochasticity, presence of

predators). Therefore, to test the resource allocation hypothesis, we need to first test whether the

productions of primary and secondary defences are costly and, then, whether their development

is linked. To investigate whether those traits are costly, and whether a specific trait (chemical

defence or warning signal) is prioritised over others, we manipulated specific constituents of the

early-life resources of male wood tiger moths. The diet used in the study varied in the amount of

protein, which is an essential compound in different aspects of the larval life history of this

species, and which has been shown to influence the efficacy of the species' warning signal,

immunity and life-history traits (Lindstedt et al., 2020).

Surprisingly, dietary protein levels had no effect on the amount of melanin in the fore- or

hindwings, suggesting that this trait may not be directly influenced by early-life environment.

However, Lindstedt et al. (2020) recently found that the protein content of the diet fed to larval

wood tiger moths directly influenced the amount of forewing melanin of adult males in the

closely related Finnish population. The apparent mismatch between Lindstedt et al. (2020) and

our results could be due to the study design: Lindstedt et al. (2020) used larvae from selection

lines for low and high larval melanisation, such that the phenotypic variation was larger than the

natural variation (and, consequently, the average phenotypes were either rare or missing

completely), likely making the possible costs of melanin easier to detect. In this study, by

contrast, we used the natural variation of phenotypes. Interestingly, Lindstedt et al. did not find

differences in the amount of hindwing melanin between individuals fed with high- and low-

protein diets. Thus, hindwing melanisation and patterning seems to be under strong genetic

control, which is unsurprising because moth hindwings commonly have an important signalling

function (Sargent 1978, Kang et al. 2017; Rönkä et al. 2018). It is also possible that the Finnish

and Estonian populations differ in their degree of plasticity of melanogenesis, but this hypothesis

requires further investigation. Other studies in invertebrates have shown that melanisation is

influenced by the amount of protein content in the diet (Lee et al., 2008; Ethier et al., 2015),

while in vertebrates melanin is less affected by dietary changes (Hill and Brawner, 1998; Lee et

al., 2008). For example, in the African cotton leafworm, *Spodoptera littoralis*, a diet with high

protein content led to more melanised cuticles, faster growth and better antibacterial activity and

survival (Lee et al., 2008). In the forest moth *Malacosoma disstria,* individuals raised under low-

nitrogen availability had decreased melanic pigmentation and smaller size, highlighting the high

costs of melanisation (Ethier et al., 2015). In contrast, in vertebrates such as the tawny owl, *Strix*

aluco, the genetic control of melanin deposition appears to be strictly regulated (Roulin &

Dijkstra, 2003; Mundy & Kelly, 2003; Bize et al., 2006; Hoekstra, 2006; Emaresi et al., 2011).

However, even in vertebrates, this is not always the case and environmental conditions may alter

melanised traits in some organisms (Fargallo et al., 2006).

The amount of melanin in the wood tiger moth wings increases with latitude and altitude

(Hegna et al. 2013), as predicted by the thermal melanism hypothesis, which states that darker

individuals have an advantage under low temperatures because they can warm up faster than

light individuals (Trullas et al. 2007). Convincing support for the fitness advantage of dark

colouration in cold environments has been found in many invertebrates (e.g., Kingsolver 1995;

Ellers & Boggs 2004; but see Rosa and Saastamoinen 2020), but also in cold-blooded vertebrates

(Vences et al. 2002; Stuart-Fox et al. 2017; Azócar et al. 2020). Interestingly, more melanin in

the wings results in a better capacity to absorb radiation and warm-up for flight at the expense of

increased vulnerability to predator attacks due to a reduction in the size of the warning signal

(yellow or white pigment in the hindwings) (Hegna et al. 2013). Similar results were found also

in wood tiger moth larvae, where higher amounts of melanin in the body resulted in improved

thermoregulation at the cost of increased predation (Lindstedt, Lindström, & Mappes, 2008;

2009; Nielsen & Mappes, 2020). It is possible that the Estonian population, situated at lower

latitudes than the Finnish population, is free from the (potentially genetic) constraints imposed

by a colder climate. Moreover, while the melanin pattern in male wood tiger moths seems

genetically regulated and the environmental component is low, our study shows that any dietary

resources beyond what is required for the melanin patterns are allocated elsewhere, such as to

more effective chemical defences.

We hypothesised that the thoracic fluid of males raised on a high-protein treatment would

elicit a stronger predator response than the fluids from moths raised on a low-protein diet, as

shown in a previous study (Furlanetto 2017) where the abundance of pyrazines was higher in

fluids from moths raised on the high-protein artificial diet. An increase in protein content means

an increase in the concentration of nitrogen, an essential element in the synthesis of the pyrazine

molecule (Hodge, Mills & Fisher, 1972; Wong & Bernhard, 1988). Pyrazines have a

characteristic repulsive odour (Rothschild, Moore & Brown, 1984; Guilford et al. 1987; Kaye et

al. 1989; Moore, Brown & Rothschild, 1990) and a deterrent effect on birds (Marples & Roper,

1996; Lindström, Rowe & Guilford, 2001; Siddall & Marples, 2011). In this study, predators

found the chemical secretions from male moths raised on the high protein diet more unpalatable

(i.e., tasted worse) than those from males raised on the low protein diet, which confirms that the

chemical defence is costly. However, we did not find any differences between the predators'

hesitation time (i.e., latency) to "attack" the bait soaked with fluids from male moths raised on

high-protein and the latency to attack baits with water (control), which suggests that the

predators may not perceive a difference in the odour (volatile compounds) of both. These results

agree with previous studies suggesting that the defensive fluids of wood tiger moths may contain

several repulsive compounds, not only pyrazines, that influence defence efficacy in concert (see

also Winters et al. 2021; Ottocento et al. 2022) and affect predator response in different ways

(Rojas et al. 2019; Winters et al. 2021). Moreover, as Ottocento et al. show, the relative amount

of the two methoxypyrazines (SBMP, IBMP) present in the wood tiger moth defensive fluids is

more relevant to predator deterrence than the total amount of pyrazines. Unfortunately, however,

it is not possible to quantify the amount of pyrazines and conduct bird assays with the defensive

fluids of the same individual. Interestingly, the same study (Ottocento et al., 2022) showed that

predator responses are stronger towards the chemical defences of individuals originating from

populations with high predation pressure (e.g., Scotland; Rönkä et al 2020) than towards

individuals from populations with lower predation pressure, even if the total amount of

pyrazines in the defensive fluids does not differ among populations.

Our results show that highly melanised male moths from both diet treatments have more

effective chemical defences than those with less melanin, hinting at a positive correlation

between costly melanin pigments and chemical defences. The positive correlation between the

amount of melanin and the strength of chemical defence implies that there is a negative

correlation between primary and secondary defences. It is possible that more melanised males

may invest more in chemical defences than less melanised males to compensate for their less

effective warning signal. While this is not what we predicted, particularly in a scenario of low

resource availability in early life, previous theoretical work has suggested that such negative

correlation between primary and secondary defences can exist because individuals with the best

secondary defences could survive a predation attempt even if bearing a weak warning signal

(Speed & Ruxton 2005). Empirical examples of aposematic species where the bearers of less

efficient warning signals have seemingly better chemical defences also exist (e.g., Darst,

Cummings, & Cannatella, 2006; Wang, 2011). Although in this species the chemical defence

compounds are not sequestered directly from plants but produced *de novo* (Burdfield-Steel et al.,

2018), the main elements that the moths require to build their defences are still collected from

their food intake. Environmental conditions may therefore have a greater influence on chemical

defences than the pigmentary composition of the hindwings, which is less sensitive to the

variability of resources than other fitness-related traits.

When looking at how early-life resource availability influences wood tiger moths' size, we

hypothesised that males raised on a low-protein diet would have both smaller pupae and smaller

hind- and forewings than those raised on a high-protein diet, as environmental fluctuations and

environmental stress (in this case due to the low amount of protein in the diet) may strongly

affect both insect wing (Bitner-Mathé & Klaczko, 1999) and pupal size (Nguyen et al., 2019).

Our findings, however, reveal no differences in size between males raised on high- and low-

protein diets. This could be because individuals raised on a low-protein diet had a longer

development time than those raised on a high-protein diet, which might facilitate the acquisition

of more resources at the larval stage, even if of poorer quality, allowing these individuals to

ultimately reach the same size as those on a richer diet, as reported by Lindstedt et al. (2017). A

longer developmental time, however, may also lead to increased predation risk (Clancy &

Price,1987; Häggström, & Larsson 1995), higher vulnerability to environmental perturbations

(Tammaru et al., 2001), reduced fecundity (Saastamoinen, Hirai, & van Nouhuys, 2013), and

difficulties to escape a risky environment (Cowan, Houde & Rose, 1996).

In summary, experimental evidence suggests that the production and maintenance of chemical

defences are affected and limited by resource available in the early life, but melanin synthesis is

not. This implies that at least in wood tiger moths, melanin synthesis in adult wings seems to be

less environmentally regulated than pyrazine production, implying that these are not limited by

the same resource pool. Our findings also confirm that resource-dependent variation in chemical

defences is perceived by natural predators, which has not always been shown when variation in

chemical defences is investigated (White and Umbers, 2021). Studying the response of natural

predators, which are the selective agents for defended prey, is key when aiming to understand

how variation in primary and secondary defences is maintained in aposematic species.

Acknowledgements

We are grateful to Helinä Nisu, who assisted with the catching and training of wild birds. We thank Konnevesi Research Station for the use of the facilities. Kaisa Suisto and other greenhouse workers were invaluable in the rearing of the moths. We thank the members of the 'Ecology and Evolution of Interactions' group for the useful discussion on an early version of the manuscript.

Funding

This study was funded by the Academy of Finland the project No. 345091 (to JM) and Societas pro Fauna et Flora Fennica (grant to CO).

Author contributions

JM, BR and EB-S conceived and designed the study. MF, EB-S and BR raised the larvae under the different treatments. MF, ON, CO and SW did the colour analyses. CO, BR and EB-S did the unpalatability assays with birds. CO and BR wrote a first draft of the manuscript with input from JM and EB-S. All authors critically reviewed the manuscript, approved the submitted version, and agreed to be held accountable for the content therein.

Data Accessibility Helsinki repository

Conflicts of Interest - None

Permits - Wild birds were used with permission from the Central Finland Centre for Economic Development, Transport and Environment and licence from the National Animal Experiment Board (ESAVI/9114/ 04.10.07/2014) and the Central Finland Regional Environment Centre (VARELY/294/2015). All experimental birds were used according to the ASAB/ABS Guidelines for the treatment of animals in behavioural research and teaching (Association for the Study of

Animal Behaviour, 2020).

References

Azócar, D. L. M., Nayan, A. A., Perotti, M. G., & Cruz, F. B. (2020). How and when

melanic coloration is an advantage for lizards: the case of three closely-related species of

Liolaemus. Zoology, 141, 125774

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects

Models Using lme4. Journal of Statistical Software, 67(1), 1–48.

https://doi.org/10.18637/jss.v067.i01

Bazzaz, F. A., Chiariello, N. R., Coley, P. D., & Pitelka, L. F. (1987). Allocating resources to

reproduction and defense. BioScience, 37, 58-67.

Bilandžija H., Laslo M., Porter M. L. & Fong D. W. (2017) Melanization in response to

wounding is ancestral in arthropods and conserved in albino cave species. Scientific

Reports 7:17148 https://doi.org/10.1038/s41598-017-17471-2

Bitner-Mathé, B. C., & Klaczko, L. B. (1999). Heritability, phenotypic and genetic correlations

of size and shape of Drosophila mediopunctata wings. Heredity, 83(6), 688-696.

Bize, P., Gasparini, J., Klopfenstein, A., Altwegg, R., & Roulin, A. (2006). Melanin-based

coloration is a nondirectionally selected sex-specific signal of offspring development in the

alpine swift. Evolution, 60(11), 2370-2380.

Blount, J. D., Rowland, H. M., Mitchell, C., Speed, M. P., Ruxton, G. D., Endler, J. A., &

Brower, L. P. (2023). The price of defence: toxins, visual signals and oxidative state in an

aposematic butterfly. Proceedings of the Royal Society B, 290(1991), 20222068.

Boggs, C. L. (1992). Resource allocation: exploring connections between foraging and life

history. Functional Ecology, 6(5), 508-518.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). Modeling zero-inflated count data with

glmm TMB. bioRxiv, 132753. https://doi.org/10.1101/132753

Burdfield-Steel E., Pakkanen H., Rojas B., Galarza J. A., Mappes J. (2018). De novo synthesis of

chemical defenses in an aposematic moth. Journal of Insect Science 18.

Burdfield-Steel E., Brain M., Rojas B., Mappes J., (2019). The price of safety: food deprivation

in early life influences the efficacy of chemical defence in an aposematic moth. Oikos 128,

245-253. doi.org/10.1111/oik.05420

Brien M. N., Orteu A., Yen E. C., Galarza J. A. , Kirvesoja J., Pakkanen H., Wakamatsu K.,

Jiggins C. D., Mappes J. (2022) Colour polymorphism associated with a gene duplication

in male wood tiger moths. bioRxiv 2022.04.29.490025 doi.org/10.1101/2022.04.29.490025

Chinery, M. (1993) Collins guide to insects of Britain and Western Europe. 3rd ed. Harper-

Collins. London

Clancy, KM. & Price, P.W. (1987) Rapid herbivore growth enhances enemy attack: Sublethal

plant defenses remain a paradox. Ecology 68: 733-737

Cowan Jr, J. H., Houde, E. D., & Rose, K. A. (1996). Size-dependent vulnerability of marine fish

larvae to predation: an individual-based numerical experiment. ICES Journal of Marine

Science, 53(1), 23-37.

Cuthill, I. C., Allen, W. L., Arbuckle, K., Caspers, B., Chaplin, G., Hauber, M. E., ... & Caro, T.

(2017). The biology of color. Science, 357(6350), eaan0221.

Darst, C., Cummings, M.E. & Cannatella, D.C. (2006) A mechanism for diversity in warning

signals: conspicuousness versus toxicity in poison frogs. Proceedings of the National

Academy of Sciences 103, 5852–5857.

Ellers, J., & Boggs, C. L. (2004). Functional ecological implications of intraspecific differences

in wing melanization in Colias butterflies. Biological Journal of the Linnean Society,

82(1), 79-87.

Ellers J. & Boggs C. (2002) The evolution of wing color in Colias butterflies: heritability, sex

linkage, and population divergence. Evolution, Vol. 56, No. 4 (Apr., 2002), pp. 836-840

Emaresi, G., Bize, P., Gasparini, J., Piault, R., & Roulin, A. (2011). Plumage polymorphism in

melanin-based coloration: a case study in the tawny owl Strix aluco. Ecology and

conservation of European forest-dwelling raptors, 242-252.

Ethier, J., Gasse, M., Lake, K., Jones, B. C., Evenden, M. L., & Despland, E. (2015). The costs

of colour: plasticity of melanin pigmentation in an outbreaking polymorphic forest moth.

Entomologia Experimentalis et Applicata, 154(3), 242-250.

Fabricant, S. A., Kemp, D. J., Krajíček, J., Bosakova, Z., & Herberstein, M. E. (2013).

Mechanisms of color production in a highly variable shield-back stinkbug, Tectocoris

diopthalmus (Heteroptera: Scutelleridae), and why it matters. PLoS One, 8(5), e64082.

Farahpour, F., Saeedghalati, M., Brauer, V. S., & Hoffmann, D. (2018). Trade-off shapes

diversity in eco-evolutionary dynamics. elife, 7, e36273.

Fargallo A. J., Laaksonen T., Korpimaki E., Wakamatsu K. (2006) A melanin based trait reflects

environmental growth conditions of nestling male Eurasian kestrels. Evolutionary Ecology.

21:157–171

Furlanetto M. (2017) Investigating the cost of visual signals and chemical defences in *Arctia*

plantaginis. Unpublished MSc Thesis, University of Padua – University of Jyväskylä

Galarza, JA. Comparative transcriptomics of albino and warningly-coloured caterpillars (2021)

Ecol Evol. 11, 7507– 7517. https://doi.org/10.1002/ece3.7581

Glazier, D. S. (2002). Resource-allocation rules and the heritability of traits. Evolution, 56, 1696-

1700.

Grill C. P., Moore A. J. (1998) Effects of a larval antipredator response and larval diet on adult

phenotype in an aposematic ladybird beetle. Oecologia 114, 274–282.

Guilford T., Nicol C., Rothschild M., Moore B. P. (1987) The biological roles of pyrazines:

evidence for a warning odour function. Biological Journal of the Linnean Society. 31, 2,

113-128

Ham, A., Ihalainen, E., Lindström, L., Mappes, J. (2006) Does colour matter? The importance of

colour in avoidance learning, memorability and generalisation. Behavioural Ecology and

Sociobiology 60: 482–491

Häggström, H. & Larsson, S. (1995) Slow larval growth on a suboptimal willow results in

high predation mortality in the leaf beetle *Galerucella lineola*. Oecologia, 104, 308–315.

http://www.jstor.org/stable/4221112

Hegna R. H., Galarza J. A., Mappes J. (2015) Global phylogeography and geographical variation

in warning coloration of the wood tiger moth (*Parasemia plantaginis*) Journal of

Biogeography 42, 1469–1481

Hegna R. H., Nokelainen O., Hegna J.R., Mappes J. (2013) To quiver or to shiver: increased

melanization benefits thermoregulation, but reduces warning signal efficacy in the wood

tiger moth. Proceedings of the Royal Society 280: 2012-2812

Hill, G. E., & Brawner III, W. R. (1998). Melanin–based plumage coloration in the house finch

is unaffected by coccidial infection. Proceedings of the Royal Society of London. Series B:

Biological Sciences, 265(1401), 1105-1109.

Hodge J. E., Mills F. D., Fisher B. E. (1972) Compounds of browned flavor derived from sugar-

amine reactions. Cereal Science Today.17, 2, 34-40

Hoekstra, H. E. (2006). Genetics, development and evolution of adaptive pigmentation in

vertebrates. Heredity, 97(3), 222-234.

Jablonski, N. G., & Chaplin, G. (2017). The colours of humanity: the evolution of pigmentation

in the human lineage. Philosophical Transactions of the Royal Society B: Biological

Sciences, 372(1724), 20160349.

Kang, C., Zahiri, R., & Sherratt, T. N. (2017). Body size affects the evolution of hidden colour

signals in moths. Proceedings of the Royal Society B: Biological Sciences, 284, 20171287.

Kaye H., Mackintosh N. J., Rothschild M., Moore, B. P. (1989) Odour of pyrazine potentiates an

association between environmental cues and unpalatable taste. Animal Behaviour 37, 563-

568

Kingsolver, J. G. (1995). Viability selection on seasonally polyphenic traits: wing melanin

pattern in western white butterflies. Evolution, 49(5), 932-941.

Knudson, C., Geyer, C. J., & Benson, S. (2022). Package 'glmm'.

Lee K. P., Simpson S. J. & Wilson K. (2008) Dietary protein-quality influences melanization and

immune function in an insect. Functional Ecology. 22, 1052–1061

Lindstedt, C., Lindström, L., & Mappes, J. (2008). Hairiness and warning colours as components

of antipredator defence: additive or interactive benefits?. Animal Behaviour 75, 1703-

1713.

Lindstedt C. Lindström L, Mappes J, McMillan W. O. (2009) Thermoregulation constrains

effective warning signal expression. Evolution 63-2: 469–478

Lindstedt C., Talsma J. H. R., Ihalainen E., Lindström L., Mappes J. (2010) Diet quality affects

warning coloration indirectly: excretion costs in a generalist herbivore. Evolution 64-1:

68–78

Lindstedt C., Eager H., Ihalainen E., Kahilainen A., Stevens M., Mappes J. (2011) Direction and

strength of selection by predators for the color of the aposematic wood tiger moth.

Behavioral Ecology 580-587

Lindstedt C., Suisto K., Mappes J. (2020) Appearance before performance? Nutritional

constraints on life-history traits, but not warning signal expression in aposematic moths.

Journal of Animal Ecology 2020,494–505.

Lindström, L., Rowe, C., & Guilford, T. (2001). Pyrazine odour makes visually conspicuous

prey aversive. Proceedings of the Royal Society of London. Series B: Biological Sciences,

268(1463), 159-162.

Majerus MEN (1998) Melanism: evolution in action. Oxford University Press, Oxford

Marples, N. M., & Roper, T. J. (1996). Effects of novel colour and smell on the response of

naive chicks towards food and water. Animal Behaviour, 51(6), 1417-1424.

McCauley, E., Murdoch, W. W., Nisbet, R. M., & William S. C. Gurney. (1990). The

Physiological Ecology of Daphnia: Development of a Model of Growth and Reproduction.

Ecology, 71(2), 703–715. https://doi.org/10.2307/1940324

McNamara, M. E., Rossi, V., Slater, T. S., Rogers, C. S., Ducrest, A. L., Dubey, S., & Roulin, A.

(2021). Decoding the evolution of melanin in vertebrates. Trends in ecology & evolution,

36(5), 430-443.

Moore B. P., Brown W. V., Rothschild M. (1990) Methylalkylpyrazines in aposematic insects,

their host plants and mimics. Chemoecology 1, 43-51

Müller F., Über die vortheile der mimicry bei schmetterlingen. Zool Anz 1, 54–55. German

(1878).

Mundy, N. I., Kelly, J., Theron, E., & Hawkins, K. (2003). Evolutionary genetics of the

melanocortin-1 receptor in vertebrates. Annals of the New York Academy of Sciences,

994(1), 307-312.

Nguyen, B., Ponton, F., Than, A., Taylor, P. W., Chapman, T., & Morimoto, J. (2019).

Interactions between ecological factors in the developmental environment modulate pupal

and adult traits in a polyphagous fly. Ecology and Evolution 9, 6342-6352.

Nielsen, M. E., & Mappes, J. (2020). Out in the open: behavior's effect on predation risk and

thermoregulation by aposematic caterpillars. Behavioral Ecology, 31(4), 1031-1039.

Nokelainen O. (2013) Many forms of the wood tiger moth (*Parasemia plantaginis*): selective

heterogeneity favours polymorphic warning signals, University of Jyväskylä.

Nokelainen O., Hubbard N., Lown A.E., Wood L.E., Stevens M. (2017) Through predators'

eyes: phenotype–environment associations in shore crab coloration at different spatial

scales. Biological Journal of the Linnean Society XX:1–14.

Ojala, K., Julkunen-Tiitto R., Lindstrom L., Mappes J. (2005) Diet affects the immune defense

and life-history traits of an Arctiid moth *Parasemia plantaginis*. Evolutionary Ecology

Research 7:1153–1170.

Ottocento, C., Winters, A. E., Rojas, B., Mappes, J., & Burdfield-Steel, E. (2022). Not just the

sum of its parts: Geographic variation and nonadditive effects of pyrazines in the chemical

defence of an aposematic moth. Journal of Evolutionary Biology.

Poulton E. B. (1890) The colours of animals, their meaning and use, especially considered in the

case of insects. New York, D. Appleton and Company

R Core Team. (2022). R: A language and environment for statistical computing. R Foundation

for Statistical Computing. https://www.R-project.org/

Rigby M. C. & Jokela J. (2000) Predator avoidance and immune defence: costs and trade-offs in

snails. Proceedings of the Royal Society 267, 171-176

Rojas B., Burdfield-Steel E., Pakkanen H., Suisto K., Maczka M., Schulz S., Mappes J. (2017).

How to fight multiple enemies: target-specific chemical defences in an aposematic moth.

Proceedings of the Royal Society B 284:20171424

Rojas, B., Mappes, J., & Burdfield-Steel, E. (2019). Multiple modalities in insect warning

displays have additive effects against wild avian predators. Behavioral Ecology and

Sociobiology 73, 1-12.

Rönkä K., Mappes J., Kaila L., Wahlberg N. (2016) Putting Parasemia in its phylogenetic place:

a molecular analysis of the subtribe Arctiina (Lepidoptera). Systematic Entomology 41:44–

853.

Rönkä, K., De Pasqual, C., Mappes, J., Gordon, S., & Rojas, B. (2018). Colour alone matters: no

predator generalization among morphs of an aposematic moth. Animal Behaviour, 135,

153-163.

Rönkä, K., Valkonen, J. K., Nokelainen, O., Rojas, B., Gordon, S., Burdfield-Steel, E., &

Mappes, J. (2020). Geographic mosaic of selection by avian predators on hindwing

warning colour in a polymorphic aposematic moth. Ecology Letters, 23(11), 1654-1663.

Roff, D. A., & Fairbairn, D. J. (2007). The evolution of trade-offs: where are we?. Journal of

evolutionary biology, 20(2), 433-447. https://doi.org/10.1111/j.1420-9101.2006.01255.x

Rosa, E., & Saastamoinen, M. (2020). Beyond thermal melanism: association of wing

melanization with fitness and flight behaviour in a butterfly. Animal Behaviour, 167, 275-

288.

Rothschild M., Moore B. P., Brown W. V. (1984) Pyrazines as warning odour components in the

monarch butterfly, *Danaus piexippus*, and in moths of the genera Zygaena and Amata

(Lepidoptera). Biological Journal of the Linnean Society 23:375-380

Roulin, A., & Dijkstra, C. (2003). Genetic and environmental components of variation in

eumelanin and phaeomelanin sex-traits in the barn owl. Heredity, 90(5), 359-364.

Saastamoinen, M., Hirai, N., & van Nouhuys, S. (2013). Direct and trans-generational

responses to food deprivation during development in the Glanville fritillary butterfly.

Oecologia, 171, 93-104.

Sargent, T. D. (1978). On the maintenance of stability in hindwing diversity among moths

of the genus Catocala (Lepidoptera: Noctuidae). Evolution, 424-434

Siddall, E. C., & Marples, N. M. (2011). The effect of pyrazine odor on avoidance learning

and memory in wild robins Erithacus rubecula. Current Zoology 57, 208-214.

Schroderus, E., Jokinen, I., Koivula, M., Koskela, E., Mappes, T., Mills, S.C., Oksanen, T.A. and

Poikonen, T., (2010). Intra-and intersexual trade-offs between testosterone and immune

system: implications for sexual and sexually antagonistic selection. The American

Naturalist, 176(4), pp.E90-E97.

Speed, M.P. & Ruxton, G.D. (2005) Warning displays in spiny animals: one(more) evolutionary

route to aposematism. Evolution, 59, 2499–2508.

Stearns, S. C. (1989). Trade-offs in life-history evolution. Functional ecology, 3(3), 259-268.

Stearns, S. C. (1992) The Evolution of Life Histories. Oxford University Press, London.

681 Stevens J. D., Hansell M. H., Freel A. J. & Monaghan A. P. (1999) Developmental trade-offs in

682   caddis flies: increased investment in larval defence alters adult resource allocation.

683   Proceedings of the Royal Society 266, 1049-1054

684 Stuart-Fox, D., Newton, E., & Clusella-Trullas, S. (2017). Thermal consequences of colour and

685   near-infrared reflectance. Philosophical Transactions of the Royal Society B: Biological

686   Sciences, 372(1724), 20160345

687 Sugumaran M. (2002) Comparative biochemistry of eumelanogenesis and the protective roles of

688   phenoloxidase and melanin in insects Pigment Cell Res. 2002 Feb;15(1):2-9. doi:

689   10.1034/j.1600-0749.2002.00056.x. PMID: 11837452.

690 Summers K., Speed M. P., Blount J. D., Stuckert A. M. M., (2015) Are aposematic signals

691   honest? A review. Journal of Evolutionary Biology 1-17

692 Tammaru, T., Tanhuanpää, M., Ruohomäki, K., & Vanatoa, A. (2001). Autumnal moth–why

693   autumnal?. Ecological Entomology, 26(6), 646-654.

694 Therneau, T. 2020. Coxme: Mixed effects cox models. R package version.2.

695   http://CRAN.R-project.org/package=coxme

696 True, J. R. (2003). Insect melanism: the molecules matter. Trends in ecology & evolution,

697   18(12), 640-647.

698 Trullas S. C., Wyk J. H.V., Spotila J. R. (2007) Thermal melanism in ectotherms. J. therm. Biol.

699   32, 235–245

700 Troscianko, J., and M. Stevens. 2015. Image Calibration and Analysis Toolbox - a free software

701   suite for objectively measuring reflectance, colour and pattern. Methods in Ecology and

702   Evolution 6:1320–1331.

van't Hof Arjen E., Reynolds Louise A., Yung Carl J., Cook Laurence M. and Saccheri Ilik J. (2019). Genetic convergence of industrial melanism in three geometrid mothsBiol. Lett.152019058220190582

Vences, M., Galán, P., Vieites, D. R., Puente, M., Oetter, K., & Wanke, S. (2002). Field body temperatures and heating rates in a montane frog population: the importance of black dorsal pattern for thermoregulation. Annales Zoologici Fennici 39: 209-220.

Wang, I.J. (2011). Inversely related aposematic traits: reduced conspicuousness evolves with increased toxicity in a polymorphic poison-dart frog. Evolution 65: 1637–1649.

Watson A., Goodger D. T. (1986) Catalogue of the Neotropical tiger moths. Trustes of British Museum (Natural History), London. 71

White, T. E. and Umbers, K. D. L. (2021) 'Meta-analytic evidence for quantitative honesty in aposematic signals', Proceedings of the Royal Society B: Biological Sciences. Royal Society, 288(1949), p. 20210679. doi: 10.1098/rspb.2021.0679.

Winters AE, Lommi J, Kirvesoja J, Nokelainen O and Mappes J (2021) Multimodal aposematic defenses through the predation sequence. Front. Ecol. Evol. 9:657740. doi: 10.3389/fevo.2021.657740

Wong J. M., Bernhard R. A. (1988) Effect of Nitrogen Source on Pyrazine Formation. Journal of Agriculture and Food Chemistry 36: 129-133