1 **Phylogenomic analysis of the Lake Kronotskoe species flock of Dolly Varden charr reveals**
2 **genetic and developmental signatures of sympatric radiation**
3

4 Katherine C. Woronowicz[1], Evgeny V. Esin[2], Grigorii N. Markevich[3], Crisvely Soto Martinez[4],
5 Sarah K. McMenamin[4], Jacob M. Daane[5], Matthew P. Harris[1]*, Fedor N. Shkil[2,6]*
6
Author affiliations
1. Department of Orthopaedics, Boston Children's Hospital and Department of Genetics,
Harvard Medical School, Boston, MA 02115
2. AN Severtsov Institute of Ecology and Evolution, RAS; Leninskiy-33, 119071 Moscow, Russian
Federation
3. Kronotsky Nature Biosphere Reserve; Ryabikova-48, 68400 Yelizovo, Russian Federation
4. Department of Biology, Boston College, Chestnut Hill, MA 02467
5. Department of Biology and Biochemistry, University of Houston, Houston, TX 77204
6. NK Koltzov Institute of Developmental Biology, RAS; Vavilova-26, 119334 Moscow, Russian
Federation

Significance Statement

Dolly Varden Charr (*Salvelinus malma*) radiation in Lake Kronotskoe provides a unique case study of the genetics of adaptation and morphological evolution. We provide first genomic and experimental analyses of this radiation and show that major axes of change may be shaped by developmental constraints.

Abstract

Recent adaptive radiations provide evolutionary case studies, which provide the context to parse the relationship between genomic variation and the origins of distinct phenotypes. Sympatric radiations of the charr complex (genus *Salvelinus*) present a trove for phylogenetics as charrs have repeatedly diversified into multiple morphs with distinct feeding specializations. However, species flocks normally comprise only two to three lineages. Dolly Varden charr inhabiting Lake Kronotske represent the most extensive radiation described for the charr genus, containing at least seven lineages, each with defining morphological and ecological traits. Here, we perform the first genome-wide analysis of this species flock to parse the foundations of adaptive change. Our data support distinct, reproductively isolated lineages with little evidence of hybridization. We also find that specific selection on thyroid signaling and craniofacial genes forms a genomic basis for the radiation. Thyroid hormone is further implicated in subsequent lineage partitioning events. These results delineate a clear genetic basis for the diversification of specialized lineages, and highlight the role of developmental mechanisms in shaping the forms generated during adaptive radiation.

Introduction

The Salmonid fishes of the genus *Salvelinus* represent an exceptional example of parallel evolution and trophic adaptation to new environments (Klemetsen 2010). Charr are remarkably variable with naturally occurring, morphologically distinct populations that exhibit diverse life histories. These include complete and partial anadromy, as well as freshwater riverine and lacustrine residency (Taylor 2016, Lecaudey, Schliewen et al. 2018, Osinov, Volkov et al. 2021). Notably, freshwater resident populations frequently establish species flocks showing stereotypical morphologies, each associated with an ecological niche (Nordeng 1983, Walker, Greer et al. 1988, Sandlund, Gunnarsson et al. 1992, Maitland, Winfield et al. 2007, Simonsen, Siwertsson et al. 2017, Markevich, Esin et al. 2018, Esin, Bocharova et al. 2020, Jacobs, Carruthers et al. 2020). Even within sympatric populations, lineages with varying diets, depth preferences, and disparate spawning intervals are commonly observed (Jonsson and Jonsson 2001, Klemetsen 2010). Within independent lacustrine *Salvelinus* radiations, specific morphological adaptations have repeatedly evolved (Klemetsen 2010). Such a propensity for

repeated, independent radiations make the genus *Salvelinus* an attractive system to dissect the mechanisms that facilitate rapid generation of morphological variation.

Salvelinus charrs have repeatedly radiated into multiple sympatric morphs with distinct evolved feeding specializations analogous to those of both African and South American cichlids (Barluenga, Stölting et al. 2006, Malinsky, Challis et al. 2015, Malinsky, Svardal et al. 2018) and African and Asian barbs (Myers 1960, Nagelkerke and Sibbing 2000, Levin, Simonov et al. 2020). Variable traits include jaw size, mouth position, eye size, pigmentation, and others . The variability within and between, charr radiations proffers important genetic case studies to uncover mechanisms of rapid morphological diversification. Such parallel events suggests that these populations experience similar selective pressures, and/or that there are genetic biases shaping the morphologies. The contributions of standing variation from ancestral populations has been suggested for other radiations, including adaptation of stickleback populations to freshwater environments (Schluter and Conte 2009). In contrast, the genetic mechanisms that enable charr to generate distinct morphologies have not yet been well defined. It has been argued that shifts in developmental timing may underlie these common transitions to specialized morphologies among charrs (Esin, Markevich et al. 2018). In this model, development biases radiations by constraining a common axis of change. Early and pleiotropic developmental shifts may give charr an exceptional capacity for adaptive radiation with only a small number of genetic modifications.

The Lake Kronotskoe radiation of Dolly Varden (*Salvelinus malma*) is unique among charrs, as it contains at least seven reproductively isolated phenotypes associating with different ecological niches (Markevich, Esin et al. 2018). This radiation is currently the broadest observed within resident lacustrine populations of this genus, and indeed, among salmonids more broadly (Markevich, Esin et al. 2018). Recent work has characterized genome-wide changes in species flocks of Alpine whitefishes (Coregonus spp.), however, those flocks comprise up to six species with varying morphologies (De-Kayne, Selz et al. 2022). Lake Kronotskoe is one of several volcanogenic lakes of the Kamchatka peninsula (**Figure 1A**) which formed approximately 12,000 years ago after a lava flow dammed the ancestral river (**Figure 1B**) (Braitseva, Melekestsev et al. 1995). The resulting landlocked Dolly Varden population diversified within the new lacustrine environment, and encompassed prominent changes in craniofacial form supporting new feeding strategies (**Figure 1C, D**) (Markevich, Esin et al. 2018, Esin, Bocharova et al. 2020). A major axis of change is proportional jaw length, seen in piscivorous Longhead and deep-water benthivorous Bigmouth morphs, as well as the modulation of frontonasal proportions found in the Nosed morphs. The deep-water omnivorous Smallmouth morph has an increase in eye size in relative proportion to the cranium as well as reduced jaw size (**Figure 1C, D**). Enlarged eye size is widespread among deep-dwelling resident charr lineages in other lakes and may suggest specialization for foraging in low light conditions (Klemetsen 2010). The varied Lake Kronotskoe charrs show strict natal homing with

some morphs spawning in tributaries and other morphs spawning within the lake (Markevich,
Zlenko et al. 2021). The morphs also exhibit differential annual timing windows for
reproduction, which reinforces reproductive isolation and facilitates sympatric radiation (Esin,
Markevich et al. 2021).
Here, we employ phylogenomics to dissect the genetic context of the Dolly Varden
radiation within Lake Kronotskoe. We parse the shared genetic foundation for the
morphological and physiological adaptations of these specialized lineages.

Genome wide assessment of variability within and between Lake Kronotskoe Dolly Varden
morphs

To investigate variation throughout the charr genome, we performed genome-wide targeted
capture of coding and non-coding loci of Dolly Varden charr from Lake Kronotskoe using a
custom, pan-Salmonidae capture array. Targeted capture of small population pools allows
identification of lineage-defining variation via cross-clade sequence comparisons. Similar
approaches have been recently used to assess variation in Belonifomes (Daane, Blum et al.
2021), rockfishes (Treaster, Deelen et al. 2022) and notothenioids (Daane, Dornburg et al.
2019), allowing for clade-wide analysis of genomic variation and phylogenetic parsing of
selective signals.
The pan-Salmonidae capture-array consists of 558,882 bait sequences designed against
conserved Atlantic Salmon (*Salmo salar*) and Rainbow Trout (*Oncorhynchus mykiss*) genomic
sequences and targets 97Mb of total sequence **(Figure 2 – figure supplement 1)**. Coding regions
constitute 83.8% of targeted regions with conserved non-coding, ultraconserved non-coding,
and miRNA hairpins comprising the remaining 16.2% of targeted regions. With our capture
methodology and analysis pipeline, bait sequences successfully hybridize with targeted
elements harboring up to 15% deviation in sequence identity (Mason, Li et al. 2011), thereby
allowing recovery and subsequent analysis of sequence variability. Pairwise analyses between
homologous loci were used to distinguish fixed and variable loci between morphs. Genes
associating with variation were identified by making strategic comparisons within a phylogeny
focused on character diversification (Daane, Rohner et al. 2015). For outgroup analysis, we
sampled anadromous Dolly Varden charr from the local Kamchatka river basin, which is
adjacent to, but isolated from, that of Lake Kronotskoe (**Figure 1B**). We also sequenced a single
S. leucomaenis individual which was collected from the same catchment as the Dolly Varden
charr to serve as an outgroup.
For each group, we recovered approximately 90% of the targeted elements sequenced
to a mean depth of 25-52 reads (**Figure 2 – Supplementary Table 1**). We detected substantial
variation within each lineage as well as the putative ancestral population (avg π in Dolly Varden
= 0.003) supporting pairwise analysis to detect and compare shared and unique variation within
each lineage. Loci underrepresented in our capture were limited and represented as general
gene classes (**Supplementary Table 2**). These elements recovered with low coverage are similar

to those observed in other broad capture approaches (Daane, Dornburg et al. 2019, Daane,
Blum et al. 2021, Treaster, Deelen et al. 2022).

Genetic differentiation and relationships among Dolly Varden morphs

As a first approach, we conducted Principal Components Analysis (PCA) to visualize the
relationships among our lacustrine and anadromous Dolly Varden samples and outgroup *S.*
leucomaenis. PC1 separates *S. leucomaenis* from Dolly Varden and from the members of the
Dolly Varden species flock (**Fig. 2A**), while PC2 broadly groups lineages according to ecological
niche. Notably, all three Nosed morphs cluster quite tightly.
We investigated relationships among Lake Kronotskoe charrs through reconstruction of
the phylogeny given our substantial sequence data. We used IQ-TREE to derive a phylogeny
from a dataset containing 622,831 variant sites, of which 22,701 variants were informative
(Nguyen, Schmidt et al. 2015). Prior interpretations of this radiation argued for multiple-step
diversification based on changes in resource utilization and accompanying feeding
specializations leading to the extant morphs (Markevich, Esin et al. 2018). Our phylogenomic
data support the existence of the described distinct morphs, with each node having high
bootstrap support (**Figure 2B**). Our analysis clusters Longhead and White morphs within a
lineage that is an outgroup to the clade consisting of the Bigmouth, Smallmouth, and Nosed
morphs. Nosed lineages differentiate as a distinct clade with Nosed 2 as an outgroup to Nosed
lineages 1 and 3. As the Nosed 2 population associates so closely with Nosed 3 via PCA (**Figure**
2A), we focused subsequent analyses on disentangling the differentiation between Nosed
morphs 1 and 3.
The anadromous and resident lacustrine Dolly Varden morphs are clearly genetically
differentiated within ecologically specialized lineages, having pairwise mean Fst consistent with
distinct populations (**Figure 2C**) (**Supplementary File 1**). The greatest pairwise differentiation is
found between the anadromous Dolly Varden charr and lacustrine deep-water Bigmouth
morph (Fst = 0.127). Notably, each lacustrine lineage is more differentiated from the
anadromous Dolly Varden population than from any other lacustrine lineage. Further,
lacustrine lineages are similarly differentiated from the putative ancestral riverine Dolly Varden
population. Within the species-flock, the least differentiated pairing is between the lacustrine
Nosed 1 and Nosed 3 morphs (Fst = 0.047) consistent with their phylogenetic relationship.
Although informative of broad patterns of divergence, these values are likely underestimates of
genetic differentiation due to the conserved-element-based dataset biasing analysis to regions
having an inherent constraint on variation.
Introgression analysis within the clade further supports the existence of distinct
lineages, although some incomplete lineage sorting was detected. Significant introgression was
identified in 80% of the 20 possible trios (DSuite; Holm-Bonferoni FWER < 0.01) in patterns
which deviated from the topology of the phylogeny (**Supplementary Table 3**) (Malinsky,
Matschiner et al. 2021). Despite observed significance, the introgression values were relatively

small compared to the recent timeline of the Lake Kronotskoe radiation. The Bigmouth and
Nosed 1 morphs showed the greatest excess of allele sharing (D_{tree} = 4.2%) (**Supplementary**
Table 3). We further calculated ƒ$_4$-admixture ratios and used ƒ-branch statistics to disentangle
the interrelatedness of admixture signals among morphs that share common internal
phylogenetic branches (**Figure 2D**). The greatest proportions of shared alleles were between
the Bigmouth morph and the Nosed 1 morph (ƒ-branch = 9.0%), between the Bigmouth morph
and the Nosed 3 morph (ƒ-branch = 8.3%), and between the White morph and the ancestor of
the Nosed 1 and Nosed 3 morphs (ƒ-branch = 5.9%). The ƒ-branch statistic further supported the
interpretation that Lake Kronotskoe lineages differentiated while maintaining relative genetic
isolation.
To determine the extent to which Nosed 1 and Nosed 3 share common introgressed loci
with Bigmouth, we calculated D in sliding windows (40 variant windows, 20 variants step-size)
for Smallmouth, Bigmouth, Nosed 1 (mean = 0.046, SD = 0.22) and Smallmouth, Bigmouth,
Nosed 3 (mean = 0.041, SD = 0.21). Anadromous Dolly Varden served as the outgroup for all
analyses of introgression. Among sliding windows with D >= 0.8 (32 of 10,617 windows for S, B,
N1; 28 of 10,790 windows for S, B, N3), we observed shared, isolated regions of the genome
with signatures of admixture (**Supplementary File 2**). Among these, we detected a large interval
with evidence for shared introgression between Bigmouth and Nosed morphs 1 and 3 (mean D
for the interval = 0.53 for S, B, N1; 0.54 for S, B, N3) with of more than one third of windows
with D > 3 SDs from the mean (**Figure 2 – figure supplement 2**). However, most introgressed
intervals were more spatially restricted. One locus potentially associating with craniofacial
differentiation spanned an interval centered on *zyg11b* which has been implicated in
craniofacial microsomia (**Figure 2E**)(Tingaud-Sequeira, Trimouille et al. 2020). Another locus
included multiple genes of interest including, including *zswim*5, which is expressed in cranial
neural crest (Wong, Rebbert et al. 2016), *rnf152*, which is involved in regulating neural crest
formation (Yoon, Kim et al. 2022), and *mc4r*, which is a crucial regulator of appetite and
metabolism via leptin and thyroid signaling modulation (**Figure 2F**)(Decherf, Seugnet et al.
2010)*. This suite of craniofacial and metabolic genes may have functioned like a superlocus
which was spread via shared introgression between the Bigmouth and Nosed lineages and
subsequently reinforced the differentiation of distinct lineages.

Patterns of differentiation between river and lake populations of Dolly Varden charr
To understand shared variation that differentiates the Lake Kronotskoe residents from the
river-caught anadromous population of Dolly Varden, pairwise Fst was calculated per coding
(25,373 genes) and conserved non-coding elements (CNE)(22,575 CNEs) from our targeted
capture (**Figure 3A, B**). Fst values were computed by grouping all variation contained within the
species flock and comparing against the variation contained in the riverine Dolly Varden charr
population (lake versus river). There are 327 genes (**Supplementary File 3**) and 80 CNEs
(**Supplementary File 4**) with Fst > 0.5. Each CNE was assigned to flanking target genes by GREAT

(McLean, Bristor et al. 2010). Some intervals harbor multiple highly differentiating CNEs. For
example, the close proximity of two CNEs led to two hits for *cholinesterase-like* and *dync2h1*
(**Figure 3B**). Interestingly, craniofacial (orange) and thyroid (blue) related genes (5 craniofacial
genes of top 20, 2 thyroid genes of top 20) and CNEs (12 craniofacial CNEs of top 20, 3 thyroid
CNEs of top 20) are enriched across the most differentiating elements (**Figure 3A, B**).
Craniofacial genes assigned to highly differentiating CNEs by GREAT analyses included
both those with predicted roles in neural crest (*tfap2a*) (Rothstein and Simoes-Costa 2020), as
well as frontonasal (*meis2, pitx2*) (Evans and Gage 2005, Fabik, Kovacova et al. 2020) and
splanchnocranium (*faf1*) development (Ma, Zhu et al. 2017). Of note, *pitx2* is also involved in
release of thyroid stimulating hormone from the pituitary (Castinetti, Brinkmeier et al. 2011).
dync2h1 and *tfap2a* appeared more than once in our analyses as putative regulated loci,
suggesting compounding alterations to the regulatory environment at these loci may contribute
to the genetic landscape that distinguishes the riverine Dolly Varden from the lacustrine
morphs. Some CNEs are flanked on both sides by genes implicated in craniofacial development
(*ofcc1* and *tfap2a*, *meis2* and *spred1*, *fam172a* and *nr2f1a*). Genes connected to ion transport
(*myh6, khnyn, or11a1, endod1, cacnb2,* and *hrh1*), and vesicular transport (*man2a1, shld2,*
gdpd2, onecut2, and *clint1a*) according to KEGG and Reactome reconstructions were also well
represented among the top differentiating genes and as targets of the top differentiating CNEs.
The majority of nucleotide changes identified were single SNPs with unknown functional
significance. However, high Fst is reflective of potential selection or bottleneck at the locus.
GO terms were assigned to all genes (2,825 GO terms) and CNEs (1,419 GO terms) that
differentiate anadromous Dolly Varden from Lake Kronotskoe residents (Fst > 0.5). Among GO
terms appearing 6 or more times, specific themes emerged (**Fig. 3C**). Highly differentiating
genes are associated with roles in ion channel regulation and protein trafficking. Highly
differentiating CNEs are associated with brain, kidney, and cartilage development. Other GO
terms shared among highly differentiating coding and noncoding elements include functions
such as DNA binding, transcription factors, and regulation of RNA polymerase II
(**Supplementary file 5**).
Within the candidate genes, there are signals related to regulation of thyroid signaling in
development. For example, the gene *slc26a10,* encodes a sulfate transporter that functions in
thyroid hormone synthesis and also acts downstream of Thyroid hormone receptor alpha
(THRa) (Richard, Guyot et al. 2020). We found that *slc26a10* has segregating nonsynonymous
changes in conserved residues that are highly differentiated between riverine and lacustrine
morphs (**Figure 4A, B**). However, the function of this gene is not well understood, and
alignment-based analyses predict this amino acid substitution to have little effect on function
(neutral by PROVEAN, tolerated by SIFT) (Ng and Henikoff 2003, Choi and Chan 2015). By Fst
score, the next candidate gene, *catl1l,* encodes Cathepsin L1-like and is orthologous to
Cathepsin L1. This gene may be involved in the processing of thyroid prohormone (Friedrichs,

Tepel et al. 2003). Next, splicing factor 3b subunit 4 (*sf3b4)* is a key developmental regulator of
frontonasal patterning and growth and is associated with craniofacial Nager syndrome in
humans (Bernier, Caluseriu et al. 2012, Petit, Escande et al. 2014). Within this locus, we find
synonymous variants at high Fst in all lacustrine morphs, and fixed within benthic groups
(**Figure 5A, B**). These elevated footprints of clustered, highly differentiating variants may
indicate the presence of further modifications within poorly conserved non-coding regions in
linkage to these sites which were not targeted by our bait sequences.

Thyroid hormone signaling activity is associated with shifts in craniofacial form
As modulation of thyroid signaling has been implicated in phenotypic specialization of charrs in
different environments (Esin, Markevich et al. 2021), the pattern of fixation in thyroid-
associated genes and non-coding elements in lacustrine morphs compared with riverine Dolly
Varden warranted further investigation. To address if changes in thyroid metabolism are
associated with different morphs in Lake Kronotskoe, we assessed levels of circulating thyroid
hormone in adult riverine and lacustrine charr individuals from the Lake. Intriguingly, we found
a significant decrease in T_3 (the most genomically active form of thyroid hormone) hormone in
specific lacustrine populations (**Figure 6A**). The pattern of reduced T_3 abundance across the
species flock correlates with a clear change in craniofacial proportions: Nosed and Smallmouth
morphs, with sub-terminally positioned mouths, have significantly decreased T_3 levels. This is in
stark contrast to riverine Dolly Varden charr and the lacustrine White morph, which have
comparatively 'wild-type' craniofacial form, and the piscivorous Longhead and deep-water
benthivorous Bigmouth morphs, which are highly specialized (**Figure 1D**).
Our findings identify a disproportionate number of differentiated loci in Nosed lineages
that are known to regulate thyroid signaling (**Fig. 6B**). We assessed specific mutations identified
as fixed or nearly fixed (Fst > 0.9) within Nosed morphs that differentiate them from Longhead
or White morphs. A candidate locus that differentiates the low T_3 Nosed 1 and Nosed 3 morphs
from the Longhead and White outgroups is a *leptin* homolog (**Fig. 6C**). All salmonids have two
leptin A ohnologs derived from the shared salmonid whole genome duplication. LeptinA2
paralogs encode an N-terminal sequence extended by 66 amino acids compared to LeptinA1 or
B orthologs within salmonids. The nonsynonymous change differentiating Nosed lineages lies
within a conserved residue of the unique sequence of this leptin paralog. As leptin reciprocally
modulates thyroid hormone activity, both endocrine signaling pathways affect global metabolic
activity. Notably, *mc4r*, which shows evidence of allele sharing between the Bigmouth and
Nosed lineages, serves as a relay through which leptin stimulates the thyroid axis, suggesting
that these hormonal axes may be modulated at multiple levels in differentiated lineages
(Decherf, Seugnet et al. 2010). Exogenous leptin and thyroid hormone treatments are
associated with enlarged craniofacial dimensions marking an intriguing relationship between
these pathways and Nosed lineage diversification (Yagasaki, Yamaguchi et al. 2003,

Zimmermann-Belsing, Brabant et al. 2003, Copeland, Duff et al. 2011, Shkil, Kapitanova et al.
2012, Keer, Cohen et al. 2019).
We also identified a highly differentiated variant within a conserved non-coding
enhancer of *otx2b* (**Fig. 6D**). Nosed 1 and Nosed 3 morphs are fixed or nearly fixed for a variant
allele which lies in a predicted *Pax1* transcription factor binding site. *otx2b* is involved in
development of the skull and the anterior pituitary gland, which regulates hormonal signaling
including the thyroid axis (Diaczok, Romero et al. 2008, Bando, Gergics et al. 2020). Due to the
significant overlapping domains of *pax1* and *otx2* expression in the pharyngeal arches and the
oral endoderm from which the pituitary gland arises, the differentiating variant identified
provides a plausible regulatory shift associated with evolution of craniofacial morphology (Liu,
Wang et al. 2013, Liu, Lin et al. 2020).

We asked whether the subterminal mouth positions exhibited by the low-T_3 lineages
(Smallmouth, Nosed 1, and Nosed 3) could be caused by the reduced plasma thyroid hormone
levels in these morphs. We used transgenic thyroid ablation to determine whether
experimental hypothyroidism (McMenamin, Bain et al. 2014) caused any parallel shifts in
subterminal mouth position in the zebrafish system. Indeed, hypothyroid zebrafish showed a
significant shift in the maxilla position, moving ventrally from a supra-terminal position (**Fig. 7A,**
B). Further, to test whether the top candidate genes were altered in a hypothyroid context, we
extracted mRNA from the heads of control and hypothyroid larval zebrafish at 7 and 14dpf and
quantified expression levels by RT-qPCR (**Fig. 7C**). While there is strong genetic differentiation
between the lake morphs and the riverine Dolly Varden for *slc26a10* and *sf3b4*, we did not
detect any significant difference in gene expression levels. We also quantified gene expression
for *otx2b* and its putative regulators, *pax1a* and *pax1b*. While genes differentiated the Nosed
morphs, we did not detect significant differences in gene expression levels in the head.
However, *lepa* was significantly upregulated in the heads of 14dpf hypothyroid larvae,
indicating that under normal developmental conditions, thyroid hormone suppresses *lepa*
expression in the head.

Work in zebrafish previously identified skeletal elements that are phenotypically
sensitive to thyroid hormone titer (Keer, Cohen et al. 2019, Keer, Storch et al. 2022); we used
geometric morphometrics to evaluate these skeletal elements in charrs. Among charr lineages,
we compared the shapes of the dentary, anguloarticulare, hyomandibula and parasphenoid.
Mirroring the pattern of zebrafish, charrs significantly differ in the shape of TH-sensitive bones
(Procrustes Anova for dentary and anguloarticulare $F_{40;944}=40.84$ p<0.0001; $F_{80;1760}=18.88$
p<0.0001, respectively) and display subtle differences in the shape of TH-insensitive bone
(Procrustes Anova for hyomandibula $F_{60;1212}=6.55$ p<0.001)(**Figure 7D, Figure 7 – figure**
supplements 2, 3). The parasphenoid, which forms the neurocranial base, is not variable
(**Figure 7 – figure supplement 4**). These data were supported by pairwise calculation of

Procrustes distances between morphs, which demonstrate significant differences (p<0.001) in
the shape of jaw bones between all morphs, excluding pairs: L-DV and N1-S, and absence of
differences in the shape of parasphenoid between most of morphs, excluding pairs formed by
B, S and piscivorous morphs (L and W). Performing principal component analysis (PCA) on the
dentary and anguloarticulare revealed that the distribution of morphs along PC1 (a component
explaining 66.9% and 57.7% of the variance, respectively) (**Fig. 7B**) corresponded to their
distribution along the T_3 value axis (**Fig. 6A**).

Discussion
Lake Kronotskoe harbors a unique radiation of Dolly Varden charr that provides a powerful new
case to study the genetic and developmental foundations supporting vertebrate radiations.
Previous models centered on morphology, ecology and feeding behavior suggest two lacustrine
clades: a deep-water clade consisting of Smallmouth and Bigmouth morphs, and a shallow-
water (pelagic and littoral) clade composed of Longhead, White, and Nosed morphs (Markevich,
Esin et al. 2018). Our data redefine the evolutionary relationships among the lake morphs and
support differentiated true-breeding lineages.
Lake Kronotskoe arose from a volcanogenic event and presently drains via a waterfall
considered impassable by charr (Viktorovskii 1978). Pairwise Fst analyses found that each lake
lineage is more differentiated from riverine Dolly Varden than from any other lineage.
Furthermore, each lake lineage is differentiated from riverine Dolly Varden charr at roughly
equivalent levels. We determined that the Lake Kronotskoe lineages maintain reproductive
isolation through low hybridization. By contrast, the much older Lake Malawi cichlid radiation,
has ƒ-branch values commonly exceeding 5% and up to 14.2% (Malinsky, Svardal et al. 2018)
and *Coregonus* salmonids also have extensive introgression (De-Kayne, Selz et al. 2022). These
new data suggest that the species flock in Lake Kronotskoe was established by a single founding
population having shared genetic signatures that quickly established and maintained
thoroughly reproductively isolated populations.
We found specific genetic differentiation between lacustrine resident lineages and
riverine Dolly Varden populations with selective signatures in genes regulating craniofacial
development and thyroid function. It is important to note that while PROVEAN and SIFT
predicted the serine substitution shared among the lake morphs would have little consequence
upon *slc26a10* function, such alignment-based prediction methods tend to underestimate
potential effects if variation is shared in other lineages (Ng and Henikoff 2003, Choi and Chan
2015). Indeed, the multiple sequence alignment showed that zebrafish also encode serine at
that residue.
While the relationship to morphological radiation is less clear, the abundance of ion
transport and protein trafficking genes among the highly differentiating genes, suggests that
these processes may also be important drivers of charr evolution. Crucially, ion homeostasis is
central to effective osmoregulation during freshwater adaptation (McCormick, Regish et al.

2019). Proper ion channel expression is also a factor in chondrocyte maturation and
homeostasis (Dicks, Maksaev et al. 2023, Brylka, Alimy et al. 2024). Whether and how protein
trafficking contributes to adaptations in Lake Kronotskoe is less obvious.

The data show that CNEs are particularly influential in the evolution of kidney function
and cartilage morphology within the Lake Kronotskoe radiation. Differentiating CNEs are
associated with development of the brain, kidneys, and cartilage. The kidneys are indispensable
for osmoregulation and obligate freshwater populations of salmonids place different demands
on their kidneys their anadromous counterparts (Tipsmark, Sørensen et al. 2010). Cartilage
templates lay foundations for many craniofacial structures. In concert, variation of these three
traits may contribute to differential behavior, physiology, and morphology characterized within
this adaptive radiation.

Genes that function in thyroid hormone regulation are differentially selected, suggesting
modulation of the hormone may underlie stereotypical phenotypic shifts in lacustrine charr
morphology. As many specialized morphologies are hypothesized to arise from heterochronic
shifts in development (Simonsen, Siwertsson et al. 2017), the thyroid axis may prove to be a
common mechanism underlying the adaptive potential and may contribute to the remarkable
similarity of morphologies exhibited by lacustrine charr species flocks (Esin, Markevich et al.
2021, Esin, Markevich et al. 2021). Indeed, the pairs of morphs in each of the Lake Kronotskoe
clades exhibit alternative heterochronic tendencies, which could result from alterations in
thyroid signaling. For example, the Smallmouth morph with proportionally large eyes and blunt,
rounded rostra shows hallmarks of paedomorphosis, while the sister Bigmouth morph
possesses peramorphic traits such as overdeveloped lower jaw. The enlarged jaws and
frontonasal protrusion of the Longhead morph, as well as the drastically modulated frontonasal
proportion of the Nosed 3 morph are peramorphic features in comparison to their sister
lineages. Similar TH-induced craniofacial changes also arose during the radiation of charrs
dwelling in other lakes and rivers. For example, in Lake El'gygytgyn, there resides an extremely
low TH-content small-mouth charr (*S. elgyticus*) has big eyes and blunt, rounded rostra, while
the closely related boganida charr (*S. boganidae*) with a high TH-content has elongated jaws
(Esin, Markevich et al. 2021, Esin, Shkil et al. 2024). The piscivorous stone charr *S. malma*
lineage, dwelling in sympatry with typical Dolly Varden the Kamchatka river and characterized
by a high TH-level, displays an accelerated rate of ossification of the tooth-bearing bones,
reduced eye size, elongated head, and big mouth as its definitive morphological traits (Esin,
Markevich et al. 2020).

Indeed, thyroid hormone-induced adaptive morphologies are found in phylogenetically
distant fishes, the large African barbs (g. *Labeobarbus*; Cypriniformes; Teleostei), inhabiting
Lake Tana (Nagelkerke and Sibbing 2000). The age of the Lake Tana species flock of barbs is
comparable with the Lake Kronotskoe species flock, yet genetic differences between Lake Tana
morphs are comparatively subtle (de Graaf, Megens et al. 2010, Nagelkerke, Leon-Kloosterziel

et al. 2015), and ecomorphological differentiation is a result of heterochronic shifts presumably

induced by thyroid axis alterations (Shkil, Lazebnyi et al. 2015). Such similarities suggest that

genetic modification of thyroid signaling may be a widespread mechanism facilitating rapid

freshwater teleost adaptive radiations; thyroid modifications could provide a pleiotropic

foundation from which more specialized morphologies may be further elaborated. Our

experimentally induced hypothyroidism lend support to this possibility: ablating thyroid follicles

in the zebrafish creates a shift towards a subterminal mouth position, recapitulating the

morphology of the low-T3 charr lineages. Further, the craniofacial elements variable among the

Lake Kronotske charrs are the same bones that are known to be sensitive to thyroid hormone

alterations in a zebrafish context (Keer, Cohen et al. 2019, Keer, Storch et al. 2022).

Furthermore, the significant increase in *lepa* expression in hypothyroid zebrafish suggests that

endocrine signaling axes may synergize, further expanding the array of potential adult

morphologies attainable along a shared axis of change.

In anadromous salmonids, smoltification in preparing for migration out to sea requires

orchestration of hormonal and physiological switches. A hypothesis stemming from lacustrine

populations is that selective pressure on the ancestral riverine charr population was relaxed

upon colonization of the lake, as the newly resident population of charr became obligate

freshwater residents. Such an initial shift in developmental programs may constitute a common

node among lacustrine-locked charr, biasing the direction of adaptation to generate similar

forms among independent lineages, and thereby laying the foundation upon which more

trophically specialized morphologies may arise. In this context, exclusion of the smoltification

stage from the charr life cycle may permit lacustrine adaptive diversification. Endocrine

signaling pathways, including the growth, hypothalamic-pituitary-interrenal, and thyroid axes

drive physiological, morphological and behavioral changes during smoltification. In landlocked

salmonid populations, the growth and hypothalamic-pituitary-interrenal axes, which ordinarily

induce physiological changes for a transition to seawater, are upregulated, while the thyroid

axis likely maintains its developmental, physiological, and adaptive significance (McCormick,

Regish et al. 2019). In support of this model, thyroid hormone signaling is selectively modified

during freshwater colonization and subsequent adaptive radiations of the threespine

stickleback, *Gasterosteus aculeatus* (Kitano, Lema et al. 2010).

The lineage-specific pattern of highly differentiating loci identified in Nosed morphs,

suggests that an initial developmental state of extensive modifications to thyroid signaling and

craniofacial development shared among all lineages, was further refined in these lineages. The

fixation of variation in *leptina2* and a predicted *otx2b* regulatory region found in Nosed morphs

over other lake groups suggests further modulation of the thyroid signaling in these lineages.

The data are supported by findings of altered T_3 levels within these lineages as adults and the

presence of *mc4r* within an interval of excess allele sharing with the Bigmouth morph. Thus,

beginning with an initial suite of shared genetic variants, lineage-specific, secondary

elaborations may have accumulated and further catalyzed the exceptional species flock
diversification in Lake Kronotskoe.
Such repeated and parallel derivations of morphotypes across the *Salvelinus* complex
suggest that there is an underlying genetic framework biasing the radiations of resident
lacustrine populations. Our data suggest that shared modulation of the thyroid signaling axis in
tandem with craniofacial regulators may enforce such biases. The patterns of variation
identified in the Lake Kronotskoe radiation point to a fundamental genetic groundwork for
craniofacial evolution and a common axis for morphological change.

Methods
Field material collection
Charrs that passes the spawning season, adults without spawning changes in color and head
and reproductive states, were sampled in Lake Kronotskoe. Adult riverine Dolly Varden charrs
were collected in the nearest watercourse draining the opposing slope of Valaginskiy range.
Blood, pectoral fin tissue and pictures were collected. Blood for thyroid hormone test was
carefully collected from the caudal vessel with a Vacuette serum tube. The distal part of the
right pectoral fin (finclip, 0.2-0.3 cm^2) was taken with scissors and fixed in pure alcohol for DNA
analysis. Fish were photographed, treated with the antibacterial and antifungal solution
(Melafix and Pimafix, API) for 30 min, and released if the fish did not display any signs of injury
and/or infection in 48 hrs. All catches were carried out in accordance with the Russian Federal
Act on Specially Protected Natural Areas (OOPT) N33-ФЗ 14/03/1995, article 10, and Plan of the
research activities of Kronotsky Nature Reserve. The procedures with fish were approved by the
Bioethics Commission of the AN Severtsov Institute of Ecology and Evolution, Russian Academy
of Science.
456
Phylochip targeted sequence enrichment design
We aimed to create a pan-Salmoniformes targeted sequence capture design that can enrich
sequencing libraries for conserved genetic regions across a broad diversity of available salmon
genomes. This design targets protein-coding exons as well as a set of conserved non-protein
coding elements (CNEs), miRNA hairpins, and ultraconservative non-coding elements (UCNEs).
The majority of capture baits were derived from the Atlantic salmon genome (*Salmo salar*,
ICSASG_v2)(Davidson, Koop et al. 2010), with inclusion of regions from the genome of rainbow
trout (*Oncorhynchus mykiss*, AUL_PRJEB4421_v1)(Berthelot, Brunet et al. 2014) that were
either not represented in the Atlantic salmon genome or were <85% identity to a capture target
within the rainbow trout genome. As these fish bracket both sides of the salmon phylogeny
(**Supp. Fig. 1A**), the 'Phylochip' design strategy enables DNA from the majority of salmonids
target regions to be efficiently enriched using this one capture design.
As the Atlantic salmon genome was not annotated at the time of capture design,
annotated coding sequences were isolated from the rainbow trout and northern pike (*Esox*

lucius, GCF_000721915.2_ASM72191v2)(Rondeau, Minkley et al. 2014) genomes. These were
then identified within the Atlantic salmon genome via BLASTN (ncbi-blast-2.2.30+; parameters '-
max_target_seqs 1 -outfmt 6'), and these hits used in the capture design. Genes from rainbow
trout that were not identified in Atlantic salmon or that had <85% identity to the best BLAST hit
within the Atlantic salmon genome were also retained in the capture design. CNEs were defined
from the constrained regions in the Ensembl compara 11-way teleost alignment (Ensembl
release-84)(Herrero, Muffato et al. 2016). To reserve space in the capture design, only CNEs
≥75bp in length were included in the capture baits. These CNEs were extracted from the
Japanese medaka (*Oryzias latipes*, MEDAKA1), three-spined stickleback (*Gasterosteus*
aculeatus, BRAOD S1), and zebrafish (*Danio rerio*, GRCz10.84) genomes using Bedtools (v2.23.0)
intersectBed (Quinlan and Hall 2010). miRNA hairpins were extracted from miRbase and
ultraconservative elements (UCNEs) from UCNEbase (Kozomara and Griffiths-Jones 2010,
Dimitrieva and Bucher 2013). As with protein coding exons, these elements were identified
within each reference genome using BLASTN (ncbi-blast-2.2.30+; parameters '-max_target_seqs
1 -outfmt 6'). miRNA hairpins were padded to be at least 100 bp to improve capture specificity.
From these targeted regions, the specific SeqCap EZ Developer (cat #06471684001)
oligonucleotide capture baits were made in collaboration with the Nimblegen design team.
Capture baits are strategically designed to standardize oligo annealing temperature, remove
low complexity DNA regions and to reduce the oligo sequence redundancy. The capture design
targeted sequence from 558,882 genomic regions (97,049,118 total bp) across the two
salmonid genomes. This included including 460,210 protein coding exons, 93,973 CNEs, 1,082
miRNAs and 3,617 UCNEs (**Supp. Fig. 1**).
493
DNA extraction and preparation of sequencing libraries
Tissue from finclips was digested and genomic DNA was column purified using QIAGEN DNeasy
Blood & Tissue Kit (QIAGEN 69506). Genomic DNA was extracted from finclips of 1 *S.*
*leucomae*nis, 3 riverine Dolly Varden charr, 8 Bigmouth morphs, 10 Longhead morphs, 5
Nosed1 morphs, 7 Nosed2 morphs, 5 Nosed3 morphs, 6 Smallmouth morphs, and 6 White
morphs. Pools of genomic DNA were produced for each lineage such that genomic DNA from
every individual in a lineage pool was equally represented. The pooled samples were sheared to
a target size of 200bp in Tris-HCl EDTA shearing buffer (10mM Tris-HCl, 0.1mM EDTA, pH 8.0).
Mechanical shearing was performed using a Covaris E220 ultrasonicator (duty cycle, 10%;
intensity, 5; cycles/burst, 200; time, 400 seconds; temperature, 8°C) and Covaris microTUBE
Snap-Cap tubes (Covaris 520045). Sequencing libraries were produced using the KAPA
HyperPrep Kit (Roche 07137923001) using 500ng of starting material for each library. Library
preparation was conducted by following the SeqCap EZ HyperCap Workflow Version 1.2. The
sequencing library for the Nosed2 samples utilized enzymatic shearing using the KAPA
HyperPlus Kit (Roche 07962401001) and 100ng of starting material (SeqCap EZ HyperCap

Workflow Version 3.0). Fragment size and DNA concentration were quantfied using Agilent
2100 BioAnalyzer and High Sensitivity DNA Chips (Agilent 5067-4626). Paired-end, 150bp
Illumina HiSeq sequencing was performed on a pool consisting of multiple barcoded libraries.
512
Trimming Adapters and Read Mapping
Illumina adapter sequences were removed from reads using Trimmomatic v0.36 (Bolger, Lohse
et al. 2014). Trimmed and masked reads were aligned to the *Salvelinus alpinus* reference
genome (RefSeq Assembly accession: GCF_002910315.2) using NextGenMap v0.5.5 (Sedlazeck,
Rescheneder et al. 2013). The flag - - strata 1 was so only the highest scoring read mappings
were recorded in the alignment file.
519
Variant Calling and Filtering
The variants used to reconstruct the phylogeny and to conduct the principal components
analyses were derived from sample of *S. leucomaenis*, anadromous Dolly Varden, and all 7
members of the species flock. samtools v1.15.1 was used to fix mates, mark duplicates, and
filter reads below the minimum mapping quality set to -q 30 (Danecek, Bonfield et al. 2021).
bcftools v1.13 was used to call and filter variants (Danecek, Bonfield et al. 2021). Only the
variants with quality scores >= 20, depth of coverage on a per-sample basis between 10 and
500 reads, fraction of missing genotypes F_MISSING <= 0.72. SNPs within 2bp of indels and
other variant types were excluded, and minor allele frequency > 0.05. The quality filtered VCF
file contained 623,619 variants. The set of variants used for introgression analyses,
quantification of Fst, π, and GOterm enrichment analyses were called and filtered from
alignments of the anadromous Dolly Varden, Bigmouth, Longhead, Nosed 1, Nosed 3,
Smallmouth, and White lineages (*S. leucomaenis*, and Nosed 2 were excluded). The calling and
filtering criteria are identical to the conditions described above except for the depth thresholds.
Those were filtered on a per site basis for coverage between 70 and 3500 reads. This VCF file
contained 526,811 variants.
536
537   **Coverage of Targeted elements**
538   Coverage statistics were derived using the BEDTools v2.21.1 coverage function (Quinlan and
539   Hall 2010). Alignment files were intersected with a bed file containing the positions of each
540   targeted element. From this intersection, the average depth of coverage was quantified per
541   base.
542
Deriving Phylogeny
The phylogeny was derived using IQ-TREE v1.6.12 (Nguyen, Schmidt et al. 2015) . The input
consisted of 622,831 nucleotide sites, including 22,701 parsimony informative variants. The
ModelFinder function (Kalyaanamoorthy, Minh et al. 2017) determined the base substitution

model of best fit to be a transversion model with empirical base frequencies and a proportion
of invariable sites (TVM + F + I). 1,000 ultrafast bootstrap replicates (Hoang, Chernomor et al.
2017) quantified support for the phylogeny.
550
Principal Component Analysis of Sequence Variation
552 PLINK v1.90b7 was utilized to conduct principal component analysis (Purcell, Neale et al. 2007).
553 Linkage pruning was conducted using 50kb windows, 10bp step size, and $R^2 > 0.1$. The linkage
554 pruned dataset consisted of 65,488 variant sites. The PLINK eigenvector and eigenvalue outputs
555 were plotted in R.
556
Introgression Analysis
558 Introgression was quantified for all trios in the phylogeny using Dsuite v0.4 (Malinsky,
559 Matschiner et al. 2021) Dtrios. Riverine Dolly Varden was specified as the outgroup. The f-
560 branch statistic was depicted as a matrix by taking the output from Dsuite Fbranch and running
561 the Dsuite dtools.py script to generate a plot. Dinvestigate was used to generate sliding
562 windows of 40 variants per window and 50% overlap.
563
Calculating pairwise Fst and Tajima's Pi
To quantify genetic differentiation, pairwise Fst was calculated using PoPoolation2 v1201
(Kofler, Pandey et al. 2011). The software package allows Fst to be quantified in sliding windows
or in a genewise manner. Sequencing alignment data were converted into the mpileup format
using SAMtools v1.13 (Li, Handsaker et al. 2009). The PoPoolation2 program mpileup2sync (--
min-qual 20) generated the sync file used as input to calculate Fst in sliding windows.
Popoolation2 calculated Fst based on allele frequency (Hartl, Clark et al. 1997). The
PoPoolation2 program fst-sliding (--min-coverage 20 --min-count 3 --max-coverage 200) was
used to calculate Fst in non-overlapping sliding windows. To assess which genes were most
differentiating between populations, the function create-genewise-sync was utilized to
intersect the sync file with a gtf containing all targeted regions in the *S. alpinus* genome, and
filtered according to the same depth and minor allele count criteria as sliding windows
analyses. Prior to filtering for depth, Fst was calculated for 59,478 genes and 22,590 CNEs.
Noncoding elements were associated with putative regulatory targets by following the GREAT
workflow to establish basal regulatory windows. A BED file of CNE loci was intersected with
Intervals spanning 5kb upstream of and 1kb downstream from transcriptional start sites, with
up to a 1 Mb extension (McLean, Bristor et al. 2010). To quantify nucleotide diversity, Tajima's
Pi was calculated using PoPoolation v1.2.2 (Kofler, Orozco-terWengel et al. 2011). The software
also enables quantification of Tajima's Pi in sliding windows. The same depth criteria that were
used for Fst sliding window quantifications were used to calculate Tajima's Pi in sliding windows
for individual lineages.

ELISA for thyroid hormone in blood

ELISA for thyroid hormone in blood
Serum samples were transferred to 2 ml specimen tubes and centrifuged at 12 000 g for 10 min
with Eppendorf MiniSpin. Serum was then collected into Eppendorf 1.5 ml tubes and placed in
in freezer at 24-26C. The total triiodothyronine (T3, bioactive form of thyroid hormone)
concentration in plasma was evaluated by enzyme-linked immunosorbent assay (Monobind
Total Triiodothyronine (tT3) test system, Monobind Inc, USA) and measured the hormone in
accordance with the manufacture protocol using StatFax 303 Plus strip reader (Awareness
Technology Inc, USA).

Zebrafish thyroid follicle ablations
Danio rerio were all of the line *Tg(tg:nVenus-v2a-nfnB)* (McMenamin, Bain et al. 2014). Briefly,
clutches of transgenic embryos were sorted for nVenus expression at 4 dpf then treated
overnight with either 1% DMSO (for control euthyroid fish) or with 1% DMSO and 10 mM
metronidazole, which induces conditional thyroid ablation in the *nfnB*-expressing thyroid
follicle cells. Thyroid ablation was visually confirmed at 5dpf.

Quantification of maxilla position
AMIRA (version 6.0.0) was used to visualize μCT scans of adult zebrafish skulls (Blythe, Nguyen
et al. 2022, Nguyen, Lanni et al. 2022). A line was drawn intersecting the parasphenoid at its
proximal- and distal-most points to approximate the long-axis of the body. A perpendicular line
was drawn from the dorsal-most position of the maxilla to intersect the parasphenoid axis. For
each individual, this distance was normalized to the standard length. Samples were selected to
be roughly equivalent in size with standard lengths ranging from 19.0mm to 20.5mm. n = 10
DMSO control individuals and n = 11 MTZ treated individuals.

Gene expression quantification
Euthyroid controls and hypothyroid siblings were decapitated at 7 or 14 dpf posterior to the
operculae, and three sets of 20 heads for each condition were stored in RNA*later*™ Stabilization
Solution (Thermo Fisher, Waltham MA, USA) at -20°C. RNA was extracted using a *Quick*-RNA™
Microprep Kit (Thermo Fisher, Waltham MA, USA) and cDNA libraries synthetized using
SuperScript™ IV Reverse Transcriptase (Thermo Fisher, Waltham MA, USA). Using primer
sequences (**Table 1**) for *actinB1*, *lepa*, *otx2b*, *pax1b*, *pax1a*, *sf3b4*, and *slc26a10*, qPCR was
performed with PowerUp™ SYBR™ Green Master Mix (Thermo Fisher, Waltham MA, USA) on a
QuantStudio™ 3 Real-Time PCR System (Thermo Fisher, Waltham MA, USA) with three technical
and biological replicates. Results were analyzed using DataConnect Software. Relative gene
expression was calculated using the ΔΔCT method with *actinB1* serving as the housekeeping
gene (Livak and Schmittgen 2001).

Target	Forward Primer	Reverse Primer
lepa	TGACGGGCAAAATTTACTTCCA	AGTGTGGATAGATCTCGGCG
otx2b	CAAGCAACCACCTTACACGG	GAGGAGTCGCTGGGTATCC
pax1b	AGTACACCCAGGCTTCATCA	TGTCCACCGTAAACACCGTA
pax1a	TTGGGGTGTCAATAGAGCGA	GTCGACGAAGGCTGAGGG
sf3b4	ACAGGACAACACCAGGGTTAT	GGGCTTGCCGTAAAGTTTGA
slc26a10	CTGCTTCACAAGAGACTGCC	AAAGCAAACGCCATCCCTTG
actinB1	CGACCAGAAGCGTACAGAGA	AATCCCAAAGCCAACAGAGA

Table 1. Sequences for primers used in qPCR.

Geometric morphometrics

Pictures of dry osteological samples (14-27 of each morph) were used for landmarking

(Saltykova, Markevich et al. 2015). Using TPSdig v2.0 (Rohlf 2015), we digitized landmarks (LM),

most of which have been used for the homologous bones of zebrafish (Keer, Storch et al. 2022)

and charrs (Guðbjörg Ósk, Laura-Marie von et al. 2024): six LM for dentary, ten LM for

anguloarticulare; eleven LM for hyomandibula; and eight LM for parasphenoid (**Figure 7 –**

figure supplement 1). Shape analysis was performed in MorphoJ v1.06d (Klingenberg, 2008).

We implemented Generalized Procrustes superimposition and assessed variation in the shape

with Principal component analysis (PCA). For better visualization of shape variability along

PC1/PC2, we created a wireframe mesh connecting landmarks. To estimate the shape

differences between the morphs, we implemented the Procrustes ANOVA and Canonical

Variate (CV) analysis with a calculation of pairwise Procrustes distances (10,000 permutation

rounds).

Acknowledgements

We owe thanks to Dr. Stacy Nguyen for providing uCT data and to Dr. E.A. Saltykova for

providing images of dry osteological samples.

Competing interests

No competing interests declared.

Funding

This work was supported by National Institutes Health [R35GM146467 to SKM, 5F32DE029362

to KCW] and the National Science Foundation [NSF 1845513 to SKM]

Data availability

We will make relevant sequencing files available on Zenodo.

Figure 1

655
656
Figure 1. Lake Kronotskoe radiation of Dolly Varden Charr. (A) A map of the Kamchatka
Peninsula. Lake Kronotskoe is centered within the highlighted box. **(B)** A map of Lake
Kronotskoe geography and locations from which anadromous (blue dots) and resident
lacustrine (red dots) Dolly Varden morphs were collected. **(C)** The Lake Kronotskoe morphs
from left to right: Longhead (L), Nosed lineages (N), Smallmouth (S), White (W), Bigmouth (B).
(D) Detailed images of representative adult heads for the seven sequenced lineages of the
species flock.

Figure 2

665
666
Figure 2. Phylogenetics and population differentiation among the sequenced lineages. (A)
PCA plot of the seven lineages from the lacustrine species flock, and outgroups anadromous
Dolly Varden and *S. leucomaenis*. PC1 distinguishes *S. Leucomaenis* from anadromous Dolly
Varden and from the lake morphs. Note the tight clustering of the three Nosed morphs. **(B)**
Phylogenic relationship of Lake Kronotskoe Dolly Varden lineages; anadromous *S. leucomaenis*
from the Kamchatka River serve as the outgroup. **(C)** Distribution of genome-wide, pairwise Fst
values calculated for non-overlapping 10kb sliding windows (**Supplementary File 5**). **(D)** The
branch-specific statistic f_b shows evidence for elevated gene flow between the Bigmouth and
Nosed 1 morphs, the Bigmouth and Nosed 3 morphs, and the ancestor of the Nosed lineages
and the White morph. White cells represent combinations for which p-values are not
significant. Gray cells represent arrangements which are not topologically feasible for
calculating ƒ-branch scores. **(E)** Sliding window plots of D for trios consisting of Smallmouth (S),
Bigmouth (B), and Nosed 1 (N1), of S, B, and Nosed 3 (N3), and of N3, N1, and B. Positive D
values indicate an excess of the ABBA pattern (red arrows), while negative values indicate an

excess of the BABA pattern. The three plots show a common pattern of excess of allele sharing
overlapping with *zyg11b* between B and N1 and B and N3, while there is no excess of allele
sharing between B and N1 over B and N3. Horizontal lines signify 3SDs from the mean. Genes
contained by the highlighted peak represented by red bars below plots. **(F)** three genes of
interest, *zswim5*, *rnf152*, and *mc4r*, possibly constitute a supergene contained by a shared peak
between B and N1 and B and N3.
687

Figure 2 – Supplementary Figure 1

689
690

Figure 2 – Supplementary Figure 1. Design of pan-Salmoniformes targeted capture array. (A)
Conserved bait sequences were derived from Atlantic salmon (*Salmo salar*) and rainbow trout
(*Oncorhynchus mykiss*) reference genomes. **(B)** Relative contributions of *S. salar* and *O. mykiss*
derived bait sequences to the capture array. **(C)** Classifications and relative abundances of
conserved elements targeted for capture.

697 **Figure 2 – Supplementary Figure 2**

698
699
Figure 2 – Supplementary Figure 2. Sliding window plots of D showing a *large* interval of
excess allele sharing. Sliding window plots for trios consisting of Smallmouth (S), Bigmouth (B),
and Nosed 1 (N1), of S, B, and Nosed 3 (N3), and of N3, N1, and B. Positive D values indicate an
excess of the ABBA pattern (red arrows), while negative values indicate an excess of the BABA
pattern. The three plots show a common pattern of excess of allele sharing overlapping with
between B and N1 and B and N3, while there is no excess of allele sharing between B and N1
over B and N3. Horizontal lines signify 3SDs from the mean.
707

708   **Figure 2 – Supplementary Table 1**

Morphotype	No. Individuals Pooled	No. Reads (million)	No. Reads Mapped	Median Depth	Mean Depth	"Conservome" Coverage	
						2x	10x
Bigmouth	8	72.3	35.3	66 reads	88 reads	91.6%	88.5%
Dolly Varden	3	64.1	32.4	61 reads	82 reads	91.6%	88.3%
Longhead	10	46.1	22.8	41 reads	56 reads	91.1%	85.7%
Nosed 1	5	68.8	34.9	65 reads	87 reads	91.7%	88.5%
Nosed 2	7	49.5	17.5	16 reads	20 reads	88.7%	65.5%
Nosed 3	5	68.4	34.1	63 reads	84 reads	91.6%	88.5%
Smallmouth	6	43.1	21.1	38 reads	52 reads	90.8%	84.9%
S. leucomaenis	1	55.9	25.5	50 reads	67 reads	90.5%	85.9%
White	6	36.9	18.5	34 reads	46 reads	90.6%	83.8%

709
710

711   **Figure 2 – Supplementary Table 1.** Summary of reads aligned and targeted element coverage.

713 **Figure 2 – Supplementary Table 2**

GO name	GO accession	bin size
activation of MAPK activity	GO:0000187	15
anaphase-promoting complex	GO:0005680	51
anaphase-promoting complex-dependent catabolic process	GO:0031145	41
ATP-dependent chromatin remodeling	GO:0043044	133
ATPase activity	GO:0016887	262
calcium ion binding	GO:0005509	1185
calcium, potassium:sodium antiporter activity	GO:0008273	52
carbohydrate binding	GO:0030246	111
carbohydrate metabolic process	GO:0005975	204
cell adhesion	GO:0007155	541
chemokine activity	GO:0008009	72
cullin-RING ubiquitin ligase complex	GO:0031461	23
DNA binding	GO:0003677	2972
DNA integration	GO:0015074	1571
DNA-templated transcription, initiation	GO:0006352	36
ferric iron binding	GO:0008199	24
fructose-bisphosphate aldolase activity	GO:0004332	27
hexose metabolic process	GO:0019318	34
homophilic cell adhesion via plasma membrane adhesion molecules	GO:0007156	362
iron ion transport	GO:0006826	23
isomerase activity	GO:0016853	45
MAP kinase activity	GO:0004707	52
nucleic acid binding	GO:0003676	2731
oligopeptide transport	GO:0006857	20
phosphatidylinositol metabolic process	GO:0046488	49
phosphatidylinositol phosphate kinase activity	GO:0016307	47
receptor tyrosine kinase binding	GO:0030971	28
regulation of mitotic metaphase/anaphase transition	GO:0030071	44
sensory perception of sound	GO:0007605	69
skeletal muscle fiber development	GO:0048741	44
SWI/SNF complex	GO:0016514	147
transcription coactivator activity	GO:0003713	164
transcription factor TFIID complex	GO:0005669	38
transmembrane transport	GO:0055085	1148
transmembrane transporter activity	GO:0022857	306
transporter activity	GO:0005215	218
transposition, DNA-mediated	GO:0006313	1551
ubiquitin protein ligase binding	GO:0031625	28
vesicle-mediated transport	GO:0016192	250

715 **Figure 2 – Supplementary Table 2. Table of significantly underrepresented GO terms.** This
716 table contains the set of significantly underrepresented GO terms among all sample libraries
717 (Benjamini-Hochberg FDR < 0.05).

     Figure 2 – Supplementary Table 3

Pop 1	Pop 2	Pop 3	D_{tree} (%)	f_4-ratio (%)	p-value	Z-score
S	N3	W	3.9	8.6	0	13.0*
S	N1	W	3.7	8.2	0	11.4*
S	N3	L	2.8	6.3	0	11.4*
S	N1	L	2.2	4.9	1.18E-16	8.3*
S	B	N1	4.2	9.0	1.56E-15	8.0*
S	B	W	2.2	4.7	1.26E-14	7.7*
B	N3	W	1.8	4.1	2.44E-14	7.6*
L	W	N1	2.2	4.5	2.73E-13	7.3*
L	W	B	1.7	3.9	2.76E-13	7.3*
B	N1	W	1.7	3.7	2.11E-12	7.0*
S	B	N3	3.6	8.3	2.31E-11	6.7*
L	W	N3	1.8	3.8	4.65E-09	5.9*
B	N3	L	1.8	4.0	6.52E-07	5.0*
S	B	L	1.1	2.4	1.01E-04	3.9*
B	N1	L	1.2	2.6	3.59E-04	3.6*
N1	N3	L	0.7	1.5	1.13E-03	3.3*
L	W	S	0.5	1.4	2.86E-03	3.0
N1	N3	S	0.4	0.9	5.72E-02	1.9
N1	N3	W	0.2	0.5	2.79E-01	1.1
N3	N1	B	0.3	0.7	2.86E-01	1.1

720
721
722   **Figure 2 – Supplementary Table 3.** Table of D_{tree} scores, f_4-admixture ratios, and Z-scores for
723   each of the 20 trios contained within the Lake Kronotskoe species flock. 16 trios were found to
724   have a significant, though minimal, contribution of introgressed alleles (asterisks)(Holm-
725   Bonferoni, FWER < 0.01). B, Bigmouth; L, Longhead; N1, Nosed 1; N3, Nosed 3; S, Smallmouth;
726   W, White.
727

Figure 3

729
730

Figure 3. Coding and non-coding conserved elements differentiating anadromous Dolly Varden charr from the Lake Kronotskoe species flock. A) Table showing the top 20 genes differentiating riverine Dolly Varden from Lake Kronotskoe inhabitants along with distribution of Ft values of genes. B) Table of the top 20 differentiating CNEs and the distribution of Fst values. C) Overview of shared and unique GO terms associated with genes and CNEs with Fst > 0.5. Venn diagram values represent GO terms which occur six or more times. Table summarizes broad categories of most frequent GO term associations. The distributions and the numbers of elements represent all non-zero Fst values; *orange*, genes with known roles in modulating craniofacial morphology (C); *blue*, genes with known roles in thyroid function (T). Asterisk denotes gene classified as (C), (T), or both.

741   **Figure 4**

742
743
Figure 4. Differentiation of *slc26a10* in pairwise comparisons between anadromous Dolly
Varden charr and the Lake Kronotskoe species flock. **(A)** The *slc26a10* locus shows high
differentiation (Fst). The gene locus contains one highly differentiating non-coding and one
coding variant. Dolly Varden *slc26a10* is homologous to human pendrin (*SLC26A4*), a known
thyroid regulator. The broader locus has low nucleotide diversity as illustrated by sliding
window plots of Tajima's Pi. Horizontal dotted gray line represents Fst = 0.9. Plots are included
for 10kb sliding windows along LG17, the coding elements within the broader locus, the
number of variants per sliding window, Fst, and Tajima's Pi per sliding window in pairwise
comparisons between riverine Dolly Varden charr and the Lake Kronotskoe species flock. **(B)**
Slc26a10 contains a fixed amino acid substitution in a conserved proline that differentiates
Dolly Varden charr from each of the major clades of the lacustrine morphs.

Figure 5

Figure 5. Fixation of variation in *sf3b4* between anadromous Dolly Varden charr and the Lake Kronotskoe species flock. **(A)** Three highly differentiating synonymous variants lie within *sf3b4*. Fst is plotted in non-overlapping 10kb sliding windows along the length of the chromosome LG30. *Sf3b4* is contained within a 200kb interval of low nucleotide diversity as illustrated by sliding window plots of Tajima's Pi. Horizontal dotted gray line represents Fst = 0.9. The broad locus encompassing *sf3b4* is shown in detail including plots of the number of variants, Fst, and Tajima's Pi in 10kb non-overlapping sliding windows. **(B)** 3 synonymous variants in *sf3b4* are fixed in lacustrine Dolly Varden charr; *dark orange,* non-reference allele fixed*; light orange,* non-reference allele predominant (alt. allele freq. > 50%) in sequence pool; *Light blue*, a lineage for which the reference allele is predominant (ref. allele freq. > 50%) at the locus.

769 **Figure 6**

Figure 6. Thyroid hormone (T3) levels and associating highly differentiating candidates in lacustrine Dolly Varden morphs related to thyroid function and craniofacial morphology. **(A)** Serum T3 levels were significantly less in Nosed 1, Nosed 3, and Smallmouth lineages relative to anadromous Dolly Varden; mean +/- 1 SD, with boundary values indicated (min/max). **(B)** Top 20 candidate regions differentiating Nosed lineages from Longhead and White lineages. Sites uniquely differentiating Longhead morphs from Nosed 1 and Nosed 3 morphs, and White morphs from Nosed 1 and Nosed 3 morphs identify loci associated with thyroid function (blue, T) or craniofacial developmental (orange, C) modulating elements. **(C)** Schematic of Leptin A paralogs in salmonids detailing previously unknown gene *leptin a2* having a unique 66 amino acid N-terminus extension compared to its paralog. Nosed lineages contain a fixed, non-synonymous SNP in this conserved N-terminal sequence. **(D)** An *otx2b* CNE contains fixed or

783   highly differentiating variants within a predicted Pax1 binding domain that associates with lake
784   morphs exhibiting significantly different thyroid signaling activities; percentage of reference
785   and alternate allele reads indicated per morph. Low level detection of variant or reference
786   alleles noted (1 read for reference allele (asterisk), 2 reads for non-reference allele (dagger)).
787

Figure 7

Figure 7. Effect of thyroid follicle ablation upon morphology and gene expression. (A) In thyroid ablated adult zebrafish (hypothyroid) there is a significant shift in the position of the maxilla towards a more terminal position. **(B)** In hypothyroid individuals, there is a significant reduction in the distance from the dorsal-most position of the maxilla to the long-axis of the body. All distances are relative to the standard length. n = 11 for each condition. **(C)** In hypothyroid individuals, there is a significant increase of lepa expression in the head at 14dpf. n = 3 pools of 20 heads for each condition. **(D)** Plotting Principal Components 1 and 2 reveals significant differences in dentary shape (Proscrustes ANOVA $F_{40;944}$=40.84 p<0.0001) between morphs. Scale bar = 10mm.

801   **Figure 7 – Supplementary Figure 1**

802
803
804   **Figure 7 – Supplementary Figure 1. Landmarks used to geomorphic morphometric analyses.**
805   Landmarks were assigned to each of four bones. The dentary (red), the parasphenoid (blue),
806   the hyomandibula (green), and the anguloarticulare (yellow).
807
808

809   **Figure 7 – Supplementary Figure 2 Figure 7 – Supplementary Figure 2**

810
811
812   **Figure 7 – Supplementary Figure 2. Geometric morphometric analyses of anguloarticulare**
813   **among Lake Kronotskoe morphs.** Procrustes analyses identified highly significant differences in
814   shape (Procrustes ANOVA $F_{80;1760}$=18.88 p<0.0001). Scale bar = 10mm.
815
816

817　**Figure 7 – Supplementary Figure 3**

818
819
820　**Figure 7 – Supplementary Figure 3. Geometric morphometric analyses of hyomandibula**
821　**among Lake Kronotskoe morphs.** Procrustes analyses identified significant differences in shape
822　(Procrustes ANOVA $F_{60;1212}$=6.55 p<0.001) with Smallmouth morphs possessing the most
823　distinct shape.
824
825

826   **Figure 7 – Supplementary Figure 4**

827
828
829   **Figure 7 – Supplementary Figure 4. Geometric morphometric analyses of parasphenoid**
830   **among Lake Kronotskoe morphs.** Procrustes analyses identified no significant differences in
831   shape. Scale bar = 10mm.

References

Bando, H., P. Gergics, B. L. Bohnsack, K. P. Toolan, C. E. Richter, J. A. Shavit and S. A. Camper (2020). "Otx2b mutant zebrafish have pituitary, eye and mandible defects that model mammalian disease." Human Molecular Genetics **29**(10): 1648-1657.

Barluenga, M., K. N. Stölting, W. Salzburger, M. Muschick and A. Meyer (2006). "Sympatric speciation in Nicaraguan crater lake cichlid fish." Nature **439**(7077): 719-723.

Bernier, F. P., O. Caluseriu, S. Ng, J. Schwartzentruber, K. J. Buckingham, A. M. Innes, E. W. Jabs, J. W. Innis, J. L. Schuette, J. L. Gorski, P. H. Byers, G. Andelfinger, V. Siu, J. Lauzon, B. A. Fernandez, M. McMillin, R. H. Scott, H. Racher, F. C. Consortium, J. Majewski, D. A. Nickerson, J. Shendure, M. J. Bamshad and J. S. Parboosingh (2012). "Haploinsufficiency of SF3B4, a component of the pre-mRNA spliceosomal complex, causes Nager syndrome." American journal of human genetics **90**(5): 925-933.

Berthelot, C., F. Brunet, D. Chalopin, A. Juanchich, M. Bernard, B. Noël, P. Bento, C. Da Silva, K. Labadie, A. Alberti, J.-M. Aury, A. Louis, P. Dehais, P. Bardou, J. Montfort, C. Klopp, C. Cabau, C. Gaspin, G. H. Thorgaard, M. Boussaha, E. Quillet, R. Guyomard, D. Galiana, J. Bobe, J.-N. Volff, C. Genêt, P. Wincker, O. Jaillon, H. R. Crollius and Y. Guiguen (2014). "The rainbow trout genome provides novel insights into evolution after whole-genome duplication in vertebrates." Nature Communications **5**(1): 3657.

Blythe, J., S. Nguyen and S. McMenamin (2022). Thyroid hormone mediates juvenile and adult craniofacial shape change in zebrafish. FaceBase Consortium.

Bolger, A. M., M. Lohse and B. Usadel (2014). "Trimmomatic: a flexible trimmer for Illumina sequence data." Bioinformatics **30**(15): 2114-2120.

Braitseva, O. A., I. V. Melekestsev, V. V. Ponomareva and L. D. Sulerzhitsky (1995). "Ages of calderas, large explosive craters and active volcanoes in the Kuril-Kamchatka region, Russia." Bulletin of Volcanology **57**(6): 383-402.

Brylka, L. J., A.-R. Alimy, M. E. A. Tschaffon-Müller, S. Jiang, T. M. Ballhause, A. Baranowsky, S. von Kroge, J. Delsmann, E. Pawlus, K. Eghbalian, K. Püschel, A. Schoppa, M. Haffner-Luntzer, D. J. Beech, F. T. Beil, M. Amling, J. Keller, A. Ignatius, T. A. Yorgan, T. Rolvien and T. Schinke (2024). "Piezo1 expression in chondrocytes controls endochondral ossification and osteoarthritis development." Bone Research **12**(1): 12.

Castinetti, F., M. L. Brinkmeier, D. F. Gordon, K. R. Vella, J. M. Kerr, A. H. Mortensen, A. Hollenberg, T. Brue, E. C. Ridgway and S. A. Camper (2011). "PITX2 and PITX1 Regulate Thyrotroph Function and Response to Hypothyroidism." Molecular Endocrinology **25**(11): 1950-1960.

Choi, Y. and A. P. Chan (2015). "PROVEAN web server: a tool to predict the functional effect of amino acid substitutions and indels." Bioinformatics **31**(16): 2745-2747.

Copeland, D. L., R. J. Duff, Q. Liu, J. Prokop and R. L. Londraville (2011). "Leptin in teleost fishes: an argument for comparative study." Frontiers in physiology **2**: 26-26.

Daane, J. M., N. Blum, J. Lanni, H. Boldt, M. K. Iovine, C. W. Higdon, S. L. Johnson, N. R. Lovejoy and M. P. Harris (2021). "Modulation of bioelectric cues in the evolution of flying fishes." Curr Biol **31**(22): 5052-5061 e5058.

Daane, J. M., A. Dornburg, P. Smits, D. J. MacGuigan, M. Brent Hawkins, T. J. Near, H. William
Detrich Iii and M. P. Harris (2019). "Historical contingency shapes adaptive radiation in Antarctic
fishes." Nature Ecology & Evolution **3**(7): 1102-1109.
Daane, J. M., N. Rohner, P. Konstantinidis, S. Djuranovic and M. P. Harris (2015). "Parallelism
and Epistasis in Skeletal Evolution Identified through Use of Phylogenomic Mapping Strategies."
Molecular Biology and Evolution **33**(1): 162-173.
Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham, T. Keane,
S. A. McCarthy, R. M. Davies and H. Li (2021). "Twelve years of SAMtools and BCFtools."
GigaScience **10**(2): giab008.
Davidson, W. S., B. F. Koop, S. J. M. Jones, P. Iturra, R. Vidal, A. Maass, I. Jonassen, S. Lien and S.
W. Omholt (2010). "Sequencing the genome of the Atlantic salmon (Salmo salar)." Genome
Biology **11**(9): 403.
de Graaf, M., H. J. Megens, J. Samallo and F. Sibbing (2010). "Preliminary insight into the age
and origin of the Labeobarbus fish species flock from Lake Tana (Ethiopia) using the mtDNA
cytochrome b gene." Mol Phylogenet Evol **54**(2): 336-343.
De-Kayne, R., O. M. Selz, D. A. Marques, D. Frei, O. Seehausen and P. G. D. Feulner (2022).
"Genomic architecture of adaptive radiation and hybridization in Alpine whitefish." Nature
Communications **13**(1): 4479.
De-Kayne, R., O. M. Selz, D. A. Marques, D. Frei, O. Seehausen and P. G. D. Feulner (2022).
"Hybridization and a mixture of small and large-effect loci facilitate adaptive radiation." bioRxiv:
2022.2002.2018.481029.
Decherf, S., I. Seugnet, S. Kouidhi, A. Lopez-Juarez, M. S. Clerget-Froidevaux and B. A. Demeneix
(2010). "Thyroid hormone exerts negative feedback on hypothalamic type 4 melanocortin
receptor expression." Proc Natl Acad Sci U S A **107**(9): 4471-4476.
Diaczok, D., C. Romero, J. Zunich, I. Marshall and S. Radovick (2008). "A Novel Dominant
Negative Mutation of OTX2 Associated with Combined Pituitary Hormone Deficiency." The
Journal of Clinical Endocrinology & Metabolism **93**(11): 4351-4359.
Dicks, A. R., G. I. Maksaev, Z. Harissa, A. Savadipour, R. Tang, N. Steward, W. Liedtke, C. G.
Nichols, C.-L. Wu and F. Guilak (2023). "Skeletal dysplasia-causing TRPV4 mutations suppress
the hypertrophic differentiation of human iPSC-derived chondrocytes." eLife **12**: e71154.
Dimitrieva, S. and P. Bucher (2013). "UCNEbase--a database of ultraconserved non-coding
elements and genomic regulatory blocks." Nucleic acids research **41**(Database issue): D101-
D109.
Esin, E. V., E. S. Bocharova, E. A. Borisova and G. N. Markevich (2020). "Interaction among
morphological, trophic and genetic groups in the rapidly radiating Salvelinus fishes from Lake
Kronotskoe." Evolutionary Ecology **34**(4): 611-632.
Esin, E. V., G. N. Markevich, N. O. Melnik, D. V. Kapitanova and F. N. Shkil (2021). "Natural toxic
impact and thyroid signalling interplay orchestrates riverine adaptive divergence of salmonid
fish." J Anim Ecol **90**(4): 1004-1019.
Esin, E. V., G. N. Markevich, N. O. Melnik, D. V. Zlenko and F. N. Shkil (2021). "Ambient
temperature as a factor contributing to the developmental divergence in sympatric salmonids."
PLoS One **16**(10): e0258536.

Esin, E. V., G. N. Markevich and M. Y. Pichugin (2018). "Juvenile divergence in adaptive traits
among seven sympatric fish ecomorphs arises before moving to different lacustrine habitats."
Journal of Evolutionary Biology **31**(7): 1018-1034.
Esin, E. V., G. N. Markevich, D. V. Zlenko and F. N. Shkil (2021). "Thyroid-Mediated Metabolic
Differences Underlie Ecological Specialization of Extremophile Salmonids in the Arctic Lake
El'gygytgyn." Frontiers in Ecology and Evolution **9**.
Esin, E. V., F. N. Shkil, D. V. Zlenko, D. A. Medvedev, N. B. Korostelev and G. N. Markevich
(2024). "Quaternary history and radiation of Salvelinus fish in the ancient Arctic Lake
El'gygytgyn." Hydrobiologia **851**(10): 2389-2404.
Evans, A. L. and P. J. Gage (2005). "Expression of the homeobox gene Pitx2 in neural crest is
required for optic stalk and ocular anterior segment development." Human Molecular Genetics
14(22): 3347-3359.
Fabik, J., K. Kovacova, Z. Kozmik and O. Machon (2020). "Neural crest cells require Meis2 for
patterning the mandibular arch via the Sonic hedgehog pathway." Biology Open **9**(6).
Friedrichs, B., C. Tepel, T. Reinheckel, J. Deussing, K. von Figura, V. Herzog, C. Peters, P. Saftig
and K. Brix (2003). "Thyroid functions of mouse cathepsins B, K, and L." The Journal of Clinical
Investigation **111**(11): 1733-1745.
Guðbjörg Ósk, J., E. Laura-Marie von, I. Finnur, T. Samuel, S. Sigurdur Sveinn, P. Arnar and S.
Sarah Elizabeth (2024). "Diversity in the internal functional feeding elements of sympatric
morphs of Arctic charr (Salvelinus alpinus)." bioRxiv: 2023.2002.2017.528955.
Hartl, D. L., A. G. Clark and A. G. Clark (1997). Principles of population genetics, Sinauer
associates Sunderland, MA.
Herrero, J., M. Muffato, K. Beal, S. Fitzgerald, L. Gordon, M. Pignatelli, A. J. Vilella, S. M. J.
Searle, R. Amode, S. Brent, W. Spooner, E. Kulesha, A. Yates and P. Flicek (2016). "Ensembl
comparative genomics resources." Database : the journal of biological databases and curation
2016: bav096.
Hoang, D. T., O. Chernomor, A. von Haeseler, B. Q. Minh and L. S. Vinh (2017). "UFBoot2:
Improving the Ultrafast Bootstrap Approximation." Molecular Biology and Evolution **35**(2): 518-
522.
Jacobs, A., M. Carruthers, A. Yurchenko, N. V. Gordeeva, S. S. Alekseyev, O. Hooker, J. S. Leong,
D. R. Minkley, E. B. Rondeau, B. F. Koop, C. E. Adams and K. R. Elmer (2020). "Parallelism in eco-
morphology and gene expression despite variable evolutionary and genomic backgrounds in a
Holarctic fish." PLoS Genet **16**(4): e1008658.
Jonsson, B. and N. Jonsson (2001). "Polymorphism and speciation in Arctic charr." Journal of
Fish Biology **58**(3): 605-638.
Kalyaanamoorthy, S., B. Q. Minh, T. K. F. Wong, A. von Haeseler and L. S. Jermiin (2017).
"ModelFinder: fast model selection for accurate phylogenetic estimates." Nature Methods
14(6): 587-589.
Keer, S., K. Cohen, C. May, Y. Hu, S. McMenamin and L. P. Hernandez (2019). "Anatomical
Assessment of the Adult Skeleton of Zebrafish Reared Under Different Thyroid Hormone
Profiles." The Anatomical Record **302**(10): 1754-1769.
Keer, S., J. D. Storch, S. Nguyen, M. Prado, R. Singh, L. P. Hernandez and S. K. McMenamin
(2022). "Thyroid hormone shapes craniofacial bones during postembryonic zebrafish
development." Evolution & Development **24**(1-2): 61-76.

Kitano, J., S. C. Lema, J. A. Luckenbach, S. Mori, Y. Kawagishi, M. Kusakabe, P. Swanson and C. L. Peichel (2010). "Adaptive Divergence in the Thyroid Hormone Signaling Pathway in the Stickleback Radiation." Current Biology **20**(23): 2124-2130.

Klemetsen, A. (2010). "The Charr Problem Revisited: Exceptional Phenotypic Plasticity Promotes Ecological Speciation in Postglacial Lakes." Freshwater Reviews **3**(1): 49-74.

Kofler, R., P. Orozco-terWengel, N. De Maio, R. V. Pandey, V. Nolte, A. Futschik, C. Kosiol and C. Schlötterer (2011). "PoPoolation: A Toolbox for Population Genetic Analysis of Next Generation Sequencing Data from Pooled Individuals." PLOS ONE **6**(1): e15925.

Kofler, R., R. V. Pandey and C. Schlötterer (2011). "PoPoolation2: identifying differentiation between populations using sequencing of pooled DNA samples (Pool-Seq)." Bioinformatics (Oxford, England) **27**(24): 3435-3436.

Kozomara, A. and S. Griffiths-Jones (2010). "miRBase: integrating microRNA annotation and deep-sequencing data." Nucleic Acids Research **39**(suppl_1): D152-D157.

Lecaudey, L. A., U. K. Schliewen, A. G. Osinov, E. B. Taylor, L. Bernatchez and S. J. Weiss (2018). "Inferring phylogenetic structure, hybridization and divergence times within Salmoninae (Teleostei: Salmonidae) using RAD-sequencing." Molecular Phylogenetics and Evolution **124**: 82-99.

Levin, B. A., E. Simonov, Y. Y. Dgebuadze, M. Levina and A. S. Golubtsov (2020). "In the rivers: Multiple adaptive radiations of cyprinid fishes (Labeobarbus) in Ethiopian Highlands." Scientific Reports **10**(1): 7192.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin and G. P. D. P. Subgroup (2009). "The Sequence Alignment/Map format and SAMtools." Bioinformatics **25**(16): 2078-2079.

Liu, X., H. Wang, G. Li, H.-Z. Huang and Y.-Q. Wang (2013). "The function of *DrPax1b* gene in the embryonic development of zebrafish." Genes & Genetic Systems **88**(4): 261-269.

Liu, Y.-H., T.-C. Lin and S.-P. L. Hwang (2020). "Zebrafish Pax1a and Pax1b are required for pharyngeal pouch morphogenesis and ceratobranchial cartilage development." Mechanisms of Development **161**: 103598.

Livak, K. J. and T. D. Schmittgen (2001). "Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method." Methods **25**(4): 402-408.

Ma, Z., P. Zhu, M. Pang, L. Guo, N. Chang, J. Zheng, X. Zhu, C. Gao, H. Huang, Z. Cui, J. W. Xiong, J. Peng and J. Chen (2017). "A novel inducible mutagenesis screen enables to isolate and clone both embryonic and adult zebrafish mutants." Sci Rep **7**(1): 10381.

Maitland, P. S., I. J. Winfield, I. D. McCarthy and F. Igoe (2007). "The status of Arctic charr Salvelinus alpinus in Britain and Ireland." Ecology of Freshwater Fish **16**(1): 6-19.

Malinsky, M., R. J. Challis, A. M. Tyers, S. Schiffels, Y. Terai, B. P. Ngatunga, E. A. Miska, R. Durbin, M. J. Genner and G. F. Turner (2015). "Genomic islands of speciation separate cichlid ecomorphs in an East African crater lake." Science **350**(6267): 1493-1498.

Malinsky, M., M. Matschiner and H. Svardal (2021). "Dsuite - Fast D-statistics and related admixture evidence from VCF files." Mol Ecol Resour **21**(2): 584-595.

Malinsky, M., H. Svardal, A. M. Tyers, E. A. Miska, M. J. Genner, G. F. Turner and R. Durbin (2018). "Whole-genome sequences of Malawi cichlids reveal multiple radiations interconnected by gene flow." Nat Ecol Evol **2**(12): 1940-1955.

Markevich, G., E. Esin and L. Anisimova (2018). "Basic description and some notes on the
evolution of seven sympatric morphs of Dolly Varden Salvelinus malma from the Lake
Kronotskoe Basin." <u>Ecol Evol</u> **8**(5): 2554-2567.
Markevich, G. N., D. V. Zlenko, F. N. Shkil, U. K. Schliewen, L. A. Anisimova, A. A. Sharapkova and
E. V. Esin (2021). "Natural Barriers and Internal Sources for the Reproductive Isolation in
Sympatric Salmonids from the Lake–River System." <u>Evolutionary Biology</u> **48**(4): 407-421.
Mason, V. C., G. Li, K. M. Helgen and W. J. Murphy (2011). "Efficient cross-species capture
hybridization and next-generation sequencing of mitochondrial genomes from noninvasively
sampled museum specimens." <u>Genome research</u> **21**(10): 1695-1704.
McCormick, S. D., A. M. Regish, W. R. Ardren, B. T. Bjornsson and N. J. Bernier (2019). "The
evolutionary consequences for seawater performance and its hormonal control when
anadromous Atlantic salmon become landlocked." <u>Sci Rep</u> **9**(1): 968.
McCormick, S. D., A. M. Regish, W. R. Ardren, B. T. Björnsson and N. J. Bernier (2019). "The
evolutionary consequences for seawater performance and its hormonal control when
anadromous Atlantic salmon become landlocked." <u>Scientific Reports</u> **9**(1): 968.
McLean, C. Y., D. Bristor, M. Hiller, S. L. Clarke, B. T. Schaar, C. B. Lowe, A. M. Wenger and G.
Bejerano (2010). "GREAT improves functional interpretation of cis-regulatory regions." <u>Nature</u>
<u>biotechnology</u> **28**(5): 495-501.
McMenamin, S. K., E. J. Bain, A. E. McCann, L. B. Patterson, D. S. Eom, Z. P. Waller, J. C. Hamill, J.
A. Kuhlman, J. S. Eisen and D. M. Parichy (2014). "Thyroid hormone–dependent adult pigment
cell lineage and pattern in zebrafish." <u>Science</u> **345**(6202): 1358-1361.
Myers, G. S. (1960). "The Endemic Fish Fauna of Lake Lanao, and the Evolution of Higher
Taxonomic Categories." <u>Evolution</u> **14**(3): 323-333.
Nagelkerke, L. and F. A. Sibbing (2000). "The large barbs (Barbus SPP., cyprinidae, teleostei) of
Lake Tana (Ethiopia), with a description of a new species, Barbus osseensis." <u>Netherlands</u>
<u>Journal of Zoology 50 (2000) 2</u> **50**.
Nagelkerke, L. A., K. M. Leon-Kloosterziel, H. J. Megens, M. De Graaf, O. E. Diekmann and F. A.
Sibbing (2015). "Shallow genetic divergence and species delineations in the endemic
Labeobarbus species flock of Lake Tana, Ethiopia." <u>J Fish Biol</u> **87**(5): 1191-1208.
Ng, P. C. and S. Henikoff (2003). "SIFT: Predicting amino acid changes that affect protein
function." <u>Nucleic Acids Res</u> **31**(13): 3812-3814.
Nguyen, L.-T., H. A. Schmidt, A. von Haeseler and B. Q. Minh (2015). "IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies." <u>Molecular</u>
<u>Biology and Evolution</u> **32**(1): 268-274.
Nguyen, S. V., D. Lanni, Y. Xu, J. S. Michaelson and S. K. McMenamin (2022). "Dynamics of the
Zebrafish Skeleton in Three Dimensions During Juvenile and Adult Development." <u>Frontiers in</u>
<u>Physiology</u> **13**.
Nordeng, H. (1983). "Solution to the "Char Problem" based on Arctic Char (Salvelinus alpinus) in
Norway." <u>Canadian Journal of Fisheries and Aquatic Sciences</u> **40**(9): 1372-1387.
Osinov, A. G., A. A. Volkov and N. S. Mugue (2021). "Charrs of the genus Salvelinus
(Salmonidae): hybridization, phylogeny and evolution." <u>Hydrobiologia</u> **848**(3): 705-726.
Petit, F., F. Escande, A. S. Jourdain, N. Porchet, J. Amiel, B. Doray, M. A. Delrue, E. Flori, C. A.
Kim, S. Marlin, S. P. Robertson, S. Manouvrier-Hanu and M. Holder-Espinasse (2014). "Nager

syndrome: confirmation of SF3B4 haploinsufficiency as the major cause." Clinical Genetics
86(3): 246-251.
Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira, D. Bender, J. Maller, P. Sklar, P. I.
de Bakker, M. J. Daly and P. C. Sham (2007). "PLINK: a tool set for whole-genome association
and population-based linkage analyses." Am J Hum Genet **81**(3): 559-575.
Quinlan, A. R. and I. M. Hall (2010). "BEDTools: a flexible suite of utilities for comparing genomic
features." Bioinformatics **26**(6): 841-842.
Quinlan, A. R. and I. M. Hall (2010). "BEDTools: a flexible suite of utilities for comparing genomic
features." Bioinformatics **26**(6): 841-842.
Richard, S., R. Guyot, M. Rey-Millet, M. Prieux, S. Markossian, D. Aubert and F. Flamant (2020).
"A Pivotal Genetic Program Controlled by Thyroid Hormone during the Maturation of
GABAergic Neurons." iScience **23**(3): 100899-100899.
Rohlf, F. J. (2015). "The tps series of software." Hystrix, the Italian Journal of Mammalogy **26**(1):
9-12.
Rondeau, E. B., D. R. Minkley, J. S. Leong, A. M. Messmer, J. R. Jantzen, K. R. von Schalburg, C.
Lemon, N. H. Bird and B. F. Koop (2014). "The Genome and Linkage Map of the Northern Pike
(Esox lucius): Conserved Synteny Revealed between the Salmonid Sister Group and the
Neoteleostei." PLOS ONE **9**(7): e102089.
Rothstein, M. and M. Simoes-Costa (2020). "Heterodimerization of TFAP2 pioneer factors drives
epigenomic remodeling during neural crest specification." Genome Research **30**(1): 35-48.
Saltykova, E., G. Markevich and K. Kuzishchin (2015). "Divergent skull morphology between
trophic separated lacustrine forms of Dolly Varden charr from Lake Kronotskoe, Kamchatka,
Russia." Environmental Biology of Fishes **98**(2): 559-570.
Sandlund, O. T., K. Gunnarsson, P. M. Jónasson, B. Jonsson, T. Lindem, K. P. Magnússon, H. J.
Malmquist, H. Sigurjónsdóttir, S. Skúlason and S. S. Snorrason (1992). "The Arctic Charr
Salvelinus alpinus in Thingvallavatn." Oikos **64**(1/2): 305-351.
Schluter, D. and G. L. Conte (2009). "Genetics and ecological speciation." Proceedings of the
National Academy of Sciences **106**(supplement_1): 9955-9962.
Sedlazeck, F. J., P. Rescheneder and A. von Haeseler (2013). "NextGenMap: fast and accurate
read mapping in highly polymorphic genomes." Bioinformatics **29**(21): 2790-2791.
Shkil, F. N., D. V. Kapitanova, V. B. Borisov, B. Abdissa and S. V. Smirnov (2012). "Thyroid
hormone in skeletal development of cyprinids: effects and morphological consequences."
Journal of Applied Ichthyology **28**(3): 398-405.
Shkil, F. N., O. E. Lazebnyi, D. V. Kapitanova, B. Abdissa, V. B. Borisov and S. V. Smirnov (2015).
"Ontogenetic mechanisms of explosive morphological divergence in the Lake Tana (Ethiopia)
species flock of large African barbs (Labeobarbus; Cyprinidae; Teleostei)." Russian Journal of
Developmental Biology **46**(5): 294-306.
Simonsen, M. K., A. Siwertsson, C. E. Adams, P.-A. Amundsen, K. Præbel and R. Knudsen (2017).
"Allometric trajectories of body and head morphology in three sympatric Arctic charr
(Salvelinus alpinus (L.)) morphs." Ecology and Evolution **7**(18): 7277-7289.
Taylor, E. B. (2016). "The Arctic char (Salvelinus alpinus) "complex" in North America revisited."
Hydrobiologia **783**(1): 283-293.

Tingaud-Sequeira, A., A. Trimouille, S. Marlin, E. Lopez, M. Berenguer, S. Gherbi, B. Arveiler, D.
Lacombe and C. Rooryck (2020). "Functional and genetic analyses of ZYG11B provide evidences
for its involvement in OAVS." Molecular Genetics & Genomic Medicine **8**(10): e1375.
Tipsmark, C. K., K. J. Sørensen and S. S. Madsen (2010). "Aquaporin expression dynamics in
osmoregulatory tissues of Atlantic salmon during smoltification and seawater acclimation."
Journal of Experimental Biology **213**(3): 368-379.
Treaster, S., J. Deelen, J. M. Daane, J. Murabito, D. Karasik and M. P. Harris (2022). Nature
Portfolio.
Viktorovskii, R. M. (1978). Mekhanizmȳ vidoobrazovaniya u gol'tsov Kronotskogo Ozera, Изд-во
Наука.
Walker, A. F., R. B. Greer and A. S. Gardner (1988). "Two ecologically distinct forms of arctic
charr Salvelinus alpinus (L.) in Loch Rannoch, Scotland." Biological Conservation **43**(1): 43-61.
Wong, T. C. B., M. Rebbert, C. Wang, X. Chen, A. Heffer, V. E. Zarelli, I. B. Dawid and H. Zhao
(2016). "Genes regulated by potassium channel tetramerization domain containing 15 (Kctd15)
in the developing neural crest." The International journal of developmental biology **60**(4-6):
159-166.
Yagasaki, Y., T. Yamaguchi, J. Watahiki, M. Konishi, H. Katoh and K. Maki (2003). "The role of
craniofacial growth in leptin deficient (ob/ob) mice." Orthodontics & Craniofacial Research **6**(4):
233-241.
Yoon, G. H., K. Kim, D. S. Park and S. C. Choi (2022). "RNF152 negatively regulates Wnt/β-
catenin signaling in Xenopus embryos." BMB Rep.
Zimmermann-Belsing, T., G. Brabant, J. Holst and U. Feldt-Rasmussen (2003). "Circulating leptin
and thyroid dysfunction." European Journal of Endocrinology Eur J Endocrinol **149**(4): 257-271.
1112