1   Reduced steroid activation of elephant shark GR and MR after inserting four amino acids from

2   the DNA-binding domain of lamprey corticoid receptor-1

4   Yoshinao Katsu[1,2,] *, Jiawen Zhang[2], Michael E. Baker[3,4,] *

5   [1] Faculty of Science

6   Hokkaido University

7   Sapporo, Japan

8   [2] Graduate School of Life Science

9   Hokkaido University

10   Sapporo, Japan

11   [3] Division of Nephrology-Hypertension

12   Department of Medicine, 0693

13   University of California, San Diego

14   9500 Gilman Drive

15   La Jolla, CA 92093-0693

16   Center for Academic Research and Training in Anthropogeny (CARTA) [4]

17   University of California, San Diego

18   La Jolla, CA 92093

*Correspondence to

Y. Katsu; E-mail: ykatsu@sci.hokudai.ac.jp

M. E. Baker; E-mail: mbaker@health.ucsd.edu

Abstract. Atlantic sea lamprey contains two corticoid receptors (CRs), CR1 and CR2, that have

identical amino acid sequences, except for a four amino acid insert (Thr-Arg-Gln-Gly) in the

CR1 DNA-binding domain (DBD). Steroids are stronger transcriptional activators of CR2 than

of CR1 suggesting that the insert reduces the transcriptional response of lamprey CR1 to steroids.

The DBD in elephant shark mineralocorticoid receptor (MR) and glucocorticoid receptor (GR),

which are descended from a CR, lack these four amino acids, suggesting that a CR2 is their

common ancestor. To determine if, similar to lamprey CR1, the presence of this insert in

elephant shark MR and GR decreases transcriptional activation by corticosteroids, we inserted

these four CR1-specific residues into the DBD of elephant shark MR and GR. Compared to

steroid activation of wild-type elephant shark MR and GR, cortisol, corticosterone, aldosterone,

11-deoxycorticosterone and 11-deoxycortisol had lower transcriptional activation of these

mutant MR and GR receptors, indicating that the absence of this four-residue segment in the

DBD in wild-type elephant shark MR and GR increases transcriptional activation by

corticosteroids.

Key words: DNA-binding domain, corticoid receptor, mineralocorticoid receptor, glucocorticoid

receptor, evolution, elephant shark, Atlantic sea lamprey

Running title: DNA-binding domain regulates steroid activation of elephant shark

glucocorticoid and mineralocorticoid receptors.

Introduction

The sea lamprey (*Petromyzon marinus*) belongs to an ancient group of jawless vertebrates known as cyclostomes, which are basal vertebrates that evolved about 550 million years ago (1–4). Sea lamprey contains a corticoid receptor (CR) (5–8), which belongs to the nuclear receptor family of transcription factors (9,10), which also contains the mineralocorticoid receptor, glucocorticoid receptor, progesterone receptor, estrogen receptor and androgen receptor (9–13). A CR in an ancestral cyclostome is the common ancestor to the mineralocorticoid receptor (MR) and the glucocorticoid receptor (GR) in vertebrates (5,7). The MR and GR first appear as separate steroid receptors in cartilaginous fish (5–7,14,15). These two closely related steroid receptors regulate important physiological responses in vertebrates. The MR regulates electrolyte transport in the kidney and colon (12,16–21), as well as regulating gene transcription in a variety of non-epithelial tissues (22–26). The GR has diverse physiological actions in vertebrates including in development, metabolism, the stress response, inflammation and glucose homeostasis (27–31).

The importance of the MR and GR in vertebrate physiology stimulated our interest in understanding the evolution of the MR and GR from the CR (7,8,32). Unfortunately, as a result of complexities in the sequencing and assembly of the lamprey's highly repetitive and GC rich genome (3,33,34) until recently, only a partial CR sequence was available in GenBank (5). Fortunately, the recent sequencing of the lamprey germline genome (35) provided contiguous DNA encoding the complete sequences two CR isoforms, CR1 and CR2, which differ only in a

novel four amino acid insertion, Thr, Arg, Gln, Gly, (TRQG) in the DNA-binding domain

(DBD) (8) (Figure 1).

With the two CR sequences in hand, we characterized the response of HEK293 cells

transfected with each CR isoform to corticosteroids (8) and found that the loss of these four

amino acids in the DBD of CR2 increased fold transcriptional activation by about two-fold in the

presence of corticosteroids such as 11-deoxycorticosterone, 11-deoxycortisol, cortisol,

corticosterone and aldosterone.

```
lamprey CR1:       CLICSDEASGCHYGVLTCGSCKVFFKRAVEGTRQGQHNYLCAGRNDCIIDKIRRKNCPACRLRKCIQAGM
lamprey CR2:       CLICSDEASGCHYGVLTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLRKCIQAGM
elephant shark GR:CLVCSDEASGCHYGVLTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRFRKCLQAGM
elephant shark MR:CLVCSDEASGCHYGVLTCGSCKVFFKRAVEG----QQNYLCAGRNDCIIDKIRRKNCPACRLRKCLKAGM
human GR:          CLVCSDEASGCHYGVLTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRYRKCLQAGM
human MR:          CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
```

77
78

Figure 1. DNA-binding domains of lamprey CR1 and CR2 and elephant shark and human

MR and GR. The DNA-binding domain of lamprey CR1 has an insertion of four amino acids

that is absent in CR2 and in elephant shark and human MR and GR. Differences between the

sequence of lamprey CR and elephant shark and human MR and GR are shown in red.

Accessions are: XP_032811370 and XP_032811371 for lamprey CR1 and CR2, respectively,

XP_042195980 and XP_007902220 for elephant shark GR and MR, respectively, NP_000167

and AAA59571 for human GR and MR respectively.

Elephant shark mineralocorticoid receptor (MR) and glucocorticoid receptor (GR), which

are descended from the CR, lack these four amino acids in their DBD, suggesting that they

evolved from a CR2 ancestor. To determine if, similar to lamprey CR2, the absence of this insert

in the DBD of elephant shark MR and GR has a functional consequence, we inserted these four

residues into their DBDs to convert them to a CR1-like sequence. Here we report that cortisol,

corticosterone, aldosterone, 11-deoxycorticosterone and 11-deoxycortisol have lower

transcriptional activation of these mutant elephant shark MR and GR receptors, indicating that

compared to mutant elephant shark MR and GR, the absence of this four-residue segment in the

DBD in both wild-type elephant shark MR and GR increases activation by corticosteroids.

Results

Steroid-dependent activation of mutated elephant shark MR and GR.

To gain a quantitative measure of steroid activation of mutated elephant shark MR and

GR, we determined the concentration dependence of transcriptional activation by corticosteroids

of FLAG-tagged wild-type and mutated elephant shark GR transfected into HEK293 cells with

either an MMTV-luciferase promoter (Figure 2A-E) (9,36,37) or a TAT3 luciferase promoter

(9,38,39) (Figure 2F-J). A parallel study was done with wild-type and mutated elephant shark

MR (Figure 3). Luciferase levels were used to calculate an EC50 value and fold-activation for

each steroid for FLAG-tagged wild-type and mutated elephant shark GR and MR (Table 1). An

EC50 for 11-deoxycortisol activation of FLAG-tagged wild-type GR and elephant shark GR-

DBD-TRQG was not determined because fold-activation by 11-deoxycortisol of FLAG-tagged

wild-type GR and elephant shark GR-DBD-TRQG did not saturate (Figure 2D, 2J).

The results shown in Figure 2 and summarized in Table 1 reveal that addition of the

TRQG sequence to the DBD in elephant shark GR reduces fold-activation of this receptor by

corticosteroids in HEK293 cells transfected with either the MMTV promoter or the TAT3

promotor.   Fold-activation varied depended on the steroid and whether the MMTV promoter or

TAT3 promoter was used in the assay. In assays with the MMTV promoter, fold-activation was

higher for FLAG-tagged wild-type GR compared to the GR-DBD-TRQG mutant for all tested

corticosteroids (Figure 2A-E). Fold-activation also was higher in the assays with the MMTV

promoter than with the TAT3 promoter. In assays with the TAT3 promoter, fold-activation was

higher for FLAG-tagged wild-type GR compared to the GR-DBD-TRQG mutant for 11-

deoxycorticosterone and slightly higher for corticosterone and aldosterone (Figure 2G,H,I).

Cortisol had similar fold-activation for FLAG-tagged wild-type GR and GR-DBD-TRQG in

assays with TAT3 (Figure 2F). There was little activation by 11-deoxycortisol (Figure 2I).

In assays with the MMTV promoter, cortisol, 11-deoxycorticosterone, corticosterone and

aldosterone have a right-shift in the EC50 for FLAG-tagged wild-type elephant shark GR

compared to GR-DBD- TRQG (Table 1, Figure 2A-2C, 2E). In the presence of the TAT3

promoter, 11-deoxycorticosterone has a small right shift in the EC50 for FLAG-tagged wild-type

GR compared to GR-DBD- TRQG (Table 1, Figure 2G).   In the presence of the TAT3 promoter,

cortisol (Figure 2F), corticosterone (Figure 2G) and aldosterone (Figure 2J) have similar EC50s

for FLAG-tagged wild-type elephant shark GR compared to GR-DBD-TRQG. Corticosteroids

also had lower transcriptional activation of FLAG-tagged elephant shark MR-DBD- TRQG

compared to FLAG-tagged wild-type elephant shark MR (Figure 3, Table 1).

FLAG-tagged elephant shark MR and elephant shark MR-DBD-TRQG also had higher

fold-activation with MMTV promoter compared to the TAT3 promoter, although it varied with

the corticosteroid. For example, fold-activation of DOC for elephant shark MR was 12.4 in the

presence of MMTV and 8.0 in the presence of TAT3. And fold-activation of aldosterone for

elephant shark MR was 12 in the presence of MMTV and 10.7 in the presence of TAT3.

We also studied progesterone activation of FLAG-tagged elephant shark MR with TRQG

in its DBD. Progesterone is an antagonist for human MR (40–42), and an agonist for fish MR

(14,41,43,44), Progesterone is a transcriptional activator of elephant shark MR, but not elephant

shark GR (14,45). We find that addition of the TRQG sequence to FLAG-tagged elephant shark

MR reduces fold-activation by progesterone in HEK293 cells transfected with either the MMTV

promoter or the TAT3 promotor (Figure 3).

Fig. 2. Concentration-dependent transcriptional activation by corticosteroids of FLAG-

tagged wild-type and mutated elephant shark GR. Plasmids for FLAG-tagged wild-type

elephant shark GR and mutated elephant shark GR (with a TRQG insert) were expressed in

HEK293 cells with either an MMTV-luciferase promoter or a TAT3-luciferase promoter (9,36–

38). Cells were treated with increasing concentrations of either aldosterone, cortisol,

corticosterone, 11-deoxycortisol, 11-deoxycorticosterone or vehicle alone (DMSO). Results are

expressed as means ± SEM, number (n) of wells for each point =3. Y-axis indicates fold-

activation compared to the activity of vector with vehicle (DMSO) alone as 1. Figure 2A-E.

FLAG-tagged elephant shark GRs with MMTV-luc. Figure 2F-J. FLAG-tagged elephant shark

GRs with TAT3-luc.

Elephant shark MR

Fig. 3. Concentration-dependent transcriptional activation by corticosteroids and

progesterone of FLAG-tagged wild-type and mutated elephant shark MR. Plasmids for

FLAG-tagged wild-type elephant shark MR and mutated elephant shark MR (with a TRQG

insert) were expressed in HEK293 cells with either an MMTV-luciferase promoter or a TAT3-

luciferase promoter. Cells were treated with increasing concentrations of either aldosterone,

cortisol, corticosterone, 11-deoxycortisol, 11-deoxycorticosterone, progesterone or vehicle alone

(DMSO). Results are expressed as means ± SEM, number (n) of wells for each point =3. Y-axis

indicates fold-activation compared to the activity of vector with vehicle (DMSO) alone as 1.

Figure 3A-F. FLAG-tagged elephant shark MRs with MMTV-luc. Figure 3G-L. FLAG-tagged

elephant shark MRs with TAT3-luc.

Table 1. Steroid Activation of FLAG-tagged Wild-Type and Mutated Elephant Shark MR and
GR in HEK293 Cells with either an MMTV Promoter or a TAT3 Promoter.
167

MMTV-luc			Cortisol	DOC	CORT	S	Aldo	Prog
Elephant Shark GR -wt	EC50 (nM)		25.2	13.0	6.0	**	14.1	-
	Fold-Activation (± SEM) *		420 (± 15.1)	228 (± 8.6)	572 (± 28.5)	69.5 (± 1.8)	473 (± 11.4)	-
Elephant Shark GR -DBD-TRQG	EC50 (nM)		13.9	8.7	3.8	**	7.0	-
	Fold-Activation (± SEM) *		191 (± 3.8)	123 (± 3.7)	254 (± 27.8)	31.3 (± 1.2)	175 (± 10.0)	-
Elephant Shark MR -wt	EC50 (nM)		0.43	0.03	0.57	0.08	0.05	0.4
	Fold-Activation (± SEM) *		32.1 (± 2.0)	19.0 (± 1.4)	18.8 (± 0.8)	19.0 (± 0.4)	20.3 (± 0.8)	8.5 (± 0.3)
Elephant Shark MR -DBD-TRQG	EC50 (nM)		0.5	0.03	0.94	0.13	0.06	0.34
	Fold-Activation (± SEM) *		14.9 (± 1.5)	9.7 (± 0.9)	11.9 (± 0.5)	9.8 (± 0.3)	9.7 (± 0.4)	5.0 (± 0.2)
TAT3-luc			Cortisol	DOC	CORT	S	Aldo	Prog
Elephant Shark GR -wt	EC50 (nM)		20.5	12.5	4.9	**	10.1	-
	Fold-Activation (± SEM) *		40 (± 0.4)	13.6 (± 0.3)	32.2 (± 2.0)	3.1 (± 0.2)	43 (± 0.5)	-
Elephant Shark GR -DBD-TRQG	EC50 (nM)		23.6	15.5	4.4	**	10.8	-
	Fold-Activation (± SEM) *		41 (± 1.6)	9.3 (± 0.2)	28.9 (± 1.3)	2.1 (± 0.1)	38 (± 0.6)	-
Elephant Shark MR -wt	EC50 (nM)		1.82	0.03	0.38	0.1	0.03	0.27
	Fold-Activation (± SEM) *		33.4 (± 1.3)	15.9 (± 0.4)	17.2 (± 0.4)	17.7 (± 0.7)	18.6 (± 1.1)	8.3 (± 0.4)
Elephant Shark MR	EC50 (nM)		7.0	0.04	0.97	0.27	0.06	0.43

-DBD-TRQG	Fold-Activation (± SEM) *	21.7 (± 1.5)	9.5 (± 0.3)	11.9 (± 1.2)	9.9 (± 0.2)	11.7 (± 0.4)	4.8 (± 0.2)

wt = wild type sequence, DBD-TRQG = TRQG motif insertion in DBD

DOC = 11-deoxycorticosterone, CORT = Corticosterone, S = 11-deoxycortisol, Aldo =

aldosterone, Prog = Progesterone

*Fold activation values were calculated based on the 1000 nM values.

**Curve did not saturate.

FLAG-tagged GR and MR protein expression

To confirm the protein expression levels of FLAG-tagged GR and MR cultured cells

transfected with each construct were collected and treated with SDS sample buffer. And then, 20

μg of protein was subjected to SDS-PAGE on a 10% polyacrylamide gel and transferred to an

Immobilon membrane. The gels were also Coomassie blue, CBB stained. Subsequently, FLAG-

tagged proteins were detected using an antibody against the FLAG tag. We detected a single

band the MR and multiple bands the GR (Figure 4). Both MR and GR showed no marked

difference in protein levels between the wild type and the mutant containing 4 amino acid

insertion in the DBD. The multiple bands in GR may be due to protein phosphorylation.

Figure 4. **Expression of FLAG-tagged elephant shark GR and MR.** HEK293 cell lysates

transfected FLAG-tagged constructs were treated with sample buffer and applied to a 10% SDS-

polyacrylamide gel, and then transferred onto a membrane. The expressed FLAG-tagged

proteins were detected with anti-FLAG antibody. (A) FLAG-tagged elephant shark GR or MR.

(B) CBB stain.

Discussion

The sequencing of lamprey germline genome (35) led to the discovery of two CRs, which

differ only in a small four residue deletion in the DBD of CR2 (8). In lamprey this deletion in

CR2 increases transcriptional activation by corticosteroids. Both elephant shark MR and GR

lack this four amino acid insert in their DBDs, which raised the question of whether adding this

four-residue insert to elephant shark MR and GR would reduce corticosteroid-mediated

transcription, as seen in lamprey CR1 (8) because, in contrast to lamprey CR1 and CR2, elephant

shark MR and GR have different sequences in the LBD (61% identical to each other) and NTD

(21% identical to each other). Both domains are important in their response to corticosteroids

(14). Thus, these sequence differences could obscure the effect of the four-residue insert in the

DBD of elephant shark MR and GR. However, as reported here (Figures 2 and 3, Table 1), we

find that like lamprey CR1, addition of four residues -TRQG- to wild-type elephant shark MR

and GR leads to a reduction in transcriptional activation by a panel of corticosteroids consisting

of cortisol, corticosterone, aldosterone, 11-deoxycorticosterone, 11-deoxycortisol and

progesterone. This effect of the TRQG insert in elephant shark MR and GR is found for

HEK293 cells transfected with either the MMTV or TAT3 promoters (Table 1). The loss of the

TRQG insert in elephant shark GR and MR appears to have provided increased transcriptional

activity for their GR and MR.

Although elephant shark MR lacks an insert in the DBD, a four amino acid (KCSW)

insert is present at this site in the DBD in human MR (46–48), rat MR (46) and *Xenopus laevis*

MR (Figure 5). The insertion of KCSR in the DBD of *Xenopus laevis* MR (Figure 4) indicates

that introduction of a mutation at this position occurred in a basal terrestrial vertebrate.

Transcriptional activation of these terrestrial vertebrate MRs by steroids has not been reported.

We are beginning to investigate corticosteroid and progesterone activation of the human MR

variant.

Figure 5. Alignment of Isoforms of the Human, Rat and Frog Mineralocorticoid Receptor

DNA-Binding Domain.

Receptor (species): accession no.	Sequence
MR-wt (*Homo sapiens*): AAA59571	626 FFKRAVEG----QHNYLCAGRN 643
MR-X1 (*Homo sapiens*): XP_11530277	626 FFKRAVEG**KCSW**QHNYLCAGRN 647
MR-wt (*Rattus norvegicus*): NP_001382006	627 FFKRAVEG----QHNYLCAGRN 644
MR-X1 (*Rattus norvegicus*): XP_038953451	627 FFKRAVEG**KCSW**QHNYLCAGRN 648
MR-wt (*Xenopus laevis*): NP_001084074	623 FFKRAVEG----QHSYLCAGRN 640
MR (*Xenopus laevis*): XP_018098693	623 FFKRAVEG**KCSR**QHSYLCAGRN 644

Wt: wild type, X1: isoform X1, **bold** indicates inserted amino acids, GAPs (-) are introduced to

optimize the sequence alignment.

Meijsing et al (49) found that the function of the DBD in human GR was more than just

docking of the GR to DNA. They showed that the human GR DBD also had an allosteric effect

on transcription by human GR. Our finding that the DBD in elephant shark MR and GR and in

sea lamprey CR has a role in regulating corticosteroid-mediated transcription indicates that a

regulatory function of the DBD appeared early in the evolution of corticosteroid receptors.

Methods

Construction of plasmid vectors

Full-length glucocorticoid receptor (GR) and mineralocorticoid receptor (MR) sequences

of elephant shark, *Callorhinchus milii*, were registered in Genbank (accession number:

XP_042195980 for GR and XP_007902220 for MR). The insertion of 4-amino acids (Thr-Arg-

Gln-Gly) into the DBD of elephant shark MR and GR was performed using KOD-Plus-

mutagenesis kit (TOYOBO).   N-terminal FLAG-tag motif was introduced into both wild type

and mutants containing the 4-amino acid insert in the DBD using PCR. The nucleic acid

sequences of all constructs were verified by sequencing.

Western blotting

FLAG-tagged GR and MR constructs were transfected as described in "Transactivation

assays and statistical analysis" (50). FLAG-tagged proteins were separated by SDS-PAGE in

10 % gel, blotted onto an Immobilon membrane (Millipore Corp. MA), and probed with anti-

FLAG (clone 2H8, TransGenic Inc. (50)) antibody.

Chemical reagents

Cortisol, corticosterone, 11-deoxycorticosterone, 11-deoxycortisol, aldosterone and

progesterone were purchased from Sigma-Aldrich. For reporter gene assays, all hormones were

dissolved in dimethyl-sulfoxide (DMSO); the final DMSO concentration in the culture medium

did not exceed 0.1%.

Transactivation assays and statistical analyses

Transfection and reporter assays were carried out in HEK293 cells, as described

previously (8). The cells were transfected with 100 ng of receptor gene, reporter gene containing

the *Photinus pyralis* luciferase gene and pRL-tk, as an internal control to normalize for variation

in transfection efficiency; pRL-tk contains the *Renilla reniformis* luciferase gene with the herpes

simplex virus thymidine kinase promoter. Each assay had a similar number of cells, and assays

were done with the same batch of cells in each experiment. All experiments were performed in

triplicate. Promoter activity was calculated as firefly (*P. pyralis*)-lucifease activity/sea pansy (*R.*

reniformis)-lucifease activity. The values shown are mean ± SEM from three separate

experiments, and dose-response data, which were used to calculate the half maximal response

(EC50) for each steroid, were analyzed using GraphPad Prism. We note that our assays were

done at 37 degrees, which is not the physiological temperature for the GR and MR in elephant

sharks.

Author Contributions

Conceptualization: Yoshinao Katsu, Michael E. Baker.

Data curation: Yoshinao Katsu, Jiawen Zhang

Formal analysis: Yoshinao Katsu, Michael E. Baker.

Investigation: Yoshinao Katsu, Jiawen Zhang

Methodology: Yoshinao Katsu.

Supervision: Yoshinao Katsu, Michael E. Baker.

Writing – original draft: Yoshinao Katsu, Michael E. Baker.

Writing – review & editing: Yoshinao Katsu, Michael E. Baker.

Funding: This work was supported by Grants-in-Aid for Scientific Research from the Ministry

of Education, Culture, Sports, Science and Technology of Japan (19K067309 and 23K05839 to

Y.K.), and the Takeda Science Foundation (to Y.K.).

274
References

1. Osório J, Rétaux S. The lamprey in evolutionary studies. Dev Genes Evol. 2008
May;218(5):221-35. doi: 10.1007/s00427-008-0208-1. Epub 2008 Feb 15. PMID: 18274775.

2. Shimeld SM, Donoghue PC. Evolutionary crossroads in developmental biology: cyclostomes
(lamprey and hagfish). Development. 2012 Jun;139(12):2091-9. doi: 10.1242/dev.074716.
PMID: 22619386.

3. Smith JJ, Kuraku S, Holt C, Sauka-Spengler T, Jiang N, Campbell MS, Yandell MD,
Manousaki T, Meyer A, Bloom OE, Morgan JR, Buxbaum JD, Sachidanandam R, Sims C,
Garruss AS, Cook M, Krumlauf R, Wiedemann LM, Sower SA, Decatur WA, Hall JA,
Amemiya CT, Saha NR, Buckley KM, Rast JP, Das S, Hirano M, McCurley N, Guo P,
Rohner N, Tabin CJ, Piccinelli P, Elgar G, Ruffier M, Aken BL, Searle SM, Muffato M,
Pignatelli M, Herrero J, Jones M, Brown CT, Chung-Davidson YW, Nanlohy KG, Libants
SV, Yeh CY, McCauley DW, Langeland JA, Pancer Z, Fritzsch B, de Jong PJ, Zhu B, Fulton
LL, Theising B, Flicek P, Bronner ME, Warren WC, Clifton SW, Wilson RK, Li W.
Sequencing of the sea lamprey (Petromyzon marinus) genome provides insights into
vertebrate evolution. Nat Genet. 2013 Apr;45(4):415-21, 421e1-2. doi: 10.1038/ng.2568.
Epub 2013 Feb 24. PMID: 23435085; PMCID: PMC3709584.

4. Nakatani Y, Shingate P, Ravi V, Pillai NE, Prasad A, McLysaght A, Venkatesh B.
Reconstruction of proto-vertebrate, proto-cyclostome and proto-gnathostome genomes
provides new insights into early vertebrate evolution. Nat Commun. 2021 Jul 23;12(1):4489.
doi: 10.1038/s41467-021-24573-z. Erratum in: Nat Commun. 2021 Jul 29;12(1):4704.
PMID: 34301952; PMCID: PMC8302630.

5. Thornton JW. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by
ligand exploitation and serial genome expansions. Proc Natl Acad Sci U S A. 2001 May
8;98(10):5671-6. doi: 10.1073/pnas.091553298. Epub 2001 May 1. PMID: 11331759;
PMCID: PMC33271.

6. Carroll SM, Bridgham JT, Thornton JW. Evolution of hormone signaling in elasmobranchs by
exploitation of promiscuous receptors. Mol Biol Evol. 2008;25(12):2643-2652.
doi:10.1093/molbev/msn204.

7. Baker ME, Katsu Y. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: Evolution
of the mineralocorticoid receptor: sequence, structure and function. J Endocrinol.
2017;234(1):T1-T16. doi:10.1530/JOE-16-0661.

8. Katsu Y, Lin X, Ji R, Chen Z, Kamisaka Y, Bamba K, Baker ME. N-terminal domain
influences steroid activation of the Atlantic sea lamprey corticoid receptor. J Steroid
Biochem Mol Biol. 2023 Jan 13;228:106249. doi: 10.1016/j.jsbmb.2023.106249. Epub ahead
of print. PMID: 36646152.

9. Evans RM. The steroid and thyroid hormone receptor superfamily. Science.
1988;240(4854):889-895. doi:10.1126/science.3283939.

10. Bridgham JT, Eick GN, Larroux C, et al. Protein evolution by molecular tinkering:
diversification of the nuclear receptor superfamily from a ligand-dependent ancestor. PLoS
Biol. 2010;8(10):e1000497. Published 2010 Oct 5. doi:10.1371/journal.pbio.1000497.

11. Beato M, Klug J. Steroid hormone receptors: an update. Hum Reprod Update. 2000 May-
Jun;6(3):225-36. doi: 10.1093/humupd/6.3.225. PMID: 10874567.

12. Whitfield GK, Jurutka PW, Haussler CA, Haussler MR. Steroid hormone receptors:
evolution, ligands, and molecular basis of biologic function. J Cell Biochem. 1999;Suppl 32-
33:110-22. doi: 10.1002/(sici)1097-4644(1999)75:32+<110::aid-jcb14>3.0.co;2-t. PMID:
10629110.

13. Baker ME. Steroid receptors and vertebrate evolution. Mol Cell Endocrinol.
2019;496:110526. doi:10.1016/j.mce.2019.110526.

14. Katsu Y, Shariful IMD, Lin X, Takagi W, Urushitani H, Kohno S, Hyodo S, Baker ME. N-
terminal Domain Regulates Steroid Activation of Elephant Shark Glucocorticoid and
Mineralocorticoid Receptors. J Steroid Biochem Mol Biol. 2021 Feb 27:105845. doi:
10.1016/j.jsbmb.2021.105845. Epub ahead of print. PMID: 33652098.

15. Fonseca E, Machado AM, Vilas-Arrondo N, Gomes-Dos-Santos A, Veríssimo A, Esteves P,
Almeida T, Themudo G, Ruivo R, Pérez M, da Fonseca R, Santos MM, Froufe E, Román-
Marcote E, Venkatesh B, Castro LFC. Cartilaginous fishes offer unique insights into the
evolution of the nuclear receptor gene repertoire in gnathostomes. Gen Comp Endocrinol.
2020 Sep 1;295:113527. doi: 10.1016/j.ygcen.2020.113527. Epub 2020 Jun 8. PMID:
32526329.

16. Shibata S. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: Mineralocorticoid
receptor and NaCl transport mechanisms in the renal distal nephron. J Endocrinol.
2017;234(1):T35-T47. doi:10.1530/JOE-16-0669.

17. Lifton RP, Gharavi AG, Geller DS. Molecular mechanisms of human hypertension. Cell.
2001;104(4):545-556. doi:10.1016/s0092-8674(01)00241-0.

18. Lombes M, Kenouch S, Souque A, Farman N, Rafestin-Oblin ME. The Mineralocorticoid
Receptor Discriminates Aldosterone from Glucocorticoids Independently of the 11 Beta-
Hydroxysteroid Dehydrogenase. Endocrinology. 1994;135(3):834-840.
Doi:10.1210/Endo.135.3.8070376.

19. Pascual-Le Tallec L, Lombès M. The mineralocorticoid receptor: a journey exploring its
diversity and specificity of action. Mol Endocrinol. 2005 Sep;19(9):2211-21. doi:
10.1210/me.2005-0089. Epub 2005 Mar 31. PMID: 15802372.

20. Rossier BC, Baker ME, Studer RA. Epithelial sodium transport and its control by
aldosterone: the story of our internal environment revisited. Physiol Rev. 2015;95(1):297-
340. doi:10.1152/physrev.00011.2014.

21. Hanukoglu I, Hanukoglu A. Epithelial sodium channel (ENaC) family: Phylogeny, structure-
function, tissue distribution, and associated inherited diseases. Gene. 2016;579(2):95-132.
doi:10.1016/j.gene.2015.12.061.

22. Jaisser F, Farman N. Emerging Roles of the Mineralocorticoid Receptor in Pathology:
Toward New Paradigms in Clinical Pharmacology. Pharmacol Rev. 2016;68(1):49-75.
doi:10.1124/pr.115.011106.

23. Hawkins UA, Gomez-Sanchez EP, Gomez-Sanchez CM, Gomez-Sanchez CE. The
ubiquitous mineralocorticoid receptor: clinical implications. Curr Hypertens Rep.
2012;14(6):573-580. doi:10.1007/s11906-012-0297-0.

24. de Kloet ER, Joëls M. Brain mineralocorticoid receptor function in control of salt balance
and stress-adaptation. Physiol Behav. 2017 Sep 1;178:13-20. doi:
10.1016/j.physbeh.2016.12.045. Epub 2017 Jan 13. PMID: 28089704.

25. Gomez-Sanchez EP. Brain mineralocorticoid receptors in cognition and cardiovascular
homeostasis. Steroids. 2014 Dec;91:20-31. doi: 10.1016/j.steroids.2014.08.014. PMID:
25173821; PMCID: PMC4302001.

26. Gomez-Sanchez CE, Gomez-Sanchez EP. The Mineralocorticoid Receptor and the Heart.
Endocrinology. 2021 Nov 1;162(11):bqab131. doi: 10.1210/endocr/bqab131. PMID:
34175946.

27. de Kloet ER. From receptor balance to rational glucocorticoid therapy. Endocrinology.
2014;155(8):2754-2769. doi:10.1210/en.2014-1048.

28. Chrousos GP. Stress and sex versus immunity and inflammation. Sci Signal. 2010 Oct
12;3(143):pe36. doi: 10.1126/scisignal.3143pe36. PMID: 20940425.

29. Cain DW, Cidlowski JA. Immune regulation by glucocorticoids. Nat Rev Immunol.
2017;17(4):233-247. doi:10.1038/nri.2017.1.

30. Weikum ER, Knuesel MT, Ortlund EA, Yamamoto KR. Glucocorticoid receptor control of
transcription: precision and plasticity via allostery. Nat Rev Mol Cell Biol. 2017;18(3):159-
174. doi:10.1038/nrm.2016.152.

31. Gross KL, Lu NZ, Cidlowski JA. Molecular mechanisms regulating glucocorticoid
sensitivity and resistance. Mol Cell Endocrinol. 2009;300(1-2):7-16.
doi:10.1016/j.mce.2008.10.001.

32. Close DA, Yun SS, McCormick SD, Wildbill AJ, Li W. 11-deoxycortisol is a corticosteroid
hormone in the lamprey. Proc Natl Acad Sci U S A. 2010 Aug 3;107(31):13942-7. doi:
10.1073/pnas.0914026107. Epub 2010 Jul 19. PMID: 20643930; PMCID: PMC2922276.

33. Smith JJ, Antonacci F, Eichler EE, Amemiya CT. Programmed loss of millions of base pairs
from a vertebrate genome. Proc Natl Acad Sci U S A. 2009 Jul 7;106(27):11212-7. doi:
10.1073/pnas.0902358106. Epub 2009 Jun 26. PMID: 19561299; PMCID: PMC2708698.

34. Smith JJ, Saha NR, Amemiya CT. Genome biology of the cyclostomes and insights into the
evolutionary biology of vertebrate genomes. Integr Comp Biol. 2010 Jul;50(1):130-7. doi:
10.1093/icb/icq023. Epub 2010 Apr 19. PMID: 21558194; PMCID: PMC3140258.

35. Smith JJ, Timoshevskaya N, Ye C, Holt C, Keinath MC, Parker HJ, Cook ME, Hess JE,
Narum SR, Lamanna F, Kaessmann H, Timoshevskiy VA, Waterbury CKM, Saraceno C,
Wiedemann LM, Robb SMC, Baker C, Eichler EE, Hockman D, Sauka-Spengler T, Yandell
M, Krumlauf R, Elgar G, Amemiya CT. The sea lamprey germline genome provides insights
into programmed genome rearrangement and vertebrate evolution. Nat Genet. 2018
Feb;50(2):270-277. doi: 10.1038/s41588-017-0036-1. Epub 2018 Jan 22. Erratum in: Nat
Genet. 2018 Apr 19;: Erratum in: Nat Genet. 2018 Nov;50(11):1617. PMID: 29358652;
PMCID: PMC5805609.

36. Beato M, Arnemann J, Chalepakis G, Slater E, Willmann T. Gene regulation by steroid
hormones. J Steroid Biochem. 1987;27(1-3):9-14. doi: 10.1016/0022-4731(87)90288-3.
PMID: 2826895.

37. Cato AC, Skroch P, Weinmann J, Butkeraitis P, Ponta H. DNA sequences outside the
receptor-binding sites differently modulate the responsiveness of the mouse mammary
tumour virus promoter to various steroid hormones. EMBO J. 1988 May;7(5):1403-10.
PMID: 2842149; PMCID: PMC458390.

38. Iñiguez-Lluhí JA, Pearce D. A common motif within the negative regulatory regions of
multiple factors inhibits their transcriptional synergy. Mol Cell Biol. 2000 Aug;20(16):6040-
50. doi: 10.1128/mcb.20.16.6040-6050.2000. PMID: 10913186; PMCID: PMC86080.

39. Lombès M, Binart N, Oblin ME, Joulin V, Baulieu EE. Characterization of the interaction of
the human mineralocorticosteroid receptor with hormone response elements. Biochem J.
1993 Jun 1;292 (Pt 2)(Pt 2):577-83. doi: 10.1042/bj2920577. PMID: 8389140; PMCID:
PMC1134249.

40. Geller DS, Farhi A, Pinkerton N, et al. Activating mineralocorticoid receptor mutation in
hypertension exacerbated by pregnancy. Science. 2000;289(5476):119-123.
doi:10.1126/science.289.5476.119.

41. Fuller PJ, Yao YZ, Jin R, et al. Molecular evolution of the switch for progesterone and
spironolactone from mineralocorticoid receptor agonist to antagonist. Proc Natl Acad Sci U
S A. 2019;116(37):18578-18583. doi:10.1073/pnas.1903172116, (n.d.).

42. Baker ME, Katsu Y. Progesterone: An enigmatic ligand for the mineralocorticoid receptor.
Biochem Pharmacol. 2020;177:113976. doi:10.1016/j.bcp.2020.113976.

43. Baker ME. Divergent evolution of progesterone and mineralocorticoid receptors in terrestrial
vertebrates and fish influences endocrine disruption. Biochem Pharmacol. 2022
Apr;198:114951. doi: 10.1016/j.bcp.2022.114951. Epub 2022 Feb 8. PMID: 35149051.

44. Sugimoto A, Oka K, Sato R, Adachi S, Baker ME, Katsu Y. Corticosteroid and progesterone
transactivation of mineralocorticoid receptors from Amur sturgeon and tropical gar. Biochem
J. 2016;473(20):3655-3665. doi:10.1042/BCJ20160579, (n.d.).

45. Katsu Y, Kohno S, Oka K, et al. Transcriptional activation of elephant shark
mineralocorticoid receptor by corticosteroids, progesterone, and spironolactone. Sci Signal.
2019;12(584):eaar2668. Published 2019 Jun 4. doi:10.1126/scisignal.aar2668, (n.d.).

46. Bloem LJ, Guo C, Pratt JH. Identification of a splice variant of the rat and human
mineralocorticoid receptor genes. J Steroid Biochem Mol Biol. 1995 Nov;55(2):159-62. doi:
10.1016/0960-0760(95)00162-s. PMID: 7495694.

47. Wickert L, Selbig J, Watzka M, Stoffel-Wagner B, Schramm J, Bidlingmaier F, Ludwig M.
Differential mRNA expression of the two mineralocorticoid receptor splice variants within
the human brain: structure analysis of their different DNA binding domains. J
Neuroendocrinol. 2000 Sep;12(9):867-73. doi: 10.1046/j.1365-2826.2000.00535.x. PMID:
10971811.

48. Wickert L, Watzka M, Bolkenius U, Bidlingmaier F, Ludwig M. Mineralocorticoid receptor
splice variants in different human tissues. Eur J Endocrinol. 1998 Jun;138(6):702-4. doi:
10.1530/eje.0.1380702. PMID: 9678540.

49. Meijsing SH, Pufall MA, So AY, Bates DL, Chen L, Yamamoto KR. DNA binding site
sequence directs glucocorticoid receptor structure and activity. Science. 2009 Apr
17;324(5925):407-10. doi: 10.1126/science.1164265. PMID: 19372434; PMCID:
PMC2777810.

50. Sasaki F, Okuno T, Saeki K, Min L, Onohara N, Kato H, Shimizu T, Yokomizo T. A high-
affinity monoclonal antibody against the FLAG tag useful for G-protein-coupled receptor
study. Anal Biochem. 2012 Jun 15;425(2):157-65. doi: 10.1016/j.ab.2012.03.014. Epub 2012
Mar 27. PMID: 22465329.

446