Toxin/Antitoxin Systems Induce Persistence and Work in Concert with Restriction/Modification Systems to Inhibit Phage

Laura Fernández-García[1,2], Sooyeon Song[1,3,4], Joy Kirigo[1], Maiken E. Petersen[1], María Tomás[2], and Thomas K. Wood[1*]

[1]Department of Chemical Engineering, Pennsylvania State University, University Park, Pennsylvania, 16802-4400, USA

[2]Microbiology Translational and Multidisciplinary (MicroTM)-Research Institute Biomedical A Coruña (INIBIC) and Microbiology Department of Hospital A Coruña (CHUAC); University of A Coruña (UDC), A Coruña, Spain

Departments of Animal Science[3] and Agricultural Convergence Technology[4], Jeonbuk National University, 587 Baekje-Daero, Deojin-Gu, Jeonju-Si, Jellabuk-Do, 54896, South Korea

*For correspondence. E-mail tuw14@psu.edu

Tel. (+)1 814-863-4811; Fax (1) 814-865-7846

Running title: Phage inhibition induces persistence.

Keywords: Phage inhibition, persistence, CRISPR-Cas, toxin/antitoxin systems

SUMMARY

Myriad bacterial anti-phage systems have been described and often the mechanism of programmed cell death is invoked for phage inhibition. However, there is little evidence of "suicide" under physiological conditions for these systems. Instead of death to stop phage propagation, we show here that persister cells are formed and survive using the *Escherichia coli* C496_10 tripartite toxin/antitoxin system MqsR/MqsA/MqsC that degrades mRNA. Specifically, MqsR/MqsA/MqsC inhibited T2 phage by 10^6-fold and reduced T2 titers by 500-fold. Further evidence of the formation of persister cells during T2 phage attack in the presence of MqsR/MqsA/MqsC include single-cell physiological changes: reduced metabolism (via flow cytometry) and heterogeneous resuscitation. Critically, we found restriction-modification systems (primarily EcoK McrBC) work in concert with the toxin/antitoxin system to inactivate phage, likely while the cells are in the persister state. Hence, phage attack invokes a stress response similar to antibiotics, starvation, and oxidation, which leads to persistence, and this dormant state likely allows restriction/modification systems to clear phage DNA.

INTRODUCTION

Phage attack, resulting in up to 40% cell death[13], is the universal constraint on bacterial immortality; hence, bacteria have developed numerous phage-inhibition systems. For phage inhibition, there is little evidence that these systems cause cell death; instead, it is much more likely these systems invoke dormancy[31, 44]. Bacterial cell dormancy has several potential benefits: (i) allowing time for other host-encoded, phage-inhibition systems to function, (ii) slowing viral replication, and (iii) accommodating for spacer accumulation for CRISPR-Cas[44]. In fact, the very existence of spacers in CRISPR-Cas type III and VI systems; i.e., those shown to induce dormancy by targeting host RNA, demonstrate cells must survive phage attack to obtain spacers. Also, growth-inhibited cells have more spacer acquisition[9]. In contrast, programmed cell death during phage attack is paradoxical from the perspective of individual cells and has yet to be proven beneficial for kin.

Given that cells enter a state of dormancy known as persistence to survive numerous stresses (e.g., antibiotics[15], starvation[17], oxidation[16]) through guanosine pentaphosphate and tetraphosphate (p)ppGpp signaling[19], which leads to dimerization of their ribosomes to cease translation[21, 41, 50], we reasoned that phage attack is another form of extreme stress, so it should lead to persistence. Also, phage-inhibition systems include toxin proteins of toxin/antitoxin systems, which are known to induce persistence by up to 14,000-fold[7, 16, 46]. For example, after transcription shutoff by T4 phage activates the Hok/Sok toxin/antitoxin system, Hok damages the cell membrane to cease cell metabolism to halt phage propagation, in the first link of a toxin/antitoxin system to phage inhibition (1996)[32]. Similarly, phage attack activates various toxins to reduce cell metabolism including those of the CBASS (phospholipases, nucleases, and pore-forming proteins[22]), RADAR (RdrB converts ATP to ITP[11]), and Thoeris (ThsA degrades NAD^{+}[23]) phage inhibition systems.

Therefore, based on these two insights (extreme stress produces persisters and phage-inhibition systems utilize toxins to reduce metabolism), we investigated whether persister cells are formed as a result of inducing the *bona fide Escherichia* spp. phage inhibition system, the tripartite toxin/antitoxin/chaperone system *E. coli* C496_10 MqsR/MqsA/MqsC[4, 45], which utilizes RNase MqsR[5, 35] upon phage attack. We

40   confirmed that the MqsR/MqsA/MqsC, produced from its natural promoter via the same system used to

find it dramatically inhibits T2 phage[45], and discovered this system converts *E. coli* into the persister state

to inhibit phage. In addition, we discovered that restriction/modification systems assist the

MqsR/MqsA/MqsC toxin/antitoxin system to inhibit T2 phage. To our knowledge, this is the is the first

connection of phage inhibition with persistence, and the first non-CRISPR-Cas system to be linked to

45   restriction/modification systems.

RESULTS

Given ours and others' extensive experience with *E. coli* persister cells[16, 20, 38-41, 49], we chose a phage

exclusion system of *Escherichia* spp. to investigate the hypothesis that persister cells are formed upon phage

attack. We also used the same expression system that was used to demonstrate phage inhibition and used

50   T2 phage since it was inhibited best by the MqsR/MqsA/MqsC system[10, 45], and produced

MqsR/MqsA/MqsC in a background that lacks this system (K-12).

MqsR/MqsA/MqsC inhibits T2 phage. As published previously[45], T2 phage was efficiently inhibited by

the MqsR/MqsA/MqsC TA system in our hands; i.e., MqsR/MqsA/MqsC increased cell viability by 10^6-

fold compared to an empty plasmid control (**Table S1**). Corroborating the phage inhibition, the number of

55   T2 phage decreased 500-fold in the presence of MqsR/MqsA/MqsC: $6 \pm 2 \times 10^5$ pfu/mL were obtained for

MqsR/MqsA/MqsC whereas $2.6 \pm 0.5 \times 10^8$ pfu/mL were obtained for the empty plasmid when T2 was

added initially at 7.14×10^6 pfu (MOI of 0.1). Therefore, MqsR/MqsA/MqsC actively inhibits T2 phage

under these conditions.

Cells surviving T2 phage attack with MqsR/MqsA/MqsC are persisters. Critically, treatment with T2

60   phage induced persistence in 8% of the cells producing MqsR/MqsA/MqsC (**Table S1**); i.e., these cells

survived treatment of ampicillin at 10X the minimum inhibitory concentration (MIC). In contrast, cells

bearing the empty plasmid were completely eradicated by ampicillin after exposure to T2 phage indicating

no cells entered the persister state without MqsR/MqsA/MqsC. Normally, persister cells are formed at less

than 1% of the population[27, 28], so the number of persisters created by MqsR/MqsA/MqsC during T2 phage

65   attack is striking (~10%). Similar results were seen with a different class of antibiotic, ciprofloxacin (10X MIC), in that 0.28% cells with the MqsR/MqsA/MqsC phage inhibition system were persisters after T2 attack (**Table S1**), whereas no cells survived T2 and ciprofloxacin in the absence of MqsR/MqsA/MqsC (**Table S1**). Moreover, the increase in persistence by MqsR/MqsA/MqsC was not due to a difference in growth since $mqsRAC^+$ had virtually no impact on the specific growth rate in the absence of phage (**Fig.**

70   **S1**, $\mu = 1.7 \pm 0.1$/h, $\mu = 1.62 \pm 0.06$/h, and $\mu = 1.74 \pm 0.09$/h for MG1655, MG1655/pCV1, and MG1655/pCV1-mqsRAC, respectively).

Corroborating that MqsR/MqsA/MqsC converts *E. coli* into persister cells, we assayed single cell resuscitation on LB gel pads after 1 h of T2 attack (0.1 MOI) and found significant heterogeneity in waking with four phenotypes including 36 ± 13% immediate waking, 28 ± 5% lags in waking, 18 ± 7% elongation,

75   6% waking then death) whereas 97 ± 3% of the sample of exponentially-growing cells wake uniformly without elongation, lags, or death (**Fig. 1**, **Table S2**). Heterogeneity in resuscitation of single-cell persisters was discovered by us[18] and confirmed by other laboratories[12, 34, 37, 48]; hence, after T2 attack, cells become persistent based on their antibiotic tolerance to two different classes of antibiotics and based on their heterogeneity in resuscitation. Further evidence of persistence after T2 phage attack includes metabolic

80   staining with flow cytometry revealed that after 1 h of T2 attack, *E. coli* cells had significantly reduced metabolism compared to exponentially-growing cells (**Fig. S2**).

Persistence cell formation is rapid upon phage attack. To investigate how rapidly persister cells form upon phage attack, we investigated induction of the three loci required to inactivate ribosomes: *raiA, rmf,* and *hpf*[41]. Using qRT-PCR, we found no induction of these loci for the cells producing the

85   MqsR/MqsA/MqsC phage inhibition systems, which indicates the cell uses existing proteins to become dormant rapidly (i.e., in msec), rather than relying on expression of ribosome inactivation genes once phage attacks. However, for cells that lack this phage inhibition system, *raiA* was induced 4-fold at 15 min after phage attack, which indicates these cells are stressed as they unsuccessfully try to thwart T2 phage (**Table S3**). Bacteriophage T2 takes 60 minutes for its burst size, with a latent period of around 20 min and a rise

90   period of 10 min[26].

Restriction/modification assists MqsR/MqsA/MqsC during persistence. Since cells with MqsR/MqsA/MqsC weather T2 attack by rapidly becoming persistent and resuscitate, which indicates inactivation of phage, we hypothesized that another system works while the cells are dormant to clear T2 phage. Therefore, we investigated whether cells that lack the restriction enzyme systems encoded by *hsdR,* *mrr,* or *mcrB* are susceptible to T2 attack and found that inactivation of all of these restriction/modification systems reduces cell viability, with inactivation of the McrBC system most important for defending against T2 since its inactivation reduces cell viability by 100,000-fold (**Table S4A**). Critically, we found that the MqsR/MqsA/MqsC toxin/antitoxin works with the McrBC restriction/modification system since cells deficient in McrBC had 10-fold less survival in the presence of MqsR/MqsA/MqsC (**Table S4B**). Therefore, MqsR/MqsA/MqsC works in concert with restriction/modification systems to inhibit T2 phage.

DISCUSSION

Our results demonstrate phage inhibition systems form persister cells. Moreover, this persistence allows at least some of the cells to survive phage attack by providing time for phage DNA degradation by restriction/modification systems. Hence, our results suggest toxin/antitoxin systems act first to reduce metabolism, followed by restriction/modification systems, to undermine phage attack. Similarly, CRISPR-Cas has been shown to act first, followed by restriction/modification systems in *Listeria* spp[47]. Our results showing synergy between toxin/antitoxin and restriction/modification systems are significant since both of these phage defense systems are found in nearly all bacteria (unlike many other phage inhibition systems such as CRISPR-Cas, CBASS and retrons that are found in 39%, 13% and 11% of procaryote genomes, respectively[43]), which suggests this interaction is widespread and may be dominant for phage inhibition. Our results are also provide the first reproducible link between toxin/antitoxin systems and persister cells under physiological conditions (rather than through overproduced toxins and discredited deletion strains[42]).

These results also have implications for other phage inhibition systems in that, rather than causing cell death as frequently claimed with little evidence[1, 24, 25, 29] or claimed after non-physiological overexpression[10], these anti-phage systems probably induce dormancy; i.e., persistence, which allows the

cell to thwart the phage in the same manner as we studied here. Along with the formation of persister cells, some cells indubitably succumb to phage attack; however, our results indicate the primary purpose of the anti-phage systems is not what has been termed 'abortive infection' in which the phage inhibition system kills the cells[29], but instead, these systems protect the cells and allow them to survive phage attack. Similar results showing dormancy have been seen with type VI CRISPR-Cas and restriction/modification systems[47].

Corroborating our findings, many phage-inhibition systems reduce energy metabolites like ATP (RADAR[11] and Detocs[36]) and NAD$^+$ (Thoeris[23]), and the reduction of energy metabolites like ATP consistently induces persistence[8, 21]. Furthermore, reduction of ATP by toxin/antitoxin systems leads to persistence[6].

Central to persister cell formation is accumulation of (p)ppGpp[21, 41, 50]; hence, it is expected that if cells prevent phage propagation by becoming persisters, they will do so via an increase in (p)ppGpp. Therefore, to counter this bacterial defense, if our hypothesis/data are correct in this report, phages should try to reduce (p)ppGpp levels. Critically, this has been recently shown with *Pectobacterium* spp. phage PcCB7V, which encodes the nucleotide pyrophosphatase MazG that is enriched against TIR-STING anti-phage systems and reduces (p)ppGpp[14]. In addition, these authors demonstrated that (p)ppGpp is increased during phage attack[14].

Our model (**Fig. 2**) is that by invoking persistence (i.e., dormancy) through phage attack, back-up phage inhibition systems like restriction modification systems can degrade the phage genome since restriction enzymes do not require any resources from the cell for activity; i.e., Gibbs free energy is less than zero for phage DNA degradation due to the increase in entropy, so degradation of the phage genome can occur while the cells sleep. This general concept of allowing back-up DNA-targeting enzymes to function while the cell is dormant has been proposed for the type III/VI CRISPR-Cas systems[30, 44] and demonstrated while this work was being prepared for type VI CRISPR-Cas in *Listeria* spp.[47]; however, the benefit of dormancy has not been suggested for the plethora of recently-discovered, non-CRSPR-Cas phage inhibition systems. Here we demonstrate that (i) dormancy in the form of persistence is a key aspect of phage inhibition of non-

CRISPR-Cas systems and that (ii) restriction/modification is utilized during the dormant state.

Since the cells are becoming dormant (i.e., persisters) rather than dying, our results also suggest that after phage attack, the cells which survive phage, resuscitate upon detecting nutrients, as we have shown using single cell microscopy for persister cells[40, 41, 50], by using chemotaxis proteins and sugar transport membrane proteins, by reducing cAMP and ppGpp internal signals, and by increasing HflX activity to convert dimerized ribosomes (100S ribosomes) into active 70S ribosomes.

Finally, our results show that, like myriad other stresses such as starvation[17], oxidation[16], and antibiotic challenge[21], bacteria form persister cells to survive the extreme stress of phage attack. Hence, the formation of persister cells is a consistent and conserved means to weather stress, which we suggest as our **Conserved** **Bacterial** **Stress** **Survival** (**CBaSS**) theory.

<center>MATERIALS AND METHODS</center>

Bacteria and growth conditions. Bacteria and plasmids are shown in **Table S5**, and cells were cultured at 37°C in LB supplemented with 30 µg/ml of chloramphenicol (to retain plasmids). The complete MqsR/MqsA/MqsC plasmid-based system (pCV1-*mqsRAC* and pCV1 empty plasmid control) was verified via Plasmidsaurus sequencing.

Persister formation assay. Single colonies were cultured overnight in LB-chloramphenicol, then diluted 100X and grown to a turbidity of 0.5 at 600 nm. T2 phage was then added (multiplicity of infection ~ 0.1) for one hour, then the cells were washed twice with phosphate buffered saline (PBS, 8 g of NaCl, 0.2 g of KCl, 1.15 g of Na_2HPO_4, and 0.2 g of KH_2PO_4 in 1000 mL of dH_2O) and cultured for three hours with LB-ampicillin (100 µg/mL, 10 MIC) or LB-ciprofloxacin (5 µg/mL, 10 MIC) to kill non-persister cells[21]. Cells were washed twice with PBS and enumerated on LB-Cm plates to measure persistence. To determine phage titers, after 1 h of phage contact, supernatants were diluted with PBS, and plaques were enumerated on top agar, TA, (1% tryptone, 0.5% NaCl, 1.5% agar lower layer and 0.4% agar top layer) double layer plates.

Persister resuscitation assay. Single-cell microscopy of resuscitating persister cells was performed as described previously[18] using LB agarose gel pads that were observed up to 3 h via a light microscope (Zeiss

Axioscope.A1, bl_ph channel at 1000 ms exposure). The persister cells were generated by the addition of T2 phage at 0.1 MOI for 1 h. The microscope was maintained at 37°C by placing it in a vinyl glove box (Coy Labs), which was warmed by an anaerobic chamber heater (Coy Labs, 8535-025).

170 **Metabolic activity via flow cytometry**. To ascertain metabolic activity, *E. coli* MG1655/pCV1-*mqsRAC* was grown to a turbidity of 0.5 at 600 nm and T2 phage was added (MOI 0.1) for 1 h. Cells were washed, resuspended in PBS, stained for metabolic activity (BacLight™ RedoxSensor™ Green Vitality Kit, Thermo Fisher Scientific Inc., Waltham, MA, USA), which measures the cellular redox state. The fluorescence signal of 100,000 cells was analyzed by flow cytometry (Beckman Coulter LSRFortessa) using the FL1

175 $(530 \pm 30$ nm).

qRT-PCR. To quantify transcription after T2 phage attack (15 min at MOI 0.1) in the presence of MqsR/MqsA/MqsC, RNA was isolated by cooling rapidly using ethanol/dry ice in the presence of RNA Later (Sigma) using the High Pure RNA Isolation Kit (Roche). The following qRT-PCR thermocycling protocol was used with the iTaq™ universal SYBR® Green One-Step kit (Bio-Rad): 95 °C for 5 min; 40

180 cycles of 95 °C for 15 s, 60°C for 1 min for two replicate reactions for each sample/primer pair. The annealing temperature was 60°C for all primers. Primers are shown in **Table S6**.

ACKNOWLEDGEMENTS

This work was supported by both a Fulbright Scholar Fellowship and a Xunta de Galicia Postdoctoral grant for LFG and a National Research Foundation of Korea (NRF) grant from the Korean Government

185 (NRF-2020R1F1A1072397) for SYS. This study was also funded by grants PI22/00323 and PMP/00092 awarded to MT within the State Plan for R+D+I 2013-2016 (National Plan for Scientific Research, Technological Development and Innovation 2008-2011) and co-financed by the ISCIII-Deputy General Directorate for Evaluation and Promotion of Research - European Regional Development Fund "A way of Making Europe" and Instituto de Salud Carlos III FEDER. We are grateful for the MqsR/MqsA/MqsC

190 plasmids provided by Prof. Michael Laub.

REFERENCES

1. **Arias CF, Acosta FJ, Bertocchini F, Herrero MA, Fernández-Arias C.** 2022. The coordination of anti-phage immunity mechanisms in bacterial cells. Nature Communications **13:**7412.
2. **Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H.** 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol. Syst. Biol. **2:**2006 0008.
3. **Blattner FR, III GP, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J, Glasner JD, Rode CK, Mayhew GF, Gregor J, Davis NW, Kirkpatrick HA, Goeden MA, Rose DJ, Mau B, Shao Y.** 1997. The complete genome sequence of *Escherichia coli* K-12. Science **277:**1453-1462.
4. **Bordes P, Sala AJ, Ayala S, Texier P, Slama N, Cirinesi A-M, Guillet V, Mourey L, Genevaux P.** 2016. Chaperone addiction of toxin–antitoxin systems. Nature Communications **7:**13339.
5. **Brown BL, Grigoriu S, Kim Y, Arruda JM, Davenport A, Wood TK, Peti W, Page R.** 2009. Three dimensional structure of the MqsR:MqsA complex: A novel toxin:antitoxin pair comprised of a toxin homologous to RelE and an antitoxin with unique properties. PLoS Pathog. **5:**e1000706.
6. **Cheng H-Y, Soo VWC, Islam S, McAnulty MJ, Benedik MJ, Wood TK.** 2014. Toxin GhoT of the GhoT/GhoS toxin/antitoxin system damages the cell membrane to reduce adenosine triphosphate and to reduce growth under stress. Environmental Microbiology **16:**1741-1754.
7. **Chowdhury N, Kwan BW, Wood TK.** 2016. Persistence Increases in the Absence of the Alarmone Guanosine Tetraphosphate by Reducing Cell Growth. Sci Rep **6:**20519.
8. **Conlon BP, Rowe SE, Gandt AB, Nuxoll AS, Donegan NP, Zalis EA, Clair G, Adkins JN, Cheung AL, Lewis K.** 2016. Persister formation in *Staphylococcus aureus* is associated with ATP depletion. Nature Microbiology **1:**16051.
9. **Dimitriu T, Kurilovich E, Łapińska U, Severinov K, Pagliara S, Szczelkun MD, Westra ER.** 2022. Bacteriostatic antibiotics promote CRISPR-Cas adaptive immunity by enabling increased spacer acquisition. Cell Host & Microbe **30:**31-40.e35.
10. **Duncan-Lowey B, McNamara-Bordewick NK, Tal N, Sorek R, Kranzusch PJ.** 2021. Effector-mediated membrane disruption controls cell death in CBASS antiphage defense. Molecular Cell **81:**5039-5051.e5035.
11. **Duncan-Lowey B, Tal N, Johnson AG, Rawson S, L.Mayer M, Doron S, Millman A, Melamed S, Fedorenko T, Kacen A, Amitai G, Sorek R, Kranzusch PJ.** 2022. Cryo-EM structure of the RADAR supramolecular anti-phage defense complex. bioRxiv.
12. **Goormaghtigh F, Van Melderen L.** 2019. Single-cell imaging and characterization of *Escherichia coli* persister cells to ofloxacin in exponential cultures. Science Advances **5:**eaav9462.
13. **Hampton HG, Watson BNJ, Fineran PC.** 2020. The arms race between bacteria and their phage foes. Nature **577:**327-336.
14. **Ho P, Chen Y, Biswas S, Canfield E, Feldman DE.** 2022. Bacteriophage anti-defense genes that neutralize TIR and STING immune responses. bioRxiv**:**2022.2006.2009.495361.
15. **Hobby GL, Meyer K, Chaffee E.** 1942. Observations on the Mechanism of Action of Penicillin. Exp Biol Med **50:**281-285.
16. **Hong SH, Wang X, O'Connor HF, Benedik MJ, Wood TK.** 2012. Bacterial persistence increases as environmental fitness decreases. Microbial Biotechnol **5:**509-522.
17. **Kim J-S, Chowdhury N, Yamasaki R, Wood TK.** 2018. Viable But Non-Culturable and Persistence Describe the Same Bacterial Stress State. Environ Microbiol **20:**2038-2048.
18. **Kim J-S, Yamasaki R, Song S, Zhang W, Wood TK.** 2018. Single Cell Observations Show Persister Cells Wake Based on Ribosome Content. Environ Microbiol **20:**2085-2098.
19. **Korch SB, Henderson TA, Hill TM.** 2003. Characterization of the *hipA7* allele of *Escherichia coli* and evidence that high persistence is governed by (p)ppGpp synthesis. Molecular Microbiology **50:**1199-1213.
20. **Kwan BW, Chowdhury N, Wood TK.** 2015. Combatting Bacterial Infections by Killing Persister Cells with Mitomycin C. Environmental Microbiology **17** 4406–4414.

21. **Kwan BW, Valenta JA, Benedik MJ, Wood TK.** 2013. Arrested protein synthesis increases persister-like cell formation. Antimicrob Agents Chemother **57:**1468-1473.

22. **Lau RK, Enustun E, Gu Y, Nguyen JV, Corbett KD.** A conserved signaling pathway activates bacterial CBASS immune signaling in response to DNA damage. The EMBO Journal **n/a:**e111540.

23. **Leavitt A, Yirmiya E, Amitai G, Lu A, Garb J, Herbst E, Morehouse BR, Hobbs SJ, Antine SP, Sun Z-YJ, Kranzusch PJ, Sorek R.** 2022. Viruses inhibit TIR gcADPR signaling to overcome bacterial defense. Nature.

24. **LeRoux M, Laub MT.** 2022. Toxin-Antitoxin Systems as Phage Defense Elements. Annual Review of Microbiology **76:**21-43.

25. **LeRoux M, Srikant S, Teodoro GIC, Zhang T, Littlehale ML, Doron S, Badiee M, Leung AKL, Sorek R, Laub MT.** 2022. The DarTG toxin-antitoxin system provides phage defence by ADP-ribosylating viral DNA. Nature Microbiology **7:**1028-1040.

26. **Levinthal C, Visconti N.** 1953. Growth and Recombination in Bacterial Viruses. Genetics **38:**500-511.

27. **Lewis K.** 2007. Persister cells, dormancy, and infectious disease. Nat. Rev. Microbiol. **5:**48-56.

28. **Lewis K.** 2008. Multidrug tolerance of biofilms and persister cells. Curr. Top. Microbiol. Immunol. **322:**107-131.

29. **Lopatina A, Tal N, Sorek R.** 2020. Abortive Infection: Bacterial Suicide as an Antiviral Immune Strategy. Annual Review of Virology **7:**371-384.

30. **Makarova KS, Anantharaman V, Aravind L, Koonin EV.** 2012. Live virus-free or die: coupling of antivirus immunity and programmed suicide or dormancy in prokaryotes. Biology Direct **7:**40.

31. **Meeske AJ, Nakandakari-Higa S, Marraffini LA.** 2019. Cas13-induced cellular dormancy prevents the rise of CRISPR-resistant bacteriophage. Nature **570:**241-245.

32. **Pecota DC, Wood TK.** 1996. Exclusion of T4 Phage by the *hok/sok* Killer Locus from Plasmid R1. J Bacteriol **178:**2044-2050.

33. **Pfaffl MW.** 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res. **29**.

34. **Pu Y, Li Y, Jin X, Tian T, Ma Q, Zhao Z, Lin S-y, Chen Z, Li B, Yao G, Leake MC, Lo C-J, Bai F.** 2019. ATP-Dependent Dynamic Protein Aggregation Regulates Bacterial Dormancy Depth Critical for Antibiotic Tolerance. Molecular Cell **73:**143-156.

35. **Ren D, Bedzyk LA, Thomas SM, Ye RW, Wood TK.** 2004. Gene expression in *Escherichia coli* biofilms. Appl Microbiol Biotechnol **64:**515-524.

36. **Rousset F, Yirmiya E, Nesher S, Brandis A, Mehlman T, Itkin M, Malitsky S, Millman A, Melamed S, Sorek R.** 2023. A conserved family of immune effectors cleaves cellular ATP upon viral infection. bioRxiv:2023.2001.2024.525353.

37. **Şimşek E, Kim M.** 2019. Power-law tail in lag time distribution underlies bacterial persistence. Proceedings of the National Academy of Sciences **116:**17635-17640.

38. **Song S, Gong T, Yamasaki R, Kim JS, Wood TK.** 2019. Identification of a potent indigoid persister antimicrobial by screening dormant cells. Biotechnol Bioeng **116:**2263-2274.

39. **Song S, Kim J-S, Yamasaki R, Oh S, Benedik MJ, Wood TK.** 2021. *Escherichia coli* cryptic prophages sense nutrients to influence persister cell resuscitation. Environ Microbiol **23:**7245-7254.

40. **Song S, Wood TK.** 2020. Persister Cells Resuscitate via Ribosome Modification by 23S rRNA Pseudouridine Synthase RluD. Environ. Microbiol. **22:**850–857.

41. **Song S, Wood TK.** 2020. ppGpp Ribosome Dimerization Model for Bacterial Persister Formation and Resuscitation. Biochem. Biophys. Res. Com. **523:**281-286.

42. **Song S, Wood TK.** 2020. A Primary Physiological Role of Toxin/Antitoxin Systems Is Phage Inhibition. Front. Microbiol. **11:**1895.

43. **Tesson F, Hervé A, Mordret E, Touchon M, d'Humières C, Cury J, Bernheim A.** 2022. Systematic and quantitative view of the antiviral arsenal of prokaryotes. Nature Communications **13:**2561.

44. **van Beljouw SPB, Sanders J, Rodríguez-Molina A, Brouns SJJ.** 2022. RNA-targeting CRISPR–Cas systems. Nature Reviews Microbiology.

45. **Vassallo CN, Doering CR, Littlehale ML, Teodoro GIC, Laub MT.** 2022. A functional selection reveals previously undetected anti-phage defence systems in the *E. coli* pangenome. Nature Microbiology **7:**1568-1579.

46. **Wang X, Lord DM, Cheng H-Y, Osbourne DO, Hong SH, Sanchez-Torres V, Quiroga C, Zheng K, Herrmann T, Peti W, Benedik MJ, Page R, Wood TK.** 2012. A Novel Type V TA System Where mRNA for Toxin GhoT is Cleaved by Antitoxin GhoS. Nat. Chem. Biol. **8:**855-861.

47. **Williams MC, Reker AE, Margolis SR, Liao J, Wiedmann M, Rojas ER, Meeske AJ.** 2023. Restriction endonuclease cleavage of phage DNA enables resuscitation from Cas13-induced bacterial dormancy. Nature Microbiology.

48. **Windels EM, Ben Meriem Z, Zahir T, Verstrepen KJ, Hersen P, Van den Bergh B, Michiels J.** 2019. Enrichment of persisters enabled by a ß-lactam-induced filamentation method reveals their stochastic single-cell awakening. Communications Biology **2:**426.

49. **Wood TK, Knabel SJ, Kwan BW.** 2013. Bacterial persister cell formation and dormancy. Appl. Environ. Microbiol. **79:**7116–7121.

50. **Yamasaki R, Song S, Benedik MJ, Wood TK.** 2020. Persister Cells Resuscitate Using Membrane Sensors that Activate Chemotaxis, Lower cAMP Levels, and Revive Ribosomes. iScience **23:**100792.

<div align="center">**Figure Legends**</div>

Fig. 1. Heterogenous single-cell resuscitation after phage attack. Representative images (from seven independent cultures) of resuscitation of cells producing MqsR/MqsA/MqsC (**A**) and exponential cells (**B**) after 1 h as determined with light microscopy (Zeiss Axio Scope.A1) using LB agarose gel pads. The persister cells were generated by the addition of T2 phage at 0.1 MOI for 1 hour. Cells with the empty plasmid are not shown due to the cellular debris that stems from complete eradication by T2 phage. Black arrows indicate cells with immediate waking, yellow arrows indicate cells with delayed waking, red arrows indicate cells that wake then die, and blue arrows indicate cells that elongate. Black arrows were not added to exponential cultures pictures because all of them divide. Data for percentages are shown in **Table S5**.

Fig. 2. Schematic illustrating phage inhibition systems invoke dormancy. Our results indicate that after phage attack, bacteria become persister cells (i.e., dormant), through RNase decay via toxin MqsR of the MqsR/MqsA/MqsC tripartite toxin/antitoxin system. During this period of inactivity, restriction/modification systems, which do not require cell resources, degrade phage DNA. Upon favorable conditions (nutrients provided), the persister cells resuscitate, having survived the phage attack.

0h   1h

Figure 1

Figure 2

Supporting Information

Toxin/Antitoxin Systems Induce Persistence and Work in Concert with Restriction/Modification Systems to Inhibit Phage

Laura Fernández-García[1,2], Sooyeon Song[1,3,4], Joy Kirigo[1], María Tomás[2], and Thomas K. Wood[1*]

[1]Department of Chemical Engineering, Pennsylvania State University,
University Park, Pennsylvania, 16802-4400, USA

[2]Microbiology Translational and Multidisciplinary (MicroTM)-Research Institute Biomedical A Coruña (INIBIC) and Microbiology Department of Hospital A Coruña (CHUAC); University of A Coruña (UDC), A Coruña, Spain

Departments of Animal Science[3] and Agricultural Convergence Technology[4], Jeonbuk National University, 587 Baekje-Daero, Deojin-Gu, Jeonju-Si, Jellabuk-Do, 54896, South Korea

*For correspondence. E-mail tuw14@psu.edu

Tel. (+)1 814-863-4811; Fax (1) 814-865-7846

Table S1. **MqsR/MqsA/MqsC inhibits T2 phage.** Cells were contacted with T2 phage at 0.1 MOI for 1 h and with antibiotics (10 MIC) for 3 h. Bold results indicate million-fold difference for inhibiting T2 phage. 'Std Dev' indicates standard deviation.

Strain	Condition	CFU/ml average	Std Dev	Survival %
MG1655/pCV1	No phage	2×10^9	5×10^8	100%
	Phage T2	**3.4×10^2**	8×10^1	100%
	No phage + 100 µg/ml ampicillin	1×10^8	1×10^8	7%
	No phage + 5 µg/ml ciprofloxacin	2.4×10^6	1×10^6	0.2%
	Phage T2 + 100 µg/ml ampicillin	0.00	0.00	0%
	Phage T2 + 5 µg/ml ciprofloxacin	0.00	0.00	0%
MG1655/pCV1-*mqsRAC*	No phage	1.5×10^9	2×10^8	100%
	Phage T2	**4×10^8**	1×10^8	100%
	No phage + 100 µg/ml ampicillin	2.5×10^8	0.7×10^8	16%
	No phage + 5 µg/ml ciprofloxacin	3×10^6	1×10^6	0.2%
	Phage T2 + 100 µg/ml ampicillin	4×10^7	1×10^7	8%
	Phage T2 + 5 µg/ml ciprofloxacin	1×10^6	4×10^5	0.3%

Table S2. Percentage of persister cell waking after treatment with T2 phage (0.1 MOI) for 1 h.

Sample	No of cells	% of waking in 0.5 h (No)	% of waking in 1 h (No)	% of lagged waking (No)	% Elongated cells (No)	% of dead (No)	Unmodified cells (No)
1	33	-	88 % (29/33)	9 % (3/33)	9% (3/33)	1 % (1/33)	
2	25	20 % (5/25)	24 % (6/25)	-	16% (4/25)	12 % (3/25)	40 % (10/25)
3	27	41 % (11/27)	37 % (10/27)	-	11% (3/27)	7 % (2/27)	7 % (2/27)
4	65	58 % (38/65)	28 % (18/65)	-	20% (13/65)	1 % (1/65)	8 % (5/65)
5	23	26 % (6/23)	22 % (5/23)	30 % (7/23)	35% (8/23)	4 % (1/23)	17 % (4/23)

Table S3. Phage attack does not induce ribosome inactivation proteins. Fold changes for the wild-type producing MqsR/MqsA/MqsC after 15 min of T2 attack are relative to the wild-type strain with the empty plasmid with 15 min of T2 attack as determined by qRT-PCR. Cycle numbers (C_t) are indicated for each sample including that for the target genes as well as that of the house-keeping gene, *rrsG,* which was used to normalize the data. Fold were calculated as described earlier[33]:

$$2\text{\textasciicircum}-(C_{t\ target\_cas2+}-C_{t\ purM\_cas2+})/2\text{\textasciicircum}-(C_{t\ target\_cas2-}-C_{t\ purM\_cas2-})$$

Gene	*rrsG*		*rmf*		*hpf*		*raiA*	
Strain	MG1655 /pCV1	*MG1655 /pCV1- mqsRAC*	MG1655 /pCV1	*MG1655 /pCV1- mqsRAC*	MG1655 /pCV1	*MG1655 /pCV1- mqsRAC*	MG1655 /pCV1	*MG1655 /pCV1- mqsRAC*
C_T	4.6 ± 0.5	4.48 ± 0.05	16.9 ± 0.6	17.01 ± 0.2	15.3 ± 0.6	15.2 ± 0.1	12.8 ± 0.2	14.7 ± 0.1
ΔC_T			12.3 ± 0.1	12.6 ± 0.2	10.7 ± 0.2	10.7 ± 0.1	8.2 ± 0.3	10.2 ± 0.2
$\Delta\Delta C_T$			0.3 ± 0.1		0.02 ± 0.02		2.0 ± 0.1	
fold			1.2		1.05		4.0	

Fernández-García et al., Pennsylvania State University   4

Table S4. MqsR/MqsA/MqsC inhibits T2 phage through cooperation with restriction/modification systems. (**A**) Survival after 30 min of T2 phage infection at 0.1 MOI. (**B**) BW25113 and BW25113 *mcrB* cells producing MqsR/MqsA/MqsC from its native promoter were contacted with T2 phage at 0.1 MOI for 1 h and with antibiotics (10 MIC) for 3 h. 'Std Dev' indicates standard deviation.

A

Strain	CFU/ml average	Std Dev
BW25113	1.40E+06	3.33E+05
BW25113 *Δmrr*	7.75E+04	2.13E+04
BW25113 *Δhsd*	3.65E+05	1.77E+05
BW25113 *ΔmcrB*	1.67E+02	1.11E+02

B

Strain	Condition	CFU/ml average	Std Dev
wild-type/pCV1-*mqsRAC*	No phage	1.7×10^9	3×10^8
	Phage T2	3.8×10^8	3×10^7
	No phage + 100 µg/ml Ampicillin	2×10^8	3×10^7
	Phage T2 + 100 µg/ml Ampicillin	**1.7×10^6**	5×10^5
ΔmcrB/pCV1-*mqsRAC*	No phage	7.6×10^8	2×10^8
	Phage T2	1.3×10^7	6×10^6
	No phage + 100 µg/ml Ampicillin	1.9×10^7	5×10^6
	Phage T2 + 100 µg/ml Ampicillin	**2×10^5**	2×10^5

Table S5. *E. coli* **bacterial strains and plasmids utilized.** Cm^R is chloramphenicol resistance, and Kan^R is kanamycin resistance.

Strains and Plasmids	Features	Source
Strains		
MG1655	F⁻ lambda- *ilvG- rfb*-50 *rph*-1	3
BW25113	*rrnB3 ΔlacZ4787 hsdR514 Δ(araBAD)567 Δ(rhaBAD)568 rph-1*	2
BW25113 *mcrB*	BW25113 *mcrB* Ω Kan^R	2
BW25113 *hsdR*	BW25113 *hsdR* Ω Kan^R	2
BW25113 *mrr*	BW25113 *mrr* Ω Kan^R	2
Plasmids		
pCV1	Cm^R	45
pCV1-*mqsRAC*	P_{mqsRAC}::*mqsRAC;* Cm^R	45

Table S6. qRT-PCR primers

Gene	Primer pair	Length	Sequence (5'→3')	T$_m$ (ºC)	Product size
rrsG *(housekeeping)*	*rrsG* Fow	20 bp	TATTGCACAATGGGCGCAAG	56.9	234 bp
	rrsG Rev	20 bp	ACTTAACAAACCGCCTGCGT	50.0	
raiA	*raiA* Fow	25 bp	CCAGCAAACAAATGGAAATTACTCC	61.3	122 bp
	raiA Rev	22 bp	TTGTGGCTCTTTGGACAGAATG	60.8	
rmf	*rmf* Fow	24 bp	AACGGGCACATCAACGTGGTTATC	64.6	100 bp
	rmf Rev	20 bp	CCTCCCAGCCATTGTGACCT	64.5	
hpf	*hpf* Fow	20 bp	TCGAGATCACCGAGGCACTG	64.5	119 bp
	gpf Rev	24 bp	GGTGTGGGTGACTTTCTCCACTTT	64.6	
hflX	*hflX* Fow	22 bp	TTCCGCCGGTGTCGAAGCATTG	66.4	124 bp
	hflX Rev	20 bp	CGAAGCACCCGTCGCTTTGA	64.5	

Supplemental Figure S1. Growth of four independent colonies of each strain in LB at 37ºC. 'MG1655' is the wild-type host (red), 'MG1655/pCV1' is the host with the empty plasmid (negative control) (orange) and 'MG1655/pCV1-mqsRAC' is the host with the MqsR/MqsA/MqsC tripartite toxin/antitoxin system (blue).

Supplemental Figure S2. Metabolic activity via flow cytometry. MG1655/pCV1-mqsRAC was grown to a turbidity of 0.5 at 600 nm and T2 phage was added (MOI 0.1) for 1 h, then stained for metabolic activity using the BacLight™ RedoxSensor™ Green Vitality Kit ('persister' cells) and compared to exponentially-growing cells stained with the same kit ('Exponential culture'). RedoxSensor Green measures the cellular redox state. 'Unstained' indicates results with an overnight, unstained sample. '%Max' indicates the relative number of cells (100,000 cells total), and 'FITC-A' is indicative of the intensity of the RedoxSensor Green fluorescence. One representative image of two independent cultures shown.