# Specialized cingulo-opercular network activation in judging ambiguity of facial expressions in a slow reveal fMRI paradigm

Jordan E. Pierce[1]*, Nathan M. Petro[2,3], Elizabeth Clancy[4], Caterina Gratton[5], Steven E. Petersen[6], Maital Neta[1]

[1]Center for Brain, Biology, and Behavior, University of Nebraska-Lincoln, Lincoln, NE, USA

[2]Institute for Human Neuroscience, Boys Town National Research Hospital, Boys Town, NE, USA

[3]Center for Pediatric Brain Health, Boys Town National Research Hospital, Boys Town, NE, USA

[4]University of Guelph, Guelph, Ontario, Canada

[5]Florida State University, Tallahassee, FL, USA

[6]Washington University in St. Louis, St. Louis, MO, USA

*corresponding author, email: jpierce14@unl.edu

address: B76 East Stadium, Lincoln, NE 68588

**Abstract**

Certain facial expressions express emotional valence clearly, while other expressions such as surprise are ambiguous and can be interpreted as expressing either positive or negative valence. In two studies, we presented a valence bias task in which masked clear and ambiguous emotional expressions were slowly revealed over several seconds. This design allows for investigation of the decision-making mechanisms supporting ambiguity processing as they unfold over the course of a trial. In the main study, the task was administered to 32 young adults in the fMRI environment and BOLD time courses were extracted from regions of interest in three cortical networks previously associated with task control, decision making, and performance reporting. In a follow-up study, the task was administered to a larger, online sample (n = 81) using a more extended slow reveal design with additional unmasking frames. Positive judgments of surprised faces were accompanied by stronger activation in the cingulo-opercular network and slower response times. These results support the initial negativity hypothesis, which posits that the default response to ambiguity is negative, by demonstrating that making positive judgments is an effortful, controlled process that varies according to individual biases.

**Keywords:** emotion, ambiguity, facial expressions, fMRI, valence bias

**1. Introduction**

Emotional expressions allow an individual to non-verbally communicate to others how they are feeling in the moment. Certain facial expressions express positive or negative emotional valence relatively clearly, such as a happy or fearful expression, and quickly convey important social cues about potential rewards or threats in the environment. Other expressions, however, may convey a signal that is ambiguous in valence, such as a surprised expression, which could be interpreted as either positive or negative depending on the context (Kim et al., 2004; Neta et al., 2011; Neta & Kim, 2022). Thus, if individuals are asked to evaluate ambiguous surprised expressions without context, valence judgments range from consistently positive appraisals to a mix of positive and negative to consistently negative appraisals in a trait-like manner that is termed "valence bias" (Harp et al., 2022; Kim et al., 2003; Neta et al., 2021).

Much of the work on this topic over the last 15 years has come to focus on determining the underlying processes that support valence bias. This work has demonstrated that the initial response to ambiguity tends to be negative, and that positive judgments may require more time to implicitly reappraise the stimulus and overcome the initial negativity (Neta et al., 2021; Neta & Tong, 2016; Neta & Whalen, 2010). For example, the presentation of low spatial frequency images of facial expressions (that can be quickly processed by the visual system) biased responses to ambiguous expressions towards negative judgments (Neta & Whalen, 2010). Another study demonstrated that when participants were encouraged to delay their response to ambiguity, their judgments shifted towards positivity (Neta & Tong, 2016). Finally, by using a mouse-tracking approach, studies revealed that only positive judgments of ambiguous stimuli are characterized by an initial motoric attraction to the competing (negative) response option (Brown et al., 2017; Neta et al., 2021). Notably, the additional processing time and response competition

for positive judgments is thought to be related to a regulatory process that helps to override the initial negativity (Neta, in press; Neta et al., 2022; Petro et al., 2018).

These findings have emerged alongside neuroimaging work which has explored the neural mechanisms underlying responses to ambiguity more broadly. For example, the decision-making process that might be particularly key for resolving ambiguity has been explored using an extended perceptual task (Gratton et al., 2017; Neta et al., 2017; Ploran et al., 2007). In this "slow reveal" task paradigm, individuals must identify an object that is unmasked slowly over several seconds or report a word that is revealed one letter at a time. An analysis of several such functional magnetic resonance imaging (fMRI) studies indicated that the cingulo-opercular (CO) and fronto-parietal (FP) control networks of the brain exhibited distinct temporal characteristics that were differentiated by the onset (early vs. late) and duration (extended vs. transient) of the hemodynamic response (Gratton et al., 2017, see also Coste & Kleinschmidt, 2016; Dosenbach et al., 2008). Specifically, early activation in the left FP network supported evidence accumulation and increased throughout the trial until a response was made, while later extended activation in the right FP network reflected processing after the response occurred. Finally, late, transient activation in the CO network corresponded to performance reporting at the end of a trial (Gratton et al., 2017).

The performance reporting function of the CO network – which includes regions such as the dorsal anterior cingulate cortex, medial prefrontal cortex, anterior insula, and frontal operculum – was investigated further using the same slow reveal paradigm for object identification with the addition of an ambiguity condition (Neta et al., 2017). Performance accuracy and brain activation were measured on trials showing objects with unique, easily identifiable outlines (e.g., butterfly) or objects with ambiguous, easily mistaken outlines (e.g.,

toothbrush) that were gradually unmasked. Ambiguous stimuli and error trials (self-reported misidentifications) yielded interactive effects in the late response of the CO network, suggesting overlapping but distinct information processing for these functions as related to performance reporting after object recognition (see also Neta et al., 2014). This is consistent with work demonstrating that the CO shows increased activation when judging the valence of ambiguous emotional stimuli (Neta et al., 2013).

In the current study, the slow reveal paradigm was combined with the valence bias task to probe the underlying process that supports emotional valence judgments. Participants were shown non-ambiguous – happy, angry, and fearful – and ambiguous – surprised – expressions. The expressions were slowly unmasked and participants were instructed to categorize each face as expressing positive or negative valence. Perceptual uncertainty may impact recognition of certain facial expressions as a function of the similarity of morphological features, potentially resulting in slower or inaccurate judgments of their emotional valence. Importantly, the ambiguity inherent in a surprised expression was predicted to contribute to further uncertainty – slowing valence responses for this condition. These slower responses then might allow participants the opportunity to engage in a regulatory process that recruits greater task control network activation and yields more positive surprised valence ratings. Given the prevalent decision-making role for CO in response to ambiguity across many different domains of scientific inquiry (Neta et al., 2013, 2014; Poudel et al., 2020; Sterzer et al., 2002; Thompson-Schill et al., 1997), the CO network was of particular interest here in putatively supporting judgments of facial expressions and, more specifically, supporting a more positive valence bias.

Finally, we attempted to replicate and extend the behavioral findings from the MRI study in a follow-up online study that used a more extended unmasking phase. Specifically, an

increased number of frames during the slow reveal was intended to provide greater sensitivity to differences in the relative timing of the valence decision-making process for ambiguous expressions. Given that positive judgments of surprised faces are often slower and putatively require overriding the default negativity, we predicted that when positive judgments occurred, they would be made later in the reveal process than negative judgments.

## 2. Methods

### 2.1 Participants

For the MRI study, thirty-seven participants were recruited from the Lincoln community via publicly posted flyers. Participants reported no history of neurological or psychiatric disorders and no use of psychotropic medications, were right-handed, had normal or corrected-to-normal vision, and were naive to the purpose of the study. One participant could not complete the MRI scan due to an unremovable piercing and four participants data were excluded due to technical issues. The final sample included 32 participants (20 female/12 male, mean age = 20.6 years (SD=1.7)). All participants provided written informed consent and all procedures were approved by the Institutional Review Board at the University of Nebraska-Lincoln.

### 2.2 Procedure

Participants received verbal task instructions prior to entering the MRI scanner, then were positioned on their back with padding to secure their head in a comfortable, static position where they could view the experimental paradigm display via a mirror attached to the head coil. Responses were collected via an MRI-compatible button box with the right index and middle finger (with counterbalanced assignment to response choices). At the start of the functional scans, task instructions were presented on the screen and participants were given a verbal

reminder. Participants were instructed to categorize each face as either positive or negative twice during the trial: once as the picture was being unmasked ("noise" condition) – as soon as they had a reasonably confident judgment – and again when the picture was fully revealed ("reveal" condition). They were told that the two responses could be the same or different. The participants then completed three practice trials containing only angry and happy faces that were not used in the main experiment. Following the practice trials, the participants were given the opportunity to ask for clarification before beginning the task.

*2.3 Task Design*

On each trial, a grayscale face was occluded by a black mask that was slowly degraded over the course of five 2-second frames (**Figure 1**). Each trial began with a 2 sec black screen, then the black mask was degraded by randomly removing pixels following a Gaussian distribution over the five "noise" frames such that 0.75, 1.25, 3.0, 7.0, and 14.0% of the face image was visible at each successive step. Each trial ended with the presentation of the fully revealed face stimulus for 2 seconds. Thus, the duration of each trial was 14 seconds. During the inter-trial interval (ITI), a fixation dot was presented at the center of the screen for 2, 4, or 6 seconds (equally presented across trial conditions).

There was a total of 240 trials, with an equal distribution of faces displaying a happy, angry, fearful, or surprised expression (60 of each) across ten separate runs. Given that temporal context can shift responses to surprised faces (Neta et al., 2011), expressions were presented in pseudorandom order such that the surprised faces were preceded by an equal number of clearly positive and negative faces. The face stimuli were drawn from the NimStim Set of Facial Expressions (Tottenham et al., 2009), the Karolinska Directed Emotional Faces database (Goeleven et al., 2008), the Umea University Database of Facial Expressions, (Samuelsson et al.,

2012), and the Ekman Pictures of Facial Affect (Ekman, 1976). The luminance of all faces was equated using the SHINE toolbox (Willenbockel et al., 2010) for MATLAB (Mathworks Inc, Natick, MA).

**Figure 1**. Task design for the slow reveal of emotional facial expressions. In the main study, each trial lasted 14 seconds, consisting of seven 2-second frames: one black frame, five "noise" frames with a decreasing number of masking pixels, and a final "reveal" frame. In between trials, a fixation dot appeared for a variable ITI of 2, 4, or 6 seconds. Here a smiley icon is used in place of the photograph shown to participants.

*2.4 MRI Parameters*

The MRI data were collected at the University of Nebraska-Lincoln Center for Brain, Biology, & Behavior on a Siemens 3T Skyra scanner using a 32-channel head coil. Structural images were acquired using a T1-weighted 3D multi-echo MPRAGE sequence (TR = 2530 ms, TE = 1.69 ms, 4 averages, slices = 176 interleaved, voxel size = 1.0 mm³, matrix = 256 × 256, FOV = 256 mm, flip angle = 7°, total acquisition time = 6:00). Blood oxygen level-dependent

(BOLD) activity was measured using a multi-band EPI scanning sequence (TR = 1000 ms, TE = 29.8 ms, multi-band factor = 3, slices = 51 interleaved, voxel size = 2.50 mm$^3$, matrix = 84 × 84 mm, FOV = 168 mm, flip angle = 60°, 454 volumes, total acquisition time = 7:48 per run), and slices were acquired parallel to the AC-PC plane, positioned to cover the entire brain.

*2.5 Behavioral analysis*

Average valence judgments for each emotion were calculated as the number of negative responses out of the number of total trials for each participant, multiplied by 100 (i.e., percent negative). A 4 x 2 (emotion by trial phase) repeated measures ANOVA was conducted on the average judgments to compare responses for happy, angry, fearful, and surprised faces during the first "noise" phase and the second "reveal" phase. Subsequently, incorrect judgments for clearly valenced emotional expressions (happy judged as negative and angry or fearful judged as positive) were considered errors and not included in the response time analyses; for surprised faces, both positive and negative judgments were retained as separate conditions.

The average response frame (out of 5) was calculated for each of five emotion conditions (happy, angry, fearful, positive surprised, and negative surprised) for each participant. Response time (RT) was calculated for responses in both the noise phase and the reveal phase. The noise RT was calculated from the beginning of the first noise frame, giving a maximum possible value of 10 seconds (2 sec x 5 frames), while the reveal RT had a maximum possible value of 2 seconds. One-way ANOVAs by emotion (5 levels) were conducted on noise and reveal RTs. Huynh-Feldt adjusted degrees of freedom were used to account for unequal variance between conditions when indicated by a significant Mauchly's test of sphericity. All statistical analyses were conducted in SPSS version 28 (IBM, Cary, NC, USA).

*2.6 FMRI Analysis*

*2.6.1 Preprocessing*

Functional data were analyzed using the AFNI software package (Cox, 1996, 2012).

Preprocessing consisted of de-spiking of time series outliers, slice timing correction, alignment

of functional volumes to each other and the individual anatomical image, standardization to the

Talairach atlas space (Talairach & Tournoux, 1988), smoothing with a 6-mm FWHM kernel, and

scaling of each voxel to a mean of 100. Volumes with a motion shift where the Euclidean norm

of the derivative was greater than 0.3 were censored (0.45% of volumes removed across all

participants).

*2.6.2 HRF Time Courses by Emotional Expression*

Preprocessed functional data were entered into a general linear model in which individual

responses were sorted by the emotion shown and valence judgment made (see Supplemental

Material and Figure S1 for ROI analysis combined across all emotions). Thus, there were

regressors for positive judgments of happy faces, negative judgments of angry faces, negative

judgments of fearful faces, positive judgments of surprised faces, and negative judgments of

surprised faces, combined across response frames 2-5. The "TENT" function was used to

estimate the amplitude of the hemodynamic response at each TR from 0-21 seconds after

stimulus onset without assuming a predetermined shape for the HRF in each voxel. Regressors of

no interest included error trials (negative judgments of happy faces and positive judgments of

angry and fearful faces), anticipatory responses (frame 1 – when there was insufficient visual

information to make a decision), polynomials for each run (two terms), and six motion

parameters estimated during alignment (x, y, z shift/rotation).

Subsequently, the beta values at each time point (i.e., the estimated hemodynamic

response function (HRF)) were extracted from 16 regions of interest (ROIs; **Figure 2**) based on a

previous study using a slow reveal task (Gratton et al., 2017). Each ROI was defined as a 5-mm sphere centered on the peak coordinates of each region. Following the approach of Gratton et al., 2017, data from these 16 ROIs were combined into three ROI clusters: cingulo-opercular (CO), left frontoparietal (L FP), and right frontoparietal (R FP). The CO cluster consisted of four ROIs (as labelled in Gratton et al., 2017): left anterior insula/frontal operculum (AI/FO), right AI/FO, pre-supplemental motor area (SMA) 1, and pre-SMA 2. The L FP cluster consisted of eight ROIs: left inferior parietal sulcus (IPS) 1, left IPS 2, left IPS 3, left frontal, left middle frontal gyrus (MFG) 1, left MFG 2, right IPS 2, and right frontal. The R FP cluster consisted of four ROIs: right IPS 1, right inferior frontal gyrus (IFG), right MFG 1, and right MFG 2. Additionally, responses from 5 visual ROIs based on another slow reveal study (Ploran et al., 2007) were averaged together in a "Sensory" ROI cluster as a control condition in which no differences were expected by emotion (see Supplemental Table S1 for all regions and their coordinates).

To compare the responses for each emotion, beta values were entered into a linear mixed model with fixed effects of emotion (5 levels), time (11 timepoints; with linear and quadratic terms), and their interaction, with a random intercept by subject. For this analysis, only the second half of the HRF time course was examined (11-21 seconds) in order to focus on the period that contains the hemodynamic peak (see Figure 3) and exclude early time points with minimal activation across conditions. Given the behavioral differences between emotional expressions, RT from the reveal phase was also included as a covariate. Emotion and time were modeled as repeated measures with a diagonal covariance structure and Satterthwaite estimates were used for determining degrees of freedom for the fixed effects. Separate models were fit for each of the three ROI clusters of interest (CO, L FP, and R FP) and the control cluster (Sensory).

For positive surprised trials, 10 participants were removed due to a low trial count (< 4 surprised trials with a "positive" response, remaining N=22); all other emotion conditions included all participants (N=32).

**Figure 2**. Location of the ROIs extracted based on Gratton et al., 2017 and Ploran et al, 2007. Colors indicate the ROI cluster: CO (green), left FP (blue), right FP (red), and Sensory (purple). Coordinates for each ROI are provided in Supplemental Table S1.

*2.6.3 Late CO Activation and Behavior*

From the initial analysis, there was a second, late peak observed in the CO responses to surprised faces (see Fig. 3) that we predicted might reflect a response to the final reveal of the (ambiguous) face. To explore this possibility, beta values were extracted for each emotional expression for the period from 16-19 seconds in the trial, corresponding to the period 4-7 seconds after the final reveal image was displayed (i.e., when an HRF peak for the reveal image would occur). The average activation in this period was correlated with participants' average reveal RT for each emotion to determine if CO activation was dependent on RT differences.

Additionally, the activation for positive and negative surprised trials was correlated with participants' overall valence bias (percent of negative judgments) for the reveal phase to determine if CO activation was driven by extended processing for positive judgments.

*2.6.4 HRF Time Courses as a Function of Switching versus Staying*

To further examine the differences in HRF time courses by emotional expression, we conducted an exploratory analysis to compare trials on which participants made the same behavioral judgment during the noise and reveal phases of the trial and trials on which participants switched their response during the reveal phase. The latter "switch" condition included clear valence trials on which participants made an error, presumably recognized the error, and corrected their response when seeing the fully revealed face. Importantly, the "switch" condition also included ambiguous trials (which do not have an objectively accurate response) on which participants changed their response at the reveal phase, perhaps because they decided that their initial judgment was inappropriate. Conversely, the "stay" condition included only trials on which participants did not change their valence judgment, presumably because they were sufficiently confident in their initial response and did not consider it an error.

For this analysis, clear valence errors (e.g., negative judgment of a happy face) were excluded for the stay condition and clear valence correct trials (e.g., positive judgment of a happy face) were excluded for the switch condition given an insufficient number of trials (< two trials for any participant). Additionally, within each of the remaining conditions, individual participants were excluded if they had fewer than three trials in a particular response condition. This requirement led to a different number of participants being included across conditions (stay condition: positive happy/negative angry/negative fearful/negative surprised: N=32; positive surprised: N=17; switch condition: negative happy: N=15; positive angry: N=24; positive fearful:

N=31; positive surprised: N=14; negative surprised: N=19). Nonetheless, this approach provides the best estimate of the HRFs using the maximum amount of available data for this initial exploratory analysis of potential factors contributing to the HRF differences by emotion.

As with the prior analysis by emotional expression, a linear mixed model was fit with fixed effects of emotion (5 levels), time (11 timepoints; with linear and quadratic terms), and their interaction, with a random intercept by subject. Emotion and time were modeled as repeated measures with a diagonal covariance structure and Satterthwaite estimates were used for determining degrees of freedom for the fixed effects. Separate models were fit for switch and stay conditions in the CO cluster (models for the L FP and R FP cluster are reported in the Supplemental Figure S2 and Tables S10-13).

### 2.6.5 Emotional Expression HRF Peak by Response Frame

Finally, to explore the effects of *response frame* on HRF timing and shape, a general linear model was fit according to the frame during which an individual participant made their judgment for each trial, as well as the emotional expression and response. Thus, there were regressors for positive happy, negative angry, negative fearful, positive surprised, and negative surprised trials for each response frame 2-5. Again, the "TENT" function was used to estimate the amplitude of the HRF at each TR from 0-21 seconds after stimulus onset. Regressors of no interest included error trials (negative happy, positive angry, and positive fearful responses), anticipatory responses (frame 1), polynomials for each run (two terms) and six motion parameters estimated during alignment (x, y, z shift/rotation). It should be noted that participants made most behavioral responses in frames 3 and 4 (~40% each) and fewer responses in frames 2 and 5 (~10% each), potentially impacting the stability of the HRF curve estimates across frames. All participants' data were included for any condition in which a behavioral response was made.

Beta values were extracted for each of the 16 ROIs and combined into the three ROI clusters of interest, focusing on the CO cluster (results for the L FP and R FP clusters are reported in the Supplemental Material). The average HRF for each emotion and frame was then upsampled from 22 time points to 2,200 using linear interpolation to aid visualization and minimize any impact of outlying time points, and peaks were identified using the "findpeaks" function in MATLAB. Peaks are reported that occurred at least five seconds into the trial (to allow the HRF time to develop), had an amplitude greater than 0.05 (to focus on positive activations), and were at least three seconds from another peak (to identify temporally distinct events).

Data are available online at https://osf.io/g39ke/.

## 3. Results

### 3.1 Behavior

Responses for valence judgments from the noise and reveal phases of the task were averaged separately for each of the four emotional expressions, resulting in a mean percent negative rating (**Table 1**). A 4 x 2 (emotion by trial phase) repeated measures ANOVA indicated significant effects of emotion ($F(2.22, 68.78) = 1785.13$, $p < .001$, $\eta^2 = .983$), phase ($F(1, 31) = 20.59$, $p < .001$, $\eta^2 = .399$), and their interaction ($F(2.13, 66.20) = 45.15$, $p < .001$, $\eta^2 = .593$). As expected, happy faces were judged as mostly positive while angry and fearful faces were judged as mostly negative. Additionally, as predicted, there were fewer errors (e.g., a happy face judged as negative) made during the reveal phase than the noise phase for each of the clearly valenced expressions ($t$'s $> 4.2$, $p$'s $< .001$). Ratings for surprised faces, on the other hand, did not significantly differ between the noise and reveal phases ($t(31) = 0.87$, $p > .05$) and were judged

as somewhat negative during both phases, with more variability (i.e., individual valence bias) during the reveal phase than for the clear valence emotions.

**Table 1**. Average Percent Negative Ratings by Emotion.

| Emotion | Noise | Reveal |
|---|---|---|
| *Main Study* | | |
| Happy | 6.70 (8.64) | 0.56 (1.08) |
| Angry | 88.79 (8.51) | 99.38 (0.98) |
| Fearful | 84.83 (9.98) | 98.14 (2.85) |
| Surprised | 87.89 (9.18) | 86.68 (11.41) |
| *Follow-up Study* | | |
| Happy | 12.15 (14.38) | 0.98 (1.98) |
| Angry | 77.65 (17.32) | 98.36 (3.01) |
| Fearful | 69.77 (17.04) | 96.39 (5.45) |
| Surprised | 76.64 (15.62) | 81.42 (17.61) |

Values are given as mean (SD).

Next, the noise frame (1-5) during which responses occurred was compared for correct happy, angry, and fearful face trials as well as separately for surprised face trials that were judged as positive versus negative. Descriptively, most responses for all emotions occurred during frames 3 and 4; happy faces that were correctly judged as positive and surprised faces that were judged as negative tended to have more early responses (frame 3) compared to correctly judged angry and fearful faces which had later responses (frame 4), as can be seen in the average response frame in the noise phase (**Table 2**). For a statistical test of these effects, a one-way ANOVA was conducted on reaction times (RTs) for the noise phase and resulted in a main effect of emotion ($F(1.88, 56.43) = 10.75$, $p < .001$, $\eta^2 = .264$). Specifically, participants made slower responses for angry and fearful face trials and faster responses for happy and negative surprised

face trials. For the reveal phase of the task, however, RTs showed a different pattern by emotion ($F$(2.31, 60.03) = 90.29, $p < .001$, $\eta^2 = .776$), with positive surprised trials resulting in the slowest responses, while happy trial responses were still faster than all other emotions (**Tables 2/S2**).

**Table 2**. Average Response Time by Emotion.

| Emotion | Average Frame | Noise RT (ms) | Reveal RT (ms) |
|---|---|---|---|
| *Main Study* | | | |
| Happy | 3.25 (0.44) | 5417 (868) | 625 (118) |
| Angry | 3.61 (0.45) | 6124 (882) | 697 (120) |
| Fearful | 3.50 (0.48) | 5918 (949) | 699 (137) |
| Positive Surprised | 3.40 (0.69) | 5792 (1376) | 1018 (216) |
| Negative Surprised | 3.31 (0.51) | 5559 (1019) | 746 (168) |
| *Follow-up Study* | | | |
| Happy | 4.82 (0.79) | 8515 (1588) | 742 (175) |
| Angry | 5.15 (0.74) | 9219 (1522) | 791 (202) |
| Fearful | 4.96 (0.72) | 8871 (1453) | 824 (191) |
| Positive Surprised | 4.82 (0.95) | 8695 (1917) | 1079 (259) |
| Negative Surprised | 4.81 (0.75) | 8637 (1481) | 831 (204) |

Values are given as mean (SD). The main study included five noise frames and the follow-up study included seven noise frames.

*3.2 FMRI*

*3.2.1 HRF Time Courses by Emotional Expression*

The fMRI analysis of the HRF responses in the three ROI clusters of interest (as well as the Sensory control cluster) focused on differences as a function of emotional expression. For each ROI cluster separately, a linear mixed model was fit with fixed effects of emotion (5

levels), time (11 timepoints; with linear and quadratic terms) and their interaction, with reveal

RT as a covariate and a random intercept by subject. Time points were included from the second

half of the trial (11-21 seconds) when the BOLD response peaked, and emotions included clear

correct trials (positive happy, negative angry, negative fearful) and both positive and negative

surprised trials. See Supplemental Tables S4-S7 for full model results.

For the CO cluster, there were significant effects of emotion ($F$(4, 306.27) = 7.92, $p <$

.001), time point (quadratic) ($F$(1, 696.42) = 49.37, $p < .001$), RT ($F$(1, 925.87) = 65.66, $p <$

.001), and the interaction of emotion and time ($F$(4, 372.26) = 19.57, $p < .001$), and a non-

significant effect of time point (linear) ($F$(1, 679.04) = 3.35, $p = .068$). The emotion effect and

the interaction were driven by a stronger response for positive surprised trials from 13-21

seconds. Negative surprised trials also had a larger response than happy, angry, and fearful trials

from 16 seconds until the end of the trial (**Figure 3A**).

For the L FP cluster, there were significant effects of emotion ($F$(4, 300.36) = 3.63, $p =$

.007), time point (linear) ($F$(1, 640.71) = 17.14, $p < .001$), time point (quadratic) ($F$(1, 713.64) =

77.13, $p < .001$), RT ($F$(1, 904.89) = 16.90, $p < .001$) and the interaction of emotion and time

($F$(4, 350.84) = 5.10, $p < .001$). Similar to the CO cluster, positive surprised trials had a stronger

response than the other emotions from 15-21 seconds (**Figure 3B**).

For the R FP cluster, there were significant effects of emotion (F(4, 281.99) = 15.62, p <

.001), time point (linear) (F(1, 630.80) = 431.91, p < .001), time point (quadratic) (F(1, 716.23) =

507.72, $p < .001$), RT ($F$(1, 862.81) = 57.92, $p < .001$), and the interaction of emotion and time

(F(4, 353.78) = 11.91, p < .001). The interaction was again driven by a stronger, more extended

response for positive surprised trials compared to all other emotions beginning at 15 seconds

(**Figure 3C**).

As a control, responses from five visual ROIs from Ploran et al. (2007) were averaged together in a "Sensory" ROI cluster. This cluster showed significant effects of time point (linear) $(F(1, 594.89) = 742.64, p < .001)$, time point (quadratic) $(F(1, 723.89) = 944.14, p < .001)$, and RT $(F(1, 681.56) = 7.85, p = .005)$ with non-significant effects of emotion $(F(4, 306.11) = 0.60, p = .660)$ and the interaction of emotion and time $(F(4, 459.01) = 0.34, p = .850)$. All faces elicited a strong response in these regions that did not differ by emotional expression (**Figure 3D**).

**Figure 3**. Estimated HRF time courses by emotional expression in each of the ROI clusters. Emotions included correct clear valence trials (happy judged as positive, angry/fearful judged as negative) and ambiguous (surprised) trials judged as either negative or positive. In all three

clusters of interest, positive surprised trials yielded the strongest response in the last half of the HRF; there were no effects of emotion in the control (Sensory) cluster. The gray bar represents the timing of each frame of the trial from 0 = black screen to 5 = final noise frame and R = fully revealed image.

### 3.2.2 Late CO Activation and Behavior

Given the a priori interest in the CO network and evidence of a second, late peak for some conditions, the average BOLD activation was extracted for each emotion for the period from 16-19 seconds, which corresponds to 4-7 seconds after the fully revealed image was presented. Accordingly, we predicted that activation in this period would relate to behavioral responses for the reveal phase. A one-way ANOVA indicated that positive surprised trials had the strongest activation in this window (**Figure 4**), followed by negative surprised, while the clear valence correct trials (happy, angry, fearful) showed a relatively weak response ($F(3.24, 67.95) = 37.07$, $p < .001$, $\eta^2 = .638$). Clear valence error trial activation was also extracted and did not significantly differ from positive (Bonferroni-corrected $p = 1.00$) or negative ($p = .064$) surprised trial activation.

The correlation between each emotion's late peak CO activation and its reveal RT was significant for fearful ($\rho = .432$, $p = .014$, 95% CI [.088, .684]) and negative surprised trials ($\rho = .400$, $p = .023$, 95% CI [.049, .663]), marginal for happy ($\rho = .315$, $p = .079$, 95% CI [-.048, .605]), and non-significant for angry ($\rho = .206$, $p = .258$, 95% CI [-.164, .525]) and positive surprised trials ($\rho = .132$, $p = .567$, 95% CI [-.330, .543]). This suggests that slower RTs may have contributed to some increased CO activation, but were not driving the marked increase for positive surprised trials. Additionally, the correlation between late peak CO activation and reveal valence bias was significant for both positive surprised ($\rho = -.471$, $p = .027$, 95% CI [-.750, -.048]) and negative surprised trials ($\rho = -.635$, $p < .001$, 95% CI [-.809, -.359]), indicating that

individuals who made more positive judgments during the reveal phase had greater activation in the CO network regardless of their response for a given trial during the noise phase. This finding may suggest that individuals with a more positive valence bias have a greater tendency to engage in CO-mediated regulatory processing of all ambiguous stimuli, even if they ultimately judge a specific face to have negative valence on a given trial.

**Figure 4**. BOLD activation in the CO Cluster during a late trial window. *Left*: Average activation for each emotional expression during the late window (16-19 sec; 4-7 sec after reveal image), including clear valence error trials. *Right*: Correlation between late CO activation during surprised trials and the average valence bias (percent of negative judgments for surprised trials) during the reveal phase. For both negative and positive judgments during noise, participants with a more positive valence bias overall exhibited greater CO activation.

*3.2.3 HRF Time Courses as a Function of Switching versus Staying*

To explore in greater detail the nature of the HRF time courses for each emotional expression and assess whether error recognition (objective or subjective) at the full reveal was driving activation, each participant's trials were sorted further according to whether they made the same response during the noise and reveal phases of the trial (i.e., "stay") or switched to the

opposite response, putatively indicating some type of error evaluation occurred (i.e., "switch").

As with the overall responses by emotional expression, a linear mixed model was fit with fixed

effects of emotion, time (linear and quadratic), and their interaction, and a random effect of

subject (intercept). Again, we focus on the CO cluster (**Figure 5**) given this network was

previously associated with post-trial performance reporting (Gratton et al., 2017), including in

response to ambiguity (Neta et al., 2017; Poudel et al., 2020), and error processing (Hester et al.,

2004; Iannaccone et al., 2015; Neta et al., 2014), but the results from other clusters are shown in

Supplemental Figure S2 and Tables S10-S13.

For switch trials in the CO cluster, there were significant effects of emotion ($F(4, 199.82)$

$= 26.09$, $p < .001$), time point (linear) ($F(1, 476.73) = 181.64$, $p < .001$), time point (quadratic)

($F(1, 510.66) = 240.32$, $p < .001$), and the interaction of emotion and time ($F(4, 204.77) = 45.33$,

$p < .001$; see Supplemental Table S8). Descriptively, surprised trials for which participants

responded "positive" during the noise phase showed the weakest response, while clear valence

error trials (negative happy, positive angry/fearful) had a stronger response. Critically, surprised

trials for which participants responded "negative" during noise and "positive" at reveal exhibited

a moderate response from 11 to 14 seconds, but showed the strongest response from 16 seconds

to the end of the trial (i.e., after the image had been revealed and the positive judgment made).

For stay trials, there were significant effects of emotion ($F(4, 220.56) = 6.79$, $p < .001$),

time point (quadratic) ($F(1, 684.90) = 42.72$, $p < .001$), and the interaction of emotion and time

($F(4, 281.60) = 16.21$, $p < .001$), and a non-significant effect of time point (linear) ($F(1, 687.95)$

$= 1.86$, $p = .173$; see Supplemental Table S9). Correct clear valence trials (positive happy,

negative angry/fearful) and negative surprised trials showed a relatively weak response.

Importantly, positive surprised trials showed the strongest response, particularly after 14

seconds, consistent with the switch trial results in demonstrating that the largest response occurred when the final valence judgment of surprised faces was positive.

*3.2.4 Emotional Expression HRF Peak by Response Frame*

Finally, the HRFs for each emotional expression were analyzed according to the frame during which a behavioral response was made to examine how the timing of the noise judgment impacted the HRF timing and shape. In the CO cluster during clear valence trials, the HRF generally had a similar, standard shape with a shift in peak time relative to the timing of the noise behavioral response (**Figure 6**; see Supplemental Table S14 for exact peak times). For surprised trials, however, the HRF time courses for early frames showed two peaks: the first which corresponded to the clear valence trial peaks and the second which occurred after the reveal image was shown. This double-peak effect may indicate that, specifically for surprised trials, the participants processed the valence of the face not only during the noise phase, but perhaps also completed some additional processing or performance reporting after the full image was revealed. This second level of processing is likely attributed to the ambiguity of the surprised expression. (The responses by frame for the other ROI clusters are shown in Supplemental Figure S3.)

**Figure 5**. Estimated HRF curves in the CO ROI Cluster for each emotion based on whether a participant switched their behavioral response from the noise to the reveal phase or stayed with the same response. Labels refer to the initial response made during the noise phase. The gray bar represents the timing of each frame of the trial from 0 = black screen to 5 = final noise frame and R = fully revealed image.

**Figure 6.** Estimated HRF curves for each emotional expression by response frame for the CO cluster. Averaged responses across participants according to the noise frame (2-5) during which a behavioral response was made for correct clear valence trials (happy, angry, fearful) and ambiguous (surprised) trials for negative and positive responses. Vertical lines at the top of each plot indicate the identified peak(s) for each curve.

*3.3 Follow-up Online Study*

In order to further tease apart the behavioral response pattern observed in the main study, particularly for surprised faces that are judged to have positive versus negative valence, a follow-up study was conducted online. The final sample consisted of 81 naïve participants (39 F/42 M) with a mean age of 31.1 years (range 18-39). In addition to collecting a larger sample of participants, the slow reveal paradigm here was modified to include seven (instead of five) frames of each facial expression in the "noise" phase of the trial (a full description of the methods is provided in the Supplemental Material). The longer period during which the face was unmasked was intended to allow for greater variability in response frame. With this extended window for noise responses and based on the initial negativity hypothesis, it was predicted that for surprised faces positive judgments would be made later than negative judgments.

Responses for valence judgments from the noise and reveal phases of the task were averaged separately for each of the four emotional expressions, resulting in an average percent negative rating (**Table 1**). A 4 x 2 (emotion by phase) repeated measures ANOVA indicated significant effects of emotion ($F(2.02, 161.77) = 1499.61$, $p < .001$, $\eta^2 = .949$), phase ($F(1, 80) = 83.27$, $p < .001$, $\eta^2 = .510$), and their interaction ($F(2.45, 196.05) = 133.82$, $p < .001$, $\eta^2 = .626$). Consistent with the results from the main study, happy faces were judged as mostly positive while angry and fearful faces were judged as mostly negative, with fewer errors (e.g., a happy face judged as negative) made during the reveal phase than the noise phase for each of the clearly valenced expressions ($t$'s $> 7.10$, $p$'s $< .001$). Unlike the main study, surprised faces were judged as significantly more negative during the reveal phase than the noise phase ($t(80) = 3.59$, $p < .001$, Cohen's $d = .399$), albeit with a smaller difference between phases than the clear valence emotions.

Next, the frame (1-7) during which responses occurred during the noise phase was compared for correct happy, angry, and fearful trials as well as separately for surprised trials that were judged as positive and negative. Descriptively, most responses occurred during frame 5 for all emotions (**Table 2**), but happy faces that were correctly judged as positive and surprised faces that were judged as negative yielded earlier responses (frames 3 and 4) than angry and fearful faces that were correctly judged as negative (frames 6 and 7). A one-way ANOVA by emotion on RT during the noise phase resulted in a main effect of emotion ($F(1.99, 158.93) = 12.26$, $p <$ .001, $\eta^2 = .133$). Participants made slower responses for angry and fearful face trials and faster responses for happy and negative surprised face trials. As in the main study, the reveal phase of the task showed a different RT pattern by emotion ($F(1.91, 135.92) = 81.95$, $p < .001$, $\eta^2 = .536$), with positive surprised trials resulting in the slowest responses, while happy trial responses were faster than all other emotions (**Tables 2/S3**).

Surprised trials for which participants stayed with the same response for the reveal phase were subsequently analyzed according to the percentage of responses in each frame (given that some participants did not make any positive stay responses, the sample size was reduced to 68 for this analysis). There was a significant effect of frame ($F(2.43, 279.70) = 34.79$, $p < .001$, $\eta^2 = .342$) and a significant interaction between frame and the valence of the response ($F(4.09, 273.82) = 3.92$, $p = .004$, $\eta^2 = .055$). Positive surprised stay trials were rated later than negative surprised stay trials. Specifically, pairwise comparisons indicated that during frames 3 and 4 more negative responses were made, while during frame 5 more positive responses were made (**Figure 7**). This effect was marginal for RT (stay positive: 9026 (2230) ms; stay negative: 8672 (1553) ms. $t(67)=1.53$, $p=.066$, Cohen's $d=.185$).

**Figure 7**. Percent of responses during each noise frame for trials on which participants stayed with the same response for the reveal phase.

## 4. Discussion

In the current work, a valence bias task was presented in which clear and ambiguous emotional facial expressions were slowly revealed over several seconds and participants were asked to make a valence judgment (positive/negative) about each face both during the unmasking and after the full image was presented. This design allows for examination of the mechanisms of processing ambiguity as they unfold over the course of a trial. Overall, ambiguous (surprised) trials on which the participant judged the face to have positive valence exhibited the most unique response characteristics.

In the main study, BOLD HRF time courses were extracted from 16 ROIs in three cortical networks previously associated with task control, decision making, and performance reporting. The results demonstrated greater activation for positive judgments of surprised faces –

notably in the CO network – and especially in the latter part of the trial, which included a second activation peak, following the presentation of the reveal image. In a follow-up study, the task was administered to a larger, online sample using a more extended slow reveal design with additional noise frames. Across both studies, participants responded more slowly when judging surprised faces as positive versus negative. Taken together, these findings support the initial negativity hypothesis (Neta, in press; Neta & Whalen, 2010), which posits that the default response to ambiguity is negative and that a positive evaluation requires greater control, effort, or regulation to overcome this initial negativity.

*4.1 Unfolding of Valence Judgments Differs by Emotion*

Behaviorally, on clear valence trials participants made more errors (e.g., judging a happy face to be negative) during the noise phase when the face was partially obscured than during the reveal phase when the full image was visible. Conversely, surprised trials showed no difference in average valence judgments between the two task phases in the fMRI study, suggesting that participants evaluated the surprised faces according to their own internal valence bias with minimal influence of the slow reveal paradigm. Participants in the online study, however, judged surprised faces as more positive during the noise phase than during the reveal phase. The elevated number of positive judgments (i.e., lower percent negative) for surprised faces may indicate that more participants in this larger sample made more subjective errors due to the perceptual uncertainty during the noise phase – as was the case for clear valence trials. An alternate explanation could be that the extended slow task design allowed participants time to adopt a positive interpretation of the surprised face during noise (see Neta & Tong, 2016), that was nonetheless overturned during the subsequent quick judgment at reveal. Unfortunately, the

current design does not allow for investigation of participants' motivation when switching their response, so future work will be needed to better understand this pattern of results.

Response times exhibited no significant difference between surprised faces judged as positive versus negative during the noise phase in the fMRI study or in the initial analysis in the follow-up study. However, given that participants occasionally switched their valence judgment from the noise to the reveal phase for surprised trials (possibly indicating a subjective or motoric error), we then compared response times for surprised trials on which a participant stayed with the same response during both phases in the follow-up online study. For these "stay" surprised trials, more negative responses were made in an earlier frame and more positive responses were made in a later frame during the noise phase. This result mirrors the surprised RT effects during the reveal phase and supports the idea that positive judgments of ambiguity require longer processing time to overcome an initial negativity, consistent with prior work (Neta & Tong, 2016; Petro et al., 2021), even in a context with perceptual uncertainty.

*4.2 Disentangling Perceptual and Valence Ambiguity*

Performance in the current task required two valence judgments about an emotional face that was slowly unmasked and then fully revealed. When considering the behavioral responses from both studies, we found evidence that distinct factors may contribute to decision-making processes during the noise and reveal phases of the task. During the noise phase, participants responded to the happy faces faster than the angry and fearful faces, indicating that perceptual features may be contributing to response times. Assuming that participants first recognized the expression or some subset of its physical features (Calvo & Nummenmaa, 2016) prior to making a decision about its valence (Storbeck et al., 2006), it may be that differences in the morphological features of the expressions impacted response times most strongly during the

noise phase. In other words, happy faces are easiest to recognize, likely due to the unique appearance of a smile (Calvo et al., 2012). Relatedly, other emotional expressions that are perceptually similar (e.g., both surprise and fear have widened eyes) may need more time/information to recognize the expression and categorize its valence (Calvo & Lundqvist, 2008; Calvo & Nummenmaa, 2016; Tottenham et al., 2009), which likely explains the longer response times for fearful faces. This is in contrast to the reveal phase, when participants still responded faster to happy than angry or fearful faces, but now responded the slowest to surprised faces that they judged as positive, as predicted. Thus, when the face was not obscured and there was no perceptual uncertainty, the morphological similarities had a weaker effect on the decision process than the valence ambiguity of surprise.

*4.3 Task Control Networks Support Ambiguity Processing*

The CO, R FP, and L FP control networks make variable contributions to decision making and task control that can be distinguished in part by their temporal characteristics. Previous research has investigated how these networks accumulate evidence, handle errors, and report task performance during the gradual reveal of objects or words in identification tasks (Gratton et al., 2017; Neta et al., 2017; Ploran et al., 2007; Wheeler et al., 2008). When examining HRF time courses in response to emotional faces (see Supplemental Material), we replicated prior work characterizing temporal characteristics of task control networks using slow reveal tasks (Gratton et al., 2017). This replication supports the domain-general decision-making functions of these regions and demonstrated the temporal effects using a more complex and socially relevant stimulus set that required a distinct type of decision to be made (i.e., categorizing emotional valence).

Critically, when the HRF was examined in these networks according to the emotional expression of the face, there was a marked increase in BOLD activation for surprised trials, particularly when a participant judged the face's valence as positive, as compared to trials with clearly valenced expressions. The surprised trial response was most notable in the CO network, with a strong response late in the trial for positive judgments and to a lesser extent negative judgments as well. In contrast, a sensory cluster composed of early visual regions showed no difference in activation by emotion and was engaged earlier in the trial as sensory information became available, well before a valence decision was made.

Further examination of the ambiguity-related late BOLD response in the CO network indicated a relationship with individual behavior such that greater activation during the noise phase of surprised trials was associated with a more positive valence bias (i.e., one's tendency to judge the expressions as positive). Therefore, individuals who have a more positive valence bias may process the ambiguous faces more thoroughly or effortfully regardless of their ultimate decision on a specific trial. This is consistent with other work demonstrating that physiological responses to emotional ambiguity track one's general tendencies (i.e., valence bias) rather than their transient response on a given trial (Neta et al., 2009). Additionally, an examination of the clear valence error trial activation (combined across happy, angry, and fearful) demonstrated a moderate response that did not significantly differ from surprised trials. This is consistent with previous work showing that ambiguity, RT, and error commission represent separable responses within the CO network (Neta et al., 2014).

*4.4 Ambiguity Elicits a Second BOLD Peak following the Second Valence Judgment*

To investigate how the HRFs differed as a function of *when* the first valence decision occurred, activation in the CO network was analyzed according to the frame during which a

participant responded. Clear valence expression trials showed a single HRF peak that was shifted in time relative to the response frame. Surprised trials showed a similar first peak, yet unexpectedly also exhibited a second HRF peak late in the trial. This peak was most evident for responses in frames 2 and 3, whereas for responses in frames 4 and 5, the two peaks seemed to overlap temporally. Similarly, the two peaks were less distinct in the prior analyses combined across frames, but nonetheless are visually apparent in the CO surprised trial HRFs (Fig. 3A).

The late peak presumably corresponds to a second judgment following the full reveal of the face image, which evidently was not necessary for clear valence expressions. Interestingly, this second peak occurred for both positive and negative surprised trials, indicating that this emotional expression not only modulated the strength of the CO activation, but also the shape of the HRF, as a result of distinct decision-making demands for ambiguous stimuli. Prior work using the slow reveal paradigm did not exhibit such a pattern for object recognition (Gratton et al., 2017; Neta et al., 2017), likely because once the image was fully revealed, the perceptual ambiguity was resolved. In other words, a picture of a toothbrush might look like many things when only partial visual information is available, but once the full image is presented, the object is clearly identifiable. In contrast, in the current work, the full reveal of an ambiguous (surprised) face is not sufficient for ambiguity resolution regarding the valence, so it elicits a second peak that may indicate participants were not simply confirming their first judgment, but perhaps making a distinct second decision about its valence. Nonetheless, the number of responses for each expression and frame was fairly small, so these results must be considered preliminary.

The present findings regarding the CO network's response to ambiguity support previous evidence of its functional role in performance reporting following a task decision (Gratton et al., 2017, 2018; Neta et al., 2014). Activation in this network began to increase later than sensory or

FP ROIs and as a function of the behavioral response timing. Alternate explanations of the function of CO regions, including salience detection, conflict monitoring, and alertness (Botvinick et al., 2001; Coste & Kleinschmidt, 2016; Dignath et al., 2020; Han et al., 2019; Seeley et al., 2007), do not align as well with our results given the specific late increase in activation for ambiguous over other emotional faces. It could be argued, however, that ambiguity might signal increased trial conflict, carry greater affective salience, and/or necessitate heightened attention. Nonetheless, the late onset of the CO response (i.e., after a decision has been made) supports a performance reporting role that is engaged at the end of an evaluation rather than an ongoing monitoring or attentional function (see also Neta et al., 2014).

*4.5 Evidence in Support of the Initial Negativity Hypothesis*

The increased activation for positive surprised trials in the CO network supports the proposal regarding the primacy of a negative judgment of ambiguity, with a positive judgment requiring additional top-down processing to overcome this initial assessment (Neta & Whalen, 2010; Petro et al., 2018). We examined two possible behavioral explanations for the unique effects for positive judgments of surprise. The first was reaction time, which had a significant effect in the HRF mixed models such that slower responses were associated with greater activation in each network. Nonetheless, the effect of emotional expression was significant even when accounting for RT. This finding was supported further by the correlations between reveal RTs and late CO activation, which were significant for fearful and negative surprised trials, but not for positive surprised trials. This pattern of results demonstrated that while processing time may contribute to CO activation, the large response for positive surprised trials was not merely an artefact of slower processing, but likely reflects a specific control process necessary to resolve the ambiguity.

Secondly, the increased activation in the CO network for positive surprised trials was analyzed according to whether a participant switched their response from the noise to reveal phase or stayed with the same response. We found that making a positive judgment of a surprised face during the reveal phase resulted in the greatest activation in the CO network regardless of the initial behavioral response (i.e., negative surprised switch trials and positive surprised stay trials had the largest late activation). This result implies that resolving the ambiguity of a surprised face with a positive judgment creates unique decision-making or regulatory demands, or perhaps entails repeated performance reporting following an uncertain initial decision. It also indicates that the large response for positive surprised trials was not a type of error signal, as even those trials on which participants stayed with a positive judgment during the reveal phase showed increased activation. Furthermore, clear valence trials where participants made an erroneous response during the noise phase and switched their valence judgment to the correct response during the reveal phase showed only a moderate response that was initially stronger than the surprised trials but weaker than the negative surprised switch trials in the later part of the trial.

Collectively, the current results demonstrate clear support for the initial negativity hypothesis, which proposes that most people initially form a quick negative appraisal of an ambiguous image, perhaps in service to preparing a response to a potential threat. Then, according to the individual's internal biases, they may reevaluate the stimulus to find a more positive appraisal (see also Neta et al., 2022). This process seems to be supported in part by the CO network, given the general increased activation for positive surprised trials and the relationship between a positive valence bias and the extent of this activation.

*4.6 Limitations and Future Directions*

The current results must be considered with respect to some limitations. First, positive judgments for surprised trials did not occur frequently and differed in frequency across participants, potentially impacting the nature of this response relative to other emotional expressions. This issue is further amplified in the exploratory analyses examining switch vs. stay trials and responses by frame, which divided positive surprised trials in an effort to investigate possible explanations for the unique responses. Further work with a greater number of trials or individuals with a more positive valence bias is needed to determine whether or how the infrequency of this condition impacts the HRF characteristics.

Second, the task design does not allow for separation of the HRFs for the noise and reveal phases, so it is not possible to precisely distinguish the contribution of each phase. This may impact the interpretation of the secondary, late peak for ambiguous faces, which was unexpected. Yet it is reasonable to conclude based on its timing that the later peak is driven by the presentation of the fully revealed face. Future work could explore the response to distinct phases of ambiguity processing in greater detail. For example, a task design with a delay between the noise and reveal phases or catch trials during which no reveal image appeared could help illuminate the nature of the two apparent peaks.

Finally, the average surprised valence judgments during the reveal phase in both studies were relatively negative compared to prior reports (e.g., Harp et al., 2022; Neta et al., 2009; Petro et al., 2018). It may be that the greater number of clearly negative facial expression conditions (angry and fearful vs. happy) created a negative context (Neta et al., 2011) that biased participants towards making a negative response. Importantly, however, surprised faces were immediately preceded by an equal number of clearly positive and negative faces. This point warrants further investigation to determine whether this negativity is a result of a negatively

biased context from the angry and fearful faces or if perhaps the nature of the slow reveal paradigm itself somehow shifts valence bias due to increased perceptual uncertainty or slow pacing.

*4.7 Conclusion*

In these two studies, we took a novel approach using a perceptual decision-making task to explore the mechanisms that support the unfolding of judgments about ambiguous and clear facial expressions. We demonstrated that cortical task control networks support processing of emotional facial expressions as they are slowly revealed. Positive judgments of surprised faces were accompanied by stronger activation in the CO network and slower response times, supporting the initial negativity hypothesis. Surprised faces also resulted in a second activation peak in the HRF following the full reveal, which likely reflects the continued valence ambiguity of these stimuli. Ambiguous stimuli often occur in daily life within contexts that may not provide sufficient evidence for a single clear interpretation. The present work suggests that making positive judgments of such ambiguity is an effortful, controlled process that varies according to individual biases.

Funding: This publication was made possible by Nebraska Tobacco Settlement Biomedical Research Development Funds and the National Institutes of Health (NIMH111640; PI: Neta).

The authors declare that there are no competing financial interests.

## 5. References

Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S., & Cohen, J. D. (2001). Conflict monitoring and cognitive control. *Psychological Review*, *108*(3), 624.

Brown, C. C., Raio, C. M., & Neta, M. (2017). Cortisol responses enhance negative valence perception for ambiguous facial expressions. *Scientific Reports*, *7*(1), 15107. https://doi.org/10.1038/s41598-017-14846-3

Calvo, M. G., Fernández-Martín, A., & Nummenmaa, L. (2012). Perceptual, categorical, and affective processing of ambiguous smiling facial expressions. *Cognition*, *125*(3), 373–393. https://doi.org/10.1016/j.cognition.2012.07.021

Calvo, M. G., & Lundqvist, D. (2008). Facial expressions of emotion (KDEF): Identification under different display-duration conditions. *Behavior Research Methods*, *40*(1), 109–115. https://doi.org/10.3758/BRM.40.1.109

Calvo, M. G., & Nummenmaa, L. (2016). Perceptual and affective mechanisms in facial expression recognition: An integrative review. *Cognition and Emotion*, *30*(6), 1081–1106. https://doi.org/10.1080/02699931.2015.1049124

Coste, C. P., & Kleinschmidt, A. (2016). Cingulo-opercular network activity maintains alertness. *NeuroImage*, *128*, 264–272. https://doi.org/10.1016/j.neuroimage.2016.01.026

Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research*, *29*(3), 162–173.

Cox, R. W. (2012). AFNI: what a long, strange trip it's been. *Neuroimage*, *62*(2), 743–747.

Dignath, D., Eder, A. B., Steinhauser, M., & Kiesel, A. (2020). Conflict monitoring and the affective-signaling hypothesis—An integrative review. *Psychonomic Bulletin & Review*, *27*(2), 193–216. https://doi.org/10.3758/s13423-019-01668-9

Dosenbach, N. U. F., Fair, D. A., Cohen, A. L., Schlaggar, B. L., & Petersen, S. E. (2008). A
dual-networks architecture of top-down control. *Trends in Cognitive Sciences*, *12*(3), 99–
105. https://doi.org/10.1016/j.tics.2008.01.001

Ekman, P. (1976). Pictures of facial affect. *Consulting Psychologists Press*.

Goeleven, E., De Raedt, R., Leyman, L., & Verschuere, B. (2008). The Karolinska directed
emotional faces: A validation study. *Cognition & Emotion*, *22*(6), 1094–1118.

Gratton, C., Neta, M., Sun, H., Ploran, E. J., Schlaggar, B. L., Wheeler, M. E., Petersen, S. E., &
Nelson, S. M. (2017). Distinct Stages of Moment-to-Moment Processing in the
Cinguloopercular and Frontoparietal Networks. *Cerebral Cortex*, *27*(3), 2403–2417.
https://doi.org/10.1093/cercor/bhw092

Gratton, C., Sun, H., & Petersen, S. E. (2018). Control networks and hubs. *Psychophysiology*,
*55*(3), e13032. https://doi.org/10.1111/psyp.13032

Han, S. W., Eaton, H. P., & Marois, R. (2019). Functional Fractionation of the Cingulo-opercular
Network: Alerting Insula and Updating Cingulate. *Cerebral Cortex*, *29*(6), 2624–2638.
https://doi.org/10.1093/cercor/bhy130

Harp, N. R., Freeman, J. B., & Neta, M. (2022). Mindfulness-based stress reduction triggers a
long-term shift toward more positive appraisals of emotional ambiguity. *Journal of
Experimental Psychology: General*. https://doi.org/10.1037/xge0001173

Hester, R., Fassbender, C., & Garavan, H. (2004). Individual Differences in Error Processing: A
Review and Reanalysis of Three Event-related fMRI Studies Using the GO/NOGO Task.
*Cerebral Cortex*, *14*(9), 986–994. https://doi.org/10.1093/cercor/bhh059

Iannaccone, R., Hauser, T. U., Staempfli, P., Walitza, S., Brandeis, D., & Brem, S. (2015). Conflict monitoring and error processing: New insights from simultaneous EEG–fMRI. *NeuroImage*, *105*, 395–407. https://doi.org/10.1016/j.neuroimage.2014.10.028

Kim, H., Somerville, L. H., Johnstone, T., Alexander, A. L., & Whalen, P. J. (2003). Inverse amygdala and medial prefrontal cortex responses to surprised faces. *NeuroReport*, *14*(18).

Kim, H., Somerville, L. H., Johnstone, T., Polis, S., Alexander, A. L., Shin, L. M., & Whalen, P. J. (2004). Contextual modulation of amygdala responsivity to surprised faces. *Journal of Cognitive Neuroscience*, *16*(10), 1730–1745.

Neta, M. (in press). Valence bias: Individual differences in response to ambiguity. In J. Gross & B. Ford (Eds.), *Handbook of Emotion Regulation* (3rd ed.).

Neta, M., Berkebile, M. M., & Freeman, J. B. (2021). The dynamic process of ambiguous emotion perception. *Cognition and Emotion*, *35*(4), 722–729. https://doi.org/10.1080/02699931.2020.1862063

Neta, M., Davis, F. C., & Whalen, P. J. (2011). Valence resolution of ambiguous facial expressions using an emotional oddball task. *Emotion (Washington, D.C.)*, *11*(6), 1425–1433. https://doi.org/10.1037/a0022993

Neta, M., Harp, N. R., Tong, T. T., Clinchard, C. J., Brown, C. C., Gross, J. J., & Uusberg, A. (2022). Think again: The role of reappraisal in reducing negative valence bias. *Cognition and Emotion*, 1–16. https://doi.org/10.1080/02699931.2022.2160698

Neta, M., Kelley, W. M., & Whalen, P. J. (2013). Neural responses to ambiguity involve domain-general and domain-specific emotion processing systems. *Journal of Cognitive Neuroscience*, *25*(4), 547–557. https://doi.org/10.1162/jocn_a_00363

Neta, M., & Kim, M. J. (2022). Surprise as an Emotion: A Response to Ortony. *Perspectives on Psychological Science*, 17456916221132788. https://doi.org/10.1177/17456916221132789

Neta, M., Nelson, S. M., & Petersen, S. E. (2017). Dorsal Anterior Cingulate, Medial Superior Frontal Cortex, and Anterior Insula Show Performance Reporting-Related Late Task Control Signals. *Cerebral Cortex*, *27*(3), 2154–2165. https://doi.org/10.1093/cercor/bhw053

Neta, M., Norris, C. J., & Whalen, P. J. (2009). Corrugator Muscle Responses Are Associated With Individual Differences in Positivity-Negativity Bias. *Emotion*, *9*(5), 640–648. https://doi.org/10.1037/a0016819

Neta, M., Schlaggar, B. L., & Petersen, S. E. (2014). Separable responses to error, ambiguity, and reaction time in cingulo-opercular task control regions. *NeuroImage*, *99*, 59–68. https://doi.org/10.1016/j.neuroimage.2014.05.053

Neta, M., & Tong, T. T. (2016). Don't like what you see? Give it time: Longer reaction times associated with increased positive affect. *Emotion*, *16*(5), 730.

Neta, M., & Whalen, P. J. (2010). The primacy of negative interpretations when resolving the valence of ambiguous facial expressions. *Psychological Science*, *21*(7), 901–907.

Petro, N. M., Basyouni, R., & Neta, M. (2021). Positivity effect in aging: Evidence for the primacy of positive responses to emotional ambiguity. *Neurobiology of Aging*, *106*, 232–240. https://doi.org/10.1016/j.neurobiolaging.2021.06.015

Petro, N. M., Tong, T. T., Henley, D. J., & Neta, M. (2018). Individual differences in valence bias: FMRI evidence of the initial negativity hypothesis. *Social Cognitive and Affective Neuroscience*, *13*(7), 687–698.

Ploran, E. J., Nelson, S. M., Velanova, K., Donaldson, D. I., Petersen, S. E., & Wheeler, M. E. (2007). Evidence accumulation and the moment of recognition: Dissociating perceptual recognition processes using fMRI. *Journal of Neuroscience*, *27*(44), 11912–11924.

Poudel, R., Riedel, M. C., Salo, T., Flannery, J. S., Hill-Bowen, L. D., Eickhoff, S. B., Laird, A. R., & Sutherland, M. T. (2020). Common and distinct brain activity associated with risky and ambiguous decision-making. *Drug and Alcohol Dependence*, *209*, 107884.

Samuelsson, H., Jarnvik, K., Henningsson, H., Andersson, J., & Carlbring, P. (2012). The Umeå University Database of Facial Expressions: A Validation Study. *J Med Internet Res*, *14*(5), e136. https://doi.org/10.2196/jmir.2196

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., Reiss, A. L., & Greicius, M. D. (2007). Dissociable Intrinsic Connectivity Networks for Salience Processing and Executive Control. *The Journal of Neuroscience*, *27*(9), 2349. https://doi.org/10.1523/JNEUROSCI.5587-06.2007

Sterzer, P., Russ, M. O., Preibisch, C., & Kleinschmidt, A. (2002). Neural correlates of spontaneous direction reversals in ambiguous apparent visual motion. *Neuroimage*, *15*(4), 908–916.

Storbeck, J., Robinson, M. D., & McCourt, M. E. (2006). Semantic Processing Precedes Affect Retrieval: The Neurological Case for Cognitive Primacy in Visual Processing. *Review of General Psychology*, *10*(1), 41–55. https://doi.org/10.1037/1089-2680.10.1.41

Talairach, J., & Tournoux, P. (1988). Co-planar stereotaxic atlas of the human brain-3-dimensional proportional system. *An Approach to Cerebral Imaging*.

Thompson-Schill, S. L., D'Esposito, M., Aguirre, G. K., & Farah, M. J. (1997). Role of left

inferior prefrontal cortex in retrieval of semantic knowledge: A reevaluation.

*Proceedings of the National Academy of Sciences*, *94*(26), 14792–14797.

Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., Marcus, D. J.,

Westerlund, A., Casey, B., & Nelson, C. (2009). The NimStim set of facial expressions:

Judgments from untrained research participants. *Psychiatry Research*, *168*(3), 242–249.

https://doi.org/10.1016/j.psychres.2008.05.006

Wheeler, M. E., Petersen, S. E., Nelson, S. M., Ploran, E. J., & Velanova, K. (2008).

Dissociating early and late error signals in perceptual recognition. *Journal of Cognitive

Neuroscience*, *20*(12), 2211–2225.

Willenbockel, V., Sadr, J., Fiset, D., Horne, G. O., Gosselin, F., & Tanaka, J. W. (2010).

Controlling low-level image properties: The SHINE toolbox. *Behavior Research

Methods*, *42*(3), 671–684. https://doi.org/10.3758/BRM.42.3.671

Supplemental Material

# Specialized cingulo-opercular network activation in judging ambiguity of facial expressions in a slow reveal fMRI paradigm

Jordan E. Pierce, Nathan M. Petro, Elizabeth Clancy, Caterina Gratton, Steven E. Petersen, Maital Neta

**Supplemental Methods**

*HRF Response Characteristics by ROI Cluster*

The data from all runs were entered into a general linear model with regressors based on each individual participant's noise responses (from frames 2-5), collapsing across all emotion expressions. The "TENT" function was used to estimate the amplitude of the hemodynamic response at each TR from 0-21 seconds after stimulus onset (22 timepoints; TR = 1 second) without assuming a predetermined shape for the response in each voxel. Regressors of no interest included anticipatory responses (frame 1), error trials (incorrect judgment for a clearly valenced stimulus), polynomials for each run (two terms), and six motion parameters estimated during alignment (x, y, z shift/rotation).

To compare the response timing and shape in each of the three ROI clusters of interest, several parameters characterizing the HRF were extracted. Following the approach of Gratton et al. (2017), peak time, onset time, offset time, and full width-half maximum (FWHM) were quantified for each ROI cluster in each participant. First, time courses from each ROI were averaged to create an estimated HRF for each ROI cluster for each participant. Next, these average HRFs were expanded from 22 time points to 2,200 time points using linear interpolation in MATLAB to obtain better characterization of the underlying HRF and limit the influence of

any potential outlying single data points. From these interpolated time courses, peak time was defined as the time point at which the curve reached its maximum amplitude. Onset time was defined as the time point at which the curve exceeded 25% of the peak amplitude (calculated by stepping back from the peak), while offset time was defined as the time point at which the curve dropped below 25% of the peak amplitude after the peak time. FWHM was defined as the time between the points at which the curve rose above and fell below 50% of the peak amplitude. A repeated measures ANOVA was conducted to compare the response characteristics between ROI cluster (3 levels) for peak time, onset, offset, FWHM, and peak amplitude. Huynh-Feldt adjusted degrees of freedom were used to account for unequal variance between conditions when indicated by a significant Mauchly's test of sphericity and Bonferroni corrected $p$-values were used for pairwise comparisons.

**Follow-up Online Study**

*Participants*

Data was collected from 135 participants recruited online using Gorilla Experiment Builder (Anwyl-Irvine et al., 2020). Participants were required to be at least 18 years old and residing in the United States. Participants' data were excluded if at least half of the task was not completed, participants did not respond to attention checks (press 'A' or 'L' when cued), or if clear valence expressions were not judged correctly at the reveal (e.g., judging most happy faces to be negative). This resulted in a final sample of 81 participants (39 F/42 M) with a mean age of 31.1 years (range 18-39), and race/ethnicity reported as 75% White, 10% Asian, 5% Black, and 5% Hispanic/Latino. Participants received monetary compensation for their time. All task materials and procedures were approved by the UNL Institutional Review Board.

*Task Design and Procedure*

Each trial began with the presentation of a white fixation cross in the center of a black screen for 250 ms, followed by a black screen for 2 seconds. Then, a grayscale face was presented that was occluded by a black mask that was slowly degraded over the course of seven 2-second frames. The black mask was degraded by randomly removing pixels following a Gaussian distribution over the seven "noise" frames such that 3.5, 5, 6.5, 8.5, 11, 13.5, and 16% of the face was visible at each successive step. The trial ended with the presentation of the fully revealed face stimulus for 2 seconds. Thus, the duration of each trial was 18 seconds from the onset of the black screen to the offset of the face stimulus.

As in the main study, participants were instructed to judge each face as either positive or negative twice during the trial, once when the picture was emerging and again when the picture was fully revealed, and were told that the two responses could be the same or different. The participants completed ten practice trials containing only angry and happy faces that were not used in the main experiment, for which they received feedback as to whether they correctly identified the image's valence during the noise and reveal stages. There was a total of 120 trials (30 each of happy/angry/fearful/surprised), with faces selected from the NimStim Set of Facial Expressions and the Karolinska Directed Emotional Faces database. Participants used a keyboard button ("A" or "L") to rate each image with the mapping to positive/negative counterbalanced across participants.

*Data Analysis*

As in the main study, the average valence judgments for each emotion were calculated as the number of negative responses out of the number of total trials for each participant, multiplied by 100 (i.e., percent negative). A 4 x 2 (emotion by trial phase) repeated measures ANOVA was conducted on the average judgments to compare responses for happy, angry, fearful, and

surprised faces during the first "noise" phase and the second "reveal" phase. Subsequently, incorrect judgments for clearly valenced emotional expressions (happy judged as negative and angry or fearful judged as positive) were considered errors and not included in the analyses.

Next, the average response frame (out of 7) was calculated for each of the five emotion conditions (happy, angry, fearful, positive surprised, and negative surprised) for each participant. The noise RT was calculated from the beginning of the first frame, giving a maximum possible value of 14 seconds (2 sec x 7 frames), while the reveal RT had a maximum possible value of 2 seconds. One-way ANOVAs by emotion were conducted on noise and reveal RTs; Huynh-Feldt adjusted degrees of freedom were used to account for unequal variance between conditions when indicated by a significant Mauchly's test of sphericity.

To further examine the behavioral relationship between positive and negative responses on surprised trials, the percent of responses made during each of the seven frames was compared depending on whether the participant stayed with the same response during the noise and reveal phases. Given the low number of "switch" surprised trials, only noise trials that were judged as having the same valence during the reveal phase were included. These "stay" surprised trials are presumably those for which the participant made a subjectively "correct" judgment during the noise phase and, thus, confirmed this judgment during the reveal phase. As some participants did not make any positive judgments of surprised faces during the reveal phase, this resulted in a reduction of the sample size to 68 for this analysis.

**Supplemental Results**

*Response Characteristics by ROI Cluster*

This analysis focused on differences in response characteristics of the estimated HRF curve across three ROI clusters (CO, L FP, R FP) created from 16 ROIs from Gratton et al. (2017). Based on the average HRF in each ROI cluster (**Figure S1**), onset time, offset time, peak time, peak amplitude, and FWHM were calculated for each participant and compared across ROI clusters using one-way ANOVAs. There was a significant effect of ROI cluster in all measures (Figure S1). For onset time, there was a significant effect of ROI cluster ($F(1.75, 54.16) = 5.22$, $p = .011$, $\eta^2 = .144$), with the L FP cluster response rising earlier than the R FP response (Bonferroni corrected $p = .026$). For peak time ($F(1.51, 46.76) = 10.00$, $p < .001$, $\eta^2 = .244$) and offset time, ($F(2, 62) = 6.10$, $p = .004$, $\eta^2 = .164$), the pairwise comparisons indicated that the R FP response peaked later than L FP ($p = .015$) and ended later than both other regions (R FP vs L FP: $p = .028$; R FP vs CO: $p = .011$). For FWHM, there was also a significant effect ($F(1.68, 52.17) = 6.29$, $p = .006$, $\eta^2 = .169$), such that the CO had a smaller FWHM than the R FP cluster ($p < .001$). Finally, there was a significant effect for peak amplitude ($F(2, 62) = 3.88$, $p = .026$, $\eta^2 = .111$), such that the R FP response had a larger amplitude than the L FP response ($p = .035$).

Across all emotions, the R FP network had a stronger, later, more extended response, while the L FP and CO networks had a more transient response, consistent with the findings of Gratton et al. (2017). The L FP network also became activated earlier and with a smaller amplitude than the other networks. These results imply that the control networks are performing similar decision making and performance reporting functions in the slow reveal paradigm regardless of the specific stimulus presented or task decision that must be made.

**Table S1**. MNI Coordinates for each region of interest

| Region | x | y | z |
|---|---|---|---|
| *Cingulo-opercular cluster* | | | |
| Left AI/FO | 31 | -25 | -1 |
| Right AI/FO | -33 | -26 | 1 |
| Pre-SMA 1 | -1 | -15 | 51 |
| Pre-SMA 2 | -7 | -24 | 37 |
| *Left frontoparietal cluster* | | | |
| Left IPS 1 | 42 | 41 | 45 |
| Left IPS 2 | 32 | 50 | 45 |
| Left IPS 3 | 24 | 67 | 49 |
| Left frontal | 43 | -4 | -33 |
| Left MFG 1 | 45 | -25 | 22 |
| Left MFG 2 | 43 | -34 | 14 |
| Right IPS 2 | -32 | 53 | 49 |
| Right frontal | -44 | -9 | 28 |
| *Right frontoparietal cluster* | | | |
| Right IPS 1 | -44 | 45 | 50 |
| Right IFG | -53 | -15 | 16 |
| Right MFG 1 | -44 | -32 | 30 |
| Right MFG 2 | -43 | -42 | 18 |
| *Sensory cluster (control)* | | | |
| Right cuneus | 19 | 102 | -4 |
| Left cuneus | -16 | 102 | -5 |
| Right lingual gyrus | -1 | 95 | -13 |
| Left lingual gyrus | 10 | 102 | -12 |
| R middle occipital gyrus | -30 | 81 | 14 |

AI = anterior insula; FO = frontal/operculum; IFG = inferior frontal gyrus; IPS = inferior parietal sulcus; MFG = middle frontal gyrus; SMA = supplemental motor area

**Table S2**. Pairwise RT comparisons for the main study.

| Emotion | | Difference | 95% CI | S.E. | p-value |
|---|---|---|---|---|---|
| *Noise Phase* | | | | | |
| **Happy** | **Angry** | **-714.2** | **(-886.1, -542.3)** | **84.2** | **<.001** |
| **Happy** | **Fearful** | **-508.4** | **(-673.0, -343.9)** | **80.6** | **<.001** |
| **Happy** | **Positive Surprised** | **-417.4** | **(-776.1, -58.7)** | **175.6** | **.024** |
| Happy | Negative Surprised | -148.3 | (-356.8, 60.1) | 102.1 | .156 |
| **Angry** | **Fearful** | **205.8** | **(104.1, 307.5)** | **49.8** | **<.001** |
| Angry | Positive Surprised | 296.9 | (-64.0, 657.7) | 176.7 | .103 |
| **Angry** | **Negative Surprised** | **565.9** | **(426.4, 705.4)** | **68.3** | **<.001** |
| Fearful | Positive Surprised | 91.1 | (-249.7, 431.9) | 166.9 | .589 |
| **Fearful** | **Negative Surprised** | **360.1** | **(254.0, 466.2)** | **51.9** | **<.001** |
| Positive Surprised | Negative Surprised | 269.0 | (-69.6, 607.7) | 165.8 | .115 |
| *Reveal Phase* | | | | | |
| **Happy** | **Angry** | **-75.7** | **(-93.8, -57.7)** | **8.8** | **<.001** |
| **Happy** | **Fearful** | **-84.2** | **(-111.7, -56.7)** | **13.4** | **<.001** |
| **Happy** | **Positive Surprised** | **-387.4** | **(-450.6, -324.3)** | **30.7** | **<.001** |
| **Happy** | **Negative Surprised** | **-142.5** | **(-187.6, -97.3)** | **22.0** | **<.001** |
| Angry | Fearful | -8.5 | (-26.4, 9.5) | 8.7 | .341 |
| **Angry** | **Positive Surprised** | **-311.7** | **(-368.3, -255.2)** | **27.5** | **<.001** |
| **Angry** | **Negative Surprised** | **-66.8** | **(-106.5, -27.0)** | **19.3** | **.002** |
| **Fearful** | **Positive Surprised** | **-303.3** | **(-358.3, -248.3)** | **26.7** | **<.001** |
| **Fearful** | **Negative Surprised** | **-58.3** | **(-95.1, -21.5)** | **17.9** | **.003** |
| **Positive Surprised** | **Negative Surprised** | **245.0** | **(181.5, 308.4)** | **30.9** | **<.001** |

Note: Bold font indicates a significant difference (p<.05).

**Table S3**. Pairwise RT comparisons for the follow-up study.

| Emotion | | Difference | 95% CI | S.E. | p-value |
|---|---|---|---|---|---|
| *Noise Phase* | | | | | |
| **Happy** | **Angry** | **-704.5** | **(-844.0, -564.9)** | **70.1** | **<.001** |
| **Happy** | **Fearful** | **-356.3** | **(-501.3, -211.2)** | **72.9** | **<.001** |
| Happy | Positive Surprised | -180.5 | (-477.8, 116.7) | 149.4 | .230 |
| Happy | Negative Surprised | -122.2 | (-278.7, 34.3) | 78.6 | .124 |
| **Angry** | **Fearful** | **348.2** | **(215.6, 480.9)** | **66.6** | **<.001** |
| **Angry** | **Positive Surprised** | **524.0** | **(217.5, 725.6)** | **154.0** | **.001** |
| **Angry** | **Negative Surprised** | **582.3** | **(438.9, 501.3)** | **72.0** | **<.001** |
| Fearful | Positive Surprised | 175.7 | (-111.6, 463.1) | 144.4 | .227 |
| **Fearful** | **Negative Surprised** | **234.0** | **(105.8, 362.3)** | **64.4** | **<.001** |
| Positive Surprised | Negative Surprised | 58.3 | (-255.6, 372.2) | 157.7 | .713 |
| *Reveal Phase* | | | | | |
| **Happy** | **Angry** | **-48.9** | **(-71.0, -26.8)** | **11.1** | **<.001** |
| **Happy** | **Fearful** | **-86.9** | **(-107.1, -66.7)** | **10.1** | **<.001** |
| **Happy** | **Positive Surprised** | **-336.3** | **(-391.0, -281.5)** | **27.5** | **<.001** |
| **Happy** | **Negative Surprised** | **-95.9** | **(-127.6, -64.1)** | **15.9** | **<.001** |
| **Angry** | **Fearful** | **-38.0** | **(-54.7, -21.2)** | **8.4** | **<.001** |
| **Angry** | **Positive Surprised** | **-287.3** | **(-341.8, -232.9)** | **27.3** | **<.001** |
| **Angry** | **Negative Surprised** | **-46.9** | **(-78.8, -15.1)** | **16.0** | **.004** |
| **Fearful** | **Positive Surprised** | **-249.4** | **(-303.5, -195.2)** | **27.2** | **<.001** |
| Fearful | Negative Surprised | -9.0 | (-36.1, 18.2) | 13.6 | .512 |
| **Positive Surprised** | **Negative Surprised** | **240.4** | **(180.4, 300.4)** | **30.1** | **<.001** |

Note: Bold font indicates a significant difference (p<.05).

**Figure S1**. Estimated HRF curves of the three ROI clusters. The upper left panel shows the interpolated HRF curve averaged from each of the ROIs within the three network clusters across participants. The gray bar represents the timing of each frame of the trial from 0 = black screen to 5 = final noise frame and R = fully revealed image. The L FP cluster had the earliest onset and smallest amplitude, while the R FP cluster had a later onset, offset, and peak time with the largest amplitude, and the CO cluster had the smallest FWHM. The bar graphs illustrate the metrics derived from these curves to characterize differences in shape and timing across clusters.

**HRF Emotional Expression Analyses**
Linear Mixed Models [Emotion reference = Negative Surprised]

**Table S4.** CO ROI Cluster Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| **Intercept** | **1, 318.669** | **3.908** | **.049** |
| **Emotion** | **4, 306.265** | **7.918** | **<.001** |
| Time (linear) | 1, 679.043 | 3.350 | .068 |
| **Time (quadratic)** | **1, 696.416** | **49.373** | **<.001** |
| **Emotion * Time** | **4, 372.264** | **19.573** | **<.001** |
| **Reveal RT** | **1, 925.871** | **65.655** | **<.001** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept** | **-.112** | **.036** | **322.064** | **-3.130** | **.002** | **-.182, -.042** |
| **Happy** | **.077** | **.015** | **324.562** | **5.300** | **<.001** | **.049, .106** |
| **Angry** | **.062** | **.014** | **342.901** | **4.360** | **<.001** | **.034, .091** |
| **Fearful** | **.043** | **.013** | **311.277** | **3.267** | **.001** | **.017, .069** |
| Positive Surprised | .028 | .029 | 177.188 | .954 | .341 | -.030, .085 |
| **Time (linear)** | **.013** | **.003** | **552.516** | **3.895** | **<.001** | **.007, .020** |
| **Time (quadratic)** | **-.002** | **.000** | **696.416** | **-7.027** | **<.001** | **-.002, -.001** |
| **Happy * Time** | **-.017** | **.002** | **361.973** | **-7.722** | **<.001** | **-.021, -.013** |
| **Angry * Time** | **-.014** | **.002** | **390.001** | **-6.163** | **<.001** | **-.018, -.009** |
| **Fearful * Time** | **-.010** | **.002** | **369.662** | **-4.942** | **<.001** | **-.015, -.006** |
| Positive Surprised * Time | .003 | .004 | 167.975 | .780 | .437 | -.005, .011 |
| **Reveal RT** | **.000** | **.000** | **925.871** | **8.103** | **<.001** | **.000, .000** |

Estimate of Random Effects Covariance Parameters:

| Parameter | | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.009** | **.002** | **3.864** | **<.001** | **.005, .015** |

Note: Bold font indicates a significant effect (p<.05).

**Table S5**. L FP ROI Cluster Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| Intercept | 1, 248.664 | .139 | .710 |
| **Emotion** | **4, 300.360** | **3.634** | **.007** |
| **Time (linear)** | **1, 640.706** | **17.136** | **<.001** |
| **Time (quadratic)** | **1, 713.639** | **77.132** | **<.001** |
| **Emotion * Time** | **4, 350.836** | **5.095** | **<.001** |
| **Reveal RT** | **1, 904.890** | **16.900** | **<.001** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| Intercept | -.009 | .036 | 256.566 | -.257 | .797 | -.079, .061 |
| **Happy** | **.040** | **.015** | **340.395** | **2.689** | **.008** | **.011, .069** |
| **Angry** | **.038** | **.013** | **302.590** | **2.797** | **.005** | **.011, .064** |
| Fearful | .005 | .013 | 283.211 | .413 | .680 | -.020, .031 |
| Positive Surprised | .029 | .026 | 178.416 | 1.092 | .276 | -.023, .080 |
| **Time (linear)** | **.015** | **.003** | **487.322** | **4.757** | **<.001** | **.009, .022** |
| **Time (quadratic)** | **-.002** | **.000** | **713.639** | **-8.782** | **<.001** | **-.003, -.002** |
| **Happy * Time** | **-.008** | **.002** | **388.469** | **-3.620** | **<.001** | **-.013, -.004** |
| **Angry * Time** | **-.006** | **.002** | **381.373** | **-2.965** | **.003** | **-.011, -.002** |
| Fearful * Time | -.003 | .002 | 387.589 | -1.403 | .161 | -.007, .001 |
| Positive Surprised * Time | .002 | .004 | 171.487 | .614 | .540 | -.005, .009 |
| **Reveal RT** | **.000** | **.000** | **904.890** | **4.111** | **<.001** | **.000, .000** |

Estimate of Random Effects Covariance Parameters:

| Parameter | | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.011** | **.003** | **3.849** | **<.001** | **.007, .019** |

Note: Bold font indicates a significant effect (p<.05).

**Table S6**. R FP ROI Cluster Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| **Intercept** | 1, 240.547 | 28.876 | <.001 |
| **Emotion** | 4, 281.994 | 15.618 | <.001 |
| **Time (linear)** | 1, 630.802 | 431.905 | <.001 |
| **Time (quadratic)** | 1, 716.230 | 507.716 | <.001 |
| **Emotion * Time** | 4, 353.784 | 11.907 | <.001 |
| **Reveal RT** | 1, 862.805 | 57.919 | <.001 |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept** | -.223 | .038 | 247.748 | -5.828 | <.001 | -.298, -.147 |
| **Happy** | .105 | .015 | 324.014 | 6.964 | <.001 | .075, .134 |
| **Angry** | .029 | .013 | 278.442 | 2.148 | .033 | .002, .055 |
| **Fearful** | .039 | .013 | 277.223 | 2.876 | .004 | .012, .065 |
| **Positive Surprised** | -.071 | .026 | 165.515 | -2.707 | .008 | -.123, -.019 |
| **Time (linear)** | .067 | .003 | 445.112 | 19.106 | <.001 | .060, .073 |
| **Time (quadratic)** | -.006 | .000 | 716.230 | -22.533 | <.001 | -.007, -.005 |
| **Happy * Time** | -.011 | .002 | 410.815 | -4.480 | <.001 | -.016, -.006 |
| Angry * Time | -.003 | .002 | 394.083 | -1.419 | .157 | -.008, .001 |
| Fearful * Time | -.004 | .002 | 389.169 | -1.677 | .094 | -.008, .001 |
| **Positive Surprised * Time** | .013 | .004 | 173.781 | 3.429 | <.001 | .006, .020 |
| **Reveal RT** | .000 | .000 | 862.805 | 7.610 | <.001 | .000, .000 |

Estimate of Random Effects Covariance Parameters:

| Parameter | | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | .013 | .003 | 3.879 | <.001 | .008, .022 |

Note: Bold font indicates a significant effect (p<.05).

**Table S7**. Sensory ROI Cluster Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| **Intercept** | **1, 92.594** | **62.540** | **<.001** |
| Emotion | 4, 306.107 | .604 | .660 |
| **Time (linear)** | **1, 594.887** | **742.642** | **<.001** |
| **Time (quadratic)** | **1, 723.891** | **944.137** | **<.001** |
| Emotion * Time | 4, 459.008 | .341 | .850 |
| **Reveal RT** | **1, 681.560** | **7.851** | **.005** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept** | **.885** | **.109** | **95.892** | **8.129** | **<.001** | **.669, 1.101** |
| Happy | -.049 | .036 | 320.077 | -1.379 | .169 | -.120, .021 |
| Angry | -.041 | .038 | 298.696 | -1.081 | .281 | -.115, .033 |
| Fearful | -.043 | .034 | 275.697 | -1.258 | .210 | -.111, .024 |
| Positive Surprised | -.030 | .063 | 191.277 | -.473 | .637 | -.154, .095 |
| **Time (linear)** | **.228** | **.009** | **382.954** | **25.031** | **<.001** | **.210, .246** |
| **Time (quadratic)** | **-.024** | **.001** | **723.891** | **-30.727** | **<.001** | **-.025, -.022** |
| Happy * Time | .005 | .006 | 471.234 | .761 | .447 | -.008, .017 |
| Angry * Time | .003 | .007 | 481.062 | .433 | .665 | -.011, .016 |
| Fearful * Time | .006 | .007 | 467.424 | .910 | .363 | -.007, .019 |
| Positive Surprised * Time | .010 | .010 | 233.667 | .930 | .353 | -.011, .030 |
| **Reveal RT** | **.000** | **.000** | **681.560** | **-2.802** | **.005** | **.000, .000** |

Estimate of Random Effects Covariance Parameters:

| Parameter | | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.209** | **.053** | **3.920** | **<.001** | **.127, .344** |

Note: Bold font indicates a significant effect (p<.05).

## Switch vs. Stay Analysis

**Table S8**. CO ROI Cluster Switch Trials Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| **Intercept** | **1, 76.644** | **8.388** | **.005** |
| **Emotion** | **4, 199.821** | **26.090** | **<.001** |
| **Time (linear)** | **1, 476.728** | **181.641** | **<.001** |
| **Time (quadratic)** | **1, 510.659** | **240.315** | **<.001** |
| **Emotion * Time** | **4, 204.771** | **45.333** | **<.001** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept** | **-.079** | **.037** | **113.871** | **-2.169** | **.032** | **-.152, -.007** |
| **Happy** | **.258** | **.037** | **177.630** | **6.974** | **<.001** | **.185, .331** |
| **Angry** | **.277** | **.030** | **176.152** | **9.388** | **<.001** | **.219, .335** |
| **Fearful** | **.204** | **.031** | **204.402** | **6.631** | **<.001** | **.143, .265** |
| **Positive Surprised** | **.088** | **.042** | **162.642** | **2.096** | **.038** | **.005, .170** |
| **Time (linear)** | **.129** | **.007** | **265.246** | **18.084** | **<.001** | **.115, .143** |
| **Time (quadratic)** | **-.008** | **.001** | **510.659** | **-15.502** | **<.001** | **-.009, -.007** |
| **Happy * Time** | **-.062** | **.006** | **188.119** | **-10.506** | **<.001** | **-.073, -.050** |
| **Angry * Time** | **-.059** | **.005** | **170.156** | **-12.773** | **<.001** | **-.068, -.050** |
| **Fearful * Time** | **-.050** | **.005** | **193.080** | **-10.321** | **<.001** | **-.059, -.040** |
| **Positive Surprised * Time** | **-.037** | **.006** | **174.004** | **-5.767** | **<.001** | **-.049, -.024** |

Estimate of Random Effects Covariance Parameters:

| Parameter |  | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.017** | **.004** | **3.759** | **<.001** | **.010, .028** |

Note: Bold font indicates a significant effect (p<.05).

**Table S9**. CO ROI Cluster Stay Trials Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| **Intercept** | **1, 57.877** | **74.890** | **<.001** |
| **Emotion** | **4, 220.562** | **6.794** | **<.001** |
| Time (linear) | 1, 687.945 | 1.860 | .173 |
| **Time (quadratic)** | **1, 684.896** | **42.716** | **<.001** |
| **Emotion * Time** | **4, 281.603** | **16.205** | **<.001** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept** | **.108** | **.019** | **64.923** | **5.671** | **<.001** | **.070, .147** |
| **Happy** | **.037** | **.014** | **273.689** | **2.680** | **.008** | **.010, .064** |
| **Angry** | **.051** | **.014** | **333.131** | **3.681** | **<.001** | **.024, .079** |
| **Fearful** | **.031** | **.013** | **301.785** | **2.452** | **.015** | **.006, .056** |
| **Positive Surprised** | **.137** | **.032** | **90.230** | **4.306** | **<.001** | **.074, .201** |
| **Time (linear)** | **.009** | **.003** | **528.267** | **2.631** | **.009** | **.002, .015** |
| **Time (quadratic)** | **-.002** | **.000** | **684.896** | **-6.536** | **<.001** | **-.002, -.001** |
| **Happy * Time** | **-.014** | **.002** | **349.255** | **-6.249** | **<.001** | **-.018, -.009** |
| **Angry * Time** | **-.012** | **.002** | **384.360** | **-5.330** | **<.001** | **-.016, -.007** |
| **Fearful * Time** | **-.009** | **.002** | **352.404** | **-4.172** | **<.001** | **-.013, -.005** |
| **Positive Surprised * Time** | **.012** | **.005** | **114.409** | **2.520** | **.013** | **.003, .022** |

Estimate of Random Effects Covariance Parameters:

| Parameter |  | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.008** | **.002** | **3.858** | **<.001** | **.005, .013** |

Note: Bold font indicates a significant effect (p<.05).

**Figure S2**. Estimated HRF curves in the L FP and R FP clusters for each emotion based on whether a participant switched their behavioral response from the noise to the reveal phase or stayed with the same response. Labels refer to the initial response made during the noise phase.

**Table S10**. L FP ROI Cluster Switch Trials Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| **Intercept** | **1, 63.187** | **6.893** | **.011** |
| **Emotion** | **4, 175.651** | **7.055** | **<.001** |
| **Time (linear)** | **1, 456.623** | **54.569** | **<.001** |
| **Time (quadratic)** | **1, 530.513** | **79.901** | **<.001** |
| **Emotion * Time** | **4, 227.527** | **10.129** | **<.001** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| Intercept | .007 | .038 | 116.587 | .169 | .866 | -.070, .083 |
| **Happy** | **.107** | **.034** | **181.715** | **3.178** | **.002** | **.041, .174** |
| **Angry** | **.144** | **.032** | **187.751** | **4.455** | **<.001** | **.080, .208** |
| **Fearful** | **.089** | **.032** | **194.302** | **2.774** | **.006** | **.026, .152** |
| Positive Surprised | .023 | .035 | 187.727 | .638 | .524 | -.047, .092 |
| **Time (linear)** | **.064** | **.007** | **324.373** | **9.110** | **<.001** | **.050, .077** |
| **Time (quadratic)** | **-.004** | **.000** | **530.513** | **-8.939** | **<.001** | **-.005, -.003** |
| **Happy * Time** | **-.027** | **.005** | **206.814** | **-5.189** | **<.001** | **-.038, -.017** |
| **Angry * Time** | **-.028** | **.005** | **205.207** | **-5.996** | **<.001** | **-.037, -.019** |
| **Fearful * Time** | **-.020** | **.004** | **190.365** | **-4.375** | **<.001** | **-.029, -.011** |
| **Positive Surprised * Time** | **-.020** | **.005** | **196.719** | **-3.742** | **<.001** | **-.030, -.009** |

Estimate of Random Effects Covariance Parameters:

| Parameter | | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.019** | **.005** | **3.792** | **<.001** | **.011, .032** |

Note: Bold font indicates a significant effect (p<.05).

**Table S11**. L FP ROI Cluster Stay Trials Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| **Intercept** | **1, 43.474** | **26.381** | **<.001** |
| **Emotion** | **4, 249.565** | **5.998** | **<.001** |
| **Time (linear)** | **1, 645.773** | **10.403** | **.001** |
| **Time (quadratic)** | **1, 708.775** | **60.634** | **<.001** |
| **Emotion * Time** | **4, 327.337** | **3.982** | **.004** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept** | **.082** | **.023** | **51.096** | **3.514** | **<.001** | **.035, .129** |
| Happy | .025 | .014 | 297.207 | 1.774 | .077 | -.003, .054 |
| **Angry** | **.034** | **.014** | **296.794** | **2.523** | **.012** | **.008, .061** |
| Fearful | -.001 | .013 | 287.483 | -.113 | .910 | -.027, .024 |
| **Positive Surprised** | **.105** | **.027** | **102.964** | **3.926** | **<.001** | **.052, .158** |
| **Time (linear)** | **.012** | **.003** | **479.340** | **3.613** | **<.001** | **.005, .018** |
| **Time (quadratic)** | **-.002** | **.000** | **708.775** | **-7.787** | **<.001** | **-.002, -.001** |
| **Happy * Time** | **-.007** | **.002** | **380.136** | **-2.986** | **.003** | **-.011, -.002** |
| **Angry * Time** | **-.005** | **.002** | **380.071** | **-2.268** | **.024** | **-.009, -.001** |
| Fearful * Time | -.002 | .002 | 396.031 | -.810 | .418 | -.006, .003 |
| Positive Surprised * Time | .004 | .004 | 125.697 | 1.036 | .302 | -.004, .012 |

Estimate of Random Effects Covariance Parameters:

| Parameter | | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.013** | **.003** | **3.880** | **<.001** | **.008, .022** |

Note: Bold font indicates a significant effect (p<.05).

**Table S12**. R FP ROI Cluster Switch Trials Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| **Intercept** | **1, 75.156** | **4.379** | **.040** |
| **Emotion** | **4, 189.618** | **12.624** | **<.001** |
| **Time (linear)** | **1, 429.312** | **250.535** | **<.001** |
| **Time (quadratic)** | **1, 511.999** | **215.660** | **<.001** |
| **Emotion * Time** | **4, 240.188** | **10.233** | **<.001** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept** | **-.162** | **.037** | **122.624** | **-4.336** | **<.001** | **-.235, -.088** |
| **Happy** | **.109** | **.038** | **171.729** | **2.866** | **.005** | **.034, .185** |
| **Angry** | **.199** | **.030** | **186.158** | **6.514** | **<.001** | **.138, .259** |
| **Fearful** | **.148** | **.031** | **201.777** | **4.760** | **<.001** | **.087, .210** |
| Positive Surprised | .046 | .041 | 146.697 | 1.099 | .274 | -.036, .127 |
| **Time (linear)** | **.125** | **.008** | **262.559** | **16.492** | **<.001** | **.110, .140** |
| **Time (quadratic)** | **-.008** | **.001** | **511.999** | **-14.685** | **<.001** | **-.009, -.007** |
| **Happy * Time** | **-.024** | **.006** | **196.352** | **-3.877** | **<.001** | **-.036, -.012** |
| **Angry * Time** | **-.031** | **.005** | **219.684** | **-6.239** | **<.001** | **-.041, -.021** |
| **Fearful * Time** | **-.025** | **.005** | **220.415** | **-5.080** | **<.001** | **-.035, -.015** |
| **Positive Surprised * Time** | **-.022** | **.006** | **169.107** | **-3.598** | **<.001** | **-.035, -.010** |

Estimate of Random Effects Covariance Parameters:

| Parameter | | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.016** | **.004** | **3.759** | **<.001** | **.010, .028** |

Note: Bold font indicates a significant effect (p<.05).

**Table S13**. R FP ROI Cluster Stay Trials Mixed Model.

Tests of Fixed Effects:

|  | df | F | p |
|---|---|---|---|
| Intercept | 1, 41.488 | .245 | .623 |
| **Emotion** | **4, 206.444** | **6.282** | **<.001** |
| **Time (linear)** | **1, 633.952** | **421.496** | **<.001** |
| **Time (quadratic)** | **1, 692.027** | **499.357** | **<.001** |
| **Emotion * Time** | **4, 318.821** | **15.443** | **<.001** |

Estimates of Fixed Effects:

| Parameter | Estimate | SE | df | t | p | 95% CI |
|---|---|---|---|---|---|---|
| Intercept | -.010 | .024 | 49.353 | -.428 | .671 | -.058, .038 |
| **Happy** | **.068** | **.014** | **269.800** | **4.703** | **<.001** | **.040, .097** |
| Angry | .012 | .013 | 280.765 | .881 | .379 | -.015, .038 |
| Fearful | .020 | .013 | 277.779 | 1.474 | .142 | -.007, .046 |
| Positive Surprised | .009 | .026 | 72.178 | .331 | .741 | -.043, .060 |
| **Time (linear)** | **.064** | **.003** | **448.517** | **18.476** | **<.001** | **.057, .071** |
| **Time (quadratic)** | **-.006** | **.000** | **692.027** | **-22.346** | **<.001** | **-.007, -.005** |
| **Happy * Time** | **-.009** | **.003** | **402.263** | **-3.630** | **<.001** | **-.014, -.004** |
| Angry * Time | -.001 | .002 | 389.610 | -.240 | .811 | -.005, .004 |
| Fearful * Time | -.002 | .002 | 385.648 | -.798 | .425 | -.006, .003 |
| **Positive Surprised * Time** | **.023** | **.004** | **118.870** | **5.433** | **<.001** | **.015, .031** |

Estimate of Random Effects Covariance Parameters:

| Parameter | | Estimate | SE | Wald Z | p | 95% CI |
|---|---|---|---|---|---|---|
| **Intercept (subject)** | **Variance** | **.014** | **.004** | **3.879** | **<.001** | **.009, .024** |

Note: Bold font indicates a significant effect (p<.05).

**Figure S3**. Estimated HRF curves for each emotional expression by response frame for the L FP and R FP ROI clusters. Averaged responses across participants according to the noise frame (2-5) during which a behavioral response was made for correct clear valence trials (happy, angry, fearful) and ambiguous (surprised) trials for negative and positive responses. Vertical lines at the top of each plot indicate the identified peak(s) for each curve.

**Table S14**. Peak times (seconds) for each ROI Cluster by Emotional Expression and Frame

| Cluster | Emotion | Frame 2 | Frame 3 | Frame 4 | Frame 5 |
|---|---|---|---|---|---|
| CO | Happy | 9.7 | 11.3 | 13.0 | 14.9 |
| | Angry | 9.5 | 11.4 | 13.0 | 15.8 |
| | Fearful | 8.9 | 11.3 | 13.2 | 16.0 |
| | Negative Surprised | 9.6, 17.7 | 11.1, 17.6 | 13.1, 17.4 | 15.9 |
| | Positive Surprised | 10.0, 18.1 | 12.4, 17.2 | 16.0 | 18.0 |
| L FP | Happy | --- | 11.2 | 13.1 | 8.6, 15.0 |
| | Angry | 5.9, 9.1 | 11.2 | 13.2 | 15.9 |
| | Fearful | 8.6 | 11.3 | 13.3 | 16.2 |
| | Negative Surprised | 9.5, 17.8 | 11.0 | 13.4 | --- |
| | Positive Surprised | 8.9, 13.1, 17.4 | 11.5, 18.0 | 14.2 | 16.2, 19.8 |
| R FP | Happy | 11.0, 17.7 | 12.6, 16.2 | 14.0 | 8.1, 16.2 |
| | Angry | 9.3 | 13.3, 16.8 | 14.4 | 16.6 |
| | Fearful | 13.0, 17.2 | 12.2 | 14.1 | 16.6 |
| | Negative Surprised | 10.0, 14.6, 17.6 | 13.0, 17.4 | 14.2, 17.3 | 17.7 |
| | Positive Surprised | 8.3, 17.9 | 11.5, 18.1 | 17.6 | 18.0 |

**References**

Anwyl-Irvine, A. L., Massonnié, J., Flitton, A., Kirkham, N., & Evershed, J. K. (2020). Gorilla

    in our midst: An online behavioral experiment builder. *Behavior Research Methods*,

    *52*(1), 388–407. https://doi.org/10.3758/s13428-019-01237-x