Boosting the toolbox for live imaging of translation

Maelle Bellec[1,2], Ruoyu Chen[3,4], Jana Dhayni[5], Antonello Trullo[1], Damien Avinens[6], Hussein Karaki[5], Flavia Mazzarda[5] Helene Lenden-Hasse[1], Cyril Favard[6], Ruth Lehmann[4], Edouard Bertrand[*5], Mounia Lagha[1], Jeremy Dufourt[*1,6]

Affiliations

[1]Institut de Génétique Moléculaire de Montpellier, Univ Montpellier, CNRS, Montpellier, France.

[2]Department of Developmental Genetics, Max Planck Institute for Heart and Lung Research, Bad Nauheim, Germany.

[3]Vilcek Institute of Graduate Studies, NYU School of Medicine, New York, United States.

[4]Whitehead Institute for Biomedical Research and Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA.

[5]Institut de Génétique Humaine, University of Montpellier, CNRS, 34396 Montpellier, France.

[6]Institut de Recherche en Infectiologie de Montpellier, CNRS UMR 9004, University of Montpellier, 1919 Route de Mende, Montpellier, 34293, Cedex 5, France.

* Correspondence to: jeremy.dufourt@irim.cnrs.fr

* Correspondence to: edouard.bertrand@igh.cnrs.fr

Abstract

Live imaging of translation based on a tag recognition by a single chain antibody is a powerful technique to assess translation regulation in living cells. However, especially in a multicellular organism, this approach is challenging and requires optimization in terms of expression level and detection sensitivity of the system. Here, we improved existing fluorescent tools and developed new ones to image and quantify nascent translation in the living *Drosophila* embryo and in mammalian cells. We tested and characterized five different Green Fluorescent Protein variants fused to the single chain fragment variable (scFv) and uncover photobleaching, aggregation and intensity disparities. Using different strengths of germline and somatic drivers, we determined that the availability of the scFv is critical in order to detect translation throughout development. We introduced a new translation imaging method based on a nanobody/tag system named ALFA-array, allowing the sensitive and simultaneous detection of the translation of several distinct mRNA species. Finally, we developed a largely improved RNA imaging system based on an MCP-tdStaygold fusion.

Introduction

During development, multipotent cells progressively acquire specific fates leading to the formation of a variety of tissues. This process relies on a spatio-temporally regulated gene expression program, contributing to the reliability and reproducibility of gene expression patterns to generate the final body plan. In that process, the regulation of mRNA translation is a key step since the dynamics of translation in space and time critically affects the outcome of gene expression. Most approaches to study the regulation of translation in whole organisms rely on a bulk population of cells with poor spatial and temporal information. Consequently, the precision and heterogeneity in the regulation of translation at the cellular and molecular level remains poorly investigated. Live imaging methods of translation have been developed in cultured cells[1–5] since 2016 and recently introduced in *Drosophila*[6–8]. These methods are based on the fluorescent labeling of nascent peptide chains. The labeling system is made of two components, a genetically-encoded antibody such as a single-chain variable fragment antibody (scFv) or Fab fragment[9–11] fused to a fluorescent protein (FP; referred to as the detector), and a N-terminal peptide epitope inserted in multiple copies in frame with the gene of interest[12]. These live imaging approaches made it possible to visualize translation up to the single-molecule scale in cultured cells but also in an entire organism[7,8]. Such technological breakthroughs provided important insights in our understanding of gene regulation through translation and, for instance, revealed the existence of motor-driven polysome transport and spatial heterogeneity of translation at the subcellular level[1,7,13–17].

The discovery of the *Aequorea victoria* green-fluorescent protein[18,19] and especially its optimized version mEGFP[20] have revolutionized imaging of biological processes[21]. Since then, several new FPs have been and continue to be discovered and

engineered[22,23]. The constant race for better FPs generally implies the optimization for

fast folding, pH resistance as well as improvement of photo-physics[24].

Introduction of the bacteriophage-derived MS2 RNA binding sites, which are

recognized by MS2 Coat Protein (MCP) fused to FP, provided a technique to visualize

transcription in living cell. This technological advance has been subjected to numerous

improvements from its initial development in 1998[25]. These improvements include

increased sensitivity down to single molecules[26], implementation in a living organism[27],

the generation of new stem loops to visualize single mRNAs at a high temporal

resolution[7,28], artefact-free RNA[29] and double mRNA species imaging with an

orthologous system[30]. However, several limitations exist with the present tools that

hinder quantitative imaging. For example, protein aggregation has been observed for

some systems derived from MS2 Coat Protein fused to GFP[31–33] and for scFv fused to

specific GFP variants[34]. Furthermore, the expression level of the detector is critical to

ensure accurate detection, particularly for translation, as most of the time one mRNA

undergoes multiple rounds of translation. However, no such optimization has been

made for the SunTag system in the context of a developing organism, although photo-

physical properties of molecules can be variable between different systems (i.e. *in vitro*

experiments, cell culture, tissues, organisms…)[24,35,36].

Here, we tested five different GFP variants for their brightness, photobleaching and

propensity to aggregation in fixed and live *Drosophila* embryos. We also generated

inducible scFv-FP constructs that, when combined with different drivers, extend the

use of the SunTag system to a wider range of genes and biological questions.

Furthermore, we developed an orthologous method, called ALFA-array, to visualize

nascent translation in *Drosophila* embryo and mammalian cells. This system is based

on a small and soluble nanobody and we show its usefulness by combining it with the

SunTag to visualize translation of two different mRNA species in *Drosophila* embryos.

Results

New fluorescent protein fused to scFv to detect nascent translation.

The SunTag method is a bipartite system based on an antibody-derived single chain

variable fragment (scFv) detector fused to a fluorescent protein (FP; the detector)

which binds a peptide named Supernova Tag (suntag) derived from the yeast GCN4

protein and are experimentally fused in multiple copies at the N-terminus of the gene

of interest[34] (Figure 1a). To improve the system, we created new scFvs with different

green fluorescent proteins under the control of *nanos* (*nos*) enhancer-promoter (EPr).

This EPr combination drives expression in the *Drosophila* germline[37–40] allowing the

maternal deposition of mature scFv-FP into laid eggs (Figure S1A). We generated a

set of four different constructs, in addition to the previously published scFv fused to the

superfolder GFP (sfGFP)[41], named scFv-sfGFP[7]. The new generation of green

fluorescent protein variants includes: i) the monomeric superfolder GFP2 (msGFP2)[42],

ii) the mGreenLantern (GL)[43], iii) mNeonGreen (NG)[44] and iv) the monomeric *A. victoria*

fluorescent protein 1 (mAvic1)[22] (Figure 1b, upper panel). Live imaging of these five

scFv-FPs during embryonic development until the end of nuclear cycle 14 (n.c. 14)

showed that all the scFv-FPs are well expressed, localize mainly in the nucleus, due

to the presence of a nuclear localization signal, and do not form aggregates (Figure

S1b and movie Sup 1-5). Quantification of the fluorescent intensity during the first forty

minutes of n.c.14, showed that the apparent brightness of these four new scFv was

similar to the scFv-sfGFP, while scFv-mAvic1 produced brighter signals (Figure S1c)

and scFv-GL was prone to photobleaching (Figure S1c). We confirmed that the

distribution of scFv-FP was homogenous in the embryo and dots/aggregates were not detected by light-sheet imaging of the two brightest scFv-FPs, scFv-mAvic1 and scFv-msGFP2 (Figure S1d and movie Sup 6-7).

We next tested the capacity of these new scFv-FPs to detect mRNA in translation in *Drosophila* embryos. We crossed the different scFv-FP lines into the background of a *twist* gene endogenously tagged with MS2 and suntag (*twi_suntag_MS2_CRISPR*) (Figure 1b, lower panel) and performed single-molecule Fluorescent *in situ* Hybridization (smFISH) on fixed embryos to detect the tagged mRNAs. Actively translated mRNAs (polysomes) were then visualized as co-localized GFP and mRNA spots. Imaging the border of *twist* mRNA transcription pattern showed that the translation signal was present only in the described pattern of *twist* expression (mesoderm)[45] (Figure 1c), supporting that the signal is specific. For all scFv-FPs, we observed a clear colocalization between mRNA (red signal) and polysomes (green signal) in addition to mRNA molecules that were not in translation (Figure S1e). Notably, a fainter signal was detected in fixed samples for polysomes labeled with scFv-mAvic1 at this stage of embryogenesis, as compared to its high apparent brightness in live samples (Figure S1f), suggesting that mAvic1 is more sensitive to the fixation/smFISH protocol. Immuno-smFISH with an antibody against GFP did not improve the signal but increased the background. Surprisingly, we found that staining with only a secondary antibody could label nascent foci of translation (Figure S2a), and only when the scFv-FP was present (Figure S2b). This result suggested that secondary antibodies alone could bind scFv-FP and that experiments using both simultaneously should be avoided. We then looked at the behavior of these scFv-FPs in live embryos in the presence of *twi_suntag_MS2_CRISPR*. We could observe bright green fluorescent dots in movement within the mesoderm (Figure 1d, movie sup 8-12). To

characterize the detection capacity of each FP in live embryos, we measured the mean intensity of all detected spots for 4 min during n.c. 14 with a high temporal resolution (Figure 1e, movie sup 8-12). The translation foci detected by scFv-mAvic1, scFv-sfGFP and scFv-msGFP2 showed a stable signal compared to scFv-NG and scFv-GL (Figure 1e). Collectively, these results show that the four new scFv-FPs are able to bind nascent suntag peptides and therefore detect mRNA in translation in live and fixed samples.

New scFv-FP prevents aggregate formation

To show whether scFv-FPs expressed under the maternal *nos* EPr could detect translation during embryogenesis, at least until gastrulation, we imaged fixed embryos at the gastrulation stage. We observed tagged Twist protein localized to nuclei in the ventral gastrulation furrow region, consistent with Twist being a mesodermal transcription factor (Figure S2c, upper panel). To test, whether the presence of a NLS in the scFv-FP construct might be responsible for this nuclear localization, we imaged the localization of the Insulin-like peptide 4 (Ilp4), an excreted protein expressed in the mesoderm[46]. We used the tagged Ilp4 with suntag and observed scFv containing the NLS in the cytoplasm (Figure S2c, lower panel). Thus, at least in these cases, the presence of a NLS in the scFv-FP construct appears not to significantly interfere with the native localization of tagged proteins.

Strikingly, at the gastrulation stage, bright GFP spots were observed only when using scFv-sfGFP in *twi_suntag_MS2_CRISPR* embryos but not with the other scFv-FPs (Figure S3a). As only the FP sequences are different between these constructs, this discrepancy might be due to a difference in protein stability, i.e. scFv-sfGFP being more stable than the other variants, or being more prone to aggregation. To distinguish

between these possibilities, we used smFISH to label *twi_suntag_MS2_CRISPR*

mRNA in the presence of the different scFv-FPs to determine whether the bright GFP

spots observed during gastrulation co-localized with mRNA and represent true

translation spots. With scFv-sfGFP we saw large green spots that were not co-localized

with mRNA (Figure S3b upper left panels, white arrowhead). We did not observe these

large spots with any of the four new scFv-FPs (Figure S3b) suggesting an aggregation

of the scFv-sfGFP in the *Drosophila* embryo at the gastrulation stage in the presence

of *twi_suntag_MS2_CRISPR*.

We next asked whether aggregation formation was specific for the tagged *twi* in

developing embryos or could also occur in cultured cells. We expressed scFv-sfGFP

and scFv-msGFP2 in *Drosophila* S2R+ cells along with a SunTag-luciferase reporter

(SunTagRLuc-MS2; see methods and Figure S3c). The SunTagRLuc-MS2 plasmid

was co-transfected with an scFv-FP plasmid as well as an MCP-Halotag plasmid to

track single mRNA molecules and their translation during live-cell imaging. As

expected, scFv-FPs were able to label polysomes by forming GFP foci that co-localized

with mRNAs (Figure S3d). With the scFv-sfGFP, however, we frequently observed

GFP aggregates which were not associated with any mRNA molecules. These

aggregates were morphologically distinct from translating foci and were more abundant

and bigger than the GFP aggregates observed in gastrulating embryos (Figure S3d-

e). This result suggests an aggregation of the SunTag peptides bound by the scFv-

sfGFP after being translated and released from polysomes. Although few aggregates

were also present in cells expressing scFv-msGFP2 with the SunTagRLuc-MS2

reporter (Figure S3d), this occurred with a significantly lower frequency (~55% of cells

without aggregates compared to ~15% using scFv-sfGFP) and smaller aggregate size,

suggesting that msGFP2 is less prone to aggregate formation.

These results support that the scFv-FPs containing newer generation monomeric FPs

are more suitable for translation analysis, as they are less likely to form aggregates

compared to sfGFP which is able to form weak dimers[42,47].

Monitoring translational arrest during *Drosophila* embryogenesis

The low amount of translation spots observed during gastrulation suggested that *twi*

translation maybe have ceased at this stage of development (Figure S3a-b). However,

a translational arrest of *twist* mRNAs has not been described previously. This

unexpected observation led us to investigate whether *twi* transcripts were undergoing

a translational arrest during n.c.14 (Figure 2a) using the four new scFv-FPs.

Interestingly, we found that at mid-late n.c. 14 (Figure 2a), the

twi_suntag_MS2_CRISPR RNA was mostly translated at the border of the mesoderm

(Figure 2b and S4a-c). Using scFv-msGFP2 fluorescence intensity quantification, the

percentage of mRNA in translation was slightly lower in the center as compared to the

border of the mesoderm (44 ± 4% at the center and 61 ±1% at the border see methods).

Strikingly, the polysomes intensities were much lower in the center compared to the

mesoderm border, suggesting reduced translation (Figure 2b and S4a-c). A

translational arrest was recently described for *Hunchback* (*Hb*) using the SunTag

system in the *Drosophila* embryo[8]. In this study the authors showed that after being

translated in the entire Hb domain (anterior half of the embryo), *hb* translation became

restricted to the border of its expression pattern, similarly to what we observed for *twist*.

At least two models can explain this result: i) *twi* translation is actively repressed at first

in regions where the RNA and protein expression is highest (as suggested for *hb*[8]), or

ii) the amount of scFv-FP becomes limiting during mid-late n.c. 14 at first in regions

where the RNA and protein expression is highest and prevents the detection of

translation (Figure 2c upper panel). To test these hypotheses, we used the *twi_suntag_transgene*, which exhibits a stochastic expression pattern, belated transcriptional activation[7] and thus lower mRNA level as compared to *twi_suntag_MS2_CRISPR.* If *twi* translation is actively repressed, a similar translational arrest should be observed for the *twi_suntag_transgene*. In contrast, if the amount of scFv-FP is limiting, translation of the transgene should be detectable over a longer period of time (Figure 2c). Using live imaging, we observed translation spots of the *twi-suntag_transgene* at a stage when *twi_suntag_MS2_CRISPR* translation spots vanished (Figure 2d, S4d and Movie S13-15). The translation dots of the *twi_suntag_transgene* colocalized with mRNA dots in late n.c. 14 and gastrulation stage as confirmed with smFISH (Figure S4e). These results suggest that the amount of scFv-FP might be limiting when *twi* translation is monitored from its endogenous locus (*twi_suntag_MS2_CRISPR* allele)*,* causing a seeming translational arrest at gastrulation.

To measure the availability of free scFv-FP during n.c.14 we performed Fluorescence Correlation Spectroscopy (FCS) in the mesoderm of embryos expressing scFv-msGFP2 with or without *twi_suntag_MS2_CRISPR.* In the absence of suntag translation (i.e. without *twi_suntag_MS2_CRISPR),* only free scFv-msGFP2 were detected and sufficient fluorescence signal was recovered to fit FCS data until the mid-late n.c.14 (Figure S4f upper panel and S4g) representing on average ~30 free scFv-msGFP2 available in the confocal volume in absence of translation (Figure S4g left panel). In contrast, in the presence of suntag translation (i.e. with *twi_suntag_MS2_CRISPR),* fluorescent signals were sufficient to fit FCS data only until early-mid n.c.14 (Figure S4f lower panel and g). These results demonstrate that from the mid-late n.c.14 stage, there is an insufficient amount of free scFv-msGFP2

available to detect translation. We next used an antibody against the suntag peptides

(anti-GCN4) in the absence of scFv-FP and saw translation of

twi_suntag_MS2_CRISPR mRNAs at mid-late n.c. 14 and gastrulation stage (Figure

2e and S5a). Moreover, the percentage of mRNA in translation was homogenous

within the mesoderm (72 ±4% at the border and 68 ±2% at the center) and polysome

intensity between center and border was similar (Figure 2e), in contrast to the results

obtained with scFv-msGFP2 (Figure 2b). Altogether these results demonstrate that

there is no translational arrest of *twi* mRNAs during n.c.14. Instead, the amount of

scFv-FP expressed under the maternally active *nos* EPr becomes insufficient to detect

translation during gastrulation, when a large pool of tagged mRNAs is translated. Our

experiments suggest that anti-GCN4 antibodies should be used as a control to

distinguish between true endogenous repression of translation and mere

experimentally-dependent depletion of the labelling reagent.

Inducible scFv-FP allows imaging of translation during embryogenesis

To overcome the limited amount of scFv-FP and allow the sustained imaging of *twi*

translation from egg laying to gastrulation, we used the UAS/Gal4 system to increase

the expression of scFv-FP during oogenesis. We generated four lines with the different

scFv-FPs downstream of a *UASp* promoter (Figure 3a and S6a) and crossed these

with the maternal *nos-Gal4* driver (Figure 3b). No aggregation and a higher scFv-FP

signal were observed in live embryos (Figure S6a, Movie S16-19 and see methods).

Next, we asked if using this system would allow detection of *twi* translation until late

n.c. 14 and gastrulation stage. In *twi_suntag_MS2_CRISPR* embryos with maternally

provided *nos-Gal4>UASp-scFv-msGFP2* we saw that *twi* mRNAs were translated

throughout n.c.14 (Figure 3c), even in the most ventral part of the embryo, at the center

of the *twi* domain of expression (Figure 3c). However, we were not able to detect small

dots of scFv-FP, most likely corresponding to individual mature proteins, probably due

to the lower signal-to-noise ratio generated by the increased levels of *scFv-msGFP2*.

We also observed translation dots at the gastrulation stage in fixed

twi_suntag_MS2_CRISPR embryos with *nos-Gal4>UASp-scFv-msGFP2* (Figure 3d

left panels) but not with *EPr-scFv-msGFP2* (Figure 3d right panels). Similar results

were observed with live imaging, where *twi_suntag_MS2_CRISPR* mRNAs were

translated until gastrulation with *nos-Gal4>UASp-scFv-msGFP2* (Figure 3e and Movie

S20) but not with *EPr-scFv-msGFP2.* Control embryos lacking

twi_suntag_MS2_CRISPR mRNAs did not show any translation spot at any stage

(Figure 3e and Movie S20).

To measure the availability of free scFv-FP when using the *nos-Gal4>UASp-scFv-*

msGFP2 expression system, we again performed FCS in the mesoderm of embryos*.*

In absence of suntag translation (i.e. without *twi_suntag_MS2_CRISPR*), sufficient free

scFv-msGFP2 signal was recovered to fit FCS data until very late n.c.14/gastrulation

stages (Figure S6b, upper panel) representing on average ~650 free scFv-msGFP2

available in the confocal volume in absence of translation (Figure S6c, left panel). In

the presence of suntag translation (i.e. with *twi_suntag_MS2_CRISPR)*, sufficient

fluorescent signals were also recovered to fit the FCS data until very late

n.c.14/gastrulation stages (Figure S6b lower panel). The brightness and number of

molecules evolve from a majority of fast moving and lacklustre molecules at early-mid

n.c.14, corresponding to free scFv-msGFP2, to a majority of slow moving and brighter

molecules corresponding most probably to polysomes at late n.c.14/gastrulation

stages (Figure S6c). Altogether, these results demonstrate that the use of *nos-*

Gal4>UASp-scFv-msGFP2 allows detection of translation until gastrulation. The *UASp*

promoter drives expression in the germline but is less efficient in somatic tissues[48]. On the contrary, the *UASt* containing *HSP70* promoter drives expression in the somatic tissue[49]. Recently, the *UASt* promoter has been modified to obtain a promoter, called *UASz*, to drive expression in both somatic and germline tissues[50]. Due to the high amount of free scFv observed with the *UASp-FP* lines, we generated a new promoter derived from the *UASz*, in which we removed the translational enhancer *IVS-Syn21*[50] to prevent excessive expression of free scFv-FP (Figure S7a, b). This new promoter named *UASy* was fused to the *scFv-msGFP2* and crossed with the *nos-Gal4* driver line. As all the other constructs generated in this study, *nos-Gal4>UASy-scFv-msGFP2* alone did not lead to aggregates during early embryogenesis (Figure S7b and Movie S21). Unfortunately, the translation of *twi_suntag_MS2_CRISPR* mRNA at the end of n.c. 14 showed a similar pattern – although less pronounced – as with *scFv-msGFP2* where translation was mostly present at the border of *twi* pattern (Figure S7c), likely due to the weaker expression of the *UASy* compared to *UASp* promoters and subsequent depletion of scFv in the central mesoderm. Maternal expression of the scFv-FP constructs limits our ability to observe translational dynamics at later stages in development. To overcome this, we crossed *UASy-scFv-msGFP2* with an early zygotic *nullo-Gal4* driver to express throughout the embryo. We observed scFv-msGFP2 expression at the beginning of gastrulation (Movie S22) and a strong and homogeneous expression in somatic tissues of later embryos (Figure S7d). We conclude that the *nos-Gal4>UASp-scFv-FP* represents a good tool to image translation from early embryogenesis to gastrulation stage and that the new *UASy-scFv-msGFP2* enables the imaging of translation in somatic cells at later stages.

Development of the ALFA-array system to monitor translation in *Drosophila*

After having optimized the scFv-FP-based detector, we sought to generate an

optimized orthologous system to the SunTag to detect the translation of two different

RNA species in *Drosophila*. To this end, we searched for a highly specific

nanobody/tag pair and decided to use the recently developed ALFA-Tag[51]. The ALFA-

Tag is a small peptide that harbors several advantages: its synthetic sequence is

absent from the proteome of model organisms, its sequence is smaller than that of the

SunTag sequence (15 amino acid compared to 19, respectively Figure 4a), it is

monovalent, hydrophilic and was effective in *Drosophila* for protein manipulation[52].

Finally, the nanobody anti-ALFA detector (NB-ALFA) is smaller than the scFv (Figure

4b), is quite soluble and displays a very strong affinity for the ALFA-Tag (measured at

26 pM)[51]. Taking advantage of these properties, we multimerized the ALFA-Tag

sequences to create a 12x and 32x_ALFA-array. Each ALFA-tag sequence was

flanked by a proline to reduce potential influence of neighboring secondary structures,

and a spacer of 5 amino acid (GSGSG) to minimize steric hindrance of neighboring

peptide binding sites (Figure 4a, c). The NB-ALFA was fused to the msGFP2 and to

the GB1 solubilization tag (also used for the scFv) to avoid potential aggregation

caused by the multimerization of ALFA-tag sequences. Finally, we generated a

Drosophila line with the *NB-ALFA-msGFP2* under the control of *nos* EPr (*NB-ALFA-*

msGFP2). The system was also tested in mammalian cells (see below).

To study translation in *Drosophila* embryos using the 32x_ALFA-array system, we

chose to tag the *Ilp4* gene, which codes for a small protein. We reasoned that if we

could visualize the translation of a very short transcript/protein, the tool would work

even better for longer ones. A 32x_ALFA-array was inserted between the *Ilp4* initiation

codon and *Ilp4* gene body, appended with *Ilp4* UTRs and *Ilp4* EPr (*Ilp4_32x_ALFA-*

array, Figure 4d). With immuno-smFISH we found that the *Ilp4_32x_ALFA-array* was

expressed in the mesoderm of early *Drosophila* embryos (Figure S8a), consistent with previous reports for *Ilp4*[46]. The anti-ALFA formed bright dots specifically colocalizing with *Ilp4_32x_ALFA-array* mRNAs and thus representing foci of translation (Figure S8b-c). Importantly, the *Ilp4_32x_ALFA-array* was not recognized by the anti-GCN4 antibody (Figure S8a).

We first confirmed by live imaging that the *NB-ALFA-msGFP2* strain was expressed during embryonic development until the end of n.c.14, that it was mainly localized to the nucleus and that it did not form any aggregates (Figure S8d and Movie S23). The *Ilp4_32x_ALFA-array* was crossed with *NB-ALFA*-msGFP2 and the resulting embryos showed bright GFP spots overlapping with *Ilp4_32x_ALFA-array* mRNAs (Figure 4d), showing the ability of NB-ALFA-msGFP2 to bind the nascent 32x_ALFA-array *in vivo.* Next, we monitored translation in live embryos and observed bright GFP dots appearing on the ventral side of the embryo (Figure 4e and Movie S24). The signal was restricted to the mesoderm during n.c.14 (Figure S8e and Movie S25) and became more restricted to the ventral furrow during gastrulation (Figure S8f and Movie S25), demonstrating the specificity of the signals. Thus, the *Ilp4_32x_ALFA-array* – NB-ALFA-msGFP2 system is a useful alternative tool to detect mRNA translation during early embryogenesis.

Additionally, staining with a secondary antibody alone could label the translation foci only when the NB-ALFA-msGFP2 is present (Figure S9a-b). This was also observed with the scFv-FPs (see above Figure S2a-b), suggesting that secondary antibodies can directly bind both the NB-ALFA-msGFP2 and the scFv-FP (Figure S2a-b and Figure S9a-b). Knowing that the only common part between NB-ALFA and scFv is the GB1 solubility domain, we tested whether this domain was responsible for binding secondary antibodies. We removed the GB1 from the scFv-msGFP2 and NB-ALFA-

msGFP2, expressed them in *Drosophila* S2R+ cells and performed Western-blot

analyses (Figure S9c). All the scFv and NB-ALFA constructs (with and without the

presence of the GB1) could be detected by an anti-GFP antibody in combination with

a suitable secondary antibody. However, when using secondary antibodies alone, the

constructs with GB1 were still detected, while no band was observed with the

constructs without GB1 (Figure S9c). These results demonstrate the direct binding of

the GB1 solubilization domain by secondary antibodies.

To test if we could use the ALFA-array system in combination with the SunTag to image

the translation of two distinct mRNA species in the same embryo, the *Ilp4_32x_ALFA-*

array was crossed with the *twi_suntag_transgene* (Figure 4f). Using immuno-smFISH

in n.c.14 embryos, we observed the two correct expression patterns of Twi and Ilp4,

including in the posterior region where only Twi is present (Figure S10a). With higher

resolution images, we could visualize single molecules of *Ilp4* mRNA in translation well

distinct from *twi* mRNA translation dots (Figure 4g). Furthermore, we were able to

distinguish the localization of the Ilp4 protein in the cytoplasm from the Twi protein in

the nucleus (Figure 4h). Finally, we then combined four different transgenes, to

express scFv-mScarlet, NB-ALFA-msGFP2, *Ilp4_32x_ALFA-array* and

twi_suntag_MS2_CRISPR in the same embryo. We were able to live image

simultaneously the translation of *Ilp4* and *twi* mRNAs (Figure 4i, S10b and Movie S26)

in live embryos (as a note the scFv-mScarlet is prone to photobleaching precluding

long time imaging as described in[7]). Furthermore, we were again able to clearly

distinguish the localization of Ilp4 protein in the cytoplasm from Twi protein in the

nucleus of live embryos (Figure 4j, S10c and Movie S27). Together, these results

demonstrate that the 32x_ALFA-array can be used in conjunction with the SunTag

system to study the translation of two different mRNAs during *Drosophila* early

embryogenesis in live and fixed samples.

Development of the ALFA-array system to monitor translation in human cells

To test the performance of the ALFA-tag labeling system to image translation in a

different model, we turned to mammalian cells. We first generated HeLa cells

expressing a NB-ALFA-sfGFP recombinant nanobody. To validate the strong binding

properties of the nanobody for the ALFA tag *in vivo* (measured at 26 pM in vitro[51]) we

took advantage of the available H2B_mCherry_ALFAtag constructs[52]. Using acceptor

photobleaching Förster Resonance Energy Transfer (FRET), we observed that the NB-

ALFA-sfGFP was closely associated to the ALFA-Tag in the nuclei of live HeLa cells

(Figure S11 a and b). We then performed Fluorescence Recovery After

Photobleaching (FRAP) of the NB-ALFA-sfGFP, knowing that H2B itself has a very

slow recovery time lasting hours[53]. This experiment showed little recovery of NB-ALFA-

sfGFP in 5 minutes (Figure S11 c and d), indicating that it is stably bound to its target

in vivo.

Next, we evaluated the performance of the ALFA-array system to image translation in

HeLa cells. We inserted a 12x_ALFA-array repetition at the N-terminus of the

DYNC1H1 gene by CRISPR-mediated knock-in (Figure 5a), as previously done for the

SunTag[4]. Heterozygous clones were obtained and imaged with and without a thirty-

minute puromycin treatment to inhibit translation. In untreated cells, bright foci were

observed in the cytoplasm, which disappeared following puromycin addition (Figure

5b). In addition, smiFISH against the endogenous DYNC1H1 mRNAs revealed that the

bright ALFA-tag foci colocalized with single mRNAs (Figure 5c), confirming that these

corresponded to DYNC1H1 polysomes. The quality of the signals obtained with only

12 repeats of the ALFA-tag demonstrates the efficiency of this labeling system to image

translation. To compare more directly the ALFA-array with the SunTag system, we

used CRISPR knock-in to insert a 32x_ALFA-array repetition at the N-terminus of the

POLR2A gene, and compared the signals with those obtained with a previously

developed 32xSuntag-POLR2A cell line[4]. In both cell lines, translation spots were

visible and co-localized with POLR2A mRNAs (detected with POLR2A probes) (Figure

S11e). Only a fraction of the mRNAs had a nanobody signal, consistent with the

heterozygous nature of the cell lines. We then measured the signal of the translation

spots obtained with the NB-ALFA-sfGFP to the ones obtained with the scFv-sfGFP.

We observed that the NB-ALFA-sfGFP system yielded ~2.5 times brighter signals,

demonstrating the high performance of the system.

Finally, an optimal system to visualize RNA is very useful to image translation at the

level of single molecules[1–5]. While the MS2/MCP system provides a very nice way to

image single RNAs, it is still limited by the low brightness of the signals and its rapid

photobleaching. To improve RNA imaging, we took advantage of the recently

developed Staygold fluorescent protein, which is bright and exceptionally

photostable[54]. Staygold is an obligate dimeric fluorescent protein, and we thus made

a tandem dimer version (tdStaygold) by degenerating its sequence and separating two

monomers with a (SAGG)x5 linker. This dimeric version of Staygold was then fused to

MCP and used to image RNA in a previously developed HeLa cell line expressing an

HIV-1 reporter RNAs tagged with 128 MS2 sites[55] In this system, the MS2 tag is intronic

and enables to label nascent RNAs as well nucleoplasmic pre-mRNAs. Live cells were

imaged using a frame rate of one 3D image stack every ten seconds for up to 6 hours

(Figure 5d). Single RNAs were observed with both NLS-MCP-tdStaygold and the

previously used NLS-MCP-eGFP[55]. However, the signal bleached rapidly with NLS-

MCP-eGFP, and single RNAs were no longer detectable after 120 minutes. In contrast,

the signal showed little photobleaching with NLS-MCP-tdStaygold and single RNAs

were efficiently detected through all the movie, and kept their initial brightness (Figure

5e-f). The NLS-MCP-tdStaygold developed here thus shows largely improved

performance for RNA imaging.

Discussion

Amplification of biological signals is crucial to study many biological processes with a

high spatio-temporal resolution and single-mRNA-molecule sensitivity. Live imaging of

translation at single molecule resolution is a relatively recent technology implemented

in 2016 in tissue culture and in 2021 in multicellular organisms. In this study, we provide

a powerful toolbox to image translation in living cells and organisms. We will discuss

some of the limitations and challenges that need to be considered to obtain robust and

reliable results.

Our data show that the choice of the fluorescent protein is a key element for the signal

quality of translation imaging-based systems. The development of fluorescent proteins

is a very rapidly evolving field and FP-based tools need to be updated to optimize their

use. Often, the choice of the FP needs to be empirically defined for each cell type,

tissue or organism studied[56–61]. One important aspect in this choice is the monomericity

of the FP to prevent aggregation issues. Although the sfGFP (weak dimer) was the

standard FP for fusion with the scFv in mammalian cell culture, we found that it might

present aggregation issue in *Drosophila*. Moreover, During the revision process of our

manuscript, three monomeric versions of StayGold proteins were published[62–64], and

future optimization should consider testing these FPs for imaging translation, taking

into account the specific imaging setup[65]. Local translation of specific mRNAs is an

important biological process to control gene expression in space and time. To obtain quantitative answers it is necessary to precisely resolve the location of mRNAs and where their translation occurs. In the present experimental designs, with the orthologous scFv and the NB-ALFA, the use of secondary antibodies can confound signal interpretation as they appear to bind the scFv and NB-ALFA fusion. We demonstrate that this recognition is due to the GB1 solubilization domain sequence added to limit aggregation and is in line with a recent study in plants, which showed a strong affinity between GB1 and IgG[66]. As the GB1 is required to for the solubility of the detector, combining translation imaging with immunofluorescence, will require the use of fluorescent primary antibodies.

An important aspect in imaging live embryos or tissues is signal amplification to increase the signal-to-noise ratio. Indeed, in systems like the MS2/MCP, llama-Tag/Fluorescent protein[67] or scFv/SunTag and their derivatives, the signal appears by local specific accumulation of fluorescent detector, over a diffuse background of molecules freely diffusing in the cells. To improve the signal-to-noise ratio, development of Quenchbody (Q-body) could in theory allow background free detection of translation as well as single mRNA[68]. Currently, the signal-to-noise ratio is optimized by reducing the amount of free fluorescent detector and therefore the background. This is achieved by using weak promoters or targeting the free fluorescent detector to other cellular compartments such as the nucleus, or specific organelles for cytoplasmic analysis. However, one should be careful that the availability of these fluorescent detector does not become limiting as more and more RNAs and proteins are produced during development. As we illustrate here, the available concentration of the detector can be a major challenge in developing organisms, especially when the source of the detector is controlled by an endogenous promoter and when the amount of template

mRNA in translation is important. This limitation can also be particularly important when

a burst of translation occurs such as during viral infection[14,15]. We provide a direct

example for detector limitation during early embryogenesis. Here, early activated

genes such as *twist* are highly and rapidly transcribed and translated leading to rapid

depletion of free detector. We demonstrate the importance of distinguishing between

detector depletion and cessation of translation by validating the results with direct

immunostaining. Indeed, a simple relation shows that the minimal amount of detector

to image mRNA translation is a function of the size of the protein to be translated, the

translation speed and the kinetic constant of the detector binding to the target (k_{on}).

The highest the speed, the shortest the protein, the higher the concentration of the

detector has to be to image translation reliably. For example, in the case of *twi*

considering the upper limit of translation speed described of 35 AA.s-1[7], the minimal

number of scFv-msGFP2 needed to accurately image translation represents on

average 3 free detectors available in the confocal volume. Here we observe only ~10

times more free detectors at early-mid n.c.14 with *EPr-scFv-msGFP2*. We also show

that increasing the expression of the detector can overcome the detection problem, as

using *nos-Gal4>UASp-scFv-msGFP2* we observe on average 650 free scFv-msGFP2

available in the confocal volume at early-mid n.c.14 corresponding to ~22 times more

scFv-msGFP2 compared to no overexpression. However, it will reduce the signal-to-

noise ratio and has thus to be taken into consideration for the specific experimental

design. In order to reach conditions that can most accurately capture the dynamics of

translation, diverse *UAS* promoters together with a variety of available Gal4 drivers

should be tested depending on the tissue of interest and the expression level of the

tagged gene.

Studies on mRNA localization and local translation have speculated that different

mRNAs species may be translationally co-regulated[69–71]. To obtain independent

translational readouts of different mRNA simultaneously, one can combine orthogonal

tagging systems such as hybrid scFvs such as HA or FLAG frankenbody[72–74] or

MoonTag nanobodies[16]. In Boersma et al., the authors inserted the MoonTag and the

SunTag[16] into different translational reading frames of the same RNA. This allowed to

observe in real time heterogeneities in the start site selection of individual ribosomes

on a single mRNA molecule. Here, we introduce a new antibody-based detector/tag

pair to image translation in living cells and organisms based on multimerization of the

ALFA-Tag. The ALFA-array system has a number of advantages: i) it comes from

synthetic design and its sequence is thus absent from organisms; ii) the tag is small,

soluble, hydrophilic and lacks aldehydes-sensitive amino acids altered by fixation; iii)

the detection is through a nanobody, which is small and soluble, has a very high affinity

for the ALFA-Tag (K_d of 27 pM), favorably comparable to the MoonTag (Kd of 30 nM)[16].

The new translation imaging system described in this study should be applicable to

many different organisms to visualize protein localization and control of translation.

Moreover, the availability of several orthogonal systems will allow to better analyze of

the complexity of translation dynamics by simultaneously comparing several mRNA in

diverse cellular environments and conditions. The wide range of applications of the

ALFA-array system, coupled with other antibody-based translation detector, provides

to the scientific community a highly versatile toolbox to study translation at the single

molecule level in cell lines and multicellular organisms. The combination of nascent

protein visualization with improved RNA imaging such as the one provided by the NLS-

MCP-tdStaygold will facilitate future scientific breakthroughs.

Online methods

Reagents created in this study

Reagent type	Genotype	Insertion type and chromosome number	Additional information
Drosophila line	*nosEPr > scFv-msGFP2*	Landing site insertion vk33 – Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*nosEPr > scFv-GL*	Landing site insertion vk33 – Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*nosEPr > scFv-mNeonGreen*	Landing site insertion vk33 – Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*nosEPr > scFv-mAvic1*	Landing site insertion vk33 – Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*nosEPr > scFv-msGFP2*	Landing site insertion vk27 – Chr. II	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*nosEPr > scFv-GL*	Landing site insertion vk27 – Chr. II	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*nosEPr > scFv-mNeonGreen*	Landing site insertion vk27 – Chr. II	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*nosEPr > scFv-mAvic1*	Landing site insertion vk27 – Chr. II	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*UASp > scFv-msGFP2*	Landing site insertion vk33 – Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*UASp > scFv-GL*	Landing site insertion vk33 – Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*UASp > scFv-mNeonGreen*	Landing site insertion vk33 – Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*UASp > scFv-mAvic1*	Landing site insertion vk33 – Chr. III	mini-white marker, plasmid

			also available (in pBPhi vector)
Drosophila line	*UASy > scFv-msGFP2*	Landing site insertion - vk33 Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*nosEPr > NB-ALFA-msGFP2*	Landing site insertion - vk33 Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
Drosophila line	*Ilp4_32x_ALFA-array*	Landing site insertion - vk33 Chr. III	mini-white marker, plasmid also available (in pBPhi vector)
DNA plasmid	NB-ALFA in pUC57-Simple	N/A	Synthetized by Genescript, codon optimized
DNA plasmid	ALFA-array in pUC57-Simple	N/A	Synthetized by Genescript, with linker GSGSG and codon optimized
DNA plasmid	pMT-SuntagRLucFKBP24MS2v5-SV40	N/A	Used in S2 cell experiments
DNA plasmid	pAWF-stdMCP-stdHalotag	N/A	Used in S2 cell experiments
DNA plasmid	pAWF-scFv-msGFP2	N/A	Used in S2 cell experiments
DNA plasmid	pAWF-scFv-sfGFP	N/A	Used in S2 cell experiments
DNA plasmid	12x_ALFA-array for CRISPR tagging DYNC1H1	N/A	Used for HeLa cells experiments
DNA plasmid	NB-ALFA-sfGFP in pHR	N/A	Used for HeLa cells experiments
DNA plasmid	pAWF-anti-ALFA-msGFP2ΔGB1	N/A	Used in S2 cell experiments
DNA plasmid	pAWF-anti-ALFA-msGFP2	N/A	Used in S2 cell experiments
DNA plasmid	pAWF-scFv-msGFP2ΔGB1	N/A	Used in S2 cell experiments
DNA plasmid	pHAGE Ubc NLS-HA MCP-tdStayGold	N/A	Used for HeLa cells experiments

Fly stocks, handling and genetics

The *yw* fly stock was used as a wild type. The germline driver *nos-Gal4:VP16* (BL4937)

and the *His2av-mRFP* (Bl23651) fly stocks come from Bloomington Drosophila Stock

Center. The *nullo-Gal4* fly stock comes from R.L. laboratory. The

twi_suntag_MS2_CRISPR, twi_suntag_transgene and *scFv-mScarlet* lines were

from[7]. Briefly, the 32x suntag sequence from[4] (containing thirty-three suntag repeats)

was inserted in frame into *twi* gene[7] and 128X MS2 from[28] placed just after the stop

codon. All live experiments were done at 21 °C except when using - or comparing to -

strains containing UAS-GAL4 which were done at 25 °C.

Cloning and transgenesis

scFv-FP under *nanos* enhancer-promoter (EPr) lines were generated using

NEBuilder® HiFi DNA Assembly Master Mix with primers listed in supplementary table

1 and inserted into scFv-sfGFP vector[7] digested with BamHI/XhoI. The transgenic

construct was inserted into the VK00033 (BL9750) and VK00027 (BL9744) landing site

using PhiC31 targeted insertion[75]. *UASp-scFv-FP* lines were done by inserting

fragment of the scFv-FP plasmid digested with XhoI/HpaI in the UASp-scFv-sfGFP

vector[6] digested with XhoI/HpaI. The transgenic construct was inserted in the VK00033

(BL9750) landing site using PhiC31 targeted insertion[75]. Fluorescent proteins were

extracted from plasmids: mAvicFP1 from pNCST-mAvicFP1 (addgene 129509),

mGreenLantern from pcDNA3.1-mGreenLantern (addgene 161912), msGFP2

(addgene 160461), mNeonGreen is a kind gift from S.Piatti. The Ilp4 transgene

(regulatory regions and coding sequence) was synthesized (GenScript Biotech) into

pUC57-simple[7]. The 32x_ALFA-array repeats were synthesized (GenScript Biotech)

into pUC57-simple (see supplementary materials). The 32x_ALFA-array was inserted

between Kpn1 and EcoRV restriction sites of the Ilp4 puc57-simple vector. Ilp4-ALFA-

array fragment was inserted into pBPhi using PmeI and Nhe1 and injected into BL9750

using PhiC31 targeted insertion[75] (BestGene, Inc.). The NB-ALFA were synthesized

(GenScript Biotech) into pUC57-simple (see supplementary materials), then msGFP2, GB1 and a NLS was added using NEBuilder® HiFi DNA Assembly Master Mix with primers listed in supplementary table 1 and inserted into scFv-sfGFP vector[7] digested with BamHI/XhoI. The transgenic construct was inserted into the VK00033 (BL9750) landing site using PhiC31 targeted insertion[75]. All cloning for plasmids used in *Drosophila* S2 cells were performed using In-Fusion® HD Cloning Plus Kit (Takara Bio) and PCR was carried out with CloneAmp HiFi PCR premix (Takara Bio). Primers used are listed in supplementary table 1. To inducibly express SunTag in S2 cells, we generated pMT-SuntagRLucFKBP24MS2v5-SV40 by inserting SuntagRLucFKBP24MS2v5 fragment (addgene 119946) into pMT backbone that was PCR-amplified from pMT-EGFPActin5C (addgene 15312) with In-Fusion® HD Cloning Plus Kit. For constitutive expression in S2 cells, pAWF (DGRC stock # 1112) was used as an expression vector. pAWF-stdMCP-stdHalotag was generated by inserting stdMCP-stdHalotag (addgene 104999) into PCR-linearized pAWF vector using In-Fusion® HD Cloning Plus Kit. To generate pAWF-scFv-FPs, sequences containing scFv fused to various fluorescent proteins and GB1-NLS were amplified from *scFv-FP* under *nanos* enhancer promoter plasmids generated in this article and ligated with PCR-linearized pAWF vector using In-Fusion® HD Cloning Plus Kit. All plasmid sequences are listed in supplementary materials. The sequence of anti-ALFA-msGFP2-GB1-NLS was inserted into pAWF vector (DGRC #1112) to generate plasmid pAWF-anti-ALFA-msGFP2. To remove GB1 and generate pAWF-anti-ALFA-msGFP2 ΔGB1, a pair of primers flanking the GB1 sequence were used to amplify the entire pAWF-anti-ALFA-msGFP2 sequence except for the GB1 and ligated with KLD enzyme mix (NEB, M0554S). Same primer pairs were used to generate pAWF-scFv-msGFP2ΔGB1 from pAWF-scFv-msGFP2.

Single molecule fluorescence *in situ* hybridization (smFISH) and immuno-smFISH in *Drosophila*

A pool of 0-4h after egg-laying (AEL) embryos were dechorionated with bleach for 3 min and thoroughly rinsed with H_2O. They were fixed in 1:1 heptane:formaldehyde-10% for 25 min on a shaker at 450 rpm; formaldehyde was replaced by methanol and embryos were shaken by hand for 1 min. Embryos that sank to the bottom of the tube were rinsed three times with methanol and kept at -20°C for further use.

Embryos were washed 5 min in 1:1 methanol:ethanol, rinsed twice with ethanol 100%, washed 5 min twice in ethanol 100%, rinsed twice in methanol, washed 5 min once in methanol, rinsed twice in PBT-RNasin (PBS 1x, 0.1% tween, RNasin® Ribonuclease Inhibitors). Next, embryos were washed 4 times for 15 min in PBT-RNasin supplemented with 0.5% ultrapure BSA and then once 20 min in Wash Buffer (10% 20X SCC, 10% Formamide). They were then incubated overnight at 37 °C in Hybridization Buffer (10% Formamide, 10% 20x SSC, 400 µg/ml *E. coli* tRNA (New England Biolabs), 5% dextran sulfate, 1% vanadyl ribonucleoside complex (VRC) and smFISH Stellaris probes against *suntag*[7] or 32x_ALFA-array coupled to Quasar 570 (supplementary table 1) or probes against 32X MS2 coupled to Cy3 (from E. Bertrand)[55] and/or GCN4 primary antibody (Novus Biologicals C11L34) (1/200) and/or affinity purified rabbit anti-ALFA polyclonal antibody (NanoTag Biotechnologies) (1/200). Probe sequences are listed in Supplementary table 1. The next day, secondary antibody (1/500 anti-mouse Alexa 488-conjugated (Life Technologies, A21202); anti-mouse Alexa 647-conjugated (Invitrogen, A32728); anti-rabbit Alexa 555-conjugated (Life Technologies, A31572) or anti-rabbit Alexa 488-conjugated (Life Technologies, A21206)) was added if necessary, during 1h at 37° in Wash Buffer. DAPI

was added at 45min (1/1000). Embryos were then washed three times 15min in 2x

SCC, 0.1% Tween at room temperature before being mounted in ProLong® Diamond

antifade reagent.

Images were acquired using a Zeiss LSM880 confocal microscope with an Airyscan

detector in Super Resolution (SR) mode. GFP/Alexa-488 was excited using a 488 nm

laser, Cy3/Quasar 570 were excited using a 561 nm laser, Alexa 647 was excited using

a 633 nm laser. Laser powers were: 27μW for 488nm laser (9μW for UAS>nos

images), 80μW for 561nm laser and 46μW for 633nm laser. Laser powers were taken

under a 10X objective. Figures were prepared with ImageJ (National Institutes of

Health), Photoshop (Adobe Systems), and Illustrator (Adobe Systems).

Percentage of mRNA in translation and polysome intensities were quantified with

custom-made algorithms developed in PythonTM[7]. Quantification was done on three

area at the center and three area at the border of the mesoderm from three mid-late

embryos for *scFv-msGFP2>twi_suntag_MS2_CRISPR/+* (related to Figure 2b) and

three area at the center and three area at the border of the mesoderm from two mid-

late embryos for *twi_suntag_MS2_CRISPR/+* stained with anti-GCN4 antibody (related

to Figure 2e). Outliers were removed using the ROUT method[76], embedded in

GraphPad Prism 9.3.1 software, with Q set to 0.1%. scFv-FP intensities (related to

Figure S1f) were quantified using images from two to three early-mid *scFv-*

FP>twi_suntag_MS2_CRISPR/+ embryos, outliers were removed using the ROUT

method[76], embedded in GraphPad Prism 9.3.1 software, with Q set to 0.1%.

Cell culture

Drosophila S2R+ cells (DGRC Stock 150) were maintained at 25°C in Schneider's

medium containing 10% Fetal Bovine Serum and 1% Penicillin-streptomycin. For the

transfection, 0.7x10^6 cells in 350ul were seeded into each well of a 24-well plate. Cells are transfected with plasmid mix of pMT-SuntagRLucFKBP24MS2v5-SV40 or pMT-ALFAx32-RLucFKBP24MS2v5-SV40 (200ng), pAWF-MCP-Halotag (5ng), pAWF-scFv-FP-GB1-NLS or pAWF-scFv-FP-NLS (10ng) using effectene transfection reagent. 24h after transfection, cells were first incubated with Schneider's medium containing 200 nM Janelia Fluor® 594 HaloTag® Ligand (Promega GA1110) for 15min and then switched to medium with 1 mM $CuSO_4$ for 1h or 3h to induce expression of pMT construct.

HeLa cells were maintained in DMEM supplemented with 10% FBS, 10 U/ml penicillin/streptomycin and 2.9 mg/ml glutamine in a humidified CO_2 incubator at 37°C. For translational inhibition, cells were treated with puromycin at 100 μg/ml for 30 min. Cells were transfected with JetPrime (Polyplus) and selected on 150 μg/ml hygromycin. For each stable cell line, several individual clones were picked and screened by smFISH with sets of fluorescent oligonucleotide probes against the integrated sequence. For CRISPR recombination, clones were additionally analyzed by genomic PCR using amplicons specific for the non-recombined or the recombined allele. The CRISPR guide and repair plasmids have been previously[4]. For the repair plasmid, the 12x_ALFA-array replaced the SunTag 56x tag.

The NB-ALFA-sfGFP was introduced into HeLa cells by retroviral infection. HEK293T cells were transiently transfected with a cocktail of plasmids coding for retroviral components and producing the genomic NB-ALFA-sfGFP retroviral RNAs. Viral particles were collected and used to infect recipient HeLa cells, which were then sorted by FACS. Only lowly expressing cells were selected.

For FRAP and FRET experiment, 150k HeLa cells expressing NB-ALFA-sfGFP were seeded in fluorodish 35mm. The day after, cells were transfected with a mix of 3ug of

H2B_mCherry_ALFAtag plasmid, 4 μl of Jet Prime Reagent and 200 μl of Jet Prime

Buffer. The mix was vortexed for 15 seconds and incubated at RT for 10 minutes before

being added to the cells. After 24h, medium containing transfection reagents was

washed with PBS and replaced with fresh medium. Cells were incubated for an

additional 24h before use. One hour before the FRAP and FRET experiment, cells

were washed three times with PBS, and incubated in imaging media.

For testing NLS-MCP-tdStaygold, stable expression of fluorescent MCP fusions was

achieved by lentivirus-mediated transduction of a self-inactivating vector containing an

internal ubiquitin promoter as described in[55]. The MCP contained the deltaFG deletion,

the V29I mutation, and an SV40 NLS. Cells expressing fluorescent MCP fusions were

grown as a pool of clones and FACS-sorted to select cell populations expressing

homogeneous levels of fluorescence. The reporter cell lines additionally expressed the

HIV-1 reporter gene pIntro-MS2x128 previously described in[55].

Western blot

S2 cells (about 10^6 cells at 3x10^6 cell/ml density) were transfected with 300ng of

plasmids pAWF-scFv-msGFP2, pAWF-scFv-msGFP2ΔGB1, pAWF-anti-ALFA-

msGFP2, pAWF-anti-ALFA-msGFP2ΔGB1, or empty pAWF vector using Effectene

transfection reagent (Qiagen, # 301425). Cells were harvested 48h after transfection,

lysed with NuPAGE™ LDS Sample Buffer (Invitrogen, NP0007), and denatured by

heating at 70°C for 10 min. Proteins in the lysate samples were resolved by

electrophoresis using NuPAGE™ 4 to 12%, Bis-Tris gel, and transferred onto a PVDF

membrane using Trans-Blot Turbo Transfer System (Bio-Rad). The PVDF membrane

was blocked using Intercept (PBS) Blocking Buffer (LI-COR, 927-70001) for 1 h before

being incubated with primary antibodies (rabbit anti-GFP, Invitrogen, A11122; mouse

anti-αTubulin, Sigma, T6199-100UL) with 1:3000 dilution overnight at 4°C. When primary antibody was omitted, the membrane was incubated in the blocking buffer overnight at 4°C. The membrane was extensively washed before being incubated with secondary antibodies (Goat anti-Rabbit-HRP, Invitrogen 65-6120; Goat anti-Mouse-HRP, sigma 12-349) at 1:3000 dilution for 1 h at room temperature. The membrane was extensively washed before applying Pierce™ ECL Western Blotting Substrate (Thermo Scientific 32106) and visualized in the BioRad gel imaging system.

Single molecule fluorescence *in situ* hybridization (smiFISH) in mammalian cells

Cells were grown on glass coverslips (0.17 mm), washed with PBS, fixed in 4% PFA for 20 min, and permeabilized in 70% ethanol overnight at 4°C. Cells were hybridized as previously described for the smiFISH procedure[77], using probes were sets of unlabeled oligonucleotides hybridizing against the target RNA and additionally contained a common supplementary sequence that was pre-annealed to a fluorescent oligonucleotide probe termed the FLAP[77]. We used sets of oligonucleotides for the DYNC1H1 and POLR2A mRNAs[4]. Slides were mounted in Vectashield with DAPI (Vector Laboratories).

Fixed cells were imaged at room temperature on an Axioimager Z1 wide-field microscope (63×, NA 1.4; ZEISS) equipped with an sCMOs Zyla 4 0.2 camera (Andor Technology) and controlled by MetaMorph (Universal Imaging). 3D image stacks were collected with a Z-spacing of 0.3 μm. Figures were prepared with ImageJ (National Institutes of Health), Photoshop (Adobe Systems), and Illustrator (Adobe Systems).

Live cell imaging of mammalian cells

Cells were plated on FluoroDish (35mm, FD35-100, World Precision Instruments,

Sarasota,FL,USA) one day before the imaging. Before imaging media was replaced

for imaging purpose with a non-fluorescent media (FluoroBrite DMEM, from

thermofischer scientific cat no.A1896701, was supplemented with 10% FBS, 1%

penicillin/streptomycin, 1% glutamine and rutin; Evrogen). The dish was mounted in a

temperature-controlled chamber with 5 % CO_2 and imaged on an inverted OMXv3

Deltavision microscope in time-lapse mode. A 100x, NA 1.4 objective was used, with

an intermediate × 2 lens and an Evolve 512 × 512 EMCCD camera (Photometrics). 9

stacks with a z-spacing of 0.6 μm were acquired. Illuminating light and exposure time

were set at 1% of laser full power, exposure of 40 ms per plane. For the time-lapse

mode, one stack was recorded every 10s for 3h to 6h. Both the MCP-eGFP and MCP-

tdStaygold cell lines were imaged under the same conditions.

Signal quantification in mammalian cells

For translation imaging using the Suntag or ALFA-arrays, signal quantification was

done using ImageJ. The mean peak intensity of polysomes and mRNAs was calculated

from maximal image projection and then subtracted from the cytoplasmic background

mean intensity. Quantification were done in duplicates.

For MCP fusions, quantification was performed using big-FISH, a Python-based

software for the analysis of RNA single molecules[78]. Spots were detected using a

Laplacian of Gaussian threshold and the program returned the number of the RNA

molecule, mean and standard deviation of the fluorescence intensity. The threshold

was manually set for each frame for optimal detection of single RNAs and transcription

sites were removed from the analysis. Background nucleoplasmic intensities from free

fluorescent MCP proteins was subtracted from each frame. The code for the analysis

described in this paper is available on GitHub: https://github.com/fish-quant.

Fluorescence correlation spectroscopy

FCS experiments were performed on a Zeiss LSM780 microscope using a 40x/1.2

water objective. GFP was excited using the 488 nm line of an Argon laser with a

constant excitation intensity. Fluorescence intensity fluctuations were acquired for 1 s

and auto-correlation functions (ACFs) generated by Zen software were loaded in the

PyCorrFit program[79] and fit using a simple 3D gaussian model:

$$G(\tau) = \frac{1}{N}\left(1 + \frac{\tau}{\tau_D}\right)^{-1}\left(1 + \frac{\tau}{s^2\tau_D}\right)^{-1/2} \tag{1}$$

With N the number of molecules, τ_D the diffusion time and s the structural parameter

(shape of the PSF) fixed to 5. The brightness B is obtained by dividing the average

value of the fluorescence intensity ($\langle F(t)\rangle$) by the number of molecules N obtained by

the fit of the ACF with equation (1).

Multiple measurements per embryo, in multiple position and embryos at 20 °C were

used to generate multiple ACF, used to extract brightness and molecule numbers at

different time point during n.c.14.

Fluorescence recovery after photobleaching

Fluorescence recovery after photobleaching (FRAP) was performed in HeLa cells

expressing NB-ALFA-sfGFP wich was transfected with H2B_mCherry_ALFAtag

plasmid[52] and imaged 48 hours later at 37°C. The following settings was used: a Zeiss

LSM980 using a 40x/1.3 Oil objective. Images (512 × 512 pixels), 16bits/pixel, pixel

size 0.12µm, zoom 3.5x, were acquired every ≈10 s during 30 frames. Excitation: GFP

with a laser at 488nm and mCherry with a laser at 561nm. Laser power was measured

using the 40x/1.3 Oil objective at 1mW for 488nm and 561nm laser at 100% and acquisition laser power was measured at 50µW and 2.2µW for 488nm and 561nm diode laser respectively. A circular ROI (3.7µm diameter) was bleached using hundred pulses of 488 nm and 561 lasers at maximal power (100%) after 1 pre-bleach frames. To discard any source of fluorescence intensity recovery other than molecular diffusion, the measured fluorescence recovery in the bleached ROI region (I_{bl}) was corrected by an unbleached ROI (I_{unbl}) of the same nucleus and another ROI outside of the cells (I_{out}) following the simple equation:

$$I_{blcorr}(t) = \frac{I_{bl}(t) - I_{out}(t)}{I_{unbl}(t) - I_{out}(t)} \tag{2}$$

Förster resonance energy transfer

Acceptor photobleaching Förster resonance energy transfer (FRET) was performed in HeLa cells expressing NB-ALFA-sfGFP wich was transfected with H2B_mCherry_ALFAtag plasmid[52] and imaged 48 hours later at 37°C. The following settings was used: a Zeiss LSM980 using a 40x/1.3 Oil objective. Images (512 × 512 pixels), 16bits/pixel, pixel size 0.12µm, zoom 3.5x, were acquired every ≈8 s during 30 frames. Excitation: GFP with a laser at 488nm and mCherry with a laser at 561nm. Laser power was measured using the 40x/1.3 Oil objective at 1.6mW for 514nm and 1mW for 561nm diode laser at 100% and acquisition laser power was measured at 50µW and 2.2µW for 488nm and 561nm diode laser respectively. A circular ROI (2.5 µm diameter), was bleached using hundred pulses of 514nm and 561nm lasers at maximal power (100%) after 1 pre-bleach frames. To discard any source of fluorescence intensity fluctuation other than molecular diffusion, the measured fluorescence increase for the GFP and fluorescence recovery for the mCherry in the

bleached ROI region (I_{bl}) were corrected by an unbleached ROI (I_{unbl}) of the same

nucleus and another ROI outside of the cells (I_{out}) following the simple equation (2)

Light-Sheet Microscopy

For light-sheet imaging (related to Movie S6-7 and Figures S1d), the MuViSPIM[80]

(Luxendo, Brüker company GMBH) was employed. This setup provides two-sided

illumination with two Nikon 10x/0.3 water objectives and two-sided detection with two

Olympus 20x/1.0 W objectives. The lightsheet is created by scanning a Gaussian

beam, as in digital scanned laser light-sheet microscopy (DSLM). We used the line

mode which is an in-built synchronization between rolling shutter readout mode of

sCMOS cameras and digital light sheets canning. This allows the rejection of out-of-

focus and scattered light, thus improving the signal to noise ratio. Images are acquired

by two ORCA Flash 4.0 (C91440) from Hamamatsu and processed by LuxControl

v1.10.2. A 50ms exposure time was used for the green channel with a 488nm laser.

Maximum intensity projections were processed with Fiji[81].

Live imaging of the scFv-FP and NB-ALFA-msGFP2 alone

For all live imaging experiments, embryos were dechorionated with tape and mounted

between a hydrophobic membrane and a coverslip as described previously[82].

Movies for scFv-FP (related to Figure S1 and Movie S1-5) and NB-ALFA-msGFP2

under *nanos* promoter (related to Figure S8 and Movie S23) were acquired using a

Zeiss LSM880 confocal microscope with a Plan-Apochromat 40x/1.3 oil objective lens,

with green excitation using a 488nm laser (6µw measured with 10x objective lens). A

GaAsP detector was used to detect fluorescence with the following settings: 512x 512-

pixels images, zoom 2x, each Z-stack comprised of 25 planes spaced 1µm apart.

Movies for *UASp>scFv-FP* (related to Figure S6, and Movie S16-19) and *UASy>scFv-*

msGFP2 with *nos-Gal4* driver (related to Figure S7 and Movie S21) were acquired

using a Zeiss LSM880 confocal microscope with a Plan-Apochromat 40x/1.3 oil

objective lens, with green excitation using a 488nm laser (1µw measured with 10x

objective lens). A GaAsP detector was used to detect fluorescence with the following

settings: 512x 512-pixels images, zoom 2x, each Z-stack comprised of 25 planes

spaced 1µm apart.

Live imaging of the scFv-FP and anti-ALFA-msGFP2 in presence of tagged genes

For all live imaging experiments, embryos were dechorionated with tape and mounted

between a hydrophobic membrane and a coverslip as described previously[82].

Movies of *scFv-FP>twi_suntag_MS2_CRISPR/+* (related to Figure 1 and Movie S8-

12) were acquired using a Zeiss LSM880 with confocal microscope in fast Airyscan

mode with a Plan-Apochromat 40x/1.3 oil objective lens. GFP was excited using a

488nm (7uw measured with 10x objective lens) laser with the following settings: 132x

132-pixel images, each Z-stack comprised of 15 planes spaced 0.5µm apart and zoom

8x. Intensity measurement of each polysomes was done on a MIP of 5 planes using a

custom-made software written in Python, that performs a spot detection on the MIP

raw data stack. For each time frame, the software performs a Laplacian of Gaussian

2D filter and the filtered images are than thresholded with a user defined threshold.

The threshold is expressed in terms of the average and standard deviation of intensity

values of each image in order to have a threshold value self-re-scaled with respect to

the characteristic of the images itself. Spots smaller in size than a user-defined value

are removed (considered as fake detection). Finally, for each of the detected spots, we

use a 2D expansion of the spots silhouettes in order to define a region around the

spots devoided of any other spots and that we can use to estimate background and

measure it all along the time evolution for corrections purposes.

Movies of *scFv-FP>twi_suntag_MS2_CRISPR/+* and *scFv-*

msGFP2>twi_suntag_transgene (related to Figure 2, Figure S4 and Movie S13-15)

were acquired using a Zeiss LSM880 with confocal microscope in fast Airyscan mode

with a Plan-Apochromat 40x/1.3 oil objective lens. GFP was excited using a 488nm

(4.7μW measured with 10x objective lens) laser with the following settings: 568x 568-

pixel images, each Z-stack comprised of 61 planes spaced 0.5μm apart and zoom 2x.

Movies were then processed to remove frame outside of the embryos or containing the

membrane signal to correct drifting, and processed stacks were maximum intensity

projected using custom-made software, developed in Python[TM7].

Movies of *nos-Gal4>UASp-scFv-msGFP2* x *twi_suntag_MS2_CRISPR/+* and *scFv-*

msGFP2 x *twi_suntag_MS2_CRISPR/+* (related to Figure 3e and Movie S20) were

acquired using a Zeiss LSM880 with confocal microscope in fast Airyscan mode with

a Plan-Apochromat 40x/1.3 oil objective lens. GFP was excited using a 488nm (5uw

measure with 10x objective lens) laser with the following settings: 396x 396-pixel

images, each Z-stack comprised of 30 planes spaced 0.5μm apart and zoom 4x.

Live tilescans of *nullo-Gal4*>UASy-*scFv-msGFP2-NLS* (related to Movie S22) were

acquired using a Zeiss LSM880 with confocal microscope in fast Airyscan mode with

a Plan-Apochromat 40x/1.3 oil objective lens. GFP was excited using a 488nm (13μW

measured with 10x objective lens) laser with the following settings: 1440 x 2746-pixel

images using two tiles, each Z-stack comprised of 9 planes spaced 2μm apart and

zoom 0.8x.

Live tilescans of *UASy-scFv-msGFP2-NLS/ his2Av-RFP* in presence or not of *nullo-*

Gal4 driver (related to Supplementary Figure S7d) were acquired using a Zeiss

LSM880 with confocal microscope in fast Airyscan mode with a Plan-Apochromat

40x/1.3 oil objective lens. GFP was excited using a 488nm (13μW measured with 10x

objective lens) and RFP was excited using a 561nm laser with the following settings:

1024 x 3072-pixel images using three tiles, each Z-stack comprised of 14 planes

spaced 2µm apart.

Movies of NB-ALFA-msGFP2 under *nanos* EPr crossed with *Ilp4_32x_ALFA-array*

(related to Figure 4e and Movie S24) were acquired using a Zeiss LSM880 with

confocal microscope in fast Airyscan mode with a Plan-Apochromat 40x/1.3 oil

objective lens. GFP was excited using a 488nm (6.4μW measured with 10x objective

lens) laser with the following settings: 256x 256-pixel images, each Z-stack comprised

of 6 planes spaced 1µm apart and zoom 6x.

Live tilescans of NB-ALFA-msGFP2 under *nanos* EPr crossed with *Ilp4_32x_ALFA-*

array (related to Figure S8e-f and Movie S25) were acquired using a Zeiss LSM880

with confocal microscope in fast Airyscan mode with a Plan-Apochromat 40x/1.3 oil

objective lens. GFP was excited using a 488nm (13μW measured with 10x objective

lens) laser with the following settings: 1612 x 4860-pixel images using 3 tiles, each Z-

stack comprised of 9 planes spaced 2µm apart.

Movies of *NB-ALFA-msGFP2*, *scFv-mScarlet*, *twi_suntag_MS2_CRISPR* crossed with

Ilp4_32x_ALFA-array (related to Figure 4i and Movie S26) and NB-ALFA-msGFP2,

scFv-mScarlet crossed with *Ilp4_32x_ALFA-array* (related to Figure S10b and Movie

S27) were acquired using a Zeiss LSM880 with confocal microscope in fast Airyscan

mode with a Plan-Apochromat 40x/1.3 oil objective lens. GFP and mScarlet were

excited using a 488nm and 561nm laser respectively with the following settings: 372x

372-pixel images, one Z-plane and zoom 6x.

S2 cell culture imaging

Cells were pipetted into a μ-Slide (Ibidi #80621) and live-imaged using a Zeiss LSM980 confocal microscope with an Airyscan detector in SR mode with a 63X Plan-Apochromat (1.4NA) oil objective lens. GFP was excited using a 488nm laser and MCP-Halotag labelled by Janelia Fluor® 594 was excited using a 561nm laser. Cells with mRNAs visible in cytoplasm and with comparable GFP signal were chosen for acquiring images. Images of individual cells were acquired with following setting: z-stack of 5 planes with 0.5um interval, 8x zoom, 396x396 pixel and 8 bit per pixel. GFP foci co-localized with mRNA foci (Janelia Fluor® 594 signal) were considered as translation sites while GFP foci or larger amorphous or ring-like GFP structures that were not co-localized with mRNA were considered as aggregates.

Acknowledgments

We are grateful to NanoTag Biotechnologies for sharing ALFA-Tag and anti-ALFA nanobody plasmids and advices. We are grateful to Dr. Karim Mazjoub's lab for sharing space in his experimental room. We thank Drs. Xavier Pichon and Dr. Vera Slaninova for their help in setting up the mammalian ALFA-Tag system and Dr. Vera Slaninova for helping developing the MCP-tdStaygold fusion. For imaging, we acknowledge Sylvain de Rossi and Montpellier Ressources Imagerie (MRI), a microscopy facility of The National Infrastructure France-BioImaging (FBI), supported by the French National Research Agency (ANR-10-INBS-04). We acknowledge M. Verbrugghe for help with fly handling, and the Montpellier *Drosophila* Facility (BioCampus). We thank Dr. Radhan Ramadass for assistance with software. We acknowledge Dr Etienne Schwob and Dr. Eric Kremer for their help in preparing the manuscript. We acknowledge HFSP (CDA Grant to M.L.) for the early sponsoring of the implementation of *Drosophila*

translation imaging project. Funding: This work was supported by the ERC SyncDev

starting grant to M.L., the ANR MemoRNP to M.L., and by IGMM internal capital, D.A.

was granted by the CNRS consortium GDR ImaBio. M.L., J.D., E.B. and C.F. are

sponsored by CNRS. M.B. was supported by a FRM fellowship and then by ERC

SyncDev. H.L was supported by ERC SyncDev. J.Dh. was supported by ANR grant

ULTIM. E.B. was supported by grants from MSDAvenir and ANR ULTIM.

Author contributions:

J.D. conceived the project. J.D. conceptually supervised the work with the help of M.L.

at its early stages. J.D., M.B., M.L., E.B., R.L. and R.C. designed the experiments.

J.D., M.B., and C.F. performed *Drosophila* embryos experiments. A.T. developed

software. H.L. helped with cloning and fly handling. M.L. acquired funding. E.B.

supervised mammalian cells experiments, which were performed by J.Dh.. R.L.

supervised *Drosophila* S2 cells experiments performed by R.C.. J.D., M.B., R.C., A.T.

E.B. and C.F. analyzed and interpreted the results. C.F. analyzed and supervised FCS

experiments with the help of J.D and M.B. J.D. and M.B. wrote the manuscript. M.L.,

R.C., E.B., R.L. and C.F. edited the manuscript. H.K., F.M. and E.B. developed MCP-

tdStaygold fusion. E.B. and J.Dh. developed the mammalian ALFA tag system. J.D.,

M.B., R.C., D.A., H.K., F.M and J.Dh. performed all experiments done for the review

process. All authors discussed and approved the manuscript.

Competing interests:

The authors declare that they have no competing interests.

Data and materials availability:

All relevant data supporting the key findings of this study are available within the article

and its Supplementary Information files or from the corresponding author upon

reasonable request. The intensity measurement software is available through this link:

https://github.com/ant-trullo/Av_Ints_software/. *Drosophila* stocks will be deposited to

VDRC stock center and can be asked to M.L. (mounia.lagha@igmm.cnrs.fr). Materials

used for S2 cell experiments can be asked to R.L. (lehmann@wi.mit.edu). Materials

used for mammalian cell experiments can be asked to E.B.

(edouard.bertrand@igh.cnrs.fr).

References:

1. Yan, X., Hoek, T. A., Vale, R. D. & Tanenbaum, M. E. Dynamics of Translation
of Single mRNA Molecules In Vivo. *Cell* **165**, 976–989 (2016).
2. Wang, C., Han, B., Zhou, R. & Zhuang, X. Real-Time Imaging of Translation on
Single mRNA Transcripts in Live Cells. *Cell* **165**, 990–1001 (2016).
3. Morisaki, T. *et al.* Real-time quantification of single RNA translation dynamics
in living cells. *Science (1979)* **352**, 1425–1429 (2016).
4. Pichon, X. *et al.* Visualization of single endogenous polysomes reveals the
dynamics of translation in live human cells. *J Cell Biol* **214**, 769–781 (2016).
5. Wu, B., Eliscovich, C., Yoon, Y. J. & Singer, R. H. Translation dynamics of
single mRNAs in live cells and neurons. *Science (1979)* **352**, 1430–1435 (2016).
6. Formicola, N. *et al.* Tyramine induces dynamic RNP granule remodeling and
translation activation in the Drosophila brain. *Elife* **10**, (2021).
7. Dufourt, J. *et al.* Imaging translation dynamics in live embryos reveals spatial
heterogeneities. *Science (1979)* **372**, 840–844 (2021).
8. Vinter, D. J., Hoppe, C., Minchington, T. G., Sutcliffe, C. & Ashe, H. L.
Dynamics of hunchback translation in real-time and at single-mRNA resolution
in the Drosophila embryo. *Development* **148**, (2021).
9. Lecerf, J.-M. *et al.* Human single-chain Fv intrabodies counteract *in situ*
huntingtin aggregation in cellular models of Huntington's disease. *Proceedings*
of the National Academy of Sciences **98**, 4764–4769 (2001).
10. Wörn, A. *et al.* Correlation between in vitro stability and in vivo performance of
anti-GCN4 intrabodies as cytoplasmic inhibitors. *J Biol Chem* **275**, 2795–2803
(2000).
11. Colby, D. W. *et al.* Development of a human light chain variable domain (V(L))
intracellular antibody specific for the amino terminus of huntingtin via yeast
surface display. *J Mol Biol* **342**, 901–912 (2004).
12. Tanenbaum, M. E., Gilbert, L. A., Qi, L. S., Weissman, J. S. & Vale, R. D. A
protein-tagging system for signal amplification in gene expression
and fluorescence imaging. *Cell* **159**, 635–646 (2014).

13. Chouaib, R. *et al.* A Dual Protein-mRNA Localization Screen Reveals
Compartmentalized Translation and Widespread Co-translational RNA
Targeting. *Dev Cell* **54**, 773-791.e5 (2020).
14. Boersma, S. *et al.* Translation and Replication Dynamics of Single RNA
Viruses. *Cell* **183**, 1930-1945.e23 (2020).
15. Bruurs, L. J. M. *et al.* Heterogeneity in viral replication dynamics shapes the
antiviral response. *bioRxiv* 2022.06.08.495262 (2022)
doi:10.1101/2022.06.08.495262.
16. Boersma, S. *et al.* Multi-Color Single-Molecule Imaging Uncovers Extensive
Heterogeneity in mRNA Decoding. *Cell* **178**, 458-472 e19 (2019).
17. Safieddine, A. *et al.* A choreography of centrosomal mRNAs reveals a
conserved localization mechanism involving active polysome transport. *Nat*
Commun **12**, 1352 (2021).
18. Chalfie, M., Tu, Y., Euskirchen, G., Ward, W. W. & Prasher, D. C. Green
fluorescent protein as a marker for gene expression. *Science* **263**, 802–805
(1994).
19. Prasher, D. C., Eckenrode, V. K., Ward, W. W., Prendergast, F. G. & Cormier,
M. J. Primary structure of the Aequorea victoria green-fluorescent protein. *Gene*
111, 229–233 (1992).
20. Zacharias, D. A., Violin, J. D., Newton, A. C. & Tsien, R. Y. Partitioning of
lipid-modified monomeric GFPs into membrane microdomains of live cells.
Science (1979) **296**, (2002).
21. Rodriguez, E. A. *et al.* The Growing and Glowing Toolbox of Fluorescent and
Photoactive Proteins. *Trends Biochem Sci* **42**, 111–129 (2017).
22. Lambert, G. G. *et al.* Aequorea's secrets revealed: New fluorescent proteins with
unique properties for bioimaging and biosensing. *PLoS Biol* **18**, (2020).
23. Bindels, D. S. *et al.* mScarlet: a bright monomeric red fluorescent protein for
cellular imaging. *Nat Methods* **14**, 53–56 (2017).
24. Leake, M. C. & Quinn, S. D. A guide to small fluorescent probes for single-
molecule biophysics. *Chemical Physics Reviews* **4**, 011302 (2023).
25. Bertrand, E. *et al.* Localization of ASH1 mRNA particles in living yeast. *Mol*
Cell **2**, 437–445 (1998).
26. Fusco, D. *et al. Single MRNA Molecules Demonstrate Probabilistic Movement*
in Living Mammalian Cells HHS Public Access Author Manuscript. Curr Biol
vol. 13 http://images.cellpress.com/supmat/supmatin.htm. (2003).
27. Forrest, K. M. & Gavis, E. R. Live imaging of endogenous RNA reveals a
diffusion and entrapment mechanism for nanos mRNA localization in
Drosophila. *Curr Biol* **13**, 1159–1168 (2003).
28. Tantale, K. *et al.* A single-molecule view of transcription reveals convoys of
RNA polymerases and multi-scale bursting. *Nat Commun* (2016)
doi:10.1038/ncomms12248.
29. Tutucci, E. *et al.* An improved MS2 system for accurate reporting of the mRNA
life cycle. *Nat Methods* **15**, 81–89 (2018).
30. Hocine, S., Raymond, P., Zenklusen, D., Chao, J. A. & Singer, R. H. Single-
molecule analysis of gene expression using two-color RNA labeling in
live yeast. *Nat Methods* **10**, 119–121 (2013).
31. Weil, T. T., Parton, R. M. & Davis, I. Making the message clear: visualizing
mRNA localization. *Trends Cell Biol* **20**, 380–390 (2010).
32. Tutucci, E. *et al.* An improved MS2 system for accurate reporting of the mRNA
life cycle. *Nat Methods* **15**, 81–89 (2018).

1026	33.	Pichon, X., Robert, M.-C., Bertrand, E., Singer, R. H. & Tutucci, E. New
1027		Generations of MS2 Variants and MCP Fusions to Detect Single mRNAs in
1028		Living Eukaryotic Cells. *Methods Mol Biol* **2166**, 121–144 (2020).
1029	34.	Tanenbaum, M. E., Gilbert, L. A., Qi, L. S., Weissman, J. S. & Vale, R. D. A
1030		protein-tagging system for signal amplification in gene expression
1031		and fluorescence imaging. *Cell* **159**, 635–646 (2014).
1032	35.	Wörn, A. *et al.* Correlation between in vitro stability and in vivo performance of
1033		anti-GCN4 intrabodies as cytoplasmic inhibitors. *J Biol Chem* **275**, 2795–2803
1034		(2000).
1035	36.	Donahue, C. E. T., Siroky, M. D. & White, K. A. An Optogenetic Tool to Raise
1036		Intracellular pH in Single Cells and Drive Localized Membrane Dynamics. *J Am*
1037		*Chem Soc* **143**, (2021).
1038	37.	Chen, D. & McKearin, D. M. A discrete transcriptional silencer in the bam gene
1039		determines asymmetric division of the Drosophila germline stem cell.
1040		*Development* **130**, 1159–1170 (2003).
1041	38.	Ali, I. *et al.* Cis-regulatory elements affecting the Nanos gene promoter in the
1042		germline stem cells. *J Biotechnol* **145**, 323–329 (2010).
1043	39.	Doren, M. Van, Williamson, A. L. & Lehmann, R. Regulation of zygotic gene
1044		expression in Drosophila primordial germ cells. *Current Biology* **8**, 243–246
1045		(1998).
1046	40.	Garcia, H. G., Tikhonov, M., Lin, A. & Gregor, T. Quantitative Imaging of
1047		Transcription in Living Drosophila Embryos Links Polymerase Activity to
1048		Patterning. *Current Biology* (2013) doi:10.1016/j.cub.2013.08.054.
1049	41.	Pédelacq, J. D., Cabantous, S., Tran, T., Terwilliger, T. C. & Waldo, G. S.
1050		Engineering and characterization of a superfolder green fluorescent protein. *Nat*
1051		*Biotechnol* **24**, (2006).
1052	42.	Valbuena, F. M. *et al.* A photostable monomeric superfolder green fluorescent
1053		protein. *Traffic* **21**, (2020).
1054	43.	Campbell, B. C. *et al.* mGreenLantern: a bright monomeric fluorescent protein
1055		with rapid expression and cell filling properties for neuronal imaging. *Proc Natl*
1056		*Acad Sci U S A* **117**, (2020).
1057	44.	Shaner, N. C. *et al.* A bright monomeric green fluorescent protein derived from
1058		Branchiostoma lanceolatum. *Nat Methods* **10**, (2013).
1059	45.	Pan, D. J., Huang, J. D. & Courey, A. J. Functional analysis of the Drosophila
1060		twist promoter reveals a dorsal-binding ventral activator region. *Genes Dev* **5**,
1061		1892–1901 (1991).
1062	46.	Brogiolo, W. *et al.* An evolutionarily conserved function of the drosophila
1063		insulin receptor and insulin-like peptides in growth control. *Current Biology* **11**,
1064		(2001).
1065	47.	Ivorra-Molla, E. *et al.* A monomeric StayGold fluorescent protein. *Nature*
1066		*Biotechnology 2023* 1–4 (2023) doi:10.1038/s41587-023-02018-w.
1067	48.	Rørth, P. Gal4 in the Drosophila female germline. *Mech Dev* **78**, 113–118
1068		(1998).
1069	49.	Brand, A. H. & Perrimon, N. Targeted gene expression as a means of altering
1070		cell fates and generating dominant phenotypes. *Development* **118**, 401–415
1071		(1993).
1072	50.	DeLuca, S. Z. & Spradling, A. C. Efficient Expression of Genes in the
1073		Drosophila Germline Using a UAS Promoter Free of Interference by Hsp70
1074		piRNAs. *Genetics* **209**, 381–387 (2018).

1075   51.   Götzke, H. *et al.* The ALFA-tag is a highly versatile tool for nanobody-based
1076   bioscience applications. *Nat Commun* **10**, 4403 (2019).
1077   52.   Alessandra Vigano, M. *et al.* Protein manipulation using single copies of short
1078   peptide tags in cultured cells and in Drosophila melanogaster. *Development*
1079   *(Cambridge)* **148**, (2021).
1080   53.   Mueller, F., Morisaki, T., Mazza, D. & McNally, J. G. Minimizing the Impact of
1081   Photoswitching of Fluorescent Proteins on FRAP Analysis. *Biophys J* **102**, 1656
1082   (2012).
1083   54.   Hirano, M. *et al.* A highly photostable and bright green fluorescent protein. *Nat*
1084   *Biotechnol* **40**, 1132–1142 (2022).
1085   55.   Tantale, K. *et al.* Stochastic pausing at latent HIV-1 promoters generates
1086   transcriptional bursting. *Nat Commun* **12**, 4503 (2021).
1087   56.   Lambert, T. J. FPbase: a community-editable fluorescent protein database.
1088   *Nature Methods 2019 16:4* **16**, 277–278 (2019).
1089   57.   Zhang, H. *et al.* Quantitative assessment of near-infrared fluorescent proteins.
1090   *Nature Methods 2023 20:10* **20**, 1605–1616 (2023).
1091   58.   Cranfill, P. J. *et al.* Quantitative assessment of fluorescent proteins. *Nat.*
1092   *Methods* **13**, 557–562 (2016).
1093   59.   Schneider, K. L., Wollman, A. J. M., Nyström, T. & Shashkova, S. Comparison
1094   of endogenously expressed fluorescent protein fusions behaviour for protein
1095   quality control and cellular ageing research. *Sci Rep* **11**, (2021).
1096   60.   Botman, D., de Groot, D. H., Schmidt, P., Goedhart, J. & Teusink, B. In vivo
1097   characterisation of fluorescent proteins in budding yeast. *Scientific Reports 2019*
1098   *9:1* **9**, 1–14 (2019).
1099   61.   Heppert, J. K. *et al.* Comparative assessment of fluorescent proteins for in vivo
1100   imaging in an animal model system. *Mol. Biol. Cell* **27**, 3385–3394 (2016).
1101   62.   Piatkevich, K. *et al.* Bright and stable monomeric fluorescent proteins derived
1102   from StayGold. (2023) doi:10.21203/RS.3.RS-3188559/V1.
1103   63.   Ando, R. *et al.* StayGold variants for molecular fusion and membrane-targeting
1104   applications. *Nature Methods 2023* 1–9 (2023) doi:10.1038/s41592-023-02085-
1105   6.
1106   64.   Ivorra-Molla, E. *et al.* A monomeric StayGold fluorescent protein. *Nature*
1107   *Biotechnology 2023* 1–4 (2023) doi:10.1038/s41587-023-02018-w.
1108   65.   Hirano, M. *et al.* StayGold Photostability under Different Illumination Modes.
1109   *bioRxiv* 2023.11.23.568490 (2023) doi:10.1101/2023.11.23.568490.
1110   66.   Song, S.-J., Diao, H.-P., Moon, B., Yun, A. & Hwang, I. The B1 Domain of
1111   Streptococcal Protein G Serves as a Multi-Functional Tag for Recombinant
1112   Protein Production in Plants. *Front Plant Sci* **13**, (2022).
1113   67.   Bothma, J. P., Norstad, M. R., Alamos, S. & Garcia, H. G. LlamaTags: A
1114   Versatile Tool to Image Transcription Factor Dynamics in Live Embryos. *Cell*
1115   **173**, 1810-1822.e16 (2018).
1116   68.   Dong, J. & Ueda, H. Recent Advances in Quenchbody, a Fluorescent
1117   Immunosensor. *Sensors (Basel)* **21**, 1–18 (2021).
1118   69.   Mateju, D. *et al.* Single-Molecule Imaging Reveals Translation of mRNAs
1119   Localized to Stress Granules. *Cell* **183**, 1801-1812.e13 (2020).
1120   70.   Ramat, A. *et al.* The PIWI protein Aubergine recruits eIF3 to activate translation
1121   in the germ plasm. *Cell Res* **30**, 421–435 (2020).
1122   71.   Kang, J.-Y. *et al.* LLPS of FXR1 drives spermiogenesis by activating translation
1123   of stored mRNAs. *Science (1979)* **377**, eabj6647 (2022).

1124   72.   Liu, Y. *et al.* Visualizing looping of two endogenous genomic loci using
1125   synthetic zinc-finger proteins with anti-FLAG and anti-HA frankenbodies in
1126   living cells. *Genes to Cells* **26**, 905 (2021).
1127   73.   Zhao, N. *et al.* A genetically encoded probe for imaging nascent and mature HA-
1128   tagged proteins in vivo. *Nature Communications 2019 10:1* **10**, 1–16 (2019).
1129   74.   Koch, A., Aguilera, L., Morisaki, T., Munsky, B. & Stasevich, T. J. Quantifying
1130   the dynamics of IRES and cap translation with single-molecule resolution in live
1131   cells. *Nat Struct Mol Biol* **27**, 1095–1104 (2020).
1132   75.   Venken, K. J. T., He, Y., Hoskins, R. A. & Bellen, H. J. P[acman]: A BAC
1133   transgenic platform for targeted insertion of large DNA fragments in D.
1134   melanogaster. *Science (1979)* **314**, 1747–1751 (2006).
1135   76.   Motulsky, H. J. & Brown, R. E. Detecting outliers when fitting data with
1136   nonlinear regression - A new method based on robust nonlinear regression and
1137   the false discovery rate. *BMC Bioinformatics* **7**, 1–20 (2006).
1138   77.   Tsanov, N. *et al.* SmiFISH and FISH-quant - A flexible single RNA detection
1139   approach with super-resolution capability. *Nucleic Acids Res* **44**, (2016).
1140   78.   Imbert, A. *et al.* FISH-quant v2: a scalable and modular tool for smFISH image
1141   analysis. *RNA* **28**, 786–795 (2022).
1142   79.   Muller, P., Schwille, P. & Weidemann, T. PyCorrFit-generic data evaluation for
1143   fluorescence correlation spectroscopy. *Bioinformatics* **30**, 2532–2533 (2014).
1144   80.   Krzic, U., Gunther, S., Saunders, T. E., Streichan, S. J. & Hufnagel, L.
1145   Multiview light-sheet microscope for rapid in toto imaging. *Nat Methods* **9**,
1146   730–733 (2012).
1147   81.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.
1148   *Nat Methods* **9**, 676–682 (2012).
1149   82.   Dufourt, J. *et al.* Temporal control of gene expression by the pioneer factor
1150   Zelda through transient interactions in hubs. *Nat Commun* **9**, (2018).

1151

1152

1153

1154

1155   **Figure legends**

1156

Figure 1: Generation of different scFv-Fluorescent proteins (scFv-FP) to monitor

nascent translation in *Drosophila*.

a. Schematic of the SunTag system. Once the mRNA (black) is translated by the

ribosomes (grey), the suntag epitopes (light blue) will be bound by the single chain

variable fragment (scFv) (dark blue) fused to a fluorescent protein (FP) (green).

Therefore, the nascent peptide will be detected by the accumulation of fluorescent

signal.

b. (top) Schematic of the different constructs of scFv-FP generated in this study. scFv-

msGFP2, scFv-GreenLantern, scFv-mAvic1 and scFv-NeonGreen were created for

this study and scFv-sfGFP in our previous study[7]. (bottom) Schematic of the

CRISPR/Cas9 targeted insertion strategy to obtain endogenous *twist* gene tagged with

suntag and 128xMS2[7].

c. (top) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern

in purple. Anterior side is on the left, posterior on the right, dorsal on the top and ventral

side on the bottom. Black square represents the imaged area of the bottom panels.

(bottom) One Z-plane of confocal images from smFISH with direct fluorescent protein

signal in green (scFv-msGFP2, scFv-GreenLantern, scFv-mAvic1, scFv-NeonGreen

and scFv-sfGFP) and *MS2* probes (red) on *scFv-FP x twi_suntag_MS2_CRISPR*

embryos in early-mid n.c. 14. Scale bar are 5µm.

d. Snapshots from representative fast-mode acquired confocal movies of

twi_suntag_MS2_CRISPR/+ embryos carrying either scFv-msGFP2, scFv-

GreenLantern, scFv-mAvic1, scFv-NeonGreen or scFv-sfGFP proteins. Green dots

represent nascent translation of *twi*. Scale bars are 5µm. (See related Movies S8-12).

e. Quantification across time during n.c. 14 of different scFv-FP signals from movies

represented in (d), n=5 movies from at least 3 embryos for each scFv-FP. Error bars

represent SEM.

1183

1184 **Figure 2: scFv-FP concentration under *nos* EPr creates an artefactual**

1185 **translation arrest of endogenous *twi* mRNA during early embryogenesis**

a. Schematic of nuclear elongation and cellularization during n.c. 14. In this study, we defined one early-mid stage (corresponding to 0 to 35min of the n.c. 14 from the mitosis) and one mid-late stage (corresponding to 35 to 55min of the n.c. 14 from the mitosis).

b. (top) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern in purple. Anterior side is on the left, posterior on the right, dorsal on the top and ventral side on the bottom. Black square represents the imaged area of the bottom panels. (bottom) A single Z-plane of confocal images from smFISH with direct fluorescent protein signal in green (scFv-msGFP2) and *MS2* probes (red) on *scFv-msGFP2 x twi_suntag_MS2_CRISPR* embryos at mid-late n.c. 14. Gray squares represent the zoomed images in the right panels. The two different zoomed images represent border (top) and internal (bottom) zone of the imaged pattern. Scale bars are 10µm on the larger images, and 5µm on the zoomed images. Staging is given by DAPI staining on a sagittal view of the imaged embryo (black and white image at the bottom). Quantification of the scFv-msGFP2 signal intensity on single molecule mRNA at the border at mid-late n.c. 14 stage (dark green, 9 images from three embryos, n=2570) and center (light green, 9 images from three embryos, n=2737) of the mesoderm is represented on the right.

c. Schematic representing *twi_suntag_MS2_CRISPR* mRNA expression from n.c. 13 to late n.c. 14 in red. From the observations on *twi* CRISPR translation (b), two hypotheses can be considered. Hypothesis 1 is that the decrease of translation observed at mid-late n.c. 14 is due to an active repression of translation. Hypothesis 2 is that the amount of scFv-FP becomes limiting at mid-late n.c. 14, leading to an arrest of the detected green signal. These two hypotheses are represented in the context of *twi_suntag_MS2_CRISPR* expression (above) and *twi_suntag_transgene* expression

(below). *twi_suntag_transgene* mRNA is expressed more stochastically and later than

twi_suntag_MS2_CRISPR mRNA. In hypothesis 1, a decrease of translation should

be simultaneously observed for the two constructs (mid-late n.c. 14). In hypothesis 2,

no arrest of translation should be observed with *twi_suntag_transgene* as the amount

of mRNA is lower.

d. Snapshots each 15 minutes from movies of *scFv-msGFP2 x*

twi_suntag_MS2_CRISPR or *scFv-msGFP2 x twi_suntag_transgene* embryos on the

ventral side. T0 correspond to early n.c.14. White squares represent the zoomed

images in the center of the panel. Note a persistence of translation for the

twi_suntag_transgene at T0+45min (white arrowhead), absent in

twi_suntag_MS2_CRISPR embryos. Scale bars are 10µm on the larger images, and

5µm on the zoomed images. (See related Movies S13-14).

e. (top) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern

in purple. Anterior side is on the left, posterior on the right, dorsal on the top and ventral

side on the bottom. Black square represents the imaged area of the bottom panels.

(bottom) Single Z-planes of confocal images from immuno-smFISH with anti-GCN4

antibody (cyan) and MS2 probes (red) on *twi_suntag_MS2* embryos at mid-late n.c.

14. Gray squares represent the zoomed images in the right panels. The two different

zoomed images represent border (top square) and internal (bottom square) zone of

the imaged pattern. Scale bars are 10µm on the larger images, and 5µm on the

zoomed images. Staging is given by DAPI staining on a sagittal view of the imaged

embryo (black and white image at the bottom). Quantification of the anti-GCN4 signal

intensity on single molecule mRNA at the border (dark blue, 6 images from two

embryos, n=3211) and center (light blue, 6 images from two embryos, n=4001) at mid-

late n.c. 14 stage of the mesoderm is represented on the right.

1236

Figure 3: Increasing scFv-FP expression allows translation detection at later stages

a. Schematic of *UASp-scFv-msGFP2* construct and activation by the Gal4 protein. 10 UAS sequences (purple) are placed upstream the P-promoter (blue) and scFv-msGFP2 (green) sequences. Gal4 proteins (yellow) will bind the UAS sequences to activate transcription of the scFv-msGFP2.

b. Schematic of the expression of the *scFv-msGFP2* in *nosGal4>UASp-scFv-msGFP2* female strain during oogenesis until egg deposition.

c. (left) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern in purple. Anterior side is on the left, posterior on the right, dorsal on the top and ventral side on the bottom. Black square represents the imaged area of the right panels. Single Z-planes of confocal images from smFISH with direct scFv-msGFP2 fluorescent protein signal (green) and *MS2* probes (red) on *nosGal4>UASp-scFv-msGFP2 x twi_suntag_MS2_CRISPR* embryos in early, early-mid and mid-late n.c. 14. Gray squares represent the zoomed images in for each panel. Nuclei are counterstained with DAPI (grey, bottom images) for staging. Scale bars are 10µm on the larger images, and 5µm on the zoomed images.

d. (top) Schematic of a *Drosophila* embryo on the ventral side with gastrulation furrow represented with invaginating cells. Black square represents the imaged area of the bottom panels. (bottom) Single Z-planes of confocal images from smFISH with direct scFv-msGFP2 fluorescent protein signal (green) and *MS2* probes (red) on *nosGal4>UASp-scFv-msGFP2 x twi_suntag_MS2_CRISPR* and *scFv-msGFP2 x twi_suntag_MS2_CRISPR* embryos at gastrulation stage and on the ventral side.

Scale bars are 10µm on the larger images, and 5µm on the zoomed images. Note that

translation dots are visible only with *nosGal4>UASp-scFv-msGFP2*.

e. (top) Schematic of a *Drosophila* embryo on the ventral side with gastrulation furrow

represented with invaginating cells. Black square represents the imaged area of the

bottom panels. (bottom) Single Z-planes of confocal images from movie of *scFv-*

msGFP2 x yw, scFv-msGFP2 x twi_suntag_MS2_CRISPR and *nosGal4>UASp-scFv-*

msGFP2 x twi_suntag_MS2_CRISPR embryos at gastrulation stage (furrow

represented with grey dashed line). Scale bars are 10µm. (See related Movie S20).

1268

1269

Figure 4: Development of the ALFA-array system to detect nascent translation

in *Drosophila*

a. Peptide sequence of one suntag with its linker (24 amino-acids) and peptide

sequence of the ALFA-tag with its linker (20 amino-acids).

b. Schematic of the scFv and the nanobody ALFA (NB-ALFA) fused to the fluorescent

protein msGFP2 and their size in amino-acid (aa).

c. Schematic of the ALFA-array system developed in this study to detect translation.

32 ALFA-tag sequences were multimerized (blue amino-acid sequence) and inserted

at the 5' of the gene of interest sequence (yellow). Upon translation, nanobodies ALFA

fused to msGFP2 (NB-ALFA_msGFP2) will bind to the ALFA-tag peptides. As a

bipartite system, the NB-ALFA is expressed under *nos* EPr fused to *msGFP2*,

Streptococcal protein G (*GB1*) domain and a *Nuclear Localisation Signal* (*NLS*).

d. Representation of the transgene mRNA containing 32x_ALFA-array in frame with

the *insulin-like peptide 4* (*Ilp4*) gene sequence. Schematic of a sagittal view of a

Drosophila embryo with *Ilp4* expression pattern in yellow. Anterior side is on the left,

posterior on the right, dorsal on the top and ventral side on the bottom. Black square represents the imaged area of the bottom panels. One Z-plane of confocal images from smFISH with direct Nb-ALFA-msGFP2 fluorescent protein signal (green) and 32x_*ALFA-array* probes (red) on *NB-ALFA_msGFP2>Ilp4_32x_ALFA-array* embryos in early-mid n.c. 14 on the ventral side. Scale bars are 10µm on the larger images, and 5µm on the zoomed images.

e. Snapshots from representative fast-mode acquired confocal movies of *Ilp4_32x_ALFA-array/+* n.c. 14 embryos carrying *NB-ALFA_msGFP2*. Bright white foci represent nascent translation of *Ilp4* transgene. Yellow squares represent zoomed images of the top panels. Nascent translation of *Ilp4* transgene is indicated by yellow arrowheads. Scale bars are 10µm on the larger images, and 2µm on the zoomed images. (See related Movies S24).

f. Representation of the transgene mRNAs containing 32x_ALFA-array fused to *Ilp4* coding sequence or suntag fused to *twist* coding sequence used in (g) and (h).

g. Schematic of a *Drosophila* embryo on the ventral side with *twi* expression pattern in purple and *Ilp4* in yellow. Black square represents the imaged area of the bottom panels. One Z-plane of confocal images from immuno-smFISH with anti-ALFA (green) labeling nascent translation of *Ilp4_32x_ALFA-array*, probes against *32x_ALFA-array* (red) labeling *Ilp4_32x_ALFA-array* mRNA molecules and anti-GCN4 antibody labeling nascent translation of *twi_suntag_transgene* (magenta) in n.c. 14 embryos. Scale bars are 1µm.

h. Maximum intensity projection of a whole embryo confocal images from immuno-staining with anti-ALFA antibody (green) and anti-GCN4 antibody (magenta) on *Ilp4_32x_ALFA-array; twi_suntag_transgene* gastrulating embryo. Scale bars are 100µm.

i. Schematic of a *Drosophila* embryo on the ventral side with *twi* expression pattern in purple and *Ilp4* in yellow. Black square represents the imaged area of the bottom panels. Single Z-plane of confocal images from dual-color live imaging of *twi_suntag_MS2_CRISPR/+; NB-ALFA_msGFP2/scFv-mScarlet x Ilp4_32x_ALFA-array* embryos in n.c. 14. Scale bars are 1µm. (see related Movie S26).

j. (top) Schematic of a *Drosophila* embryo on the ventral side with gastrulation furrow represented with invaginating cells. Confocal snapshots of a whole live embryo expressing *twi_suntag_MS2_CRISPR/+; NB-ALFA_msGFP2/scFv-mScarlet x Ilp4_32x_ALFA-array*, after gastrulation. Note the cytoplasmic staining from *Ilp4* translation (green) and nuclear staining from *twi* translation (purple). Scale bars are 100µm.

Figure 5: Development of the ALFA-array system to detect nascent translation in mammalian cells

a. (1) Schematic of the CRISPR/Cas9 targeted insertion strategy to obtain endogenous *DYNC1H1* gene tagged with 12X ALFA-tag. (2) Schematic of the ALFA-array system developed in mammalian cells. 12 ALFA-tag sequences were multimerized (blue amino-acid sequence) and inserted at the 5' of the gene of interest sequence (DYNC1H1, yellow). (3) The NB-ALFA is expressed under *spleen focus-forming virus (SFFV)* promoter fused to *sfGFP* and *Streptococcal protein G* (*GB1*) domain.

b. Left: micrographs of HeLa cells expressing a DYNC1H1 allele endogenously tagged 12x_ALFA-array tag and NB-ALFA-sfGFP, and treated (right) or not (left) with puromycin. Green: NB-ALFA-sfGFP signal; blue: DAPI. Scale bars are 5µm. Arrows: NB-ALFA-sfGFP spots. Right: quantification of the number of NB-ALFA-sfGFP spots

per cell, with and without puromycin treatment. Error bars: standard deviation (n=40 cells).

c. Micrographs of HeLa cells expressing a DYNC1H1 allele endogenously tagged 12x_ALFA-array tag and NB-ALFA-sfGFP, and hybridized in situ with a set of probes against DYNC1H1 mRNAs. On the merged panel: NB-ALFA-sfGFP signal (green); smFISH signals (red); DAPI (blue). Scale bar is 5µm. Arrows: DYNC1H1 polysomes.

d. Top: schematic of the MCP fusion used to image RNA. Bottom: maximum projection intensity images of cells stably expressing MCP-tdStayGold (on the left) or MCP-eGFP (on the right). All images are taken under the same conditions on an OMX widefield microscope. Six time point are selected (0 min, 30 min, 60 min, 120 min, 150 min and 180 min) and they are displayed with the same dynamic range. Red arrows point at transcription sites and green arrows point at individual RNA molecules. Scale bar: 10 µm.

e. Intensities of single RNA molecules over time. Graph shows intensities of RNA molecules (mean and s.d. error; five cells). Because of bleaching single molecule signal could no longer be detected and measured after 120 min in the case of MCP-eGFP.

f. The graph shows the number of single RNA molecule detected over time using FISHquant (five cells).

Figure Supp 1: Characterization of the different scFv-Fluorescent Proteins (scFv-FP)

a. Schematic of the expression pattern of scFv-FP under control of the *nanos* enhancer/promoter during *Drosophila* oogenesis.

b. Snapshots from movies on embryos carrying scFv-sfGFP, scFv-msGFP2, scFv-GreenLantern, scFv-mAvic1 or scFv-NeonGreen in early and late n.c. 14. Scale bars are 10µm. (see related Movies S1-5).

c. Quantification across time during n.c. 14 of the different scFv-FP signal from movies represented in (b), n=3 movies from 3 embryos for scFv-sfGFP and scFv-GreenLantern and n=4 movies from 4 embryos for scFv-msGFP2, scFv-mAvic1 and scFv-NeonGreen. Error bars represent SEM.

d. Snapshots from light-sheet movies on embryos carrying scFv-mAvic1 or scFv-msGFP2 in n.c. 14 and at gastrulation stage. Ventral and dorsal view are represented for each embryo. Scale bars are 100µm. (See related Movies S6-7).

e. Single Z-planes of confocal images from smFISH with direct fluorescent protein signal in green (scFv-msGFP2, scFv-GreenLantern, scFv-mAvic1, scFv-NeonGreen and scFv-sfGFP) and *MS2* probes (red) on *scFv-FP x twi_suntag_MS2_CRISPR* embryos in early-mid n.c. 14 on the ventral side. Images represent zoomed images from Figure 1c. Scale bars are 1µm.

f. Quantification of the different scFv-FP signal intensity on single molecule mRNA (n=1646, n=3658, n=3080, n=6820, n=2924, of scFv-sfGFP, scFv-msGFP2, scFv-mAvic1, scFv-GL and scFv-NG respectively) in the mesoderm of early-mid n.c. 14 embryo from immuno-smFISH (cropped area inside *twi* pattern, typical images in Figure 1c).

Figure Supp 2: Localization of the tagged protein and scFv recognition by secondary antibody

a. (top) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern in purple. Black square represents the imaged area of the bottom panels. Single Z-

planes of confocal images from immuno-smFISH with direct fluorescent protein signal

in green (scFv-msGFP2, scFv-GreenLantern, scFv-mAvic1, scFv-NeonGreen and

scFv-sfGFP) and secondary anti-mouse antibody (magenta) on *scFv-FP x*

twi_suntag_MS2_CRISPR embryos in n.c. 14 on the ventral side. Note the co-

localization between the direct fluorescent protein signal and the secondary antibody

signal as illustrated by the schematic on the top right. Scale bar are 1μm.

b. (top) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern

in purple. Black square represents the imaged area of the bottom panels. Single Z-

planes of confocal images from immuno-smFISH with MS2 probes (red) and

secondary anti-mouse antibody (magenta) on *twi_suntag_MS2_CRISPR* embryos in

n.c. 14 on the ventral side. Note that in absence of scFv-FP the secondary antibody

does not detect any signal as illustrated by the schematic on the bottom. Scale bars

are 1μm.

c. (top) Schematic of a *Drosophila* embryo on the ventral side with gastrulation furrow

represented with invaginating cells. Black square represents the imaged area of the

bottom panels. (bottom) Single Z-planes of confocal images from smFISH with direct

scFv-msGFP2 fluorescent protein signal (green) and *MS2* or *Ilp4* probes (red) on *scFv-*

msGFP2 > twi_suntag_MS2_CRISPR and *scFv-msGFP2 > Ilp4_suntag* embryos at

gastrulation stage and on the ventral side. Note a nuclear signal in

twi_suntag_trangene embryos and a cytoplasmic signal in *Ilp4_suntag* embryos. Scale

bars are 10μm.

1404

Figure Supp 3: The scFv-sfGFP can form aggregates in *Drosophila* embryos and

S2 cells

a. Schematic of a *Drosophila* embryo on the ventral side with gastrulation furrow

represented with invaginating cells. Black square represents the imaged area of the

bottom panels. Snapshots from representative confocal movies of

twi_suntag_MS2_CRISPR/+ embryos at the gastrulation stage carrying either scFv-

sfGFP, scFv-NeonGreen, scFv-mAvic1 or scFv-msGFP2 proteins. Scale bars are

10µm.

b. Schematic of a *Drosophila* embryo on the ventral side with gastrulation furrow

represented with invaginating cells. Black square represents the imaged area of the

bottom panels. Maximum intensity projection of 10 Z-planes of confocal images from

smFISH with direct fluorescent protein signal in green (scFv-FP) and *MS2* probes (red)

on *scFv-FP x twi_suntag_MS2_CRISPR* embryos at gastrulation stage on the ventral

side. Grey squares represent zoomed images. Note the presence of scFv-sfGFP signal

not overlapping with RNA (white arrowheads). Scale bars are 5µm.

c. Schematic of the different constructs transfected in S2 cells. *scFv-FPs* and *tdMCP-*

stdHalotag were placed downstream of the *Actin5C* promoter. Reporter containing

suntag Renilla Luciferase (RLuc) and 24XMS2 was placed under the control of the

inducible metallothionein (MT) promoter.

d. Single Z-planes of confocal images of S2 cells expressing scFv-FPs (green) and

MCP-Halotag (red) with induction of *SuntagRLuc24XMS2* reporter showing

aggregation size of the scFv-FPs. A zoom on single mRNA in translation is shown on

no aggregate's panels (white squares). Scale bars are 5µm.

e. Percentage of the number of big (dark blue) and small (blue) aggregates (scFv-FP

signal not colocalizing with an mRNA) and no (light blue) aggregates (scFv-FP signal

colocalizing with an mRNA) in cells expressing *MCP-Halotag*, *SuntagRLuc24XMS2*

reporter and *scFv-sfGFP2* or *scFv-msGFP2* (n=100 cells for each condition).

1432

Figure Supp 4: *twi* translation detection in developing embryos until gastrulation

a. (top) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern in purple. Anterior side is on the left, posterior on the right, dorsal on the top and ventral side on the bottom. Black square represents the imaged area of the bottom panels. (bottom) A single Z-plane of confocal images from smFISH with direct fluorescent protein signal in green (scFv-mAvic1) and *MS2* probes (red) on *scFv-mAvic1 x twi_suntag_MS2_CRISPR* embryos at mid-late n.c. 14. Gray squares represent the zoomed images in the right panels. The two different zoomed images represent border (top square) and internal (bottom square) zone of the imaged pattern. Scale bars are 10µm on the larger images, and 5µm on the zoomed images. Staging is given by DAPI staining on a sagittal view of the imaged embryo (black and white image at the bottom).

b. (top) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern in purple. Anterior side is on the left, posterior on the right, dorsal on the top and ventral side on the bottom. Black square represents the imaged area of the bottom panels. (bottom) A single Z-plane of confocal images from smFISH with direct fluorescent protein signal in green (scFv-NeonGreen) and *MS2* probes (red) on *scFv-NeonGreen x twi_suntag_MS2_CRISPR* embryos at mid-late n.c. 14. Gray squares represent the zoomed images in the right panels. The two different zoomed images represent border (top square) and internal (bottom square) zone of the imaged pattern. Scale bars are 10µm on the larger images, and 5µm on the zoomed images. Staging is given by DAPI staining on a sagittal view of the imaged embryo (black and white image at the bottom).

c. (top) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern in purple. Anterior side is on the left, posterior on the right, dorsal on the top and ventral side on the bottom. Black square represents the imaged area of the bottom panels.

(bottom) A single Z-plane of confocal images from smFISH with direct fluorescent

protein signal in green (scFv-GreenLantern) and *MS2* probes (red) on *scFv-*

GreenLantern x twi_suntag_MS2_CRISPR embryos at mid-late n.c. 14. Gray squares

represent the zoomed images in the right panels. The two different zoomed images

represent border (top square) and internal (bottom square) zone of the imaged pattern.

Scale bars are 10µm on the larger images, and 5µm on the zoomed images. Staging

is given by DAPI staining on a sagittal view of the imaged embryo (black and white

image at the bottom).

d. Snapshots each 15 minutes from movies of *scFv-msGFP2 x*

twi_suntag_MS2_CRISPR embryos on the ventral side. T0 correspond to early n.c.14.

White squares represent the zoomed images in the center of the panel. Note the

absence of translation for *twi_suntag_MS2_CRISPR* at T0+45min, Scale bars are

10µm on the larger images, and 5µm on the zoomed images. (See related Movies

S15).

e. Single Z-planes of confocal images from smFISH with direct fluorescent protein

signal in green (scFv-msGFP2) and *suntag* probes (red) on *scFv-msGFP2 x*

twi_suntag_transgene embryos in late n.c. 14 (top panels) and gastrulation stage

(bottom panels) on the ventral side. Nuclei are counterstained with DAPI (blue). Scale

bars are 10µm.

f. Representative examples of autocorrelation function from FCS performed in

embryos without translation (*scFv-msGFP2* only, upper panel) or with translation

(*scFv-msGFP2 x twi_suntag_MS2_CRISPR*) at early-mid (green), mid-late (blue) and

late-gastrulation (purple) n.c. 14 stages. Dark curves represent fitting of the data using

a diffusion model (see methods).

g. Dot plots representing the brightness as a function of the number of molecules obtained from FCS analysis (f). In the absence of translation: early-mid n=120, mid-late n=160; in the presence of translation early-mid n=722.

Figure Supp 5: anti-GCN4 immunostaining detect translation until gastrulation

a. (top) Schematic of a *Drosophila* embryo on the ventral side with gastrulation furrow represented with invaginating cells. Black square represents the imaged area of the bottom panels. (bottom) Single Z-planes of confocal images from immuno-smFISH with anti-GCN4 antibody (cyan) and MS2 probes (red) on *twi_suntag_MS2_CRISPR* embryos. Grey squares represent the zoomed images in the bottom panels. Scale bars are 10µm on the larger images, and 5µm on the zoomed images.

Figure Supp 6: Generation of UASp-Gal4 system to overexpress scFv-Fluorescent Proteins

a. (top) Schematic of the different constructs expressing the different scFv-FP (scFv-GreenLantern, scFv-NeonGreen, scFv-mAvic1, or scFv-msGFP2) under the control of the UASp promoter. (bottom) Snapshots from movies of embryos carrying *UASp-scFv-msGFP2, UASp-scFv-GreenLantern, UASp-scFv-mAvic1* or *UASp-scFv-NeonGreen* and *nosGal4* at early n.c.14 and gastrulation stage. Scale bars are 10µm. (See related Movies S16-19).

b. Representative examples of autocorrelation function from FCS performed in embryos overexpressing the scFv-msGFP2 without translation (nosGal4>UASp-*scFv-msGFP2* only, upper panel) or with translation (nosGal4>UASp-*scFv-msGFP2 x twi_suntag_MS2_CRISPR*) at early-mid (green), mid-late (blue) and late-gastrulation

(purple) n.c. 14 stages. Dark curves represent fitting of the data using a diffusion model

(see methods).

c. Dot plots representing the brightness as a function of the number of molecules

obtained from FCS analysis (f). In the absence of translation: early-mid n=280, mid-

late n=280 and late-gastrulation n=200; in the presence of translation early-mid n=683,

mid-late n=877 and late-gastrulation n=489.

1511

Figure Supp 7: Generation of UASy-Gal4 system to express scFv-Fluorescent

Proteins

a. Schematic of the two constructs UASz and UASy containing ten *UAS* with *hsp70*

promoter and with *IVS-syn21* sequence (UASz) or without (UASy). Both of these

constructs contain the *scFv-msGFP2*.

b. Snapshots from movies of embryos carrying *UASy-scFv-msGFP2* and *nanos-Gal4*

at early n.c.14 and gastrulation stage. Scale bars are 5µm. (See related Movies S21).

c. (left) Schematic of a sagittal view of a *Drosophila* embryo with *twi* expression pattern

in purple. Anterior side is on the left, posterior on the right, dorsal on the top and ventral

side on the bottom. Black square represents the imaged area of the right panels. (right)

A single Z-plane of confocal images from smFISH with direct fluorescent protein signal

in green (scFv-msGFP2) and *MS2* probes (red) on *nosGal4>UASy-scFv-msGFP2 x*

twi_suntag_MS2_CRISPR embryos in early-mid and mid-late n.c. 14. Staging is given

by DAPI staining on a sagittal view of the embryo (black and white image at the

bottom). Blue and white squares (border and internal zone of the pattern, respectively)

represent the two different zoomed images on the right side of each panel. Scale bars

are 10µm on the larger images, and 5µm on the zoomed images.

d. Maximum intensity projected snapshots of a whole embryos confocal live tilescan

expressing *UASy-scFv-msGFP2* and *Histone-H2A-RFP* (left panels) or

nulloGal4>UASy-scFv-msGFP2 and Histone-H2A-RFP (right panels), after

gastrulation. Scale bars are 100µm.

1533

Figure Supp 8: Efficient detection of *Ilp4* transgene translation with the new

AlfaTag system

a. (left) Schematic of a sagittal view of a *Drosophila* embryo with *Ilp4* expression

pattern in yellow. Anterior side is on the left, posterior on the right, dorsal on the top

and ventral side on the bottom. (right) Maximum intensity projection of a whole embryo

tilescan confocal images from immuno-smFISH with anti-ALFA antibody (green),

32x_ALFA-array probes (red) and anti-GCN4 antibody (magenta) on *Ilp4_32x_ALFA-*

array n.c.14 embryos. Scale bars are 100µm.

b. (top) Schematic of a sagittal view of a *Drosophila* embryo with *Ilp4* expression

pattern in yellow. Black square represents the imaged area of the bottom panels.

(bottom) Single Z-planes of confocal images from immuno-smFISH with anti-ALFA

antibody (green) and *32x_ALFA-array* probes (red) on *Ilp4_32x_ALFA-array* in early-

mid n.c. 14 embryos at the border of *Ilp4* expression pattern. Scale bars are 10µm.

c. (top) Schematic of a sagittal view of a *Drosophila* embryo with *Ilp4* expression

pattern in yellow. Black square represents the imaged area of the bottom panels.

(bottom) Single Z-planes of confocal images from immuno-smFISH with anti-ALFA

antibody (green) and *32x_ALFA-array* probes (red) on *Ilp4_32x_ALFA-array* in early-

mid n.c. 14 embryos to visualize single mRNA molecules in translation. Scale bars are

5µm.

d. (top) Schematic of the expression pattern of NB-ALFA under control of the *nos EPr* during *Drosophila* oogenesis. (bottom) Snapshots from movies of embryos carrying *nos>NB-ALFA-msGFP2* in n.c. 11, 12, 13 and 14. Scale bars are 10µm. (See related Movies S23).

e. (top) Schematic of a sagittal view of a *Drosophila* embryo with *Ilp4* expression pattern in yellow. Black square represents the imaged area of the bottom panel. (bottom) Snapshots from movies of *NB-ALFA-msGFP2 x Ilp4_32x_ALFA-array* embryos in n.c. 14. The top image is a tile scan of the whole embryo and the bottom image a zoomed area represented with a white square at the border of *Ilp4* expression pattern. Note the green signal located in nuclei outside the pattern and in the cytoplasm within *Ilp4* pattern (corresponding to tagged Ilp4 proteins). Scale bars are 100µm in the tilescan and 10µm in the zoomed image. (See related Movies S25).

f. (top) Schematic of a sagittal view of a *Drosophila* embryo with *Ilp4* expression pattern in yellow. Black square represents the imaged area of the bottom panel. (bottom) Snapshots from movies of *NB-ALFA_msGFP2 > Ilp4_32x_ALFA-array* embryos at gastrulation stage. The top image is a tile scan of the whole embryo and the bottom image a zoomed area represented with a white square at the gastrulation furrow. Scale bars are 100µm in the tilescan and 10µm in the zoomed image. (See related Movies S24).

Figure Supp 9: The GB1 domain is recognized by secondary antibodies

a. (top) Schematic of a sagittal view of a *Drosophila* embryo with *Ilp4* expression pattern in yellow. Black square represents the imaged area of the bottom panels. Single Z-planes of confocal images from immuno-smFISH with direct fluorescent protein signal in green (NB-ALFA-msGFP2) and secondary anti-mouse antibody

(magenta) on *NB-ALFA-msGFP2 x llp4_32x_ALFA-array* embryos in n.c. 14 on the ventral side. Note the co-localization between the direct fluorescent protein signal and the secondary antibody signal as illustrated by the schematic on the bottom. Scale bars are 1µm.

b. (top) Schematic of a sagittal view of a *Drosophila* embryo with *llp4* expression pattern in yellow. Black square represents the imaged area of the bottom panels. Single Z-planes of confocal images from immuno-smFISH with ALFA probes (red) and secondary anti-mouse antibody (magenta) on *llp4_32x_ALFA-array* embryos in n.c. 14 on the ventral side. Note that in absence of NB-ALFA the secondary antibody does not detect any signal as illustrated by the schematic on the bottom. Scale bars are 1µm.

c. Western-blot experiments on *Drosophila* S2R+ cells transfected with scFv-msGFP2, scFv-msGFP2ΔGB1, empty vector, NB-ALFA-msGFP2 or NB-ALFA-msGFP2ΔGB1. The membrane on the left was hybridized with either anti-GFP primary and anti-rabbit secondary antibodies or with anti-rabbit secondary alone. Bands corresponding to the scFv and the NB fused to msGFP2 (with and without the GB1) are highlighted with black stars. The red stars highlight the binding of the secondary anti-rabbit-HRP to the scFv-msGFP2 and NB-msGFP2 in the presence of the GB1. The membrane on the right was hybridized with either anti-mouse secondary alone or with anti-Tubulin primary and anti-mouse secondary antibodies. The green stars highlight the binding of the secondary anti-mouse-HRP to the scFv-msGFP2 and NB-msGFP2 in the presence of the GB1.

Figure Supp 10: Specificity of detection of two distinct mRNA species in fixed and live samples

a. Maximum intensity projection of a whole embryo tilescan confocal images from immuno-smFISH with anti-ALFA antibody (green), *32x_ALFA-array* probes (red) and anti-GCN4 antibody (magenta) on *Ilp4_32x_ALFA-array; twi_suntag_transgene* embryos in n.c. 14. Scale bars are 100µm.

b. Schematic of a *Drosophila* embryo on the ventral side with *twi* expression pattern in purple and *Ilp4* in yellow. Black square represents the imaged area of the bottom panels. One Z-plane of confocal images from live imaging of *NB-ALFA_msGFP2/scFv-mScarlet x Ilp4_32x_ALFA-array* embryos in n.c. 14. Note the absence of signal with scFv-mScarlet illustrating specificity of the scFv and the NB-ALFA. Scale bars are 1µm (see related Movies S27).

c. (top) Schematic of a *Drosophila* embryo on the ventral side with gastrulation furrow represented with invaginating cells. Confocal snapshots of a live embryo expressing *NB-ALFA_msGFP2/scFv-mScarlet x Ilp4_32x_ALFA-array*, after gastrulation. Scale bars are 100µm.

Figure Supp 11: NB-ALFA exhibits a long binding time with its ALFA-Tag

a. Images from a FRET experiment on HeLa cells expressing NB-ALFA-sfGFP (green) and transfected with an H2B_mCherry_ALFAtag plasmid (magenta). Images before and after photobleaching are represented (note the increase in fluorescence intensity in the GFP channel after having bleached the mCherry). The bleached area is represented with dashed white circles. Scale bars are 5µm.

b. FRET recovery curve from experiment in (a) of the fluorescence intensity as a function of time. Recovery of the NB-ALFA-sfGFP is in green (FRET) and of the H2B (bleached) in magenta.

c. Images from a FRAP experiment on HeLa cells expressing a NB-ALFA-sfGFP

(green) and transfected with an H2B_mCherry_ALFAtag plasmid (magenta). Images

before and after photobleaching are represented. The bleached area is represented

with dashed white circles. Scale bars are 5μm.

d. FRAP recovery curve from experiment in (c) of the fluorescence intensity as a

function of time. Recovery of the NB-ALFA-sfGFP is in green and of the H2B in

magenta.

e. Left: images are micrographs of Hela with a POLR2A allele tagged with 32X_ALFA-

array (top), or a 32X_suntag (bottom), with in the merge the antibody signal in green

and the smiFISH signal of POLR2A mRNA in red. DNA is stained with DAPI. Scale

bar: 3 μm. Arrows: POLRA polysomes. Right: quantification of polysome and single

RNA intensities (arbitrary light units; n=5 cells).

1638

1639 **Movie Sup legends**

1640
Movie S1: Maximum intensity projection of confocal Z-stack live imaging of an embryo

containing *scFv-sfGFP*. scale bar 10μm.

Movie S2: Maximum intensity projection of confocal Z-stack live imaging of an embryo

containing *scFv-mAvic1*. scale bar 10μm.

Movie S3: Maximum intensity projection of confocal Z-stack live imaging of an embryo

containing *scFv-msGFP2*. scale bar 10μm.

Movie S4: Maximum intensity projection of confocal Z-stack live imaging of an embryo

containing *scFv-NeonGreen*. scale bar 10μm.

Movie S5: Maximum intensity projection of confocal Z-stack live imaging of an embryo

containing *scFv-GreenLantern*. scale bar 10μm.

Movie S6: Maximum intensity projection of LightSheet Z-stack live imaging of *scFv-*

mAvic1 embryo. Upper part: dorsal view, lower part: ventral view Scale bar 100µm.

Movie S7: Maximum intensity projection of LightSheet Z-stack live imaging of *scFv-*

msGFP2 embryo. Upper part: dorsal view, lower part: ventral view Scale bar 100µm.

Movie S8: Maximum intensity projection of confocal Z-stack fast live imaging of an

embryo containing *scFv-NeonGreen>twi_suntag_MS2_CRISPR/+.* Translation foci

are in green, scale bar 1µm.

Movie S9: Maximum intensity projection of confocal Z-stack fast live imaging of an

embryo containing *scFv-sfGFP >twi_suntag_MS2_CRISPR/+.* Translation foci are in

green, scale bar 1µm.

Movie S10: Maximum intensity projection of confocal Z-stack fast live imaging of an

embryo containing *scFv-GreenLantern >twi_suntag_MS2_CRISPR/+.* Translation foci

are in green, scale bar 1µm.

Movie S11: Maximum intensity projection of confocal Z-stack fast live imaging of an

embryo containing *scFv-msGFP2>twi_suntag_MS2_CRISPR/+.* Translation foci are in

green, scale bar 1µm.

Movie S12: Maximum intensity projection of confocal Z-stack fast live imaging of an

embryo containing *scFv-mAvic1>twi_suntag_MS2_CRISPR/+.* Translation foci are in

green, scale bar 1µm.

Movie S13: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *scFv-msGFP2>twi_suntag_MS2_CRISPR/+.* Translation foci are in

white, scale bar 10µm.

Movie S14: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *scFv-msGFP2>twi_suntag_transgene/+.* Translation foci are in

white, scale bar 10µm.

Movie S15: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *scFv-mAvic1> twi_suntag_transgene/+.* Translation foci are in

white, scale bar 10µm.

Movie S16: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *nos-Gal4>UASp-scFv-GreenLantern /+*. scale bar 10µm.

Movie S17: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *nos-Gal4>UASp-scFv-msGFP2 /+*. scale bar 10µm.

Movie S18: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *nos-Gal4>UASp-scFv-NeonGreen /+*. scale bar 10µm.

Movie S19: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *nos-Gal4>UASp-scFv-mAvic1 /+*. scale bar 10µm.

Movie S20: Comparison of maximum intensity projection of confocal Z-stack live

imaging of an embryo containing *scFv-msGFP2>twi_suntag_MS2_CRISPR/+ or nos-*

Gal4>UASp-scFv-mAvic1 ; twi_suntag_MS2_CRISPR/+. Translation foci are in white.

Translationnal arrest is seen *scFv-msGFP2>twi_suntag_MS2_CRISPR/+ but not* in

nos-Gal4>UASp-scFv-mAvic1 ; twi_suntag_MS2_CRISPR/+. scale bar 10µm.

Movie S21: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *nos-Gal4>UASy-scFv-msGFP2 /+*. scale bar 10µm.

Movie S22: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *nullo-Gal4>UASy-scFv-msGFP2 /+*. scale bar 100µm.

Movie S23: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *NB-ALFA-msGFP2*. scale bar 10µm.

Movie S24: Maximum intensity projection of confocal Z-stack live imaging of an

embryo containing *NB-ALFA-msGFP2> Ilp4_32x_ALFA-array/+*. Scale bar 10µm.

Movie S25: Maximum intensity projection of confocal tile-scan with Z-stack live
imaging of an embryo containing *NB-ALFA-msGFP2> Ilp4_32x_ALFA-array/+*. Scale
bar 100µm.

Movie S26: Single plane of a confocal live imaging of an embryo containing
twi_suntag_MS2_CRISPR/+; NB-ALFA_msGFP2/scFv-mScarlet x Ilp4_32x_ALFA-
array during n.c.14. Scale bar 5µm.

Movie S27: Single plane of a confocal live imaging of an embryo containing *NB-*
ALFA_msGFP2/scFv-mScarlet x Ilp4_32x_ALFA-array during n.c.14. Scale bar 5µm.

1708

Figure 2

a

GAL4   GAL4   GAL4

Loxp 5xUAS Loxp 5xUAS *P*-Pr *scFv-msGFP2*

UASp

b

Egg laying

strong expression and maternal deposition of scFv-msGFP2

nosGal4>UASp-scFv-msGFP2

c

time into n.c. 14

twi expression pattern

Dorsal

Ventral

early   early-mid   mid-late

scFv-msGFP2

MS2

merge

nc14 staging

nosGal4>UASp-scFv-msGFP2 x twi_suntag_MS2_CRISPR

d

Gastrulation furrow

scFv-msGFP2

MS2

merge

nosGal4>UASp-scFv-msGFP2 x twi_suntag_MS2_CRISPR

scFv-msGFP2 x twi_suntag_MS2_CRISPR

e

Gastrulation furrow

Gastrulation furrow   Gastrulation furrow   Gastrulation furrow

nosGal4>UASp-scFv-msGFP2 x yw

scFv-msGFP2 x twi_suntag_MS2_CRISPR

nosGal4>UASp-scFv-msGFP2 x twi_suntag_MS2_CRISPR

a

suntag V4 sequence (19aa) and linker (5aa):
EELLSKNYHLENEVARLKK GSGSG

ALFA-tag V1 sequence (15aa) and linker (5aa):
PSRLEEELRRRLTEP GSGSG

b
scFv_msGFP2
563 aa

NB-ALFA_msGFP2
456 aa

c
mRNA 5'UTR 32x multimerized ALFA-tag PSRLEEELRRRLTEP-...-PSRLEEELRRRLTEP- gene of interest 3'UTR

protein

nos EPr nos 5'UTR Nanobody ALFA msGFP2 GB1 NLS 3'UTR

d
mRNA 5'UTR ALFA-array Ilp4 3'UTR

Ilp4 expression pattern

NB-ALFA-msGFP2   ALFA-array   merge

NB-ALFA-msGFP2 x Ilp4_32x_ALFA-array

e
n.c. 14

NB-ALFA-msGFP2 x Ilp4_32x_ALFA-array

f
mRNA 5'UTR ALFA-array Ilp4 3'UTR
mRNA 5'UTR suntag twist 3'UTR

g
Ilp4 expression pattern   twi expression pattern

anti-ALFA   ALFA-array   anti-GCN4   merge

Ilp4_32x_ALFA-array ; twi_suntag_transgene

h
anti-ALFA
anti-GCN4
anti-ALFA anti-GCN4
merge

Ilp4_32x_ALFA-array ; twi_suntag_transgene

i
Ilp4 expression pattern   twi expression pattern

NB-ALFA-msGFP2   scFv-mScarlet   merge

twi_suntag_MS2_CRISPR/+;NB-ALFA-msGFP2/
scFv-mScarlet x Ilp4_32x_ALFA-array

j
Gastrulation furrow

NB-ALFA-msGFP2   scFv-mScarlet   merge

twi_suntag_MS2_CRISPR/+;NB-ALFA-msGFP2/scFv-mScarlet x
Ilp4_32x_ALFA-array

a

1.

coordinates
Chr14   101.959.597   102.057.848

DYNC1H1
5'UTR   3'UTR

CRISPR/Cas9
targeted insersion
12x_ALFA-array

2.

mRNA

12x multimerized ALFA-tag

5'UTR   PSRLEEELRRRLTEP-...............   DYNC1H1   3'UTR

3.

SFFV promoter   Nanobody ALFA   sfGFP   GB1 NLS

b

12X_ALFA-array-DYNC1H1

Control   Puromycin

spot per cell
Cont.   Puro.

c

12X_ALFA-array-DYNC1H1

NB-ALFA-sfGFP-GB1   DYNC1H1 smiFISH   Merge

d

SV40-NLS   MCP   StayGold   linker (SAGG)X5   StayGold

SV40-NLS   MCP   eGFP

MCP-tdStayGold
0min   30min   60min
120min   150min   180min

MCP-eGFP
0min   30min   60min
120min   150min   180min

e

Single molecule RNA florescence intensity (a.u.)

NLS-MCP-eGFP
NLS-MCP-tdStayGold

Time (min)

f

Number of detected RNA (% t0)

NLS-MCP-eGFP
NLS-MCP-tdStayGold

Time (min)

a

scFv-FP x twi_suntag_MS2_CRISPR

b

twi_suntag_MS2_CRISPR

c

scFv-msGFP2 x twi_suntag_MS2_CRISPR

scFv-msGFP2 x Ilp4_suntag

scFv-FP x twi_suntag_MS2_CRISPR

a

Gastrulation furrow

anti-GCN4   MS2   merge

twi_suntag_MS2_CRISPR

a

scFv-msGFP2 scFv-GreenLantern

early n.c.14   gastrulation   early n.c.14   gastrulation

nosGal4>UASp-scFv-FP

scFv-mAvic1   scFv-NeonGreen

early n.c.14   gastrulation   early n.c.14   gastrulation

nosGal4>UASp-scFv-FP

b

Absence of translation (*nosGal4>UASp-scFv-msGFP2* only)

early-mid n.c.14   mid-late n.c.14   late-gastrulation

× Experimental data — Fit

Presence of translation (*nosGal4>UASp-scFv-msGFP2 x twi_suntag_MS2_CRISPR*)

early-mid n.c.14   mid-late n.c.14   late-gastrulation

× Experimental data — Fit

c

Absence of translation   Presence of translation

- early-mid n.c.14
- mid-late n.c.14
- late-gastrulation

Molecular Brightness (kHz) vs Number of Molecules

a

UASz

GAL4 GAL4 GAL4 *hsp70* Pr

10xUAS *IVS-syn21*

UASy

GAL4 GAL4 GAL4 *hsp70* Pr

10xUAS *scFv-msGFP2*

b

scFv-msGFP2

early n.c.14 gastrulation

nosGal4>UASy-scFv-msGFP2

c

twi expression pattern

Dorsal

Ventral

n.c.14

early-mid mid-late

scFv-msGFP2

MS2

nosGal4>UASy-scFv-msGFP2 x twi_suntag_MS2_CRISPR

d

H2Av-RFP

scFv-msGFP2

merge

UASy-scFv-msGFP2/ his2Av-RFP

H2Av-RFP

scFv-msGFP2

merge

nulloGal4>UASy-scFv-msGFP2/ his2Av-RFP

a

Ilp4 expression pattern

twi expression pattern

anti-ALFA

ALFA-array

anti-GCN4

anti-ALFA
ALFA-array

merge

Ilp4_32x_ALFA-array ; twi_suntag_transgene

b

Ilp4 expression pattern

twi expression pattern

NB-ALFA-msGFP2   scFv-mScarlet   merge

NB-ALFA-msGFP2/scFv-mScarlet x Ilp4_32x_ALFA-array

c

Gastrulation furrow

msGFP2   scFv-mScarlet   merge

NB-ALFA-msGFP2/scFv-mScarlet x Ilp4_32x_ALFA-array