The ancestral shape of the access proton path of mitochondrial ATP synthases revealed by a split subunit-a

Jonathan E. Wong[1,2], Alena Zíková[1,2], Ondřej Gahura[1,3*]

[1]Institute of Parasitology, Biology Centre, Czech Academy of Sciences, 37005 České Budějovice, Czech Republic

[2]Faculty of Science, University of South Bohemia, 37005 České Budějovice, Czech Republic

[*]Corresponding author: gahura@paru.cas.cz

Abstract

The passage of protons across membranes through F_1F_o-ATP synthases spins their rotors and drives synthesis of ATP. While the principle of torque generation by proton transfer is known, the mechanisms and routes of proton access and release and their evolution are not fully understood. Here, we show that the entry site and path of protons in the lumenal half-channel of mitochondrial ATP synthases are largely defined by a short N-terminal α-helix of subunit-a. In *Trypanosoma brucei* and other Euglenozoa, the α-helix is part of another polypeptide chain that is a product of subunit-a gene fragmentation. This α-helix and other elements forming the proton pathway are widely conserved across eukaryotes and in Alphaproteobacteria, the closest extant relatives of mitochondria, but not in other bacteria. The α-helix blocks one of two proton routes found in *Escherichia coli*, resulting in the single proton entry site in mitochondrial and alphaproteobacterial ATP synthases. Thus, the shape of the access half-channel predates eukaryotes and originated in the lineage from which mitochondria evolved by endosymbiosis.

1 Introduction

F_1F_o-ATP synthases are rotary molecular machines that convert the proton-motive force across energy-transducing membranes of bacteria, chloroplasts and mitochondria to the chemical energy of ATP. The flow of protons down their electrochemical gradient through the membrane-intrinsic F_o subcomplex spins the enzyme's asymmetric rotor, whose rotation changes the conformation of three catalytic sites on the membrane-extrinsic F_1-ATPase subcomplex, resulting in synthesis of ATP. The transmembrane part of the rotor, the c-ring, consists of several copies of subunit-c. Proton translocation is based on sequential protonation of a glutamate residue of each protomer of the c-ring, rotation of the c-ring with the neutralized glutamates exposed to the hydrophobic environment of the phospholipid bilayer, and release of protons on the other side of the membrane. The second element crucial for proton passage is the stator subunit-a (also referred to as ATP6 or subunit-6). Canonical subunits-a contain six α-helices (h_1-h_6), five of which are largely embedded in the membrane. While h_1 is a vertical transmembrane helix, h_3-h_6 are horizontal, positioned at an angle ~20° relative to the plane of the membrane. Helices h_5-h_6 form a hairpin bending around the c-ring. The sites of proton binding and release from the c-ring glutamates are separated by a universally conserved arginine of h_5. The positively charged arginine prevents protons from leaking without c-ring rotation and induces their dissociation at the release site (reviewed in Walker 2017; Kuhlbrandt 2019; Guo and Rubinstein 2022). The proton-binding glutamates on the c-ring are accessible from the surface of the complex by two offset solvent-filled half-channels. In mitochondrial ATP synthases, the access half-channel opens into the lumen of cristae, hence it is called the lumenal half-channel. It is shaped by subunit-a and adjacent membrane elements, including lipids (Klusch, et al. 2017). Based on the high rate of proton translocation, it was proposed that delivery of protons to the c-ring is facilitated by a Grotthuss mechanism through a chain of coordinated water molecules and adjacent amino acid residues (Feniouk, et al. 2004). Recently,

the proton-hopping mechanism was supported by observations of cryoEM densities

corresponding to water molecules in the inner part of the channel in mammalian (Spikes, et al.

2020) and trypanosomal (Gahura, et al. 2022) mitochondrial ATP synthases. Molecular

dynamics simulations are also consistent with the role of coordinated water molecules in proton

translocation (Ivontsin, et al. 2022). Direct evidence for the Grotthuss mechanism of proton

transfer in both the inlet and outlet channels was provided by a combination of mutagenesis and

single-molecule rotation studies with the ATP synthase from *Escherichia coli* (Yanagisawa and

Frasch 2021).

Here, we explore the lumenal half-channel forming elements across species and show that

the N-terminal region of subunits-a form a short α-helix common to alphaproteobacterial and

vast majority of mitochondrial ATP synthases. In Euglenozoa, this helix is a part of a different

polypeptide chain encoded by a nuclear gene that is presumably the product of fragmentation

of the mitochondrial gene for subunit-a. We document that the helix defines the proton entry

site and contributes to the shaping of the inner part of the half-channel in modern eukaryotes

and their common ancestors.

Results and discussion

The short N-terminal α-helix of subunit-a is a widely conserved structural element of

mitochondrial ATP synthases

When examining the proton translocation elements of mitochondrial ATP synthases of two

divergent eukaryotes, *Saccharomyces cerevisiae* (Guo, et al. 2017) and *Trypanosoma brucei*

(Gahura, et al. 2022), we noticed a remarkable similarity between the N-terminal regions of

subunit-a in the yeast and subunit ATPEG3 in the parasite. The N-termini of both proteins form

a short single-turn α-helix, hereafter referred to as h_N, which is located at the layer

corresponding to the lumenal surface of the membrane and interacts with the horizontal helix

h_5 of subunit-a in the same manner in both structures. Downstream of h_N, yeast subunit-a and

trypanosomal ATPEG3 follow a similar path; however, after ten residues the yeast chain turns

back towards the first transmembrane helix h_1, thereby diverging from ATPEG3, which

continues to the periphery of the complex. The canonical trypanosomal subunit-a, which

contains h_1 and four characteristic horizontal membrane-intrinsic helices h_3-h_6, begins nine

amino acid residues upstream of h_1 in close proximity to the site of divergence between yeast

subunit-a and ATPEG3 (Fig. 1A). Based on the striking structural and sequence (see below)

similarity, we conclude that subunit-a in *T. brucei* has been split and ATPEG3 represents its N-

terminal fragment. To acknowledge the origin of ATPEG3, we introduce an alternative name

subunit-a_N.

The split of subunit-a in *T. brucei* occurred in the region corresponding to the loop

involved in ATP synthase dimerization in *S. cerevisiae* (Guo, et al. 2017). In *T. brucei*, the loop

is structurally replaced by a two-stranded β-sheet, formed by the N-terminus of subunit-a and

a β-strand of subunit-a_N (Fig. 1A,B). Unlike in yeast, this structure does not contribute to the

ATP synthase dimerization interface, which is provided by subunits g and e in this organism

(Gahura, et al. 2022). After splitting, subunit-a_N was extended at its C-terminus. The extension

interacts exclusively with lineage-specific subunits (ATPTB1, ATPTB6, ATPTB11,

ATPTB12) or with lineage-specific extensions of subunits shared with other ATP synthases

(subunit-i/j and k; Gahura, et al. 2022). It extends to the periphery of the membrane part of the

F_o-moiety, where it co-constitutes a euglenozoan-specific subcomplex, separated from the

conserved part by a lipid-filled cavity (Muhleip, et al. 2019; Gahura, et al. 2022). Consistently,

fragmented subunit-a is also discernible in the structure of ATP synthase from *Euglena gracilis*

in contrast to other mitochondrial and bacterial subunit-a exemplified by several selected

species (Fig. 1C; Muhleip, et al. 2019; Gahura, et al. 2021).

The sequence of h_N and flanking residues in yeast is highly similar to the N-terminal

sequence of subunit-a in all major groups of Alphaproteobacteria (Fig. 2, Supplementary table

S1 and Supplementary data 1&2), the closest extant relatives to mitochondria (Munoz-Gomez,

et al. 2022). We predicted the structure of subunit-a from *Rickettsia rickettsii* using AlphaFold

(Jumper, et al. 2021). The model resembles structures of bacterial and mitochondrial homologs

resolved experimentally and includes h_N located at exactly the same position as the h_N in *T.*

brucei and *S. cerevisiae* (Fig. 1C). Similarity of this region in ATP synthases from mitochondria

of two deeply branching eukaryotic lineages and the group of bacteria from which mitochondria

evolved strongly indicates that the observed architecture of the N-terminal segment of subunit-

a is ancestral.

We compared subunit-a sequences from 94 species covering all major eukaryotic

lineages, and structures from selected species determined by cryoEM or predicted by

AlphaFold. The sequence comparison documented that the split of subunit-a observed in *T.*

brucei and *E. gracilis* is restricted to Euglenozoa, because subunits-a from all representatives

of the sister lineage Heterolobosea contain a conserved sequence of h_N. Consistently, the

subunit-a_N was found in euglenids, diplonemids and kinetoplastids, the three lineages of

Euglenozoa (Sinha and Wideman 2023; Supplementary table S1, Supplementary data 3).

Further, the analysis revealed high sequence similarity in the N-terminal region, with most

sequences showing a degenerate sequence motif SPLEQF (Fig. 2, Supplementary table S1),

supporting the notion that presence of h_N represents an ancestral state. The only organisms

without the SPLEQF motif were most alveolates, chlorophycean algae *Polytomella* and

Chlamydomonas, and all metazoans (Fig. 2, Supplementary table S1). In alveolates and

chlorophycean algae, the N-terminal region including h_1 is divergent or absent, as documented

by cryoEM structures of ATP synthases from *Tetrahymena* (Flygaard, et al. 2020), *Polytomella*

(Murphy, et al. 2019) and *Toxoplasma* (Muhleip, et al. 2021). In *Polytomella*, subunit-a lacks

the entire N-terminal region including the transmembrane helix h_1, which is positionally replaced by the small transmembrane subunit Asa10 (Murphy, et al. 2019). However, Asa10 is unlikely to be a fragment of subunit-a, because it crosses the membrane in the opposite direction (Sinha and Wideman 2023). The missing helix h_N in *Polytomella* is positionally replaced by another conserved component, subunit-8 (Murphy, et al. 2019). Subunit-a in *Toxoplasma* is even more reduced and contains horizontal helices h_5 and h_6 only (Fig. 1C). On the contrary, subunit-a in *Tetrahymena* features a highly extended and structurally divergent N-terminus, which interacts predominantly with ciliate-specific ATP synthase components. Overall, the regions occupied by N-terminal parts of subunit-a in other ATP synthases are structurally highly divergent in *Toxoplasma* and *Tetrahymena*. The only alveolate with a canonical N-terminus of subunit-a is *Colponema* (Fig. 2), which supports the view that it represents a lineage sister to all other Alveolata (Tikhonenkov, et al. 2020).

In Metazoa, the N-terminal sequence of subunit-a differs from h_N in other groups, but invariably contains the motif FxxF (Fig. 2 and Supplementary table S1 and Supplementary Data 1). In experimentally determined and predicted structures of mammalian (Pinke, et al. 2020; Spikes, et al. 2020) and *Drosophila* subunit-a, respectively, this motif adopts the canonical fold of h_N (Fig. 1C), indicating that the ancestral architecture is retained despite sequence divergence. The only two exceptions are ctenophores and nematodes, represented by *Mnemiopsis leidyi* and *Caenorhabditis elegans*, respectively, whose subunit-a begins with helix h_1, like subunit-a in trypanosomes. In *C. elegans*, this observation is in line with a recent cryoelectron tomography study that showed that the nematode ATP synthase diverged architecturally from canonical ATP synthases from yeasts and mammals and might lack subunits 8, k and i/j (Buzzard, et al. 2023), which are otherwise widely conserved in eukaryotes (Sinha and Wideman 2023). Due to high conservancy of h_N across eukaryotes, we wondered whether an equivalent of h_N could be retained in nematodes, possibly as a product of subunit-a

gene split analogous to that observed in *T. brucei*. Such evolutionary independent event of

fission of a mitochondrial gene would not be unprecedented. The gene encoding the

mitoribosomal subunit uS3m has been fragmented in Ciliata (Swart, et al. 2012; Tobiasson and

Amunts 2020) and some Heterolobosea (Fu, et al. 2014), and the gene encoding Cox2, a subunit

of cytochrome oxidase, has been independently split at least three times in evolution,

specifically in alveolates, chlorophycean algae (Perez-Martinez, et al. 2001; Waller and Keeling

2006; Rodriguez-Salinas, et al. 2012), and the hymenopteran insect *Campsomeris* (Szafranski

2017a). Therefore, we searched the proteomes of Nematoda using several approaches (blast,

motif search, structure similarity search in AlphaFold database) to identify the protein

constituting the missing region including h_N, but we found no candidate, presumably due to the

short length of the region and limited sequence conservancy.

In summary, the vast majority of eukaryotes with the exception of most Alveolata, a group

of chlorophycean algae, and possibly Nematoda and Ctenophora have retained the ancestral

architecture of the region corresponding to the N-terminus of subunit-a.

Loss of transmembrane helix h_1 predetermines the transfer of subunit-a from the

mitochondrial to the nuclear genome

While the gene encoding subunit-a_N in Euglenozoa was transferred to the nuclear genome, the

subunit-a gene has been retained in the mitochondrial genome. Likewise, subunit-a in

nematodes, also starting with the first transmembrane helix h_1, is encoded mitochondrially. In

contrast, the gene of subunit-a in *Polytomella* and *Chlamydomonas*, which lacks h_1, has been

transferred to the nuclear genome. This suggests that while loss of h_1 allows the transfer of the

gene from the mitochondrial to the nuclear genome, the removal of the N-terminal fragment

upstream of h_1 is not sufficient for the relocation. Consistently, the reduced subunit-a in

Toxoplasma and related species are also nuclear encoded (Huet, et al. 2018; Muhleip, et al.

2021; Fig. 2). The last known organism with subunit-a encoded in the nuclear genome is the

ctenophore *M. leidyi*. The transmembrane helix prediction tools DeepTMHMM (Hallgren et

al., bioRxiv), MEMSAT (Jones, et al. 1994), and HMMTOP (Tusnady and Simon 2001) do not

predict h1 in the *M. leidyi* subunit-a. Alphafold does model h_1 in *M. leidyi*, but with low

confidence scores. Thus, it is unclear whether canonical h_1 is present in the *M. leidyi* subunit-a

or not. Together, these observations indicate that barriers associated with the insertion of h_1-

containing subunit-a to the inner mitochondrial membrane from the intermembrane space (or

with translocation of the protein into mitochondria prior its insertion from inside) may be a

factor limiting the transfer of the subunit-a gene from mitochondrial DNA to the nuclear

genome. We expanded a previous analysis of hydropathy of subunit-a (Muhleip, et al. 2021) to

all species included in our dataset. It revealed that some subunits-a (e.g., from *Nicotiana* or

Tetrahymena) retained in mitochondrial genomes show relatively low hydrophobicity similar

to those transferred to the nuclear genome (Fig. 2, Supplementary table S1), suggesting that the

reduced hydrophobicity is not the only prerequisite for the intracellular subunit-a gene transfer.

Regardless of the factor that allowed the subunit-a gene to be transferred, this is a very rare

event that has most likely occurred only three times in the evolution of eukaryotes described to

date, namely in the ancestors of the present-day myzozoans, ctenophores and

Chlamydomonadales, an order of chlorophycean algae including *Polytomella* and

Chlamydomonas (Fig. 2).

h_N of subunit-a contributes to the coordination of the chain of proton translocating water

molecules

The wide conservancy of h_N and its flanking residues (Fig. 2), whether within subunit-a or as

part of a fission product, implies the functional importance of this region. The h_N contributes to

coordination of the water chain in the lumenal proton half-channel in structurally characterized

ATP synthases from *Trypanosoma* (Gahura, et al. 2022), *S. cerevisiae* (Guo, et al. 2017) and

mammals (Pinke, et al. 2020; Spikes, et al. 2020) (Fig. 1B). One of the water molecules in the

Grotthuss water chain is coordinated by positionally equivalent His19 of h_N of *T. brucei*

subunit-a_N and Ser6 of h_N bovine subunit-a (Pinke, et al. 2020; Spikes, et al. 2020; Gahura, et

al. 2022). In the vast majority of mitochondrial subunits-a, this position is occupied by another

hydrogen bonding residue, glutamine, (Gln15 in *S. cerevisiae*; Figs. 1B&2 and Supplementary

table S1). Thus, h_N most likely contributes to coordination of water molecules in most

eukaryotic lineages and Alphaproteobacteria.

The shape of the inner access half-channel in bacteria resembles that of mitochondrial

ATP synthase (Montgomery, et al. 2021), and a density consistent with presence of a water

molecule in the inlet half-channel was observed in the cryoEM map of the *E. coli* ATP synthase

(Sobti, et al. 2020). Early mutagenesis studies revealed several subunit-a residues crucial for

enzymatic function of ATP synthase in *E. coli* (Cain and Simoni 1988; Lightowlers, et al. 1988;

Vik, et al. 1988; Cain and Simoni 1989; Howitt, et al. 1990; Eya, et al. 1991; Hartzog and Cain

1994; Hatch, et al. 1995). Most of these residues are localized in the proton half-channels or at

the interface between the subunit-a and the c-ring, implying their role in proton translocation.

To assess to what extent the inner constricted portion of the lumenal proton half-channel differs

between bacterial and diverse eukaryotic ATP synthases, we examined the conservancy of these

residues and compared their positions with the residues proposed to be involved directly in the

proton transfer or shown to coordinate ordered proton-transporting water molecules in

mitochondrial ATP synthases (Spikes, et al. 2020; Gahura, et al. 2022); Fig. 1B&D,

Supplementary figure 1). The only two invariable residues of subunit-a are the arginine of h5

separating the two half-channels and a glutamine of h6 (Arg210 and Gln245 in *E. coli*

numbering), juxtaposing with each other. Asn214 and Glu196 lining the proton channel in *E.*

coli are widely, yet not universally, conserved in mitochondrial ATP synthases. Functionally

important His245 in *E. coli* is replaced by an acidic residue, in most cases glutamate (Glu203,

Glu233 and Asp202 in bovine, *S. cerevisiae* and *T. brucei*, respectively), in vast majority of

mitochondrial subunits-a. This residue is assumed to pass protons from the chain of water

molecules to the c-ring either directly or by binding other, as yet unidentified, water molecules

(Spikes, et al. 2020). Other functional residue, Glu219 in *E. coli*, is in most mitochondrial ATP

synthases replaced by histidine (His168/155 in bovine/*T. brucei*), which binds one of the

molecules in the water chain in the mammalian and trypanosomal enzymes (Spikes, et al. 2020;

Gahura, et al. 2022). Thus, the positions of these two key histidine and glutamate residues are

swapped between *E. coli* and mitochondrial ATP synthases. Nevertheless, due to the properties

of these residues, both configurations are compatible with the Grotthus mechanism of proton

translocation. Thus, the segment of the half-channel proximal to the c-ring most likely does not

differ between bacterial and mitochondrial ATP synthases, supporting the view that the

Grotthuss mechanism represents an ancestral and conserved mode of proton transfer in ATP

synthases. In vast majority of mitochondrial ATP synthases and their alphaproteobacterial

ancestors, h_N is a key component of the network of interactions involved in proton translocation.

The entry point of the proton access half-channel in mitochondrial and

alphaproteobacterial ATP synthases is defined by h_N of subunit-a

Notably, the *E. coli* ATP synthase (Sobti, et al. 2019) and ATP synthases in bacteria from

various groups outside Alphaproteobacteria (*Mycobacterium;* Guo, et al. 2021; Montgomery,

et al. 2021, and *Acinetobacter*; Demmer, et al. 2022) contain an α-helix at the N-terminus of

the subunit-a, which is longer and sequentially dissimilar to h_N in mitochondrial ATP synthases

and positionally shifted further from the loop connecting helices h_3 and h_4 (Fig. 1C). As

proposed based on the structure of the *Acinetobacter* ATP synthase, the position of the N-

terminal helix determines whether protons enter to the half-channel using the "front" or "rear"

entry site as viewed from the c-ring (Demmer, et al. 2022). In *Acinetobacter*, the protons have

been proposed to enter from the side facing the c-ring through the front entry located between

the N-terminal helix and h_5 of subunit-a. In *E. coli*, the protons are funneled from the periplasm

to an entry point located between the N-terminal helix of subunit-a and the N-terminus of

subunit-b (Yanagisawa and Frasch 2021), termed the rear entry (Demmer, et al. 2022). The

aqueous funnel is lined with polar and charged residues, which are contributed by the N-

terminal helix (Asp10, His14, His15) or h_1 (Glu131, His132) and are not conserved in

mitochondrial ATP synthases, and narrows to Glu219, presumably the first residue involved in

the proton hopping. Notably, molecular dynamics simulations strongly suggest that protons can

be delivered to Glu219 in *E. coli* also using another aqueous cavity (Ivontsin, et al. 2022),

corresponding to the front entry in *A. baumanii*. Because *A. baumanii* and *E. coli* are closely

related species and their subunits-a and adjacent regions are highly similar in both sequence

and structure (Fig. 1, Supplementary data 1), we argue that both proton entry sites are most

likely present also in *A. baumanii* (Fig. 3A).

Unlike in *E. coli* and *A. baumanii*, in yeast and bovine mitochondrial ATP synthases

protons use exclusively the rear opening, formed by a narrow cleft between h_N and the C-

terminal tail of subunit-f(Spikes, et al. 2020; Demmer, et al. 2022) (Fig. 3A,B). In the yeast and

bovine structures, as well as in the structure from *T. brucei* and the predicted structure from

Rickettsia, the position of the front entry is blocked by h_N (Fig 3B,C), specifically by the

phenylalanine residue of the SPLEQF motif that is nearly invariably conserved in all h_N

sequences included in our analysis (Fig 2 and Supplementary table S1). Therefore, we reason

that the rear entry is common to all h_N-containing ATP synthases. In addition, the conserved

phenylalanine clashes with the side chain of Glu219 in *E. coli* (Fig. 3B), which is the residue,

at which the front and rear paths converge. Localization of coordinated water molecules in the

part of the trypanosomal inlet channel that is more distal from the c-ring than Glu219 in *E. coli*

(Fig. 1B) suggests that mitochondrial ATP synthases might use an extended water chain

compared to bacteria.

Taken together, h_N contributes to shaping of both the surface-exposed and the inner

portions of the access proton half-channel, underlying its almost universal occurrence in

mitochondrial ATP synthases.

Concluding remarks

By combination of analyses of available ATP synthase structures with sequence comparison

and structure prediction across species, we documented that the proton access path and proton

transfer mechanism in the access half-channel is conserved in majority of mitochondrial ATP

synthases. The key structural determinant, so far largely overlooked, is h_N of subunit-a, which

defines the entry to the access half channel and contributes to hydrogen bonding of water

molecules in the Grotthuss chain (Fig. 1C). Unlike ATP synthase from *E. coli* and *A. baumanii*,

which feature two possible proton entry sites, mitochondrial and alphaproteobacterial

counterparts have only one access to the proton inlet half-channel, co-defined by h_N of subunit-

a. Functional consequences of this difference remain to be elucidated.

Uniquely, F_1F_o-ATP synthase in *T. brucei* mitochondria contains two functionally central

components, the subunits α and a, that are both constituted of two separate polypeptide chains.

While the subunit-α is cleaved into two fragments posttranslationally (Gahura, et al. 2018;

Montgomery, et al. 2018), the subunit-a is split on the gene level. Our finding of a split subunit-

a in *T. brucei* complements recently reported cases of mitochondrial gene fragmentation

revealed by structures of mitochondrial ribosomes (uS4m and uS7m in *Polytomella*; Tobiasson,

et al. 2022) and respiratory chain supercomplexes (Cox3 and Nad5 in *Tetrahymena*; Mühleip,

et al. 2022; Zhou, et al. 2022), and several occurrences reported earlier (reviewed in Szafranski

2017b). The transfer of the same gene from the mitochondrial to the nuclear genome

independently in several lineages facilitates the inference of properties of encoded proteins that are incompatible with such events. We inferred that the transmembrane helix h_1 might represent such an obstacle in the case of subunit-a.

Material and Methods

Subunit-a sequences were searched by hmmsearch using the alignment of subunit-a sequences from *Bos taurus*, *Saccharomyces cerevisiae*, *Rickettsia rickettsii* and *Colponema vietnamica* as a query. The sequences from Ciliata and Apicomplexa were searched by phmmer using sequences from *Tetrahymena* and *Toxoplasma*, respectively, as queries. Both searches were performed on the HMMER web server (Potter, et al. 2018) in the Uniprot database. Amino acid sequence alignments were constructed by MUSCLE (Edgar 2004) and corrected manually using available structures as references. Some sequences of subunits a and a_N from Euglenozoa were obtained from (Sinha and Wideman 2023). Structures were predicted by AlphaFold (Jumper, et al. 2021), and superposed by the Matchmaker tool and visualized in ChimeraX (Goddard, et al. 2018). Hydrophobicity was calculated as grand average of hydropathy using hydropathy indices from Kyte and Doolittle 1982.

Acknowledgements

We thank Anzhelika Butenko for the initial search of subunit-a homologs. This work was supported by Czech Science Foundation grant 20-04150Y to O.G. and by European Regional Development Fund (ERDF) and Ministry of Education, Youth and Sport (MEYS) project CZ.02.1.01/0.0/0.0/16_019/0000759 and by the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Program (grant agreement no. 101044951) to A.Z.

Author contributions

Conceptualization: O.G. and A.Z.; investigation and data analyses: J.E.W. and O.G.; writing –

original draft: O.G.; writing – review and editing: J.E.W, A.Z., O.G.; funding acquisition: O.G.

and A.Z.

Declaration of interests

They authors declare they have no competing interests.

References

Burki F, Roger AJ, Brown MW, Simpson AGB. 2020. The New Tree of Eukaryotes. Trends

Ecol Evol 35:43-55.

Buzzard E, McLaren M, Bragoszewski P, Brancaccio A, Ford H, Daum B, Kuwabara P,

Collinson I, Gold VAM, unpublished data,

https://www.biorxiv.org/content/10.1101/2023.02.02.526626v2

Cain BD, Simoni RD. 1988. Interaction between Glu-219 and His-245 within the a subunit of

F_1F_o-ATPase in *Escherichia coli*. Journal of Biological Chemistry 263:6606-6612.

Cain BD, Simoni RD. 1989. Proton Translocation by the F1FOATPase of Escherichia coli.

Journal of Biological Chemistry 264:3292-3300.

Demmer JK, Phillips BP, Uhrig OL, Filloux A, Allsopp LP, Bublitz M, Meier T. 2022.

Structure of ATP synthase from ESKAPE pathogen *Acinetobacter baumannii*. Sci Adv

8:eabl5966.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 32:1792-1797.

Eya S, Maeda M, Futai M. 1991. Role of the carboxyl terminal region of H^+-ATPase F_oF_1 a

subunit from *Escherichia coli*. Arch Biochem Biophys 284:71-77.

Feniouk BA, Kozlova MA, Knorre DA, Cherepanov DA, Mulkidjanian AY, Junge W. 2004.

The proton-driven rotor of ATP synthase: ohmic conductance (10 fS), and absence of voltage

gating. Biophys J 86:4094-4109.

Flygaard RK, Muhleip A, Tobiasson V, Amunts A. 2020. Type III ATP synthase is a symmetry-

deviated dimer that induces membrane curvature through tetramerization. Nat Commun

11:5342.

Fu CJ, Sheikh S, Miao W, Andersson SG, Baldauf SL. 2014. Missing genes, multiple ORFs,

and C-to-U type RNA editing in *Acrasis kona* (Heterolobosea, Excavata) mitochondrial DNA.

Genome Biol Evol 6:2240-2257.

Gahura O, Hierro-Yap C, Zikova A. 2021. Redesigned and reversed: architectural and

functional oddities of the trypanosomal ATP synthase. Parasitology 148:1151-1160.

Gahura O, Muhleip A, Hierro-Yap C, Panicucci B, Jain M, Hollaus D, Slapnickova M, Zikova

A, Amunts A. 2022. An ancestral interaction module promotes oligomerization in divergent

mitochondrial ATP synthases. Nat Commun 13:5989.

Gahura O, Subrtova K, Vachova H, Panicucci B, Fearnley IM, Harbour ME, Walker JE, Zikova

A. 2018. The F_1-ATPase from *Trypanosoma brucei* is elaborated by three copies of an

additional p18-subunit. FEBS J 285:614-628.

Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE. 2018.

UCSF ChimeraX: Meeting modern challenges in visualization and analysis. Protein Sci 27:14-

25.

Guo H, Bueler SA, Rubinstein JL. 2017. Atomic model for the dimeric F_o region of

mitochondrial ATP synthase. Science 358:936-940.

Guo H, Courbon GM, Bueler SA, Mai J, Liu J, Rubinstein JL. 2021. Structure of mycobacterial

ATP synthase bound to the tuberculosis drug bedaquiline. Nature 589:143-147.

Guo H, Rubinstein JL. 2022. Structure of ATP synthase under strain during catalysis. Nat

Commun 13:2232.

Hartzog PE, Cain BD. 1994. Second-site suppressor mutations at glycine 218 and histidine 245

in the alpha subunit of F_1F_o ATP synthase in *Escherichia coli*. Journal of Biological Chemistry

269:32313-32317.

Hatch LP, Cox GB, Howitt SM. 1995. The essential arginine residue at position 210 in the alpha

subunit of the *Escherichia coli* ATP synthase can be transferred to position 252 with partial

retention of activity. J Biol Chem 270:29407-29412.

Howitt SM, Lightowlers RN, Gibson F, Cox GB. 1990. Mutational analysis of the function of

the a-subunit of the F_oF_1-ATPase of *Escherichia coli*. Biochim Biophys Acta 1015:264-268.

Huet D, Rajendran E, van Dooren GG, Lourido S. 2018. Identification of cryptic subunits from

an apicomplexan ATP synthase. Elife 7.

Ivontsin L, Mashkovtseva E, Nartsissov Y. 2022. Insights on the proton translocation pathways

in F_oF_1-ATP synthase using molecular dynamics simulations. Arch Biochem Biophys

717:109135.

Jones DT, Taylor WR, Thornton JM. 1994. A model recognition approach to the prediction of

all-helical membrane protein structure and topology. Biochemistry 33:3038-3049.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates

R, Zidek A, Potapenko A, et al. 2021. Highly accurate protein structure prediction with

AlphaFold. Nature 596:583-589.

Klusch N, Murphy BJ, Mills DJ, Yildiz O, Kuhlbrandt W. 2017. Structural basis of proton

translocation and force generation in mitochondrial ATP synthase. Elife 6.

Kuhlbrandt W. 2019. Structure and Mechanisms of F-Type ATP Synthases. Annu Rev Biochem

88:515-549.

Kyte J, Doolittle RF. 1982. A simple method for displaying the hydropathic character of a

protein. J Mol Biol 157:105-132.

Lightowlers RN, Howitt SM, Hatch L, Gibson F, Cox G. 1988. The proton pore in the

Escherichia coli F_oF_1-ATPase: substitution of glutamate by glutamine at position 219 of the

alpha-subunit prevents F_o-mediated proton permeability. Biochim Biophys Acta 933:241-248.

Montgomery MG, Gahura O, Leslie AGW, Zikova A, Walker JE. 2018. ATP synthase from

Trypanosoma brucei has an elaborated canonical F_1-domain and conventional catalytic sites.

Proc Natl Acad Sci U S A 115:2102-2107.

Montgomery MG, Petri J, Spikes TE, Walker JE. 2021. Structure of the ATP synthase from

Mycobacterium smegmatis provides targets for treating tuberculosis. Proc Natl Acad Sci U S A

118.

uhleip A, Flygaard RK, Baradaran R, Haapanen O, Gruhl T, Tobiasson V, Marechal A, Sharma

V, Amunts A. 2023. Structural basis of mitochondrial membrane bending by the I-II-III$_2$-IV$_2$

supercomplex. Nature 615:934-938.

Muhleip A, Kock Flygaard R, Ovciarikova J, Lacombe A, Fernandes P, Sheiner L, Amunts A.

2021. ATP synthase hexamer assemblies shape cristae of *Toxoplasma* mitochondria. Nat

Commun 12:120.

Muhleip A, McComas SE, Amunts A. 2019. Structure of a mitochondrial ATP synthase with

bound native cardiolipin. Elife 8.

Munoz-Gomez SA, Susko E, Williamson K, Eme L, Slamovits CH, Moreira D, Lopez-Garcia

P, Roger AJ. 2022. Site-and-branch-heterogeneous analyses of an expanded dataset favour

mitochondria as sister to known Alphaproteobacteria. Nat Ecol Evol 6:253-262.

Murphy BJ, Klusch N, Langer J, Mills DJ, Yildiz O, Kuhlbrandt W. 2019. Rotary substates of

mitochondrial ATP synthase reveal the basis of flexible F_1-F_o coupling. Science 364.

Perez-Martinez X, Antaramian A, Vazquez-Acevedo M, Funes S, Tolkunova E, d'Alayer J,

Claros MG, Davidson E, King MP, Gonzalez-Halphen D. 2001. Subunit II of cytochrome c

oxidase in Chlamydomonad algae is a heterodimer encoded by two independent nuclear genes.

J Biol Chem 276:11302-11309.

Pinke G, Zhou L, Sazanov LA. 2020. Cryo-EM structure of the entire mammalian F-type ATP

synthase. Nat Struct Mol Biol 27:1077-1085.

Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. 2018. HMMER web server: 2018

update. Nucleic Acids Res 46:W200-W204.

Rodriguez-Salinas E, Riveros-Rosas H, Li Z, Fucikova K, Brand JJ, Lewis LA, Gonzalez-

Halphen D. 2012. Lineage-specific fragmentation and nuclear relocation of the mitochondrial

cox2 gene in chlorophycean green algae (Chlorophyta). Mol Phylogenet Evol 64:166-176.

Sinha SD, Wideman JG. 2023. The persistent homology of mitochondrial ATP synthases.

iScience 26.

Sobti M, Ishmukhametov R, Bouwer JC, Ayer A, Suarna C, Smith NJ, Christie M, Stocker R,

Duncan TM, Stewart AG. 2019. Cryo-EM reveals distinct conformations of *E. coli* ATP

synthase on exposure to ATP. Elife 8.

Sobti M, Walshe JL, Wu D, Ishmukhametov R, Zeng YC, Robinson CV, Berry RM, Stewart

AG. 2020. Cryo-EM structures provide insight into how *E. coli* F_1F_o ATP synthase

accommodates symmetry mismatch. Nat Commun 11:2615.

Spikes TE, Montgomery MG, Walker JE. 2020. Structure of the dimeric ATP synthase from

bovine mitochondria. Proc Natl Acad Sci U S A 117:23519-23526.

Swart EC, Nowacki M, Shum J, Stiles H, Higgins BP, Doak TG, Schotanus K, Magrini VJ,

Minx P, Mardis ER, et al. 2012. The *Oxytricha trifallax* mitochondrial genome. Genome Biol

Evol 4:136-154.

Szafranski P. 2017a. Evolutionarily recent, insertional fission of mitochondrial cox2 into

complementary genes in bilaterian Metazoa. BMC Genomics 18:269.

Szafranski P. 2017b. Intercompartmental Piecewise Gene Transfer. Genes (Basel) 8.

Tikhonenkov DV, Mikhailov KV, Gawryluk RMR, Belyaev AO, Mathur V, Karpov SA,

Zagumyonnyi DG, Borodina AS, Prokina KI, Mylnikov AP, et al. 2022. Microbial predators

form a new supergroup of eukaryotes. Nature 612:714-719.

Tikhonenkov DV, Strassert JFH, Janouskovec J, Mylnikov AP, Aleoshin VV, Burki F, Keeling

PJ. 2020. Predatory colponemids are the sister group to all other alveolates. Mol Phylogenet

Evol 149:106839.

Tobiasson V, Amunts A. 2020. Ciliate mitoribosome illuminates evolutionary steps of

mitochondrial translation. Elife 9.

Tobiasson V, Berzina I, Amunts A. 2022. Structure of a mitochondrial ribosome with

fragmented rRNA in complex with membrane-targeting elements. Nat Commun 13:6132.

Tusnady GE, Simon I. 2001. The HMMTOP transmembrane topology prediction server.

Bioinformatics 17:849-850.

Vik SB, Cain BD, Chun KT, Simoni RD. 1988. Mutagenesis of the alpha subunit of the F_1F_0-

ATPase from *Escherichia coli*. Mutations at Glu-196, Pro-190, and Ser-199. Journal of

Biological Chemistry 263:6599-6605.

Walker JE. 2017. Structure, Mechanism and Regulation of ATP Synthases. In: Wikstrom M,

editor: The Royal Society of Chemistry.

Waller RF, Keeling PJ. 2006. Alveolate and chlorophycean mitochondrial cox2 genes split

twice independently. Gene 383:33-37.

Yanagisawa S, Frasch WD. 2021. pH-dependent 11 degrees F_1Fo ATP synthase sub-steps

reveal insight into the F_o torque generating mechanism. Elife 10.

Zhou L, Maldonado M, Padavannil A, Guo F, Letts JA. 2022. Structures of *Tetrahymena*'s

respiratory chain reveal the diversity of eukaryotic core metabolism. Science 376:831-839

Figures and legends

Figure 1. Split subunit-a of mitochondrial ATP synthase in *Trypanosoma brucei* retains

ancestral structure and shapes the access proton half-channel (A) Comparison of the

structures of subunit-a in *S. cerevisiae* and *E. coli* and subunits a and a_N in *T. brucei*. The c-

rings are shown as transparent surface. In the structure from *S. cerevisiae*, a fragment of the

subunit-a from the other ATP synthase protomer is shown (su-a'). **(B)** The close-up view of the

structures in (A). Conserved residues of h_N and other selected conserved residues are shown as

sticks. The residues labeled in red are directly involved in proton translocation or coordinate

water molecules in the Grotthuss chain. Water molecules inferred from cryoEM density are

shown as yellow balls in the *T. brucei* structure. **(C)** Structures of subunit-a of mitochondrial

and bacterial ATP synthases from selected species. The structure from *Rickettsia* was predicted

by AlphaFold.

Figure 2. Helix h_N is conserved in most mitochondrial ATP synthases and

Alphaproteobacteria. For individual lineages mapped on the phylogenetic tree based on

(Burki, et al. 2020; Tikhonenkov, et al. 2022) following features are shown: number of species

included in the analyses (in parentheses); genome, in which subunit-a/a_N is encoded ("mt" –

mitochondrial, "n" – nuclear); presence or absence of helices h_N and h_1 of subunit-a/a_N; selected

representative species used in the analyses presented in the following columns (for species

shown in bold a cryoEM structure of ATP synthase has been reported); sequence of h_N and

flanking residues (universally conserved phenylalanine is shown in red, other conserved

residues of the SPLEQF motif, or the FXXF motif in Metazoa, are shown in bold); average

hydrophobicity of the entire subunit-a (FL), region h_1-h_6, and region h_5-h_6 (blue – more

hydrophobic, red – less hydrophobic).

Figure 3. The proton entry point and path in the access half-channel is co-determined by

h_N. (A) Comparison of proton paths in the access half-channel of the bovine mitochondrial ATP

synthases with *Escherichia coli* and *Acinetobacter baumanii* counterparts. The c-ring is shown

as white surface. Blue and red arrows indicate the path through the front and rear entry,

respectively. **(B)** Superposition of the structures from *E. coli* and bovine mitochondria shows

that h_N is positionally shifted compared to the N-terminal helix of subunit-a in *E. coli* and that

the conserved phenylalanine occupies the position of Glu219 in *E. coli*. Selected functionally

important residues are shown as sticks in panels (A) and (B). **(C)** The position of the proposed

front proton entry to the access half-channel in *A. baumanii* ATP synthase(Demmer, et al. 2022)

is blocked by conserved phenylalanine in h_N of mitochondrial and alphaproteobacterial ATP

synthases. A view of the front entry (red arrowhead) to the half channel in the *A. baumanii*

subunit-a (surface representation) from the c-ring shown individually and superposed with

subunit-a from other ATP synthases (cartoon representation): *S. cerevisiae*, the AlphaFold

model of *Rickettsia*, and subunits a and a_N from *T. brucei*. The conserved phenylalanine is

shown as sticks.